



**HAL**  
open science

# A note on the antisymmetry in the speed of a random walk in reversible dynamic random environment

Oriane Blondel

► **To cite this version:**

Oriane Blondel. A note on the antisymmetry in the speed of a random walk in reversible dynamic random environment. 2022. hal-03828998v1

**HAL Id: hal-03828998**

**<https://hal.science/hal-03828998v1>**

Preprint submitted on 25 Oct 2022 (v1), last revised 9 Jan 2023 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A note on the antisymmetry in the speed of a random walk in reversible dynamic random environment

Oriane Blondel

October 17, 2022

## Abstract

In this short note, we prove that  $v(-\varepsilon) = -v(\varepsilon)$ . Here,  $v(\varepsilon)$  is the speed of a one-dimensional random walk in a dynamic *reversible* random environment, that jumps to the right (resp. to the left) with probability  $1/2 + \varepsilon$  (resp.  $1/2 - \varepsilon$ ) if it stands on an occupied site, and vice-versa on an empty site. We work in any setting where  $v(\varepsilon), v(-\varepsilon)$  are well-defined, *i.e.* a weak LLN holds. The proof relies on a simple coupling argument that holds only in the discrete setting.

## 1 Introduction

We consider the so-called “ $\varepsilon$ -random walk”: a random walk in one-dimensional dynamic random environment with two values that jumps to the right (resp. to the left) with probability  $1/2 + \varepsilon$  (resp.  $1/2 - \varepsilon$ ) if it stands on an occupied site, and vice-versa on an empty site (Figure 1).




Figure 1: Jump rates for the  $\varepsilon$ -random walk.

A lot of energy has been devoted to describing the behavior of this  $\varepsilon$ -RW in various random environments, mainly to find whether it satisfies the usual limit theorems (LLN, CLT...). This is generally a hard problem, since the environment seen from the  $\varepsilon$ -RW is highly non-stationary (even when the environment is). Additionally, it belongs to the class of nestling random walks, which do not have an a-priori preferred direction. There are known settings in which the LLN does not hold, mainly due to the presence of traps [Sol75], [BHT18, Section 9]. Cases in which the LLN has been shown include perturbative regimes (small  $\varepsilon$ ) [ABF18], well-known environments like the exclusion process [HS15, HKT20] or the contact process [MV15], uniform mixing hypotheses [Z04, AdHR11, RV13], or fast enough decay of correlations [BHT18].

When the random walk does satisfy a law of large numbers, let us call  $v(\varepsilon)$  its asymptotic speed (we will simply say that  $v(\varepsilon)$  is well-defined). We observed in [ABF16] that, if the environment is given by a reversible Markov process with positive spectral gap, and  $|\varepsilon|$  is smaller than the spectral gap,  $v$  –in addition to being well-defined– satisfies the antisymmetry property

$$v(-\varepsilon) = -v(\varepsilon). \quad (1)$$

This property also holds in higher dimensions for random walks with a certain symmetry property (Assumption 3 in [ABF16]). Simulations moreover suggest that this property extends out of the perturbative regime of [ABF16] (where the law of large numbers was not yet even established; see Figure 3 in [ABF16]). The proof relies on an expansion of the speed in  $\varepsilon$ , in which the terms of even degrees are shown to cancel due to reversibility.

It is worth pointing out that this antisymmetry does not seem to follow from obvious symmetry properties of the system. Indeed, it is tempting at first sight to think that the  $(-\varepsilon)$ -random walk reversed in time should have the same distribution as the  $\varepsilon$ -random walk, from which (1) would immediately follow. But closer inspection reveals that this is not true and those two processes have very different trajectories in general (see for instance Figure 4 in [ABF16]). Rather, (1) holds iff the speed of the  $\varepsilon$ -random walk is the same in the reversible environment and in its image under the mirror symmetry  $x \mapsto -x$ . While this property is obvious for reversible systems invariant under mirror symmetry (e.g. simple symmetric exclusion process), there exist reversible, translation invariant processes which have no mirror-invariance (e.g. the East model).

To state our main result, let  $(X_t^\varepsilon)_{t \geq 0}$  be the trajectory of a  $\varepsilon$ -RW started from 0. A proper construction of this object is given in Section 2.

**Theorem 1.** *Assume the environment is a translation invariant and reversible Markov process. Moreover, assume that  $X^\varepsilon$  and  $X^{-\varepsilon}$  satisfy a weak law of large numbers, i.e. there exist  $v(\varepsilon), v(-\varepsilon)$  such that*

$$\frac{1}{t} X_t^{\pm\varepsilon} \xrightarrow[t \rightarrow \infty]{\mathbb{P}} v(\pm\varepsilon). \quad (2)$$

Then

$$v(-\varepsilon) = -v(\varepsilon). \quad (3)$$

Checking that the weak law of large numbers holds is generally not easy. In [BHT18], it was established under the condition that the environment has polynomially decaying covariances with a high enough exponent. This includes Markov processes with positive spectral gap. The East model [JE91, FMRT13] is an example of such a model with no mirror symmetry.

The proof presented here relies on a coupling valid only for the discrete time  $\varepsilon$ -RW, which in turn allows to deduce the result for the continuous time. The coupling relies heavily on the fact that we consider nearest-neighbor trajectories in dimension 1. However, as mentioned above, Theorem 1 is expected to hold also in higher dimensions and for random walks allowing longer range jumps under appropriate symmetry assumptions [ABF16]. Outside the perturbative region, this is still an open problem.

## 2 Setting and construction of the random walk

We now give explicit constructions of the  $\varepsilon$ -RW, in discrete and continuous time. In the following,  $\mathbb{L} = \mathbb{R}_+$  when we consider the continuous time setting,  $\mathbb{L} = \mathbb{N}$  in the discrete time setting.

### 2.1 Environment

The environment is given by a collection  $\eta = (\eta_t(x))_{(x,t) \in \mathbb{Z} \times \mathbb{L}}$  such that  $\eta_t(x) \in \{0, 1\}$  for all  $(x, t) \in \mathbb{Z} \times \mathbb{L}$ . We assume

1. **(Stationarity)** The distribution of the environment is invariant under space-time shifts, *i.e.*

$$\forall x \in \mathbb{Z}, t \in \mathbb{L}, \quad (\eta_{t+s}(x + \cdot))_{s \in \mathbb{L}} = (\eta_s(\cdot))_{s \in \mathbb{L}} \quad \text{in distribution.} \quad (4)$$

2. **(Reversibility)** The environment is given by a reversible process  $\eta = (\eta_t)_{t \in \mathbb{L}}$  on  $\{0, 1\}^{\mathbb{Z}}$ , *i.e.*

$$\text{for all } T \in \mathbb{L} \setminus \{0\}, \quad (\eta_{T-t})_{t \in [0, T] \cap \mathbb{L}} = (\eta_t)_{t \in [0, T] \cap \mathbb{L}} \quad \text{in distribution.} \quad (5)$$

### 2.2 $\varepsilon$ -RW in discrete time

Let us consider a collection  $U = (U_{x,n})_{x \in \mathbb{Z}, n \in \mathbb{N}}$  of iid  $\mathcal{U}([0, 1])$  random variables. With the environment  $\eta$  and the collection  $U$  we associate a set  $(A_{x,n}^\varepsilon)_{x \in \mathbb{Z}, n \in \mathbb{N}}$  (that will prescribe the directions taken by the RW) in the following way:

$$A_{x,n}^\varepsilon = \begin{cases} +1 & \text{if } (\eta_n(x) = 1 \text{ and } U_{x,n} \leq \frac{1}{2} + \varepsilon) \text{ or } (\eta_n(x) = 0 \text{ and } U_{x,n} \leq \frac{1}{2} - \varepsilon), \\ -1 & \text{else.} \end{cases} \quad (6)$$

For  $x \in \mathbb{Z}$ , we denote by  $(X_n^{x,\varepsilon})_{n \in \mathbb{N}}$  the  $\varepsilon$ -RW started at  $x$ . It is built iteratively as follows:

1.  $X_0^{x,\varepsilon} = x$ ;
2. if  $X_n^{x,\varepsilon} = y$ ,  $X_{n+1}^{x,\varepsilon} = y + A_{x,n}^\varepsilon$ .

Note that by translation invariance of the environment,  $X_n^{x,\varepsilon} - x \stackrel{(d)}{=} X_n^{0,\varepsilon}$  for all  $x \in \mathbb{Z}$ . Also,  $X_n^{x,\varepsilon}$  has the same parity as  $n + x$ .

### 2.3 $\varepsilon$ -RW in continuous time

We could use a similar construction, but it will be more convenient to alter it a little. Let  $U = (U_n)_{n \in \mathbb{N}}$  be a collection of iid  $\mathcal{U}([0, 1])$  random variables and  $T = (T_n)_{n \in \mathbb{N}}$  a PPP(1) on  $\mathbb{R}_+$ . Assume  $(U, T, \eta)$  to be independent. The continuous time  $\varepsilon$ -RW started at  $x$ ,  $(X_t^{x,\varepsilon})_{t \in \mathbb{R}_+}$  is built as follows:

1.  $X_0^{x,\varepsilon} = x$ ;
2. if  $X_{t^-}^{x,\varepsilon} = y$  and  $y \notin \{T_n, n \in \mathbb{N}\}$ ,  $X_t^{x,\varepsilon} = y$ ;
3. if  $X_{t^-}^{x,\varepsilon} = y$  and  $t = T_n$ ,

$$X_t^{x,\varepsilon} = \begin{cases} y + 1 & \text{if } (\eta_t(y) = 1 \text{ and } U_n \leq \frac{1}{2} + \varepsilon) \text{ or } (\eta_t(y) = 0 \text{ and } U_n \leq \frac{1}{2} - \varepsilon), \\ y - 1 & \text{else.} \end{cases} \quad (7)$$

### 3 Proof of Theorem 1 in the discrete setting

A key remark is that the reversibility assumption allows to construct jointly a  $\varepsilon$ -RW and a backwards  $(-\varepsilon)$ -RW, that is a  $(-\varepsilon)$ -RW evolving on the reversed environment process  $(\eta_{N-n})_{n \leq N}$  for  $N \in \mathbb{N}$ .

In order to simplify the notations, let us fix  $\varepsilon \in [-1, 1]$ , and  $N \in \mathbb{N}$ . We let  $X = X^{0, \varepsilon}$ , and for  $x \in \mathbb{Z}$  we let  $(\widehat{X}_n)_{n \leq N}$  be constructed as follows.

1.  $\widehat{X}_N = x$ ;
2. for  $n < N$ , if  $\widehat{X}_{N-n} = y$ ,  $\widehat{X}_{N-n-1} = y - A_{y, N-n}^\varepsilon$ .

Note that the construction uses the same collection  $A^\varepsilon$  as the construction of  $X$ .

Reversibility of the environment clearly gives us the following property.

**Lemma 1.**  $(\widehat{X}_n)_{n \leq N} \stackrel{(d)}{=} (X_n^{x, -\varepsilon})_{n \leq N}$ .

*Proof.* This follows from the reversibility of  $\eta$ . □

It is also not difficult to check that  $\widehat{X}$  and  $X$  cannot cross, in the following sense.

**Lemma 2.** *Let  $X$  and  $\widehat{X}$  be built using the same collection  $A^\varepsilon$ . Then, if  $\widehat{X}_0$  and  $X_0$  have the same parity,*

$$(\widehat{X}_0 - X_0)(\widehat{X}_N - X_N) \geq 0. \quad (8)$$

*Proof.* Without loss of generality, let us assume  $\widehat{X}_N > X_N$ . Assume by contradiction that  $X_0 > \widehat{X}_0$ . With our parity assumption, for any  $n \leq N$ ,  $\widehat{X}_n - X_n \in 2\mathbb{Z}$ , and that difference only takes steps in  $\{-2, 0, 2\}$ . Let us look at the largest  $n < N$  such that  $\widehat{X}_n = X_n$ . Let  $y = X_n$ . Under our assumptions, necessarily  $\widehat{X}_{n+1} = y + 1$  and  $X_{n+1} = y - 1$ . In particular,  $A_{y, n}^\varepsilon = -1$ , and therefore  $\widehat{X}_{n-1} = y + 1 \geq X_{n-1}$ . Now, if  $\widehat{X}_{n-1} > X_{n-1}$ , we reproduce the previous argument to show  $\widehat{X}_{n-2} \geq X_{n-2}$  (unless  $n-1$  is zero, in which case we stop). If  $\widehat{X}_{n-1} = X_{n-1} = y + 1$ ,  $A_{y+1, n-1}^\varepsilon = -1$ , and we also end up with  $\widehat{X}_{n-2} = y + 2 \geq X_{n-2}$ . We can reproduce this argument until we reach time 0 to get a contradiction. □

With this non-crossing property, it becomes easy to show the antisymmetry property.

*Proof of Theorem 1 (discrete setting).* Without loss of generality, assume by contradiction that  $\delta := v(\varepsilon) + v(-\varepsilon) > 0$ . Fix  $N$  large enough that

$$\mathbb{P} \left( \left| \frac{X_N^{\pm \varepsilon}}{N} - v(\pm \varepsilon) \right| \geq \delta/4 \right) \leq 1/3. \quad (9)$$

Define  $\tilde{x} = \lfloor (v(\varepsilon) - \delta/2)N \rfloor$ . If  $\tilde{x}$  has the same parity as  $N$ , let  $x = \tilde{x}$ . Let  $x = \tilde{x} + 1$  else. Define  $X, \widehat{X}$  as above. Consider the event

$$E = \left\{ \frac{X_N}{N} \geq v(\varepsilon) - \delta/4 \text{ and } \frac{\widehat{X}_0}{N} - x \geq v(-\varepsilon) - \delta/4 \right\}. \quad (10)$$

By Lemma 1,  $\mathbb{P}(E) \geq 1/3$ , but by Lemma 2,  $E$  is empty, since  $X_N > \widehat{X}_N = x$  and  $X_0 < \widehat{X}_0$  on  $E$ . We have our contradiction. □

**Remark 1.** *While the non-crossing property is elementary to check in the discrete time setting, it is not easy to find a coupling with the same property in the continuous time setting. One could think of associating independent Poisson clocks to every vertex of  $\mathbb{Z}$ , a direction  $A_{x, n}^\varepsilon$  with every clock ring and build  $X, \widehat{X}$  in a similar fashion. However, the jumps starting from neighboring sites would not be simultaneous, and it is easy to find examples where crossings occur. Fortunately, once we have the result in the discrete time setting, we do not need to work hard to push it to the continuous time setting.*

### 4 Proof of Theorem 1 in the continuous setting

Recall the construction of Section 2.3. Define for  $n \in \mathbb{N}$ ,  $\sigma_n = \eta_{T_n}$ . Theorem 1 will be proved when we have checked the three points of the following lemma.

**Lemma 3.** 1. *The process  $\sigma$  is translation invariant and reversible.*

2. If  $X^\varepsilon$  is the  $\varepsilon$ -RW in continuous time defined on  $\eta$  as in Section 2.3,  $(X_{T_n}^\varepsilon)_{n \in \mathbb{N}}$  is equal in distribution to  $\tilde{X}^\varepsilon$  the  $\varepsilon$ -RW in discrete time defined on  $\sigma$ .
3.  $X^\varepsilon$  satisfies the weak LLN (2) iff  $\tilde{X}^\varepsilon$  does, with the same limit speed  $v(\varepsilon)$ .

*Proof of Lemma 3.* 1. This follows from the translation invariance and reversibility of  $\eta$  and the PPP  $T$ , along with the independence of  $T$  and  $\eta$ .

2. This follows from the independence of  $U, T, \eta$ , and a construction in the discrete case analog to the one we use in Section 2.3.

3. This is an immediate consequence of the fact that  $T_n/n \xrightarrow[n \rightarrow \infty]{} 1$  almost surely. □

*Proof of Theorem 1 (continuous setting).* We assume that both  $X^\varepsilon$  and  $X^{-\varepsilon}$  satisfy the weak LLN (2). By the third point of Lemma 3,  $\tilde{X}^\varepsilon$  and  $\tilde{X}^{-\varepsilon}$  satisfy (2) with respective speeds  $v(\varepsilon), v(-\varepsilon)$ . Therefore, the discrete time version of Theorem 1 applies to  $\tilde{X}^\varepsilon, \tilde{X}^{-\varepsilon}$  and  $v(-\varepsilon) = -v(\varepsilon)$ . □

## References

- [ABF16] Luca Avena, Oriane Blondel, and Alessandra Faggionato. A class of random walks in reversible dynamic environments: Antisymmetry and applications to the East model. *Journal of Statistical Physics*, 165(1):1–23, Oct 2016.
- [ABF18] L. Avena, O. Blondel, and A. Faggionato. Analysis of random walks in dynamic random environments via  $l_2$ -perturbations. *Stochastic Processes and their Applications*, 128(10):3490 – 3530, 2018.
- [AdHR11] L. Avena, F. den Hollander, and F. Redig. Law of large numbers for a class of random walks in dynamic random environments. *Electron. J. Probab.*, 16:no. 21, 587–617, 2011.
- [BHT18] Oriane Blondel, Marcelo R. Hilario, and Augusto Teixeira. Random Walks on Dynamical Random Environments with Non-Uniform Mixing. *arXiv e-prints*, page arXiv:1805.09750, May 2018.
- [FMRT13] A. Faggionato, F. Martinelli, C. Roberto, and C. Toninelli. The East model: recent results and new progresses. *Markov Process. Relat. Fields*, 19(3):407–452, 2013.
- [HKT20] Marcelo R. Hilário, Daniel Kiouss, and Augusto Teixeira. Random walk on the simple symmetric exclusion process. *Commun. Math. Phys.*, 379(1):61–101, 2020.
- [HS15] François Huveneers and François Simenhaus. Random walk driven by the simple exclusion process. *Electron. J. Probab.*, 20:no. 105, 42, 2015.
- [JE91] J. Jäckle and S. Eisinger. A hierarchically constrained kinetic ising model. *J.Phys.B*, 84:115–124, 1991.
- [MV15] Thomas Mountford and Maria E. Vares. Random walks generated by equilibrium contact processes. *Electron. J. Probab.*, 20:no. 3, 17, 2015.
- [RV13] Frank Redig and Florian Völlering. Random walks in dynamic random environments: a transference principle. *Ann. Probab.*, 41(5):3157–3180, 2013.
- [Sol75] Fred Solomon. Random walks in a random environment. *Ann. Probability*, 3:1–31, 1975.
- [Z04] Francis and Ofer Zeitouni. A law of large numbers for random walks in random mixing environments. *Ann. Probab.*, 32(1B):880–914, 2004.