



**HAL**  
open science

# Dog behaviours in veterinary consultations: Part I. Effect of the owner's presence or absence

C Girault, Nathalie Priymenko, M Helsly, C Duranton, F Gaunet

## ► To cite this version:

C Girault, Nathalie Priymenko, M Helsly, C Duranton, F Gaunet. Dog behaviours in veterinary consultations: Part I. Effect of the owner's presence or absence. the Veterinary journal, 2022. hal-03829678

**HAL Id: hal-03829678**

**<https://hal.science/hal-03829678v1>**

Submitted on 25 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Original Article**

2  
3

4 **Dog behaviours in veterinary consultations: Part I. Effect of the owner's presence or**  
5 **absence.**

6  
7

8 C. Girault<sup>a\*</sup>, N. Priymenko<sup>b</sup>, M. Helsly<sup>a</sup>, C. Duranton<sup>c,d</sup>, F. Gaunet<sup>d</sup>

9

10 <sup>a</sup> *Ecole Nationale Vétérinaire de Toulouse, ENVT, 23 chemin des Capelles, BP 87614, 31076*  
11 *Toulouse Cedex, France*

12 <sup>b</sup> *TOXALIM, Université de Toulouse, INRA, ENVT, 23 chemin des Capelles, BP 87614, 31076*  
13 *Toulouse cedex, France*

14 <sup>c</sup> *AVA Association, Cuy-Saint-Fiacre, France*

15 <sup>d</sup> *Laboratoire de Psychologie Cognitive, AMU, CNRS, Fédération 3C Aix-Marseille Université,*  
16 *UMR 7290, Marseille, France*

17

18

19

20

21 \* Corresponding author. Tel.: +33 6 95 10 02 76

22 *E-mail address:* ciska.girault@gmail.com (C. Girault).

23 **Abstract**

24       Veterinary practices can be stressful places for dogs. Decreasing stress during veterinary  
25 consultations is therefore a major concern, since animal welfare matters both for owners and  
26 veterinarians. Stress can be expressed through behaviour modifications; monitoring dogs'  
27 behaviour is thus one way to assess stress levels. We also know that the owner can affect dog  
28 behaviour in different ways. The aim of this study was therefore to assess the effect of the  
29 presence of owners on the behaviour of their dogs in veterinary consultations. We studied 25  
30 dogs-owner dyads at two standardised veterinary consultations, conducted at intervals of five  
31 to seven weeks; the owner was present for the first consultation and absent for the second  
32 (O/NoO group, n= 12), or vice versa (NoO/O group, n= 13). A consultation consisted in three  
33 phases: exploration, examination, greeting. Dog behaviours were compared between the two  
34 conditions using a video recording.

35  
36       Despite some limitations (e.g. no male owners, the exclusion of aggressive dogs, a  
37 limited sample size, minimally invasive veterinary examinations, restricted owner-dog  
38 interactions), our results showed that the presence or absence of the owner had no significant  
39 effect on the stress-related behaviour of the dog or the veterinarian's ability to handle the  
40 animal during the examination phase ( $P > 0.05$ ). Nevertheless, the behaviour of the dogs  
41 towards people was affected before, during, and after the veterinary examination: in the  
42 presence of their owner, dogs were more willing to enter the consultation room ( $P > 0.05$ );  
43 they appeared more relaxed during the exploration phase ( $P > 0.05$ ); and they seemed less  
44 eager to leave the room after the examination ( $P > 0.1$ ). During the examination, dogs looked  
45 in direction of their owner in both condition (respectively owner present and behind the door;  
46  $P > 0.05$ ). These results suggest that allowing the owner to stay in the room during veterinary  
47 consultations is a better option for dogs' welfare.

48

49 *Keywords:* Dog Behaviour; Owner Presence; Owner Absence; Stress; Veterinary

50 Consultation.

## 51 **Introduction**

52           Improving animal welfare during veterinary consultations is a key concern of  
53 veterinarians, researchers and owners alike. Dogs frequently experience stress in these  
54 situations (Lind, 2017; Edwards, 2019) which can be assessed by monitoring behaviour  
55 (Beerda et al., 1997; Maximino et al., 2010; Koolhaas et al., 2011), for example, when  
56 entering the veterinary practice (Stanford, 1981; Mariti et al., 2017), during time spent in the  
57 waiting room (Mariti et al., 2015; Csoltova et al., 2017; Mariti et al., 2017), and during the  
58 examination itself (Döring et al., 2009; Mariti et al., 2017). In a study by Döring et al. (2009),  
59 80% of dogs showed stress-related behaviours on the examination table: 56.3% panted, 61.5%  
60 shivered, and 71.9% displayed avoidance behaviour. Glardon et al. (2010) found that around  
61 25% of dogs could not be handled during the examination. Studies also report physiological  
62 signs of stress, such as increased plasma cortisol levels, pulse rates, and blood pressure (Kallet  
63 et al., 1997; Vonderen et al., 1998).

64

65           Chronic stress can cause impaired welfare which has negative effects on health, leading  
66 to reduced immune system performance (Herbert and Cohen, 1993; Stowell et al., 2001;  
67 Gimsa et al., 2018), increased tumours (Riley, 1975; Dai et al., 2020), and premature aging  
68 (Epel et al., 2004). Acute stress, as expected in veterinary consultations, can lead to  
69 dysregulation of the stress system if it is extremely intense or recurrent (Chrapusta et al.,  
70 1997; De Kloet et al., 2005; Vaessen et al., 2015). Stress also modifies the behaviour of dogs  
71 and may increase aggression. When faced with a threatening situation, dogs tend to react in  
72 three different ways: freeze; fight; and/or flight (Bracha, 2004). Aggression is dangerous for  
73 veterinarians and owners: in Australia in 2006, 48% of veterinarians declared that they had  
74 been bitten by a dog at work between one and five times in the previous 12 months (Fritschi  
75 et al., 2006).

76

77           The ways in which dogs tend to react depend on their temperament (Goodloe and  
78 Borchelt, 1998; Serpell and Hsu, 2001; Svartberg, 2002; Bray et al., 2017); coping style  
79 (Koolhaas et al., 1999; Horváth et al., 2007; Diverio et al., 2017); genetics (Wilsson and  
80 Sundgren, 1997; Saetre et al., 2006; Meyer et al., 2012; Arvelius et al., 2014; Persson et al.,  
81 2015); and previous experiences (Seligman et al., 1979; Döring et al., 2009; Douglas et al.,  
82 2012). According to Döring et al. (2009), even one past aversive experience increases stress-  
83 related behaviour in dogs, thus modifying their behaviour at future visits to the veterinarian.

84

85           Many factors can be stressful for dogs in a veterinary practice (Edwards, 2019):  
86 transportation between home and the practice (Beerda et al., 1997); the novelty of the place  
87 (Beerda et al., 1997); the “white coat effect” (Kallet et al., 1997; Belew et al., 1999); the  
88 presence of new people and animals (Scotney, 2010); and unusual sounds and bustle (Beerda  
89 et al., 1997; Wells et al., 2002). Even smells such as those released by stressed people and  
90 animals can be stressful for dogs (Graham et al., 2005; Siniscalchi et al., 2011; Siniscalchi et  
91 al., 2016). In addition, dogs can be fearful when entering a veterinary practice due to previous  
92 experiences (Döring et al., 2009; Ziv, 2017). Veterinarians may also use gestures or postures  
93 that are stressful for dogs (Mariti et al., 2017; Edwards, 2019), such as bending over them  
94 (Vas et al., 2005; Györi et al., 2010; McGreevy et al., 2012); touching them (Payne et al.,  
95 2015 ); placing them on the examination table (Döring et al., 2009); restraining them by force  
96 (Beerda et al., 1997); holding their collar or closing their mouth (Kuhne et al., 2014); looking  
97 at them directly in the eyes (Györi et al., 2010); or bringing their face close to the dog’s head  
98 (Rezac et al., 2015). As a result, veterinary practices may be fearful places for dogs.

99

100 Other stressful factors originate from the owner (Lind, 2017). Studies focused on dog-  
101 owner attachment have shown that dogs can behave differently depending on whether their  
102 owner is present or absent (Topál et al., 1998). In particular, when dogs are left in a novel  
103 place without a familiar caregiver, they show higher activity (Tuber et al., 1996); higher  
104 glucocorticoid levels (Tuber et al., 1996; Palestrini et al., 2005); higher heart rate levels  
105 (Palestrini et al., 2005); and higher anxiety (Prato-Previde et al., 2003; Palestrini et al., 2005;  
106 Parthasarathy and Crowell-Davis, 2006), even if an unknown person is present (Parthasarathy  
107 and Crowell-Davis, 2006). Miklosi et al. (2003) showed that dogs looked at their owner when  
108 facing an unsolvable task, and Kerepesi et al. (2015) demonstrated that dogs moved closer to  
109 their owner but not towards other individuals, even familiar ones, in situations provoking  
110 anxiety or fear. Other studies have shown that dogs react in the same way as their owner  
111 when confronted with a strange object (Merola et al., 2012) or an unknown person (Duranton  
112 et al., 2016).

113

114 Owners can thus potentially modify the behaviour of their dog in a veterinary  
115 consultation. This hypothesis has been observed anecdotally by veterinarians in the field.  
116 Some believe that the very presence of owners can calm down their dog, whereas others  
117 maintain that dogs are easier to handle in the absence of their owner. Only a study by Stellato  
118 et al. (2020) focused on this question, and showed positive effects of owner presence on  
119 behavioural and physiological measures of fear in dogs during veterinary consultation.

120

121 In Part I of our study, we examined the effect of owner presence or absence on the  
122 behaviours of dogs during a veterinary consultation, regardless of the owners' actions. Based  
123 on the abovementioned literature, we expected the owner's presence to decrease stress-related  
124 behaviours of dogs during the veterinary examination and hence facilitate their handling by

125 the veterinarian. In Part II of the study (see Helsly et al., under review), using the raw data  
126 from consultations with the owner present, we explored whether owners' actions affect dog  
127 behaviours by observing dog-owner dyads.

128

## 129 **Materials and methods**

### 130 *Participants*

131 A total of 32 dog-owner dyads were recruited. Four dogs were excluded due to the  
132 owner's withdrawal between the two appointments, another one due to aggressive behaviour  
133 and two due to protocol deviation. The final study included 25 owner-dog dyads. All  
134 participants were volunteers and were recruited via social media. The owners (all women) and  
135 dogs meeting the following criteria were selected: adult dogs between 12 months and 10 years  
136 old and unfamiliar with the researchers, measuring less than 70 cm at the withers in order to  
137 be easily lifted and examined on the table, in good general health, and nonaggressive towards  
138 humans to avoid the use of a muzzle that could modify their behaviour; owners not working  
139 as a veterinarian, assistant, or veterinary student.

140 Participant demographics are shown in Table 1.

141

### 142 *Experimental procedure*

143 The study protocol was approved by the ethical committee SSA (Science et Santé  
144 Animale) n°115 (SSA\_2018\_008) on 18 July 2018. The experiment took place in an  
145 examination room at the National Veterinary School of Toulouse (ENVT), France (Fig. 1).  
146 All dogs underwent two videotaped veterinary consultations at an interval of five to seven  
147 weeks (Table 1), one in the presence of their owner and the other in their absence. Prior to  
148 each consultation, all owners were told how to behave with their dogs. The veterinary  
149 consultations were all carried out by the same two researchers: two female veterinary



150 students, one in the role of the veterinarian (C. G.) and the other in the role of the assistant  
151 (M. H.). During the consultations, the researchers conducted the veterinary examination using  
152 as neutral a disposition as possible: the researchers stayed still and did not talk to the dogs, pet  
153 them, make eye contact with them, or punish them. The veterinary consultation was divided  
154 into three main phases: phase 1: exploration; phase 2: examination; and phase 3: greeting. We  
155 consider a “consultation” to include all events between the times when the dog entered and  
156 exited the examination room, whereas an “examination” includes only the phase where the  
157 dog was examined (see “Standardised protocol for the veterinary consultations” below). Dogs  
158 were randomly distributed into two groups using AB/BA crossover design: in the O/NoO  
159 group ( $n = 12$ ), the owner was present for the first veterinary consultation and absent for the  
160 second, and vice versa in the NoO/O group ( $n = 13$ ). Consultations were arranged by  
161 appointment according to the availability of owners.

162

163         Raw data collected for this study was also used in Part II (Helsly et al., under review).  
164 Part II only focuses on data collected during consultations with the owner present. Four  
165 additional dogs were included in Part II but not in Part I because the dyads did not attend the  
166 second consultation with the owner absent.

167

### 168 *Data collection and analysis*

169         Consultations were videotaped from when the dog entered the consultation room until  
170 the end of the greeting phase. We used two cameras (a Canon Legria HF S21 and a Panasonic  
171 HC-WX970 with a Panasonic vW-W4907H wide-viewing angle) situated in two corners of  
172 the room and facing the examination table. The recorded videos from the two cameras were  
173 synchronised and assembled into a single video (Fig. 1).

174

175 The variables we studied, described below, differed depending on the phase of the  
176 consultation. A summary of all studied variables and their availability for the three phases is  
177 described in Appendix B: Supplementary material. Dog behaviour was analysed using the  
178 Solomon Coder beta 17.03.22 program<sup>1</sup> and the behavioural repertoire adapted from the  
179 literature (Beerda et al., 1998; Mills et al. 2006; Deldalle and Gaunet, 2014; Csoltova et al.,  
180 2017; see Table 2). Stress-related behaviours are described in Table 2. As the phase durations  
181 were variable, the durations of behaviours were converted into a time percentage (behaviour  
182 duration/phase duration) for all behaviours. Behavioural indices were further computed as  
183 detailed in Table 3. The Emotional State of the dog is a subjective rate assessing stress, scored  
184 by using a three-point scale defined as follows: relaxed, aroused, anxious, see definitions in  
185 Table 4. The dogs' apparent comfort in entering the room and the greeting intensity of the  
186 dogs towards their owner and of owners towards dogs were evaluated using a five-point scale  
187 defined in Table 5. The level of physical restraint was scored using a three-point scale defined  
188 as follows: low, medium, high. All definitions are given in Table 6. The success and difficulty  
189 of the manipulations were evaluated using a scale defined in Table 7.

190

#### 191 *Standardised protocol for the veterinary consultations*

192 Owner: present condition

193 Exploration phase (phase 1): The owner entered the room with her dog on a leash and  
194 sat on a chair (Fig. 2). The leash was dropped, and the dog explored the room freely for 2  
195 minutes and 30 seconds. Neither the owner nor the researcher spontaneously interacted with  
196 the dog; the owner could nevertheless respond to the dog's requests (physical, verbal and  
197 visual interactions were allowed). The researchers asked questions similar to those asked in a  
198 standard veterinary consultation.

---

<sup>1</sup> See : Solomon Coder, András Péter, <https://solomoncoder.com> (accessed 10 September 2020).

199

200 Examination phase (phase 2): The dog was put on the examination table by the  
201 researchers; the owner stood one metre away from the table in a designated spot, facing the  
202 dog. The veterinarian began a standardised veterinary examination following this predefined  
203 sequence: examination of eyes; ears; teeth; mouth mucosa; palpation of lymph nodes;  
204 examination of scapular skin fold; abdominal palpation; heart and lung auscultation;  
205 measurement of rectal temperature; and paw palpation. The assistant held the dog using a  
206 standardised restraint (Fig. 3): one hand on the chest and another on the base of the tail, using  
207 the minimal necessary strength to keep the dog sitting or standing on the table. If a particular  
208 manipulation failed because the dog was not cooperative for five seconds (for example, the  
209 dog struggled, resisted, or tried to escape), the veterinarian did not repeat the manipulation  
210 and continued with the next one. During the examination phase, the owner was only allowed  
211 to talk to or look at her dog (verbal and visual interactions were allowed). If the dog showed  
212 any sign of aggression toward the researchers or behaviour endangering them (for example,  
213 growling, showing teeth, or trying to bite), the procedure was terminated, and the dog was  
214 excluded from the study.

215

216 Greeting phase (phase 3): This phase started when the dog was on the floor and the  
217 owner in the room and lasted exactly 20 second. The dog was indeed taken down from the  
218 table by the researchers, and the leash was given back to the owner. The researchers then  
219 stepped aside and filled in forms in order to allow the owner and the dog to interact freely  
220 (physical, verbal and visual interactions were allowed).

221

222 Owner: absent condition

223           The procedure was the same as with ‘Owner: present condition’, except that the owner  
224 waited outside the room during the exploration and examination phases and only entered the  
225 room for the greeting phase. During the exploration phase, the owner accompanied the dog to  
226 the open door, gave the leash to the assistant and was free to interact with the dog in order to  
227 encourage the dog to enter the room. The dog explored the room freely for 2 minutes and 30  
228 seconds while still on the leash. During this phase, the researchers did not interact with the  
229 dog and spoke in a neutral tone. The procedure of the examination phase was the same as  
230 described in ‘Owner: present condition’. During the greeting phase, the dog was taken down  
231 from the table by the researchers. Then, the assistant went outside to fetch the owner, and the  
232 leash was given back to the owner when she came back in the room without specific  
233 instructions. The end of this phase was the same as described in ‘Owner: present condition’  
234 after having given the leash back to the owner.

235

### 236 *Interobserver agreement*

237           Three assessors participated in the video analysis. The two researchers coded all the  
238 behaviours in all the videos: half of the behaviours were coded by one researcher and the  
239 other half by the second researcher. To assess the reproducibility of the behavioural analysis,  
240 a third assessor who was unaware of the study hypotheses and aims coded 30% of the  
241 behaviours in a random subset of 30% of the videos. Considering that a concordance, and not  
242 only a correlation, was needed to assess the interobserver reproducibility, Lin’s concordance  
243 correlation test was used (Lawrence et al., 1989; Barnhart et al., 2002; Barnhart et al., 2007).  
244 Interobserver agreement between the two assessors was determined by calculating  $\rho C$  values  
245 and rated according to Landis and Koch (1977) ( $\rho C = 0 - 0.2$ : slight agreement,  $\rho C = 0.21 -$   
246  $0.4$ : fair agreement,  $\rho C = 0.41 - 0.60$ : moderate agreement,  $\rho C = 0.61 - 0.8$ : substantial  
247 agreement,  $\rho C > 0.81$ : excellent agreement). Lin’s concordance correlation coefficients were

248 excellent ( $\rho C > 0.98$ ) for whining, tail between the legs, and moving and gazing at the owner,  
249 and substantial for contact with the assistant ( $\rho C = 0.71$ ) and sniffing ( $\rho C = 0.69$ ). Physical  
250 restraint was evaluated by the assistant, the success and difficulty of the manipulations were  
251 assessed by the veterinarian, and the other scores were rated by the three assessors.

252

### 253 *Statistical analysis*

254 The exploration, examination, and greeting phases were analysed separately.  
255 Comparisons were carried out for each behaviour or behavioural index with the owner absent  
256 or present conditions. A paired Student's *t* test was performed for the O/NoO and NoO/O  
257 groups using R software<sup>2</sup>.

258

## 259 **Results**

260 On average, the exploration phase lasted  $155.43 \pm 9.54$  seconds and the examination  
261 phase  $142.53 \pm 16.4$  seconds. The greeting phase, standardised in the study, lasted 20 seconds.  
262 The sample is described in Appendix A: Supplementary material.

263

### 264 *Exploration phase (phase 1)*

#### 265 Entering the room

266 Dogs appeared to enter the room more readily when the owner was present rather than  
267 absent, and the difference was significant ( $P < 0.05$ ; see Table 8, Video 1).

268

#### 269 Emotional state

270 Dogs had a significantly lower score for emotional state during the exploration phase  
271 when the owner was present rather than absent ( $P < 0.05$ ; see Table 8, Video 2).

---

<sup>2</sup> See: The R Project for Statistical Computing. <http://www.R-project.org>. (accessed 24 May 2017)

272

273 *Examination phase (phase 2)*

274 Stress-related behaviour

275 No differences were observed regarding the stress-related behaviours or Total Stress  
276 (defined in Table 3) during the examination phase in the presence or absence of the owners  
277 ( $P>0.05$ ).

278

279 Behaviour towards the veterinarian and assistant

280 The dogs' contact with the veterinarian and/or assistant lasted significantly longer  
281 during the examination phase if the owner was present rather than absent ( $P< 0.05$ ; Table 9).

282

283 Behaviour (gaze) towards the owner and/or door

284 During the examination phase, dogs looked straight ahead toward the assigned place  
285 of the owner significantly more if the owner was present rather than absent. ( $P< 0.05$ ; Table  
286 9, Video 3). Furthermore, dogs looked at the door significantly less when their owner was  
287 present rather than absent ( $P< 0.05$ ; Video 3).

288

289 Dogs' handling

290 No difference was observed regarding the restraint and the success and difficulty  
291 scores of manipulations in the presence or absence of the owners ( $P> 0.05$ ).

292

293 *Greeting phase (phase 3)*

294 Intensity of greeting

295 When the dog and owner were reunited after the examination phase, dogs greeted their  
296 owners for a significantly shorter period of time ( $P< 0.05$ ) and showed a lower reunion score

297 ( $P < 0.05$ ) if the owner was present rather than absent during the examination phase (Video 4).  
298 Nevertheless, the scores of owner behaviour towards their dog were similar regardless of  
299 whether the owner was present or not in the previous phase ( $P > 0.05$ ).

300

301 Door-directed gaze

302 Finally, dogs gazed at the door slightly less often when the owner was present rather  
303 than absent ( $P < 0.1$ ; Table 10, Video 5).

304

## 305 **Discussion**

306 This experiment aimed at investigating whether the presence or absence of a dog's  
307 owner influences dogs' behaviours in veterinary consultations. Overall results suggest that  
308 allowing the owner to stay in the room during veterinary consultations is a better option for  
309 dogs' welfare: During the exploration phase, in the presence of their owner, dogs were more  
310 willing to enter the consultation room and appeared more relaxed. During the examination  
311 phase, dogs looked in direction of their owner in both condition: when their owner was  
312 present (standing in front of the dog), dogs looked straight ahead more often and at the door  
313 less often than in the absence of their owner. When the owner was absent (she had left the  
314 room through the door), the dogs looked at the door more often and less straight ahead than in  
315 the presence of the owner. Physical contacts engaged by dogs with the researchers lasted  
316 longer when their owners were present rather than absent. Our results also indicate that the  
317 presence or absence of the owner had no significant effect on the stress-related behaviour of  
318 the dog or the veterinarian's ability to handle the animal during the examination phase.  
319 Finally, during the greeting phase, our results showed that the dogs exhibited less greeting  
320 behaviour and tended to look at the door less often if the owner had never left the room than if  
321 the owner had been absent.

322

323           Only a study by Stellato et al. (2020) focused on the effect of the presence or absence  
324 of the owner on dog behaviour in veterinary practices, comparing two standardized veterinary  
325 consultation with owner present or absent. They showed that dogs had a lower rate of  
326 vocalisation, higher rate of yawning and lower mean axillary temperature in the presence of  
327 the owner. Thus, they encouraged owner to remain with their dog during routine veterinary  
328 examination. Note that we did not find such a difference for vocalisation and yawning, but we  
329 did find significant differences for other parameters. A study by Csoltova et al. (2017)  
330 focused on the active or passive support of the owner during veterinary examinations. The  
331 authors compared behavioural and physiological measures of dogs during a veterinary  
332 examination in two conditions: the active presence of the owner (talking and petting), and the  
333 passive presence of the owner (sitting quietly next to the examination table). They found heart  
334 rate and internal temperature variations showing a beneficial effect of dog-owner interaction  
335 on the dogs' well-being but no significant behavioural changes. As mentioned in the  
336 introduction, owners can have various effects on the behaviour of their dog. Studies have  
337 shown that dogs adjust their behaviour to their owner's overall emotional body posture (Vas  
338 et al., 2005; Custance and Meyer 2012), to the owner's behaviours (Millot, 1994; Merola et  
339 al., 2012; Horn et al., 2012; Durantou and Gaunet, 2015); and to the owner's facial  
340 expressions (Deputte and Doll, 2011). Other studies detailed in Part II have shown that  
341 physical contact does not have the same effect on dog behaviour as talking (Helsly et al.,  
342 under review). The effect of the presence of the owner is thus hard to accurately predict, as it  
343 can depend on the owner's behaviour and mood.

344

345           In the present study, several elements can explain the absence of any significant  
346 differences in stress-related behaviours during the examination phase. First, physical contact



347 between the owners and dogs was not allowed during the examination phase to control  
348 parameters not being studied. Yet the Csoltova study showed a beneficial effect of contact  
349 during veterinary examinations, and other studies detailed in Part II have shown that talking  
350 and physical contact do not have the same effect (Helsly et al., under review). Another  
351 element in our study involved the subjective evaluation of stress (emotional state). This  
352 evaluation showed higher levels of stress during the exploration phase (phase 1) when the  
353 owner was absent than when the owner was present. In contrast, during the examination phase  
354 (phase 2), no difference was found in the subjective evaluation by the judges (emotional state)  
355 or the video analyses (stress-related behaviours). A study by Firnkes et al. (2017)  
356 demonstrated that some stress-related behaviours (“licking of lips” and “looking away”)  
357 decrease even though the stressor intensity increases. We thus postulate that the dogs reached  
358 a high threshold of stress in the ‘Owner: absent condition’ that prevented them from  
359 displaying additional stress-related behaviours. Potentially, too many stressors may hide the  
360 display of stress-related behaviour at some point. Alternatively, the absence of any difference  
361 may show that dogs are not actually more stressed when the owner is absent during a  
362 veterinary examination. In the examination phase, dogs engaged in more physical contact with  
363 the researchers when their owner was present than when their owner was absent. This  
364 engagement with researchers suggests that in an interventionist situation, if we had imposed  
365 physical contact on the dogs to keep them on the table, the presence of the owner would act as  
366 a social reference for the dog towards the veterinarian and assistant. That is, the presence of  
367 the owner could help dogs to handle this difficulty. Even if no significant differences were  
368 shown in terms of stress-related behaviours during examination phase, other results suggest  
369 that the owner does play a role. For instance, dogs looked in the direction of their owner both  
370 in the absence and in the presence of the owner. This also highlights the importance of the  
371 owner’s presence to help dogs to deal with the situation (e.g. social referencing in Merola et

372 al., 2012; Duranton et al., 2016; Salamon et al., 2020; and also Part II (Helsly et al., under  
373 review, for additional information). Furthermore, greeting behaviours during the two reunion  
374 conditions differed. Once dogs were placed back on the floor and the leash was given back to  
375 the owner, dogs showed more greetings when the owner came back than when the owner had  
376 never left the room. This is in accordance with studies showing that greeting behaviour is  
377 more marked when dogs spend a stressful time without their owner (Konok et al., 2011; Rehn  
378 and Keeling, 2011). Dogs tended to look at the door more often during the greeting phase  
379 when the owner had come back than when the owner had never left. Dogs are known to look  
380 at a desirable object (Gaunet, 2008; Gaunet, 2010; Gaunet and Deputte, 2011), and thus they  
381 may have been more motivated to leave the room when their owner was absent, since the time  
382 they spent in the room was more stressful, in accordance with the emotional state evaluation  
383 during the exploration phase. Finally, the presence of the owner had no detrimental effect but  
384 some beneficial effects on dogs. Dogs were neither more stressed nor more difficult to handle  
385 during examination phase and appeared less stressed during the exploration phase. In  
386 conclusion, the presence of owners appears to be more beneficial than their absence during  
387 veterinary examinations.

388

389 Our sample size was small, the veterinary examination was minimally invasive; puppy,  
390 old dogs and dogs measuring more than 70cm were not included and aggressive dogs were  
391 excluded from our study for safety reasons. Glardon et al. (2010) estimated that 16% of dogs  
392 display aggressive behaviour during veterinary examinations. If these dogs had been included  
393 in the study, the results may have been different, since aggressive dogs are less tolerant of  
394 manipulations. Furthermore, the manipulations used in this study were standard but minimally  
395 invasive. Dogs' tolerance of manipulation would have differed if we had employed more  
396 invasive or painful procedures (Holton et al., 2001; Hansen, 2003), and the presence or

397 absence of the owner could have had a different effect in these cases. Owner and researcher  
398 behaviour could have differed from a real-life scenario, as owners were not allowed to touch  
399 their dogs during the examination phase, and researchers kept a neutral disposition and did not  
400 initiate interaction toward dogs or respond to their requests for attention. In addition, the dogs  
401 involved in the current study were healthy and did not need any invasive manipulation.  
402 Owners may be more stressed in a real veterinary examination and therefore have a different  
403 effect on dog behaviour than in the controlled experimental conditions. Furthermore, while it  
404 has been shown that dogs react differently to men and women, we were not able to study the  
405 gender effects of owners and researchers on the dogs' behaviours, as both the owners and  
406 researchers were all women (Hennessy et al., 1998; Wells and Hepper, 1999; Deputte and  
407 Doll, 2011). In the present study, we focused on canine behaviour, although physiological  
408 measures such as plasma or salivary cortisol, heart rate, and infrared thermography can also  
409 be used. These measures could provide additional information about the stress experienced by  
410 dogs during veterinary consultations and help highlight the differences that cannot be  
411 observed by behaviour analysis alone, such as behaviours with high interindividual variation  
412 (Firnkes et al., 2017). We thus encourage further studies to focus on neurochemical and  
413 physiological differences in conjunction with behavioural differences, to use a larger sample  
414 of dogs and to study gender effect.

415

## 416 **Conclusions**

417 In summary, despite some limitations (e.g. no male owners, no dog above 70cm, no old  
418 dogs nor puppies, no aggressive dogs, 25 dogs by condition, minimally invasive  
419 examinations, restricted owner-dog interactions), the current study shows that the presence of  
420 the owner in veterinary consultations (constituted by exploration, examination and greeting  
421 phases) may help to reduce the stress-related behaviours of dogs before veterinary

422 examinations. During examination phase, dogs looked at their owners and appear to seek  
423 social information from owner, whereas when their owners were absent, they looked for them.  
424 Behaviours like greeting their owners and door-related behaviours suggest that even if no  
425 significant differences were shown for stress-related behaviours in terms of the absence or  
426 presence of owners, dogs were **less** stressed during examination phase when their owner was  
427 present. Given these results, it seems more appropriate to allow owners to attend veterinary  
428 examinations with their dog, as only positive effects were observed in terms of the dogs'  
429 behaviour and well-being, in spite of the previously mentioned limitations.

430

#### 431 **Conflict of interest statement**

432 None of the authors has any financial or personal relationships that could inappropriately  
433 influence or bias the content of the paper.

434

#### 435 **Acknowledgements**

436 We are grateful to Elodie Losserand who helped us during the study. We thank all the dog  
437 owners who voluntarily participated in this study. We are also grateful to Caniplex, Bagat-en-  
438 Quercy, France, for providing thank-you gifts to all the volunteers. Preliminary results were  
439 presented during a confidential veterinary thesis defence at the Ecole Nationale Vétérinaire de  
440 Toulouse on 13 December 2017.

441

#### 442 **Supplementary material**

443 Supplementary data associated with this article can be found, in the online version, at  
444 doi: ...

445

#### 446 **References**

447 Arvelius, P., Asp, H.E., Fikse, W.F., Strandberg, E., Nilsson, K. 2014. Genetic analysis of a  
448 temperament test as a tool to select against everyday life fearfulness in Rough Collie.  
449 *Journal of Animal Science* 92, 4843-4855.  
450

451 Barnhart, H.X., Haber, M., Song, J., 2002. Overall concordance correlation coefficient for  
452 evaluating agreement among multiple observers. *Biometrics* 58, 1020–1027.  
453

454 Barnhart, H.X., Haber, M.J., Lin, L.I., 2007. An overview on assessing agreement with  
455 continuous measurements. *Journal of Biopharmaceutical Statistics* 17, 529–69.  
456

457 Beerda, B., Schilder, M.B., Van Hooff, J.A., De Vries, H.W., 1997. Manifestations of chronic  
458 and acute stress in dogs. *Applied Animal Behaviour Science* 52, 307–319.  
459

460 Beerda, B., Schilder, M.B., Van Hooff, J.A., De Vries, H.W., Mol, J.A., 1998. Behavioural,  
461 saliva cortisol and heart rate responses to different types of stimuli in dogs. *Applied*  
462 *Animal Behaviour Science* 58, 365–381.  
463

464 Belew, A.M., Barlett, T., Brown, S.A., 1999. Evaluation of the white-coat effect in cats.  
465 *Journal of Veterinary Internal Medicine* 13, 134-142.  
466

467 Bracha, H.S., 2004. Freeze, flight, fight, fright, faint: Adaptationist perspectives on the acute  
468 stress response spectrum. *CNS Spectrums* 9, 679–685.  
469

470 Bray, E.E., Sammel, M.D., Cheney, D.L., Serpell, J.A., Seyfarth, R.M. 2017. Effects of  
471 maternal investment, temperament, and cognition on guide dog success. *Proceedings*  
472 *of the National Academy of Sciences* 114, 9128-9133.  
473

474 Chrapusta, S. J., Wyatt, R. J., Masserano, J. M., 1997. Effects of single and repeated  
475 footshock on dopamine release and metabolism in the brains of Fischer rats. *Journal of*  
476 *Neurochemistry* 68, 2024-2031.  
477

478 Csoltova, E., Martineau, M., Boissy, A., Gilbert, C., 2017. Behavioral and physiological  
479 reactions in dogs to a veterinary examination: Owner-dog interactions improve canine  
480 well-being. *Physiology and Behavior* 177, 270–281.  
481

482 Custance, D., Mayer, J., 2012. Empathic-like responding by domestic dogs (*Canis familiaris*)  
483 to distress in humans: an exploratory study. *Animal Cognition* 15, 851-859.  
484

485 Dai, S., Mo, Y., Wang, Y., Xiang, B., Liao, Q., Zhou, M., Zeng, Z., 2020. Chronic stress  
486 promotes cancer development. *Frontiers in Oncology* 10, 1492.  
487

488 De Kloet, E. R., Joëls, M., Holsboer, F., 2005. Stress and the brain: from adaptation to  
489 disease. *Nature Reviews Neuroscience* 6, 463-475.  
490

491 Deldalle, S., Gaunet, F., 2014. Effects of 2 training methods on stress-related behaviors of the  
492 dog (*Canis familiaris*) and on the dog–owner relationship. *Journal of Veterinary*  
493 *Behavior: Clinical Applications and Research* 9, 58–65.  
494

495 Deputte, B.L., Doll, A., 2011. Do dogs understand human facial expressions? *Journal of*  
496 *Veterinary Behavior: Clinical Applications and Research* 1, 78-79.

497

498 Diverio, S., Menchetti, L., Riggio, G., Azzari, C., Iaboni, M., Zasso, R., Santoro, M.M. 2017.  
499 Dogs' coping styles and dog-handler relationships influence avalanche search team  
500 performance. *Applied Animal Behaviour Science* 191, 67-77.

501

502 Döring, D., Roscher, A., Scheipl, F., Küchenhoff, H., Erhard, M.H., 2009. Fear-related  
503 behaviour of dogs in veterinary practice. *The Veterinary Journal* 182, 38–43.

504

505 Douglas, C., Bateson, M., Walsh, C., Bédué, A., Edwards, S.A., 2012. Environmental  
506 enrichment induces optimistic cognitive biases in pigs. *Applied Animal Behaviour*  
507 *Science* 139, 65-73.

508

509 Duranton, C., Gaunet, F., 2015. *Canis sensitivus*: Affiliation and dogs' sensitivity to others'  
510 behavior as the basis for synchronization with humans? *Journal of Veterinary*  
511 *Behavior: Clinical Applications and Research* 10, 513–524.

512

513 Duranton, C., Bedossa, T., Gaunet, F., 2016. When facing an unfamiliar person, pet dogs  
514 present social referencing based on their owners' direction of movement alone.  
515 *Animal Behaviour* 113, 147-156.

516

517 Edwards, P.T., Smith, B.P., McArthur, M.L., Hazel, S.J., 2019. Fearful fido: Investigating  
518 dog experience in the veterinary context in an effort to reduce distress. *Applied*  
519 *Animal Behaviour Science* 213, 14-25.

520

521 Epel, E.S., Blackburn, E., Lin, J., Dhabhar, F.S., Adler, N.E., Morrow, J.D., Cawthon, R.M.,  
522 2004. Accelerated telomere shortening in response to life stress. *Proceedings of the*  
523 *National Academy of Sciences of the United States of America* 101, 17312–17315.

524

525 Firnkes, A., Bartels, A., Bidoli, E., Erhard, M., 2017. Appeasement signals used by dogs  
526 during dog–human communication. *Journal of Veterinary Behavior: Clinical*  
527 *Applications and Research* 19, 35–44.

528

529 Fritschi, L., Day, L., Shirangi, A., Robertson, I., Lucas, M., Vizard, A., 2006. Injury in  
530 Australian veterinarians. *Occupational Medicine* 56, 199-203.

531

532 Gaunet, F., 2008. How do guide dogs of blind owners and pet dogs of sighted owners (*Canis*  
533 *familiaris*) ask their owners for food? *Animal Cognition* 11, 475-483.

534

535 Gaunet, F., 2010. How do guide dogs and pet dogs (*Canis familiaris*) ask their owners for  
536 their toy and for playing? *Animal Cognition* 13, 311-323.

537

538 Gaunet, F., Deputte, B. L., 2011. Functionally referential and intentional communication in  
539 the domestic dog: effects of spatial and social contexts. *Animal Cognition* 14, 849-  
540 860.

541

542 Gimsa, U., Tuchscherer, M., Kanitz, E., 2018. Psychosocial stress and immunity - what can  
543 we learn from pig studies? *Frontiers in Behavioral Neuroscience* 12, 64.

544  
545 Glardon, J., Hartnack, S., Horisberger, L., 2010. Analyse du comportement des chiens et des  
546 chats pendant l'examen physique en cabinet vétérinaire. *Schweizer Archiv für*  
547 *Tierheilkunde* 152, 69–75.  
548  
549 Godbout, M., Palestrini, C., Beauchamp, G., Frank, D., 2007. Puppy behavior at the  
550 veterinary clinic: A pilot study. *Journal of Veterinary Behavior: Clinical Applications*  
551 *and Research* 2, 126–135.  
552  
553 Goodloe, L.P., Borchelt, P.L., 1998. Companion dog temperament traits. *Journal of Applied*  
554 *Animal Welfare Science* 1, 303-338.  
555  
556 Graham, L., Wells, D L., Hepper, P.G., 2005. The influence of olfactory stimulation on the  
557 behaviour of dogs housed in a rescue shelter. *Applied Animal Behaviour Science* 91,  
558 143-153.  
559  
560 Győri, B., Gácsi, M., Miklósi, A., 2010. Friend or foe: Context dependent sensitivity to  
561 human behaviour in dogs. *Applied Animal Behaviour Science* 128, 69–77.  
562  
563 Hansen, B.D., 2003. Assessment of pain in dogs: veterinary clinical studies. *Institute for*  
564 *Laboratory Animal Research Journal* 44, 197–205.  
565  
566 Helsly, M., Priymenko, N., Girault, C., Duranton, C., Gaunet, F. Dog behaviours in veterinary  
567 consultations: Part II. The relationship between the behaviours of dogs and owners.  
568 Under review.  
569  
570 Hennessy, M.B., Williams, M.T., Miller, D.D., Douglas, C.W., Voith, V.L., 1998. Influence  
571 of male and female petters on plasma cortisol and behaviour: can human interaction  
572 reduce the stress of dogs in a public animal shelter? *Applied Animal Behaviour*  
573 *Science* 61, 63-77.  
574  
575 Herbert, T.B., Cohen, S., 1993. Stress and immunity in humans: a meta-analytic review.  
576 *Psychosomatic Medicine* 55, 364–379.  
577  
578 Holton, L., Pawson, P., Nolan, A., Reid, J., Scott, E.M., 2001. Development of a behaviour-  
579 based scale to measure acute pain in dogs. *The Veterinary Record* 148, 525–531.  
580  
581 Horn, L., Virányi, Z., Miklósi, Á., Huber, L., Range, F., 2012. Domestic dogs (*Canis*  
582 *familiaris*) flexibly adjust their human-directed behavior to the actions of their human  
583 partners in a problem situation. *Animal Cognition* 15, 57-71.  
584  
585 Horváth, Z., Igyártó, B.Z., Magyar, A., Miklósi, Á. 2007. Three different coping styles in  
586 police dogs exposed to a short-term challenge. *Hormones and Behavior* 52, 621-630.  
587  
588 Kallet, A.J., Cowgill, L.D. and Kass, P.H., 1997. Comparison of blood pressure  
589 measurements obtained in dogs by use of indirect oscillometry in a veterinary clinic  
590 versus at home. *Journal of the American Veterinary Medical Association* 210, 651–  
591 654.  
592

593 Kaminski, J., Neumann, M., Bräuer, J., Call, J., Tomasello, M., 2011. Dogs, *Canis familiaris*,  
594 communicate with humans to request but not to inform. *Animal Behaviour* 82, 651–  
595 658.

596

597 Kerepesi, A., Dóka, A., Miklósi, Á., 2015. Dogs and their human companions: the effect of  
598 familiarity on dog–human interactions. *Behavioural Processes* 110, 27-36.  
599

600 Konok, V., Dóka, A., Miklósi, A., 2011. The behavior of the domestic dog (*Canis familiaris*)  
601 during separation from and reunion with the owner: A questionnaire and an  
602 experimental study. *Applied Animal Behaviour Science* 135, 300–308.  
603

604 Koolhaas, J.M., Korte, S.M., De Boer, S.F., Van Der Vegt, B.J., Van Reenen, C.G., Hopster,  
605 H., De Jong I.C., Ruis M.A.W., Blokhuis, H.J., 1999. Coping styles in animals:  
606 current status in behavior and stress-physiology. *Neuroscience and Biobehavioral*  
607 *Reviews* 23, 925-935.  
608

609 Koolhaas, J. M., Bartolomucci, A., Buwalda, B., de Boer, S. F., Flügge, G., Korte, S. M.,  
610 Meerlo, P., Murisong, R., Olivier, B., Palanzak, P., Richter-Levine, G., Sgoifok, A.,  
611 Steimerj, T., Stiedl, O., van Dijkh, G., Wöhrd, M., Fuchs, E., 2011. Stress revisited: a  
612 critical evaluation of the stress concept. *Neuroscience and Biobehavioral Reviews* 35,  
613 1291-1301.  
614

615 Kuhne, F., Hößler, J. C., Struwe, R., 2014. Emotions in dogs being petted by a familiar or  
616 unfamiliar person: Validating behavioural indicators of emotional states using heart  
617 rate variability. *Applied Animal Behaviour Science* 161, 113-120.  
618

619 Landis, J.R., Koch, G.G., 1977. The measurement of observer agreement for categorical data.  
620 *Biometrics*, 159-174.  
621

622 Lawrence, I., Lin, L.I., 1989. A concordance correlation coefficient to evaluate  
623 reproducibility. *Biometrics* 45, 255–268.  
624

625 Lind, A.K., Hydbring-Sandberg, E., Forkman, B., Keeling, L.J., 2017. Assessing stress in  
626 dogs during a visit to the veterinary clinic: Correlations between dog behavior in  
627 standardized tests and assessments by veterinary staff and owners. *Journal of*  
628 *Veterinary Behavior* 17, 24-31.  
629

630 Mariti, C., Raspanti, E., Zilocchi, M., Carlone, B., Gazzano, A., 2015. The assessment of dog  
631 welfare in the waiting room of a veterinary clinic. *Animal Welfare* 24, 299-305.  
632

633 Mariti, C., Pierantoni, L., Sighieri, C., Gazzano, A., 2017. Guardians' perceptions of dogs'  
634 welfare and behaviors related to visiting the veterinary clinic. *Journal of Applied*  
635 *Animal Welfare Science* 20, 24–33.  
636

637 Maximino, C., de Brito, T. M., da Silva Batista, A. W., Herculano, A. M., Morato, S.,  
638 Gouveia Jr, A., 2010. Measuring anxiety in zebrafish: a critical review. *Behavioural*  
639 *Brain Research* 214, 157-171.  
640



- 641 McGreevy, P.D., Starling, M., Branson, N.J., Cobb, M.L., Calnon, D., 2012. An overview of  
642 the dog–human dyad and ethograms within it. *Journal of Veterinary Behavior: Clinical*  
643 *Applications and Research* 7, 103–117.  
644
- 645 Merola, I., Prato-Previde, E., Marshall-Pescini, S., 2012. Social referencing in dog-owner  
646 dyads? *Animal Cognition* 15, 175–185.  
647
- 648 Meyer, F., Schawalder, P., Gaillard, C., Dolf, G., 2012. Estimation of genetic parameters for  
649 behavior based on results of German Shepherd Dogs in Switzerland. *Applied Animal*  
650 *Behaviour Science* 140, 53-61.  
651
- 652 Miklósi, Á., Kubinyi, E., Topál, J., Gácsi, M., Virányi, Z., Csányi, V., 2003. A simple reason  
653 for a big difference: wolves do not look back at humans, but dogs do. *Current*  
654 *Biology* 13, 763–766.  
655
- 656 Millot, J.L. 1994. Olfactory and visual cues in the interaction systems between dogs and  
657 children. *Behavioural Processes* 33, 177-188.  
658
- 659 Mills, D.S., Ramos, D., Estelles, M.G., Hargrave, C., 2006. A triple blind placebo-controlled  
660 investigation into the assessment of the effect of Dog Appeasing Pheromone (DAP) on  
661 anxiety related behaviour of problem dogs in the veterinary clinic. *Applied Animal*  
662 *Behaviour Science* 98, 114–126.  
663
- 664 Palestirini, C., Previde, E.P., Spiezio, C., Verga, M., 2005. Heart rate and behavioural  
665 responses of dogs in the Ainsworth's Strange Situation: A pilot study. *Applied Animal*  
666 *Behaviour Science* 94, 75-88.  
667
- 668 Parthasarathy, V., Crowell-Davis, S.L. 2006. Relationship between attachment to owners and  
669 separation anxiety in pet dogs (*Canis lupus familiaris*). *Journal of Veterinary Behavior*  
670 1, 109-120.  
671
- 672 Pastore, C., Pirrone, F., Balzarotti, F., Faustini, M., Pierantoni, L., Albertini, M., 2011.  
673 Evaluation of physiological and behavioral stress-dependent parameters in agility  
674 dogs. *Journal of Veterinary Behavior: Clinical Applications and Research* 6, 188–194.  
675
- 676 Payne, E., Boot, M., Starling, M., Henshall, C., McLean, A., Bennett, P., McGreevy, P., 2015.  
677 Evidence of horsemanship and dogmanship and their application in veterinary  
678 contexts. *The Veterinary Journal* 204, 247–254.  
679
- 680 Persson, M.E., Roth, L.S.V., Johnsson, M., Wright, D., Jensen, P., 2015. Human-directed  
681 social behaviour in dogs shows significant heritability. *Genes, Brain and Behavior* 14,  
682 337-344.  
683
- 684 Prato-Previde, E., Spiezio, C., Sabatini, F., Custance, D.M. 2003. Is the dog-human  
685 relationship an attachment bond? An observational study using Ainsworth's strange  
686 situation. *Behaviour* 140, 225-254.  
687
- 688 Rehn, T., Keeling, L.J., 2011. The effect of time left alone at home on dog welfare. *Applied*  
689 *Animal Behaviour Science* 129, 129–135.

690  
691 Rezac, P., Rezac, K., Slama, P., 2015. Human behavior preceding dog bites to the face. *The*  
692 *Veterinary Journal* 206, 284–288.  
693  
694 Riley, V., 1975. Mouse mammary tumors: alteration of incidence as apparent function of  
695 stress. *Science* 189, 465–467.  
696  
697 Saetre, P., Strandberg, E., Sundgren, P.E., Pettersson, U., Jazin, E., & Bergström, T.F., 2006.  
698 The genetic contribution to canine personality. *Genes, Brain and Behavior* 5, 240-248.  
699  
700 Scotney, R.L., 2010. Environmental enrichment in veterinary practice. *The Veterinary Nurse*  
701 1, 140-149.  
702  
703 Seligman, M.E., Maier, S.F., Geer, J., 1979. Alleviation of learned helplessness in the dog. In  
704 *Origins of Madness*, Pergamon, 401-409.  
705  
706 Serpell, J.A., Hsu, Y., 2001. Development and validation of a novel method for evaluating  
707 behavior and temperament in guide dogs. *Applied Animal Behaviour Science* 72, 347-  
708 364.  
709  
710 Siniscalchi, M., Sasso, R., Pepe, A.M., Dimatteo, S., Vallortigara, G., Quaranta, A., 2011.  
711 Sniffing with the right nostril: lateralization of response to odour stimuli by dogs.  
712 *Animal Behaviour* 82, 399–404.  
713  
714 Siniscalchi, M., d’Ingeo, S., Quaranta, A., 2016. The dog nose “KNOWS” fear: Asymmetric  
715 nostril use during sniffing at canine and human emotional stimuli. *Behavioural Brain*  
716 *Research* 304, 34-41.  
717  
718 Stanford, T.L., 1981. Behavior of dogs entering a veterinary clinic. *Applied Animal*  
719 *Ethology* 7, 271–279.  
720  
721 Stellato, A.C., Dewey, C.E., Widowski, T.M., Niel, L., 2020. Evaluation of associations  
722 between owner presence and indicators of fear in dogs during routine veterinary  
723 examinations. *Journal of the American Veterinary Medical Association* 257, 1031-  
724 1040.  
725  
726 Stowell, J.R., Kiecolt-Glaser, J.K., Glaser, R., 2001. Perceived stress and cellular immunity:  
727 When coping counts. *Journal of Behavioral Medicine* 24, 323-339.  
728  
729 Svartberg, K. 2002. Shyness–boldness predicts performance in working dogs. *Applied*  
730 *Animal Behaviour Science* 79, 157-174.  
731  
732 Topál, J., Miklósi, A., Csányi, V., Dóka, A., 1998. Attachment behavior in dogs (*Canis*  
733 *familiaris*): a new application of Ainsworth’s (1969) Strange Situation Test. *Journal of*  
734 *Comparative Psychology* 112, 219–229.  
735  
736 Tuber, D.S., Hennessy, M.B., Sanders, S., Miller, J.A., 1996. Behavioral and glucocorticoid  
737 responses of adult domestic dogs (*Canis familiaris*) to companionship and social  
738 separation. *Journal of Comparative Psychology* 110, 103–108.

739  
740 Vaessen, T., Hernaes, D., Myin-Germeys, I., van Amelsvoort, T., 2015. The dopaminergic  
741 response to acute stress in health and psychopathology: a systematic review.  
742 *Neuroscience and Biobehavioral Reviews* 56, 241-251.  
743  
744 Vas, J., Topál, J., Gácsi, M., Miklósi, A., Csányi, V., 2005. A friend or an enemy? Dogs'  
745 reaction to an unfamiliar person showing behavioural cues of threat and friendliness at  
746 different times. *Applied Animal Behaviour Science* 94, 99–115.  
747  
748 Vonderen, I.K., Kooistra, H.S. and Rijnberk, A.D., 1998. Influence of veterinary care on the  
749 urinary corticoid: creatinine ratio in dogs. *Journal of Veterinary Internal*  
750 *Medicine* 12, 431–435.  
751  
752 Wells, D.L., Hepper, P.G., 1999. Male and female dogs respond differently to men and  
753 women. *Applied Animal Behaviour Science* 61, 341-349.  
754  
755 Wells, D.L., Graham, L., Hepper, P.G., 2002. The influence of auditory stimulation on the  
756 behaviour of dogs housed in a rescue shelter. *Animal Welfare* 11, 385-393.  
757  
758 Wilsson, E., Sundgren, P. E., 1997. The use of a behaviour test for selection of dogs for  
759 service and breeding. II. Heritability for tested parameters and effect of selection  
760 based on service dog characteristics. *Applied Animal Behaviour Science* 54, 235-241.  
761  
762 Ziv, G., 2017. The effects of using aversive training methods in dogs—A review. *Journal of*  
763 *Veterinary Behavior: Clinical Applications and Research* 19, 50–60.  
764

765 **Table 1**

766 Characteristics of owners and dogs.

Dog	Experimental group	Interval between visits (weeks)	Age (years)	Sex	Breed	Age of owner
1	NoO/O	5	3	FN	Mixed Shepherd	25-40
2	NoO/O	5	1	MN	Mixed Retriever	25-40
3	NoO/O	6	4	FN	Cavalier King Charles	41-60
4	O/NoO	5	2.5	FN	Beauceron	25-40
5	O/NoO	7	3	FN	Schapendoes	<25
6	O/NoO	5	5	FE	Mixed Terrier	25-40
7	O/NoO	6	5	FN	Mixed Terrier	>60 <sup>a</sup>
8	O/NoO	6	5	ME	Mixed Terrier	>60 <sup>a</sup>
9	O/NoO	6	7.5	MN	Labrador	41-60
10	O/NoO	6	2.5	ME	Boxer	25-40
11	NoO/O	5	5	MN	Mixed Terrier	25-40 <sup>a</sup>
12	O/NoO	5	6	MN	Mini Australian Shepherd	25-40 <sup>a</sup>
13	NoO/O	7	2.5	MN	Whippet	25-40
14	O/NoO	5	2	ME	Boxer	41-60
15	NoO/O	6	2	MN	French Bulldog	41-60
16	O/NoO	5	1.5	ME	German Shepherd	41-60
17	O/NoO	7	7	FN	Spitz	25-40
18	O/NoO	6	2	FE	Boxer	41-60
19	NoO/O	7	4	MN	Whippet	>60
20	NoO/O	6	3.5	FN	Cotton Tulear	25-40
21	NoO/O	5	3.5	MN	White Swiss Shepherd	25-40

22	NoO/O	7	9	FN	Australian Shepherd	41-60 <sup>a</sup>
23	O/NoO	7	3	FN	Australian Shepherd	41-60 <sup>a</sup>
24	NoO/O	6	4	FE	Groenendael	25-40
25	NoO/O	7	5	ME	Malinois	41-60

---

767 NoO/O, Owner was absent for the first veterinary consultation and present for the second;  
768 O/NoO, Owner was present for the first veterinary consultation and absent for the second; F,  
769 Female; M, Male; N, Neutered; E, Entire. <sup>a</sup> Owners participating with more than one dog.

770 **Table 2**

771 Recorded dog behaviours during the examination and greeting phases and their definitions.

Observed behaviour	Description
Non-exclusive stress-related behaviours	
Scratching <sup>a</sup> / Sniffing <sup>a</sup> / Shivering <sup>a</sup> / Shaking <sup>a</sup>	The dog scratched itself / The dog sniffed the ground or straight ahead / The dog trembled / The dog shook
Low postures <sup>a</sup>	The dog's tail was lowered, its ears faced backwards, or its legs were bent; at least two of these postures were exhibited
Mouth (exclusive behaviours)	
Yawning <sup>a</sup> / Panting <sup>a</sup> / Licking <sup>a</sup>	The dog yawned / The dog panted / The dog licked its mouth
Vocalisations (exclusive behaviours)	
Whining <sup>a</sup> / Barking <sup>a</sup>	The dog whined / The dog barked
Gaze (exclusive behaviours)	
Gaze at a person	The dog gazed with its head oriented towards the owner (Gaze O), the veterinarian (Gaze V), the assistant (Gaze A), or the veterinarian and the assistant (Gaze VA)
Gaze at an object or thing	The dog gazed with its head oriented towards the door (Gaze D) or straight ahead when on the table (Gaze Ad)
Avoidance (exclusive behaviours)	
Avoidance	The dog stepped backwards away from the veterinarian or the assistant following one of their actions
Situation (exclusive behaviours)	

Situation / somebody	Half of the dog's body (head and chest) was situated less than 50 cm from the owner (Situation O) or from the veterinarian and the assistant (Situation VA)
Situation / something	Half of the dog's body (head and chest) was situated less than 1 m from the door (Situation D), or the dog was not in one of the previous locations (Situation E)
Movement (exclusive behaviours)	
Move	The dog moved its four limbs with less than 1 second between the movement of each limb
Contact (exclusive behaviours)	
Contact	The dog intentionally touched the owner (Contact O), the veterinarian (Contact V), or the assistant (Contact A)
Tail (exclusive behaviours)	
Tail wagging	Tail wagged below the spine but was not between the legs
Tail between legs <sup>a</sup>	Tail was between the rear limbs
Tail low	Tail was below the spine but was not wagging or between legs
Tail high	Tail was above the spine, whether wagging or not

---

772 <sup>a</sup>Stress-related behaviour

773 **Table 3**

774 Behavioural indices calculated using several behaviours from Table 2.

775

Index (Unit)	Definition	Formula
Total Stress (%)	Sum of percentages of time spent yawning, panting, scratching, adopting low posture, shivering, sniffing, whining, barking and licking	Total Stress (%) = Yawning (%) + Panting (%) + Shivering (%) + Low postures (%) + Shaking (%) + Sniffing (%) + Whining (%) + Barking (%) + Licking (%)
Contact V+A (%)	Sum of percentages of time spent in contact with veterinarian (V) and/or assistant (A)	Contact V+A (%) = Contact V (%) + Contact A (%)
Gaze V+A (%)	Sum of percentages of time spent gazing at veterinarian (V) and/or assistant (A)	Gaze V+A (%) = Gaze V (%) + Gaze A (%) + Gaze VA (%)
Gaze O+Ad (%)	Sum of percentages of time spent gazing at owner (O) when present or straight ahead (Ad) when absent	Gaze O+Ad (%) = Gaze O (%) + Gaze Ad (%)
Behav. Tow. Owner (%) (dog behaviours towards owner)	Sum of percentages of time spent gazing at, having contact with, and seeking proximity to the owner	Behav. Tow. Owner (%) = Gaze O (%) + Contact O (%) + Situation O (%)

776 %, Time percentage (behaviour duration/phase duration); Yawning, The dog yawned;  
 777 Panting, The dog panted; Scratching, The dog scratched itself; Low posture, The dog's tail  
 778 was lowered, its ears faced backwards, or its legs were bent, at least two of these postures  
 779 were exhibited; Shaking, The dog shook; Sniffing, The dog sniffed the ground or straight  
 780 ahead; Whining, The dog whined; Barking, The dog barked; Licking, The dog licked its  
 781 mouth; Contact V, The dog intentionally touched the veterinarian; Contact A, The dog



782 intentionally touched the assistant; Gaze V, The dog gazed with its head oriented towards the  
783 veterinarian; Gaze A, The dog gazed with its head oriented towards the assistant; Gaze VA,  
784 The dog gazed with its head oriented towards the veterinarian and the assistant; Gaze O, The  
785 dog gazed with its head oriented towards the owner; Gaze Ad, The dog gazed with its head  
786 oriented straight ahead when on the table; Contact O, The dog intentionally touched the  
787 owner; Situation O, Half of the dog's body (head and chest) was situated less than 50 cm from  
788 the owner.

789 **Table 4**

790 Rated emotional states of dogs during exploration, examination, and greeting phases, their  
791 definition, and their score (from Mills et al., 2006).

Emotional state	Definition	Score
Relaxed	No or low frequency of movement, with no visual evidence of tension in the body	1
Aroused	Tense, with high frequency of movement, but no visual evidence of anxious behaviours	2
Anxious	Tense, with licking, yawning, crying, agitation or observable fearful posture	3

792 **Table 5**

793 Rated scores of dogs when entering the examination room (exploration phase) and reuniting  
 794 with the owner after the veterinary examination (greeting phase), along with the name of the  
 795 factor and the signification of the score.

Factor	Name of factor	Score (from 1 to 5)
Apparent comfort in entering the room	Entering Room	1 = Has to be drawn to enter, 5 = Enters voluntarily, pulls on leash
Greeting intensity by dog towards owner	Reunion / Dog	1 = Indifferent, 5 = Very happy, jumps on the owner, requests contact
Greeting intensity by owner towards dog	Reunion / Owner	1 = Indifferent, 5 = Talks to the dog, pets the dog a lot

796 **Table 6**

797 Rated levels of dog restraint performed by the assistant during the examination phase, along  
798 with their definition and score.

Restraint	Definition	Score
Low	The assistant did not need to use force to keep the dog in the right position. The dog was voluntarily almost immobile.	1
Medium	The assistant needed to increase her restraint of the dog to keep it in the same position. The dog was agitated/moved frequently.	2
High	The assistant had to hold the dog firmly to keep it on the examination table or help the veterinarian perform the clinical examination. The dog tried to escape.	3

799 **Table 7**

800 Rated manipulations performed by the veterinarian during the examination phase and the  
 801 meaning of their success and difficulty score.

Manipulation	Definition	Score and Value	
Table	Dog was picked up and lifted onto the examination table	Failure: 0 <sup>a</sup> Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Eye	Eye and mucosa observation	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Ear	Ear manipulation and observation	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Mouth	Examination of teeth and mouth mucosa	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Palpation	Abdominal and lymph node palpation	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Skin fold	Examination of scapular skin fold	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Auscultation	Cardiac and pulmonary auscultation	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Thermometer	Measuring rectal temperature	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard
Paws	Manipulating paws	Failure: 0 Success: 1	Difficulty (from 1 to 5) 1: Easy / 5: Hard

802 <sup>a</sup> Each dog received a score of 0 or 1, these scores were used to calculate the percentage of  
 803 success of all 25 dogs.

804 **Table 8**

805 Emotional state of dogs in a veterinary practice with the presence or absence of the owner  
 806 during the exploration phase. Results are expressed as the standard error of the mean.

Behaviours (units)	Owner: present condition	Owner: absent condition	t	<i>P</i>	1-β (%)
Entering room (score)	3.24 ± 0.76	2.72 ± 1.27	-2.7	0.012	100
Emotional State (score)	1.76 ± 0.63	2.22 ± 0.69	3.682	0.001	69.2

807 Entering room, Rated apparent comfort in entering the room (1 = Dog has to be drawn to  
 808 enter, 5 = Dog enters voluntarily, pulls on leash; also see Table 5); Emotional State, Rated  
 809 emotional states of dogs (1 = Relaxed: No or low frequency of movement, with no visual  
 810 evidence of tension in the body, 2 = Aroused: Tense, with high frequency of movement, but  
 811 no visual evidence of anxious behaviours, 3 = Tense, with licking, yawning, crying, agitation  
 812 or observable fearful posture; also see Table 4).

813 **Table 9**

814 Behaviour of dogs in a veterinary practice with the presence or absence of the owner during  
 815 the examination phase.

Behaviours (units)		Owner: present condition	Owner: absent condition	t	<i>P</i>	1-β (%)
Behaviour towards veterinarian and/or assistant						
Contact - V+A	Duration (% time)	2.33 ± 4.91	1.34 ± 3.86	-2.187	0.039	12.2
Behaviour towards owner and/or door						
Gaze - Door	Duration (% time)	12.46 ± 9.56	25.15 ± 13.52	3.813	0.0008	96.9
Gaze O+Ad	Duration (% time)	37.32 ± 19.33	21.64 ± 12.17	3.342	0.0008	93

816 Contact V+A, Sum of percentages of time spent in contact with veterinarian (V) and/or  
 817 assistant (A); Gaze O+Ad, Sum of percentages of time spent gazing at owner (O) when  
 818 present or straight ahead (Ad) when absent; % time, Percentage of time (behaviour  
 819 duration/phase duration; also see Table 3). Results are expressed as the standard error of the  
 820 mean. Non-significant results are not presented.

821 **Table 10**

822 Behaviour of dogs in a consultation room with the presence or absence of the owner during  
823 the greeting phase.

Behaviours (units)		Owner: present condition	Owner: absent condition	t	P	1-β (%)
Behav. Tow. Owner	Duration (% time)	129.24 ± 50.70	170.76 ± 49.70	3.455	0.002	83.3
Reunion / Dog	Score	3.11 ± 1.29	4.24 ± 0.79	5.106	0.00003	96.2
Gaze - Door	Duration (% time)	8.12 ± 8.95	12.32 ± 13.50	1.818	0.081	25.4

824 Behav. Tow. Owner, Behaviour Towards Owner, Sum of percentages of time spent gazing at,  
825 having contact with, and seeking proximity to the owner (also see Table 3); Reunion / Dog,  
826 Rated greeting intensity by dog towards owner (1 = indifferent, 5 = very happy, jumps on the  
827 owner, seeks contact; also see Table 5); % time, Percentage of time (behaviour duration/phase  
828 duration); Gaze – Door, Percentage of time spent gazing at the door. Results are expressed as  
829 the standard error of the mean. Non-significant results are not presented.



830 **Figure legends**

831

832 Fig. 1. Image of the experimental room, with the two videos assembled in a single image.

833 Left: 'Owner: absent condition'. Right: 'Owner: present condition'

834

835 Fig. 2. Layout of the experimental room. A: veterinarian's chair, B: assistant's chair, C:

836 owner's chair, X: owner's position during the veterinary examination

837

838 Fig. 3. Image of the standardised restraint by the assistant on the examination table. One hand

839 on the chest and another on the base of the tail, with the minimal necessary strength to keep

840 the dog sitting or standing on the table.

841

842

843 **Appendix A**

844 Sex of dogs

	Number of female	Number of male	Total
Number of neutered dog	9 (36 %)	8 (32 %)	17 (68%)
Number of entire dog	3 (12 %)	5 (2 %)	8 (32 %)
Total	12 (48 %)	13 (52 %)	25 (100%)

845

846 Age of dogs

	Number of dogs
1 to 3 years	12 (48 %)
4 to 6 years	10 (40 %)
7 to 9 years	3 (12 %)

9 to 12 years	0 (0%)
---------------	--------

847

848 Sex of owners

	Number of owners
Female	21 (100 %)
Male	0 (0 %)
Total	21 (100 %)

849

850 Age of owners

	Number of owners
Under 25 years	1 (5%)
25 to 40 years	10 (47 %)
41 to 60 years	8 (38 %)
Over 60 years	2 (10%)
TOTAL	21 (100%)

851 Three owners participated with more than one dog.

852

853 **Appendix B**

854 Summary of all studied variables and their availability for the three phases.

Variable	Available in	Available in	Available
	phase 1: exploration	phase 2: examination	in phase 3: greeting
Recorded behaviours			
Scratching	no	yes	yes

Sniffing	no	yes	yes
Low Postures	no	yes	yes
Mouth behaviours	no	yes	yes
Vocalisations	no	yes	yes
Gaze	no	yes	yes
Avoidance	no	yes	no
Situation	no	yes	yes
Movement	no	no	yes
Contact	no	yes	yes
Tail behaviours	no	yes	yes

#### Behavioural indices

Total Stress	no	yes	yes
Contact V+A	no	yes	yes
Gaze V+A	no	yes	yes
Gaze O+Ad	no	yes	no
Behav. Tow. Owner	no	no	yes

#### Scores

Emotional State	yes	yes	yes
Entering room	yes	no	no
Reunion/Dog	no	no	yes

Reunion/Owner	no	no	yes
Restraint	no	yes	no
Success of manipulation (eye, ear, mouth, palpation, skin fold, auscultation thermometer, paws)	N/A	yes	N/A
Difficulty of manipulation (eye, ear, mouth, palpation, skin fold, auscultation thermometer, paws)	N/A	yes	N/A

---

855 Contact V+A, Sum of percentages of time dog spent in contact with veterinarian (V) and/or assistant  
856 (A); Gaze V+A, Sum of percentages of time dog spent gazing at veterinarian (V) and/or assistant (A);  
857 Gaze O+Ad, Sum of percentages of time spent gazing at owner (O) when present or straight ahead (Ad  
858 ) when absent; Behav. Tow. Owner, Sum of percentages of time dog spent gazing at, having contact  
859 with, and seeking proximity to the owner; Emotional State, Rated emotional states of dogs (1 =  
860 Relaxed: No or low frequency of movement, with no visual evidence of tension in the body, 2 =  
861 Aroused: Tense, with high frequency of movement, but no visual evidence of anxious behaviours, 3 =  
862 Tense, with licking, yawning, crying, agitation or observable fearful posture); Entering room, Rated  
863 apparent comfort in entering the room (1 = Dog has to be drawn to enter, 5 = Dog enters voluntarily,  
864 pulls on leash); Reunion / Dog, Rated greeting intensity by dog towards owner (1 = indifferent, 5 =  
865 very happy, jumps on the owner, seeks contact); Reunion/Owner, Rated greeting intensity by owner  
866 towards dog (1 = Indifferent, 5 = Talks to the dog, pets the dog a lot); Restraint, Rated levels of dog  
867 restraint performed by the assistant (1=Low, 3=High);

868

869

870

871

872

873

874

875

876

877

878

879

880


881 Figure 1



882

883

884 Figure 2




885

886

887

888 Figure 3



889