



Headland Rip Very-low-Frequency Fluctuations and Surf Zone Eddies during High-Energy Wave Events

Arthur Mouragues, Kévin Martins, Philippe Bonneton, Bruno Castelle

► To cite this version:

Arthur Mouragues, Kévin Martins, Philippe Bonneton, Bruno Castelle. Headland Rip Very-low-Frequency Fluctuations and Surf Zone Eddies during High-Energy Wave Events. *Journal of Physical Oceanography*, 2022, <10.1175/JPO-D-22-0006.1>. <hal-03831770>

HAL Id: hal-03831770

<https://hal.science/hal-03831770v1>

Submitted on 27 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Headland rip very-low-frequency fluctuations and surf zone eddies**
2 **during high-energy wave events**

3 Arthur Mouragues,^a Kévin Martins,^a Philippe Bonneton,^a and Bruno Castelle^a

4 ^a *CNRS, UMR 5805 EPOC, University of Bordeaux, 33615 Pessac, France*

5 *Corresponding author:* Arthur Mouragues, arthur.mouragues@u-bordeaux.fr

6 ABSTRACT: A wave group-resolving model is used to investigate the driving mechanisms and the
7 spatio-temporal variability of very-low-frequency (VLF) fluctuations of an headland deflection rip,
8 measured during a 4-m oblique wave event. Surf zone eddies (SZE) occurring in the presence of a
9 strongly-sheared longshore current V at a longshore-uniform beach are first modelled. The spectral
10 signature and the variability of SZE are displayed and compared with the literature. The model
11 is then used to explore the dynamics of vorticity in the surf zone and against a headland under
12 energetic oblique wave conditions. The resulting weakly-sheared V is found to host large-scale
13 SZE propagating towards the headland at a speed decreasing seaward. Vorticity animations and
14 spectral diagrams indicate that VLF fluctuations of the deflection rip are driven by the deflection
15 of the upstream SZE. In line with measurements, periods of 40 min to 1 h dominate the spectrum
16 hundreds of meters from the headland at low tide. At high tide, vorticity spectra in the rip are much
17 narrower than in the surf zone, suggesting that the headland enforces the merging of SZE. This
18 mechanism is further analysed using idealized simulations with varying headland lengths, aiming
19 at extending traditional deflection patterns at the VLF scale. Finally, we discuss the existence of a
20 continuum in SZE driving mechanisms, going from fully wave group-driven to both wave group-
21 and shear instability-driven SZE for weakly- and strongly-sheared V , respectively. This continuum
22 suggests the importance of wave groups to produce SZE under energetic wave conditions.

23 1. Introduction

24 Surf zone eddies (SZE) are two-dimensional horizontal vorticity motions typically associated
25 with frequencies as low as a few millihertz, namely within the very-low-frequency (VLF) band
26 (*e.g.* MacMahan et al. 2004; Reniers et al. 2007). Besides broadly contributing to mixing
27 and dispersion processes in the nearshore (*e.g.* Clark et al. 2010), SZE can also affect the spatio-
28 temporal variability of longshore currents, thus impacting the transport of sediments and pollutants
29 (Deigaard et al. 1995). Several SZE driving mechanisms have been proposed (*e.g.* Bowen and
30 Holman 1989; Long and Özkan-Haller 2009; Feddersen 2014). The first one that was put forward
31 is associated with the presence of shear waves.

32 Shear waves are alongshore progressive vortical structures propagating in the surf zone with
33 periods and alongshore wavelengths of $O(100)$ s and $O(100)$ m, respectively. These motions are
34 commonly observed in the presence of a relatively strong mean (time-averaged) longshore current
35 ($O(1)$ m/s; *e.g.* Oltman-Shay et al. 1989; Dodd et al. 1992) and result from instabilities due to
36 cross-shore shear of the longshore current (Bowen and Holman 1989). They appear in the form of
37 a meandering longshore current depicted by SZE with rather large velocity fluctuations. Observed
38 root-mean-squared velocity amplitudes of such SZE can reach half of the mean longshore current
39 magnitude (Oltman-Shay et al. 1989; Lippmann et al. 1999; Noyes et al. 2004). These highly-
40 energetic motions can therefore span the entire surf zone, altering the nearshore circulation and, in
41 particular, the cross-shore distribution of the longshore current. The presence of shear instability-
42 driven SZE can lead to a substantial cross-shore mixing of momentum in the surf zone, resulting
43 in a smoother longshore current profile (Özkan-Haller and Kirby 1999). Surf zone eddies can also
44 be ejected seaward giving rise to spatially- and temporally- transient rip currents (Özkan-Haller
45 and Kirby 1999), representing a dangerous hazard for swimmers (Castelle et al. 2016).

46 Shear wave motions in the nearshore were first theoretically explained by Bowen and Holman
47 (1989), who highlighted the importance of the seaward shear of the longshore current (cross-shore
48 gradient of the longshore current magnitude) to characterize these instabilities. For a strongly-
49 sheared longshore current, the latter becomes unstable, disturbances (or unstable modes) with
50 different wavelengths (wavenumber) are generated and propagate in the same direction of the
51 longshore current at a speed proportional to the longshore current peak. This was confirmed by
52 Oltman-Shay et al. (1989) who analysed measurements collected at a longshore-uniform sandy

53 beach under moderately-energetic and highly-oblique wave conditions (offshore significant wave
54 height H_s of 1.36 m and peak angle of wave incidence θ_p of 30°). Since the works of Bowen
55 and Holman (1989) and Oltman-Shay et al. (1989), shear wave motions have been investigated in
56 many studies which have been extensively reviewed in Dodd et al. (2000). The spectral signature
57 of shear waves is very specific within the longshore wavenumber-frequency (k - f) spectrum. Most
58 shear waves-related energy is spread around a relatively linear dispersion relation located outside of
59 the region of surface gravity motions, indicating that, at a given cross-shore position, all unstable
60 modes of these vorticity motions propagate at the same speed. By analysing the cross-shore
61 variability of shear waves, Noyes et al. (2004) suggested that the celerity of such motions is related
62 to the local value of the mean longshore current. Subsequent two-dimensional horizontal (2DH)
63 modelling studies have highlighted the mechanisms for shear wave energy dissipation through
64 bottom friction and horizontal mixing (*e.g.* Dodd et al. 1992; Falqués and Iranzo 1994; Özkan-
65 Haller and Kirby 1999). In particular, Özkan-Haller and Kirby (1999) used a phase-averaged
66 model to depict complex vortex structures that result from shear instabilities. These structures are
67 essentially made of energetic vorticity fronts and detaching eddies.

68 Including the effect of wave groups, Long and Özkan-Haller (2009) showed that the production
69 of vorticity due to wave groups and shear instabilities can be of the same order. The inclusion of
70 wave groups was shown to result in broader k - f spectra and a more chaotic vorticity field. These
71 authors also showed that the vorticity due to wave groups was dominant over shear instabilities
72 for weakly-oblique incident waves. Using a phase-resolving model, Feddersen (2014) showed that
73 breaking wave vorticity forcing, which includes both breaking wave group and individual wave
74 vorticity forcing, are the dominant vorticity generation mechanisms, compared to shear instabilities,
75 except for highly-oblique large waves. Overall, these results suggest that SZE are not necessarily
76 always driven by shear instabilities but are driven by mechanisms that may change depending on
77 the longshore current profile which can substantially vary with incident wave conditions.

78 A significant number of measurement and modelling studies have investigated the spatio-temporal
79 variability of surf zone eddies in the presence of a longshore current. However, most of these
80 studies have been conducted for SZE propagating along longshore-uniform (barred or planar)
81 sandy beaches under low- to moderate- energy wave conditions, with H_s rarely exceeding 2 m. The
82 eddy regime under energetic wave conditions is therefore poorly understood. For such conditions,

83 the longshore current is generally wider, potentially resulting in a different seaward longshore
84 current shear than under less energetic wave conditions. The cross-shore structure of the longshore
85 current is yet known to play an important role in the shear wave regime such as their characteristic
86 spatial and temporal scales and the range of unstable modes (*e.g.* Bowen and Holman 1989; Dodd
87 and Thornton 1990; Baquerizo et al. 2001). Although never addressed, the variability of SZE and
88 their driving mechanisms, under high-energy conditions can be potentially different than under
89 low- to moderate-energy conditions. Lastly, there is a lack of knowledge on the behaviour of SZE
90 propagating along longshore non-uniform beaches (Dodd et al. 2000). In particular, the dynamics
91 of SZE propagating over complex morphologies with geological constraints, such as bedrocks or
92 headlands that are commonly present along rugged coastlines, is virtually unknown.

93 Along rugged or artificial coastlines, the presence of physical boundaries, such as natural head-
94 lands or man-made structures, can significantly disrupt the nearshore circulation. Under obliquely-
95 incident wave conditions, wave breaking induces a longshore current that can be deflected against
96 boundaries (*e.g.* Scott et al. 2016), creating a so-called headland deflection rip (Castelle et al. 2016).
97 This type of rip was measured during a three-week field experiment conducted at La Petite Cham-
98 bre d'Amour (PCA; Anglet, SW France) beach in October 2018 (Mouragues et al. 2020b). The
99 field site comprises complex morphological features such as bedrocks and a 500-m rocky headland
100 (Saint Martin headland in Figure 1g). During a 4-m oblique wave event, the longshore current was
101 deflected seaward against the headland, resulting in a strong tidally-modulated deflection rip ex-
102 tending hundreds of meters offshore (Figure 1a-f). At low tide, energetic very-low-frequency (VLF)
103 fluctuations of the deflection rip were measured 800 m offshore, with dominant (most energetic)
104 periods around 30 min and 1 h (see SIG1 in Figure 1; Mouragues et al. 2020a). These fluctuations
105 were successfully reproduced by Mouragues et al. (2021) using a wave group-resolving 2DH model
106 (XBeach SurfBeat mode, hereafter XB-SB, Roelvink et al. 2009) but their driving mechanisms
107 and their spatio-temporal variability were not addressed. Following Mouragues et al. (2021), the
108 present study uses XB-SB to investigate these fluctuations and their driving mechanisms.

115 The paper is organised as follows. The Section 2 briefly presents the field site and the experimental
116 setup. In Section 3, the XB-SB model, its implementation and methods for analysing rotational
117 motions which includes the computation of their k - f spectra and their bulk characteristics are
118 described. In Section 4, the model is used to simulate surf zone eddies in the presence of a




FIG. 1. Time series of water depth h_0 (a and b), 5 min-running averaged cross-shore velocity U_C (c and d) and longshore velocity U_L (e and f) measured at two instrument locations (SIG1 and AQ, respectively) on the 7th of October 2018. The panel g shows the location of the field site (PCA; grey rectangle), the two instrument locations (black and blue points) and the approximate direction of the measured current (black and blue arrows based at the points). The offshore significant wave height and peak angle of wave incidence (H_s and θ_p) are also indicated. Note that the bathymetry map and the entire array of instruments are shown in Figure 2.

longshore current measured during the SandyDuck experiment (Duck, North Carolina). This first modelling experiment ensures that the model is able to reproduce SZE at a longshore-uniform sandy beach under moderate wave conditions. Model performances are assessed by comparing bulk characteristics of eddies and their spatio-temporal variability with previous studies. The ability of the model to reproduce SZE observed during the SandyDuck setup suggests that it can be used to simulate such motions occurring over more complex morphology and under more energetic wave conditions and to further explore their variability. Hence, in Section 5, we investigate surf zone eddies and headland rip VLF fluctuations at PCA under energetic wave conditions and for different tidal levels. For this investigation, 20 hour-long model simulations with a constant tidal level are set up in order to simulate a significant number VLF fluctuations (with periods $< O(1)$ h) and to fix the cross-shore location of wave breaking, thus removing the tidal modulation of surf zone circulation which was strong in the measurement dataset (see Figure 1a-f). These model

131 setups therefore subsequently allow addressing, through statistically-significant spectrum analysis
132 and a stationary state of VLF surf zone currents, the correct exploration of such motions. The
133 mean (time-averaged) circulation patterns and the presence of SZE at low tide and high tide are
134 then emphasized. The last part of Section 5 presents the spatio-temporal (frequency) variability
135 of vorticity in the surf zone and along the headland, showing that VLF fluctuations measured and
136 modelled off the headland tip are related with SZE propagating in the upstream longshore current.
137 In Section 6, the findings of this study are discussed. The discussion includes the control of
138 morphology on VLF fluctuations, with suggestions for future research. The effect of wave group
139 variability on surf zone rotational motions is also discussed, suggesting that wave group forcing
140 may be the primary driver of VLF fluctuations of the rip during the high-energy wave event, rather
141 than shear instabilities of the longshore current. Conclusions and perspectives for future works are
142 finally drawn in Section 7.

143 2. Field experiment

144 From the 3rd to the 26th of October 2018, a field experiment was conducted at La Petite Chambre
145 d'Amour (PCA; Figure 2a) beach located in Anglet in the south of the Aquitaine coast (SW France).
146 This rugged coast is a mesotidal high-energy environment that is regularly exposed to energetic
147 Atlantic swells coming from the W-NW direction (Abadie et al. 2005). PCA is a double-barred
148 sandy beach located at the southern end of a 4-km embayment, comprising six groynes, bounded
149 by the Adour river to the North and by the Saint Martin 500-m rocky headland to the South (Figure
150 1g). The reader is referred to Mouragues et al. (2020a,b) for a detailed description of the field site
151 and of the experiment.

152 A large array of instruments were deployed to measure and study the natural variability of
153 wave-induced circulation at a high-energy geologically-constrained beach. In particular, Acoustic
154 Doppler Current Profilers (ADCPs) were installed near the headland to collect high-frequency
155 Eulerian velocity measurements (Figure 2a). On the 7th of October 2018, obliquely-incident
156 energetic waves ($H_s \approx 4.0$ m and $\theta_p \approx 20^\circ$) induced an intense deflection rip flowing against the
157 headland. At low tide, high velocities were measured at the most offshore ADCP located 800-m
158 offshore sitting in 12-m depth (Figure 1 and 2a). Time-averaged current velocities showed energetic
159 VLF fluctuations with dominant periods of around 30 min and 1 h, and associated peak velocities

up to 0.7 m/s (see *e.g.* Figure 1c,e). These fluctuations were numerically reproduced by Mouragues et al. (2021) using XB-SB which model domain is shown in Figure 2b. The analysis of modelling results further suggested that the rip actually extended up to 1600 m offshore and was strongly modulated by tides (see Figure 1a-f; Mouragues et al. 2020a, 2021). The modelling approach used in the present study is similar to that of Mouragues et al. (2021) and is described in the following section.

3. Modelling surf zone vortical motions

a. Modelling strategies

Over the past decades, several numerical modeling strategies have been employed in order to simulate surf zone eddies. These approaches mostly differ in terms of the wave scale resolved, ranging from phase-averaged models excluding wave groups (hereafter referred to as fully phase-averaged models; *e.g.* Özkan-Haller and Kirby 1999; Noyes et al. 2005) or including wave groups (hereafter referred to as wave group-resolving models; *e.g.* Reniers et al. 2007; Long and Özkan-Haller 2009) to models simulating motions at the individual wave scale (Feddersen 2014). Fully phase-averaged models have been applied to assess, in particular, the effect of bottom friction and horizontal mixing on shear waves and their cross-shore variability. Because these models use a steady forcing to drive the nearshore circulation (*i.e.* averaged over many wave groups), they can only reproduce very-low-frequency (VLF) scales ($f < 0.004$ Hz) associated with shear instabilities of the mean current, and not vorticity motions at the wave group scale. However, accounting for the variability of the wave forcing at scales at least similar to wave groups is essential for better reproducing surf zone rotational motions (Long and Özkan-Haller 2009; Feddersen 2014).

Feddersen (2014) used a Boussinesq model to simulate the dynamics of SZE that were measured during the SandyDuck experiment. Such a modelling approach allows the simulation of the vorticity field generated by shear instabilities and through wave breaking at both the individual and wave group scales. However, the sub-metric spatial resolution required to accurately reproduce motions at this scale still makes the use of these models computationally challenging over large spatial domains ($> O(10) \text{ km}^2$) and for long periods of time (temporal scale of a storm). In this context, wave group-resolving models appear as a good compromise since they are much less costly than fully phase-resolving approaches while keeping the ability of these models to




FIG. 2. Bathymetry map of the field site (PCA; (a); black lines show 1-m spaced elevation contours) which location is shown by a black rectangle in the full model domain ((b); black lines show 2-m spaced elevation contours). Colour indicates elevation relative to the mean sea level (m+MSL), blue line is the MSL contour and coloured points indicate ADCP location. (c) and (d) show the cross-shore (Δx) and longshore (Δy) mesh step size, respectively.

reproduce the vorticity generated by wave groups. Wave group-resolving models such as XB-SB use a wave forcing varying at the wave group scale to drive the nearshore circulation, allowing to simulate low-frequency surf zone motions which include infragravity and VLF motions. This approach has been used by several authors to reproduce the low frequency variability of circulation

198 along rip-channelled open beaches (Reniers et al. 2006, 2007) and to simulate vorticity motions
 199 that result from obliquely-incident wave groups along a longshore-uniform open beach (Long
 200 and Özkan-Haller 2009). The XB-SB wave group-resolving approach has here been chosen to
 201 investigate the driving mechanism and the spatio-temporal variability of VLF fluctuations of the
 202 deflection rip at PCA. XBeach is a morphodynamic model initially developed to reproduce storm
 203 response of sandy beaches where infragravity swash is dominant (Roelvink et al. 2009). It solves
 204 the coupled two-dimensional horizontal (2DH) equations for wave propagation, flow, sediment
 205 transport and bottom changes (see Roelvink et al. 2009 for a model description). The short-wave
 206 effects on currents are modelled through the radiation stress gradients approach (Longuet-Higgins
 207 and Stewart 1964). In this paper, sediment transport and bottom change modules are disabled as
 208 only hydrodynamics (longshore current and deflection rip) is investigated.

209 *b. Implementation of the XBeach model*

210 Based on the input wave spectrum, the Surfbeat approach uses a random phase selection procedure
 211 to reconstruct time series of free surface elevation, from which the short-wave energy varying at
 212 the wave-group scale can be extracted and used to force the spectral wave model (see Roelvink
 213 et al. 2009, 2018). The circulation model is forced with elevations and currents corresponding to
 214 the incident bound infragravity waves computed from the generated time series of surface elevation
 215 following Herbers et al. (1994). The free surface elevation time series are generated using a random
 216 set of phases, but the model allows them to be saved so that the exact same boundary conditions
 217 can be used between numerical tests. This method will often be used in this paper to analyse the
 218 spatial-temporal variability of hydrodynamics (*e.g.* vorticity) between different model setup (*e.g.*
 219 morphological setup, free parameter calibration).

220 The XBeach model includes several free parameters requiring calibration with measurements.
 221 For the modelling experiments carried out in this paper, three free parameters were found to
 222 significantly impact the modelled hydrodynamics : the breaking parameter γ , a bed friction Chezy
 223 coefficient C and a mixing free parameter c_s . The first two parameters are important, in particular,
 224 to the surf zone width and to the intensity of the longshore current, respectively. The third
 225 parameter alters the value of the eddy viscosity ν_h that controls the horizontal mixing. Increasing
 226 the eddy viscosity is known to have a damping effect on shear instabilities along with increasing

227 their longshore length scales (*e.g.* Falqués and Iranzo 1994; Özkan-Haller and Kirby 1999). In
 228 XBeach, ν_h can be parametrised, using a Smagorinsky model, as a function of c_s , velocity spatial
 229 gradients and mesh step sizes :

$$\nu_h = c_s^2 2^{1/2} \sqrt{\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial v}{\partial y}\right)^2 + \frac{1}{2} \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}\right)^2} \Delta x \Delta y, \quad (1)$$

230 where Δx and Δy are the mesh step sizes.

231 Calibration of these three parameters was made by finding their values that minimize dis-
 232 crepancies between model outputs and measurements (running-averaged velocities or rotational
 233 velocities). Model calibration results are described in Sections 4 and 5 and are also indicated in
 234 Table A1 (see Appendices). Finally, it should be noted that the effect of currents on waves was
 235 neglected.

236 *c. Vortical motion analysis*

237 1) WAVENUMBER-FREQUENCY SPECTRUM

238 To investigate surf zone eddy motions, wavenumber-frequency spectra ($k - f$ where k is the
 239 longshore wavenumber in m^{-1}) of the cross-shore velocity u , the longshore velocity v and of the
 240 associated vorticity $q = dv/dx - du/dy$ are computed. Along a longshore transect located at a
 241 given cross-shore position, modelled u , v and q are outputted every 10 s (sampling frequency f_s
 242 = 0.1 Hz) and every Δy m (longshore mesh size; sampling longshore wave number $k_s = 1/\Delta y$)
 243 and stored into 2D-matrices. A 2D-FFT is then applied to these matrices to estimate the energy
 244 density at each wavenumber and frequency $E(k, f)$. Wavenumber-frequency spectra of u , v and q
 245 are hereafter referred to as $E^u(k, f)$, $E^v(k, f)$ and $E^q(k, f)$, respectively.

246 Wavenumber-frequency spectra are typically partitioned into a gravity region and a non-gravity
 247 region (see Figure 3a) which are separated by the mode 0 edge wave dispersion line which is
 248 computed following Eckart (1951). The gravity region typically hosts gravity wave motions such
 249 as edge waves which can remain trapped within the surf zone and propagate down the coast or leaky
 250 waves which may exit the surf zone and propagate seaward. On the other hand, the non-gravity
 251 region may host energetic rotational motions such as shear waves. Figure 3b shows a typical
 252 $E(k, f)$ that is computed in this study and highlights the presence of vorticity motions propagating

in the direction of the mean longshore current V ($k < 0$). The spectral signature of these motions resembles the spectral signature of shear waves consisting of a near-linear dispersion line and for which a representative eddy celerity can be estimated (*e.g.* Oltman-Shay et al. 1989; Özkan-Haller and Kirby 1999; Noyes et al. 2004).




FIG. 3. (a) Schematic of a k - f spectrum ($E(k, f)$) illustrating gravity and non-gravity regions (after MacMahan et al. 2004). These regions are separated by the mode 0 edge wave (gray lines; computed with slope $\tan(\beta) = 0.03$). (b) Zoom of a modelled k - f spectrum of cross-shore velocities $E^u(k, f)$ showing eddies propagating in the direction of the longshore current. The solid gray line shows the mode 0 edge wave dispersion line. Thin dashed black lines indicate the upper and lower dispersion lines (f_{up} and f_{low} , respectively) used to compute the rotational motion frequency $f_{rot}(k)$ (gray points). The black line shows the dispersion line resulting from the fitting of $k - f_{rot}$ and whose slope is the estimated eddy celerity C_{rot} . The thick dashed black line shows the dispersion line of the local longshore current magnitude $V(x)$.

2) EDDY CELERITY

A representative eddy celerity C_{rot} can be estimated as the $E(k, f)$ ridge slope. To estimate C_{rot} , a method similar to Özkan-Haller and Kirby (1999) is employed. For each wavenumber, a rotational motion frequency $f_{rot}(k)$ is computed by integrating $E(k, f)$ over a rotational motion

269 frequency region :

$$f_{\text{rot}}(k) = \frac{\int_{f_{\text{low}}}^{f_{\text{up}}} f E(k, f) df}{\int_{f_{\text{low}}}^{f_{\text{up}}} E(k, f) df}, \quad (2)$$

270 where f_{up} and f_{low} are the upper and lower cutoff frequency lines delimiting the k - f region
 271 attributed to surf zone eddies (dashed thin black lines in Figure 3b). These cutoff lines were set
 272 to $f_{\text{up}} = 4V(x)k$ and $f_{\text{low}} = V(x)(k+0.01)$, with $V(x)$ being the value of the mean longshore current
 273 (dashed thick black line in Figure 3b) at the cross-shore position x , where the longshore transect
 274 is located. This method leads to the points k - f_{rot} (grey points in Figure 3b) that are used to fit
 275 a straight dispersion line whose slope corresponds to the estimated C_{rot} (plain thick black line
 276 in Figure 3b). For the fitting procedure, each data point k - f_{rot} is weighted by the total energy
 277 $\int_{f_{\text{low}}}^{f_{\text{up}}} E(k, f) df$. It should be noted that, as its name suggests, C_{rot} is a celerity representative of the
 278 most energetic rotational motions. As shown by the following results, interactions between eddies
 279 may occur which tends to increase discrepancies between the individual eddy celerity and C_{rot} .
 280 Note that the spectral signature of such eddy interactions is emphasized by the relative broadness
 281 of the k - f spectrum (see *e.g.* Figure 3b)

282 3) ROTATIONAL VELOCITIES

283 The total frequency spectrum $E(f)$ is computed by integrating $E(k, f)$ over all wavenumber as :

$$E(f) = \int_{-k_N}^{+k_N} E(k, f) dk, \quad (3)$$

284 where $k_N = k_s/2$ is the Nyquist wave number. The spectrum of rotational motions $E_{\text{rot}}(f)$ is
 285 calculated by integrating $E(k, f)$ over regions outside the mode 0 edge wave dispersion line
 286 ($k_{0-}(f)$ and $k_{0+}(f)$; see Figure 3a) :

$$E_{\text{rot}}(f) = E(f) - E_{\text{gw}}(f), \quad (4)$$

287 where the spectrum of irrotational motions $E_{\text{gw}}(f)$ is given by :

$$E_{\text{gw}}(f) = \int_{k_{0-}(f)}^{k_{0+}(f)} E(k, f) dk. \quad (5)$$

288 The total, cross- and longshore root-mean-square rotational velocities ($U_{\text{rms,rot}}$, $u_{\text{rms,rot}}$ and $v_{\text{rms,rot}}$,
 289 respectively) are then computed as :

$$(U_{\text{rms,rot}})^2 = \int_{f_1}^{f_2} (E_{\text{rot}}^u(f) + E_{\text{rot}}^v(f)) df, \quad (6a)$$

$$(u_{\text{rms,rot}})^2 = \int_{f_1}^{f_2} E_{\text{rot}}^u(f) df, \quad (6b)$$

$$(v_{\text{rms,rot}})^2 = \int_{f_1}^{f_2} E_{\text{rot}}^v(f) df, \quad (6c)$$

290 where f_1 and f_2 are the lower and upper cutoff frequencies. It should be noted that these rotational
 291 velocities are velocities associated with rotational motions which include motions resulting from
 292 shear instabilities of the mean longshore current and from wave group forcing (when included).

293 4. Assessment of the model at a longshore-uniform sandy beach

294 In this section, the model ability to accurately simulate the dynamics of surf zone eddies propa-
 295 gating at a longshore-uniform sandy beach is investigated. This section serves as a model validation
 296 step for simulating SZE arising over a 2D barred beach under moderate energy wave conditions.
 297 In addition, a brief sensitivity analysis to different parameters is conducted, whose results are
 298 compared with previous studies.

299 The model is setup at the longshore-uniform barred sandy beach of Duck (North Carolina), for
 300 which many shear wave studies build on (see Dodd et al. 2000 for a shear wave study review). The
 301 model is calibrated against measurements collected on the 1st of November 1997 as part of the
 302 SandyDuck experiment described in Noyes et al. (2004). This event was characterised by highly-
 303 oblique and moderate-energy wave conditions, with offshore $H_s = 1.49$ m and mean wave angle
 304 $\theta = 21^\circ$. The latter conditions produced a strong and narrow longshore current V , peaking around
 305 0.9 m/s (time-averaged) with a seaward width Δ (cross-shore distance between the peak position
 306 and the offshore near-zero current intensity) of around 150 m (Noyes et al. 2005; Feddersen 2014).
 307 This strongly-sheared current (mean seaward shear $V_{\text{max}}/\Delta \approx 0.006 \text{ s}^{-1}$) was unstable and hosted
 308 longshore-propagating surf zone eddies driven, at least partially, by shear instabilities (Noyes et al.
 309 2005; Feddersen 2014).

310 In the following sections, XB-SB is first calibrated against measurements presented in Noyes
311 et al. (2005). The calibrated model is then used to investigate the spatio-temporal variability of
312 vorticity which is compared with previous studies. The computational domain extends 1000 m and
313 1000 m in the cross-shore x and longshore y direction, respectively, with mesh resolution set to 2
314 m. The bathymetry is made of a longshore-uniform cross-shore depth profile that is representative
315 of the instrumented transect, with the shoreline and a submerged inner sandbar located around $x =$
316 110 m and $x = 160$ m.

317 The model is forced by a Jonswap spectrum with the same bulk parameters as measurements
318 (H_s , peak period T_p , θ and directional spreading σ_θ) estimated at the 8-m depth pressure gauges
319 array (Long 1996). Similar to Noyes et al. (2005), the period of simulation is set to 8 h with the
320 last 4 h used for analysis. It should be noted that a sensitivity analysis to the longshore domain
321 length and mesh resolution was conducted and indicated that these parameters have a little impact
322 on eddy bulk characteristics (not shown). Note that a mesh resolution of 1 m and 2 m gave similar
323 results while a mesh resolution of 5 m resulted in lower rotational velocities.

324 *a. Model calibration*

325 Figure 4 displays the sensitivity of the time-averaged longshore current $V(x)$ (Figure 4b) and
326 rotational velocities (Figure 4c, d) to bottom friction and horizontal mixing. Bottom friction Chezy
327 coefficient $C = 55 \text{ m}^{1/2}/\text{s}$ best replicates the measured $V(x)$ while the eddy viscosity coefficient
328 $c_s = 1$ correctly reproduces the spatial distribution of both cross-shore and longshore rotational
329 velocities. Both these parameter values are taken for the following analysis (see Table A1). Of
330 note, this eddy viscosity coefficient leads to a cross-shore profile of eddy viscosity (Figure 4a) that
331 is relatively similar to the ones shown in Özkan-Haller and Kirby (1999).

332 In line with previous studies (*e.g.* Özkan-Haller and Kirby 1999), increasing the horizontal
333 mixing (increasing c_s) leads to less energetic rotational motion amplitudes. In addition, it should
334 be noted that XB-SB is slightly better able at computing these amplitudes compared to the fully
335 phase-averaged model used in Noyes et al. (2005). This is because including wave groups leads to
336 a broader spectrum than when excluding wave groups (Long and Özkan-Haller 2009). The control
337 of wave groups on the dynamics of rotational motions will be discussed in Section 6. Lastly,
338 it should be noted that a local bottom slope-dependent wave dissipation coefficient (following

Pezerat et al. 2021) was implemented and used for the SandyDuck modeling experiment. The latter implementation allows to prevent a substantial over-dissipation of incident wave energy occurring seaward of the bar. Such a slope-dependent coefficient was not used for the Anglet modeling experiment as the cross-shore distribution of the longshore current was not measured during the field experiment.




FIG. 4. Bottom friction (C) and horizontal mixing (c_s) coefficient sensitivity to the mean longshore current and to rotational motions (including shear waves). Cross-shore profile of the horizontal viscosity ν_h (a; for $C = 55 \text{ m}^{1/2}/\text{s}$), of the mean longshore current V (b) and of root-mean-square cross- and longshore rotational velocities $u_{\text{rms,rot}}$ and $v_{\text{rms,rot}}$ (c and d, respectively; for $C = 55 \text{ m}^{1/2}/\text{s}$). Yellow points indicate observations from Noyes et al. (2005).

b. Surf zone eddy variability

The calibrated model is now used to explore the spatio-temporal variability of vorticity. Figure 5 shows wavenumber-frequency spectra of vorticity ($E^q(k, f)$) at four cross-shore positions seaward of the longshore current peak position. The spectra of modelled vorticity are in qualitative agreement with the spectra of observed velocities shown in Noyes et al. (2005), exhibiting, in particular, the presence of shear waves propagating in the direction of V . In line with previous studies, shear wave-related energy and the range of energetic frequencies decrease seaward. The most energetic longshore length and period are order of 200 m and 200 s, respectively. Because

both irrotational and rotational motions populate velocity spectra, shear wave motions are best detected using vorticity spectrum and will hereafter used to investigate rotational motions. An example of u , v and q spectra is provided in Figure B1 to illustrate this point.

Although the eddy celerity estimated from velocities can be up to 20 % higher than celerity estimated from vorticity, the entire set of estimated celerities are of the same order of magnitude and is proportional to the local value of the mean longshore current (see Figure 6), which is in agreement with past studies (Noyes et al. 2004, 2005). Eddy celerities reach the longshore current peak at its peak position and decrease seaward. For each estimated celerity, its standard deviation computed for different eddy viscosity coefficients is indicated, showing that horizontal mixing does not strongly impact eddy celerities (with a maximum deviation of 6 %), which is in line with the literature (Özkan-Haller and Kirby 1999).




FIG. 5. (a) Cross-shore profile of the modelled mean longshore current $V(x)$. Yellow points indicate observations. (b)-(e) Modeled wavenumber-frequency spectra of vorticity q ($E^q(k, f)$) at the same cross-shore positions of observations ($x = 160, 210, 260$ and 310 m). For each panel, the local value of the mean longshore current V and the estimated eddy celerity C_{rot} are indicated. The definition of multiple dispersion lines is the same as in figure 3b.




FIG. 6. **(a)** Eddy celerity C_{rot} extracted from cross-shore velocities (blue; C_{rot}^u), longshore velocities (red; C_{rot}^v) and vorticity (black; C_{rot}^q). **(b)** Cross-shore profile of rotational velocities (points) and mean longshore current (black line). For both panels, points show the averaged celerities (for different c_s shown in Figure 4) while vertical bars show the associated standard deviation. In panel (b), yellow squares show observed eddy celerities extracted from cross-shore velocities ($C_{\text{rot}}^{u,\text{obs}}$).

The spatio-temporal variability of vorticity modelled at Duck is shown in Figure 7. This figure displays snapshots and time-space diagram of the 100-s running averaged vorticity along a seaward transect, emphasising the presence of longshore-propagating vorticity fronts and detaching eddies. Snapshots indicate the presence of a vorticity front pair (positive and negative fronts) located near the longshore current peak position, propagating with the longshore current (right to left) and with longshore lengths of around 200 m. Some vorticity fronts may sometimes stretch until breaking down into two fronts with the main front continuing its course with the longshore current and the other secondary front being expelled offshore (detached eddy). Two successive fronts may also merge together which results in a larger front. It is suggested that such eddy interactions result in some discrepancies between the mean current and the estimated celerity seaward of the mean current peak position (see Figure 6; Long and Özkan-Haller 2009). To better visualize vorticity fronts and their interactions, the reader is referred to the vorticity animation (see Supplementary

materials). All these results are in good agreement with vorticity patterns previously modelled for similar setup (Özkan-Haller and Kirby 1999; Long and Özkan-Haller 2009).




FIG. 7. Left-hand panels : snapshots of vorticity at four different times ($t = 30, 36, 42$ and 48 min) depicting the stretching and the splitting of a vorticity front. The dashed black line correspond to the cross-shore position $x = 260$ m. Right-hand panel : time-space diagram of vorticity along the cross-shore position $x = 260$ m. Dashed black lines correspond to times at which vorticity field is plotted on the left-hand panels. For a better visualisation, the reader is referred to vorticity animations (see Supplementary Materials).

The previous results indicate that XB-SB is able to simulate the dynamics of surf zone eddies at a longshore-uniform sandy beach under moderate energy wave conditions. The main characteristics (length and time scales, amplitudes) of SZE are well reproduced by the model. In the next sections, the model is used to investigate SZE and VLF fluctuations of an headland deflection rip occurring at a geologically-constrained beach and under high-energy wave conditions.

5. Surf zone eddies and headland rip VLF fluctuations at a geologically-constrained beach

XB-SB was implemented on a regular grid extending 4000 m and 10000 m in the cross-shore x and the longshore y direction, respectively (Figure 2b). The computational domain comprises morphological features such as offshore bedrock and sand deposit lobe off the river mouth, the

Adour dike and the six groynes along Anglet beaches, PCA and the Saint Martin headland and the adjacent embayment GPB. The mesh step size was set to 5 m at PCA, gradually increasing to 25 m close to the offshore boundaries (Figure 2c, d).

Mouragues et al. (2021) describe how the present model was calibrated at the Anglet site using realistic forcing for waves (wave buoy Candhis 06402 moored in 50 m water depth) and water levels collected at a nearby tidal gauge during the energetic event considered here (event D2 on the 7th of October 2018). In the present study, the VLF fluctuations of the deflection rip are investigated with JONSWAP spectra generated using the mean offshore wave conditions averaged over the considered event ($H_s = 4$ m, $T_p = 10$ s and $\theta_p = -20^\circ$). The peak enhancement factor and the directional wave spreading are set to 3.3 and 24° (XBeach default values). This spectrum is used to force the model for 20 hours in order to ensure that a significant number of VLF fluctuations are modelled. In order to disregard model spin-up, the last 12 hours of the simulation are used for the investigation. The tide elevation is set constant during the entire simulation: $\zeta_{\text{tide}} = -2$ m and $\zeta_{\text{tide}} = 2$ m for the low tide and high tide simulation, respectively. As previously explained, using a constant tidal elevation throughout the simulations ensures that surf zone VLF motions are stationary.

Following the model calibration carried out by Mouragues et al. (2021), $\gamma = 0.50$ and $C = 45$ m^{1/2}/s will be used for the Anglet modelling experiment. Finally, it should be noted that modelled deflection rip velocities in Anglet were found insensitive to the value of c_s (not shown; results are similar to Mouragues et al. 2021). For consistency with the SandyDuck modelling experiment, c_s was set to 1 for all Anglet modelling experiments.

a. Mean circulation and surf zone eddies

Figure 8 shows the mean velocity field along PCA at low tide (panel a) and high tide (panel b). The breaking of highly-oblique and highly-energetic incident waves ($H_s \approx 4.0$ m and $\theta_p \approx 20^\circ$) induces a strong and wide longshore current V oriented to the south. The latter is deflected seaward against the headland, creating a deflection rip extending hundreds of meters off the headland tip. At low tide, the surf zone width is similar to the headland length and the longshore current is deflected by the headland and by the adjacent embayment flow (Mouragues et al. 2021).

434 The surf zone morphology of PCA is relatively longshore-uniform with a relatively steep upper
 435 beach face and a low tide terrace that can be exposed at low tide. For a given incident wave
 436 condition, changes in tidal elevation alter the pattern and intensity of depth-induced breaking wave
 437 energy dissipation which, in turn, results in a strong tidal modulation of the cross-shore distribution
 438 of the longshore current along PCA (Figure 8c). At low tide, the longshore current is wide and
 439 has two local maxima. The highest maxima is located seaward at $x = -450$ m and reaches 0.95
 440 m/s. At high tide, the main peak moves shoreward at $x = -250$ m and reaches 0.98 m/s. The mean
 441 seaward shear is around 0.0027 s^{-1} for both tide levels, which is less than half of the longshore
 442 current shear during the SandyDuck experiment. For each longshore current profile V , its potential
 443 vorticity V_x/h is shown and displays at least one local extremum which is a necessary condition
 444 for a shear instability to exist (Bowen and Holman 1989; Dodd et al. 1992). Because the longshore
 445 current peak is closer to the shoreline at high tide than at low tide, the location along the headland
 446 at which the longshore current is deflected offshore is also closer to the shoreline. This leads to a
 447 more concentrated, slightly more intense and narrower seaward flowing jet against the headland at
 448 high tide. Along T_H , the headland rip flow is around 50-100 m wider at low tide than at high tide
 449 (Figure 8d).

450 Wavenumber-frequency spectra of vorticity at different cross-shore positions in the surf zone
 451 and for both tide levels are shown in Figure 9. Similar to the SandyDuck experiment, the presence
 452 of SZE propagating in the same direction as the longshore current along PCA is ubiquitous. The
 453 approximative range of energetic frequencies is centred around 2 mHz (8 min) and decreases
 454 seaward with, for instance, much lower energy for $f > 5$ mHz at $x = -600$ m ($x = -370$ m) than at
 455 $x = -500$ m ($x = -270$ m) for low tide (high tide). By contrast, k-f spectra suggest that the range
 456 of energetic wavenumber remains relatively constant with cross-shore positions and is centred
 457 around 0.004 m^{-1} (250 m). The estimated celerity decreases seaward, going from $O(1)$ m/s near
 458 the longshore current peak to $O(0.1)$ m/s 500 m and 700 m offshore at high tide and low tide,
 459 respectively (see Figure 10). It should be noted that the estimation of the eddy celerity is relatively
 460 sensitive to the method to fit the straight dispersion line (Özkan-Haller and Kirby 1999). Such
 461 sensitivity is enhanced in Anglet which setup (morphology, high-energy wave conditions) is more
 462 complex than along the longshore-uniform sandy beach of Duck. This complexity, in addition to
 463 eddy interactions, may strongly increase discrepancies between the estimated eddy celerity and




FIG. 8. **(a)-(b)** Mean velocity field at low tide and at high tide. For both top panels, black line show two transects along which vorticity is plotted in other figures (T_C is the cross-shore transect of the longshore current and T_H is the cross-section transect of the deflection rip). Grey rectangle indicates area where $k-f$ spectrum are computed. **(c)** Cross-shore profile of mean longshore current V (black line) and potential vorticity associated with V (red line). **(d)** Cross-section profile of the mean deflection magnitude U_{rip} . For both bottom panels, plain (dashed) lines are for low tide (high tide).

the local mean longshore current. To increase the accuracy of the estimated eddy celerity, points (k, f_{rot}) with energy less than three orders of magnitude of the energy peak were removed from the fitting procedure. The resulting eddy celerity and the mean current have equivalent trends which is similar to SandyDuck.

Concluding this section, model results indicate the presence of SZE along PCA at both low tide and high tide. These eddies propagate in the direction of the longshore current and their celerity decreases seaward which is similar to SandyDuck (see Section 4). In the next section, the model is used to explore the relationship between such eddies and the VLF fluctuations of the deflection rip.




FIG. 9. Modeled wavenumber-frequency spectra of vorticity q ($E^q(k, f)$) at low tide (top) and high tide (bottom) extracted at different cross-shore positions x . For each panel, the local value of the mean longshore current V and the estimated eddy celerity C_{rot} are written. The definition of multiple dispersion lines is the same as in figure 3b.

b. Surf zone eddies and headland rip fluctuations

To investigate the hydrodynamic connections between the surf zone and the deflection rip, time-space and frequency-space diagrams of vorticity are computed along a transect in the surf zone T_C and against the headland T_H (see Figure 8a, b for transect location). These diagrams are shown in Figure 11 at low tide. Associated with those, Figure 12 shows several snapshots of vorticity which emphasise the length scales of vorticity fronts, their longshore advection, their merging and their offshore shedding through the deflection rip.

In the surf zone, two vorticity front pairs are present (Figure 12) and are associated with the two local maxima of longshore current at low tide (see Figure 8c). The seaward vorticity front pair is characterised by intense negative vorticity fronts that span the entire seaward longshore current region, going from around $x = -550$ m to $x = -870$ m. In this region, the frequency-space diagram




FIG. 10. **(a)** Eddy celerity C_{rot} extracted from vorticity versus local value of mean longshore current V . **(b)** Cross-shore profile of eddy celerities (points) and mean longshore current (line). Black and grey are for low tide and high tide, respectively.

along the cross-shore transect in the surf zone T_C (upper right-hand panel in Figure 11) highlights energetic frequencies that are similar to frequencies of SZE detected previously. Each cross-shore position is characterised by a relatively narrow range of energetic periods. Energetic periods of 5 to 8 min dominate the spectrum just seaward of the peak while periods of 30 min and 1 h dominate the spectrum further offshore.

In this locally alongshore-uniform surf zone situation, the most energetic period increases with the distance to the location of the longshore current peak (see upper right-hand panel in Figure 11). As suggested above, the most energetic longshore wavelength may remain relatively constant across the surf zone, indicating that the spatial structure of the corresponding eddies are conserved while their propagation speed decreases seaward. The latter mechanism can force the splitting of some vorticity fronts, leading to the generation of detached eddies that can propagate off the surf zone. This is better emphasized in the vorticity animation (see Supplementary Materials) and vorticity snapshots shown in Figure 12. Vorticity fronts span the entire surf zone and are advected by the longshore current. The most shoreward part of a front propagates faster than the seaward

part, which forces front stretching and detaching eddies. These eddies can flow off the surf zone or merge with the following front. Such mechanisms can also explained the differences between the representative eddy celerity and the mean longshore current (Figure 6).

Close to the headland, these fronts are expelled offshore through the deflection rip against the headland. Frequency-space diagrams along T_C and T_H are relatively similar, suggesting that fluctuations in the deflection rip are associated with the deflection of the upstream SZE. Along the cross-section of the rip neck T_H , energetic periods ranging from 5 min to 1 h dominate the spectrum. Far from the headland, only fluctuations with energetic periods of around 42 min dominate the spectrum at SIG1 location, which is similar with measurements (see Figure 2 and Mouragues et al. 2021).

Similar to low tide, the spatio-temporal (frequency) variability of vorticity at high tide is shown in Figure 13 and Figure 14. The frequency-space diagram of vorticity along T_C is relatively similar to the one at low tide, with energetic periods that essentially increase seaward. At high tide, the mean longshore current has also two local maxima which are associated with two vorticity front pairs in the surf zone (see around $x = -280$ m and $x = -500$ m). Just seaward of the main peak located at $x = -270$ m, intense negative vorticity fronts are advected along the stream with energetic periods of around 5 min. Further offshore, the longshore current profile displays a secondary bump (see around $x = -420$ m) and is associated with vorticity fronts propagating slower than the shoreward fronts, with energetic periods around 30 min.

The examination of vorticity animation at high tide (see Supplementary Materials) suggests that vorticity fronts merge at the location where the longshore current starts to be deflected offshore (region around $x = -400$ m and $y = 400$ m). The merged fronts are then expelled offshore through the deflection rip which leads to a much narrower range of energetic periods along T_H than along T_C , with distinct energetic periods of 23 min and 50 min along T_H (see bottom right-hand panel in Figure 13). The latter model result differs with the low tide simulation which highlighted the relatively same range of energetic periods along T_C and T_H .

This is believed to be controlled by the relative difference of mean flow patterns between low tide and high tide. At high tide, the mean flow against the headland is concentrated within a narrower region than at low tide (Figure 8). This allows all surf zone vorticity fronts to pass through the deflection rip at low tide while, at high tide, some surf zone fronts may merge together before




FIG. 11. Time-space diagram of 100-s running averaged vorticity ($q_{\text{mean}}(t)$; left-hand panels) and frequency-space diagram of instantaneous vorticity spectrum ($E^q(f)$; right-hand panels) along two transects at low tide (see Figure 8a for transect location). Top panels are for transect T_C with the black line indicating the mean longshore current peak position. Bottom panels are for transect T_H with the black (red) line indicating the mean deflection rip peak (SIG1) position.

being expelled offshore by the rip. This leads to energetic higher periods against the headland that were not necessarily present updrift in the surf zone (Figure 13). This mechanism will be further discussed in Section 6.




FIG. 12. Snapshots of vorticity at low tide at four different times showing stretched vorticity fronts, front interactions and offshore shedding through the deflection rip. Yellow point indicates SIG1 location and the green arrow in the first panel shows the main propagation direction of vorticity fronts. Dashed magenta rectangles show an example of two vorticity front pairs. For a better visualisation, the reader is referred to vorticity animations (see Supplementary Materials).

6. Discussion

In the following section, some components of the above model results are discussed and suggestions for future works are proposed. The morphological control on headland rip VLF fluctuations, through idealised morphology, is first explored. The latter will be critical to support the fact that the headland may enforce the merging of surf zone eddies as previously highlighted at PCA at high tide. Then, the role of wave group forcing and shear instability processes on surf zone rotational motions is discussed. In particular, we will point out the essential role of incident wave groups to produce surf zone eddies under high-energy oblique wave conditions. The latter will raise forward




FIG. 13. Time-space diagram of 100-s running averaged vorticity ($q_{\text{mean}}(t)$; left-hand panels) and frequency-space diagram of instantaneous vorticity spectrum ($E^q(f)$; right-hand panels) along two transects at high tide (see Figure 8b for transect location). Top panels are for transect T_C with the black line indicating the mean longshore current peak position. Bottom panels are for transect T_H with the black (red) line indicating the mean deflection rip peak (SIG1) position.

the discussion on a continuum of the driving mechanism of surf zone rotational motions under obliquely-incident waves.

a. Morphological control on headland rip VLF fluctuations

In section 5, the spatio-temporal variability of vorticity suggests that fluctuations of the rip are associated with the propagation of eddies along the deflection stream. For a given headland length, the reduction of the surf zone width can strongly modify the mean headland rip flow, with a more




FIG. 14. Snapshots of vorticity at high tide at four different times and showing stretched vorticity fronts, front interactions and offshore shedding through the deflection rip. Yellow point indicates SIG1 location and the green arrow in the first panel shows the main propagation direction of vorticity fronts. For a better visualisation, the reader is referred to vorticity animations (see Supplementary Materials).

concentrated seaward jet flow as surf zone width decreases. This forces the merging of vorticity fronts near the headland which results in energetic higher periods against the headland that were not necessarily present in the surf zone. To further analyse this mechanism, the model is run on an idealised morphology which is made of a planar surf zone morphology (constant slope of 0.03) and a physical boundary (headland) with a given length L_g . The incident wave conditions are the same for all modelling experiments ($H_s = 2$ m and $\theta = -20^\circ$). These conditions result in a 200-m wide longshore current (surf zone width $X_b \approx 200$ m) peaking around 0.8 m/s (Figure 15a). In order to obtain different mean flow patterns and further emphasize their effects on the energetic periods, L_g is varied so that the boundary length to surf zone width ratio L_g/X_b varies from 0.3 to 1.2. This allows to model the full spectrum of mean deflection patterns, going from weakly- to

strongly-deflected longshore current (Scott et al. 2016). With a constant X_b , the mean flow patterns feature different deflection rip widths (see left-hand panels in Figure 15). It should be noted that obtaining flow patterns with different surf zone and deflection rip widths could have also been done with *e.g.* varying tidal level (as previously done in Anglet; see Section 5) or varying offshore wave conditions (varying X_b). Here, we choose to vary L_g/X_b for consistency with Scott et al. (2016). These authors have varied this ratio to highlight the different deflection mean (time-averaged) flow patterns. The present model experiments aim at extending their work at the VLF scales.

For each experiment, the time-space (frequency) diagrams of vorticity along two transects (T_C and T_H) are presented (see middle and right-hand panels in Figure 15). Note that the same incident infragravity wave phases were used to run the model so that the variability of vorticity in the surf zone (transect T_C) is similar for all experiments. The latter vorticity variability is shown in panels b and c. The other middle and right-hand panels show vorticity along transect T_H for all experiments and highlight the effect of varying L_g (or L_g/X_b) and, in turn, the deflection rip width on the range of energetic periods in the deflection rip. As L_g/X_b increases, the deflection rip is compressed within a narrow region and its width is much shorter than the surf zone width (see left-hand panels). The range of energetic periods in the deflection rip strongly decreases compared to the range of energetic periods in the longshore current. For the longest boundary ($L_g/X_b = 1.2$), only periods higher than 7 min dominate the deflection rip spectrum (Figure 15r) while the longshore current features a wide range of energetic periods, going from 4 min to 50 min (Figure 15c).

Overall, these results underlines the previously-observed effect that an headland may have on the vorticity motions generated upstream in the surf zone. This is better emphasised by the continuous peak frequency $f_{p,vlf}$ computed as $\int f(E^q(f))^4 df / \int (E^q(f))^4 df$ (see right-hand panels). As L_g/X_b increases, the cut-off frequency ($f_{p,vlf}$ closest to the headland) decreases from 2.5 to 1.75 mHz for $L_g/X_b = 0.3$ and 1.2, respectively (Figure 15f and r, respectively). The peak frequency also strongly decreases with the distance from headland x_h . To further synthesise this mechanism, Figure 16 shows $T_{p,vlf} = 1/f_{p,vlf}$ at two different x_h as a function of L_g/X_b and for different H_s in order to highlight the effect of varying X_b . Both X_b feature similar period patterns. As L_g/X_b increases, periods at both positions and the period gap between both positions increase, which illustrates the effect of the headland on energetic periods.




FIG. 15. Left-hand panels : mean circulation patterns for different idealised boundary length. The boundary length to surf zone width ratio L_g/X_b is indicated for each panel ($X_b \approx 200$ m). Transects T_H and T_C are indicated in each panel. Middle panels : time-space diagram of running-averaged vorticity along T_H for different L_g/X_b . Right-hand panels : frequency-space diagram of instantaneous vorticity along T_H for different L_g/X_b . The upper middle (upper right-hand) panel shows the time (frequency)-space diagram of running-averaged (instantaneous) vorticity along T_C which is the same for each modelled case (same incident wave phase time series). Black lines on right-hand panels show the continuous peak frequency $f_{p,vlf}$ computed as $\int f(E^q(f))^4 df / \int (E^q(f))^4 df$.

b. Effects of wave-group forcing on vorticity dynamics

The effect of wave forcing, through breaking wave vorticity forcing, on the characteristics of surf zone rotational motions under obliquely incident wave conditions has been addressed by very few




FIG. 16. Very-low-frequency peak periods $T_{p,vlf}$ as a function of the boundary length to surf zone width ratio L_g/X_b . Yellow and red are for $H_s = 2$ m ($X_b = 200$ m) and for $H_s = 4$ m ($X_b = 400$ m). $T_{p,vlf}$ is shown at two different distances from the boundary x_h , corresponding to the distance of maximum vorticity variance (50 m and 65 m; coloured squares) and further away from the boundary (75 % of the deflection rip width of the narrowest deflection rip which is for $L_g/X_b = 1.2$; coloured circles). The size of squares and sizes is proportional to the vorticity variance, with large and small points associated with strongly- and weakly-fluctuating vorticity. For each H_s , the longshore current profile $V(x)$ is also shown.

studies (*e.g.* Long and Özkan-Haller 2009; Feddersen 2014). In the above modelling analysis, the term 'shear waves' was sometimes used, for simplicity, to refer to surf zone rotational motions in the presence of a longshore current V . This could mask the fact that shear instabilities of V are not the only driving mechanism of such motions. Under obliquely incident wave conditions, surf zone eddy generation mechanisms include shear instabilities of V (hereafter referred to as SI; *e.g.* Bowen and Holman 1989) and breaking wave vorticity forcing at the wave group scale (hereafter referred to as WG; *e.g.* Long and Özkan-Haller 2009) and at the individual wave scale through along-crest variation in wave dissipation (hereafter referred to as IW; *e.g.* Peregrine 1998; Feddersen 2014). Understanding the relative importance of each mechanism is essential to effectively predict surf zone rotational motions which strongly control mixing processes in the nearshore (*e.g.* Spydell et al. 2007; Clark et al. 2012).

A WG-resolving model allows simulating surf zone rotational motions driven by both WG and SI. Long and Özkan-Haller (2009) showed that the vorticity production due to WG was dominant

646 compared to SI under weakly-oblique incident wave conditions ($\theta = 8^\circ$), while production due to
 647 WG and SI was similar under strongly-oblique incident wave conditions ($\theta = 20^\circ$). In line with
 648 Long and Özkan-Haller (2009), excluding WG for the SandyDuck experiment (strong shear; Figure
 649 5a) resulted in weaker rotational motions while excluding WG for the Anglet experiment (weak
 650 shear; Figure 8c) resulted in the absence surf zone eddies (not shown).

651 This suggests the existence of a continuum in the driving mechanisms of surf zone rotational
 652 motions under obliquely incident waves, from fully WG-driven motions for weakly-sheared V to
 653 both WG- and SI-driven motions for V with stronger shear. Note that Feddersen (2014) used
 654 a wave-resolving model to investigate eddy generation due to all three mechanisms described
 655 above (SI, WG and IW). The existence of a continuum was suggested but was not fully illustrated.
 656 Importantly, this continuum would further point out the critical role of WG to produce surf zone
 657 eddies under high-energy and oblique wave conditions.

658 To illustrate such continuum, the model is run onto the barred beach of the SandyDuck experiment
 659 with varying wave angle of incidence θ . It should be noted that a barred beach was necessary,
 660 as opposed to a planar beach, to obtain a sheared-enough V so that eddies are generated when
 661 excluding WG. Simulations excluding and including WG are made for θ ranging $0-40^\circ$ leading to
 662 longshore currents with a wide range of peak magnitude V_{\max} (Figure 17a). The associated $k-f$
 663 vorticity spectra emphasise how wave group forcing broadens the spectrum (Figure 17b). Energetic
 664 frequencies and wavenumbers are concentrated within a very narrow ridge when excluding WG
 665 while the energy is much scattered when including WG. For normally-incident waves, surf zone
 666 eddies are absent when WG are excluded, emphasising the essential role of WG to produce eddies
 667 for weakly-oblique waves.

668 This is better illustrated by plotting the total vorticity variance as a function of θ ($(\sigma_{q,\text{rot}})^2$ and
 669 $(\sigma_{q,\text{sw}})^2$ including and excluding WG, respectively; Figure 17c). For $\theta < 30^\circ$, $(\sigma_{q,\text{rot}})^2$ is at least
 670 two times higher than $(\sigma_{q,\text{sw}})^2$, indicating that WG forcing is responsible for at least 50 % of
 671 vorticity variance. For θ between 0° and 25° , $(\sigma_{q,\text{rot}})^2$ increases until reaching a maximum at
 672 around $\theta = 25^\circ$. For $\theta > 25^\circ$, $(\sigma_{q,\text{rot}})^2$ decreases which highlights the decreasing contribution of
 673 WG to surf zone eddies. This decreasing trend may be due to the modification of the longshore
 674 structure of wave groups as a result of strong wave refraction, therefore modifying the longshore

variability of wave forcing and momentum injected in the surf zone. Further modeling works will be required to confirm the latter suggestion.

For θ between 0° and 25° , $(\sigma_{q,sw})^2$ linearly increases until reaching a plateau for $\theta > 25^\circ$, suggesting that the SI contribution reaches a saturation. Note that Figure 17b-d was plotted at a given cross-shore location $x = -200$ m but similar trends were obtained with other cross-shore locations. The SI contribution saturation may be explained by the fact that the local longshore current shear V_x and the local curvature V_{xx} , which controls the SI contribution to surf zone eddy field (see *e.g.* Long and Özkan-Haller 2009; Feddersen 2014), becomes relatively steady for strongly-oblique incident waves. Note that the presence of more energetic surf zone eddies when WG are included leads to more cross-shore mixing, compared to when WG are excluded, which smooths the longshore current profile (Figure 17a). Discrepancies between both variances decreases as θ increases, suggesting the decreasing contribution of WG, compared to SI, as waves becomes more oblique, which is in line with Long and Özkan-Haller (2009).

The relative balance between SI and WG contribution to surf zone eddies is well emphasised by plotting the vorticity variance ratio $((\sigma_{q,rot})^2 - (\sigma_{q,sw})^2) / (\sigma_{q,rot})^2$ as a function of the mean seaward shear τ (Figure 17d). The latter is defined as V_{max}/Δ , where Δ is the seaward width, computed from $V(x)$ including WG (see plain lines in Figure 17a). For the lowest τ ($\tau < 0.004 s^{-1}$), this ratio is always higher than 0.8, indicating that WG forcing is the main eddy generation mechanism. The mean seaward shear for the Anglet experiment (see blue text in Figure 17d) falls within the lowest range of τ , suggesting that surf zone vorticity motions and, in turn, VLF fluctuations of the deflection rip may be primarily driven by WG rather than SI. For higher τ , the SI contribution to surf zone eddies increases, with the variance ratio dropping below 0.6. This indicates that both SI and WG force surf zone eddies. Finally, for $\tau > 0.008 s^{-1}$ ($\theta \geq 35^\circ$), the ratio is close to 0, suggesting that WG do not substantially contribute to the total vorticity variance.

Overall, Figure 17d suggests the existence of a continuum in the surf zone eddy generation mechanisms, from fully WG-driven eddies for low τ to both WG- and SI-driven eddies for high τ . This is in general agreement with Feddersen (2014) who pointed out the fact that surf zone eddies are primarily controlled by breaking wave vorticity forcing, with possible exceptions for very narrow-banded highly-oblique wave conditions. Here, the total vorticity variance was used, as a first approximation, to quantify the relative importance of SI and WG contribution to surf zone




FIG. 17. Surf zone rotational motions modelled including and excluding wave group (WG) forcing for different angle of wave incidence θ ranging 0-40°. **(a)** Cross-shore profiles of mean longshore current $V(x)$. Vertical dashed line shows the cross-shore position at which quantities are plotted in other panels. **(b)** k - f vorticity spectra computed excluding and including WG for different θ . **(c)** Vorticity variance as a function of θ , excluding $(\sigma_{q,sw})^2$ and including $(\sigma_{q,rot})^2$ WG (circle and cross, respectively). **(d)** Vorticity variance ratio as a function of the mean seaward shear τ computed as V_{\max}/Δ where V_{\max} and Δ are the peak of $V(x)$ and the seaward width, respectively (see panel a). τ for the Anglet model experiments is shown. The dashed horizontal arrow emphasises the continuum of the surf zone rotational motions driving mechanisms which is discussed in the text (SI is for shear instability).

eddy generation. However, such bulk quantity does not account for the frequency and wavenumber spreading of the vorticity energy. As observations generally report strongly scattered k - f spectra (*e.g.* Özkan-Haller and Kirby 1999; Noyes et al. 2004), the inclusion of breaking wave vorticity

717 forcing is essential to accurately model the full range of surf zone eddies, even for strongly-sheared
718 current.

719 **7. Summary and future works**

720 A wave group-resolving model was implemented to investigate the driving mechanisms and the
721 spatio-temporal variability of VLF fluctuations of a deflection rip flowing against a 500-m rocky
722 headland located along Anglet beaches (SW, France). These energetic fluctuations were measured
723 800 m offshore during a 4-m oblique wave event and had dominant periods of around 1 h and 30
724 min. The model was first used to simulate surf zone eddies (SZE) in the presence of a longshore
725 current V at a longshore-uniform sandy beach under moderate wave conditions (SandyDuck). This
726 first modeling experiment ensured that the model was able to reproduce characteristics of measured
727 SZE propagating along a strongly-sheared current. The spectral signature and the spatio-temporal
728 variability of surf zone rotational motions, which included shear instability-driven SZE, were
729 displayed and qualitatively compared with past studies.

730 The model was then implemented in Anglet showing the presence of SZE propagating in the same
731 direction as the longshore current at both low tide and high tide. Due to energetic wave conditions,
732 the longshore current was relatively wide and weakly-sheared compared to the SandyDuck model-
733 ing experiment. SZE spanned the entire seaward width and propagated towards the headland at a
734 speed proportional to the local longshore current value. k - f spectra indicated that the celerity and
735 the range of energetic periods of these eddies were decreasing seaward. Space-frequency diagrams
736 of vorticity showed that spectral patterns in the surf zone and along the headland were relatively
737 similar, suggesting that VLF fluctuations of the rip are driven by the deflection of upstream SZE.

738 At low tide, most energetic periods increased with the distance from the headland, going from
739 $O(1)$ - $O(10)$ min very close to the headland to around 40 min to 1 h hundreds of meters away from
740 the headland which is line with measurements. At high tide, the range of most energetic periods in
741 the rip was much narrower than in the surf zone, suggesting that the headland enforces the merging
742 of surf zone eddies resulting in energetic higher periods against the headland. This mechanism
743 was further explored using idealised simulations with varying boundary length to surf zone width
744 ratio L_g/X_b . Increasing such ratio was shown to reduce the deflection rip width and to strongly
745 increase most energetic periods of vorticity fluctuations against the headland. These findings

746 have possible implications for sediment transport occurring along headlands. Most morphological
747 studies along idealised-isolated headlands or natural embayed beaches have shown the ability of
748 deflection rips, based on their time-averaged characteristics, to transport sediments offshore and
749 laterally (*e.g.* McCarroll et al. 2018; Valiente et al. 2020; McCarroll et al. 2021). Here, we show
750 that time-averaged deflection patterns may not be representative of its dynamics as a result of
751 potentially strong VLF fluctuations. These fluctuations may translate into fluctuations of sediment
752 flows which could impact the amount and the spatial dispersion of sediments transported by the
753 rip. Future modelling accounting for sediment transport and morphological changes will unravel
754 the deflection rip-induced sediment transport variability at the VLF scales. In addition to the
755 L_g/X_b -dependent deflection patterns, similar idealised simulations with varying headland spacing
756 L_s could be conducted to explore the distance required for longshore currents to develop SZE.

757 Lastly, the model was used to assess the effect of wave group forcing on surf zone eddies.
758 Excluding wave group forcing resulted in less energetic eddies for SandyDuck and in the absence
759 of eddies for Anglet, suggesting the existence of a continuum in the driving mechanisms of SZE
760 under obliquely incident waves. To illustrate this continuum, the SandyDuck setup simulation was
761 run with varying wave angle of incidence θ resulting in longshore currents with different mean
762 seaward shear. Ratio of the total vorticity variance indicates that wave group forcing accounts
763 for more than 50 % of the variance for θ lower than 30° . This suggests that SZE are primarily
764 controlled by breaking wave vorticity forcing rather than shear instabilities, except for strongly-
765 oblique wave conditions (strongly-sheared current). Interestingly, the contribution of wave groups
766 to the total variance reached a maximum and decreased for θ higher than 30° . Further works are
767 required to understand how wave groups generate eddies fluctuating at the VLF scale and how their
768 spatial structure impact the eddy variability. Finally, recent studies suggest that low-frequency
769 large-scale surf zone motions may be the result of a 2D turbulence inverse energy cascade, fed by
770 high-frequency short-scale vorticity motions due to along-crest energy dissipation (*e.g.* Feddersen
771 2014; Elgar and Raubenheimer 2020). Future works could address whether vorticity injected at
772 the wave group scale or at the individual wave scale due to the inverse cascade mechanism is more
773 dominant in generating surf zone eddies.

774 *Acknowledgments.* The Anglet field study has received financial support from Region Nouvelle-
775 Aquitaine (contract number 2017-1R20107) and was carried out as part of the project MEPELS
776 (contract number 18CP05), performed under the auspices of the DGA, and led by SHOM. Kévin
777 Martins greatly acknowledges the financial support from the University of Bordeaux, through an
778 International Postdoctoral Grant (Idex, 1024R-5030). We thank the two anonymous reviewers for
779 making very helpful comments and suggestions.

780 APPENDIX A


781 **Calibrated free parameters for SandyDuck and Anglet experiment**

Free parameter	Description	Duck	Anglet
$\gamma[-]$	Breaking parameter	0.3	0.5
$C[m^{1/2}.s^{-1}]$	Bottom friction Chezy parameter	55	45
$c_s[-]$	Horizontal mixing parameter	1	1

TABLE A1. Value of each calibrated free parameter for SandyDuck and Anglet experiment.

782 APPENDIX B

783 **Wavenumber-frequency spectra of velocities and vorticity during the SandyDuck** 784 **experiment**



785 FIG. B1. Modeled wavenumber-frequency spectra of cross-shore velocity u (a), longshore velocity v (b) and
 786 vorticity q (c) during the SandyDuck experiment, at the cross-shore position $x = 160$ m. The definition of
 787 multiple dispersion lines is the same as in figure 3b.

788 *Data availability statement.* Datasets for this research are included in Mouragues et al. (2020a).

789 **References**

790 Abadie, S., R. Butel, H. Dupuis, and C. Brière, 2005: Paramètres statistiques de la houle au
791 large de la côte sud-aquitaine. *Comptes Rendus Geoscience*, **337 (8)**, 769–776, [https://doi.org/](https://doi.org/10.1016/j.crte.2005.03.012)
792 10.1016/j.crte.2005.03.012.

793 Baquerizo, A., M. Caballeria, M. A. Losada, and A. Falqués, 2001: Frontshear and backshear
794 instabilities of the mean longshore current. *Journal of Geophysical Research: Oceans*, **106 (C8)**,
795 16 997–17 011, <https://doi.org/10.1029/2001JC900004>.

796 Bowen, A. J., and R. A. Holman, 1989: Shear instabilities of the mean longshore current 1. theory.
797 9.

798 Castelle, B., T. Scott, R. Brander, and R. McCarroll, 2016: Rip current types, circulation and
799 hazard. *Earth-Science Reviews*, **163**, 1–21, <https://doi.org/10.1016/j.earscirev.2016.09.008>.

800 Clark, D. B., S. Elgar, and B. Raubenheimer, 2012: Vorticity generation by short-crested
801 wave breaking. *Geophysical Research Letters*, **39 (24)**, 2012GL054 034, [https://doi.org/](https://doi.org/10.1029/2012GL054034)
802 10.1029/2012GL054034.

803 Clark, D. B., F. Feddersen, and R. T. Guza, 2010: Cross-shore surfzone tracer dispersion in an
804 alongshore current. *Journal of Geophysical Research: Oceans*, **115 (C10)**, 2009JC005 683,
805 <https://doi.org/10.1029/2009JC005683>.

806 Deigaard, R., E. D. Christensen, J. S. Damgaard, and J. Fredsøe, 1995: Numerical simulation
807 of finite amplitude shear waves and sediment transport. *Coastal Engineering 1994*, American
808 Society of Civil Engineers, Kobe, Japan, 1919–1933, [https://doi.org/10.1061/9780784400890](https://doi.org/10.1061/9780784400890.139).
809 139.

810 Dodd, N., V. Iranzo, and A. Reniers, 2000: Shear instabilities of wave-driven alongshore currents.
811 *Reviews of Geophysics*, **38 (4)**, 437–463, <https://doi.org/10.1029/1999RG000067>.

812 Dodd, N., J. Oltman-Shay, and E. Thornton, 1992: Shear instabilities in the longshore cur-
813 rent: A comparison of observation and theory. *Journal of Physical Oceanography*, 62–82,
814 [https://doi.org/10.1175/1520-0485\(1992\)022<0062:SIITLC>2.0.CO;2](https://doi.org/10.1175/1520-0485(1992)022<0062:SIITLC>2.0.CO;2).

- 815 Dodd, N., and E. B. Thornton, 1990: Growth and energetics of shear waves in the nearshore.
816 *Journal of Geophysical Research*, **95** (C9), 16 075, <https://doi.org/10.1029/JC095iC09p16075>.
- 817 Eckart, C., 1951: Surface waves in water of variable depth. *Wave Rep.* 100, 99.
- 818 Elgar, S., and B. Raubenheimer, 2020: Field evidence of inverse energy cascades in the surfzone.
819 *Journal of Physical Oceanography*, **50** (8), 2315–2321, <https://doi.org/10.1175/JPO-D-19-0327>.
820 1.
- 821 Falqués, A., and V. Iranzo, 1994: Numerical simulation of vorticity waves in the nearshore. *Journal*
822 *of Geophysical Research*, **99** (C1), 825, <https://doi.org/10.1029/93JC02214>.
- 823 Feddersen, F., 2014: The generation of surfzone eddies in a strong alongshore current. *Journal of*
824 *Physical Oceanography*, **44** (2), 600–617, <https://doi.org/10.1175/JPO-D-13-051.1>.
- 825 Herbers, T. H. C., S. Elgar, and R. T. Guza, 1994: Infragravity-frequency (0.005–0.05 hz) motions
826 on the shelf. part i: Forced waves. *JOURNAL OF PHYSICAL OCEANOGRAPHY*, 917–927.
- 827 Lippmann, T. C., T. H. C. Herbers, and E. B. Thornton, 1999: Gravity and shear wave contributions
828 to nearshore infragravity motions. *JOURNAL OF PHYSICAL OCEANOGRAPHY*, **29**, 9.
- 829 Long, C., 1996: Index and bulk parameters for frequency-direction spectra measured at cerc field
830 research facility, june 1994 to august 1995. *Misc. Pap. CERC-96-6*.
- 831 Long, J. W., and H. T. Özkan-Haller, 2009: Low-frequency characteristics of wave group–forced
832 vortices. *Journal of Geophysical Research*, **114** (C8), C08 004, [https://doi.org/10.1029/](https://doi.org/10.1029/2008JC004894)
833 [2008JC004894](https://doi.org/10.1029/2008JC004894).
- 834 Longuet-Higgins, M., and R. Stewart, 1964: Radiation stresses in water waves; a physical dis-
835 cussion, with applications. *Deep Sea Research and Oceanographic Abstracts*, **11** (4), 529–562,
836 [https://doi.org/10.1016/0011-7471\(64\)90001-4](https://doi.org/10.1016/0011-7471(64)90001-4).
- 837 MacMahan, J. H., A. J. H. M. Reniers, E. B. Thornton, and T. P. Stanton, 2004: Surf zone eddies
838 coupled with rip current morphology. *Journal of Geophysical Research: Oceans*, **109** (C7),
839 <https://doi.org/10.1029/2003JC002083>.

840 McCarroll, R., G. Masselink, N. Valiente, T. Scott, E. King, and D. Conley, 2018: Wave and
841 tidal controls on embayment circulation and headland bypassing for an exposed, macrotidal site.
842 *Journal of Marine Science and Engineering*, **6 (3)**, 94, <https://doi.org/10.3390/jmse6030094>.

843 McCarroll, R. J., G. Masselink, N. G. Valiente, E. V. King, T. Scott, C. Stokes, and M. Wiggins,
844 2021: An xbeach derived parametric expression for headland bypassing. *Coastal Engineering*,
845 **165**, 103 860, <https://doi.org/10.1016/j.coastaleng.2021.103860>.

846 Mouragues, A., P. Bonneton, B. Castelle, V. Marieu, R. J. McCarroll, I. Rodriguez-Padilla, T. Scott,
847 and D. Sous, 2020a: High-energy surf zone currents and headland rips at a geologically-
848 constrained mesotidal beach. *Journal of Geophysical Research: Oceans*, [https://doi.org/10.](https://doi.org/10.1029/2020JC016259)
849 [1029/2020JC016259](https://doi.org/10.1029/2020JC016259).

850 Mouragues, A., P. Bonneton, B. Castelle, and K. Martins, 2021: Headland rip modelling at a
851 natural beach under high-energy wave conditions. *Journal of Marine Science and Engineering*,
852 **9 (11)**, 1161, <https://doi.org/10.3390/jmse9111161>.

853 Mouragues, A., and Coauthors, 2020b: Field observations of wave-induced headland rips. *Journal*
854 *of Coastal Research*, **95 (sp1)**, 578, <https://doi.org/10.2112/SI95-113.1>.

855 Noyes, T. J., R. Guza, S. Elgar, and T. Herbers, 2004: Field observations of shear waves in
856 the surf zone. *Journal of Geophysical Research*, **109 (C1)**, C01 031, [https://doi.org/10.1029/](https://doi.org/10.1029/2002JC001761)
857 [2002JC001761](https://doi.org/10.1029/2002JC001761).

858 Noyes, T. J., R. Guza, F. Feddersen, S. Elgar, and T. Herbers, 2005: Model-data compar-
859 isons of shear waves in the nearshore. *Journal of Geophysical Research*, **110 (C5)**, C05 019,
860 <https://doi.org/10.1029/2004JC002541>.

861 Oltman-Shay, J., P. A. Howd, and W. A. Birkemeier, 1989: Shear instabilities of the mean
862 longshore current: 2. field observations. *Journal of Geophysical Research*, **94 (C12)**, 18 031,
863 <https://doi.org/10.1029/JC094iC12p18031>.

864 Özkan-Haller, H. T., and J. T. Kirby, 1999: Nonlinear evolution of shear instabilities of the
865 longshore current: A comparison of observations and computations. *Journal of Geophysical*
866 *Research: Oceans*, **104 (C11)**, 25 953–25 984, <https://doi.org/10.1029/1999JC900104>.

- 867 Peregrine, D., 1998: Surf zone currents. *Theoretical and Computational Fluid Dynamics*, **10** (1-4),
868 295–309, <https://doi.org/10.1007/s001620050065>.
- 869 Pezerat, M., X. Bertin, K. Martins, B. Mengual, and L. Hamm, 2021: Simulating storm waves
870 in the nearshore area using spectral model: Current issues and a pragmatic solution. *Ocean*
871 *Modelling*, **158**, 101 737, <https://doi.org/10.1016/j.ocemod.2020.101737>.
- 872 Reniers, A., J. MacMahan, E. Thornton, and T. Stanton, 2006: Modelling infragravity motions on a
873 rip-channel beach. *Coastal Engineering*, **53** (2-3), 209–222, [https://doi.org/10.1016/j.coastaleng.](https://doi.org/10.1016/j.coastaleng.2005.10.010)
874 2005.10.010.
- 875 Reniers, A. J. H. M., J. H. MacMahan, E. B. Thornton, and T. P. Stanton, 2007: Modeling of
876 very low frequency motions during ripex. *Journal of Geophysical Research*, **112** (C7), C07 013,
877 <https://doi.org/10.1029/2005JC003122>.
- 878 Roelvink, D., R. McCall, S. Mehvar, K. Nederhoff, and A. Dastgheib, 2018: Improving predic-
879 tions of swash dynamics in xbeach: The role of groupiness and incident-band runup. *Coastal*
880 *Engineering*, **134**, 103–123, <https://doi.org/10.1016/j.coastaleng.2017.07.004>.
- 881 Roelvink, D., A. Reniers, A. van Dongeren, J. van Thiel de Vries, R. McCall, and J. Lescinski,
882 2009: Modelling storm impacts on beaches, dunes and barrier islands. *Coastal Engineering*,
883 **56** (11-12), 1133–1152, <https://doi.org/10.1016/j.coastaleng.2009.08.006>.
- 884 Scott, T., M. Austin, G. Masselink, and P. Russell, 2016: Dynamics of rip currents associated
885 with groynes — field measurements, modelling and implications for beach safety. *Coastal*
886 *Engineering*, **107**, 53–69, <https://doi.org/10.1016/j.coastaleng.2015.09.013>.
- 887 Spydell, M., F. Feddersen, R. T. Guza, and W. E. Schmidt, 2007: Observing surf-zone dispersion
888 with drifters. *Journal of Physical Oceanography*, **37** (12), 2920–2939, [https://doi.org/10.1175/](https://doi.org/10.1175/2007JPO3580.1)
889 2007JPO3580.1.
- 890 Valiente, N. G., G. Masselink, R. J. McCarroll, T. Scott, D. Conley, and E. King, 2020: Nearshore
891 sediment pathways and potential sediment budgets in embayed settings over a multi-annual
892 timescale. *Marine Geology*, **427**, 106 270, <https://doi.org/10.1016/j.margeo.2020.106270>.