

Information throughput in healthy individuals who are comparable in age and gender with persons with stroke.

Tifenn Fauviaux, Germain Faity, Denis Mottet

▶ To cite this version:

Tifenn Fauviaux, Germain Faity, Denis Mottet. Information throughput in healthy individuals who are comparable in age and gender with persons with stroke. ACAPS 2021, Oct 2021, Montpellier, France. hal-03834663

HAL Id: hal-03834663 https://hal.science/hal-03834663v1

Submitted on 30 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tifenn FAUVIAUX¹, Germain FAITY¹, Denis MOTTET¹

1 - EuroMov Digital Health in Motion, Univ Montpellier, IMT Mines Alès, Montpellier, France EuroMov Digital Health in Motion University of Montpellier / IMT Mines Alès



Information throughput in healthy individuals who are comparable in age and gender with persons with stroke.



Stroke limits the capacity of the central nervous system to integrate sensory inputs for the execution of specific voluntary movement in response to task demands. Functional limitations are usually quantified with clinical scales, but it is unclear how clinical scores relate to the integration of sensory feedback into motor control loops.

> We use a circular steering task to quantify the effectiveness of continuous visuomotor feedbacks in goal-oriented behaviors.

Non dominant

hand

rest

Objective : Provide baseline values in a population of healthy individuals who are comparable in age and gender with persons with stroke.



19 Healthy volunteers were recruited into 2 groups: [>75 years old]; [≤75 years old].
Each participant had to follow a circular path as fast as possible with 2 to 6% of errors. They performed a total of 20 trials (20s each), organized into 2 blocks x 2 hands x 5 repetitions

Dominant hand

rest

THE CIRCULAR STEERING TASK :

We measured the Index of Effective Performance (IPe, in bit/s), which measure the effectiveness of visuomotor feedback.



IDe (bits) = $\frac{2\pi}{3}$ Standart Deviation of the radius

03

For the effectiveness of visuomotor feedbacks:

Hand side matters

trial

Non dominant

hand

rest

Age matters

Trajectory

rest

Gender does not matter





Dominant hand

rest



05 Conclusion

These preliminary results need to be supplemented with a larger number of participants to define age- and genderspecific norms for comparison with individuals with stroke.

The index of effective performance in a circular steering task quantifies the information throughput, which is a probe into the effectiveness of visuomotor feedbacks.

Reference values in healthy people will help quantifying the possibilities of improvement for each patient, to guide therapists in the rehabilitation strategy of motor control of the upper limbs after stroke.

References

Edwards, L.et al. Front. Integr. Neurosci. **2009**, 13,16 Fitts, P. M. Journal of Experimental Psychology, **1954**, 47(6), 381-391. MacKenzie. Human-computer interaction, **1992**, 7(1), 91-139.

Contact

tifenn.fauviaux@etu.umontpellier.fr germain.faity@umontpellier.fr denis.mottet@umontpellier.fr

