

Egg cooling associated with nest size in a passerine bird Marcel Lambrechts, Samuel Caro

▶ To cite this version:

Marcel Lambrechts, Samuel Caro. Egg cooling associated with nest size in a passerine bird. Journal of Thermal Biology, 2022, 110, pp.103383. 10.1016/j.jtherbio.2022.103383 . hal-03836351

HAL Id: hal-03836351 https://hal.science/hal-03836351

Submitted on 2 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Egg cooling associated with nest size in a passerine bird
Marcel M. Lambrechts* and Samuel P. Caro
CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
*Corresponding author: marcel.lambrechts@cefe.cnrs.fr. Address: CEFE-CNRS, 1919 route de
Mende, F-34293 Montpellier (cedex 5), France

18 **Abstract.** Breeders evolved adaptive responses to rapid changes in ambient temperature. In birds, nests are expected to reduce egg cooling when the incubator is temporarily off the eggs. Here we 19 present the results of two complementary laboratory experiments aiming at testing the association 20 21 between egg cooling and the thickness of the nest under and surrounding the eggs in a nondomesticated avian model species (great tit, Parus major). To simulate incubation behaviour, we 22 exposed nests with 4-egg clutches to a heat source until the eggs reached a normal incubation 23 temperature (ca. 39°C) and then recorded egg cooling eight minutes after removal of the heat 24 source, which corresponds to the time females generally leave eggs unattended during the 25 incubation period. Eggs cooled more quickly when the nest layer under the eggs was thinner and 26 when ambient temperature was cooler. We also show that the wall around the nest cup is important 27 to buffer egg cooling. It is hypothesised that in bird nests, both the thickness of the material under 28 the eggs, and the wall surrounding the nest cup interact to maintain a heat envelope around the eggs 29 for the time the incubating parent is foraging. This could explain why the thickness of the nest base 30 31 and wall are adjusted to the ambient temperature the birds experience during the nest building 32 phase, to anticipate the thermal conditions during incubation and preserve egg viability. 33 34 35 36 Key words great tit, *Parus major*, nest size, egg cooling, temperature

37

17

39 1. Introduction

Temperature is at the centre of studies that focus on biological consequences of climate change 40 (Parmesan and Yohe, 2003; White, 2008; IPCC Report, 2014; Deeming and Reynolds, 2015; 41 Gaughan et al. 2017). Temperature and its changes have a significant impact on how living beings 42 43 perform and breed (Visser et al., 2004; Cooper et al., 2005; Mainwaring et al., 2017; Andreasson et al., 2018; Marrot et al., 2018; Page et al., 2022). However, animal breeders evolved numerous 44 proximate response mechanisms to changes in the thermal environment so that embryos and 45 offspring develop in optimal breeding environments. At macro- and micro-geographic scales, they 46 can indeed adjust their breeding site; modulate their timing of reproduction and embryonic 47 development; they can alter litter or clutch sizes; or they can even pause their breeding behaviours 48 in case of extreme temperatures (Baker, 1938; Drent, 1975; Wingfield et al., 1998; Visser et al., 49 2004; Cooper et al., 2005; Caro et al., 2013; Deeming and Revnolds, 2015; Gladalski et al., 2018; 50

51 Kauffman et al., 2021; Sharpe et al., 2021).

52

53 In birds, larger nests with higher thermal insulation properties are often found in colder 54 breeding environments. One explanation is that these nests create a suitable environment for incubated eggs, the development of which entirely depends on the surrounding temperature (Drent, 55 56 1975; Deeming and Reynolds, 2015). For instance, bigger nests are expected to protect developing embryos against a decrease in egg temperature in cold environments (Gibb, 1950; Kluijver, 1950; 57 58 Drent, 1975; de Heij and Tinbergen, 2006; Deeming and Reynolds, 2015). In temperate zone passerines, the temperature of incubated eggs is usually between 30 and 40°C (Gibb, 1950; 59 60 Kluijver, 1950; Drent, 1975; Haftorn, 1981b, 1988). During incubation, the surface temperature of the nest cup (where the eggs sit) is >10°C higher than the temperature of the nest wall surrounding 61 62 the nest cup (Deeming and Pike, 2015; own observations). In addition, the temperature of the incubated eggs is often more than 10 to 20°C higher than the temperature above or outside the nest 63 (Gibb, 1950; Drent, 1975; Haftorn, 1981a,b; Haftorn 1988; own observations). When an incubating 64 parent leaves the nest, incubated eggs are suddenly exposed to a colder environment that induces 65 egg cooling (Gibb, 1950; Haftorn, 1981a,b, Haftorn 1988). Egg cooling might hamper embryo 66 development when eggs are maintained below 34 to 38°C for prolonged periods of time (Drent, 67 1975; Mainwaring, 2015; Webb, 1987). In addition, low egg temperatures increase the energy 68 needed to re-heat the eggs when the incubator returns to the nest (reviewed in Drent, 1975; 69 Monaghan and Nager, 1997). We therefore hypothesised that, especially in colder breeding 70 conditions, bigger or thicker nests are designed to store more incubation-generated heat and that this 71 heat, when released progressively, will slow down egg cooling in the absence of the incubating 72 parent. However, few intraspecific investigations quantified associations between dynamics in egg 73 temperature and aspects of nest design across a wide range of thermal conditions. 74

76 Here we report results of a laboratory study of egg cooling in association with nest thickness 77 under the eggs of a non-domesticated model species, the great tit (*Parus major*). We heated 4-egg 78 clutches placed inside nests to reach a normal incubation temperature (ca. 39°C) (Gibb, 1950; 79 Kluijver, 1950; Drent, 1975; Haftorn, 1981b; Haftorn, 1988), and then removed the heat source to monitor egg cooling for 8 minutes, which corresponds to a frequently observed off-bout duration in 80 natural conditions (Gibb, 1950; Kluijver, 1950; Haftorn, 1981a; Álvarez and Barba, 2014b; 81 Bambini et al., 2019). We performed those tests at various ambient temperatures, with nests of 82 different sizes, and by heating either entire nests, or nests in which the wall had been removed. 83 These different experimental conditions allowed us to test several factors that are expected to 84 influence egg cooling, including the role of the wall in the insulating properties of the nests. Based 85 on our findings on the basic thermal physics of nests, we provide a simple proximate explanation 86 for why colder environments might favour the construction of bigger and thicker nests. 87 88

89 2. Methods

90 The thermal properties of nests were investigated in laboratory conditions (cf. Lambrechts et al., 91 2020). We tested 101 nests collected in 2018. Those nests had contained nestlings that successfully 92 fledged. Using nests that had been deserted, either before egg hatching (17 nests) or during the stage of nestling rearing (3 nests), led to similar results (see supplementary material for more detail). 93 94 Nests had been constructed into four box types that have been described in Lambrechts et al. (2017). Briefly, some nests were constructed inside wood-concrete Schwegler B1 boxes (internal 95 diameter 10.5 cm, bottom area 113 cm², volume of the nest chamber under the entrance hole: 1525 96 cm³) whereas the other collected nests were constructed inside one of three wooden box types that 97 98 differed in size. Wooden boxes (made of European larch, Larix decidua) were either small (bottom area 36 cm², volume under the entrance hole 576 cm³), medium-sized (bottom area 121 cm², 99 100 volume under the entrance hole 1936 cm³) or large (bottom area 210 cm², volume under the 101 entrance hole 3780 cm³). Nests from different box types differed in size (Lambrechts et al., 2017; 102 Holveck et al., 2019), and therefore the use of different box types only aimed at increasing the range 103 of nest sizes. We also used eggs from deserted clutches in 2018 (cf. Lambrechts et al., 2020). Eggs 104 had different stages of development, and included eggs with yolk and eggs with embryos. Nests and 105 eggs were from nest boxes erected in eight urbanised plots situated in the city of Montpellier 106 (southern France) and one forest plot situated at ca. 25 km from Montpellier (Demeyrier et al., 107 2016; Lambrechts et al., 2016; 2017) (see table S1 for a description of the number of nests across 108 study plots and box types). Nests and eggs were transferred to the laboratory and kept at room 109 temperature before the start of the study.

110

111 We used the collected eggs to constitute three artificial 4-egg clutches, each clutch constituted of 112 eggs from the same nest of origin. We selected 4-egg clutches for the following reasons: 1) The size 113 of the incubation patch allows the female to simultaneously heat not more than four eggs via direct contact with the skin (Löhrl, 1986). Larger clutches lead to variation in surface temperature of 114 115 individual eggs, due to their position within the nest cup (Boulton and Cassey, 2012); 2) A clutch size of four eggs per nest facilitated the measurement of temperature under the eggs; 3) Using one 116 single clutch size allowed to remove effects of clutch size on egg cooling rates (e.g. reviewed in 117 Cooper et al., 2005; Nord and Nilsson, 2012; Deeming and Reynolds, 2015). General experimental 118 119 procedures consisted in placing nests, each containing one 4-egg clutch, under an infrared lamp 120 (250 W, Philips, at 38 cm above the nest bottom) for the time needed to reach a temperature observed during normal incubation (39 – 39.7°C, Haftorn, 1988; see also table 1 for a summary of 121 122 the heating duration needed across experimental conditions). Nests with eggs at incubation 123 temperature were then placed inside a closed insulated cabinet for eight minutes to monitor egg 124 cooling. By placing eggs in a closed cabinet, we could manipulate the ambient temperature and we 125 could also examine cooling in the absence of wind to simulate a nest-cavity environment 126 (Dickinson et al., 2019; Wachob, 1996). A fixed cooling period of eight minutes simulated the duration that an incubating female is off the eggs in natural conditions (Gibb, 1950; Kluijver, 1950; 127 128 Haftorn, 1981a; Álvarez and Barba, 2014b; Bambini et al., 2019). Both the ambient temperature 129 and the temperature under the eggs were simultaneously measured using an Otio thermometer with 130 two temperature sensors (Model: EM713P, Auterive, France, www.otio.com). Nests and 4-egg 131 clutches were used multiple times, under different thermal conditions (see below). Because thermic 132 properties differ between eggs and are associated with egg development (Cooper and Vos, 2013), 133 the artificial 4-egg clutches were switched between nests so that different nests received the same 134 clutches.

135

136 Both the nest characteristics and the thermal environment were manipulated in two 137 experiments (Table 1). In a first experiment we heated both entire nests and nests from which the 138 wall had been removed. Cut nests were obtained by vertically sawing the nest wall with a hacksaw, 139 only leaving the nest material under the nest cup (figure S1). This way, we obtained nests all having 140 a surface of approximately 7 cm x 7 cm (Table 1). Nests with an even smaller nest floor surface (36 141 cm^2) have been reported in our urban study site when great tits nested in a small box type 142 (Lambrechts et al., 2017). Intact and cut nests were heated, one at the time, with the eggs inside. 143 When the eggs reached a normal incubation temperature (39.0 - 39.7 °C) (Haftorn 1988), the nests 144 with eggs were placed for eight minutes inside an insulated cabinet that had an ambient temperature that differed within and between days of testing. Ambient temperature inside the cabinet ranged 145 between 11 and 27°C. Each nest (entire and cut) was tested more than once with different clutches 146

147 of eggs that were chosen randomly among the three available clutches of four eggs, to avoid

148 possible egg characteristic effects on the temperatures measured. Measurements were taken

149 between 6:00 and 22:30, with sampling duration and sampling hours differing between sampling

150 days (1 to 12 measurements of egg cooling per nest). A thermometer was always placed next to the

151 nest inside the cabinet, and this value was used as a continuous variable in the statistical tests (see

152 below). A total of 462 egg cooling measurements were performed for this experiment.

153

154 In a second experiment, we also exposed nests to two different heating treatments. This time 155 however, nests were left intact, and the treatments consisted in placing, or not, a pierced insulation

156 layer in between the nest and the infrared lamp. The insulation layer consisted in a 5cm-thick foam

157 layer covered with cardboard. It was pierced in the middle in a way that, when placed over the nest,

158 the lamp only heated the nest cup containing the eggs, and not the nest rim. Such a procedure better

159 mimics natural conditions, in which an incubator only sits on the eggs, and not on the whole nest.

160 Like in experiment 1, nests were heated until eggs reach a normal incubation temperature (39.0 –

161 39.6 °C), and ambient temperature differed within and between days of testing. Each nest was tested

162 in both conditions (with and without the insulation layer) and more than once, with different

163 clutches of eggs that were chosen randomly among three of the available clutches of four eggs. A

164 total of 96 measurements were performed for this experiment.

165

166 Statistics

To analyse the data, we used mixed model procedures (lmer, package lmerTest, Kuznetsova et al. 167 168 2016, in R version 3.5.0, R-Core-Team, 2015) always using the level of egg cooling (i.e. differences in egg temperature between the beginning and end of the eight-minute period of cooling) as the 169 170 dependent variable. We tested the effects of the treatment (entire vs. cut nests in exp. 1; with vs. without insulation layer in exp. 2), the ambient temperature during the cooling phase (in °C), nest 171 172 thickness under the eggs (in mm), and bottom surface area of the nests (in cm²) to account for the 173 size of the different box types from which nests had been collected. Nest identity and clutch identity 174 were added as crossed random intercepts.

175

176 **3. Results**

177 In experiment 1, the nest surface area did not influence egg cooling rate (p=0.122, table 2). Egg 178 cooling was however strongly associated with the ambient temperature, and the thickness of nest 179 material under the eggs (all p<0.001, table 2, figure 1 left panel), with eggs cooling more rapidly as 180 temperature and nest thickness were low. Eggs also cooled down more in nests without a wall, 181 compared to entire nests (treatment effect, p<0.001, figure 1 right panel), which suggests that the 182 wall itself contributes to the thermal protection of the eggs. Birds however do not cover the entire 183 nest when they incubate, but mainly the nest cup. Therefore, to better understand whether the nest 184 wall buffers egg cooling naturally, or whether the treatment effect observed here partly results from 185 the fact that we heated the entire nests, we designed experiment 2. In this second experiment, nests 186 were left intact and we heated the entire nest, or only its cup, using insulating material above the 187 nest rim. While the effects of ambient temperature and nest thickness observed in experiment 1 188 were confirmed in this second experiment (p < 0.001), we found no effect of the nest surface area 189 (p=0.207), and no effect of the treatment (p=0.273, table 2). This confirms that the buffering effect 190 of the nest wall on egg cooling observed in experiment 1 was intrinsic and not because the nest wall 191 had also been exposed to heat.

192

193 4. Discussion

Deeming (2016) proposed the concept of the 'bird-nest incubation unit'. In small passerines, like 194 Paridae, the incubator would require the nest surrounding the eggs to ensure that there is 195 196 maintenance of temperature. In this framework, thicker, larger or heavier nests would be more 197 efficient in maintaining egg temperature during the act of incubation to explain why larger nests are 198 found in colder environments (e.g. de Heij and Tinbergen, 2006; Crossman et al., 2011; Deeming et 199 al., 2012; Mainwaring et al., 2012, 2014; Heenan et al., 2015; but see Lambrechts and Caro, 2018). 200 Several techniques have been used to study the thermal insulation properties of passerine nests in 201 laboratory conditions, always in the absence of real eggs (e.g. Szentirmai et al., 2005; de Heij and 202 Tinbergen, 2006; Cruz et al., 2016; Akresh et al., 2017; Botero-Delgadillo et al., 2017; Gray and Deeming, 2017; Biddle et al. 2018; 2019; Deeming and Humphreys, 2020). The laboratory studies 203 204 focused on several nest-size characteristics, including nest length, nest width, nest depth, inner cup 205 diameter, inner cup depth and nest dry weight (e.g. de Heij and Tinbergen, 2006; Botero-Delgadillo 206 et al., 2017). Associations between aspects of nest morphology and nest cooling have also been reported in standardized laboratory conditions with relatively fixed ambient temperatures (e.g. Cruz 207 208 et al., 2016; Akresh et al., 2017; Botero-Delgadillo et al., 2017; but see de Heij and Tinbergen, 209 2006). Most of those studies however focused on the temperature of the nest itself, sometimes also 210 on dummy eggs, but not on the thermal dynamics of the eggs of the species that built the nest. 211

Here we hypothesized that larger nests might provide thermal benefits to the eggs once the incubating female leaves the nest. To test this hypothesis, we heated nests that contained a 4-egg clutch to monitor egg cooling after removal of the heat source. In our tests that simulated aspects of natural incubation behaviour (natural incubation temperature, time that the incubator is off the eggs during the egg cooling phase, dynamic environmental temperature within and across days), we observed that eggs cooled more quickly when the nest layer under the eggs was thinner and when the thermal conditions were cooler. A thicker nest layer under the eggs is thus able to store more 219 incubation-generated heat to be released for a longer period when the female is off the eggs. This is 220 a simple proximate explanation for why inside thicker nests, eggs cooled more slowly in our 221 different experimental set ups. That heated nests store heat to be released when suddenly exposed to 222 colder environmental conditions fits with our own observations that in thicker nests the temperature 223 under the eggs can continue to increase for up to several minutes after a heated nest is placed in 224 shade or cold conditions (M.M. Lambrechts, pers. obs.).

225

226 The important role played by the nest layers under the eggs does however not exclude the 227 possibility that the thickness of the nest wall might also contribute to the thermal insulation 228 properties of the nest, and this is what we showed with our experimental manipulation of the nest 229 wall. In experiment 1 we indeed showed that the presence of a nest wall is important to buffer egg 230 cooling, because eggs cooled down more quickly when the nest wall was removed. The presence of 231 a nest wall thus not only acts as a mechanical protection of the eggs against potentials shocks or to 232 hold eggs together in the centre of the nest, but it also maintains a heat envelope around the eggs, preventing them to cool down rapidly when the nest is not attended. In experiment 2 we further 233 234 show that this heat envelope exists and operates efficiently even if only the centre of the nest is 235 heated, like it is the case in natural conditions when the incubating bird only heats its eggs and the 236 nest layers immediately below them. The brood patch of the incubator is indeed substantially 237 smaller than the size of the whole nest. The surface of four great tit eggs touching the incubation 238 patch is *circa* 10 cm² (Löhrl, 1986), whereas the bottom surface of nests varied between *circa* 36 239 and 210 cm² in our study population. Altogether our experiments show that the nest is a complex 240 environment in which both the thickness of the material under the eggs, and the wall surrounding 241 the nest cup interact to maintain eggs at a viable range of temperatures for the time the incubating 242 parent leaves its nest for foraging and for other self-maintenance activities (see also Deeming et al, 243 2020).

244

245 Based on our laboratory results on egg cooling, we would expect that bigger or thicker nests 246 are built in colder environmental conditions, e.g. earlier in the season, at higher elevations or at higher latitudes (Briggs et al., 2019; Deeming et al., 2012). Associations between aspects of nest 247 248 size and ambient temperature have been investigated in field studies (e.g. Deeming and Reynolds, 249 2015 for a review) and in laboratory conditions (Lambrechts and Caro, 2018). Consistent with these 250 expectations, we found in previous field studies that great tits invested more in nest building when 251 nests were constructed earlier in the season (Lambrechts et al., 2017). We also found that blue tits 252 built smaller replacement nests built later in the season than first nests built earlier in the same breeding season (Lambrechts et al., 2012) (see also Deeming and Mainwaring, 2015). However, in 253 an 18-year field study that only focused on nests from first breeding attempts built in one box type 254

255 (Schwegler), we did not find a significant association between nest volume and the average ambient temperature during the week preceding the onset of egg laying (Lambrechts et al., 2016). This 256 257 would imply that our long-term field measure of nest size (i.e. height of the nest rim at the entrance 258 door, ignoring the nest cup properties) does not necessarily reflect the functional thermic properties 259 of a nest (see also Deeming and Mainwaring, 2015; Cruz et al., 2016). Field studies interested in thermic properties of nests influencing changes in egg temperature therefore should also take other 260 261 aspects of nest size into account, such as aspects associated with nest-cup characteristics (e.g. thickness or composition of the material under the eggs) (e.g. de Heij and Tinbergen, 2006; 262 Deeming et al., 2012; 2020; Mainwaring et al., 2012; Deeming and Mainwaring, 2015). In addition, 263 264 in a previous laboratory study in great and blue tits (Lambrechts and Caro, 2018), we did not find an association between ambient temperature and nest volume and nest weight. This was a bit 265 266 surprising given that nest weight and volumes probably better reflects nest thickness under the eggs. 267 which we measure here, than the sole measure of the external nest wall as done in the field study. 268 But we also observed that the captive birds only started to build when photoperiod was very long. We therefore do not exclude that a very long photoperiod might change the sensitivity to ambient 269 270 temperature in breeders (e.g. Lambrechts and Perret, 2000) therefore perhaps also influencing 271 investment in nest building.

272

Future studies could focus on which egg characteristics proximately determine their thermal properties and whether egg cooling might also be associated with aspects of nest morphology other than mass or thickness, such as nest composition (Deeming et al., 2020). Egg warming following egg cooling could also be investigated with similar approaches to investigate whether low egg temperatures might increase a proximate cost of egg warming when the female returns to the cooled eggs.

279

280 Acknowledgements

We thank all the staff members and students that kindly helped with nest collection and fieldmonitoring.

283

284 Conflicts of interest

285 The authors declare that they have no conflicts of interest.

286

288

289 References

Akresh, M. E., Ardia, D. R., King, D. I., 2017. Effect of nest characteristics on thermal properties,
clutch size, and reproductive performance for an open-cup nesting songbird. Avian Biol.

292 Res. 10, 107-118.

- Álvarez, E., Barba, E., 2014a. Behavioural responses of great tits to experimental manipulation of
 nest temperatures during incubation. Ornis Fennica 91, 220-230.
- Álvarez, E., Barba, E., 2014b. Within and between population variations of incubation rhythm of
 great tits *Parus major*. Behaviour 151, 1827-1845.
- Andreasson, F., Nord, A., Nilsson, J.-A., 2018. Experimentally increased nest temperature affects
 body temperature, growth and apparent survival in blue tit nestlings. J. Avian Biol. e01620
 doi: 10.1111/jav.01620
- 300 Baker, J.R., 1938. The evolution of breeding seasons. In: Evolution: essays on aspects of
- evolutionary biology (G.R.De Beer, ed.), pp. 161–177. Oxford University Press, London,
 UK.
- Bambini, G., Schlicht, E., Kempenaers, B., 2019. Patterns of female nest attendance and male
 feeding throughout the incubation period in blue tits *Cyanistes caeruleus*. Ibis 161, 50-65.
- Biddle, L. E., Goodman, A. M., Deeming, D. C., 2018. Infrared thermography provides insight into
 the thermal properties of bird nests. J. Therm. Biol. 76, 95-100.
- Biddle, L.E., Goodman, A.M., Deeming, D.C., 2019. Wall structure affects insulation of nests built
 by thrushes. Acta Ornithol. 54, 255-262.
- Botero-Delgadillo, E., Orellana, N., Serrano, D., Poblete, Y., Vásquez, R. A., 2017. Interpopulation
 variation in nest architecture in a secondary cavity-nesting bird suggests site-specific
 strategies to cope with heat loss and humidity. Auk 134, 281-294.
- 312 Boulton, R.L., Cassey, P., 2012. How avian incubation behaviour influences egg surface
- temperatures: relationships with egg position, development and clutch size. J. Avian Biol.
 43, 289-296.
- Briggs, K.B., Biddle, L.E., Deeming, D.C., 2019. Geographical location affects size and materials
 used in the construction of European Pied Flycatcher (Ficedula hypoleuca) nests. Avian
 Research 10, 15.
- Caro SP, Schaper SV, Hut RA, Ball GF, Visser ME, 2013. The case of the missing mechanism: How
 does temperature influence seasonal timing in endotherms? PLoS Biol 11:e1001517.
- 320 Cooper C. B., Hochachka W. M., Butcher G., Dhondt A. A. 2005. Seasonal and latitudinal trends in
- 321 clutch size: thermal constraints during laying and incubation. Ecology 86: 2018-2031
- Cooper, C. B., Voss, M. A., 2013. Avian incubation patterns reflect temporal changes in developing
 clutches. PLoS ONE 8(6), e65521, doi:10.1371/ journal.pone.0065521

- 324 Crossman, C. A., Rohwer, V. G., Martin, P. R., 2011. Variation in the Structure of Bird Nests
- between Northern Manitoba and Southeastern Ontario. PLoS ONE 6(4), e19086,
 doi:10.1371/journal.pone.0019086
- 327 Cruz, Á, Álvarez, A., Barba, E., 2016. Nest insulation capacity during incubation and after fledgling
 328 are related. Avian Biol. Res. 9, 22-27.
- de Heij, M. E., Tinbergen, J. M., 2006. Great tits with thin nests delay the onset of full incubation.
 Chapter 5 in de Heij M. E., PhD Thesis, University of Groningen
- 331 Deeming, D. C., 2016, How does the bird-nest incubation unit work? Avian Biol. Res., 103-113.
- 332 Deeming, D.C., Humphreys, E., 2020. Insulation of Common Chaffinch Fringilla coelebs nests is
 333 largely driven by animal-derived materials in the cup lining. Acta Ornithol. 55, 250-260.
- 334 Deeming, D.C., Pike, T.W., 2015. Nest surface temperature predicts fledging success of Blue Tits
 335 *Cvanistes caeruleus* but not Great Tits *Parus major*. Acta Ornithol. 50, 247-251.
- 336 Deeming, D. C., Reynolds, S. J., 2015 (eds.). Nests, eggs, and incubation: new ideas about avian

reproduction. Oxford University Press, Oxford.

344

- Deeming, D. C., Mainwaring, M. C., 2015. Functional properties of nests. In: Deeming, D.C. &
 Reynolds, S.J. (eds), *Nests, eggs, and incubation: new ideas about avian reproduction*, pp.
 29–49. Oxford University Press, Oxford.
- 341 Deeming, D. C., Mainwaring, M. C., Hartley, I. R., Reynolds, S. J., 2012. Local temperature and
- not latitude determines the design of blue tit and great tit nests. Avian Biol. Res. 5, 203-208.
- 343 Deeming, D.C., Dickinson, A.M., Broughton, R.E., Locke, E., Gray, L.A., Bennett, S.L., Gilchrist,
- 345 songbird nests as measured using temperature loggers. Physiol. Biochem. Zool. 93, 488-504.

R., Muniz, S., Goodman, A.M., Biddle, L.E., 2020. Factors affecting thermal insulation of

- 346 Demeyrier, V., Lambrechts, M.M., Perret, P., Grégoire, A., 2016. Experimental demonstration of an
- 347 ecological trap for a wild bird in a man-transformed environment. Anim. Behav. 118, 181-190.
- 348 Dickinson, A.M., Goodman, A.M., Deeming, D.C., 2019. Air movement affects insulatory values of
 and a nests constructed by Old World Warblers. J. Thermal Biol. 81, 194-200.
- 350 Drent, R. 1975. Incubation. In: Farner DS and King JR (eds), *Avian Biology*, pp. 333-420. Academic
 351 Press, New York
- Gaughan, J. B., Lees, A. M., Sejian, V., 2017. Sixty years of animal biometeorology. Int. J. Biometeol.
 https://doi.org/10.1007/s00484-017-1459-1
- 354 Gibb, J. 1950. The breeding biology of the great and blue titmice. Ibis 92, 507-539.
- 355 Glądalski, M., Bańbura, M., Kaliński, A., Markowski, M., Skwarska, J., Wawrzyniak, J., Zieliński,
- Z., Bańbura, J., 2018. Hatching delays in great tits and blue tits in response to an extreme cold
 spell: a long-term study. Int. J. Biometeol. https://doi.org/10.1007/s00484-018-1541-3
- 358 Gray, L. A., Deeming, D. C., 2017. Effects of air movement on the thermal insulation of avian nests.
- 359 Bird Study https://doi.org/10.1080/00063657.2017.1387518

Gwinner, H., Capilla-Lasheras, P., Cooper, C., Helm, B., 2018. 'Green incubation': avian offspring
benefit from aromatic nest herbs through improved parental incubation behaviour. Proc. R.

362 Soc. Lond. B 285, DOI: 10.1098/rspb.2018.0376

- 363 Haftorn, S., 1981a. Incubation rhythm in the great tit *Parus major*. Fauna norv. Ser. C. Cinclus 4, 9364 26.
- 365 Haftorn, S., 1981b. Incubation during the egg-laying period in relation to clutch-size and other aspects
 366 of reproduction in the great tit *Parus major*. Ornis Scand. 12, 169-185.
- 367 Haftorn S., 1988. Incubating female passerines do not let the egg temperature fall below the
 368 'physiological zero temperature' during their absences from the nest. Ornis Scand. 19, 97-110.
- 369 Harrison, C., 1982. An Atlas of the Birds of the Western Palaearctic. Collins, London, 322 pp.

Heenan, C. B., Goodman, B. A., White, C. R., 2015. The influence of climate on avian nest
construction across large geographic gradients. Global Ecol. Biogeogr. 24, 1203-1211.

372 Holveck, M.-J., Grégoire, A., Doutrelant, C., Lambrechts, M. M. 2019. Nest height is affected by

- lampost lighting proximity in addition to bestbox size in urban great tits. J Avian Biol 50,e01798.
- 375 IPCC Report on Climate Change, 2014. Synthesis Report. Contribution of Working Groups I, II and
 376 III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core
 377 Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151 pp.
- 378 Kauffman, K.L., Elmore, R.D., Davis, C.A., Fuhlendorf, S.D., Goodman, L.E., Hagen, C.A., Tanner,
- E.P., 2021. Role of the thermal environment in scaled quail (*Callipepla squamata*) nest site
 selection and survival. J. Thermal Biol. 95, 12.
- 381 Kluijver, H. N., 1950. Daily routines of the great tit, Parus m. major L. Ardea 38, 99-135.
- 382 Kuznetsova A, Brockhoff PB, Christensen RHB. 2016. ImerTest: Tests in Linear Mixed Effects
 383 Models. R package version 2.0-30.
- 384 Lambrechts, M. M., Aimé, C., Midamegbe, A., Galan, M.-J., Perret, P., Grégoire, A., Doutrelant,
- C., 2012. Nest size and breeding success in first and replacement clutches: an experimental
 study in blue tits *Cyanistes caeruleus*. J. Ornithol. 153, 173-179.
- 387 Lambrechts, M. M., Blondel, J., Bernard, C., Caro, S. P., Charmantier, A., Demeyrier, V.,
- 388 Doutrelant, C., Dubuc-Messier, G., Fargevieille, A., de Franceschi, C., Giovannini, P.,
- 389 Grégoire, A., Hurtrez-Boussès, S., Lucas, A., Mainwaring, M. C., Marrot, P., Mennerat, A.,
- 390 Perret, S. & Perret, P., 2016. Exploring biotic and abiotic determinants of nest size in
- 391 Mediterranean great tits (*Parus major*) and blue tits (*Cyanistes caeruleus*). Ethology 122,
- *492-501.*
- 393 Lambrechts, M. M., Caizergues, A. E., Perrier C., Charmantier, A. & Caro, S. P., 2020. Surface
- 394 temperatures of non-incubated eggs in great tits (*Parus major*) are strongly associated with
- ambient temperature. Int. J. Biometeorol., https://doi.org/10.1007/s00484-020-01958-1

- Lambrechts, M. M., Caro, S. P., 2018. Experimental manipulation of photoperiod and temperature
 do not influence nest size in blue and great tits. Auk 135, 218-227.
- Lambrechts, M. M., Charmantier, A., Demeyrier, V., Lucas, A., Perret, S., Abouladzé, M., Bonnet,
 M., Canonne, C., Faucon, V., Grosset, S., le Prado, G., Lidon, F., Noell, T., Pagano, P.,
- 400 Perret, V., Pouplard, S., Spitaliéry, R., Bernard, C., Perret, P., Blondel, J., Grégoire A., 2017.
- 401 Nest design in a changing world: great tit *Parus major* nests from a Mediterranean city
- 402 environment as a case study. Urban Ecosyst. 20, 1181-1190.
- 403 Lambrechts, M. M., Perret, P., 2000. A long photoperiod overrides non-photoperiodic factors in
 404 Blue Tits' timing of reproduction. Proc. Roy. Soc. Lond. B 267, 585-588.
- 405 Löhrl, H., 1986. Experimente zur Bruthöhlenwahl der Kohlmeise (*Parus major*). J. Orn. 127, S. 51406 59.
- 407 Mainwaring, M. C., 2015. Nest construction and incubation in a changing climate. In: Deeming,
- 408 D.C. & Reynolds, S.J. (eds), *Nests, eggs, and incubation: new ideas about avian*409 *reproduction*, pp. 65–74. Oxford University Press, Oxford.
- 410 Mainwaring, M. C., Barber, I., Deeming, D. C., Pike, D. A., Roznik, E. A., Hartley, I. R., 2017.
- 411 Climate change and nesting behaviour in vertebrates: a review of the ecological threats and
 412 potential for adaptive responses. Biol. Rev. 92, 1991-2002.
- 413 Mainwaring, M. C., Hartley, I.R., Bearhop, S., Brulez, K., du Feu, C. R., Murphy, G., Plummer, K.
- 414 E., Webber, S. L., Reynolds, S. J., Deeming, D. C. 2012. Latitudinal variation in blue tit and 415 great tit nest characteristics indicates environmental adjustment. J. Biogeogr. 39, 1669-1677.
- 416 Mainwaring, M. C., Hartley, I. R., Lambrechts, M. M., Deeming, D. C., 2014 The design and
- 417 function of birds' nests. Ecol. Evol. 20, 3909-3928.
- 418 Marrot, P., Charmantier, A., Blondel, J., Garant, D., 2018. Current spring warming as a driver of
 419 selection on reproductive timing in a wild passerine. J. Anim. Ecol. 87, 754-764.
- 420 Monaghan, P., Nager, R. G., 1997. Why don't birds lay more eggs? TREE 12, 270-273.
- 421 Nord, A., Nilsson, J.-A., 2012. Context-dependent costs of incubation in the pied flycatcher. Anim.
 422 Behav. 84, 427-436.
- 423 Page, J.L., Nord, A., Dominoni, D.M., McCafferty, D.J., 2022. Experimental warming during
- 424 incubation improves cold tolerance of blue tit (*Cyanistes caeruleus*) chicks. J. Exp. Biol.
 425 225, jeb243933.
- 426 Parmesan, C., Yohe, G., 2003. A globally coherent fingerprint of climate change impacts across
 427 natural systems. Nature 421 (6918), 37-42.
- 428 R-Core-Team., 2015. R: A language and environment for statistical computing.: R Foundation for
- 429 Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL
- 430 http://www.R-project.org/.

- 431 Sharpe, L.L., Bayter, C., Gardner, J.L., 2021. Too hot to handle? Behavioural plasticity during
 432 incubation in a small, Australian passerine. J. Thermal Biol. 98, 11.
- 433 Slagsvold, T., 1989. Experiments on clutch size and nest size in passerine birds. Oecologia 80, 297434 302.
- 435 Szentirmai, I., Székely, T., Liker, A., 2005. The influence of nest size on heat loss of penduline tit
 436 eggs. Acta Zool. Acad. Sci. Hung. 51, 59-66.
- Visser, M. E., Both, C., Lambrechts, M. M., 2004. Global climate change leads to mistimed avian
 reproduction. Adv Ecol. Res. 35, 89-110.
- Wachob, D. G., 1996. A microclimate analysis of nest-site selection by mountain chickadees. J.
 Field Ornithol. 67, 523-533.
- 441 Webb, D.R., 1987. Thermal tolerance of avian embryos: a review. The Condor 89, 874-898.
- 442 Wingfield J. C., Breuner C. W., Jacobs J. D., Lynn S., Maney D. L., Ramenofsky M., Richardson R.
- 443 D. 1998. Ecological bases of hormone-behavior interactions: The "emergency life history
- 444 stage". Am. Zool. 38, 191-206.
- White T. C. R. 2008. The role of food, weather and climate in limiting the abundance of animals.Biol. Rev. 83, 227-248.

Table 1: Laboratory tests used to study associations between nest-size characteristics and cooling of 4-egg clutches (i.e. difference in egg temperature between the beginning and end of a fixed eightminute period after heating). Experiment 1: Heating of entire nests *versus* heating of nests of which the nest wall was removed. Experiment 2: Heating of entire nests versus heating of only the nest cups of entire nests. Average values and their associated standard deviations, as well as the entire ranges of values for each of the measurements taken, are shown. Individual nests were tested multiple times varying temperature conditions across measurements.

- 456
- 457 458

	Experi	Experiment 1		ment 2
	Entire nest heated	Nests without wall heated	Entire nest heated	Nest cup heated
Sample size	336 measurements of egg cooling	126 measurements of egg cooling	48 measurements of egg cooling	48 measurements of egg cooling
Nest mass (in g)	23.4 <u>+</u> 11.1 4.7 – 61.9	$6.3 \pm 2.8 \\ 1 - 13$	23.9 ± 13.0 8.0 - 61.9	23.9 <u>+</u> 13.0 8.0 - 61.9
Nest thickness under eggs (in mm)	21.5 ± 6.9 10.0 - 44.0	20.9 ± 6.7 10.0 - 44.0	22.3 <u>+</u> 7.6 11.4 – 39.0	22.3 <u>+</u> 7.6 11.4 – 39.0
Duration of egg heating prior to cooling (in seconds)	717 <u>+</u> 266 324 – 3076	855 <u>+</u> 303 418 – 2292	668 <u>+</u> 184 324 – 1052	643 <u>+</u> 148 377 – 995
Temperature under heated eggs at the onset of egg cooling (°C)	39.1 <u>+</u> 0.1 39.0 – 39.7	39.1 <u>+</u> 0.1 39.0 – 39.5	39.1 <u>+</u> 0.1 39.0 – 39.6	39.3 <u>+</u> 0.2 39.0 – 39.6
Egg cooling (°C)	7.0 <u>+</u> 2.5 2.8 – 15.5	9.8 <u>+</u> 3.4 4.9 – 18.7	6.5 ± 1.5 3.5 - 9.9	6.7 <u>+</u> 1.6 3.4 – 9.9
Air temperature during egg cooling (°C)	19.9 ± 5.4 12.0 - 26.9	20.5 ± 5.4 11.0 - 26.5	24.2 <u>+</u> 1.5 21.3 – 27.1	24.3 ± 1.4 21.2 - 26.3
Measure day (Julian dates)	170 – 246 43 days	175 – 279 26 days	237 – 246 20 days	237 – 246 20 days

459

460

461 462

463

466 Table 2: Analyses of the variables that influence the level of egg cooling (i.e. difference in egg 467 temperature between the beginning and end of a fixed eight-minute period after heating). In experi-468 ment 1 (n=462 measures), treatment consisted in heating entire nests or nests from which the wall 469 had been removed (intercept depicts cut nests, coded as 1 in the model). In experiment 2 (n=96 470 measures), treatment consisted in placing or removing a pierced insulation layer above the nest rim 471 in order to restrict heating to the nest cup (intercept is without the insulation layer, coded as 1 in the 472 model).

473

	Variable	Estimate	S.E.	F	p-value
Experiment 1					
	(Intercept)	22.74	0.74		
	Treatment	-2.99	0.15	410.42	< 0.001
	Ambient temperature	-0.39	0.01	1019.03	< 0.001
	Nest thickness under eggs	-0.07	0.01	21.09	< 0.001
	Nest surface area	-0.00	0.00	2.44	0.122
		-0.00	0.00		
Experiment 2		-0.00			
Experiment 2	(Intercept)	21.82	1.54		
Experiment 2	(Intercept) Treatment	21.82	1.54 0.09	1.21	0.276
Experiment 2	(Intercept) Treatment Ambient temperature	21.82 0.10 -0.59	1.54 0.09 0.05	1.21 118.92	0.276 <0.001
Experiment 2	(Intercept) Treatment Ambient temperature Nest thickness under eggs	21.82 0.10 -0.59 -0.08	1.54 0.09 0.05 0.02	1.21 118.92 13.62	0.276 <0.001 <0.001

474

Figure 1: Influence of the thickness of nest material under the eggs (left panel, p<0.001, the line depicts the mixed-model output), and of the removal of the nest wall (right panel, p<0.001) on egg cooling in experiment 1. Eggs cool down more slowly when there is more material under the nest cup, and when the nest wall is present.



479