



Secondary structure and ^1H , ^{15}N & ^{13}C resonance assignments of the periplasmic domain of OutG, major pseudopilin from *Dickeya dadantii* type II secretion system

Theis Jacobsen, Régine Dazzoni, Melvin Renault, Benjamin Bardiaux, Michael Nilges, Vladimir Shevchik, Nadia Izadi-Pruneyre

► To cite this version:

Theis Jacobsen, Régine Dazzoni, Melvin Renault, Benjamin Bardiaux, Michael Nilges, et al.. Secondary structure and ^1H , ^{15}N & ^{13}C resonance assignments of the periplasmic domain of OutG, major pseudopilin from *Dickeya dadantii* type II secretion system. *Biomolecular NMR Assignments*, 2022, 16 (2), pp.231-236. 10.1007/s12104-022-10085-4 . hal-03836613

HAL Id: hal-03836613

<https://hal.science/hal-03836613>

Submitted on 16 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Secondary structure and ^1H , ^{15}N & ^{13}C resonance assignment of the periplasmic domain of OutG, major pseudopilin from *Dickeya dadantii* type II secretion system

Theis Jacobsen¹, Régine Dazzoni¹, Melvin Renault^{2,3}, Benjamin Bardiaux¹, Michael Nilges¹, Vladimir Shevchik^{2,3}, Nadia Izadi-Pruneyre^{1*}

1: Institut Pasteur, Université de Paris, CNRS UMR3528, Structural Bioinformatics Unit, F-75015 Paris, France 2: Université de Lyon 1, INSA-Lyon, F-69621 Villeurbanne, France. 3: CNRS, UMR5240, Microbiologie Adaptation et Pathogénie, F-69622 Lyon, France.

* corresponding author: nadia.izadi@pasteur.fr

Declarations

Keywords: NMR resonance assignment, Type II Secretion System, OutG, *Dickeya dadantii*, pseudopilin

Funding

This work was funded by the Institut Pasteur, the Centre National de la Recherche Scientifique (CNRS), the French Agence Nationale de la Recherche (ANR Synergy-T2SS ANR-19-CE11-0020-01) and European Union's Horizon 2020 Research and Innovation program under the Marie Skłodowska-Curie Grant agreement no 765042. The 800-MHz NMR spectrometer of the Institut Pasteur was partially funded by the Région Ile de France (SESAME 2014 NMRCHR grant no 4014526).

Conflicts of interest/Competing interests

The authors declare they have no conflict of interest.

Availability of data and material (data transparency)

The chemical shift values have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu/>) under the accession number 51296.

31

32 Code availability (software application or custom code) Not applicable.

33

34 Ethics approval

35 The experiments comply with the current laws of France.

36 The corresponding author, Nadia Izadi-Pruneyre, serves as guarantor for the article and accepts
37 full responsibility for the work and the conduct of the study.

38

39 All authors:

40 1) made substantial contributions to the conception or design of the work; or the acquisition,
41 analysis, or interpretation of data;

42 2) drafted the work or revised it critically for important intellectual content;

43 3) approved the version to be published; and

44 4) agree to be accountable for all aspects of the work in ensuring that questions related to the
45 accuracy or integrity of any part of the work are appropriately investigated and resolved

Abstract

Abstract length: 189 words.

The ability to interact and adapt to the surrounding environment is vital for bacteria that colonise various niches and organisms. One strategy developed by Gram-negative bacteria is to secrete exoprotein substrates *via* the type II secretion system (T2SS). The T2SS is a proteinaceous complex spanning bacterial envelope that translocates folded periplasmic proteins such as toxins and enzymes to the extracellular milieu. In T2SS, a cytoplasmic ATPase elongates in the periplasm the pseudopilus, a non-covalent polymer composed of protein subunits named pseudopilins, anchored in the inner membrane by a transmembrane helix. The pseudopilus polymerisation is coupled to the secretion of substrates. The T2SS of *Dickeya dadantii* secretes more than 15 substrates, essentially plant cell wall degrading enzymes. In *D. dadantii*, the major pseudopilin or the major subunit of the pseudopilus is called OutG. To better understand the mechanism of secretion of these numerous substrates *via* the pseudopilus, we have been studying the structure of OutG by NMR. Here, as the first part of this study, we report the ¹H, ¹⁵N and ¹³C backbone and sidechain chemical shift assignment of the periplasmic domain of OutG and its NMR derived secondary structure.

Biological context

The type II secretion system (T2SS) is a molecular machinery which is widely used by Gram-negative bacteria to specifically secrete exoprotein substrates (Korotkov, Sandkvist and Hol, 2012; Thomassin *et al.*, 2017; Gu *et al.*, 2017). The substrates are species-specific and are often enzymes degrading biopolymers of carbohydrates, proteins, lipids and nucleotides (Cianciotto and White, 2017). In addition, T2SSs promote secretion of toxins, adhesins or cytochromes that are involved in respiration, motility or biofilm formation and remain attached to the bacterial cells (Nivaskumar and Francetic, 2014). The T2SS machinery is made up of twelve to fifteen components referred to as A to O, and spans both the inner and outer membrane. The T2SS could be divided into three functional blocks, the outer membrane pore composed of fifteen copies of the secretin D, the assembly platform comprising the inner membrane proteins F, L, M, C and the associated cytoplasmic ATPase E, which actively assembles a non-covalent polymer of protein subunits, called the pseudopilus, in the periplasm. The pseudopilus is composed of two types of proteins referred to as pseudopilins: the major pseudopilin G composing the bulk of the pseudopilus, and the minor pseudopilins H, I, J and K that form a complex initiating the pseudopilus formation (Nivaskumar and Francetic, 2014;

Escobar *et al.*, 2021). The minor pseudopilins are present in much lower abundance compared to the major pseudopilin but are crucial for the formation and stability of the pseudopilus and for secretion of substrates. Substrates are basically exported by the Sec or Tat translocator; are then folded in the periplasm where they are recruited by T2SS; with the elongation of the pseudopilus they reach the extracellular space through a secretin pore in the outer membrane (Korotkov, Sandkvist and Hol, 2012; Thomassin *et al.*, 2017; Gu *et al.*, 2017). It remains largely unknown how the growing pseudopilus can translocate the T2SS substrates and whether and how substrates are specifically recognized by pseudopilus.

Dickeya dadantii is a phytopathogenic γ -proteobacterium, which causes soft rot in vegetables and growing plants (Hugouvieux-Cotte-Pattat *et al.*, 2020). *D. dadantii* secrete more than 15 substrates *via* the T2SS (Kazemi-Pour, Condemine and Hugouvieux-Cotte-Pattat, 2004). Most of the characterized substrates are hydrolytic enzymes that degrade the cell wall of plants and thereby release nutrients that the bacteria can take up and metabolise (Hugouvieux-Cotte-Pattat *et al.*, 2014). Occurrence of multiple, structurally dissimilar substrates makes *D. dadantii* an attractive model to study the mechanisms of T2SS.

In this study, we focused on the major pseudopilin from *D. dadantii*, named OutG, as an essential T2SS element. OutG is a 13.5 kDa protein composed of 153 residues. It has a short N-terminal prepilin signal sequence which allows correct insertion in the inner membrane, and is then cleaved by a prepilin peptidase. The mature protein OutG protein of 146 residues long is anchored in the inner membrane with an N-terminal, 24 residues long, hydrophobic transmembrane helix followed by a C-terminal globular domain of 122 residues located in the periplasm (thereafter termed to as OutGp).

Structural insight into OutG is of high importance for the understanding of the putative interactions of the pseudopilus with the substrates and other T2SS components. Here we present the resonance assignment and derived secondary structure of OutGp as a starting point for its structural and interaction studies by NMR.

Method and experiments

Expression and purification of isotope labelled OutGp

The pET20b(+) vector (Novagen) was used for the expression of OutGp in the *Escherichia coli* periplasm. The N-terminus of OutGp was successively fused to the PelB signal peptide, followed by 6His and the TEV cleavage site. The hybrid OutGp is exported by Sec translocon into the periplasm and the PelB signal peptide is cleaved off by the LepB signal peptidase. Thereafter, 6His tag was removed by TEV protease treatment during purification. In

111 this way, the final OutGp protein used in this study is composed of an N-terminal GMG
112 sequence followed by residues M25 to P146 of the mature OutG (122 residues).

113 Uniformly ^{15}N - ^{13}C double-labelled OutGp was produced in M9 minimal media using 1 g/L of
114 $^{15}\text{NH}_4\text{Cl}$ and 4 g/L ^{13}C -glucose as the only nitrogen and carbon source, respectively. Protein
115 production was induced by addition of 1mM IPTG overnight at 18°C in *E. coli* BL21 (DE3)
116 cells. After induction the cells were lysed by sonication in equilibration buffer (50 mM Tris-
117 HCl pH 8, 100 mM NaCl, 10mM Imidazole), then the polynucleotides were digested with
118 nuclease (*Benzonase*®, *Sigma*) and the cell debris were pelleted by centrifugation at 16.000 g
119 for one hour at 4°C. The supernatant was loaded onto a HiTrap HP column (*Cytiva*) previously
120 loaded with Ni^{2+} ions, by saturating the resin with 0.1 M NiSO_4 solution, then equilibrated in
121 equilibration buffer. After loading of the lysate, the column was washed with the equilibration
122 buffer, and bound proteins were eluted with a linear gradient of imidazole from 10 to 300 mM.
123 The eluted protein fractions were pooled and treated with TEV protease overnight at 14°C, for
124 the cleavage of the N-terminal His-tag. The mixture was loaded onto a HiTrap HP column
125 (*Cytiva*) and the flow-through containing OutGp without its N-terminal His-tag was
126 concentrated in a Vivaspin® 20 (*Satorius*) concentrator with a 5 kDa molecular weight cut-off.
127 The concentrated OutGp fraction was loaded onto a Sephacryl S-100 column (*Cytiva*)
128 equilibrated in 50 mM HEPES, pH 7, 100 mM NaCl, 5 mM CaCl_2 . After elution, the fractions
129 containing OutGp were pooled and concentrated to 0.43 mM for NMR data acquisition. The
130 pH was adjusted to pH 6 by adding 2 μL of HCl at 0.1M. All steps of the purification were
131 evaluated by SDS-PAGE and in the final NMR sample no contaminants were visible in the
132 final protein preparation subjected to NMR. All buffers used during the purification were
133 supplemented with EDTA-free Protease inhibitor cocktail (*Roche*).

135 **NMR spectroscopy**

136 All NMR experiments were recorded on a 600 MHz Avance III HD (*Bruker Biospin*)
137 or a 800 MHz Avance NEO (*Bruker Biospin*) spectrometer, both equipped with a cryogenically
138 cooled triple resonance ^1H [^{13}C / ^{15}N] probe. Standard set of experiments for the ^1H , ^{15}N and ^{13}C
139 for backbone and side chain resonance assignment were recorded at 25°C using NMR
140 experiments implemented in TopSpin 3.6.1 and TopSpin 4.07 (*Bruker Biospin*) for the 600
141 MHz and 800 MHz, respectively, and IBS libraries (Favier and Brutscher, 2019): 2D ^1H - ^{15}N
142 HSQC, ^1H - ^{13}C HSQC, HBCBCGCDHD, HBCBCGCDCEHE and 3D HNCA, HN(CO)CA,
143 HNCACB, CBCA(CO)HN, HNCO, HN(CA)CO, HCCH-TOCSY, C(CO)NH-TOCSY,

H(CCO)NH-TOCSY, ^{15}N -NOESY and ^{13}C -NOESY. All proton chemical shifts were referenced to 2,2-dimethyl-2-silapentane-5-sulfonate (DSS) as 0 ppm. ^{15}N and ^{13}C chemical shifts were referenced indirectly to DSS (Wishart *et al.*, 1995). The data were analysed as previously described (Lopez-Castilla *et al.*, 2017) using CcpNMR Analysis (Vranken *et al.*, 2005), and the prediction of the secondary structure was achieved by analysing the chemical shifts of HN, H α , C α , C β , C', and N in TALOS-N (Shen and Bax, 2013).

Extent of assignment and data deposition

High quality data were obtained for OutGp as shown in the ^1H - ^{15}N HSQC spectrum in Fig 1. Backbone amide peaks were observed for all non-proline residues, except for the first three N-terminal residues (G22-G24). In total 96% of the observable backbone resonances and more than 91% of their corresponding sidechain resonances were assigned. Multiple backbone amide peaks of the C-terminal part of OutGp (N₁₃₈GSNGNGNP₁₄₆) were observed indicating that these residues have multiple conformations under the experimental conditions. Consequently, only one conformation of residues N138-N141 from this region could be assigned. It was not possible to assign any resonances corresponding to the last five residues (G₁₄₂NGNP₁₄₆), due to the repetitive nature of the sequence and the presence of a proline residue. The backbone amide peaks that could not be assigned are marked with red crosses in Fig 1. The chemical shift values have been deposited in the BioMagResBank (<http://www.bmrb.wisc.edu/>) under the accession number 51296.

Secondary Structure

The secondary structure of OutGp was estimated by using two approaches, secondary chemical shifts analysis and TALOS-N prediction (Fig. 2A). Secondary chemical shifts are obtained by calculating the differences between the assigned chemical shifts and the theoretical random coil chemical shift. The value of these secondary chemical shifts for C α , C β and C' ($\Delta\delta^{13}\text{C}\alpha$, $\Delta\delta^{13}\text{C}\beta$ & $\Delta\delta^{13}\text{C}'$) is related to the secondary structure of the individual residues. Positive values of $\Delta\delta^{13}\text{C}\alpha$ and $\Delta\delta^{13}\text{C}'$ correspond to a α -helical conformation whereas negative values correspond to β -strand structure. For $\Delta\delta^{13}\text{C}\beta$ positive values reflect β -strand structure and negative values α -helical conformation (Fig. 2A). To complete this prediction, a TALOS-N analysis of the assigned HN, H α , C α , C β , C' and N chemical shifts was performed (Shen and Bax, 2013). The TALOS-N probabilities for each residue to be in an α -helical conformation or a β -strand structure is plotted in Fig. 2B.

The results of both analysis methods agree to the location of the secondary structure elements, and reveal three helical segments (α 1: K28-N54, α 2: T60-Q62 and α 3: G65-L67) and two β -strands (β 1: Q99-V101 and β 2: D110-S113), in the α 1- α 2- α 3- β 1- β 2 sequential order. Structural determination of OutGp will be performed following this study.


Acknowledgements

This work was funded by the Institut Pasteur, the Centre National de la Recherche Scientifique (CNRS), the French Agence Nationale de la Recherche (ANR Synergy-T2SS ANR-19-CE11-0020-01) and European Union's Horizon 2020 Research and Innovation program under the Marie Skłodowska-Curie Grant agreement no 765042. We thank Rémy Le Meur and Bruno Vitorge for their help in NMR experiments, and Olivera Francetic for helpful discussions. The 800-MHz NMR spectrometer of the Institut Pasteur was partially funded by the Région Ile de France (SESAME 2014 NMRCHR grant no 4014526).

References

- Cianciotto, N.P. and White, R.C. (2017) 'Expanding Role of Type II Secretion in Bacterial Pathogenesis and Beyond.' *Infect Immun* 85. <https://doi.org/10.1128/IAI.00014-17>
- Escobar, C.A., Douzi, B., Ball, G., Barbat, B., Alphonse, S., Quinton, L., Voulhoux, R. and Forest, K.T. (2021) 'Structural interactions define assembly adapter function of a type II secretion system pseudopilin.' *Structure*. Oct 7;29(10):1116-1127.e8. doi: 10.1016/j.str.2021.05.015.
- Favier, A. and Brutscher, B (2019) 'NMRLib: user-friendly pulse sequence tools for Bruker NMR spectrometers', *Journal of Biomolecular NMR*, 73, pp. 199–211. doi: 10.1007/s10858-019-00249-1.
- Gu, S., Shevchik, V.E., Shaw, R., Pickersgill, R.W. and Garnett, J.A. (2017) 'The role of intrinsic disorder and dynamics in the assembly and function of the type II secretion system.' *Biochim Biophys Acta Proteins Proteom* 1865:1255-1266. doi:10.1016/j.bbapap.2017.07.006.
- Hugouvieux-Cotte-Pattat, N., Condemine, G., Gueguen, E. and Shevchik, V.E. (2020) 'Dickeya Plant Pathogens.' in *eLS*, John Wiley & Sons, Ltd (Ed.). pp 1-10
- Hugouvieux-Cotte-Pattat, N., Condemine, G., and Shevchik, V. E. (2014) 'Bacterial pectate lyases, structural and functional diversity.' *Environmental microbiology reports* 6, 427-440 doi: 10.1111/1758-2229.12166.
- Kazemi-Pour, N., Condemine, G. and Hugouvieux-Cotte-Pattat, N. (2004) 'The secretome of the plant pathogenic bacterium *Erwinia chrysanthemi*', *Proteomics*, 4(10), pp. 3177–3186. doi: 10.1002/pmic.200300814.

- 219 Korotkov, K. V., Sandkvist, M. and Hol, W. G. J. (2012) 'The type II secretion system:
220 Biogenesis, molecular architecture and mechanism', *Nature Reviews Microbiology*, pp. 336–
221 351. doi: 10.1038/nrmicro2762.
- 222
- 223 López-Castilla, A. *et al.* (2017) '1H,15N and13C resonance assignments and secondary
224 structure of PulG, the major pseudopilin from *Klebsiella oxytoca* type 2 secretion system',
225 *Biomolecular NMR Assignments*, 11(2), pp. 155–158. doi: 10.1007/s12104-017-9738-7.
- 226
- 227 Nivaskumar, M. and Francetic, O. (2014) 'Type II secretion system: A magic beanstalk or a
228 protein escalator', *Biochimica et Biophysica Acta - Molecular Cell Research*. Elsevier B.V.,
229 1843(8), pp. 1568–1577. doi: 10.1016/j.bbamcr.2013.12.020.
- 230
- 231 Shen, Y. and Bax, A. (2013) 'Protein backbone and sidechain torsion angles predicted from
232 NMR chemical shifts using artificial neural networks', *Journal of Biomolecular NMR*, 56(3),
233 pp. 227–241. doi: 10.1007/s10858-013-9741-y.
- 234
- 235 Thomassin, J.L. *et al* (2017) 'The trans-envelope architecture and function of the type 2
236 secretion system: new insights raising new questions. *Mol Microbiol* 105:211-226. doi:
237 10.1111/mmi.13704.
- 238
- 239 Vranken, W. F. *et al.* (2005) 'The CCPN data model for NMR spectroscopy: Development of
240 a software pipeline', *Proteins: Structure, Function and Genetics*, 59(4), pp. 687–696. doi:
241 10.1002/prot.20449.
- 242
- 243 Wishart, D. S. *et al.* (1995) 'H, 13 C and 15 N chemical shift referencing in biomolecular
244 NMR', *Journal of Biomolecular NMR*, 6.
- 245
- 246
- 247
- 248






Fig. 2 – Secondary structure prediction based on the chemical shifts of OutGp.

A Secondary chemical shifts of $\text{C}\alpha$, $\text{C}\beta$ and C' resonances. The sequence of OutGp is shown at the top. **B** TALOS-N secondary structure probabilities were used to predict the secondary structure elements along the protein sequence. The probability for each residue to be in an α -helical conformation (black bars) or in β -sheet conformation (grey bars) are plotted as a function of residue number. Both the secondary chemical shifts and TALOS-N secondary structure prediction agree to a consensus of secondary structure elements illustrated between panel A and B (cylinders represent α -helices and arrows represent β -sheets).