



HAL
open science

Functional spectrum and specificity of mitochondrial ferredoxins FDX1 and FDX2

Vinzent Schulz, Somsuvro Basu, Sven-A. Freibert, Holger Webert, Linda Boss, Ulrich Mühlenhoff, Fabien Pierrel, Lars-O. Essen, Douglas Warui, Squire Booker, et al.

► **To cite this version:**

Vinzent Schulz, Somsuvro Basu, Sven-A. Freibert, Holger Webert, Linda Boss, et al.. Functional spectrum and specificity of mitochondrial ferredoxins FDX1 and FDX2. *Nature Chemical Biology*, 2022, 10.1038/s41589-022-01159-4 . hal-03836706

HAL Id: hal-03836706

<https://hal.science/hal-03836706>

Submitted on 2 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Functional spectrum and structural specificity of mitochondrial ferredoxins FDX1 and FDX2

Vinzent Schulz¹, Somsuvro Basu^{1,#}, Sven-A. Freibert¹, Holger Weibert¹, Linda Boß¹,
Ulrich Mühlenhoff¹, Fabien Pierrel², Lars-O. Essen³, Douglas M. Warui⁴, Squire Booker^{4,5,6},
Oliver Stehling^{1,7,*}, and Roland Lill^{1,7,*}

¹ Institut für Zytobiologie, Philipps-Universität Marburg, Karl-von-Frisch-Str. 14, 35032 Marburg, Germany

² Univ. Grenoble Alpes, CNRS, UMR 5525, VetAgro Sup, Grenoble INP, TIMC, 38000 Grenoble, France

³ Department of Biochemistry, Faculty of Chemistry, Philipps-Universität Marburg, Hans-Meerwein-Str. 4, 35032 Marburg, Germany

⁴ Department of Chemistry, The Pennsylvania State University, University Park, PA 16802, USA

⁵ Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, PA 16802, USA

⁶ The Howard Hughes Medical Institute, The Pennsylvania State University, University Park, PA 16802, USA

⁷ Zentrum für Synthetische Mikrobiologie Synmikro, Karl-von-Frisch-Str. 14, 35032 Marburg, Germany

Present affiliation and address: Freelance medical communications consultant; Vídeňská 281/77, 63900 Brno, Czech Republic

* Corresponding authors:

Roland Lill

E-mail: Lill@staff.uni-marburg.de, phone: +49-6421-286 6449

Oliver Stehling

E-mail: Stehling@staff.uni-marburg.de, phone: +49-6421-286 4044

Keywords: Iron-sulfur proteins, lipoyl synthase, biogenesis, heme, coenzyme Q, mitochondria, cellular respiration, elesclomol.

38 **Abstract**

39 Ferredoxins comprise a large family of iron-sulfur (Fe/S) proteins that shuttle electrons in
40 diverse biological processes. Human mitochondria contain two isoforms of [2Fe-2S]
41 ferredoxins, FDX1 (aka adrenodoxin) and FDX2, with known functions in cytochrome P450-
42 dependent steroid transformations and Fe/S protein biogenesis. Here, we show that only
43 FDX2, but not FDX1, is involved in Fe/S protein maturation. Vice versa, FDX1 is specific not
44 only for steroidogenesis, but also for heme *a* and lipoyl cofactor biosyntheses. In the latter
45 pathway, FDX1 provides electrons to kickstart the radical chain reaction catalyzed by lipoyl
46 synthase. We also identified lipoylation as a target of the toxic anti-tumor copper ionophore
47 elesclomol. Finally, the striking target specificity of each ferredoxin was assigned to small
48 conserved sequence motifs. Swapping these motifs changed the target specificity of these
49 electron donors. Together, our findings identify new biochemical tasks of mitochondrial
50 ferredoxins and provide structural insights into their striking functional specificity.

51

52 Introduction

53 Ferredoxins (FDXs) comprise a large family of redox proteins found in all kingdoms of life^{1,2}.
54 Their iron-sulfur (Fe/S) cofactors allow electron transfer to numerous targets in diverse
55 biological pathways. FDXs from pathogens may serve as drug targets^{3,4}. Mitochondria harbor
56 the [2Fe-2S] cluster-containing FDXs that are evolutionarily derived from bacteria, and receive
57 their electrons from NADPH via ferredoxin reductase (FDR)¹. The long-known mammalian
58 mitochondrial FDX1 (aka adrenodoxin) functions in the metabolism of steroid hormones, bile
59 acids, and vitamins A and D transferring electrons to all seven class I mitochondrial cytochrome
60 P450 enzymes⁵. Mammals, including humans, possess the additional [2Fe-2S] FDX2 which,
61 however, cannot take over the function of FDX1 in cortisol formation⁶. Instead, FDX2 and its
62 fungal counterparts including *S. cerevisiae* Yah1 are central components of the core iron-sulfur
63 cluster assembly (ISC) machinery, and hence are present in virtually all mitochondria or in
64 mitochondria-related, ISC-containing mitosomes and hydrogenosomes^{4,6-11}. FDX2 is required
65 twice in mitochondrial Fe/S protein maturation. Initially, it interacts with the cysteine
66 desulfurase complex NFS1-ISD11-ACP1¹²⁻¹⁴ and reduces a persulfide (-SSH) bound to the
67 scaffold protein ISCU2 to sulfide, thereby inducing *de novo* [2Fe-2S] cluster synthesis^{9,15-17}.
68 Later in mitochondrial [4Fe-4S] cluster assembly, FDX2 facilitates reductive fusion of two [2Fe-
69 2S] to a [4Fe-4S] cluster on the ISCA proteins¹⁸. The importance of human FDX2 and FDR
70 is reflected by severe genetic diseases including mitochondrial myopathy and sensory
71 neuropathies¹⁹⁻²². Moreover, FDR is a p53-dependent tumor suppressor, and its deficiency
72 leads to tumor formation and liver disease in a mouse model²³.

73 Beyond steroidogenesis and Fe/S protein biogenesis, mitochondrial FDXs were
74 reported to be involved in other metabolic reactions. For instance, yeast mitochondrial Yah1,
75 together with its reductase Arh1, provides electrons to the hydroxylase Cox15, thereby yielding
76 heme *a/a*₃ of cytochrome *c* oxidase (COX)²⁴⁻²⁶. Since *FDX2* knockdown in human cells is
77 associated with a partial COX defect, the protein was suggested to perform a similar function
78 as Yah1⁶. Additionally, Yah1-Arh1 are required for coenzyme Q (ubiquinone or yeast CoQ₆)
79 biosynthesis, supposedly to reduce the Coq6 protein that catalyzes C5-hydroxylation of a CoQ₆
80 intermediate^{27,28}. Neither human FDX1 nor FDX2 can replace Yah1 in yeast CoQ₆ synthesis,
81 leaving open whether these electron donors might support CoQ₁₀ biosynthesis in humans.
82 Conflicting results have been reported for the involvement of human FDX1 in cellular Fe/S
83 protein biogenesis. Earlier studies, using *in vivo* complementation in yeast and RNA
84 interference (RNAi) experiments in human cell culture, identified FDX2 as the sole Fe/S protein
85 biogenesis-related FDX⁶. Accordingly, FDX2 but not FDX1 was crucial for *in vitro* reconstitution
86 of both [2Fe-2S] and [4Fe-4S] cluster formation^{9,16,18,29}. In principle, the specialized functions
87 of human FDX1 and FDX2 fit their strikingly different tissue distribution identified by

88 immunostaining⁶ (Suppl. Fig. 1a; see legend for details). However, systematic transcript
89 analyses indicate significant expression of *FDX1* mRNA in a variety of tissues despite
90 undetectable protein levels, raising the question of a broader function of FDX1 (Suppl. Fig. 1b).
91 Accordingly, two reports proposed FDX1 to function also in Fe/S protein biogenesis, based on
92 RNAi depletion experiments and *in vitro* [2Fe-2S] cluster synthesis on ISCU2 using FDXs
93 chemically reduced by dithionite (DT)^{8,30}. It remained unexplained, however, why the two FDXs
94 would not functionally complement each other *in vivo* in Fe/S protein biogenesis.

95 Here, we aimed to clarify these important issues and identified so far unknown functions
96 of the human FDXs in mitochondrial metabolism, most prominently the essential role of FDX1
97 to kickstart the radical chain reaction of lipoyl synthase (LIAS). Knowledge of the functional
98 capacity of the two FDXs is crucial for understanding their impact on both mitochondrial
99 metabolism and mitochondria-cytoplasm interaction, particularly in (tumorigenic) cells that
100 shutdown their dependence on mitochondrial respiration³¹⁻³³. We also report a striking target
101 specificity of the two FDXs, and hence examined the structural basis of target recognition by
102 site-directed mutagenesis, leading to the identification of small structural segments that
103 functionally discriminate the two FDXs. Our work defining the functional spectrum of the human
104 FDXs may have possible implications for better understanding disease phenotypes and for
105 biotechnological applications of these unique electron donors³⁴.

106

107 **Results**

108 **FDX2 but not FDX1 is required for cellular Fe/S protein biogenesis**

109 *In vivo* and *in vitro* studies have provided conflicting results on the function of human FDX1 in
110 Fe/S protein biogenesis, while the involvement of FDX2 in this process is undisputed^{6,8,9,30}. Re-
111 evaluation of these experiments confirmed that substantial depletion of FDX1 by RNAi did not
112 induce any significant growth defect or reduction in the levels of various mitochondrial and
113 cytosolic-nuclear Fe/S proteins, in contrast to depletion of FDX2, which is essential for the
114 biogenesis of cellular Fe/S proteins including FDX1 (Fig. 1a; Suppl. Fig. 2a,b)⁶. To exclude
115 that our RNAi knockdown of *FDX1* was too weak to elicit a detectable phenotype, we knocked
116 out *FDX1* (or *FDX2* as a control) by CRISPR-Cas9 technology. Application of three different
117 *FDX2*-directed guide RNAs (gRNAs) did not yield viable FDX2-depleted cells after puromycin
118 selection, consistent with the indispensable function of FDX2 in Fe/S protein assembly^{6,8}. Two
119 of three *FDX1*-directed gRNAs (CC1 and CC2; Suppl. Table 1) efficiently depleted FDX1, and,
120 in contrast to the RNAi findings, profoundly decreased cell growth over time, while gRNA CC3
121 was ineffective (Fig. 1b; Extended data Fig. 1a,b). The results suggested a critical *in vivo* role
122 of FDX1 disclosed only upon strong depletion. CC1- or CC2-mediated *FDX1* knockout hardly
123 affected the levels of the mitochondrial Fe/S-cluster-containing respiratory chain subunits
124 NDUFS1, NDUFS8, NDUFV2, SDHB, and UQCRC1, in contrast to the RNAi-mediated *FDX2*
125 knockdown (Fig. 1a,b; Extended data Fig. 1b; Suppl. Fig. 2b). Consistently, the enzyme
126 activities of key mitochondrial Fe/S proteins, i.e. aconitase and succinate dehydrogenase
127 (SDH), were not altered upon *FDX1* knockout (Fig. 1c; Extended data Fig. 1c-e), unlike upon
128 *FDX2* RNAi depletion^{6,8}. Together, these findings strongly argue against a critical *in vivo* role
129 of FDX1 in cellular Fe/S protein biogenesis.

130 We next compared the capacities of FDX1 and FDX2 to *in vitro* donate electrons to the
131 initial step of Fe/S protein biogenesis, i.e. the *de novo* [2Fe-2S] cluster synthesis on the
132 scaffold protein ISCU2 by enzymatic function of the cysteine desulfurase complex NFS1-
133 ISD11-ACP1 (NIA) and frataxin (FXN)^{9,15,30}. [2Fe-2S] cluster formation was measured by
134 circular dichroism (CD) spectroscopy, and strictly depended on FDX2 addition up to equimolar
135 amounts to NIA (Fig. 1d; Extended data Fig. 2). In contrast, even a fivefold excess of FDX1
136 over NIA did not support any [2Fe-2S] cluster synthesis above control reactions performed
137 without any FDX. In summary, our *in vivo* and *in vitro* results do not indicate a direct role of
138 FDX1 in human Fe/S protein biogenesis, in contrast to previous suggestions^{8,30}. Rather, the
139 data support and extend earlier findings indicating a vital role of FDX2 as the exclusive
140 physiological electron donor in mitochondrial Fe/S protein biogenesis^{6,9,18}.

141

142

143 **A crucial role of FDX1 in heme a-dependent cytochrome c oxidase activity**

144 We next sought to find an explanation for the substantial growth retardation upon *FDX1* gene
145 knockout. The culture medium of *FDX1*-deficient cells showed a profound acidification
146 (Extended data Fig. 3a), indicating a metabolic switch from mitochondrial oxidative
147 phosphorylation to glycolysis. Since respiratory complexes I-III and aconitase were unaffected
148 in these cells (cf. Fig. 1b,c), we analyzed the activity of respiratory complex IV (cytochrome c
149 oxidase, COX). We found a diminution in the levels of COX2 and COX6A subunits and a
150 substantial decrease in COX enzyme activity (Fig. 2a,b; Extended data Fig. 3b). These effects
151 were specific because complementation of *FDX1*-deficient cells by mitochondria-targeted
152 *FDX1* recovered both COX subunit levels and activity (Extended data Fig. 3c,d).

153 COX lacks Fe/S clusters, and consistently RNAi-mediated depletion of *FDX2* only
154 weakly affected COX2 or COX6A subunit levels, in contrast to the severe decrease upon *FDX1*
155 gene knockout (Fig. 2a; Suppl. Fig. 2b). COX contains two Cu centers and heme *a/a₃* cofactors
156 that are derived from heme *b* by COX10-dependent farnesylation (yielding the intermediate
157 heme *o*) and COX15-catalyzed formylation^{24,26,35-37}. A role of *FDX1* in heme *b* synthesis as an
158 explanation for the COX defect was excluded, because heme *b*-containing complexes II and
159 III were not altered in *FDX1* knockout cells (Fig. 1b,c; Extended data Fig. 1b-d). In *S.*
160 *cerevisiae*, Cox15-dependent heme *a* formation requires the function of ferredoxin Yah1,
161 suggesting that the impaired COX function in *FDX1* knockout cells may be caused by a heme
162 *a* synthesis defect. A direct measurement of heme *a* levels by HPLC²⁴ or mass spectrometry
163 failed due to detection limits. Notably, *FDX1*-deficient cells (CC2-based knockout) retained
164 33% of wild-type COX activity, likely due to the presence of *FDX2* (Fig. 2b; Extended data
165 Fig. 3b). Consistently, RNAi depletion of *FDX2* was associated with some decrease in COX
166 subunit levels and activity (Suppl. Fig. 2b)⁶. Since this effect is also seen upon depletion of
167 other ISC factors including the ISCA-IBA57 proteins³⁸, it may be an indirect consequence of
168 *FDX2* depletion on COX. Collectively, *FDX1* performs a critical role for COX activity,
169 presumably by its function in heme *a* synthesis, yet *FDX2* may partially replace *FDX1* in this
170 process.

171 Our previous *FDX1* and *FDX2* complementation studies in Yah1-depleted Gal-*YAH1*
172 yeast cells had indicated a (weak) requirement of *FDX2* but not *FDX1* for yeast COX activity⁶,
173 challenging our results with human cells. We reasoned that this discrepancy may be due to a
174 failure of human *FDX1* to properly interact with yeast Cox15 during heme *o* to heme *a*
175 conversion. We therefore expressed human *COX15* in the regulatable yeast strain Gal-*YAH1*-
176 *cox15* Δ in which yeast *COX15* was deleted. When Yah1 was depleted by growth in the
177 presence of glucose (Glc) or glycerol (Gly)⁷, cells showed a growth defect that could be
178 rescued only by expression of *FDX2* but not by *FDX1*, consistent with the vital function of *FDX2*
179 in Fe/S protein biogenesis (Fig. 2c). However, under respiratory growth conditions (Gly-

180 containing medium) only the combined expression of FDX1 and FDX2 resulted in full growth
181 complementation, indicating independent, complementary functions of the two human FDXs.
182 Measurement of COX activity in extracts of Yah1-depleted Gal-*YAH1-cox15* Δ cells still showed
183 low residual COX activity, due to the leaky *GAL* promoter (Fig. 2d). Upon FDX1
184 complementation, COX activity increased ca. 3-fold, despite the severe growth defect under
185 this condition. In contrast, FDX2 complementation did not recover COX activity, despite normal
186 cell growth on glucose-containing medium due to restored Fe/S protein biogenesis. These
187 results for COX15-humanized yeast perfectly support our observations with human cells,
188 implying that FDX1 is the predominant human ferredoxin supporting COX15-dependent heme
189 *a* formation.

190

191 **Both human ferredoxins are dispensable for ubiquinone biosynthesis**

192 In *S. cerevisiae*, both Yah1 and Arh1 are essential for the biosynthesis of ubiquinone
193 (coenzyme Q₆; CoQ₆) by supporting Coq6-catalyzed hydroxylation of precursor
194 compounds^{27,28}. The electron donor for the biosynthesis of human ubiquinone (CoQ₁₀) is
195 unknown, and we therefore tested the potential function of the two human FDXs in this process.
196 Determination of cellular CoQ₁₀ levels in *FDX1* knockout cells did not diminish CoQ₁₀ levels,
197 not even upon prolonged tissue culture following puromycin selection (Fig. 2e). Likewise,
198 *FDX2*-deficient cells contained wild-type levels of CoQ₁₀ even after 9 days of RNAi-mediated
199 depletion (Fig. 2f, left) when cells showed a growth defect (Extended data Fig. 4a). We
200 analyzed CoQ₁₀ levels in *FDX2*-depleted *FDX1* knockout cells to test a potential redundant
201 function of the FDXs. The FDX double-deficient cells showed a more profound growth
202 retardation compared to single depletions (Extended data Fig. 4a), yet no significant changes
203 in CoQ₁₀ levels (Fig. 2f, right), despite severe diminution of mitochondrial Fe/S protein levels
204 (Extended data Fig. 4b). This result indicates that CoQ₁₀ synthesis in human cells is FDX-
205 independent, and, unlike in yeast, may involve another, still unknown oxidoreductase.

206

207 **FDX1 functions as a radical chain starter for LIAS-dependent lipoyl synthesis**

208 To search for reasons of the growth defect and metabolic switch of *FDX1* knockout cells, we
209 next performed anti-lipoyl immunostaining, and thereby identified a strong lipoylation defect of
210 the E2 subunits dihydrolipoyl transacetylase (DLAT) and dihydrolipoyl succinyl transferase
211 (DLST) of pyruvate dehydrogenase (PDH) and α -ketoglutarate dehydrogenase (KGDH),
212 respectively (Fig. 3a)³⁹. The steady-state levels of the DLAT protein and of the radical S-
213 adenosyl methionine (SAM) Fe/S enzyme lipoyl synthase (LIAS) remained unchanged.
214 Complementation of *FDX1*-deficient cells by expression of mitochondria-targeted Su9-*FDX1*

215 recovered DLAT lipoylation (Extended data Fig. 5). This finding suggests a crucial function of
216 FDX1 in lipoylation, an observation not revealed by the less severe RNAi depletion of FDX1
217 (cf. Suppl. Fig. 2b).

218 We surmised that FDX1 could act as the so far unknown electron donor starting the
219 radical chain reaction of LIAS (explained in Extended data Fig. 6a)^{40,41}. To test this idea, we
220 adapted an *in vitro* assay for bacterial lipoyl synthase (LipA) for human LIAS⁴⁰. An octapeptide
221 containing the octanoyllysyl precursor served as a substrate analogue, and the SAM-
222 dependent formation of both the 6-thiooctanoyl intermediate and lipoyl product under strictly
223 anaerobic conditions was measured and quantified by HPLC-MS. In the presence of the non-
224 physiological reductant DT, normally used to artificially start the radical chain, both the
225 intermediate and product of the LIAS reaction were formed strictly dependent on the presence
226 of DT, SAM, and LIAS (Fig. 3b,c). Strikingly, reduced FDX1 (by action of NADPH and FDXR)
227 could replace DT much more efficiently as an electron source. Both the time courses and final
228 yields for intermediate and product formation differed, comparable to reports for bacterial
229 LipA⁴⁰, assigning the second sulfur insertion step as being rate-limiting under *in vitro* conditions
230 (Extended data Fig. 6b). We observed 29- and 61-fold, respectively, lower yields of
231 intermediate and product with reduced FDX2 instead of FDX1 (Fig. 3b,c). This substantial
232 difference suggested FDX1 being the physiological electron donor for LIAS, consistent with
233 our *in vivo* experiments.

234 We further took advantage of human LIAS-expressing yeast cells lacking the
235 endogenous *LIP5* gene to verify the FDX1 function and specificity in lipoylation. Yeast Gal-
236 *YAH1-lip5Δ* cells were transformed with plasmids harboring human *LIAS*, *FDX1* and/or *FDX2*,
237 followed by growth in the presence of either glucose or galactose. As expected, anti-lipoyl
238 staining showed no lipoylation in extracts of Yah1-depleted Gal-*YAH1-lip5Δ* cells (lacking
239 human LIAS) compared to wild-type cells (Fig. 3d). Lipoylation could be restored to wild-type
240 levels by complementation with LIAS when cells were grown under Yah1-inducing conditions
241 (galactose), but only weakly upon depletion (glucose) of Yah1 that is needed for LIAS Fe/S
242 cluster maturation during ectopic expression. FDX1 but not FDX2 fully complemented the
243 severe lipoylation defect of LIAS-containing, Yah1-depleted Gal-*YAH1-lip5Δ* cells. Apparently,
244 the low residual levels of Yah1 in this condition (leaky Gal promoter) were adequate to mature
245 enough FDX1 and LIAS for sufficient lipoylation. The humanized yeast model fully supports
246 the dedicated role of human FDX1 in lipoylation.

247 To confirm this notion, we measured the enzyme activities of the lipoyl-dependent yeast
248 KGDH and the Fe/S enzyme aconitase in this yeast model (Fig. 3e). As expected, *lip5Δ*
249 deletion cells lacked KGDH activity, and aconitase activity dropped by half, likely as an indirect
250 effect of impaired mitochondrial metabolism. In Yah1-depleted Gal-*YAH1-lip5Δ* cells, absent
251 KGDH and low aconitase activities were rescued to the *lip5Δ* levels only by expression of LIAS,

252 FDX1, and FDX2. The findings indicate the specific, non-overlapping functions of FDX1 and
253 FDX2 in LIAS-dependent lipoylation and Fe/S protein biogenesis, respectively. This view
254 perfectly agrees with the growth behavior of the various LIAS-expressing Gal-YAH1-lip5Δ cells
255 in different liquid or solid media (Extended data Fig. 7).

256

257 **Lipoylation is a primary target of the elesclomol-copper complex**

258 FDX1 has been proposed to the target of the anti-cancer drug elesclomol (Ele) because
259 CRISPR-Cas9 deletion of *FDX1* desensitizes these cells against treatment with Ele³². Ele was
260 shown to be toxic for proteasome inhibitor-resistant cells depending on increased
261 mitochondrial energy metabolism, and it was proposed that Ele imparts a toxic gain of FDX1's
262 function in Fe/S protein biogenesis. The toxic effect was further shown to be mediated by the
263 complex of Ele with Cu (Ele:Cu) rather than Ele alone³², which is consistent with earlier reports
264 that Ele acts as a Cu ionophore transporting Cu to mitochondria, thereby restoring COX
265 function in Cu deficiency^{42,43}. A study published during review of this work suggested both
266 FDX1 and lipoylation as targets of Ele:Cu toxicity, but no molecular explanations were
267 described³³. Our *in vivo* and *in vitro* findings refuting FDX1 function in Fe/S protein biogenesis,
268 but showing its role in lipoylation, urged for a biochemical evaluation of the toxic role of Ele:Cu.

269 We first tested the effects of Ele and/or Cu on LIAS- and FDX1-dependent lipoyl
270 synthesis *in vitro*. Addition of Ele alone did not affect 6-thiooctanoyl and lipoyl formation, even
271 at a 14-fold excess over FDX1 (Fig. 4a,b; left). In contrast, both Cu and Ele:Cu imposed severe
272 effects on these reactions in a dose-dependent manner (Fig. 4a,b; middle and right), fully
273 consistent with reports indicating toxic effects of Ele only in the presence of Cu or other
274 metals^{32,33,42,44,45}. Notably, an excess of Ele over Cu slightly attenuated the synthesis defect.
275 The high stability constant of $K_{\text{Ele:Cu(II)}}=10^{24.2}$ L/mol⁴⁶ suggests that both Cu and Ele:Cu inhibit
276 lipoylation, with Cu being slightly more potent. Similar relative impacts of Cu and Ele:Cu have
277 been observed for superoxide dismutase enzyme activity⁴⁴. Collectively, our *in vitro* results
278 identify lipoylation rather than Fe/S protein biogenesis as a sensitive target of Ele:Cu toxicity.
279 In particular, the second sulfur insertion step appears to be most affected (Fig. 4a,b; Extended
280 data Fig. 6a). Since lipoyl-dependent PDH and KGDH are central components of mitochondrial
281 metabolism, our findings satisfactorily explain why particularly cells dependent on this process
282 are most vulnerable to Ele treatment³².

283 The *in vitro* lipoylation assay did not readily allow the discrimination between FDX1
284 and/or LIAS as potential targets of the Ele:Cu toxicity. To address this issue biochemically, we
285 first studied the impact of Ele and/or Cu on FDX1 during NADPH-FDXR-FDX1-dependent
286 electron transfer to cytochrome *c*. Reduction of cytochrome *c* was not affected by addition of
287 Ele or Ele:Cu, yet was fully abolished by Cu (Fig. 4c). The striking difference of this result to

288 the observed inhibition of lipoylation by both Cu and Ele:Cu indicates the LIAS reaction as the
289 preferential target of Ele:Cu and FDX1 as target of Cu only. Consistently, only Cu but not Ele
290 or Ele:Cu destroyed the Fe/S clusters of both FDX1 and FDX2 (Fig. 4d,e; Suppl. Fig. 3), in
291 agreement with the Cu sensitivity of yeast *Yah1*⁴⁷. Our *in vitro* findings identifying LIAS function
292 in lipoylation as a target of the toxic effect of Ele:Cu are further supported *in vivo*. Human cells
293 treated with Ele or Ele:Cu showed a strong impairment of lipoylation and a destabilization of
294 LIAS (not seen upon *FDX1* knockout), while DLAT protein and the two FDXs were hardly
295 affected (Fig. 4f). In contrast, Cu alone elicited no detectable effects. Collectively, the
296 biochemical and cell biological results document that the Ele:Cu complex strongly affects LIAS
297 function in lipoylation and may be the target of the toxic effect of Ele (see Suppl. Discussion).
298

299 **Converting FDX1 into a ferredoxin functional in Fe/S protein biogenesis**

300 To address the structural basis of the functional specificity of the two human FDXs, we solved
301 the crystal structure of mature FDX2 (residues 66-171; PDB ID: 2Y5C; Suppl. Table 2; Suppl.
302 Results) for comparison with the known FDX1 structure (PDB ID: 3P1M). Significant structural
303 differences between FDX1 and FDX2 were observed only for the loop between helix C and the
304 Fe/S cluster binding site, because the Phe-Gly (FG) dipeptide is missing in FDX2 (Fig. 5a;
305 Extended data Fig. 8; Suppl. Fig. 4; Suppl. Results). Comparison of the surface potentials⁴⁸
306 revealed a less negatively charged area of FDX2 (Fig. 5b; Suppl. Fig. 5; Suppl. Results; for
307 details see Discussion), mainly due to exchanges of semi-conserved Asp91^{FDX1} by His95^{FDX2}
308 and Glu133^{FDX1} by Arg135^{FDX2} (Extended data Fig. 8). Overall, the 3D structures of FDX1 and
309 FDX2 display only minor differences in backbone geometry.

310 Consequently, we compared multi-sequence alignments of FDX1-, FDX2-, fungal- and
311 bacterial-type FDXs (Extended data Fig. 8). Ten regions (including those structurally
312 pinpointed above) were identified that distinguish FDX1- and FDX2-type proteins (named M1
313 to M10). We first tested the importance of these regions for FDX2 function by introducing the
314 corresponding FDX1 M1-M10 sequences (Suppl. Table 3), and we examined N- and C-
315 terminal truncations of FDX2 (termed Δ N13 and Δ C12). FDX2 mutant proteins were expressed
316 in *Yah1*-depleted Gal-*YAH1* cells, and growth was analyzed (Extended data Fig. 9).
317 Exchanges of regions M3 to M7 decreased the *Yah1*-rescuing function of FDX2, with segment
318 M7 showing the most pronounced growth defect. Moreover, the C terminus (Δ C12) but not the
319 N terminus (Δ N13) of mature FDX2 was critical for growth. Hence, region M7 and the FDX2 C
320 terminus are important structural elements for FDX2 function.

321 Second, we introduced FDX2 segments M1-M10 (either alone or in combination) into
322 FDX1, and we transferred 14 or 27 C-terminal FDX2 residues into FDX1 (exC14 and exC27),
323 in an attempt to generate FDX2 functionality in FDX1 (Fig. 5c,d). Like FDX1⁶, most FDX1

324 variants did not detectably rescue the growth defect of Yah1-depleted Gal-YAH1 yeast, even
325 though all recombinantly purified proteins contained a [2Fe-2S] cluster, except for FDX1-M5
326 (deletion of the FG motif), explaining why this mutant is non-functional (Fig. 5d; Suppl. Fig. 6).
327 However, partial growth restoration was observed by introducing the FDX2 C terminus into
328 FDX1. Further growth improvement was obtained by additional exchange of segment M7,
329 particularly in combination with either M3 or M6. Consistent with the findings above (Extended
330 data Fig. 9), region M7 and the C terminus are decisive features of FDX2 functionality, while
331 regions M3 and M6 fulfil a weak auxiliary role. To extend these findings, we measured the
332 enzyme activities of aconitase and catalase (a non-Fe/S protein, yet indirectly affected by Fe/S
333 protein synthesis defects^{6,49}) in Yah1-depleted Gal-YAH1 yeast complemented with FDX1,
334 FDX2, or the growth-restoring FDX1-M3+M7+exC27 or FDX1-M6+M7+exC27 mutant proteins
335 (Fig. 5e,f). While wild-type FDX1 showed no improvement of both enzyme activities in Yah1-
336 depleted cells, both FDX1 mutant proteins restored ca. 30-50% of the activities seen in FDX2-
337 complemented cells. Apparently, small increases in Fe/S protein biogenesis activity by
338 introducing FDX2 functionality into FDX1 can lead to partial growth restoration.

339

340 **Converting FDX2 into a ferredoxin with FDX1 functionality**

341 In a reversed approach, we sought to identify regions specific for FDX1 functionality. We
342 introduced the FDX-discriminating regions M1-M10 from FDX1 into FDX2, and we deleted the
343 conserved FDX2 C terminus (Δ C12; Fig. 6a; Extended data Fig. 8; Suppl. Table 3). Because
344 simple cell-based *in vivo* assays were not readily feasible to test restoration of FDX1-specific
345 functions in FDX2, we studied two specifically FDX1-catalyzed reactions *in vitro*, namely the
346 cytochrome P450-dependent cortisol synthesis from the 11-deoxycortisol precursor^{5,6} (Suppl.
347 Fig. 7) and lipoylation (cf. Fig. 3b,c). The various FDX2 mutant proteins were recombinantly
348 expressed and purified. They showed close to wild-type UV/Vis spectral properties, except for
349 FDX2-M7 which did not bind any Fe/S cluster, explaining why this segment was essential for
350 FDX2 function (see above; Extended data Fig. 9; Suppl. Fig. 6). Cortisol formation by
351 cytochrome P450 CYP11B1 was efficiently catalyzed by FDX1, but not by FDX2⁶ or several
352 mutant proteins (Fig. 6b,c). In contrast, mutant FDX2-R135E showed a strong (35-fold)
353 increase in cortisol formation compared to wild-type FDX2. Further small improvements were
354 seen by combining this site-specific exchange with the C-terminal deletion (FDX2-
355 R135E+ Δ C12) and segment M5 (FDX2-M5+R135E+ Δ C12) to yield a 42- and 45-fold
356 stimulation, respectively. Overall, up to 9% of the FDX1-catalyzed cortisol levels were
357 generated, showing that mutated FDX2 is now well, yet not fully capable of steroid production.
358 We conclude that FDX1-specific Glu133 represents a decisive functional element for steroid
359 formation, and the C-terminal truncation provides a further minor contribution. Interestingly, the

360 fungal-type *S. cerevisiae* Yah1 and *S. pombe* Etp1^{fd} also supported cortisol synthesis at 20-
361 60% efficiencies compared to FDX1-type human FDX1 and *Bos taurus* Adx^{50,51} (Fig. 6c).
362 These fungi contain only one FDX and do not possess mitochondrial cytochrome P450
363 enzymes⁵². However, both fungal proteins contain a negatively charged Asp at position
364 Glu133^{FDX1}, re-emphasizing the importance of negative charge at this position for steroid
365 formation.

366 Analogously, the FDX2 mutant proteins were assayed by FDX1-specific lipoylation (cf.
367 Fig. 3b,c). Surprisingly, FDX2-R135E did not significantly improve lipoylation compared to wild-
368 type FDX2 (Fig. 6d,e). Instead, the C-terminally truncated FDX2-ΔC12 variants (without or with
369 additional modifications) showed robust activity increases, enabling up to 49% 6-thiooctanoyl
370 and 21% lipoyl formation compared to FDX1, equivalent to a 6-10-fold increase over FDX2-
371 catalyzed levels. Thus, the C terminus of FDX2 appears to negatively interfere with LIAS
372 interaction, an effect also evident from the ca. 3-fold higher lipoylation (compared to FDX2) by
373 *S. cerevisiae* Yah1 hosting a shorter C terminus distinct from FDX2 (Fig. 6d,e; Extended data
374 Fig. 8). This result agrees well with the *in vivo* complementation experiments with LIAS-
375 humanized yeast (cf. Fig. 3d). Collectively, our mutational studies identify different regions of
376 FDX1 (highlighted in Fig. 6a) with importance for either cortisol or lipoyl formation, suggesting
377 that the FDX1 interaction sites with CYP11B1 and LIAS are not identical.

378

379

380 Discussion

381 In this work, we have expanded the knowledge on the physiological roles of the mitochondrial
382 ferredoxins FDX1 and FDX2 in human cells by identifying new targets of their electron transfer
383 function and by dissecting the molecular basis of their target specificity (Fig. 6f). A major finding
384 of our study was the identification of the so far unknown molecular function of FDX1 in lipoyl
385 cofactor biosynthesis, half a century after the discovery of its (i.e. adrenodoxin's) crucial role
386 in steroidogenesis⁵³. Strikingly, both FDX1 and FDX2 are involved in lipoylation, yet perform
387 distinct, non-overlapping roles. While FDX2 as a member of the core mitochondrial ISC
388 machinery assists the assembly of the [4Fe-4S] protein lipoyl synthase (LIAS)²², also *FDX1*
389 knockout cells showed a severe lipoylation defect. This was surprising, because FDX1 had no
390 detectable role in Fe/S protein maturation (Fig. 1). To elucidate the molecular basis of this
391 finding, we adapted an *in vitro* reconstitution system of bacterial lipoyl synthesis for the human
392 pathway⁴⁰, and show that FDX1 serves as a dedicated electron donor to kickstart the multi-
393 step radical SAM-dependent LIAS reaction (Fig. 3b,c). Frequently, the electron donors of
394 members of the large radical SAM enzyme family are ill-defined⁴¹, or the roles of electron
395 donors in either Fe/S cluster assembly or SAM reduction were not dissected⁵⁴. *Thermotoga*

396 *maritima* [4Fe-4S] FDXs reduce MiaB, and *E. coli* LipA employs a flavodoxin and its
397 reductase^{55,56}, clearly distinguishing mitochondrial and bacterial lipoylation pathways despite
398 the close evolutionary connection of mitochondria and bacteria¹⁰. During *in vitro* reconstitution
399 of lipoylation, FDX1 was more potent than the artificial reductant DT, yet could not efficiently
400 be replaced by FDX2. The physiological relevance of the FDX specificity was further verified
401 by a human LIAS-expressing yeast model in which endogenous Yah1 was depleted and
402 replaced by FDX1 and/or FDX2. FDX1 but not FDX2 was essential for efficient lipoylation (Fig.
403 3d,e). Structural modeling using AlphaFold showed a trimeric complex between LIAS, GCSH
404 (H-protein of the glycine cleavage system;³⁹) with the octanoyl-carrying loop reaching into the
405 substrate tunnel of LIAS, and FDX1 with its [2Fe-2S] cluster being close to the catalytic cluster
406 of LIAS consistent with direct electron transfer through a tunnel in LIAS (for details see Suppl.
407 Results; Extended data Fig. 10). In contrast, FDX2 could not be modeled in a physiologically
408 relevant position in keeping with its low activity in lipoylation.

409 FDX1 was further characterized to play a decisive role in COX maturation presumably
410 by facilitating heme *a* synthesis (Fig. 6f), similar to Yah1 in yeast or FdxA in *Trypanosomes*^{4,24}.
411 This function became only evident upon *FDX1* gene knockout, again suggesting that rather
412 low levels of FDX1 satisfy the cellular needs of heme *a* production (see Suppl. Discussion). In
413 part, the low FDX1 requirement may be due to a minor activity of FDX2 in this process⁶, a
414 notion supported by residual COX activity (33%) in human FDX1 knockout cells (Fig. 2b) and
415 our humanized COX15 yeast model expressing FDX1 and/or FDX2 (Fig. 2d). Collectively,
416 FDX1, besides its well-established role in steroidogenesis, shows a functional plasticity to
417 cooperate with multiple target electron acceptors. Even in steroidogenesis, FDX1 is known to
418 interact with multiple mitochondrial cytochrome P450 (CYP) enzymes⁵. The array of possible
419 CYP partners of FDX1 might even be extended under certain (sometimes pathological)
420 conditions^{57,58}. We conclude that mammalian FDX1 has evolved to a remarkably versatile
421 enzyme donating electrons to multiple acceptors in different physiological pathways.

422 In contrast, FDX2 mainly functions in Fe/S protein biogenesis. Given the importance of
423 cellular Fe/S protein biogenesis for life, this FDX2 function is essential for virtually all eukaryotic
424 cells and tissues¹¹, and is even required at two stages within the mitochondrial ISC machinery
425 (see Introduction). In contrast to earlier reports^{8,30}, we did not find any activity of FDX1 in
426 mitochondrial Fe/S protein maturation, neither *in vivo* in cells lacking FDX1 nor *in vitro* during
427 enzymatic [2Fe-2S] cluster reconstitution on ISCU2 (Fig. 1). This notion was further supported
428 by our yeast complementation experiments and the mutual interconversion of FDX1 and FDX2
429 functions (see below). The explanation for these contradictory findings remains unclear, yet a
430 possible reason may be the indirect consequences of both lipoylation and heme *a* synthesis
431 defects in FDX1-deficient cells on mitochondrial metabolism. Collectively, our study shows a
432 high specialization of each FDX for electron transfer to distinct redox partners.

433 The high target specificity of the two human FDXs raised the question of what structural
434 or other characteristics these enzymes utilize to discriminate their targets. The reduction
435 potentials of FDX1 and FDX2 at -267 and -342 mV, respectively, cannot readily explain why
436 FDX2 cannot cooperate with typical FDX1 substrates, fitting to a study with bacterial FDXs^{6,59}.
437 To explore structural criteria, we performed mutational studies to swap conspicuous regions
438 between the two FDXs, thereby gaining Fe/S protein biogenesis function of FDX1 or lipoyl and
439 cortisol synthesis activity of FDX2. In each case, only few selected regions showed a
440 substantial activity increase over the low residual activity of each FDX, yet full activity was not
441 reached showing the complexity of substrate selection. A prominent, up to 50-fold increased
442 cortisol formation activity was observed by replacing a positive by a negative residue in FDX2
443 (mutant R135E; Fig. 6b). While specific ion pair interactions might explain the FDX2 activity
444 gain, the charge inversion also changes the electronic surface potential above the [2Fe-2S]
445 cluster binding site to better resemble that of FDX1-type proteins (Suppl. Fig. 5). In a study of
446 bacterial Fdx-CYP interactions, a similar region was found critical for interaction with a large
447 number of CYP partners⁵⁹. Strongly negative surfaces can also be found for fungal Yah1 and
448 Etp1^{fd} (containing Asp at position 135) which primarily function in Fe/S protein and heme a
449 biogenesis^{7,24,27}, yet *in vitro* can perform CYP-dependent catalysis despite the absence of
450 endogenous mitochondrial CYPs in fungi (Fig. 6c)⁵⁰. Overall, negative ionic interactions and
451 surface potentials may contribute to the specificity of FDX1-type proteins in steroidogenesis.
452 Interestingly, a different structural determinant was critical for FDX1 specificity in lipoylation.
453 Deletion of the conserved C terminus (FDX2 mutant Δ C12) created substantially increased (up
454 to tenfold) lipoylation activity, while mutant R135E was not improved (Fig. 6d,e). This indicates
455 different specificity determinants for FDX1 binding to its dedicated targets CYPs and LIAS.
456 Conversely, FDX1 obtained a gain in FDX2 functionality (e.g., partial growth recovery of Yah1-
457 depleted yeast cells) upon transfer of the conserved C terminus of FDX2 (Fig. 5d-f). A minor
458 growth improvement was obtained by (additionally) exchanging various small segments (M7
459 with M3, M6) discriminating the two FDXs. This shows that a combination of several regions is
460 crucial to define FDX2 specificity in its major function, i.e. Fe/S protein biogenesis.

461 Here, we have studied known and novel physiological functions of the two human FDXs
462 and have defined the molecular basis of their specificities. The findings may open avenues for
463 engineering FDXs to function with diverse redox partners or as redox gates in synthetic biology
464 approaches⁶⁰. An interesting open question concerns the existence of additional FDX targets.
465 Prime candidates are other mitochondrial radical SAM enzymes such as CDK5RAP involved
466 in tRNA modification or RSAD1 with a still unclear function^{41,61}. Future studies may therefore
467 further expand the broad spectrum of FDX targets.

468

469

470 Acknowledgments

471 We thank Ralf Rösser and Sahra Hanschke for excellent technical assistance, Dr. Uwe Linne
472 and Jan Bamberger from the Core Facility 'Mass spectrometry and Elemental analysis' of
473 Philipps-Universität Marburg for mass spectrometry, and Dr. Frank Hannemann and Rita
474 Bernhardt (Saarbrücken) for help with cortisol formation assays. We acknowledge the
475 contribution of the Core Facility 'Protein Biochemistry and Spectroscopy' of the Philipps-
476 Universität Marburg. RL received generous financial support from Deutsche
477 Forschungsgemeinschaft (SPP 1927) and COST Action FeSBioNet (Contract CA15133). SJB
478 acknowledges support from the National Science Foundation (MCB-1716686) and the Eberly
479 Family Distinguished Chair in Science. SJB is an Investigator of the Howard Hughes Medical
480 Institute.

481

482

483 References

- 484 1. Campbell, I.J., Bennett, G.N. & Silberg, J.J. Evolutionary Relationships Between Low
485 Potential Ferredoxin and Flavodoxin Electron Carriers. *Frontiers in Energy Research*
486 **7**(2019).
- 487 2. Hanke, G. & Mulo, P. Plant type ferredoxins and ferredoxin-dependent metabolism. *Plant*
488 *Cell Environ* **36**, 1071-84 (2013).
- 489 3. Miotto, O. et al. Genetic architecture of artemisinin-resistant Plasmodium falciparum. *Nat*
490 *Genet* **47**, 226-34 (2015).
- 491 4. Changmai, P. et al. Both human ferredoxins equally efficiently rescue ferredoxin
492 deficiency in Trypanosoma brucei. *Mol Microbiol* **89**, 135-51 (2013).
- 493 5. Ewen, K.M., Ringle, M. & Bernhardt, R. Adrenodoxin--a versatile ferredoxin. *IUBMB Life*
494 **64**, 506-12 (2012).
- 495 6. Sheftel, A.D. et al. Humans possess two mitochondrial ferredoxins, Fdx1 and Fdx2, with
496 distinct roles in steroidogenesis, heme, and Fe/S cluster biosynthesis. *Proceedings of*
497 *the National Academy of Sciences of the United States of America* **107**, 11775-80
498 (2010).
- 499 7. Lange, H., Kaut, A., Kispal, G. & Lill, R. A mitochondrial ferredoxin is essential for
500 biogenesis of cellular iron-sulfur proteins. *Proc Natl Acad Sci U S A* **97**, 1050-5 (2000).
- 501 8. Shi, Y., Ghosh, M., Kovtunovych, G., Crooks, D.R. & Rouault, T.A. Both human
502 ferredoxins 1 and 2 and ferredoxin reductase are important for iron-sulfur cluster
503 biogenesis. *Biochimica et biophysica acta* **1823**, 484-92 (2012).
- 504 9. Weibert, H. et al. Functional reconstitution of mitochondrial Fe/S cluster synthesis on Isu1
505 reveals the involvement of ferredoxin. *Nat. Commun.* **5**, 5013 (2014).
- 506 10. Freibert, S.A. et al. Evolutionary conservation and in vitro reconstitution of
507 microsporidian iron-sulfur cluster biosynthesis. *Nat Commun* **8**, 13932 (2017).
- 508 11. Braymer, J.J., Freibert, S.A., Rakwalska-Bange, M. & Lill, R. Mechanistic concepts of
509 iron-sulfur protein biogenesis in Biology. *Biochim Biophys Acta Mol Cell Res* **1868**,
510 118863 (2021).

- 511 12. Kispal, G., Csere, P., Prohl, C. & Lill, R. The mitochondrial proteins Atm1p and Nfs1p are
512 essential for biogenesis of cytosolic Fe/S proteins. *EMBO J* **18**, 3981-9 (1999).
- 513 13. Boniecki, M.T., Freibert, S.A., Muhlenhoff, U., Lill, R. & Cygler, M. Structure and
514 functional dynamics of the mitochondrial Fe/S cluster synthesis complex. *Nat Commun*
515 **8**, 1287 (2017).
- 516 14. Van Vranken, J.G. et al. The mitochondrial acyl carrier protein (ACP) coordinates
517 mitochondrial fatty acid synthesis with iron sulfur cluster biogenesis. *Elife* **5**, e17828
518 (2016).
- 519 15. Kim, J.H., Frederick, R.O., Reinen, N.M., Troupis, A.T. & Markley, J.L. [2Fe-2S]-
520 Ferredoxin binds directly to cysteine desulfurase and supplies an electron for iron-sulfur
521 cluster assembly but is displaced by the scaffold protein or bacterial frataxin. *Journal of*
522 *the American Chemical Society* **135**, 8117-20 (2013).
- 523 16. Gervason, S. et al. Physiologically relevant reconstitution of iron-sulfur cluster
524 biosynthesis uncovers persulfide-processing functions of ferredoxin-2 and frataxin. *Nat*
525 *Commun* **10**, 3566 (2019).
- 526 17. Freibert, S.A. et al. N-terminal tyrosine of ISCU2 triggers [2Fe-2S] cluster synthesis by
527 ISCU2 dimerization. *Nat Commun* **12**, 6902 (2021).
- 528 18. Weiler, B.D. et al. Mitochondrial [4Fe-4S] protein assembly involves reductive [2Fe-2S]
529 cluster fusion on ISCA1-ISCA2 by electron flow from ferredoxin FDX2. *Proc Natl Acad*
530 *Sci U S A* **117**, 20555-20565 (2020).
- 531 19. Spiegel, R. et al. Deleterious mutation in FDX1L gene is associated with a novel
532 mitochondrial muscle myopathy. *Eur J Hum Genet* **22**, 902-6 (2014).
- 533 20. Paul, A. et al. FDXR Mutations Cause Sensorial Neuropathies and Expand the Spectrum
534 of Mitochondrial Fe-S-Synthesis Diseases. *Am J Hum Genet* **101**, 630-637 (2017).
- 535 21. Gurgel-Giannetti, J. et al. A novel complex neurological phenotype due to a homozygous
536 mutation in FDX2. *Brain* **141**, 2289-2298 (2018).
- 537 22. Lill, R. & Freibert, S.A. Mechanisms of Mitochondrial Iron-Sulfur Protein Biogenesis.
538 *Annu Rev Biochem* **89**, 471-499 (2020).
- 539 23. Zhang, Y. et al. Ferredoxin reductase is critical for p53-dependent tumor suppression via
540 iron regulatory protein 2. *Genes Dev* **31**, 1243-1256 (2017).
- 541 24. Barros, M.H., Carlson, C.G., Glerum, D.M. & Tzagoloff, A. Involvement of mitochondrial
542 ferredoxin and Cox15p in hydroxylation of heme O. *FEBS Lett* **492**, 133-8 (2001).
- 543 25. Bareth, B. et al. The heme a synthase Cox15 associates with cytochrome c oxidase
544 assembly intermediates during Cox1 maturation. *Mol Cell Biol* **33**, 4128-37 (2013).
- 545 26. Swenson, S.A. et al. From Synthesis to Utilization: The Ins and Outs of Mitochondrial
546 Heme. *Cells* **9**(2020).
- 547 27. Pierrel, F. et al. Involvement of mitochondrial ferredoxin and para-aminobenzoic acid in
548 yeast coenzyme Q biosynthesis. *Chem Biol* **17**, 449-59 (2010).
- 549 28. Ozeir, M. et al. Coenzyme Q biosynthesis: Coq6 is required for the C5-hydroxylation
550 reaction and substrate analogs rescue Coq6 deficiency. *Chemistry & biology* **18**, 1134-
551 42 (2011).
- 552 29. Yan, R., Adinolfi, S. & Pastore, A. Ferredoxin, in conjunction with NADPH and
553 ferredoxin-NADP reductase, transfers electrons to the IscS/IscU complex to promote
554 iron-sulfur cluster assembly. *Biochim Biophys Acta* **1854**, 1113-7 (2015).
- 555 30. Cai, K., Tonelli, M., Frederick, R.O. & Markley, J.L. Human Mitochondrial Ferredoxin 1
556 (FDX1) and Ferredoxin 2 (FDX2) Both Bind Cysteine Desulfurase and Donate Electrons
557 for Iron-Sulfur Cluster Biosynthesis. *Biochemistry* **56**, 487-499 (2017).

- 558 31. Zhu, J. & Thompson, C.B. Metabolic regulation of cell growth and proliferation. *Nat Rev*
559 *Mol Cell Biol* **20**, 436-450 (2019).
- 560 32. Tsvetkov, P. et al. Mitochondrial metabolism promotes adaptation to proteotoxic stress.
561 *Nat Chem Biol* **15**, 681-689 (2019).
- 562 33. Tsvetkov, P. et al. Copper induces cell death by targeting lipoylated TCA cycle proteins.
563 *Science* **375**, 1254-1261 (2022).
- 564 34. Campbell, I.J. et al. Recombination of 2Fe-2S Ferredoxins Reveals Differences in the
565 Inheritance of Thermostability and Midpoint Potential. *ACS Synth Biol* **9**, 3245-3253
566 (2020).
- 567 35. Brown, K.R., Allan, B.M., Do, P. & Hegg, E.L. Identification of novel hemes generated by
568 heme A synthase: evidence for two successive monooxygenase reactions. *Biochemistry*
569 **41**, 10906-13 (2002).
- 570 36. Moraes, C.T., Diaz, F. & Barrientos, A. Defects in the biosynthesis of mitochondrial
571 heme c and heme a in yeast and mammals. *Biochim Biophys Acta* **1659**, 153-9 (2004).
- 572 37. Antonicka, H. et al. Mutations in COX15 produce a defect in the mitochondrial heme
573 biosynthetic pathway, causing early-onset fatal hypertrophic cardiomyopathy. *Am J Hum*
574 *Genet* **72**, 101-14 (2003).
- 575 38. Sheftel, A.D. et al. The human mitochondrial ISCA1, ISCA2, and IBA57 proteins are
576 required for [4Fe-4S] protein maturation. *Mol Biol Cell* **23**, 1157-66 (2012).
- 577 39. Cronan, J.E. Assembly of Lipoic Acid on Its Cognate Enzymes: an Extraordinary and
578 Essential Biosynthetic Pathway. *Microbiol Mol Biol Rev* **80**, 429-50 (2016).
- 579 40. McCarthy, E.L. & Booker, S.J. Destruction and reformation of an iron-sulfur cluster
580 during catalysis by lipoyl synthase. *Science* **358**, 373-377 (2017).
- 581 41. Landgraf, B.J., McCarthy, E.L. & Booker, S.J. Radical S-Adenosylmethionine Enzymes
582 in Human Health and Disease. *Annu Rev Biochem* **85**, 485-514 (2016).
- 583 42. Nagai, M. et al. The oncology drug elesclomol selectively transports copper to the
584 mitochondria to induce oxidative stress in cancer cells. *Free Radic Biol Med* **52**, 2142-50
585 (2012).
- 586 43. Soma, S. et al. Elesclomol restores mitochondrial function in genetic models of copper
587 deficiency. *Proc Natl Acad Sci U S A* **115**, 8161-8166 (2018).
- 588 44. Hasinoff, B.B., Yadav, A.A., Patel, D. & Wu, X. The cytotoxicity of the anticancer drug
589 elesclomol is due to oxidative stress indirectly mediated through its complex with Cu(II).
590 *J Inorg Biochem* **137**, 22-30 (2014).
- 591 45. Modica-Napolitano, J.S., Bharath, L.P., Hanlon, A.J. & Hurley, L.D. The Anticancer
592 Agent Elesclomol Has Direct Effects on Mitochondrial Bioenergetic Function in Isolated
593 Mammalian Mitochondria. *Biomolecules* **9**(2019).
- 594 46. Yadav, A.A., Patel, D., Wu, X. & Hasinoff, B.B. Molecular mechanisms of the biological
595 activity of the anticancer drug elesclomol and its complexes with Cu(II), Ni(II) and Pt(II). *J*
596 *Inorg Biochem* **126**, 1-6 (2013).
- 597 47. Vallieres, C., Holland, S.L. & Avery, S.V. Mitochondrial Ferredoxin Determines
598 Vulnerability of Cells to Copper Excess. *Cell Chem Biol* **24**, 1228-1237 e3 (2017).
- 599 48. Mellor, S.B., Vavitsas, K., Nielsen, A.Z. & Jensen, P.E. Photosynthetic fuel for
600 heterologous enzymes: the role of electron carrier proteins. *Photosynth Res* **134**, 329-
601 342 (2017).
- 602 49. Mühlenhoff, U. et al. Cytosolic monothiol glutaredoxins function in intracellular iron
603 sensing and trafficking via their bound iron-sulfur cluster. *Cell Metab* **12**, 373-85 (2010).
- 604 50. Schiffler, B. et al. The adrenodoxin-like ferredoxin of *Schizosaccharomyces pombe*
605 mitochondria. *J Inorg Biochem* **98**, 1229-37 (2004).

- 606 51. Muller, J.J. et al. Structural and thermodynamic characterization of the adrenodoxin-like
607 domain of the electron-transfer protein Etp1 from *Schizosaccharomyces pombe*. *J Inorg*
608 *Biochem* **105**, 957-65 (2011).
- 609 52. Omura, T. & Gotoh, O. Evolutionary origin of mitochondrial cytochrome P450. *J Biochem*
610 **161**, 399-407 (2017).
- 611 53. Kimura, T. & Suzuki, K. Components of the electron transport system in adrenal steroid
612 hydroxylase. Isolation and properties of non-heme iron protein (adrenodoxin). *J Biol*
613 *Chem* **242**, 485-91 (1967).
- 614 54. Swift, R.P., Rajaram, K., Elahi, R., Liu, H.B. & Prigge, S.T. Roles of Ferredoxin-
615 Dependent Proteins in the Apicoplast of *Plasmodium falciparum* Parasites. *mBio*,
616 e0302321 (2022).
- 617 55. Arcinas, A.J., Maiocco, S.J., Elliott, S.J., Silakov, A. & Booker, S.J. Ferredoxins as
618 interchangeable redox components in support of MiaB, a radical S-adenosylmethionine
619 methylthiotransferase. *Protein Sci* **28**, 267-282 (2019).
- 620 56. Cicchillo, R.M. et al. Lipoyl synthase requires two equivalents of S-adenosyl-L-
621 methionine to synthesize one equivalent of lipoic acid. *Biochemistry* **43**, 6378-86 (2004).
- 622 57. Lehnerer, M., Schulze, J., Bernhardt, R. & Hlavica, P. Some properties of mitochondrial
623 adrenodoxin associated with its nonconventional electron donor function toward rabbit
624 liver microsomal cytochrome P450 2B4. *Biochem Biophys Res Commun* **254**, 83-7
625 (1999).
- 626 58. Liao, W.L., Dodder, N.G., Mast, N., Pikuleva, I.A. & Turko, I.V. Steroid and protein ligand
627 binding to cytochrome P450 46A1 as assessed by hydrogen-deuterium exchange and
628 mass spectrometry. *Biochemistry* **48**, 4150-8 (2009).
- 629 59. Zhang, W. et al. Mechanistic Insights into Interactions between Bacterial Class I P450
630 Enzymes and Redox Partners. *ACS Catalysis* **8**, 9992-10003 (2018).
- 631 60. Atkinson, J.T. et al. Metalloprotein switches that display chemical-dependent electron
632 transfer in cells. *Nat Chem Biol* **15**, 189-195 (2019).
- 633 61. Wei, F.Y. et al. Cdk5rap1-mediated 2-methylthio modification of mitochondrial tRNAs
634 governs protein translation and contributes to myopathy in mice and humans. *Cell Metab*
635 **21**, 428-42 (2015).
- 636 62. Heigwer, F., Kerr, G. & Boutros, M. E-CRISP: fast CRISPR target site identification. *Nat*
637 *Methods* **11**, 122-3 (2014).
- 638 63. Ran, F.A. et al. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**,
639 2281-2308 (2013).
- 640 64. Sherman, F. Getting started with yeast. *Methods Enzymol.* **350**, 3-41 (2002).
- 641 65. Gietz, R.D. & Woods, R.A. Transformation of yeast by lithium acetate/single-stranded
642 carrier DNA/polyethylene glycol method. *Methods Enzymol* **350**, 87-96 (2002).
- 643 66. Diekert, K., deKroon, A.I.P.M., Kispal, G. & Lill, R. Isolation and sub-fractionation of
644 mitochondria from the yeast *Saccharomyces cerevisiae*. *Methods Cell Biol.* **65**, 37-51
645 (2001).
- 646 67. McCarthy, E.L. & Booker, S.J. Biochemical Approaches for Understanding Iron-Sulfur
647 Cluster Regeneration in *Escherichia coli* Lipoyl Synthase During Catalysis. *Methods*
648 *Enzymol* **606**, 217-239 (2018).
- 649 68. Harlow, E. & Lane, D. *Using Antibodies: A Laboratory Manual*, (Cold Spring Harbor
650 Laboratory, Cold Spring Harbor, NY, 1998).
- 651 69. Biederbick, A. et al. Role of human mitochondrial Nfs1 in cytosolic iron-sulfur protein
652 biogenesis and iron regulation. *Mol. Cell. Biol.* **26**, 5675-5687 (2006).

- 653 70. Stehling, O. et al. Investigation of iron-sulfur protein maturation in eukaryotes. in
654 *Methods in Molecular Biology*, Vol. 372 (eds. Leister, D. & Herrmann, J.M.) 325-342.
655 (Humana Press Inc., Totowa, NJ, 2007).
- 656 71. Stehling, O., Sheftel, A.D. & Lill, R. Chapter 12 Controlled expression of iron-sulfur
657 cluster assembly components for respiratory chain complexes in mammalian cells.
658 *Methods Enzymol* **456**, 209-31 (2009).
- 659 72. Sambrook, J. & Russell, D.W. *Molecular cloning: A laboratory manual*, (Cold Spring
660 Harbor Press, Cold Spring Harbor, NY, 2001).
- 661 73. Gueldener, U., Heinisch, J., Koehler, G.J., Voss, D. & Hegemann, J.H. A second set of
662 loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast.
663 *Nucleic Acids Res* **30**, e23 (2002).
- 664 74. Mühlenhoff, U., Richhardt, N., Ristow, M., Kispal, G. & Lill, R. The yeast frataxin
665 homolog Yfh1p plays a specific role in the maturation of cellular Fe/S proteins. *Hum Mol*
666 *Genet* **11**, 2025-36 (2002).
- 667 75. Molik, S., Lill, R. & Mühlenhoff, U. Methods for studying iron metabolism in yeast
668 mitochondria. *Methods Cell. Biol.* **80**, 261-280 (2007).

669

670

671 **Materials and Methods**

672 **Small interfering RNAs and guide RNAs**

673 Sets of three Silencer Select small interfering RNAs (siRNAs) directed against the mRNA of
674 FDX1 or FDX2⁶ were purchased from ThermoFisher Scientific (Waltham, USA). FDX1- and
675 FDX2-directed guide RNA sequences for CRISPR-Cas9-mediated gene knockout (Suppl.
676 Table 1) were designed using E-CRISP⁶² and cloned into plasmid PX459 (Addgene,
677 Watertown, MA, USA) according to Ref. ⁶³.

678

679 **Tissue culture and cell transfection**

680 Human embryonic kidney (HEK) 293 cells were cultured in high glucose DMEM with standard
681 supplements and transfected with siRNA by electroporation⁶ as specified in Supplementary
682 Materials and Methods. PX459-based plasmids allowing for simultaneous expression of gRNA,
683 Cas9, and the puromycin resistance marker were chemically transfected into HEK293 cells
684 using the JetPrime transfection reagent (Polyplus, Illkirch, France) according to the
685 manufacturer's instructions. In order to facilitate chemical transfection, cells were grown in
686 collagenized flasks, and transfectants were selected by puromycin (10 µg/mL) for two to three
687 days. The day of puromycin removal was considered as day zero of gene knockout, and cells
688 were cultured as lines until use for up to 17 days.

689

690 **Yeast strains, cell growth and plasmids**

691 Yeast strains and plasmids used in this study are listed in Suppl. Table 4 and 5. Cells were
692 cultivated either in minimal medium containing all recommended supplements (SC) or in rich
693 medium (YP) plus the following carbon sources: 2% w/v glucose (SD/YPD), 2% w/v galactose
694 (SGal/YPGal), 3% w/v glycerol (SGly/YPGly), 2% w/v lactate plus 0.05% w/v glucose
695 (SLac/YPLac) or 2% w/v raffinose plus 0.2% w/v galactose (SRaff + Gal)⁶⁴. Yeast cells were
696 transformed using the lithium acetate method⁶⁵.

697

698 **Yeast complementation assays with human genes**

699 For FDX1/2 interconversion experiments, Gal-YAH1 cells⁷ were transformed with plasmids
700 encoding yeast or human FDXs (p426-FDX1, p426-FDX2, p416-Yah1 and corresponding
701 FDX1/2 mutant plasmids). Single colonies were grown on SGal plates and resuspended in
702 water to OD₆₀₀ 0.5. Serial dilutions (1:5) were spotted onto plates containing different growth
703 media and incubated at 30°C for three days. Enzyme activity measurements of isolated yeast
704 mitochondria were performed as described previously⁶⁶.

705 For *in vivo* lipoyl and heme *a* formation experiments, Gal-YAH1-*lip5Δ* and Gal-YAH1-
706 *cox15Δ* cells were transformed with plasmids p416-FDX1, p415-FDX2, p414-LIAS and p414-
707 COX15. Transformants were grown overnight in SD medium, centrifuged, and cell pellets
708 diluted in water to OD₆₀₀ 0.1. Serial dilutions (1:5) were spotted onto plates containing different
709 growth media and incubated at 30°C for three days.

710

711 **Reconstitution of *de novo* Fe/S cluster synthesis on ISCU2**

712 *In vitro* enzymatic reconstitution of [2Fe-2S] cluster formation on ISCU2 was performed as
713 described^{9,10}. Ferredoxin titration reactions were prepared in an anaerobic chamber and
714 contained 0.5 mM NADPH, 0.8 mM sodium-ascorbate, 0.3 mM FeCl₂, 0.2 mM MgCl₂, 5 mM
715 GSH, 75 μM ISCU2, 7.5 μM NIA, 7.5 μM FXN, 1 μM FDXR and FDX1 or FDX2 as indicated at
716 a final volume of 300 μL in degassed reconstitution buffer (50 mM Tris/HCl pH 8.0, 150 mM
717 NaCl). Reactions were transferred to a CD-spectrometer (Jasco J-815) in a tightly sealed
718 cuvette containing a magnetic stirring bar, and the CD-signal at 431 nm was monitored at 20°C.
719 After 30 s, Fe/S cluster synthesis was initiated by injecting 0.5 mM cysteine into the sealed
720 cuvette and a time course recorded for 10 min. Time courses were normalized to the signal
721 intensity before cysteine addition.

722

723

724

725 **Chemical Fe/S cluster reconstitution on LIAS**

726 Chemical reconstitution of LIAS was done under strictly anaerobic conditions. His-affinity
727 chromatography purified LIAS was diluted to 0.1 mM in degassed LIAS buffer (50 mM HEPES
728 pH 7.5, 300 mM KCl and 20% w/v glycerol) on ice. 10 mM DTT was added, and the mixture
729 was incubated for 1 h. Subsequent addition of 0.8 mM ferric ammonium citrate (FAC) and 0.8
730 mM Li₂S, both added slowly whilst mixing gently, led to a deep brown-yellowish color.
731 Spectroscopic analysis at different time points after reconstitution revealed that the cluster
732 content of LIAS reaches its maximum within 30 min. LIAS was re-purified after 30 min
733 reconstitution by size exclusion chromatography using a HiLoad® 16/600 Superdex® 75 pg
734 column. The holoprotein was characterized as described previously⁶⁷ by UV/Vis-spectroscopy
735 (Jasco V-550), Bradford assay, amino acid analysis, SDS-PAGE, and iron and sulfide
736 determination.

737

738 **Lipoyl formation assay**

739 Experiments were done under strictly anaerobic conditions⁴⁰. Degassed reaction buffer (50
740 mM HEPES pH 7.5, 300 mM KCl, 10% w/v glycerol) was supplied with 10 mM MgCl₂, 2 mM
741 NADPH and 0.5 mM octanoyl peptide (Glu-Ser-Val-[N⁶-octanoyl]Lys-Ala-Ala-Ser-Asp).
742 Optionally, up to 2 mM CuCl₂ and/or elesclomol (Ele) were added. Ele:Cu inhibition assays
743 contained 2% v/v DMSO. Control reactions showed LIAS activity not being altered at
744 concentrations of 0–4% v/v DMSO. 20 μM FDXR, 140 μM FDX and 1 mM S-
745 adenosylmethionine (SAM) were added subsequently. The reaction was started by adding 35
746 μM reconstituted LIAS to a final volume of 25 μL and incubated at 23°C for 2.5 h, if not noted
747 otherwise. Reactions were quenched by adding 100 mM H₂SO₄, 4 mM TCEP and 40 μM of an
748 internal peptide standard (Pro-Met-Ser-Ala-Pro-Ala-Arg-Ser-Met). The amounts of octanoyl-,
749 6-thiooctanoyl- and lipoyl-peptide were quantified by HPLC-MS as described previously⁶⁷.

750

751 **Miscellaneous methods**

752 Specific rabbit antisera against human FDX1 and FDX2 were raised using recombinant full-
753 length proteins^{6,68}. For other primary antibodies see Suppl. Table 6. Peroxidase- (Biorad,
754 Germany) or biotin- (Vector Laboratories, Burlingame, USA) conjugated goat anti-rabbit and
755 anti-mouse antibodies as well as the ABC system (Vector Laboratories, Burlingame, USA)
756 were used as secondary reagents. The following published methods were used: Digitonin-
757 based fractionation of HEK293 cells, activity measurements for aconitase, succinate
758 dehydrogenase, cytochrome c reductase; cytochrome c oxidase, citrate synthase, and lactate
759 and malate dehydrogenase⁶⁹⁻⁷¹; manipulation of DNA and PCR⁷²; gene disruptions and

760 promoter exchanges in yeast^{73,74}; preparation of yeast mitochondria and cell extracts⁶⁶;
761 determination of enzyme activities in yeast extracts⁷⁵; immunological techniques⁶⁸; *in vitro*
762 cytochrome c reduction by FDXs¹⁸. Measurements of replicates were performed using distinct
763 samples. Statistical analyses were performed via ordinary one-way analyses of variance
764 (ANOVA) with multiple comparisons using Prism software.
765

766 **Figure legends**

767

768 **Fig. 1 FDX1 is not involved in mitochondrial Fe/S protein assembly.** **a** HEK293 cells were
769 transfected without (control) or with scrambled (SCR) non-targeting siRNA, or with pools of
770 *FDX1*- or *FDX2*-directed siRNAs (si) at a three-day interval⁶. Cells were harvested 9 days after
771 the first transfection. Cell extracts were analyzed by immunostaining for the indicated
772 mitochondrial proteins. Observed molecular masses (kDa) for proteins are given in
773 parentheses. C-I, C-II, C-III, respiratory complexes I, II, and III; C-V, F_1F_0 ATP synthase. **b**
774 HEK293 cells were chemically transfected with the empty plasmid PX459 (control) or with
775 plasmids containing three different *FDX1*-directed gRNAs (CC1 to CC3). Transfected cells
776 were selected by puromycin for 3 days. Samples harvested 8 days after antibiotic removal
777 were subjected to immunostaining of the indicated mitochondrial proteins. For quantitation see
778 Extended data Fig. 1b. **c** Total aconitase (Aco) and succinate dehydrogenase (SDH) enzyme
779 activities were determined in extracts from cells described in (**b**). Results were expressed as
780 a ratio to citrate synthase (CS) activity, and presented relative to the respective values of
781 control cells (set to 100%, dashed line; $n \geq 6$). **d** Enzymatic reconstitution of [2Fe-2S] cluster
782 formation on ISCU2 in the presence of indicated amounts of *FDX1* or *FDX2* was measured by
783 CD spectroscopy. Initial rates were determined by linear fitting of the kinetic traces (Extended
784 data Fig. 2; $n = 3$; one-way ANOVA for control without versus with *FDX* reactions, $*p < 0.05$,
785 $***p < 0.001$). Error bars indicate the SEM. Representative blots are shown.

786

787 **Fig. 2 The role of human ferredoxins in heme a and CoQ₁₀ biosynthesis.** **a** HEK293 cell
788 extracts from Fig. 1b were subjected to immunostaining of indicated cytochrome c oxidase
789 (complex IV, C-IV) subunits. VDAC1 served as loading control (see Fig. 1b). Observed
790 molecular masses (kDa) for proteins are given in parentheses. Representative blots are
791 shown. **b** COX activity (relative to CS activity) was analyzed in total membrane fractions
792 derived from cells treated as in Fig. 1c ($n \geq 3$; one-way ANOVA for control versus *FDX1*-CC1-3,
793 $***p < 0.001$). **c** Gal-*YAH1-cox15* Δ yeast cells were transformed with plasmids encoding
794 human *COX15* and additionally *FDX1*, *FDX2* or no *FDX* (–) as indicated. Subsequently, cells
795 initially grown in liquid SC glucose medium were plated as a 1:5 dilution series, and grown for
796 3 days at 30°C on the indicated minimal media. **d** COX activities (relative to malate
797 dehydrogenase, MDH) were determined in extracts from cells of part (**c**) grown for 3 days in
798 liquid SD medium ($n \geq 4$; one-way ANOVA for empty vector control versus *FDX*-complemented
799 cells, $***p < 0.001$). **e** HEK293 cells transfected as in Fig. 1b were harvested within the
800 indicated time intervals after puromycin selection. Total cellular coenzyme Q₁₀ (CoQ₁₀)
801 content per total cellular protein was determined and presented relative to the value of control
802 cells ($n \geq 4$). **f** HEK293 cells were chemically transfected with the *FDX1*-directed gRNA-

803 encoding plasmid CC2 or with the PX459 control plasmid. Six to seven days after puromycin
804 selection cells were repeatedly transfected by electroporation at a three-day interval with a
805 pool of *FDX2*-directed siRNAs, and harvested up to 9 days after the first transfection similar to
806 Fig. 1a. At each transfection round cell samples were removed for the analysis of total cellular
807 CoQ₁₀ content as in (e) ($n \geq 3$). CoQ₁₀ levels were normalized to amounts before starting RNAi
808 depletion. Error bars show the SEM.

809

810 **Fig. 3 FDX1 starts the radical chain reaction of lipoyl biosynthesis.** a HEK293 cell samples
811 from Fig. 1b were immunostained for the indicated proteins or the lipoyl cofactor. VDAC1
812 served as loading control (see Fig. 1b). Observed molecular masses (kDa) for proteins are
813 given in parentheses. Representative blots are shown. b, c Formation of the (b) 6-thiooctanoyl
814 and (c) lipoyl peptide from a peptide substrate analogue [Glu-Ser-Val-(N6-octanoyl)Lys-Ala-
815 Ala-Ser-Asp] by LIAS. Samples contained 0.5 mM peptide substrate, 35 μ M LIAS, 2 mM
816 NADPH, 20 μ M FDXR, 140 μ M FDX1, and 1 mM SAM, unless indicated otherwise. Dithionite
817 (DT) samples lacked FDX1, FDXR, and NADPH. Reactions were incubated at 23°C, quenched
818 with acid after 2.5 h, and products quantified by HPLC-MS using peptide standards ($n \geq 3$; one-
819 way ANOVA for FDX1 reactions versus all other conditions). d Gal-*YAH1-lip5* Δ yeast cells
820 harboring plasmids encoding human LIAS, FDX1 and/or FDX2 as indicated were grown in
821 minimal medium with glucose (Glc) or galactose (Gal) as indicated for 3 days. Mitochondria
822 were isolated and analyzed for the indicated proteins or lipoyl cofactor in PDH and KGDH by
823 immunostaining. Porin (Por1) was used as loading control. e Mitochondrial extracts were
824 analyzed for aconitase (Aco1) and KGDH activities, and results were normalized to MDH
825 activity. Wild-type (WT) and *lip5* Δ cells served as controls ($n \geq 4$; one-way ANOVA, LIAS-
826 complemented Gal-*YAH1-lip5* Δ cells versus all other Gal-*YAH1-lip5* Δ cells). * $p < 0.05$, *** $p <$
827 0.001. Error bars indicate the SEM. n.d., not detectable.

828

829 **Fig. 4 Inhibitory effect of elesclomol-copper on lipoyl synthesis *in vitro*.** a,b Formation
830 and quantitation of the 6-thiooctanoyl- (a) and lipoyl- (b) peptide was as in Fig. 3b,c in the
831 presence of the indicated concentrations of CuCl₂ and/or elesclomol (Ele) ($n = 2$; one-way
832 ANOVA for control versus Ele/Cu titrations, ** $p < 0.01$, *** $p < 0.001$). n.d., not detectable. c
833 Reduction of cytochrome *c* (80 μ M) by 1 mM NADPH, 2 nM FDXR, and 0.1 μ M FDX1 in
834 samples containing CuCl₂ and/or Ele as indicated. Cytochrome *c* reduction was monitored at
835 550 nm for 60 s. Error bars indicate the SEM ($n \geq 3$; one-way ANOVA, *** $p < 0.001$). d,e CD
836 spectra of FDX1 (d) and FDX2 (e) recorded after incubation with the indicated amounts of
837 CuCl₂ and/or Ele (see also Suppl. Fig. 3). f HEK293 cells were grown in the presence of the
838 indicated amounts of Ele and/or Cu for 72 h before harvesting and immunostaining of the

839 indicated proteins or lipoyl cofactor. Observed molecular masses (kDa) for proteins are given
840 in parentheses. Representative blots are shown.

841

842 **Fig. 5 Identification of structural elements crucial for functional specificity of FDX2. a**

843 Superposition of crystal structures of human FDX1 (gold, PDB code: 3P1M, residues 65–170)
844 and FDX2 (green, PDB code: 2Y5C, residues 69–174). The PheGly insertion in FDX1 (FG
845 loop) is shown in magenta, and the orientation of helix C is indicated by rods. The C terminus
846 is not resolved in the FDX2 structure and corresponding FDX1 residues are not shown. **b**

847 Electrostatic surface potential calculated by the APBS server
848 (<https://server.poissonboltzmann.org>) mapped to human ferredoxin surfaces (orientation as in

849 **a**; color bar covers the range from -10 kT/e to +10 kT/e). The Fe/S cluster binding sites are
850 marked by yellow circles. Red, negative charges; blue, positive charges. **c** Structure of FDX1

851 (PDB code: 3P1M, residues 65–184) highlighting regions exchanged with the respective FDX2
852 sequences. The bottom table provides the residues introduced from FDX2 into FDX1 for

853 functional interconversion. **d** Complementation of *Gal-YAH1* yeast with plasmids encoding
854 FDX2, FDX1 variants depicted in **c** or no FDX (-). Cells were grown for 3 days on minimal

855 medium (S) with glucose (Glc) or galactose (Gal) using a dilution series of resuspended cells
856 initially grown on SGal plates. Dashed lines separate different agar plates. **e** Aconitase (Aco)

857 and **f** catalase (Cat) activities were determined in extracts from *Gal-YAH1* cells complemented
858 with the indicated constructs. Cells were grown for 40 h in liquid SGlc medium prior to analysis.

859 Error bars indicate the SEM ($n = 3$; one-way ANOVA for FDX1-complemented yeast versus all
860 other conditions, $*p < 0.05$, $***p < 0.001$).

861

862 **Fig. 6 Conversion of FDX2 to generate FDX1 functionality. a** Structure of FDX2 (PDB code:
863 2Y5C, residues: 69–174) highlighting regions exchanged with the respective FDX1 sequences.

864 The bottom table provides the residues introduced from FDX1 into FDX2 for functional
865 interconversion. The site of the C-terminal truncation ($\Delta C12$) is marked by an arrow because

866 the C terminus is not resolved in the crystal structure. **b, c** Cortisol formation by different FDXs
867 (**b**) or the indicated FDX2 variants (**c**) for 10 min at 37°C. Reactions contained 11-

868 deoxycortisol, human CYP11B1, and the electron transfer chain NADPH, FDXR, and
869 ferredoxin. Cortisol content was quantified by HPLC and normalized to FDX1 reactions. (Yah1,

870 Etp1(fd), FDX2 R135E mutants: $n = 2$, Adx: $n = 3$, all other reactions: $n = 4$). **d, e** Synthesis of
871 6-thiooctanoyl (**d**) and lipoyl (**e**) by LIAS in the presence of different FDXs or indicated FDX2

872 variants. Formation and quantitation was as in Fig. 3b,c. Values were normalized to FDX1
873 reactions ($n = 3$). Error bars indicate the SEM. One-way ANOVA for FDX2 reactions versus all

874 other conditions was conducted in **b, c, d, e** ($**p < 0.01$, $***p < 0.001$). **f** Model showing the
875 distinct functions of human FDX1 and FDX2 in mitochondrial metabolism. FDX2 performs a

876 specific function in both *de novo* [2Fe-2S] cluster assembly by the core ISC complex and the
877 IBA57-ISCA1-ISCA2-dependent reductive fusion of two glutaredoxin GLRX5-derived [2Fe-2S]
878 to [4Fe-4S] clusters^{18,22}. FDX1 cannot replace FDX2 in these functions, yet is shown here to
879 perform two crucial functions in addition to the long-known mitochondrial cytochrome P450
880 (CYP)-dependent steroid transformation⁵. First, FDX1 serves as an electron donor to initiate
881 the radical SAM-dependent biosynthesis of the lipoyl cofactor by the [4Fe-4S] protein lipoyl
882 synthase (LIAS). Lipoyl is a cofactor in, e.g., pyruvate and 2-ketoglutarate dehydrogenase
883 (PDH, KGDH) complexes. The LIAS reaction is a sensitive target of the toxic elesclomol-
884 copper (Ele:Cu) anti-cancer drug. Second, FDX1 is involved in cytochrome oxidase (COX)
885 heme *a* formation by the COX15 enzyme, yet FDX2 can partially replace this FDX1 function.
886 Heme *a* is exclusively present in cytochrome *c* oxidase. Different from yeast mitochondria, a
887 non-FDX oxidoreductase (oxred) is required for the biosynthesis of coenzyme Q (CoQ) by the
888 ubiquinone (UBI) complex.
889

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5










Figure 6




Extended data for

**Functional spectrum and structural specificity
of mitochondrial ferredoxins FDX1 and FDX2**

Vinzent Schulz, Somsuvro Basu, Sven-A. Freibert, Holger Webert, Linda Boß,
Ulrich Mühlenhoff, Fabien Pierrel, Lars-O. Essen, Douglas M. Warui, Squire Booker,
Oliver Stehling, and Roland Lill



Extended data Figure 1 | Growth phenotypes and mitochondrial Fe/S protein status of *FDX1* knockout cells. **a** HEK293 cell lines were subjected to CRISPR-Cas9 *FDX1* gene knockout using CC1-CC3 guide RNAs. Cumulative growth of cells treated as in Fig. 1b was calculated from cell counts at various harvesting time points after puromycin removal. In detail, individual cell lines were sub-cultured by harvesting an entire culture vessel and subsequent re-seeding of a defined cell number into a new culture device. Remaining cells were collected, and the total protein content of this sample was determined. This protein amount was then used as a denominator for the calculation of the specific (i.e. total protein-related) enzymes activities measured in the respective sample. This tissue culture regime involving the re-seeding of aliquots of harvested cells produced sufficient cell material to conduct multiple analyses, even in case of an experimentally elicited growth retardation. Based on the cell counting performed at each harvest, the cumulative growth was calculated for each cell line from the total cell yield as well as the portion of cells needed for re-seeding during sub-culturing. Values are presented relative to those of control cells (set to 100%, dashed line; $n \geq 4$; mean \pm SEM). **b** Densitometric quantification of immunostains from cells treated as in Fig. 1b, 2a, and 3a (i.e. harvested 7 or 8 days after puromycin removal) using Image studio lite 5.2. Error bars indicate SEM ($n \geq 4$). **c-e** Total aconitase (Aco), succinate dehydrogenase (SDH) and citrate synthase (CS) activities were determined in cell samples obtained at the indicated time points after puromycin removal as in Fig. 1c. Values were presented relative to those of control cells (set to 100%, dashed line; $n \geq 3$; mean \pm SEM).







Extended data Figure 2 | Even high excess of FDX1 does not support enzymatic [2Fe-2S] cluster reconstitution on ISCU2. Enzymatic synthesis of [2Fe-2S] clusters on ISCU2 *in vitro* was monitored by the CD signal at 431 nm for 10 min. Reactions included NIA, FXN, FDXR and the indicated equivalent amounts (eq.) of (a) FDX2 or (b) FDX1 relative to NIA. Representative experiments are shown and initial rates are presented in Fig. 1d.

a

11 days after puromycin chase


14 days after puromycin chase

**b****c****d**

Extended data Figure 3 | *FDX1* knockout cells are deficient in cytochrome oxidase activity. **a**


Tissue culture media of HEK293 control and *FDX1* knockout cells (cf. Figs. 2a and Extended data Fig. 1a) were collected at the indicated time points. Cells were harvested, counted, and processed as required. The color change of the pH indicator phenol red in the culture medium from reddish toward yellowish, particularly for *FDX1*-CC1 and -CC2 knockout cells, revealed a pH drop despite lower cell numbers in these cultures (given at the top; in millions). **b** Cytochrome c oxidase (COX) activities were determined in total membrane fractions from cells harvested at the indicated time points after puromycin removal (cf. Fig. 2b). COX activities were expressed relative to citrate

synthase (CS) activities (Extended data Fig. 1e), and values are presented relative to those of control cells (set to 100%, dashed line; $n \geq 3$; mean \pm SEM). **c** HEK293 cells (see Fig. 2f) were chemically transfected with control plasmid PX459 or the PX459-derived plasmid FDX1-CC2 and selected by puromycin for 3 days as in Fig. 2a. 10 to 15 days after selection cells were transiently transfected by electroporation with plasmids coding for either a PEST-destabilized EGFP or FDX1, each fused to the N-terminal mitochondrial Su9 targeting sequence (from *Neurospora crassa* subunit 9 of mitochondrial F₁F₀ ATP synthase). Three days after transfection cells were harvested and analyzed for COX and CS activities. Data are presented relative to the values for Su9-EGFP control cells ($n = 4$; SEM). **d** Cell samples from (**c**) were subjected to immunostaining against the indicated proteins. Observed molecular masses (kDa) for proteins are given in parentheses. Representative blots are shown.





Extended data Figure 4 | Combined *FDX1* deletion and *FDX2* depletion elicits severe defects in growth and mitochondrial Fe/S proteins. **a** Cumulative growth of HEK293 cells from Fig. 2f was calculated from cell counts at the three harvests on days 3, 6 ($n \geq 4$ each), and 9 ($n = 2$) after the first electroporation. Values were presented relative to those of mock control cells (no CRISPR, no RNAi treatment; set to 100%, dashed line; mean \pm SEM). **b** HEK293 cells were transfected with *FDX1*-directed gRNA-encoding plasmids (CC1 to CC3) and subsequently with *FDX2*-directed

siRNAs similar to Fig. 2f. Cell samples were obtained at the specified time points and subjected to immunostaining of the indicated mitochondrial proteins or lipoyl cofactor. The observed molecular weights are given in parentheses. C-I, C-II, C-III, respiratory complexes I, II, and III. C-V, F₁F_o ATP synthase. The immunoblot signals from cell samples with combined FDX1-CC2 and FDX2-si deficiency are representative for at least three independent experiments, and are here presented conjointly with FDX1-CC1 / FDX2-si and FDX1-CC1 / FDX2-si treated cells, respectively.





Extended data Figure 5 | The lipoylation defect in *FDX1* knockout cells can be complemented by *FDX1*. HEK293 cells from Extended data Fig. 3c,d knocked out for *FDX1* (by FDX1-CC2 gRNA) and complemented with Su9-EGFP-PEST or Su9-FDX1 plasmids were subjected to immunostaining against the indicated proteins or lipoyl cofactor. Observed molecular masses (kDa) for proteins are given in parentheses. Representative blots are shown.




Extended data Figure 6 | *In vitro* synthesis of 6-thiooctanoyl intermediate and lipoyl product by human lipoyl synthase LIAS requires ferredoxin FDX1 as an electron donor. a Model of the multi-step reaction mechanism of lipoyl formation by human lipoyl synthase (LIAS) based on bacterial LipA¹. The LIAS enzyme contains two [4Fe-4S] clusters, the catalytic and auxiliary cluster,


needed for reductive cleavage of *S*-adenosylmethionine (SAM) and as a source for sulfur insertion into the octanoyl precursor, respectively. (1) The reaction starts with the binding of SAM and the octanoyl substrate which is covalently attached to a lysinyl residue of a lipoyl carrier domain (LCD) of the H protein of the glycine cleavage system (GCS)^{1,2}. An electron donor (orange) transfers a single electron to the catalytic cluster which mediates reductive cleavage of SAM to methionine and a 5'-deoxyadenosyl radical (5'-dA[•]). This work identified human FDX1 as the physiological electron donor, efficiently replacing dithionite that typically is used as an artificial electron donor¹. (2) The radical abstracts a hydrogen atom from the octanoyl C6 carbon, forming 5'-deoxyadenosine (5'-dAH). (3) The octanoyl C6 carbon in turn forms a covalent bond with a sulfur atom of the auxiliary cluster, concomitant with partial degradation of the cluster. (4) A second SAM molecule binds to LIAS, and upon electron supply from FDX1 (or DT) again leads to 5'-dA[•] radical formation by the catalytic cluster and abstraction of a proton from the terminal C8 carbon of the octanoyl moiety. (5) A second sulfur atom is covalently attached to the thiooctanoyl molecule, (6) leading to formation of the mature lipoyl cofactor and further degradation of the Fe/S cluster. To enable multiple reaction cycles, the auxiliary cluster must be regenerated by a still unclear mechanism. In humans, the mature lipoyl cofactor is finally transferred to the target proteins, e.g., the E2 subunits of pyruvate (PDH) and α -ketoglutarate (KGDH) dehydrogenase complexes. BCKDH, branched-chain ketoacid dehydrogenase; OADH, 2-oxoadipate dehydrogenase³. **b** Time courses of 6-thiooctanoyl intermediate and lipoyl product formation in FDX1-catalyzed reactions (see Fig. 3b,c). Samples included 0.5 mM peptide substrate, 35 μ M LIAS, 2 mM NADPH, 20 μ M FDXR, FDX1 as indicated and 1 mM SAM. Formation of the 6-thiooctanoyl intermediate proceeded significantly faster than lipoyl formation, indicating the second sulfur insertion step to be rate-limiting under the experimental conditions. The result suggested to record the data of other experiments after 150 min incubation.

a**b**


Extended data Figure 7 | Differential growth complementation of human LIAS-containing Gal-YAH1-lip5Δ cells by human FDX1 and FDX2 reflects their distinct functions. Gal-YAH1-lip5Δ yeast cells were transformed with a plasmid encoding human *LIAS* plus vectors containing no gene (-), *FDX1* and/or *FDX2* as indicated. **a** Cells were grown in liquid media for 4 days. Optical density (OD) at 600 nm was measured every 30 min. Growth in minimal lactate medium (SLac) is indicated by dashed lines, in rich yeast peptone lactate (YPLac) medium by solid lines. Grey dashed lines indicate standard errors ($n = 4$) using a microplate reader. **b** Serial dilutions of the indicated yeast strains were spotted onto agar plates containing minimal (S) or yeast peptone rich (YP) medium plus the indicated carbon sources. Plates were incubated at 30°C for 3 days. The growth results fit to distinct functions of human FDX1 and FDX2 in lipoylation/heme *a* synthesis and Fe/S protein biogenesis, respectively. Complementation of the LIAS-expressing Gal-YAH1-lip5Δ cells with FDX2 but not FDX1 supported growth due to Fe/S protein biogenesis restoration. The residual growth on non-fermentable carbon sources (lactate or glycerol) was due to the leaky *GAL* promoter allowing residual amounts of Yah1, and hence lipoate/heme *a*, being produced. Growth under these conditions was increased to normal levels by the combined expression of both human FDXs because FDX1 regenerated synthesis of both lipoate and heme *a*, and FDX2 supported Fe/S protein biogenesis.



Extended data Figure 8 | Multi-sequence alignment of mitochondrial ferredoxins. The multi-sequence alignment was generated by Multalin⁴. Secondary structure elements of the human FDX2 structure are shown above the alignment according to PROMOTIF⁵. Numbering is according to the full-length sequences of human (Hs) FDX1 and FDX2 retrieved from Uniprot (<https://www.uniprot.org>). The Fe/S cluster-coordinating cysteine residues are highlighted in yellow. Altered residues/regions of human FDX1 and FDX2 are highlighted in grey (see also Suppl. Table S3). Residues/regions mutated for interconversion of FDX1 and FDX2 functions (mutants M3, M5, M6, M7, R135E and C-terminal deletion/exchange) are additionally highlighted by colored boxes for animal FDX1/2-type sequences. Names of organisms are colored according to FDX-type: animal FDX2 (red), mitosomal (black), plant (green), fungal (orange), animal FDX1 (blue) and bacterial (grey). Two *Trypanosoma brucei* (Tb) FDXs best align between fungal and FDX1-type (FdxA) proteins or between FDX1-type and bacterial (FdxB) proteins⁶. FDXs from the following organisms were used (sequence identifiers in brackets): *Homo sapiens* (NP_001026904.2, NP_004100.1), *Canis lupus familiaris* (XP_038284595.1, XP_038367022.1), *Bos taurus* (NP_001073695.1, NP_851354.1), *Rattus norvegicus* (NP_001101472.1, NP_058822.2), *Mus musculus* (NP_001034913.1, NP_032022.1), *Danio rerio* (NP_001070132.1, XP_001922722.2), *Drosophila melanogaster* (NP_001189075.1, NP_647889.2), *Encephalitozoon cuniculi* GB-M1 (NP_585988.1), *Trachipleistophora hominis* OX=72359 GN=THOM_0371 PE=4 SV=1 (L7K0F4_TRAHO), *Arabidopsis thaliana* (NP_001329852.1), *Populus alba* (XP_034922198.1), *Oryza sativa Japonica Group* (XP_015647182.1), *Chaetomium thermophilum* var. *thermophilum* DSM 1495 (XP_006692961.1), *Chaetomium globosum* CBS 148.51 (XP_001225251.1), *Neurospora crassa* OR74A (XP_958085.1), *Aspergillus fumigatus* Af293 (XP_747954.1), *Yarrowia lipolytica* CLIB122 (XP_500417.1), *Saccharomyces cerevisiae* S288C (NP_015071.1), *Schizosaccharomyces pombe* Etp1^{fd} (2WLB_A), *Trypanosoma brucei brucei* (XP_845713.1 and XP_844647.1), *Magnetospirillum magnetotacticum* (WP_009867486.1), *Rhodospirillum rubrum* (WP_200292119.1), *Rickettsia prowazekii* (WP_004595967.1), *Yersinia pestis* (WP_015683702.1), *Shigella flexneri* (EHF1065008.1), *Salmonella enterica* (WP_001124473.1).



Extended data Figure 9 | Site-directed mutagenesis identifies FDX2 regions that functionally distinguish from FDX1. Gal-YAH1 yeast cells were transformed with vectors expressing the indicated FDXs and variants (see Supplementary Table 5). Growth of serial cell dilutions on minimal medium (S) agar plates containing glucose or galactose was at 30°C for 3 days. FDX1-FDX2-discriminating regions critical for the *in vivo* function of human FDX2 were identified using FDX2 variants in which amino acids were exchanged to those of FDX1. A loss of cell growth identified the respective altered region as being important for FDX2 function. The broken lines separate independent plates.



Extended data Figure 10 | Alphafold2 predicts a functionally meaningful structure for the trimeric LIAS-GCSH-FDX1 complex. Structures of human LIAS, GCSH, and FDX1, and of complexes were predicted with Google Colab and the top-scoring prediction is displayed (https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/beta/AlphaFold2_advanced.ipynb). **a** Overlay of the predicted LIAS structure (blue, residues: 29-372) with *Mycobacterium tuberculosis* LipA (PDB code: 5EXK, green). The *Mt* LipA catalytic (cat.) and auxiliary (aux.) cluster are displayed as spheres and its bound 6-thiooctanoyllysyl (thiooct) moiety as grey sticks. **b-c** Modeling of the LIAS-GCSH-FDX1 complex (residues: LIAS, 29-372; GCSH, 41-173; FDX1, 54-184) predicted a functionally meaningful complex, where GCSH inserts its lipoyl-carrying residue Lys107 into the octanoyl substrate entry tunnel of LIAS. The position of the [2Fe-2S] cluster of FDX1 is consistent with electron transfer to the catalytic [4Fe-4S] cluster of LIAS. Fe/S cluster-coordinating residues and GCSH Lys107 are shown as sticks. Positions of Fe/S clusters are modelled based on crystal structures of FDX1 (PDB code: 3P1M) and *Mt* LipA (PDB code: 5EXK). LIAS, blue; GCSH, grey; FDX1, gold. In **(c)** the electrostatic surface potential is mapped onto the half transparent surface of LIAS. When we instead modeled a trimeric LIAS-GCSH-FDX2 complex, binding of FDX2 was predicted at the substrate tunnel of LIAS. This would block substrate-product delivery by GCSH. GCSH interacted with FDX2, but hardly with LIAS in this case. Overall, no physiologically relevant complex could be modeled in presence of FDX2. **d-f** Electrostatic surfaces mapped onto LIAS **(d)**, FDX1 **(e)** and GCSH **(f)**. LIAS-FDX1 and LIAS-GCSH interacting regions are encircled by yellow and black dashed lines, respectively. The location of the FDX1 [2Fe-2S] cluster and helix F are encircled by yellow and light blue solid lines, respectively. Surface potentials were calculated using the APBS server (<https://server.poissonboltzmann.org>). Negative charges are colored in red, positive charges in blue. The color bar covers the range from -10 kT/e to +10 kT/e. N and C termini are indicated.

Extended data References

1. McCarthy, E.L. & Booker, S.J. Destruction and reformation of an iron-sulfur cluster during catalysis by lipoyl synthase. *Science* **358**, 373-377 (2017).
2. Landgraf, B.J., McCarthy, E.L. & Booker, S.J. Radical S-Adenosylmethionine Enzymes in Human Health and Disease. *Annu Rev Biochem* **85**, 485-514 (2016).
3. Solmonson, A. & DeBerardinis, R.J. Lipoic acid metabolism and mitochondrial redox regulation. *J Biol Chem* **293**, 7522-7530 (2018).
4. Corpet, F. Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res.* **16**, 10881-10890. (1988).
5. Hutchinson, E.G. & Thornton, J.M. PROMOTIF--a program to identify and analyze structural motifs in proteins. *Protein Sci* **5**, 212-20 (1996).
6. Changmai, P. et al. Both human ferredoxins equally efficiently rescue ferredoxin deficiency in *Trypanosoma brucei*. *Mol Microbiol* **89**, 135-51 (2013).

Supplementary information for

Functional spectrum and structural specificity of mitochondrial ferredoxins FDX1 and FDX2

Vinzent Schulz, Somsuvro Basu, Sven-A. Freibert, Holger Webert, Linda Boß,
Ulrich Mühlenhoff, Fabien Pierrel, Lars-O. Essen, Douglas M. Warui, Squire Booker,
Oliver Stehling, and Roland Lill

Content:

Supplementary Results

Supplementary Discussion

Supplementary Figures 1-7

Supplementary Materials and Methods

Supplementary Tables 1-6

Supplementary References

Supplementary Results

Crystal structure of FDX1 and 3D structural features discriminating it from FDX2

The 3D structure of mature FDX2 (residues 66-171; PDB ID: 2Y5C) was solved *a priori* by single-wavelength anomalous diffraction (SAD) phasing using the anomalous scattering of the [2Fe-2S] iron ions (Suppl. Table 2). Consistent with other [2Fe-2S] FDXs, only a C-terminally truncated protein formed crystals diffracting to 1.7 Å. FDX2 displays a compact ($\alpha+\beta$) fold typical for vertebrate-type mitochondrial FDXs (Fig. 5a). Significant structural differences between the known FDX1 structure (PDB ID: 3P1M) and that of FDX2 were observed only for the FDX2 loop between helix C and the Fe/S cluster binding site, with the Phe-Gly (FG) dipeptide of FDX1 missing in FDX2 (Fig. 5a; Extended data Fig. 8; Suppl. Fig. 4). The lack of FG leads to a distortion of helix C by 15° compared to FDX1, resulting in a more compact structure and different surface shape around the helix-loop region. These subtle differences alone were unlikely to provide a satisfactory explanation for the striking specificity of the two human FDXs. We therefore compared the surface potentials¹ of both FDXs and found that FDX2 possesses a less negatively charged area around the Fe/S cluster binding site, including helix F, which is known to participate in partner protein recognition²⁻⁴ (Fig. 5b; Suppl. Fig. 5). Most conspicuously, exchanges of Asp91^{FDX1} by His95^{FDX2} in helix C and Glu133^{FDX1} by Arg135^{FDX2} in helix F are the major contributions to the altered surface potential at the redox-active site of these human FDXs. However, since these positively charged residues of FDX2 are not fully conserved in organisms with two FDXs, there may be other specificity-discriminating structural factors (Extended data Fig. 8). In conclusion, the 3D structures of FDX1 and FDX2 display only minor differences in backbone geometry and partner interaction surfaces, with a different surface potential as a discriminating feature.

Structural prediction of the trimeric LIAS-GCSH-FDX1 complex

The structure of LIAS in a catalytic heterotrimeric complex with the physiological octanoyl-substrate carrier GCSH (H protein of glycine cleavage system⁵) and the electron donor FDX1 was modelled using Google Colab, a software based on AlphaFold^{6,7}. Predicted folds of GCSH and FDX1 are almost identical to published crystal structures of bovine GCSH (PDB code: 3KLR) and human FDX1 (PDB code: 3P1M), excluding the disordered N- and C-terminal regions. Apart from the N terminus, the predicted fold of LIAS and positioning of its Fe/S cluster-coordinating residues are similar to published crystal structures of *Mycobacterium tuberculosis* (*Mt*) and *Thermosynechococcus elongatus* (*Te*) LipA (PDB codes: 5EXJ, 5EXK, 5U0O, 5U0P)^{8,9}. Similarity is highest to a crystallographic snapshot of lipoyl synthesis showing *Mt* LipA with a 6-thiooctanoyl intermediate bound at the auxiliary cluster site (Extended data Fig. 10a).

Intriguingly, the predicted trimeric LIAS-GCSH-FDX1 complex exhibits GCSH and FDX1 suitably positioned to fulfil their respective functions (Extended data Fig. 10b). The side chain of Lys104^{GCSH} functioning as octanoyl-carrier is inserted into the substrate binding tunnel of LIAS, resembling the positioning of the 6-thiooctanoyllysyl intermediate as observed in *Mt* LipA; the respective Lys ϵ -amino moieties of both structures are displaced by 1.9 Å⁸. FDX1 binds LIAS via its cluster-coordinating loop and helix F, as reported previously for complexes of FDX1 with FDXR, CYP11A1 and CYP11B2^{2,3,10}. The Fe/S cluster is located above a LIAS tunnel potentially involved in electron transfer to the catalytic Fe/S cluster of LIAS (Extended data Fig. 10c). The predicted distance between the Fe ions of the FDX1 and LIAS clusters of 16.2 Å might allow for direct electron transfer¹¹. Notably, both *Mt* and *Te* LipA crystal structures exhibit a similar tunnel (see PDB codes: 5EXK and 5U0P), yet other structures of the same two proteins show the tunnel entrance being blocked by rearrangement of residues of the catalytic cluster-coordinating loop (PDB codes: 5EXJ and 5U0O). It is currently unknown whether the opening or closing of the tunnel might represent a means of regulating the electron flow.

In the predicted trimeric LIAS-GCSH-FDX1 complex, LIAS-binding regions of both GCSH and FDX1 are strongly negatively charged (red) and contact positively charged regions (blue) of LIAS, suggesting that electrostatic interactions may be crucial for binding (Extended data Fig. 10d-f). The region between both aforementioned LIAS tunnels exhibits the most positive surface potential of the protein.

Removal of the FDX2 C terminus was shown in this work to enable lipoylation to some extent (14.2% of FDX1 reactions, Fig. 6e). As this may suggest the FDX2 C terminus to interfere with LIAS interaction, it is interesting to note that the FDX1 C terminus makes slight contact to LIAS in the predicted structure. Modelling the trimeric complex with FDX2 WT or the FDX2 Δ C12 variant instead of FDX1 did not yield any physiologically relevant quaternary structure, in line with FDX1 being the dedicated redox partner of LIAS. In summary, the predicted structure of the trimeric LIAS-GCSH-FDX1 complex is compatible with the experimental data presented in this work and may support future investigation of the biosynthetic mechanism of lipoyl formation.


Supplementary Discussion

The results from both the RNAi-based *FDX1* knockdown and the humanized yeast system also resolve the old conundrum of conflicting *FDX1* expression levels in mammalian cells. Transcriptomics data suggest that *FDX1* is expressed in virtually all cell types, yet the FDX1 protein was detectable by immunostaining solely in tissues associated with high steroid production such as adrenal gland, testis, and placenta (Suppl. Fig. 1)¹². The importance of

lipoylation in virtually all cell types necessitates and explains the global expression of *FDX1*, yet low lipoyl synthesis capacity, i.e. low *FDX1* or *LIAS*, suffices to satisfy the needs for normal mitochondrial metabolism. In keeping with this idea, low levels of *FDX1* remaining after RNAi-based knockdown maintained lipoylation in human cells (Fig. 1a). Likewise, small residual amounts of yeast *Yah1* were sufficient to produce enough ectopic *LIAS* for normal lipoylation by *FDX1* (Fig. 3d).

FDX1 has been identified in a systematic study to be a target of the tight complex of the Cu ionophore Ele with Cu, leading to the proposal that the role of *FDX1* in Fe/S protein maturation might explain the Ele:Cu toxicity¹³⁻¹⁵. This view has been revised and extended by work published during review of our study to show that both *FDX1* and lipoylation might be the target of Ele:Cu toxicity, yet a molecular explanation was not provided¹⁶. Our definition of the physiological *FDX1* functions, particularly in lipoylation and not in Fe/S protein maturation, renders it unlikely that *FDX1* is the primary target of Ele:Cu toxicity *in vivo*, because *FDX1* is sensitive to high Cu but not Ele:Cu (Fig. 4c-e). However, *FDX1* could be damaged by the Cu accumulated in cells and mitochondria in the presence of Ele¹⁴. This seems unlikely though, because *FDX1* and *FDX2* show similar Cu sensitivity, yet their Fe/S protein biogenesis is unaffected by Ele (Fig. 4d-f; Suppl. Fig. 3). Here, we provide *in vitro* and *in vivo* evidence that the complex process of lipoylation, most prominently the final step of lipoyl formation by *LIAS*, is severely affected by both Ele:Cu and Cu (Fig. 4a,b,f). The lipoyl-dependence of the mitochondrial enzymes PDH and KGDH satisfactorily explains why particularly cells critically relying on mitochondrial metabolism are most sensitive to Ele treatment¹⁵. An interesting future question will be the deeper mechanistic dissection of how Ele:Cu affects the biochemical function of *LIAS*. Overall, our findings identify lipoylation as a highly Cu-sensitive pathway in human cells, distinguishing these cells from both fungi hosting highly Cu-sensitive Fe/S proteins and bacteria containing Cu-sensitive hydratases with solvent-exposed [4Fe-4S] clusters^{17,18}.

Supplementary Figures




Supplementary Figure 1 | Tissue-specific protein and mRNA levels for human mitochondrial FDX1 and FDX2. Expression analyses for the two human FDXs by either specific protein immunostaining or transcript measurements within the Human Protein Atlas project (HPA; see below)¹⁹ provide contradictory results for FDX1. The high expression of FDX1 RNA and protein in testis, placenta, adrenal gland and kidney is well explained by its long-known role in steroid conversion. The findings of our work now satisfactorily explain the general expression of the *FDX1* gene in all tissues as judged by transcript analyses. The newly described function of FDX1 in lipoylation represents a central biosynthetic reaction that is needed in virtually all tissues.


a Semi-quantitative estimation of FDX1 and FDX2 protein levels by immunostaining using commercially obtained tissue lysates (see Fig. S1 in Ref. ¹²) or a systematic tissue distribution

analysis within the HPA project showed detectable levels of *FDX2* in almost all analyzed tissues, whereas *FDX1* was found only in testis, placenta, adrenal gland and kidney.


b On the contrary, significant amounts of both *FDX1* and *FDX2* mRNAs were detected within the HPA project across numerous tissues, with *FDX2* levels normally being higher than those of *FDX1*. The latter expectedly is expressed most strongly in steroid-transforming organs. Quantitation of *FDX1* and *FDX2* mRNA levels is presented as protein-coding transcripts per million (pTPM) across 43 tissues/cell types. Credit: HPA project (<https://www.proteinatlas.org/ENSG00000137714-FDX1/tissue> and <https://www.proteinatlas.org/ENSG00000267673-FDX2/tissue>, release 01.10.2020). Notably, the transcript results are not fully consistent with independent datasets for tissue-specific RNA levels obtained in projects Genotype-Tissue Expression (GTEx) and Functional Annotation Of The Mammalian Genome (FANTOM5; see above websites for details).



Supplementary Figure 2 | RNAi-mediated *FDX1* and *FDX2* knockdown phenotypes. a Cumulative growth of control, *FDX1*-depleted (*FDX1*-si), and *FDX2*-depleted cells treated as in Fig. 1a was calculated from cell counts at the three harvests on days 3, 6, and 9 after the first transfection. **b** Densitometric quantitation of immunoblots of Fig. 1a using Image studio lite 5.2. Values were presented relative to those of control cells (set to 100%, dashed line in a).




Supplementary Figure 3 | Elesclomol prevents copper-mediated Fe/S cluster destruction on FDX1 and FDX2. Samples containing 140 μM FDX1 or FDX2 were titrated with CuCl_2 or elesclomol (Ele) under anaerobic conditions, and CD spectra of the [2Fe-2S] clusters were recorded after each titration step. Samples containing 1.5 mM Ele were subsequently titrated with CuCl_2 . Excess of Ele prevented Cu-mediated destruction of Fe/S clusters on FDX1 and FDX2 by Ele:Cu complex formation (see also Fig. 4d,e). Representative data of two independent experiments per condition.







Supplementary Figure 4 | Comparison of the 3D structures of different ferredoxins.

Depicted are the crystal structures of **a** FDX1 (PDB code: 3P1M, residues 65 – 184), **b** FDX2 (2Y5C, 69 – 174), **c** *Bt* Adx (1CJE, 63 – 169), **d** *Ec* Fdx (1I7H, 2 – 110), and **e** *Sp* Etp^{fd} (2WLB, 518 – 603), as well as **f** the NMR solution structure of *Sc* Yah1 (2MJD, 58 – 172). For the crystal structures, C-terminal residues are resolved only for FDX1. The PheGly dipeptide present in FDX1-like proteins (FG loop) is depicted in magenta.





Supplementary Figure 5 | The electrostatic surface potential of FDX2 is less negatively charged. Depicted are *Bt* Adx (PDB code: 1CJE, residues 63-169), FDX1 (3P1M, 65-184), FDX2 (2Y5C, 69-174), *Ec* Fdx (1I7H, 2-110), *Sp* Etp^{fd} (2WLB, 518-603) and *Sc* Yah1 (2MJD, 58-160). Surface potentials were calculated using the APBS server (<https://server.poissonboltzmann.org>). Negative charges are colored in red, positive charges in blue. The color bar covers the range from -10 kT/e to +10 kT/e. All proteins are shown in three different orientations, defined on top by the ribbon presentations of *Bt* Adx (grey). In the left column the position of the Fe/S cluster binding site is marked by yellow circles. For *Sc* Yah1, the NMR structure was employed, as no crystal structure was available. The C-terminal residues are only shown for FDX1. The structure of FDX2 R135E variant was modelled using UCSF Chimera²⁰. The exchanged residue is indicated by an arrow.





e


	Yah1	His-FDX1	FDX1	FDX1 Δ C12	His-FDX1 M5	His-FDX1 M7
c (mg/mL)	18	32	28	17	6	6




f

	His-FDX2	FDX2	FDX2 Δ C12	FDX2 M5	FDX2 M5 Δ C12	FDX2 M7
c (mg/mL)	15	21	30	7	5	3





Supplementary Figure 6 | Characterization of purified mitochondrial ferredoxins. The indicated mitochondrial FDXs were purified after recombinant expression in *E. coli*. **a-d, g, h** UV/VIS spectra were recorded and normalized to 62 μ M. Spectra were shifted by 0.1 AU_{700nm} along the y-axis for better visualization. **e, f, i, j** Photographs were taken for FDX solutions at the indicated protein concentrations (top). Purified proteins (5 μ g each) were analyzed for purity by SDS-PAGE (bottom). Typically, apoproteins lacking a Fe/S cluster showed poorer purity. Where indicated, proteins possessed an N-terminal His₆-tag. All other proteins were devoid of tags.



Supplementary Figure 7 | Reaction scheme for cortisol formation by cytochrome P450 CYP11B1. The monooxygenase CYP11B1 (also named β -hydroxylase) introduces a hydroxyl group into 11-deoxycortisol in the final step of cortisol formation. The required reduction of molecular oxygen is enabled by delivery of two electrons from FDX1²¹.

Supplementary Materials and Methods

Transfection of HEK293 cells by electroporation

HEK293 cells were harvested by trypsination, and 9×10^6 cells were resuspended in 250 μL of electroporation buffer²². 250 pmol of each of three individual siRNA were added, yielding a total of 750 pmol or 3 μM siRNA, and transfection by electroporation was performed by a Xcell Gene Pulser device (Biorad, München, Germany) using a pulse of 262 V and 525 μF with exponential decay. Cell recovery was improved by addition of up to 5 μg of the pVA-I plasmid to the transfection^{23,24}. Immediately after electroporation cells were transferred into 20 mL of high-glucose DMEM supplemented with 7.5% FCS, penicillin/streptomycin, and glutamine, and seeded into collagenized 75 cm^2 flasks. In order to prolong the time period of RNAi-mediated mRNA depletion, cells were re-transfected twice at a three-day interval, resulting in a total depletion time of 9 days.

Coenzyme Q extraction and HPLC analysis

Cell pellets (~1mg protein) were processed and analyzed essentially as described²⁵. Briefly, we added to the cell pellets 0.3 mL of 150 mM KCl, 200 μL glass beads, 10 μL of a 5 μM CoQ₈ standard solution and 3 mL methanol. The tubes were vortexed for 1 min, 2 mL petroleum ether (40 – 60 °C boiling range) was added, and vortex was repeated for 1 min. After centrifugation at 700 rpm for 1 min, the upper phase was collected, and the methanol phase was extracted again with 2 mL petroleum ether. Both petroleum ether phases were combined, dried under a nitrogen flow and the lipid extracts were resuspended in 100 μL ethanol. Samples corresponding to 0.2 mg protein were injected onto the C18 column, and separation was obtained at a flow rate of 1 mL/min with a mobile phase composed of 25% isopropanol, 45% methanol, 20% ethanol, and 10% of a solution composed of 90% (v/v) isopropanol, 10% (v/v) 1 M ammonium acetate, and 0.1% (v/v) formic acid. A precolumn electrode (5020 Guard Cell, Thermo) was set at +650 mV and electrochemical detection (ECD) was performed with a Coulochem III equipped with a 5011A analytical cell (E1, -650 mV ; E2, +650 mV). Standard solutions of CoQ₈ and CoQ₁₀ were injected in the same conditions to generate standard curves that were used to quantify CoQ₁₀ in the samples and to correct for sample loss during the organic extraction (based on the recovery of CoQ₈).

Expression and purification of recombinant proteins

An overview of the plasmid constructs used for heterologous protein expression is given in Supplementary Table 5. CYP11B1²⁶, FDXR²⁷ and LIAS²⁸ were expressed and purified as described before, with the exception that LIAS was overproduced in terrific broth instead of M9

medium. For all other proteins, plasmids were introduced into *E. coli* BL21 (DE3) cells and overproduced in terrific broth medium using appropriate antibiotics (100 µg/mL ampicillin or 30 µg/mL kanamycin). Cultures were inoculated from bacterial overnight suspensions and shaken at 37°C and 160 rpm until OD₆₀₀ ~0.8. pASK-IBA43(+) constructs were induced with 0.2 mg/L anhydrotetracycline and pET-vectors with 1 mM isopropyl-β-D-thiogalactoside (IPTG). For NIA production, two constructs encoding NFS1-ISD11 and ACP were co-expressed. Additional supplements were added for production of all FDXs (50 µM Fe(NH₄)₂(SO₄)₂) and NIA (2 mM pyridoxine hydrochloride). Subsequently, cultures were shaken at 22°C overnight and cell pellets harvested by centrifugation. Purification was done aerobically at 4°C or on ice. His-tagged constructs were purified by His-affinity and non-tagged constructs by anion exchange chromatography (AEC). Pellets were resuspended in His-buffer (35 mM Tris/HCl pH 7.4, 300 mM NaCl, 5% w/v glycerol) or AEC-buffer (35 mM Tris/HCl pH 7.5, 50 mM NaCl, 5% w/v glycerol), respectively. Subsequent to the addition of proteinase inhibitor (cOmplete™ Protease Inhibitor Cocktail) and optionally lysozyme and/or DNase I, cells were lysed by sonication. The lysate was cleared by centrifugation and the cell extract loaded onto either a His-binding (NiNTA-Agarose) or an anion exchange column (Source 30Q) pre-equilibrated with His- or AEC-buffer. His-binding columns were washed using His-buffer plus 10 mM (NIA) or 70 mM (ISCU2, FXN) imidazole and proteins eluted in His-buffer plus 250 mM imidazole. Anion exchange columns were washed with AEC-buffer and proteins eluted using a gradient from AEC- to high salt buffer (35 mM Tris/HCl pH 7.5, 1 M NaCl, 5% w/v glycerol). Eluates were concentrated to a volume of 2 mL and further purified by size exclusion chromatography using HiLoad® 16/600 Superdex® columns. Desired fractions were combined, concentrated to ~1 mL and checked for purity by SDS-PAGE. Protein concentration was determined using the Bradford assay. A correction factor for concentrations determined by the Bradford assay was determined for FDX1, FDX2, LIAS, ISCU2, FXN and NIA by amino acid analysis (Leibniz-Institut für Analytische Wissenschaften, Dortmund, Germany). Iron and sulfide content of Fe/S cluster proteins were estimated by colorimetric methods^{29,30}.

***In vitro* Fe/S cluster stability assay of ferredoxins**

FDX1 or FDX2 were diluted to a final volume of 300 µL in CD-buffer (50 mM Tris/HCl pH 7.5, 150 mM NaCl, 5% w/v glycerol) in an anaerobic chamber and transferred to a CD-cuvette which was sealed tightly. CD-spectra (300 – 650 nm) were recorded at 20°C and the proteins subsequently titrated with elesclomol and/or CuCl₂. For each addition of the reagents, the cuvette was transferred back to the anaerobic chamber to sustain anaerobicity.

FDX2 crystallization, data collection, structure determination and refinement

As the full length FDX2 did not yield crystals even without a His₆-tag, we designed a C-terminally truncated construct assuming that flexibility of the C-terminus is precluding crystal formation. We also found that the first N-terminal residues of FDX2 are not important for *in vivo* function (Extended data Fig. 9), hence for crystallization we N-terminally truncated the human FDX2 protein to the same length as the crystallized bovine Adx^{31,32}. The truncated FDX2 protein covered residues 66 – 171 and yielded crystals after one day. Crystals diffracted to 1.7 Å and the structure could be solved *a priori* by single-wavelength anomalous diffraction (SAD) phasing using the anomalous scattering of the iron atoms of the [2Fe-2S] cluster. Data collection and refinement statistics are summarized in Supplementary Table 2. The final model of FDX2 covered two *holo*-proteins within one asymmetric unit with all residues visible except the most likely cleaved start methionine.

Initial crystallization screening was done with the His-tagged version, the untagged full length and the untagged C-terminal truncated version of human FDX2 using commercially available screens (QIAGEN) with a Cartesian 4000 dispensing system (Genomic Solutions). Only the untagged C-terminal truncated human FDX2 protein yielded crystals in different conditions with either ammonium sulfate or sodium citrate as precipitant within one to a few days. Optimized crystals were grown by mixing 300 nL of purified protein solution (30 mg/ mL) with 300 nL reservoir solution and incubation against 80 µL reservoir solution at 4°C in the sitting drop setup. The optimized reservoir solution contained solely 1.6 M sodium citrate. For data collection, a single red-brown crystal of the *holo*-protein was flash-frozen in liquid nitrogen with 20% w/v glycerol as cryo protectant. Native data set from this single crystal was collected at 100 K at the European Synchrotron Radiation Facility (ESRF, Grenoble), beamline ID14-4, on a MAR CCD detector to a resolution of 1.70 Å. The data were processed using XDS and the CCP4 package³³. The crystal belonged to be the primitive monoclinic space group P2₁ and contained two molecules in the asymmetric unit with molecular masses of 11937 Da resulting in an apparent V_m of 2.79 Å³/ Da and a solvent content of 55.9%³⁴.

The crystal structure of the human FDX2 protein was determined by the Single-wavelength Anomalous Diffraction (SAD) technique using the anomalous signal of the iron from the Fe/S cluster. Initial phasing with SHELX-CDE³⁵ clearly showed four anomalous sites as expected for two FDX2 *holo*-proteins per asymmetric unit each containing a [2Fe-2S] cluster. Automated model building using ARP/wARP³⁶ yielded a model containing 202 of the 216 amino acids of the two molecules. Missing residues could be manually added into additional electron density using COOT³⁷ and refinement with REFMAC³⁸. Several rounds of refinement provided a final model covering residues 3 to 109 in both chains, 220 water molecules, two [2Fe-2S] cluster and two sulfate ions ($R_{\text{cryst}} = 17.28\%$, $R_{\text{free}} = 20.05\%$). In the final model the residues E40 in chain A and Q16 in chain B were modeled in alternative conformations. The side chains of some charged residues (E58, D62, E83, K106, R109) were flexible and could only be modeled

with lower or zero occupancy due to missing electron density. The model quality was analyzed with PROCHECK³⁹ and PROMOTIF⁴⁰. The quality of the final model is summarized in Supplementary Table 2. The structure factors have been deposited in the Protein Data Bank, www.PDB.org (PDB code: 2Y5C).

The human FDX1 structure was generated by homology modeling using the bovine adrenodoxin structure (PDB code: 1CJE_B) as template with SWISS MODEL⁴¹. Figures were generated using UCSF Chimera.

***In vitro* cortisol formation assay**

In vitro hydroxylation of 11-deoxycortisol to cortisol was performed using human mitochondrial CYP11B1 as described previously²⁶. The reaction mixture contained 0.5 μ M FDXR, 0.5 μ M CYP11B1, 400 μ M 11-deoxycortisol, 20 μ M FDX and a NADPH regenerating system consisting of 1 mM MgCl₂, 5 mM glucose-6-phosphate and 1 U glucose-6-phosphate dehydrogenase. The reaction was started by the addition of 1 mM NADPH and incubated at 37°C for 10 min. The reaction was stopped and the steroid compounds were organically extracted and analyzed by reversed phase HPLC.

Supplementary Tables

Supplementary Table 1: Guide RNA sequences used in this study

gRNA	cDNA sequence
FDX1-CC1	GCGGCCTGCTGAGGAACCGG
FDX1-CC2	GTGACAGGTTGAACAAGCCA
FDX1-CC3	GATGGCTTTGGTGCATGTGA
FDX2-CC1	GGTCTTCACTCACATACACA
FDX2-CC2	GCTGGAGAGGAGGACGCGGG
FDX2-CC3	GGCCAGCGGATCCCAGTGAG

Supplementary Table 2: Crystal structure of human FDX2 – data collection and refinement statistics

Data collection	
Wavelength, Å	1.319
Space group	P 1 2 1
Cell dimensions	
<i>a</i> , <i>b</i> , <i>c</i> , Å	51.38, 35.45, 75.51
β, °	105.12
Resolution range, Å	49.6 – 1.70 (1.79 – 1.70)
<i>R</i> _{merge} , %	5.7 (45.7)
<i>I</i> /σ(<i>I</i>)	22.0 (5.2)
Completeness, %	97.9 (96.5)
Multiplicity	7.1 (7.2)
Number of reflections	
Overall	202987 (29635)
Unique	28713 (4094)
<i>B</i> factor ^a , Å ²	18.0
Anomalous Completeness	96.9 (95.6)
Anomalous Multiplicity	3.7 (3.7)
Phasing statistics	
No. of Fe sites	4
CC for solutions ^b	>20
Refinement statistics	
<i>R</i> _{crist} / <i>R</i> _{free} (%)	17.31/ 20.17
No. nonhydrogen-atoms	
Protein ^c	1688
Water	222
<i>B</i> factors, Å ²	
Protein ^c	14.8
Water	27.0
Rmsd bonds length (Å)	0.012
Rmsd bonds angle (°)	1.55
Ramachandran plot	
Allowed regions (%)	91.8
Additional allowed regions (%)	8.2
Generously allowed regions (%)	0
Disallowed regions (%)	0

Values in parentheses are for the highest resolution shell.

^a *B* factor as defined from Wilson plot.

^b correlation coefficient CC as defined by SHELXD

^c including two [2Fe-2S] clusters and two sulfate-ions

Supplementary Table 3: Overview of mutated regions in FDX1 and FDX2

Mutant name	Residues (FDX1)	Sequence (FDX1)	Residues (FDX2)	Sequence (FDX2)
Δ N13	--	--	56 – 68	AGEEDAGGPERPG
M1	77 – 82	ETLTTK	81 – 86	QRIPVS
M2	88 – 89	SL	92 – 93	NV
M3	91 – 97	DVVVEN	95 – 101	HLAQRH
M4	100 – 101	ID	104 – 105	LE
M5	103 – 104	FG	--	--
M6	108 – 109	GT	110 – 111	AS
M7	117 – 119	LIF	119 – 121	VYV
M8	128 – 131	DAIT	130 – 133	PPPE
M9	142 – 143	YG	144 – 145	PL
M9a	142	Y	144	P
M10	145 – 157	TDR	147 – 149	QEN
R135E	133	E	135	R
exC14	171 – 184	VADARQSIDVGKTS	161 – 174	TRNFYVDGHVPKPH
exC27	158 – 184	KSM DNMTVRVPET VADARQSIDVGKTS	148 – 174	PELEGAEFTLPKTR NFYVDGHVPKPH
Δ C12	173 – 184	DARQSIDVGKTS	163 – 174	NFYVDGHVPKPH

The table shows the name of the mutations and the respective native sequences of human FDX1 and FDX2 as depicted in Extended data Fig. 8. For mutants M1 – M10, the FDX1 sequence was replaced with the respective FDX2 residues and vice versa. For mutants exC14 and exC27, the 14/27 C-terminal residues of FDX1 were exchanged with the respective FDX2 sequence. These mutations were only done with FDX1. Δ N13 and Δ C12 refer to deletions of the N- and C-terminal residues. The Δ N13 and R135E mutations were only done with FDX2. Numbering according to sequence alignment in Extended data Fig. 8.

Supplementary Table 4: Yeast strains used in this study

Strain	Genotype	Method of Generation	Source/Reference
W303-1A	<i>MATa; ura3-1; ade2-1; trp1-1; his3-11,15; leu2-3,112</i>		42
Gal-YAH1	W303-1A, <i>pYAH1::LEU2-pGAL1-10</i>		43
BY4742	<i>MATa; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0</i>	obtained from Euroscarf	44
<i>lip5Δ</i>	BY4742, <i>lip5::KanMX4</i>	obtained from Euroscarf	
GalL-YAH1 <i>lip5Δ</i>	<i>lip5Δ; trp1::hph1NT1; pYAH1::GAL-L-natNT2</i>	PCR fragments (pFA6a-hph1NT1; pYM-N27) ⁴⁵	this work
GalL-YAH1 <i>cox15Δ</i>	W303-1A; <i>cox15::HIS3; pYAH1::GAL-L-natNT2</i>	PCR fragments (pFA6a-HIS3 (Euroscarf); pYM-N27 ⁴⁵)	this work

Table 5: Plasmid constructs used in this study

Plasmid	ORF	Backbone	Use	Source/Reference
pVA-I	none	pHindB	Human cell recovery after electroporation	23
PX459	none	pSpCas9(BB)-2A-Puro	gRNA expression	10.1038/nprot.2013.143
pEGFP-PEST	<i>EGFP-PEST</i>	pEGFP-N1	transfection control	this work
Su9-FDX1	<i>Su9-FDX1</i>	pDsRed2-C1	complementation	this work
p414-LIAS	<i>F1β-LIAS (1-35Δ)</i>	p414-TDH3	complementation	this work
p414-COX15	<i>F1β-COX15 (1-67Δ)</i>	p414-TDH3	complementation	this work
p416-FDX1	<i>F1β-FDX1 (1-60Δ)</i>	p416-TDH3	complementation	this work
p415-FDX2	<i>F1β-FDX2 (1-68Δ)</i>	p415-TDH3	complementation	this work
p426-TDH3	none	p426-TDH3	complementation	46
p426-FDX1	<i>F1β-FDX1 (1-61Δ)</i>	p426-TDH3	complementation	this work
p426-FDX2	<i>F1β-FDX2 (1-68Δ)</i>	p426-TDH3	complementation	this work
p416-Yah1	<i>S.c. Yah1</i>	p416-MET25	complementation	47
pETDuet1-FDX1	<i>His₆-FDX1 (1-60Δ)</i>	pETDuet1	protein purification	12
pETDuet1-FDX2	<i>His₆-FDX2 (1-68Δ)</i>	pETDuet1	protein purification	12
pASK-IBA43(+)-FDX1	<i>FDX1 (1-62Δ)</i>	pASK-IBA43(+)	protein purification	this work
pASK-IBA43(+)-FDX2	<i>FDX2 (1-68Δ)</i>	pASK-IBA43(+)	protein purification	this work
pASK-IBA43(+)-FDX2- Δ C12	<i>FDX2 (1-68Δ, 175-186Δ)</i>	pASK-IBA43(+)	protein purification, crystallization	this work
pET15b-Yah1	<i>S.c. Yah1 (1-57Δ)</i>	pET15b	protein purification	this work
pET28a(+)-LIAS	<i>His₆-Thr-LIAS (1-35Δ)</i>	pET28a(+)	protein purification	this work
pET24b(+)-ISCU2	<i>ISCU2-His₆ (1-34Δ)</i>	pET24b(+)	protein purification	48
pETDuet1-NFS1- <i>ISD11</i>	<i>NFS1 (1-55Δ), His₆-Tev-<i>ISD11</i></i>	pETDuet1	protein purification	48
pRSFDuet1-ACP	<i>ACP (1-68Δ)</i>	pRSFDuet1	protein purification	48
pMCSG7-FXN	<i>His₆-Tev-FXN (1-80Δ)</i>	pMCSG7	protein purification	48
pETDuet1-FDXR	<i>His₆-FDXR (1-32Δ)</i>	pETDuet1	protein purification	12

Mutations of the human FDXs as described in the paper were introduced by mutagenesis PCR using p426-*FDX1*, p426-*FDX2*, pETDuet1-*FDX1*, pASK-IBA43(+)-*FDX1* and pASK-IBA43(+)-*FDX2* as the initial templates. Combinations of different mutations were achieved by employment of newly generated mutated plasmids as templates for further rounds of mutagenesis.

Supplementary Table 6: Primary antibodies used in this study

Antibody	Source
rabbit anti- <i>FDX1</i>	12
rabbit anti- <i>FDX2</i>	12
rabbit anti-UQCRCFS1	H. Schagger (Frankfurt, Germany)
rabbit anti-MT-COX2	H. Schagger (Frankfurt, Germany)
rabbit anti-COX6A	H. Schagger (Frankfurt, Germany)
rabbit anti-ATP5F1A/B	H. Schagger (Frankfurt, Germany)
mouse anti-SDHB	Abcam (Cambridge, UK)
rabbit anti-lipoic acid	Calbiochem (Merck, Darmstadt, Germany)
rabbit anti-LIAS	Protein Tech Group (Chicago, USA)
rabbit anti VDAC1	Protein Tech Group (Chicago, USA)
mouse anti-DLAT	Santa Cruz biotechnology (Dallas, USA)
mouse anti-NDUFS1	Santa Cruz biotechnology (Dallas, USA)
mouse anti-NDUFS8	Santa Cruz biotechnology (Dallas, USA)
mouse anti-NDUFV2	Santa Cruz biotechnology (Dallas, USA)
mouse anti-GFP	Takara Bio (Göteborg, Sweden)

Supplementary References

1. Mellor, S.B., Vavitsas, K., Nielsen, A.Z. & Jensen, P.E. Photosynthetic fuel for heterologous enzymes: the role of electron carrier proteins. *Photosynth Res* **134**, 329-342 (2017).
2. Strushkevich, N. et al. Structural basis for pregnenolone biosynthesis by the mitochondrial monooxygenase system. *Proc Natl Acad Sci U S A* **108**, 10139-43 (2011).
3. Muller, J.J., Lapko, A., Bourenkov, G., Ruckpaul, K. & Heinemann, U. Adrenodoxin reductase - adrenodoxin complex structure suggests electron transfer path in steroid biosynthesis. *J. Biol. Chem.* **276**, 2786-2789. (2001).
4. Coghlan, V.M. & Vickery, L.E. Site-specific mutations in human ferredoxin that affect binding to ferredoxin reductase and cytochrome P450_{scc}. *J Biol Chem* **266**, 18606-12 (1991).
5. Cronan, J.E. Assembly of Lipoic Acid on Its Cognate Enzymes: an Extraordinary and Essential Biosynthetic Pathway. *Microbiol Mol Biol Rev* **80**, 429-50 (2016).
6. Jumper, J. et al. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-589 (2021).
7. Varadi, M. et al. AlphaFold Protein Structure Database: massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439-D444 (2022).
8. McLaughlin, M.I. et al. Crystallographic snapshots of sulfur insertion by lipoyl synthase. *Proc Natl Acad Sci U S A* **113**, 9446-50 (2016).
9. Harmer, J.E. et al. Structures of lipoyl synthase reveal a compact active site for controlling sequential sulfur insertion reactions. *Biochem J* **464**, 123-33 (2014).
10. Brixius-Anderko, S. & Scott, E.E. Structural and functional insights into aldosterone synthase interaction with its redox partner protein adrenodoxin. *J Biol Chem* **296**, 100794 (2021).
11. Chiliza, Z.E., Martinez-Oyanedel, J. & Syed, K. An overview of the factors playing a role in cytochrome P450 monooxygenase and ferredoxin interactions. *Biophys Rev* **12**, 1217-1222 (2020).
12. Sheftel, A.D. et al. Humans possess two mitochondrial ferredoxins, Fdx1 and Fdx2, with distinct roles in steroidogenesis, heme, and Fe/S cluster biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 11775-80 (2010).
13. Yadav, A.A., Patel, D., Wu, X. & Hasinoff, B.B. Molecular mechanisms of the biological activity of the anticancer drug elesclomol and its complexes with Cu(II), Ni(II) and Pt(II). *J Inorg Biochem* **126**, 1-6 (2013).
14. Nagai, M. et al. The oncology drug elesclomol selectively transports copper to the mitochondria to induce oxidative stress in cancer cells. *Free Radic Biol Med* **52**, 2142-50 (2012).
15. Tsvetkov, P. et al. Mitochondrial metabolism promotes adaptation to proteotoxic stress. *Nat Chem Biol* **15**, 681-689 (2019).
16. Tsvetkov, P. et al. Copper induces cell death by targeting lipoylated TCA cycle proteins. *Science* **375**, 1254-1261 (2022).
17. Vallieres, C., Holland, S.L. & Avery, S.V. Mitochondrial Ferredoxin Determines Vulnerability of Cells to Copper Excess. *Cell Chem Biol* **24**, 1228-1237 e3 (2017).
18. Macomber, L. & Imlay, J.A. The iron-sulfur clusters of dehydratases are primary intracellular targets of copper toxicity. *Proc Natl Acad Sci U S A* **106**, 8344-9 (2009).

19. Uhlen, M. et al. Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419 (2015).
20. Pettersen, E.F. et al. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605-12 (2004).
21. Ewen, K.M., Ringle, M. & Bernhardt, R. Adrenodoxin--a versatile ferredoxin. *IUBMB Life* **64**, 506-12 (2012).
22. Chu, G., Hayakawa, H. & Berg, P. Electroporation for the efficient transfection of mammalian cells with DNA. *Nucleic Acids Res* **15**, 1311-26 (1987).
23. Svensson, C. & Akusjarvi, G. Adenovirus VA RNAI: a positive regulator of mRNA translation. *Mol Cell Biol* **4**, 736-42 (1984).
24. Stehling, O. et al. Function and crystal structure of the dimeric P-loop ATPase CFD1 coordinating an exposed [4Fe-4S] cluster for transfer to apoproteins. *Proc Natl Acad Sci U S A* **115**, E9085-E9094 (2018).
25. Acosta Lopez, M.J. et al. Vanillic Acid Restores Coenzyme Q Biosynthesis and ATP Production in Human Cells Lacking COQ6. *Oxid Med Cell Longev* **2019**, 3904905 (2019).
26. Zollner, A. et al. Purification and functional characterization of human 11beta hydroxylase expressed in Escherichia coli. *FEBS J* **275**, 799-810 (2008).
27. Weiler, B.D. et al. Mitochondrial [4Fe-4S] protein assembly involves reductive [2Fe-2S] cluster fusion on ISCA1-ISCA2 by electron flow from ferredoxin FDX2. *Proc Natl Acad Sci U S A* **117**, 20555-20565 (2020).
28. McCarthy, E.L. & Booker, S.J. Biochemical Approaches for Understanding Iron-Sulfur Cluster Regeneration in Escherichia coli Lipoyl Synthase During Catalysis. *Methods Enzymol* **606**, 217-239 (2018).
29. Molik, S., Lill, R. & Mühlenhoff, U. Methods for studying iron metabolism in yeast mitochondria. *Methods Cell. Biol.* **80**, 261-280 (2007).
30. Rabinowitz, J.C. Analysis of acid-labile sulfide and sulfhydryl groups. *Methods Enzymol* **53**, 275-7 (1978).
31. Muller, A. et al. New aspects of electron transfer revealed by the crystal structure of a truncated bovine adrenodoxin, Adx(4-108). *Structure* **6**, 269-80 (1998).
32. Pikuleva, I.A., Tesh, K., Waterman, M.R. & Kim, Y. The tertiary structure of full-length bovine adrenodoxin suggests functional dimers. *Arch Biochem Biophys* **373**, 44-55 (2000).
33. Collaborative Computational Project, N. The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D Biol Crystallogr* **50**, 760-3 (1994).
34. Matthews, B.W. Solvent content of protein crystals. *J Mol Biol* **33**, 491-7 (1968).
35. Sheldrick, G.M. A short history of SHELX. *Acta Crystallogr A* **64**, 112-22 (2008).
36. Perrakis, A., Morris, R. & Lamzin, V.S. Automated protein model building combined with iterative structure refinement. *Nat Struct Biol* **6**, 458-63 (1999).
37. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* **60**, 2126-32 (2004).
38. Murshudov, G.N., Vagin, A.A. & Dodson, E.J. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr* **53**, 240-55 (1997).
39. Laskowski, R.A., MacArthur, M.W., Moss, D.S. & Thornton, J.M. PROCHECK: a program to check the stereochemical quality of protein structures. *Journal of Applied Crystallography* **26**, 283-291 (1993).
40. Hutchinson, E.G. & Thornton, J.M. PROMOTIF--a program to identify and analyze structural motifs in proteins. *Protein Sci* **5**, 212-20 (1996).

41. Arnold, K., Bordoli, L., Kopp, J. & Schwede, T. The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics* **22**, 195-201 (2006).
42. Mortimer, R.K. & Johnston, J.R. Genealogy of principal strains of the yeast genetic stock center. *Genetics* **113**, 35-43 (1986).
43. Lange, H., Kaut, A., Kispal, G. & Lill, R. A mitochondrial ferredoxin is essential for biogenesis of cellular iron-sulfur proteins. *Proc Natl Acad Sci U S A* **97**, 1050-5 (2000).
44. Brachmann, C.B. et al. Designer deletion strains derived from *Saccharomyces cerevisiae* S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast* **14**, 115-32 (1998).
45. Janke, C. et al. A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* **21**, 947-962 (2004).
46. Mumberg, D., Müller, R. & Funk, M. Yeast vectors for controlled expression of heterologous proteins in different genetic backgrounds. *Gene* **156**, 119-122. (1995).
47. Ozeir, M. et al. Coenzyme Q biosynthesis: Coq6 is required for the C5-hydroxylation reaction and substrate analogs rescue Coq6 deficiency. *Chemistry & biology* **18**, 1134-42 (2011).
48. Freibert, S.A. et al. N-terminal tyrosine of ISCU2 triggers [2Fe-2S] cluster synthesis by ISCU2 dimerization. *Nat Commun* **12**, 6902 (2021).