



HAL
open science

Extremal weights and a tameness criterion for mod p Galois representations

Daniel Le, Bao Viet Le Hung, Brandon Levin, Stefano Morra

► **To cite this version:**

Daniel Le, Bao Viet Le Hung, Brandon Levin, Stefano Morra. Extremal weights and a tameness criterion for mod p Galois representations. 2022. hal-03840799

HAL Id: hal-03840799

<https://hal.science/hal-03840799v1>

Preprint submitted on 6 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXTREMAL WEIGHTS AND A TAMENESS CRITERION FOR MOD p GALOIS REPRESENTATIONS

DANIEL LE, BAO V. LE HUNG, BRANDON LEVIN, AND STEFANO MORRA

ABSTRACT. We study the weight part of Serre’s conjecture for generic n -dimensional mod p Galois representations. We first generalize Herzig’s conjecture to the case where the field is ramified at p and prove the weight elimination direction of our conjecture. We then introduce a new class of weights associated to n -dimensional local mod p representations which we call *extremal weights*. Using a “Levi reduction” property of certain potentially crystalline Galois deformation spaces, we prove the modularity of these weights. As a consequence, we deduce the weight part of Serre’s conjecture for unit groups of some division algebras in generic situations.

CONTENTS

1. Introduction	3
1.1. Results	3
1.2. Geometric methods: Levi reduction for deformation rings	6
1.3. Overview	7
1.4. Acknowledgements	8
1.5. Notation	8
2. Preliminaries	11
2.1. Extended affine Weyl groups	11
2.2. Combinatorics of the extended affine Weyl group	12
2.3. The weight part of a Serre-type conjecture for tame representations	17
2.4. Combinatorics of Serre weights	19
3. Breuil–Kisin modules	24
3.1. Moduli of Breuil–Kisin modules and local models	24
3.2. Mod p monodromy	28
3.3. Semicontinuity I	29
3.4. Specializations	30
3.5. Semicontinuity II	31
3.6. Specialization pairs	34
3.7. Extremal weights	35
3.8. Maximally ordinary weights	36
3.9. Connections to Emerton–Gee stacks	38
3.10. The extremal locus	39
4. Some potentially crystalline deformation rings	41
4.1. The main result on Galois deformation rings	41
4.2. Gauge bases and parabolic structures	41
4.3. Interlude: GL_2 Pappas–Zhu models	45
4.4. Analysis of the monodromy condition	45
4.5. Proof of Theorem 4.1.1	50

5. The main results	52
5.1. Weight elimination	52
5.2. Patching functors	52
5.3. Cycles from patching functors	53
5.4. Weight elimination and modularity of extremal weights	54
5.5. Global results	57
References	65

1. INTRODUCTION

In [Ser87], Serre announced his celebrated modularity conjecture for two-dimensional mod p Galois representations (now a theorem of Khare–Wintenberger [KW09]). Much of that paper is dedicated to formulating a strong version which predicts the minimal level and weight at which one would find a suitable modular form in terms of the restriction of the mod p Galois representation to decomposition groups. There has been enormous progress on generalizing Serre’s strong version to other number fields and higher dimensions as well as proving that the strong version follows from the weak version. This includes rather complete results in dimension two [CV92, Edi92, Gro90, BDJ10, Sch08, Gee11, New14, GLS14, GLS15] which have led to a much better understanding of the hypothetical mod p Langlands correspondence [BP13, BHH⁺]. Still, there has been no complete generalization of the weight part of Serre’s conjecture in any dimension larger than two, reflecting difficulties in integral p -adic Hodge theory.

Already in Serre’s original paper, the most subtle case occurs when the restriction $\bar{\rho}$ to the decomposition group at p is *not* semisimple: the *très ramifiée* extensions have fewer modular Serre weights than the *peu ramifiée* ones. Still in dimension two but for unramified extensions of \mathbb{Q}_p , the set of Serre weights attached to a nonsplit extension by [BDJ10] depends in a subtle way on the extension class even in *generic* cases (see also [DDR16, CEGM17]). In higher dimensions, [Her09, GHS18] define a conjectural weight set for unramified extensions of \mathbb{Q}_p only when $\bar{\rho}$ is generic and *semisimple*. This conjecture was proven in [LLHLMa] (for many definite unitary groups) under a somewhat exotic genericity condition—again only for semisimple $\bar{\rho}$. Further, [LLHLMa, Conjecture 9.1.5] extended (still for unramified extensions of \mathbb{Q}_p) the weight part of Serre’s conjecture to generic non-semisimple Galois representations using stacks defined by Emerton–Gee [EGa] and their local models. It predicts that $\bar{\rho}$ is modular of a Serre weight σ if $\bar{\rho}$ lies on a certain cycle on the Emerton–Gee stack attached to σ using known cases of the Breuil–Mézard conjecture. It seems that this geometric perspective is essential because already in dimension three the set of modular weights cannot be described by linear conditions on extension classes [LLHLMb]. At present, this conjecture seems difficult to access for several reasons. First, proving the conjecture using the Breuil–Mézard perspective would require showing that various Taylor–Wiles patched modules have full support which is a notoriously difficult problem. (The relevant deformation rings are not always integral domains when $\bar{\rho}$ is not semisimple.) Second, the cycle attached to σ depends on cycle multiplicities appearing in local models that are not well understood at present. Finally, a classification of possible weight sets seems complicated because the formal definition of the Emerton–Gee stack makes it difficult to understand basic geometric questions such as the intersection patterns of its irreducible components.

The goal of this paper is to establish upper and lower bounds for the set of modular Serre weights which apply to generic non-semisimple Galois representations in arbitrary dimension over possibly ramified extensions of \mathbb{Q}_p . When applicable, our bounds greatly improve the existing ones in the literature (cf. Remark 1.1.7). They arise from our study of (potentially crystalline) Emerton–Gee stacks. The new symmetries and geometric structures we observe should play a role in organizing further investigations on Serre weight conjectures. Our results are new even in the semisimple case: we formulate a generalization of Herzig’s conjecture for ramified p -adic fields which we prove is always an upper bound. We give two concrete applications of our bounds. Under a mild and explicit genericity condition, we establish the weight part of Serre’s conjecture for unit groups of certain division algebras and generalize Gross’ automorphic tameness criterion.

1.1. Results. Let p be a prime and $n \geq 2$ be an integer. Let F/F^+ be a CM extension of a totally real field $F^+ \neq \mathbb{Q}$. Assume for the sake of exposition that there is a single place v of F^+ dividing

p which splits in F . (Our results apply whenever all the places of F^+ dividing p split in F .) Let G be a definite unitary group over F^+ split over F which is isomorphic to $U(n)$ at each infinite place and split at v . A (global) Serre weight is an irreducible smooth $\overline{\mathbb{F}}_p$ -representation V of $G(\mathcal{O}_{F^+,v})$, i.e. the inflation to $G(\mathcal{O}_{F^+,v})$ of an irreducible $\overline{\mathbb{F}}_p$ -representation of $G(k_v)$, where k_v is the residue field of F^+ at v . For a mod p Galois representation $\overline{r} : G_F \rightarrow \mathrm{GL}_n(\overline{\mathbb{F}}_p)$, let $W(\overline{r})$ denote the collection of modular Serre weights for \overline{r} . That is, $V \in W(\overline{r})$ if the Hecke eigensystem attached to \overline{r} appears in a space of mod p automorphic forms on G of weight V for some prime to v level.

Fix a place \tilde{v} of F dividing v which identifies $G(k_v)$ with $\mathrm{GL}_n(k_v)$. Define $\overline{r}_v := \overline{r}|_{\mathrm{Gal}(\overline{F}_{\tilde{v}}/F_{\tilde{v}})}$. The goal of the weight part of Serre’s conjecture is to predict $W(\overline{r})$ in terms of \overline{r}_v or more precisely, the restriction of \overline{r}_v to inertia.

Our global (and local) results include genericity conditions on \overline{r}_v which will be made precise in the body of the paper. We stress that our genericity conditions are completely explicit, unlike those of [LLHLMa]. We note however that for most results the genericity conditions require p to be at least $O(en^2)$ (where e is the absolute ramification index of $F_{\tilde{v}}$).

Let K/\mathbb{Q}_p be a finite extension with residue field k . For any tame n -dimensional $\overline{\mathbb{F}}_p$ -representation \overline{r} of $I_K \subset \mathrm{Gal}(\overline{K}/K)$ which extends to $\mathrm{Gal}(\overline{K}/K)$, one associates a Deligne–Lusztig representation $V(\overline{r})$ of $\mathrm{GL}_n(k)$ (generalizing [GHS18, Proposition 9.2.1]) which is defined over a finite extension E/\mathbb{Q}_p . It is also a representation of $\mathrm{GL}_n(\mathcal{O}_K)$ by inflation. Recall also the operator \mathcal{R} (see [GHS18, §9.2]) on the set of irreducible $\overline{\mathbb{F}}_p$ -representations of $\mathrm{GL}_n(k)$ (i.e. the set of Serre weights).

If K is unramified over \mathbb{Q}_p and \overline{r} is tame and generic, then Herzig defined the collection $W^?(\overline{r}) = \{\mathcal{R}(\sigma) \mid \sigma \in \mathrm{JH}(\overline{V}(\overline{r}|_{I_K}))\}$. In the ramified setting, we make the following generalization:

Definition 1.1.1. If \overline{r} is tame and generic, we define

$$W^?(\overline{r}) \stackrel{\mathrm{def}}{=} \left\{ \mathcal{R}(\sigma) \mid \sigma \in \left(\mathrm{JH} \left(\overline{V}(\overline{r}|_{I_K}) \otimes \overline{W}(0, 1-e, 2(1-e), \dots, (n-1)(1-e)) \right) \right) \right\}.$$

where $\overline{W}(0, 1-e, 2(1-e), \dots, (n-1)(1-e))$ is the irreducible algebraic representation of (parallel) highest weight $(0, 1-e, 2(1-e), \dots, (n-1)(1-e))$.

Remark 1.1.2. (1) In [LLHLMa] (see Theorem 4.7.6), in the unramified case, we give a geometric interpretation of Herzig’s $W^?(\overline{r})$ in terms of torus fixed points on certain subvarieties of the affine flag variety. Although we don’t directly use this description here because of a lack of local model theory in the ramified case, it motivated Definition 1.1.1.

(2) When $n = 2$, Schein gave in [Sch08] an explicit description of a weight set for tamely ramified \overline{r} . The two sets agree when \overline{r} is sufficiently generic, cf. §2.4.1.

We prove the weight elimination direction generalizing [LLHL19]:

Theorem 1.1.3 (“Weight elimination”, cf. Theorem 5.1.1). *Suppose that $\overline{r} : G_F \rightarrow \mathrm{GL}_n(\overline{\mathbb{F}}_p)$ satisfies standard Taylor–Wiles hypotheses and that \overline{r}_v is tame and sufficiently generic. Then*

$$\sigma_v \in W(\overline{r}) \implies \sigma_v \in W^?(\overline{r}_v).$$

When \overline{r} is not tame, unless $n \leq 3$, we don’t have an analogue of $W^?(\overline{r})$. Historically, certain classes of Serre weights have been identified which are expected to belong to $W^?(\overline{r})$. For example, Gee–Geraghty proved very generally the modularity of *ordinary* weights, i.e., those weights for which \overline{r}_v admits ordinary crystalline lifts. For tame \overline{r} and K unramified, [GHS18] introduce a notion of *obvious weight* which roughly speaking are characterized by the property that \overline{r} admits an “obvious” crystalline lift of specified Hodge–Tate weights, namely a sum of inductions of characters. Building on what we discovered when $n = 3$ in [LLHLMb], we introduce a notion of *extremal weights*

$W_{\text{extr}}(\bar{\rho})$ which encompasses (in generic cases) both these earlier notions and prove the following theorem:

Theorem 1.1.4 (“Modularity of extremal weights”, Theorem 5.5.5). *Let $\bar{r} : G_F \rightarrow \text{GL}_n(\mathbb{F})$ be an automorphic representation satisfying standard Taylor–Wiles conditions and such that \bar{r}_v is sufficiently generic. If either \bar{r} is potentially diagonalizably automorphic or $W_{\text{extr}}(\bar{r}_v) \cap W(\bar{r})$ is non-empty, then*

$$W_{\text{extr}}(\bar{r}_v) \subset W(\bar{r}).$$

When \bar{r} is tamely ramified at v and F^+ is unramified at p , Theorem 1.1.4 was proven in [LLHL19] and played a critical role in the theory of local models in [LLHLMa] and its applications to the weight part of Serre’s conjecture and the Breuil–Mézard conjecture. We expect Theorem 1.1.4 to have similar applications when F^+ is ramified at p and hope to return to this in future work.

There are two main ingredients in the proof of Theorem 1.1.4: a geometric one (which will be discussed in the next section) and a combinatorial one. The combinatorial ingredient is a hidden Weyl group symmetry. When $\bar{\rho}$ is tame and generic, then $W_{\text{extr}}(\bar{\rho})$ is naturally a torsor for a product of $[k : \mathbb{F}_p]$ -copies of the Weyl group S_n of GL_n , as explained in [GHS18]. Wildly ramified $\bar{\rho}$ have fewer weights in general and fewer extremal weights (see Proposition 3.7.3), but it turns out that the symmetry can be restored by enhancing an extremal weight with the data of a *specialization*.

A tame inertial $\overline{\mathbb{F}}_p$ -type is a continuous tame representation $I_K \rightarrow \text{GL}_n(\overline{\mathbb{F}}_p)$ which admits an extension to G_K . Tame inertial $\overline{\mathbb{F}}_p$ -types admit a combinatorial description in terms of fundamental characters of G_K (see §2.3.3). To a generic $\bar{\rho}$, we attach a collection of tame inertial $\overline{\mathbb{F}}_p$ -types which we call (extremal) *specializations* (Definition 3.4.1). This notion is somewhat elaborate, relying on the geometry of the Emerton–Gee stack (see e.g. §3.10). The semisimplification of $\bar{\rho}$ restricted to I_K is a prototypical example of a specialization but there are always others when $\bar{\rho}$ is not tame. It is generally expected that the predicted Serre weights of a wildly ramified $\bar{\rho}$ should be a subset of those of $\bar{\rho}^{\text{ss}}$. What we discover is the same is true for the other specializations of $\bar{\rho}$ as well.

Theorem 1.1.5 (cf. Theorem 5.1.1). *Suppose $\bar{r} : G_F \rightarrow \text{GL}_n(\overline{\mathbb{F}}_p)$ satisfies standard Taylor–Wiles hypotheses and that \bar{r}_v is generic. Let \bar{r}_v^{sp} be a specialization of \bar{r}_v . Then*

$$\sigma_v \in W(\bar{r}) \implies \sigma_v \in W^?(\bar{r}_v^{\text{sp}}).$$

The proof follows from a purely local result, showing that if \bar{r}_v admits a tamely potentially crystalline lift of type $(\tau, (n-1, n-2, \dots, 0))$ then so does any extension of the specialization of \bar{r}_v^{sp} to G_K , combined with the same weight elimination combinatorics used in the tame case.

Remark 1.1.6. Theorems 1.1.4 and 1.1.5 interact in an interesting way to constrain $W(\bar{r})$. Weight elimination can be used to produce further extremal Serre weights which in turn can be used to produce further specializations, and hence further weight elimination. Each iteration sharpens the bounds on $W(\bar{r})$.

Remark 1.1.7. Theorem 1.1.4 improves on previous lower bounds for the set of modular weights given by the sets of ordinary weights [GG12] and Fontaine–Laffaille weights [BGG18]. For example, if \bar{r}_v is completely reducible and generic, then \bar{r}_v has $n!$ ordinary weights, n^f Fontaine–Laffaille weights, and $n!^f$ extremal weights where f denotes $[k_v : \mathbb{F}_p]$.

The only previously known upper bound for $W(\bar{r})$ was $W^?(\bar{r}_v^{\text{ss}})$ (when F_v^+/\mathbb{Q}_p is unramified) [LLHL19]. When \bar{r}_v has $n!^f$ specializations (which is the generic behavior geometrically on the Emerton–Gee stack), Theorem 1.1.5 gives a far better upper bound. When $n = 3$ and \bar{r}_v has 6^f specializations, Theorem 1.1.5 gives an upper bound with a set of size at most 2^f rather than 9^f .

When $n = 2$ and F_v^+/\mathbb{Q}_p is unramified, the lower and upper bounds are equal and agree with the set defined in [BDJ10]. When $n = 3$ and F_v^+/\mathbb{Q}_p is unramified, the lower and upper bounds are close and play a major role in the resolution of the Serre weight conjecture in generic cases [LLHLMb].

A byproduct of our methods is an automorphic tameness criterion in the spirit of [Gro90]. When $n = 2$ and $F = \mathbb{Q}$, Gross’s tameness criterion says that for generic modular \bar{r} , tameness of \bar{r} at p is equivalent to $W(\bar{r})$ having two distinct Serre weights (as opposed to one). Here we show a similar criterion in terms of the modularity of two extremal weights.

Theorem 1.1.8 (“Automorphic tameness criterion”, Theorem 5.5.6). *Let $\sigma_v, \sigma'_v \in W_{\text{extr}}(\bar{r}_v^{\text{ss}})$ be extremal weights of \bar{r}_v^{ss} which differ by the longest element w_0 under the Weyl group symmetry. Suppose that $\sigma_v \in W(\bar{r})$ and that \bar{r}_v is sufficiently generic. Then the following are equivalent:*

- (1) $\sigma'_v \in W(\bar{r})$; and
- (2) \bar{r}_v is tame.

Remark 1.1.9. In the case where $W(\bar{r})$ contains a lowest alcove weight, our methods also give a refined version of the tameness criterion, showing that automorphic information even detects the stratum of \bar{r}_v in the moduli of Fontaine–Laffaille representations (with respect to a natural partition). This idea plays a crucial role in [LLHM⁺].

Finally, we discuss our results on the weight part of Serre’s conjecture for division algebras. When G is an anisotropic mod center inner form of GL_n locally at v , Serre weights lift to characteristic zero, and hence the modularity of a Serre weight can be rephrased in terms of the existence of automorphic lifts of specified types. By local-global compatibility, a necessary condition for the modularity of a generic Serre weight χ_v is the existence of a lift which is potentially crystalline at v of type $(\tau(\chi_v), (n-1, n-2, \dots, 0))$ for a certain tame cuspidal type $\tau(\chi_v)$. We prove the converse under some hypotheses.

Theorem 1.1.10 (Serre weights for division algebras, Theorem 5.5.13). *Suppose that v is unramified in F^+ , that G is an anisotropic mod center inner form of GL_n locally at v , that $\bar{r} : G_F \rightarrow \text{GL}_n(\overline{\mathbb{F}}_p)$ satisfies standard Taylor–Wiles hypotheses and is potentially diagonalizably automorphic, and that $\bar{\rho}_v$ is sufficiently generic. Then $\chi_v \in W(\bar{r})$ if and only if $\bar{\rho}_v$ admits a potentially crystalline lift of type $(\tau(\chi_v), (n-1, n-2, \dots, 0))$.*

This generalizes results in [GS11] when $n = 2$ and [Dotb] (announced in 2019) when \bar{r} is tamely ramified at v . The main difficulty lies in the construction of automorphic lifts. One has access to powerful potentially Barsotti–Tate modularity lifting results when $n = 2$ [GK14] that are not available in general. Via a transfer à la Jacquet–Langlands, it suffices to construct automorphic lifts for a group which is quasisplit at places dividing p . Generalizing [LLHL19] in the tame case, we construct the desired automorphic lifts using the modularity of extremal weights. In fact, we prove a more general criterion for the existence of automorphic lifts where $\tau(\chi_v)$ is replaced by any sufficiently generic tame inertial type (Corollary 5.5.11). That the modularity of extremal weights suffices to produce the desired automorphic lifts reduces to combinatorics of the extended affine Weyl group when \bar{r} is tamely ramified at v . In general, it requires an analysis of the geometry of local models (§3.10).

1.2. Geometric methods: Levi reduction for deformation rings. We now explain how we go about proving the modularity of the extremal weights (Theorem 1.1.4). As mentioned earlier, this notion expands on the notions of ordinary and obvious weights. Gee–Geraghty [GG12] proved modularity of ordinary weights in considerable generality. Three of the authors proved modularity

of obvious weights in the unramified and generic semisimple case ([LLHL19]). Both of these results rely on producing potentially diagonalizable lifts of some prescribed type. Using these methods, we can access some but not all of the extremal weights. Instead, we adopt the strategy of [LLHLM18] and exploit the symmetry of our situation.

As described above, extremal weights when enhanced with the data of a specialization admit a Weyl group symmetry. The main point is to show that if two extremal weights σ, σ' are related by a simple reflection then the modularity of one implies the modularity of the other. To do this, we show that we can find a sequence of well-chosen tame types τ_0, \dots, τ_{2e} connecting σ to σ' where we can establish good combinatorial behavior of Serre weights and show that the Galois deformation rings are integral domains. The following is the main result on deformation rings that we use.

Theorem 1.2.1 (particular case of Theorem 4.1.1). *Let τ_i be one of the well-chosen tame inertial types described above (which will be sufficiently generic in our setup).*

Then $R_{\bar{\rho}}^{(n-1, n-2, \dots, 0), \tau}$ is either zero or is a normal domain. Furthermore, if it is nonzero, then either it is formally smooth over \mathcal{O} or the special fiber is reduced with exactly two irreducible components.

We actually prove a more general result for a larger class of deformation rings (Theorem 4.1.1). We approach the deformation spaces using the methods for studying Breuil–Kisin modules developed in [LLHLM18, LLHL19, LLHLMa]. This is the first time these methods have been adapted to the ramified setting. The key ingredient in our proof of Theorem 4.1.1 is the fact that the local models (in the sense of [LLHLMa] adapted to the ramified setting) of these Galois deformation spaces have a Levi reduction property: namely, they are formally smooth over similar local models attached to suitable Levi subgroups of GL_n . This turns out to be a general phenomenon whenever the shape of $\bar{\rho}$ relative to the type τ is suitably “decomposable”, which may be of independent interest. In the specific case of Theorem 1.2.1, the Levi subgroup we reduce to is $\mathrm{GL}_2 \times \mathrm{GL}_1^{n-2}$. Thus, we are able to show essentially that $R_{\bar{\rho}}^{(n-1, n-2, \dots, 0), \tau}$ is smooth over the completed local ring of a ramified local model of Pappas–Rapoport from which we deduce the normality and the description of the special fiber. We prove a similar Levi reduction property for the Pappas–Zhu local models, which is a key geometric input (Lemma 2.2.7) into the analysis of Serre weight combinatorics for these tame types.

Remark 1.2.2. When K/\mathbb{Q}_p is unramified, the relevant local model is a product of the Iwahori local models for GL_2 . Concretely, Theorem 1.2.1 says that $R_{\bar{\rho}}^{(n-1, n-2, \dots, 0), \tau}$ will either be power series ring over \mathcal{O} or will be formally smooth over $\mathcal{O}[[x, y]]/(xy - p)$. This observation in the case of GL_3 in [LLHLM18] was the starting point for this work.

1.3. Overview. In §2, after preliminaries on the affine Weyl group and admissible sets (§2.2), and recollections on Serre weights (§2.3), we formulate a Serre type conjecture on the weights of tame Galois representations over a possibly ramified field (cf. Definition 2.3.1) and obtain our main results on the combinatorics of Serre weights and tame inertial types for the *shapes* we will be interested in (cf. Propositions 2.4.8, 2.4.9).

§3 introduces the notion of *extremal weights* for Galois representations (§3.7). This requires preliminaries on the semicontinuity of shapes for Kisin modules (§3.3, 3.5, in different degrees of generality), the notion of *specializations* for Galois representations (§3.4) and the closely related notion of *specialization pairs* (§3.6). The non-emptiness of the set of the extremal weights is proved in §3.8 and §3.10 with different methods. In particular the geometric interpretation of this set in terms of the Emerton–Gee stack in the unramified case is in §3.9, 3.10.

§4 calculates the tamely potentially crystalline deformation rings which appear when studying extremal weights of Galois representations. We first establish structural results on Breuil–Kisin modules of certain *parabolic shapes* (§4.2) and then analyze the monodromy condition on them (§4.4, Lemma 4.4.3).

In §5, after a number of preliminaries on patching functors and cycles on potentially crystalline deformation rings (§5.2, 5.3), we prove in §5.4 the modularity of extremal weights in an axiomatic setup (Theorem 5.4.3). §5.5 contains our global applications to automorphic forms on definite unitary groups.

1.4. Acknowledgements. The origin of this work dates back to a stay at the Mathematisches Forschungsinstitut Oberwolfach in winter 2019, which provided excellent working condition, and was further carried out during several visits at the University of Arizona and Northwestern University. We would like to heartily thank these institutions for the outstanding research conditions they provided, and for their support.

We thank Andrea Dotto for explaining the application of the work [LLHL19] of the first three authors to the weight part of Serre’s conjecture for division algebras and for pointing out Remark 5.5.14.

D.L. was supported by the National Science Foundation under agreements Nos. DMS-1128155 and DMS-1703182, an AMS-Simons travel grant, and a start-up grant from Purdue University. B.L.H. acknowledges support from the National Science Foundation under grant Nos. DMS-1128155, DMS-1802037 and the Alfred P. Sloan Foundation. B.L. was supported by National Science Foundation grant DMS-1952556 and the Alfred P. Sloan Foundation. S.M. was supported by the Institut Universitaire de France and the ANR-18-CE40-0026 (CLap CLap).

1.5. Notation. For a field K , we denote by \overline{K} a fixed separable closure of K and let $G_K \stackrel{\text{def}}{=} \text{Gal}(\overline{K}/K)$. If K is defined as a subfield of an algebraically closed field, then we set \overline{K} to be this field.

If K is a nonarchimedean local field, we let $I_K \subset G_K$ denote the inertial subgroup and $W_K \subset G_K$ denote the Weil group. We fix a prime $p \in \mathbb{Z}_{>0}$. Let $E \subset \overline{\mathbb{Q}}_p$ be a subfield which is finite-dimensional over \mathbb{Q}_p . We write \mathcal{O} to denote its ring of integers, fix a uniformizer $\varpi \in \mathcal{O}$ and let \mathbb{F} denote the residue field of E . We will assume throughout that E is sufficiently large.

1.5.1. Reductive groups. Let G denote a split connected reductive group (over some ring) together with a Borel B , a maximal split torus $T \subset B$, and $Z \subset T$ the center of G . Let $d = \dim G - \dim B$. When G is a product of copies of GL_n , we will take B to be upper triangular Borel and T the diagonal torus. Let $\Phi^+ \subset \Phi$ (resp. $\Phi^{\vee,+} \subset \Phi^\vee$) denote the subset of positive roots (resp. positive coroots) in the set of roots (resp. coroots) for (G, B, T) . We use the notation $\alpha > 0$ (resp. $\alpha < 0$) for a positive (resp. negative) root $\alpha \in \Phi$. Let Δ (resp. Δ^\vee) be the set of simple roots (resp. coroots). Let $X^*(T)$ be the group of characters of T , and set $X^0(T)$ to be the subgroup consisting of characters $\lambda \in X^*(T)$ such that $\langle \lambda, \alpha^\vee \rangle = 0$ for all $\alpha^\vee \in \Delta^\vee$. Let $\Lambda_R \subset X^*(T)$ denote the root lattice for G . Let $W(G)$ denote the Weyl group of (G, T) . Let w_0 denote the longest element of $W(G)$. We sometimes write W for $W(G)$ when there is no chance for confusion. Let W_a (resp. \widetilde{W}) denote the affine Weyl group and extended affine Weyl group

$$W_a = \Lambda_R \rtimes W(G), \quad \widetilde{W} = X^*(T) \rtimes W(G)$$

for G . We use $t_\nu \in \widetilde{W}$ to denote the image of $\nu \in X^*(T)$.

The Weyl groups $W(G)$, \widetilde{W} , and W_a act naturally on $X^*(T)$. If A is any ring, then the above Weyl groups act naturally on $X^*(T) \otimes_{\mathbb{Z}} A$ by extension of scalars.

Let M be a free \mathbb{Z} -module of finite rank (e.g. $M = X^*(T)$). The duality pairing between M and its \mathbb{Z} -linear dual M^* will be denoted by $\langle \cdot, \cdot \rangle$. If A is any ring, the pairing $\langle \cdot, \cdot \rangle$ extends by A -linearity to a pairing between $M \otimes_{\mathbb{Z}} A$ and $M^* \otimes_{\mathbb{Z}} A$, and by an abuse of notation it will be denoted with the same symbol $\langle \cdot, \cdot \rangle$.

We write $G^{\vee} = G_{/\mathbb{Z}}^{\vee}$ for the split connected reductive group over \mathbb{Z} defined by the root datum $(X_*(T), X^*(T), \Phi^{\vee}, \Phi)$. This defines a maximal split torus $T^{\vee} \subseteq G^{\vee}$ such that we have canonical identifications $X^*(T^{\vee}) \cong X_*(T)$ and $X_*(T^{\vee}) \cong X^*(T)$.

Let $V \stackrel{\text{def}}{=} X^*(T) \otimes_{\mathbb{Z}} \mathbb{R} \setminus (\bigcup_{(\alpha, n)} H_{\alpha, n})$. For $(\alpha, k) \in \Phi \times \mathbb{Z}$, we have the root hyperplane $H_{\alpha, k} \stackrel{\text{def}}{=} \{x \in V \mid \langle x, \alpha^{\vee} \rangle = k\}$ and the half-hyperplanes $H_{\alpha, k}^+ = \{x \in V \mid \langle x, \alpha^{\vee} \rangle > k\}$ and $H_{\alpha, k}^- = \{x \in V \mid \langle x, \alpha^{\vee} \rangle < k\}$. An alcove is a connected component of $V \setminus (\bigcup_{(\alpha, n)} H_{\alpha, n})$.

We say that an alcove A is *restricted* if $0 < \langle \lambda, \alpha^{\vee} \rangle < 1$ for all $\alpha \in \Delta$ and $\lambda \in A$. We let A_0 denote the (dominant) base alcove, i.e. the set of $\lambda \in X^*(T) \otimes_{\mathbb{Z}} \mathbb{R}$ such that $0 < \langle \lambda, \alpha^{\vee} \rangle < 1$ for all $\alpha \in \Phi^+$. Let \mathcal{A} denote the set of alcoves. Recall that \widetilde{W} acts transitively on the set of alcoves, and $\widetilde{W} \cong \widetilde{W}_a \rtimes \Omega$ where Ω is the stabilizer of A_0 . We define

$$\widetilde{W}^+ \stackrel{\text{def}}{=} \{\tilde{w} \in \widetilde{W} : \tilde{w}(A_0) \text{ is dominant}\}.$$

and

$$\widetilde{W}_1^+ \stackrel{\text{def}}{=} \{\tilde{w} \in \widetilde{W}^+ : \tilde{w}(A_0) \text{ is restricted}\}.$$

We fix an element $\eta_0 \in X^*(T)$ such that $\langle \eta_0, \alpha^{\vee} \rangle = 1$ for all positive simple roots α and let \tilde{w}_h be $w_0 t_{-\eta_0} \in \widetilde{W}_1^+$.

When $G = \text{GL}_n$, we fix an isomorphism $X^*(T) \cong \mathbb{Z}^n$ in the standard way, where the standard i -th basis element $(0, \dots, 1, \dots, 0)$ (with the 1 in the i -th position) of the right-hand side corresponds to extracting the i -th diagonal entry of a diagonal matrix. When G is a product of copies of GL_n indexed over a set \mathcal{J} we take $\eta_0 \in X^*(T)$ to correspond to the element $(n-1, n-2, \dots, 0)_{j \in \mathcal{J}} \in (\mathbb{Z}^n)^{\mathcal{J}}$ in the identification above. In this case, given $j \in \mathcal{J}$ we write $\eta_{0, j} \in$ to denote the element which corresponds to the tuple $(n-1, \dots, 1, 0)$ at j and to the zero tuple elsewhere.

Let F_p^+ be a finite étale \mathbb{Q}_p -algebra. Then F_p^+ is isomorphic to a product $\prod_{S_p} F_v^+$ for some finite set S_p where for each $v \in S_p$, F_v^+ is finite extension of \mathbb{Q}_p . For each $v \in S_p$, let $\mathcal{O}_{F_v^+} \subset F_v^+$ be the ring of integers, k_v the residue field, $F_{v,0}^+ \subset F_v^+$ the maximal unramified subextension, f_v the unramified degree $[F_{v,0}^+ : \mathbb{Q}_p]$, and e_v the ramification degree $[F_v^+ : F_{v,0}^+]$. Let \mathcal{O}_p be the product $\prod_{v \in S_p} \mathcal{O}_{F_v^+}$ and k_p the product $\prod_{v \in S_p} k_v$.

In global applications, S_p will be a finite set of places dividing p of a number field F^+ . When working locally, S_p will have cardinality one, in which case we drop the subscripts from f_v , e_v , and k_v and denote the single extension F_v^+ of \mathbb{Q}_p by K .

If G is a split connected reductive group over \mathbb{F}_p , with Borel B , maximal split torus T , and center Z , we let $G_0 \stackrel{\text{def}}{=} \text{Res}_{k_p/\mathbb{F}_p} G/k_p$ with Borel subgroup $B_0 \stackrel{\text{def}}{=} \text{Res}_{k_p/\mathbb{F}_p} B/k_p$, maximal torus $T_0 \stackrel{\text{def}}{=} \text{Res}_{k_p/\mathbb{F}_p} T/k_p$, and $Z_0 = \text{Res}_{k_p/\mathbb{F}_p} Z/k_p$. Assume that \mathbb{F} contains the image of any ring homomorphism $k_p \rightarrow \overline{\mathbb{F}}_p$ and let \mathcal{J} be the set of ring homomorphisms $k_p \rightarrow \mathbb{F}$. Then $\underline{G} \stackrel{\text{def}}{=} (G_0)_{/\mathbb{F}}$ is naturally identified with the split reductive group $G_{/\mathbb{F}}^{\mathcal{J}}$. We similarly define $\underline{B}, \underline{T}$, and \underline{Z} . Corresponding to $(\underline{G}, \underline{B}, \underline{T})$, we have the set of positive roots $\underline{\Phi}^+ \subset \underline{\Phi}$ and the set of positive coroots $\underline{\Phi}^{\vee, +} \subset \underline{\Phi}^{\vee}$. The notations $\underline{\Lambda}_R, \underline{W}, \underline{W}_a, \underline{\widetilde{W}}, \underline{\widetilde{W}}^+, \underline{\widetilde{W}}_1^+, \underline{\Omega}$ should be clear as should the natural isomorphisms $X^*(\underline{T}) = X^*(T)^{\mathcal{J}}$ and the like. The absolute Frobenius automorphism φ on k_p induces an automorphism π of the identified groups $X^*(\underline{T})$ and $X_*(\underline{T}^{\vee})$ by the formula $\pi(\lambda)_{\sigma} = \lambda_{\sigma \circ \varphi^{-1}}$ for all $\lambda \in X^*(\underline{T})$ and

$\sigma : k_p \rightarrow \mathbb{F}$. We assume that, in this case, the element $\eta_0 \in X^*(\underline{T})$ we fixed is π -invariant. We similarly define an automorphism π of \underline{W} and \widetilde{W} .

1.5.2. *Galois Theory.* We now assume that S_p has cardinality one. We write $K \stackrel{\text{def}}{=} F_v^+$ and drop the subscripts from f_v , e_v , and k_v . Let $W(k)$ be ring of Witt vectors which is also ring of integers \mathcal{O}_{K_0} of K_0 . We denote the arithmetic Frobenius automorphism on $W(k)$ by φ , which acts as raising to p -th power on the residue field. We fix an embedding σ_0 of K_0 into E (equivalently an embedding k into \mathbb{F}) and define $\sigma_j = \sigma_0 \circ \varphi^{-j}$, which gives an identification between $\mathcal{J} = \text{Hom}(k, \mathbb{F})$ and $\mathbb{Z}/f\mathbb{Z}$.

We normalize Artin's reciprocity map $\text{Art}_K : K^\times \rightarrow W_K^{\text{ab}}$ in such a way that uniformizers are sent to geometric Frobenius elements.

Given an uniformizer $\pi_K \in \mathcal{O}_K$ and a sequence $\underline{\pi}_K \stackrel{\text{def}}{=} (\pi_{K,m})_{m \in \mathbb{N}} \in \overline{K}^{\mathbb{N}}$ satisfying $\pi_{K,m+1}^p = \pi_{K,m}$, $\pi_{K,0} \stackrel{\text{def}}{=} \pi_K$ we let K_∞ be $\bigcup_{m \in \mathbb{N}} K(\pi_{K,m})$.

Given an element $\pi_1 \stackrel{\text{def}}{=} (-\pi_K)^{\frac{1}{p^f-1}} \in \overline{K}$ we have a corresponding character $\omega_K : I_K \rightarrow W(k)^\times$ which, using our choice of embedding σ_0 gives a fundamental character of niveau f

$$\omega_f := \sigma_0 \circ \omega_{\pi_1} : I_K \rightarrow \mathcal{O}^\times.$$

Let $\rho : G_K \rightarrow \text{GL}_n(E)$ be a p -adic, de Rham Galois representation. For $\sigma : K \hookrightarrow E$, we define $\text{HT}_\sigma(\rho)$ to be the multiset of σ -labeled Hodge-Tate weights of ρ , i.e. the set of integers i such that $\dim_E(\rho \otimes_{\sigma, K} \mathbb{C}_p(-i))^{G_K} \neq 0$ (with the usual notation for Tate twists). In particular, the cyclotomic character ε has Hodge-Tate weights 1 for all embedding $\sigma : K \hookrightarrow E$. For $\mu = (\mu_j)_j \in X^*(\underline{T})$ we say that ρ has Hodge-Tate weighs μ if

$$\text{HT}_{\sigma_j}(\rho) = \{\mu_{1,j}, \mu_{2,j}, \dots, \mu_{n,j}\}.$$

The *inertial type* of ρ is the isomorphism class of $\text{WD}(\rho)|_{I_K}$, where $\text{WD}(\rho)$ is the Weil-Deligne representation attached to ρ as in [CDT99], Appendix B.1 (in particular, $\rho \mapsto \text{WD}(\rho)$ is *covariant*). An inertial type is a morphism $\tau : I_K \rightarrow \text{GL}_n(E)$ with open kernel and which extends to the Weil group W_K of G_K . We say that ρ has type (μ, τ) if ρ has Hodge-Tate weights μ and inertial type given by (the isomorphism class of) τ .

1.5.3. *Miscellaneous.* For any ring S , we define $\text{Mat}_n(S)$ to be the set of $n \times n$ matrix with entries in S . If $M \in \text{Mat}_n(S)$ and $A \in \text{GL}_n(S)$ we write

$$(1.1) \quad \text{Ad}(A)(M) \stackrel{\text{def}}{=} A M A^{-1}.$$

If X is an ind-scheme defined over \mathcal{O} , we write $X_E \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}} \text{Spec } E$ and $X_{\mathbb{F}} \stackrel{\text{def}}{=} X \times_{\text{Spec } \mathcal{O}} \text{Spec } \mathbb{F}$ to denote its generic and special fiber, respectively.

2. PRELIMINARIES

2.1. Extended affine Weyl groups. In this section, we collect some background material on Weyl groups which will be needed throughout the paper.

Recall from §1.5.1 that G is a split reductive group with split maximal torus T and Borel B . Let $W \stackrel{\text{def}}{=} W(G, T)$ be the Weyl group and $V \stackrel{\text{def}}{=} X^*(T) \otimes \mathbb{R} \cong X_*(T^\vee) \otimes \mathbb{R}$ denote the apartment of (G, T) on which $\widetilde{W} \stackrel{\text{def}}{=} X^*(T) \rtimes W$ acts. Let \mathcal{C}_0 denote the dominant Weyl chamber in V . For any $w \in W(G)$, let $\mathcal{C}_w = w(\mathcal{C}_0)$. In particular, denoting the longest element of W by w_0 , \mathcal{C}_{w_0} is the anti-dominant Weyl chamber.

Recall from §1.5.1 that \mathcal{A} denotes the set of alcoves of $X^*(T) \otimes \mathbb{R}$ and that $A_0 \in \mathcal{A}$ denotes the dominant base alcove. We let \uparrow denote the upper arrow ordering on alcoves as defined in [Jan03, §II.6.5]. Since W_a acts simply transitively on the set of alcoves, $\tilde{w} \mapsto \tilde{w}(A_0)$ induces a bijection $W_a \xrightarrow{\sim} \mathcal{A}$ and thus an upper arrow ordering \uparrow on W_a . The dominant base alcove A_0 also defines a set of simple reflections in W_a and thus a Coxeter length function on W_a denoted $\ell(-)$ and a Bruhat order on W_a denoted by \leq .

If $\Omega \subset \widetilde{W}$ is the stabilizer of the base alcove, then $\widetilde{W} = W_a \rtimes \Omega$ and so \widetilde{W} inherits a Bruhat and upper arrow order in the standard way: For $\tilde{w}_1, \tilde{w}_2 \in W_a$ and $\delta \in \Omega$, $\tilde{w}_1\delta \leq \tilde{w}_2\delta$ (resp. $\tilde{w}_1\delta \uparrow \tilde{w}_2\delta$) if and only if $\tilde{w}_1 \leq \tilde{w}_2$ (resp. $\tilde{w}_1 \uparrow \tilde{w}_2$), and elements in different right W_a -cosets are incomparable. We extend $\ell(-)$ to \widetilde{W} by letting $\ell(\tilde{w}\delta) \stackrel{\text{def}}{=} \ell(\tilde{w})$ for any $\tilde{w} \in W_a$, $\delta \in \Omega$.

Definition 2.1.1. If $\tilde{w}_1, \dots, \tilde{w}_m \in \widetilde{W}$, we say that $\tilde{w}_1\tilde{w}_2 \cdots \tilde{w}_m$ is a *reduced expression* if the inequality $\ell(\tilde{w}_1\tilde{w}_2 \cdots \tilde{w}_m) \leq \sum_{i=1}^m \ell(\tilde{w}_i)$ is an equality.

Let $(\widetilde{W}^\vee, \leq)$ be the following partially ordered group: \widetilde{W}^\vee is identified with \widetilde{W} as a group, and $\ell(-)$ and \leq are defined with respect to the *antidominant* base alcove.

Definition 2.1.2. We define a bijection $\tilde{w} \mapsto \tilde{w}^*$ between \widetilde{W} and \widetilde{W}^\vee as follows: for $\tilde{w} = t_\nu w \in \widetilde{W}$, with $w \in W$ and $\nu \in X^*(T) = X_*(T^\vee)$, then $\tilde{w}^* \stackrel{\text{def}}{=} w^{-1}t_\nu \in \widetilde{W}^\vee$.

This bijection respects notions of length and Bruhat order (see [LLHL19, Lemma 2.1.3]).

We recall some fundamental notions associated to the geometry of $X^*(T)$ and \widetilde{W} .

Definition 2.1.3. Let $\lambda \in X^*(T)$. The convex hull of the set $\{w(\lambda) \mid w \in W\}$ is defined to be

$$\text{Conv}(\lambda) \stackrel{\text{def}}{=} \bigcap_{w \in W} w(\lambda) + \overline{\mathcal{C}_{ww_0}}$$

where $\overline{\mathcal{C}_{ww_0}}$ denotes the closure of the Weyl chamber \mathcal{C}_{ww_0} .

We recall the definition of the admissible set from [KR00]:

Definition 2.1.4. For $\lambda \in X^*(T)$, define

$$\text{Adm}(\lambda) \stackrel{\text{def}}{=} \left\{ \tilde{w} \in \widetilde{W} \mid \tilde{w} \leq t_{w(\lambda)} \text{ for some } w \in W \right\}.$$

For a positive integer e , define the e -critical strips to be strips $H_\alpha^{(1-e, e)} = \{x \in V \mid 1 - e < \langle x, \alpha^\vee \rangle < e\}$ where $\alpha \in \Phi^+$.

Definition 2.1.5. An alcove $A \in \mathcal{A}$ is e -regular if A does not lie in any e -critical strip. For any $\tilde{w} \in \widetilde{W}$, we say \tilde{w} is e -regular if $\tilde{w}(A_0)$ is e -regular. Define

$$\text{Adm}^{e\text{-reg}}(\lambda) = \{\tilde{w} \in \text{Adm}(\lambda) \mid \tilde{w} \text{ is } e\text{-regular}\}.$$

Proposition 2.1.6. *If $\tilde{w} \in \widetilde{W}$ is e -regular, then there exist \tilde{w}_1 and $\tilde{w}_2 \in \widetilde{W}_1^+$ and a dominant weight $\nu \in X^*(T)$ such that $\tilde{w} = \tilde{w}_2^{-1} w_0 t_{\nu+(e-1)\eta_0} \tilde{w}_1$. Moreover, \tilde{w}_1 , \tilde{w}_2 , and ν as above are unique up to $X^0(T)$. Conversely, if \tilde{w}_1 and \tilde{w}_2 are elements of \widetilde{W}^+ , then $\tilde{w}_2^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_1$ is e -regular.*

We conclude this section by recalling from [LLHLMa, Definition 2.1.10] the various notions of genericity for elements of $X^*(T)$.

Definition 2.1.7. Let $\lambda \in X^*(T)$ be a weight and let $m \geq 0$ be an integer.

- (1) We say that λ lies m -deep in its p -alcove if for all $\alpha \in \Phi^+$, there exist integers $m_\alpha \in \mathbb{Z}$ such that $pm_\alpha + m < \langle \lambda + \eta_0, \alpha^\vee \rangle < p(m_\alpha + 1) - m$.
- (2) We say that $\lambda \in X^*(T)$ is m -generic if $m < |\langle \lambda, \alpha^\vee \rangle + pk|$ for all $\alpha \in \Phi$ and $k \in \mathbb{Z}$ (or equivalently, $\lambda - \eta_0$ is m -deep in its p -alcove).
- (3) We say that an element $\tilde{w} = wt_\nu$ (in either \widetilde{W} or \widetilde{W}^\vee) is m -small if $\langle \nu, \alpha^\vee \rangle \leq m$ for all $\alpha \in \Phi$.

2.2. Combinatorics of the extended affine Weyl group. In this section, we collect a variety of results on the combinatorics of the extended affine Weyl group. These will be applied to the analysis of the combinatorics of Serre weights in §2.4. The methods are elementary with the exception of a geometric input from Pappas–Zhu local models in the proof of Lemma 2.2.7. We begin with results concerning the partial orderings \leq and \uparrow .

Lemma 2.2.1. *Suppose that $\tilde{x}^+ \in \widetilde{W}^+$ and $w \in W$. Then $w\tilde{x}^+$ is a reduced expression.*

Proof. There are galleries in the 1-direction from $w^{-1}(A_0)$ to A_0 and from A_0 to $\tilde{x}^+(A_0)$. We conclude that $\ell(w\tilde{x}^+) = \ell(w) + \ell(\tilde{x}^+)$. \square

Lemma 2.2.2. *Suppose that $\tilde{x} \in \widetilde{W}$ and $\tilde{w}^+ \in \widetilde{W}^+$ and $\tilde{x} \leq w_0\tilde{w}^+$. Then $w_0\tilde{w}^+ \uparrow w\tilde{x}$ for any $w \in W$.*

Proof. Since $w_0\tilde{w}^+$ is a reduced factorization by Lemma 2.2.1, $\tilde{x} \leq w_0\tilde{w}^+$ implies that $\tilde{x} = s\tilde{x}'$ for $s \in W$ and $\tilde{x}' \in \widetilde{W}$ with $\tilde{x}' \leq \tilde{w}^+$. Factoring \tilde{x}' as the reduced expression $s'\tilde{x}^+$ where $s' \in W$ and $\tilde{x}^+ \in \widetilde{W}^+$, we have that $\tilde{x}^+ \leq \tilde{x}'$. Replacing s by ss' and \tilde{x}' by \tilde{x}^+ , we can thus assume without loss of generality that $\tilde{x}' = \tilde{x}^+$ is in \widetilde{W}^+ . Wang’s theorem ([Wan87, Theorem 4.3] or [LLHL19, Theorem 4.1.1]) implies that $\tilde{x}^+ \uparrow \tilde{w}^+$. Then we have that $w_0\tilde{w}^+ \uparrow w_0\tilde{x}^+ \uparrow ws\tilde{x}^+ = w\tilde{x}$ for any $w \in W$ by [Jan03, II 6.5(5)]. \square

Lemma 2.2.3. *If \tilde{x} and $\tilde{y} \in \widetilde{W}$ and $\tilde{x} \leq \tilde{y}$, then $\tilde{x}^+ \uparrow \tilde{y}^+$ where \tilde{x}^+ and \tilde{y}^+ are the unique elements in $W\tilde{x} \cap \widetilde{W}^+$ and $W\tilde{y} \cap \widetilde{W}^+$, respectively. In particular, we have $\tilde{x} \uparrow \tilde{y}^+$.*

Proof. Let \tilde{y}^+ be $w\tilde{y}$ with $w \in W$. Since $w_0 = (w_0w)w^{-1}$ and $w_0(w\tilde{y})$ are reduced expressions (the latter by Lemma 2.2.1, the former by e.g. [?, §1.8]), so is $(w_0w)w^{-1}(w\tilde{y})$ and therefore so is $(w_0w)\tilde{y}$. Since $\tilde{x}^+ \leq \tilde{x}$ (by Lemma 2.2.1) and $\tilde{x} \leq \tilde{y}$, $w_0w\tilde{x}^+ \leq w_0w\tilde{y} = w_0\tilde{y}^+$. Lemma 2.2.2 implies that $w_0\tilde{y}^+ \uparrow w_0\tilde{x}^+$ so that $\tilde{x}^+ \uparrow \tilde{y}^+$. The last claim follows from [Jan03, II 6.5(5)]. \square

Lemma 2.2.4. *If $\tilde{w}, \tilde{w}' \in \widetilde{W}_1^+$, $\lambda, \nu \in X^*(T)$ with λ dominant, then $t_\nu w_0 t_\lambda \tilde{w} \leq w_0 t_\lambda \tilde{w}'$ and $t_{-\nu} w_0 t_\lambda \tilde{w}' \leq w_0 t_\lambda \tilde{w}$ imply that $\nu \in X^0(T)$ and $\tilde{w}' = t_\nu \tilde{w}$.*

Proof. Suppose that $t_\nu w_0 t_\lambda \tilde{w} \leq w_0 t_\lambda \tilde{w}'$ and $t_{-\nu} w_0 t_\lambda \tilde{w}' \leq w_0 t_\lambda \tilde{w}$. Lemma 2.2.2 implies that $w_0 t_\lambda \tilde{w}' \uparrow t_\nu w_0 t_\lambda \tilde{w}$ and $w_0 t_\lambda \tilde{w} \uparrow t_{-\nu} w_0 t_\lambda \tilde{w}'$. Combining these, we have that $w_0 t_\lambda \tilde{w}' \uparrow t_\nu w_0 t_\lambda \tilde{w} \uparrow w_0 t_\lambda \tilde{w}'$ which implies that $w_0 t_\lambda \tilde{w}' = t_\nu w_0 t_\lambda \tilde{w}$ or equivalently that $\tilde{w}' = t_{w_0\nu} \tilde{w}$. This implies that \tilde{w} and \tilde{w}' have the same image in W . Using that \tilde{w} and \tilde{w}' are both in \widetilde{W}_1^+ , we find that $t_{w_0\nu} = \tilde{w}'\tilde{w}^{-1} \in X^0(T)$ and in particular $w_0\nu = \nu$. \square

We now begin our analysis of certain elements of the admissible set which play an important role in our modularity results. For a simple root α , let $W_{a,\alpha}$ be the subgroup of W_a generated by s_α and t_α .

Lemma 2.2.5. *Let α be a simple root. Suppose that $\tilde{w}_\alpha \tilde{w}_1 \uparrow \tilde{w}_2 \uparrow \tilde{w}_1$ for some $\tilde{w}_\alpha \in W_{a,\alpha}$. Then $\tilde{w}_2 \in W_{a,\alpha} \tilde{w}_1$.*




FIGURE 1. The alcoves which are below $\tilde{w}_1(A_0)$ in the \uparrow order are those below the thickened red lines. The alcoves corresponding to $W_{a,\alpha} \tilde{w}_1$ are drawn in dotted blue lines.

Proof. Let $x \in A_0$. Then $\tilde{w}_1(x) - \tilde{w}_2(x)$ and $\tilde{w}_2(x) - \tilde{w}_\alpha \tilde{w}_1(x)$ are nonnegative linear combinations of positive simple roots. On the other hand, $\tilde{w}_1(x) - \tilde{w}_\alpha \tilde{w}_1(x)$ is a nonnegative multiple of α . This implies that so is $\tilde{w}_1(x) - \tilde{w}_2(x)$.

There is a series of hyperplane reflections $(s_i)_{i=1}^m$ such that

$$\tilde{w}_2 \uparrow s_1 \tilde{w}_2 \uparrow s_2 s_1 \tilde{w}_2 \uparrow \cdots \uparrow s_m \cdots s_2 s_1 \tilde{w}_2 = \tilde{w}_1.$$

If the corresponding positive roots are $(\alpha_i)_{i=1}^m$, then $\tilde{w}_1(x) - \tilde{w}_2(x)$ is a positive linear combination of the roots in $\{\alpha_i\}_{i=1}^m$. The above paragraph implies that $\alpha_i = \alpha$ for all i . \square

Let e be a positive integer. Recall that the $e\eta_0$ -admissible set $\text{Adm}(e\eta_0) \subset \tilde{W}$ is the subset of elements \tilde{w} such that $\tilde{w} \leq t_w(e\eta_0)$ for some $w \in W$.

Proposition 2.2.6. *The set $w^{-1}W_{a,\alpha}t_{e\eta_0}w \cap \text{Adm}(e\eta_0)$ consists of elements*

$$t_{w^{-1}(e\eta_0 - k\alpha)} \quad \text{for } 0 \leq k \leq e$$

and

$$\tilde{w}^{-1} s_\alpha t_{e\eta_0 - (k+1)\alpha} \tilde{w} \quad \text{for } 0 \leq k \leq e - 1,$$

where $\tilde{w} \in \tilde{W}_1^+$ is an element (unique up to $X^0(T)$) with image w in W .

Proof. It is easy to check that the listed elements lie in $w^{-1}W_{a,\alpha}t_{e\eta_0}w$. Furthermore, they are all less than or equal to either $t_{w^{-1}(e\eta_0)}$ or $t_{(s_\alpha w)^{-1}(e\eta_0)}$. Indeed, set $\tilde{z}_k \stackrel{\text{def}}{=} t_{(e-1)\eta_0 - k\alpha}$ and $\tilde{z}'_k \stackrel{\text{def}}{=}$

$t_{(e-1)\eta_0 - (e-1-k)\alpha} s_\alpha$. Then

$$(2.1) \quad t_{w^{-1}(e\eta_0 - k\alpha)} = (\tilde{w}_h \tilde{w})^{-1} w_0(\tilde{z}_k \tilde{w}) = (\tilde{w}_h \tilde{w})^{-1} (w_0 s_\alpha)(\tilde{z}'_k \tilde{w})$$

$$(2.2) \quad \tilde{w}^{-1} s_\alpha t_{e\eta_0 - (k+1)\alpha} \tilde{w} = (\tilde{w}_h \tilde{w})^{-1} (w_0 s_\alpha)(\tilde{z}_k \tilde{w}) = (\tilde{w}_h \tilde{w})^{-1} w_0(\tilde{z}'_k \tilde{w})$$

where $0 \leq k \leq e$ for the elements in (2.1) and $0 \leq k \leq e-1$ for the elements in (2.2). Both $\tilde{z}_k, \tilde{z}'_k \uparrow t_{(e-1)\eta_0}$ and, if $k \neq e$, one of them is in \widetilde{W}^+ . Wang's theorem implies that, for $0 \leq k \leq e-1$, one among $\tilde{z}_k, \tilde{z}'_k$ is less than or equal to $t_{(e-1)\eta_0}$. This implies that for $0 \leq k \leq e-1$ the elements (2.1), (2.2), with the exception of $t_{(s_\alpha w)^{-1}(e\eta_0)}$, are less than or equal to $t_{w^{-1}(e\eta_0)}$ in the Bruhat ordering. The exceptional element is less than or equal to itself.

We claim that any element in $w^{-1}W_{a,\alpha} t_{e\eta_0} w$ of length at most that of $t_{w^{-1}(e\eta_0)}$ is one of the listed elements. This would provide the reverse inclusion. For each positive root β and $\tilde{w} \in \widetilde{W}$, let

$$n_\beta(\tilde{w}) = \begin{cases} \lfloor \langle \tilde{w}(x), w^{-1}(\beta^\vee) \rangle \rfloor & \text{if } w(\beta) > 0 \\ \lfloor \langle \tilde{w}(x), w^{-1}(\beta^\vee) \rangle \rfloor + 1 & \text{if } w(\beta) < 0 \end{cases}$$

for any $x \in A_0$. Let $m_\beta(\tilde{w})$ be $|n_\beta(\tilde{w})|$. Then $\ell(\tilde{w})$ is the sum $\sum_{\beta > 0} m_\beta(\tilde{w})$ ([IM65, Proposition 1.23], see also [HN14, §1.3]). Let $d(\tilde{w})$ be the sum $m_\alpha(\tilde{w}) + \sum_{\beta > 0, \beta \neq \alpha} n_\beta(\tilde{w})$. The function $d(-)$ has three favorable properties: $d(\tilde{w}_1) \leq \ell(\tilde{w}_1)$ for all $\tilde{w}_1 \in \widetilde{W}$, $\ell(\tilde{w}_1) = d(\tilde{w}_1)$ if $w\tilde{w}_1 \in \widetilde{W}^+$ (in particular for $\tilde{w}_1 = t_{w^{-1}(e\eta_0)}$), and as we shall see next, $d(\tilde{w}_1) - m_\alpha(\tilde{w}_1) = \sum_{\beta > 0, \beta \neq \alpha} n_\beta(\tilde{w}_1)$ is the same for all $\tilde{w}_1 \in w^{-1}W_{a,\alpha} t_{e\eta_0} w$.

Fix x as above such that $\langle x, w^{-1}(\alpha^\vee) \rangle = \pm \frac{1}{2}$. Then for each $\tilde{w}_1 \in w^{-1}W_{a,\alpha} t_{e\eta_0} w$, $\tilde{w}_1(x) = x + w^{-1}(e\eta_0 - \frac{k}{2}\alpha)$ for a some $k \in \mathbb{Z}$. Moreover, the map $\tilde{w}_1 \mapsto k$ defines a bijection $w^{-1}W_{a,\alpha} t_{e\eta_0} w \rightarrow \mathbb{Z}$. We claim that

$$(2.3) \quad \sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle t_{w^{-1}(e\eta_0)}(x), w^{-1}(\beta^\vee) \rangle \rfloor = \sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle \tilde{w}_1(x), w^{-1}(\beta^\vee) \rangle \rfloor.$$

Assuming (2.3) for the moment, we obtain

$$\begin{aligned} \ell(t_{w^{-1}(e\eta_0)}) - d(\tilde{w}_1) &= d(t_{w^{-1}(e\eta_0)}) - d(\tilde{w}_1) \\ &= m_\alpha(t_{w^{-1}(e\eta_0)}) - m_\alpha(\tilde{w}_1) + \sum_{\beta > 0, \beta \neq \alpha} (\lfloor \langle t_{w^{-1}(e\eta_0)}(x), w^{-1}(\beta^\vee) \rangle \rfloor - \lfloor \langle \tilde{w}_1(x), w^{-1}(\beta^\vee) \rangle \rfloor) \\ &= m_\alpha(t_{w^{-1}(e\eta_0)}) - m_\alpha(\tilde{w}_1) \\ &= |e| - |e - k|. \end{aligned}$$

If $\ell(\tilde{w}_1) \leq \ell(t_{w^{-1}(e\eta_0)})$, then since $d(\tilde{w}_1) \leq \ell(\tilde{w}_1)$, $|e| \geq |e - k|$ so that $0 \leq k \leq 2e$. These $2e + 1$ values for k correspond to the $2e + 1$ listed elements. (See Figure 2 for the case of GL_3 and $e = 3$.)

It suffices to justify (2.3). We need to show that

$$\sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle x, w^{-1}(\beta^\vee) \rangle \rfloor = \sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle x - \frac{k}{2} w^{-1}(\alpha), w^{-1}(\beta^\vee) \rangle \rfloor,$$

or equivalently, letting $y = w(x)$, that

$$\sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle y, \beta^\vee \rangle \rfloor = \sum_{\beta > 0, \beta \neq \alpha} \lfloor \langle y - \frac{k}{2} \alpha, \beta^\vee \rangle \rfloor.$$

We can ignore roots β such that $\langle \alpha, \beta^\vee \rangle = 0$. The remaining positive roots come in pairs (β_-, β_+) where $\langle \alpha, \beta_-^\vee \rangle < 0$ and $\beta_+ = s_\alpha(\beta_-)$. Fix such a pair. The fact that $\langle \alpha, \beta_-^\vee \rangle + \langle \alpha, \beta_+^\vee \rangle = 0$ implies

that

$$\langle y, \beta_+^\vee \rangle + \langle y, \beta_-^\vee \rangle = \langle y - \frac{k}{2}\alpha, \beta_+^\vee \rangle + \langle y - \frac{k}{2}\alpha, \beta_-^\vee \rangle.$$

It suffices to show that

$$(2.4) \quad \{\langle y, \beta_+^\vee \rangle\} + \{\langle y, \beta_-^\vee \rangle\} = \{\langle y - \frac{k}{2}\alpha, \beta_+^\vee \rangle\} + \{\langle y - \frac{k}{2}\alpha, \beta_-^\vee \rangle\}$$

where $\{r\}$ denotes the fractional part $r - \lfloor r \rfloor$ of $r \in \mathbb{R}$. If $\langle \alpha, \beta_+^\vee \rangle$, and therefore $\langle \alpha, \beta_-^\vee \rangle$, is even, then (2.4) is clear. Now suppose that $\langle \alpha, \beta_+^\vee \rangle$ is odd. Recall that we chose x so that $\langle y, \alpha^\vee \rangle = \pm \frac{1}{2}$. Then $\beta_+ = \beta_- + \langle \beta_+, \alpha^\vee \rangle \alpha$ implies that $\{\langle y, \beta_-^\vee \rangle\} = \{\langle y, \beta_+^\vee \rangle + \frac{1}{2}\}$. We see that the terms of each side of (2.4) are the same (resp. permuted) when k is even (resp. odd). \square

Lemma 2.2.7. *Let α be a simple root and w be an element of W . If $\tilde{w} \in w^{-1}W_{a,\alpha}t_{e\eta_0}w \cap \text{Adm}(e\eta_0)$ and $\tilde{w} \leq t_{\sigma^{-1}(e\eta_0)}$ for some $\sigma \in W$, then $\sigma \in \{w, s_\alpha w\}$.*

Proof. Suppose that \tilde{w} is as in the statement. If $\tilde{w} = t_{w^{-1}(e\eta_0)}$ or $t_{(s_\alpha w)^{-1}(e\eta_0)}$ and $\tilde{w} \leq t_{\sigma^{-1}(e\eta_0)}$, then $\tilde{w} = t_{\sigma^{-1}(e\eta_0)}$ since $\ell(\tilde{w}) = \ell(t_{\sigma^{-1}(e\eta_0)})$ and the conclusion follows. Otherwise, $\tilde{w} \leq t_{w^{-1}(e\eta_0)}, t_{(s_\alpha w)^{-1}(e\eta_0)}$ by Proposition 2.2.6 and the first part of its proof (applying Proposition 2.2.6 with w both taken to be w and $s_\alpha w$ here). But by Corollary 4.3.3, there are at most two $\sigma \in W$ with $\tilde{w} \leq t_{\sigma^{-1}(e\eta_0)}$ (the reader can check that the proof of Corollary 4.3.3 only involves geometric properties of Pappas–Zhu local models, and does not make use of any of the results of this section). The conclusion follows. \square

Proposition 2.2.8. *Let α be a simple root and w be an element of W . Suppose that*

- (1) $\tilde{x} \in w^{-1}W_{a,\alpha}t_{e\eta_0}w \cap \text{Adm}(e\eta_0)$;
- (2) $\tilde{w}_2 \in \widetilde{W}^+$ and $\tilde{w}_\lambda \in \widetilde{W}_1^+$ such that $\tilde{w}_2 \uparrow \tilde{w}_h \tilde{w}_\lambda$; and
- (3) $\tilde{w}_2 \tilde{x} \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$.

Then \tilde{w}_2 equals $\tilde{w}_h \tilde{w}_\lambda$ and the image of \tilde{w}_λ in W is in the set $\{w, s_\alpha w\}$. Moreover, we can take \tilde{w}_λ as above to have image $w \in W$ (resp. $s_\alpha w \in W$) if and only if $\tilde{x} \neq t_{(s_\alpha w)^{-1}(e\eta_0)}$ (resp. $\tilde{x} \neq t_{w^{-1}(e\eta_0)}$).

Proof. There exists a dominant weight ω (unique up to $X^0(T)$) such that $t_{-\omega} \tilde{w}_2 \in \widetilde{W}_1^+$. Then $t_{-w_0(\omega)} \tilde{w}_\lambda \in \widetilde{W}^+$ and item (2) and [LLHL19, Proposition 4.1.2] give us $t_{-w_0(\omega)} \tilde{w}_\lambda \uparrow t_{-w_0(\omega)} \tilde{w}_h^{-1} \tilde{w}_2 = \tilde{w}_h^{-1} t_{-\omega} \tilde{w}_2$. Then Wang’s theorem implies that $t_{-w_0(\omega)} \tilde{w}_\lambda \leq \tilde{w}_h^{-1} t_{-\omega} \tilde{w}_2$, and so by [LLHL19, Lemma 4.1.9] we have

$$\begin{aligned} \tilde{x} \leq \tilde{w}_2^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda &= (t_{-\omega} \tilde{w}_2)^{-1} w_0 t_{(e-1)\eta_0} t_{-w_0(\omega)} \tilde{w}_\lambda \\ &\leq (t_{-\omega} \tilde{w}_2)^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_h^{-1} t_{-\omega} \tilde{w}_2 = t_{(w_0 w_2)^{-1}(e\eta_0)} \end{aligned}$$

where $w_2 \in W$ is the image of \tilde{w}_2 . Lemma 2.2.7 implies that $w_0 w_2 \in \{w, s_\alpha w\}$.

Suppose without loss of generality that $w_0 w_2 = w$. Let \tilde{w}_2 be $t_\omega \tilde{w}_h \tilde{w}$ where $\tilde{w} \in \widetilde{W}_1^+$ has image $w \in W$ and $\omega \in X^*(T)$ is dominant (ω in the last paragraph can be chosen to coincide with ω here). By (1), we let \tilde{x} be $(\tilde{w}_h \tilde{w})^{-1} w_0 \tilde{w}_\alpha t_{(e-1)\eta_0} \tilde{w}$ for some $\tilde{w}_\alpha \in W_{a,\alpha}$. Then (3) becomes $t_\omega w_0 \tilde{w}_\alpha t_{(e-1)\eta_0} \tilde{w} \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$ which implies by Lemma 2.2.3 that

$$t_{w_0(\omega)} \tilde{w}_\alpha t_{(e-1)\eta_0} \tilde{w} \uparrow t_{(e-1)\eta_0} \tilde{w}_\lambda,$$

which upon multiplying by $t_{-(e-1)\eta_0}$ and using item (2) and [LLHL19, Proposition 4.1.2] gives

$$\tilde{w}'_\alpha \tilde{w}_h^{-1} \tilde{w}_2 \uparrow \tilde{w}_\lambda \uparrow \tilde{w}_h^{-1} \tilde{w}_2$$

for some $\tilde{w}'_\alpha \in W_{a,\alpha}$ (using that $W_{a,\alpha}$ is stable under conjugation by $X^*(T)$). Then $\tilde{w}_\lambda \in W_{a,\alpha} \tilde{w}_h^{-1} \tilde{w}_2$ by Lemma 2.2.5, or equivalently $\tilde{w}_2 \in W_{a,-w_0(\alpha)} \tilde{w}_h \tilde{w}_\lambda$.




FIGURE 2. The $G = \text{GL}_3$ $3\eta_0$ -admissible set is in red. The set $\beta^{-1}W_{a,\alpha}t_{e\eta_0}\beta$ is in blue.

That $\tilde{w}_2 \in \tilde{W}^+$, $\tilde{w}_2 \uparrow \tilde{w}_h \tilde{w}_\lambda$, and $\tilde{w}_h \tilde{w}_\lambda \in \tilde{W}_1^+$ imply respectively that

$$0 \leq [\langle \tilde{w}_2(x), -w_0(\alpha^\vee) \rangle] \leq [\langle \tilde{w}_h \tilde{w}_\lambda(x), -w_0(\alpha^\vee) \rangle] = 0$$

for any $x \in A_0$, which forces equalities throughout. Combined with the fact that $\tilde{w}_2 \in W_{a,-w_0(\alpha)} \tilde{w}_h \tilde{w}_\lambda$, we see that $\tilde{w}_2 = \tilde{w}_h \tilde{w}_\lambda$. In particular, the image of \tilde{w}_λ in W is $w_0 w_2$.

The final part follows from the first part of the proof of Proposition 2.2.6. \square

Proposition 2.2.9. *Let α be a simple root and w be an element of W . If $\tilde{w}_1 \in w^{-1}W_{a,\alpha}t_{e\eta_0}w$ and $\tilde{w}_1 \leq \tilde{w}_2 \leq t_{w^{-1}(e\eta_0)}$, then $\tilde{w}_2 \in w^{-1}W_{a,\alpha}t_{e\eta_0}w$.*

Proof. By Lemma 2.2.3, the inequalities $\tilde{w}_1 \leq \tilde{w}_2 \leq t_{w^{-1}(e\eta_0)}$ imply that $w\tilde{w}_1 \uparrow s\tilde{w}_2 \uparrow wt_{w^{-1}(e\eta_0)}$ where $s \in W$ is the unique element such that $s\tilde{w}_2 \in \widetilde{W}^+$. Since $w\tilde{w}_1 \in W_{a,\alpha}wt_{w^{-1}(e\eta_0)}$, we deduce from Lemma 2.2.5 that $s\tilde{w}_2 \in W_{a,\alpha}wt_{w^{-1}(e\eta_0)}$ or equivalently, that $\tilde{w}_2 \in s^{-1}W_{a,\alpha}wt_{w^{-1}(e\eta_0)}$.

We now narrow the possibilities of s . Since $\tilde{w}_1 \leq \tilde{w}_2 \leq t_{s^{-1}(e\eta_0)}$, where the final inequality follows from [HH17, Corollary 4.4], $s \in \{w, s_\alpha w\}$ by Lemma 2.2.7. Combining with the above paragraph, $\tilde{w}_2 \in w^{-1}W_{a,\alpha}t_{e\eta_0}w$. \square

2.3. The weight part of a Serre-type conjecture for tame representations. The aim of this section, and of the following one, is to recollect the necessary notions to formulate the weight part for Serre conjectures, and to pursue a combinatorial study of the set of conjectural modular weights in terms of the geometry of the affine Weyl group.

2.3.1. Serre weights. Recall from 1.5.1 that G is a split group defined over \mathbb{F}_p , k_p is a finite étale \mathbb{F}_p -algebra, $G_0 = \text{Res}_{k_p/\mathbb{F}_p} G/k_p$ and \mathbb{F} contains the image of any ring homomorphism $k_p \rightarrow \overline{\mathbb{F}}_p$ so that $\underline{G} \stackrel{\text{def}}{=} (G_0)_{/\mathbb{F}} \cong G_{/\mathbb{F}}^{\text{Hom}(k_p, \mathbb{F})}$. Let G be $G_0(\mathbb{F}_p)$. A *Serre weight* (of G) is an absolutely irreducible \mathbb{F} -representation of G .

Let $\lambda \in X^*(\underline{T})$ be a dominant character. We write $W(\lambda)_{/\mathbb{F}}$ for the \underline{G} -module $\text{Ind}_{\underline{B}}^G w_0 \lambda$. Let $F(\lambda)$ denote the (irreducible) socle of the G -restriction of $W(\lambda)_{/\mathbb{F}}(\mathbb{F})$.

We define:

$$X_1(\underline{T}) \stackrel{\text{def}}{=} \{\lambda \in X^*(\underline{T}), 0 \leq \langle \lambda, \alpha^\vee \rangle \leq p-1 \text{ for all } \alpha \in \underline{\Delta}\}$$

which we call the set of p -restricted weights. Then the map $\lambda \mapsto F(\lambda)$ defines a bijection from $X_1(\underline{T})/(p-\pi)X^0(\underline{T})$ to the set of isomorphism classes of Serre weights of G (see [GHS18, Lemma 9.2.4]). We say that $\lambda \in X_1(\underline{T})$ is *regular p -restricted* if $\langle \lambda, \alpha^\vee \rangle < p-1$ for all $\alpha \in \underline{\Delta}$ and say a Serre weight $F(\lambda)$ is *regular* if λ is. Similarly we say that $F(\lambda)$ is *m -deep* if λ is m -deep.

To handle the combinatorics of Serre weights it is convenient to introduce the notion of p -alcoves and the p -dot action on them. A p -alcove is a connected component of $X^*(\underline{T}) \otimes_{\mathbb{Z}} \mathbb{R} \setminus (\bigcup_{(\alpha, pn)} (H_{\alpha, pn} - \eta_0))$ and we say that a p -alcove \underline{C} is *dominant* (resp. *p -restricted*) if $0 < \langle \lambda + \eta_0, \alpha^\vee \rangle$ (resp. if $0 < \langle \lambda + \eta_0, \alpha^\vee \rangle < p$) for all $\alpha \in \underline{\Delta}$ and $\lambda \in \underline{C}$. We write \underline{C}_0 for the dominant base p -alcove, i.e. the alcove characterized by $\lambda \in \underline{C}_0$ if and only if $0 < \langle \lambda + \eta_0, \alpha^\vee \rangle < p$ for all $\alpha \in \underline{\Phi}^+$. The *p -dot action* of \widetilde{W} on $X^*(\underline{T}) \otimes_{\mathbb{Z}} \mathbb{R}$ is defined by $\tilde{w} \cdot \lambda \stackrel{\text{def}}{=} w(\lambda + \eta_0 + p\nu) - \eta_0$ for $\tilde{w} = wt_\nu \in \widetilde{W}$ and $\lambda \in X^*(\underline{T}) \otimes_{\mathbb{Z}} \mathbb{R}$. Then we have

$$\widetilde{W}^+ = \{\tilde{w} \in \widetilde{W} : \tilde{w} \cdot \underline{C}_0 \text{ is dominant}\}$$

and

$$\widetilde{W}_1^+ = \{\tilde{w} \in \widetilde{W}^+ : \tilde{w} \cdot \underline{C}_0 \text{ is } p\text{-restricted}\}$$

and $\underline{\Omega}$ is the stabilizer of \underline{C}_0 for the dot action.

We have an equivalence relation on $\widetilde{W} \times X^*(\underline{T})$ defined by $(\tilde{w}, \omega) \sim (t_\nu \tilde{w}, \omega - \nu)$ for all $\nu \in X^0(\underline{T})$ ([LLHLMa, §2.2]). For $(\tilde{w}_1, \omega - \eta_0) \in \widetilde{W}_1^+ \times (X^*(\underline{T}) \cap \underline{C}_0)/\sim$, we define the Serre weight $F_{(\tilde{w}_1, \omega)} \stackrel{\text{def}}{=} F(\pi^{-1}(\tilde{w}_1) \cdot (\omega - \eta_0))$ (this only depends on the equivalence class of (\tilde{w}_1, ω)). We call the equivalence class of (\tilde{w}_1, ω) a *lowest alcove presentation* for the Serre weight $F_{(\tilde{w}_1, \omega)}$ and note that $F_{(\tilde{w}_1, \omega)}$ is m -deep if and only if $\omega - \eta_0$ is m -deep in alcove \underline{C}_0 . (We often implicitly choose a representative for a lowest alcove presentation to make *a priori* sense of an expression, though it is *a posteriori* independent of this choice.)

2.3.2. Deligne–Lusztig representations. To a *good* pair $(s, \mu) \in \underline{W} \times X^*(\underline{T})$ we attach a Deligne–Lusztig representation $R_s(\mu)$ of G defined over E (see [LLHL19, §2.2] and [GHS18, Proposition 9.2.1, 9.2.2]), where the representation $R_s(\mu)$ is there denoted $R(s, \mu)$. We call $(s, \mu - \eta_0)$ a *lowest alcove presentation* for $R_s(\mu)$ and say that $R_s(\mu)$ is N -*generic* if $\mu - \eta_0$ is N -deep in alcove \underline{C}_0 for $N \geq 0$. If $\mu - \eta_0$ is 1-deep in \underline{C}_0 then $R_s(\mu)$ is an irreducible representation. We say that a Deligne–Lusztig representation R is N -generic if there exists an isomorphism $R \cong R_s(\mu)$ where $R_s(\mu)$ is N -generic.

2.3.3. Tame inertial types. An inertial type (for K , over E) is the $\mathrm{GL}_n(E)$ -conjugacy class of an homomorphism $\tau : I_K \rightarrow \mathrm{GL}_n(E)$ with open kernel and which extends to the Weil group of G_K . An inertial type is *tame* if one (equivalently, any) homomorphism in the conjugacy class factors through the tame quotient of I_K .

Let $s \in \underline{W}$ and $\mu \in X^*(\underline{T}) \cap \underline{C}_0$. Associated to this data we have an integer r (the order of the element $s_0 s_1 \cdots s_{f-2} s_{f-1} \in W$), n -tuples $\mathbf{a}'^{(j')} \in \mathbb{Z}^n$ for $0 \leq j' \leq fr - 1$, and a tame inertial type $\tau(s, \mu + \eta_0) \stackrel{\text{def}}{=} \sum_{i=1}^n (\omega_{fr})^{\mathbf{a}'^{(0)}}$. (See [LLHLMa, Example 2.4.1, equations (5.2), (5.1)] for the explicit construction of the n -tuples $\mathbf{a}'^{(j')} \in \mathbb{Z}^n$.) We say that $\tau(s, \mu + \eta_0)$ is a *principal series type* if $r = 1$.

If $N \geq 0$ and μ is N -deep in alcove \underline{C}_0 , the pair (s, μ) is said to be an N -*generic lowest alcove presentation* for the tame inertial type $\tau(s, \mu + \eta_0)$. We say that a tame inertial type is N -generic if it admits an N -generic lowest alcove presentation. (Different pairs (s, μ) can give rise to isomorphic tame inertial types, see [LLHL19, Proposition 2.2.15].)

If (s, μ) is a lowest alcove presentation of τ , let $\tilde{w}(\tau) \stackrel{\text{def}}{=} t_{\mu + \eta_0} s \in \widetilde{W}$. (In particular, when writing $\tilde{w}(\tau)$ we use an implicit lowest alcove presentation for τ .)

Inertial \mathbb{F} -types are defined similarly with E replaced by \mathbb{F} . Tame inertial \mathbb{F} -types have analogous notions of lowest alcove presentations and genericity. If $\bar{\tau}$ is a tame inertial \mathbb{F} -type we write $[\bar{\tau}]$ to denote the tame inertial type over E obtained from $\bar{\tau}$ using the Teichmüller section $\mathbb{F}^\times \hookrightarrow \mathcal{O}^\times$.

Assume that μ is 1-deep in \underline{C}_0 . For each $0 \leq j' \leq fr - 1$ we define $s'_{\text{or}, j'}$ to be the (necessarily unique) element of W such that $(s'_{\text{or}, j'})^{-1}(\mathbf{a}'^{(j')}) \in \mathbb{Z}^n$ is dominant. (In the terminology of [LLHLM18], cf. Definition 2.6 of *loc. cit.*, the fr -tuple $(s'_{\text{or}, j'})_{0 \leq j' \leq fr-1}$ is the *orientation* of $(\mathbf{a}'^{(j')})_{0 \leq j' \leq fr-1}$.) We will need the observation that $(s'_{\text{or}, j})^{-1}(\mathbf{a}'^{(j)})$ equals $s_j^{-1}(\mu_j + \eta_{0, j})$ modulo p for all $0 \leq j \leq f - 1$.

2.3.4. Inertial local Langlands. Let K/\mathbb{Q}_p be a finite extension with ring of integers \mathcal{O}_K and residue field k . Let τ be a tame inertial type for K . By [CEG⁺16, Theorem 3.7] there exists an irreducible smooth representation $\sigma(\tau)$ of $\mathrm{GL}_n(\mathcal{O}_K)$ over E satisfying results towards the inertial local Langlands correspondence.

Let $k_p = k$, so that $G_0(\mathbb{F}_p) \cong \mathrm{GL}_n(k)$. When $\tau \cong \tau(s, \mu)$ with $\mu - \eta \in \underline{C}_0$ a 1-deep character, the representation $\sigma(\tau)$ can and will be taken to be (the inflation to $\mathrm{GL}_n(\mathcal{O}_K)$ of) $R_s(\mu)$ (see [LLHLMa, Theorem 2.5.3], [LLHL19, Corollary 2.3.5]).

The following definition will play a key role in our generalization of Herzig’s Serre weight conjecture.

Definition 2.3.1. Let \mathcal{R} denote the bijection on regular Serre weights given by $F(\lambda) \mapsto F(\tilde{w}_h \cdot \lambda)$. If $\bar{\tau} : I_K \rightarrow \mathrm{GL}_n(\mathbb{F})$ is a 1-generic tame inertial \mathbb{F} -type for I_K we define

$$W^?(\bar{\tau}) \stackrel{\text{def}}{=} \mathcal{R} \left(\mathrm{JH} \left(\overline{\sigma([\bar{\tau}]} \right) \otimes W((1 - e)w_0 \eta_0) \right).$$

2.3.5. *Compatibilities of lowest alcove presentations.* Recall the canonical isomorphism $\widetilde{W}/W_a \xrightarrow{\sim} X^*(\underline{Z})$. Let $\zeta \in X^*(\underline{Z})$. We say that an element $\tilde{w} \in \widetilde{W}$ is ζ -compatible if it corresponds to ζ via the isomorphism $\widetilde{W}/W_a \xrightarrow{\sim} X^*(\underline{Z})$.

A lowest alcove presentation (s, μ) for a tame inertial type τ for K over E or a Deligne–Lusztig representation R is ζ -compatible if $t_{\mu+\eta_0}s \in \widetilde{W}$ is ζ -compatible. A lowest alcove presentation (s, μ) of a tame inertial \mathbb{F} -type is ζ -compatible if $t_{\mu-(e-1)\eta_0}s \in \widetilde{W}$ is ζ -compatible. (If τ is a tame inertial type and $\bar{\tau}$ is the tame inertial \mathbb{F} -type obtained by reduction, the same lowest alcove presentation of τ and $\bar{\tau}$ are compatible with elements of $X^*(\underline{Z})$ that differ by $\eta_0|_{\underline{Z}}$.) A lowest alcove presentation (\tilde{w}_1, ω) for Serre weight is ζ -compatible if the element $t_{\omega-\eta_0}\tilde{w}_1 \in \widetilde{W}$ is ζ -compatible. Finally, lowest alcove presentations (of possibly different types of objects) are compatible if they are all ζ -compatible for some $\zeta \in X^*(\underline{Z})$.

2.3.6. *L -parameters.* Recall from §1.5.1 the finite étale \mathbb{Q}_p -algebra F_p^+ . We adapt the constructions of tame inertial types and the inertial local Langlands above to arbitrary S_p . We assume that E contains the image of any homomorphism $F_p^+ \rightarrow \overline{\mathbb{Q}_p}$. Let

$$\underline{G}^\vee \stackrel{\text{def}}{=} \prod_{F_p^+ \rightarrow E} G_{/\mathcal{O}}^\vee$$

be the dual group of $\text{Res}_{F_p^+/\mathbb{Q}_p}(G_{/F_p^+})$ and ${}^L\underline{G} \stackrel{\text{def}}{=} \underline{G}^\vee \rtimes \text{Gal}(E/\mathbb{Q}_p)$ the Langlands dual group of $\text{Res}_{F_p^+/\mathbb{Q}_p}(G_{/F_p^+})$ (where $\text{Gal}(E/\mathbb{Q}_p)$ acts on the set $\{F_p^+ \rightarrow E\}$ by post-composition). An L -parameter (over E) is a $\underline{G}^\vee(E)$ -conjugacy class of an L -homomorphism, i.e. of a continuous homomorphism $\rho : G_{\mathbb{Q}_p} \rightarrow {}^L\underline{G}(E)$ which is compatible with the projection to $\text{Gal}(E/\mathbb{Q}_p)$. An inertial L -parameter is a $\underline{G}^\vee(E)$ -conjugacy class of an homomorphism $\tau : I_{\mathbb{Q}_p} \rightarrow \underline{G}^\vee(E)$ with open kernel, and which admits an extension to an L -homomorphism $G_{\mathbb{Q}_p} \rightarrow {}^L\underline{G}(E)$. An inertial L -parameter is *tame* if some (equivalently, any) representative in its equivalence class factors through the tame quotient of $I_{\mathbb{Q}_p}$.

The argument of [GHS18, Lemmas 9.4.1, 9.4.5] carries over in our setting and we have a bijection between L -parameters (resp. tame inertial L -parameters) and collections of the form $(\rho_v)_{v \in S_p}$ (resp. of the form $(\tau_v)_{v \in S_p}$) where for all $v \in S_p$ the element $\rho_v : G_{F_v^+} \rightarrow \text{GL}_n(E)$ is a continuous Galois representation (resp. the element $\tau_v : I_{F_v^+} \rightarrow \text{GL}_n(E)$ is a tame inertial type for F_v^+). (This bijection depends on a choice of isomorphisms $\overline{F_v^+} \xrightarrow{\sim} \overline{\mathbb{Q}_p}$ for all $v \in S_p$.) We have similar notions for L -parameters (resp. inertial L -parameters) over \mathbb{F} .

In this setting, given a tame inertial L -parameter τ corresponding to the collection of tame inertial types $(\tau_v)_{v \in S_p}$, we let $\sigma(\tau)$ be the irreducible smooth E -valued representation of $\text{GL}_n(\mathcal{O}_p)$ given by $\otimes_{v \in S_p} \sigma(\tau_v)$, where $\sigma(\tau_v)$ is the smooth representation corresponding to τ_v via the inertial local Langlands correspondence appearing in §2.3.4.

2.4. **Combinatorics of Serre weights.** In this section, we apply the results of §2.2 on extended affine Weyl groups to analyze the combinatorics of the Serre weight sets defined in §2.3. We assume for simplicity that $F_p^+ = K$, but the results herein do not require this. Given tame inertial types τ and $\bar{\rho}^{\text{sp}}$ over E and \mathbb{F} , respectively, with fixed compatible lowest alcove presentations, we define $\tilde{w}(\bar{\rho}^{\text{sp}}, \tau)$ to be $\tilde{w}(\tau)^{-1}\tilde{w}(\bar{\rho}^{\text{sp}}) \in t_{e\eta_0}\widetilde{W}_a$.

The following two results follow readily from [LLHLMa, Proposition 2.3.7].

Proposition 2.4.1. *Suppose we fix a $2h_{\eta_0}$ -generic lowest alcove presentation of a tame inertial type τ . The map*

$$(2.5) \quad (\tilde{w}_\lambda, \tilde{w}_2) \longmapsto F_{(\tilde{w}_\lambda, \tilde{w}(\tau)\tilde{w}_2^{-1}(0))}$$

induces a bijection between

- the set of pairs $(\tilde{w}_\lambda, \tilde{w}_2)$, modulo the diagonal action of $X^0(T)$, with $\tilde{w}_\lambda \in \widetilde{W}_1^+$ and $\tilde{w}_2 \in \widetilde{W}^+$ such that $\tilde{w}_\lambda \uparrow \tilde{w}_h^{-1}\tilde{w}_2$; and
- elements of $\text{JH}(\overline{\sigma}(\tau))$.

Moreover, these lowest alcove presentations of Serre weights are compatible with that of τ . Finally, the weight corresponding to $(\tilde{w}_\lambda, \tilde{w}_h\tilde{w}_\lambda)$ appears as a Jordan–Hölder factor with multiplicity one.

Proposition 2.4.2. *Suppose we fix an $(\max\{2, e\}h_{\eta_0})$ -generic lowest alcove presentation of a tame inertial type $\overline{\rho}^{\text{sp}}$ over \mathbb{F} . The map*

$$(\tilde{w}_\lambda, \tilde{w}_2) \longmapsto F_{(\tilde{w}_\lambda, \tilde{w}^*(\overline{\rho}^{\text{sp}})\tilde{w}_2^{-1}(0))}.$$

induces a bijection between

- pairs $(\tilde{w}_\lambda, \tilde{w}_2)$ with $\tilde{w}_\lambda \in \widetilde{W}_1^+$ and $\tilde{w}_2 \in \widetilde{W}^+$, up to the diagonal $X^0(\underline{T})$ -action, such that $\tilde{w}_2 \uparrow t_{(e-1)\eta_0}\tilde{w}_\lambda$; and
- elements of $W^?(\overline{\rho}^{\text{sp}})$.

Moreover, these lowest alcove presentations of Serre weights are compatible with that of $\overline{\rho}^{\text{sp}}$.

The following definition is central to this paper.

Definition 2.4.3 (Extremal weights). *Suppose we fix a $(\max\{2, e\}h_{\eta_0})$ -generic lowest alcove presentation for a tame inertial type $\overline{\rho}^{\text{sp}}$ over \mathbb{F} . Let w be an element of W and $\tilde{w} \in \widetilde{W}_1^+$ be an element (unique up to $X^0(\underline{T})$) whose image in W is w . The weight*

$$F_{(\tilde{w}, \tilde{w}(\overline{\rho}^{\text{sp}})\tilde{w}^{-1}(-(e-1)\eta_0))} \in W^?(\overline{\rho}^{\text{sp}})$$

is called the *extremal weight of $\overline{\rho}^{\text{sp}}$ corresponding to w* . Let $W_{\text{extr}}(\overline{\rho}^{\text{sp}})$ be the set of all extremal weights of $\overline{\rho}^{\text{sp}}$. (While the extremal weight corresponding to w depends on the choice of lowest alcove presentation, the set of all extremal weights does not.)

Remark 2.4.4. If $\overline{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ is semisimple and $2h_{\eta_0}$ -generic, and K is unramified, the notion of obvious weight for $\overline{\rho}$ corresponding to w ([LLHLMa, Definition 2.6.3]) and of extremal weight for $\overline{\rho}$ corresponding to w coincide, and the set $W_{\text{extr}}(\overline{\rho}|_I)$ is the set defined in [GHS18, Definition 7.1.3].

The following combinatorial result relates the set $W^?(\overline{\rho}^{\text{sp}})$ and the admissible set and is key to weight elimination.

Proposition 2.4.5. *Suppose we fix an $(\max\{2, e\}h_{\eta_0})$ -generic lowest alcove presentation for $\tilde{w}(\overline{\rho}^{\text{sp}})$. Let $(\tilde{w}_\lambda, \omega)$ be a compatible lowest alcove presentation of a $3h_{\eta_0}$ -deep Serre weight σ . Then $\omega = \tilde{w}(\overline{\rho})\tilde{w}^{-1}(0)$ for a unique $\tilde{w} \in \widetilde{W}^+$. Let τ be the tame inertial type over E with $\tilde{w}(\overline{\rho}^{\text{sp}}, \tau) = (\tilde{w}_h\tilde{w}_\lambda)^{-1}\omega_0\tilde{w}$ for some (necessarily compatible) lowest alcove presentation. Then*

- (1) $\sigma \in \text{JH}(\overline{\sigma}(\tau))$; and
- (2) $(\tilde{w}_h\tilde{w}_\lambda)^{-1}\omega_0\tilde{w} \in \text{Adm}(e\eta_0)$ implies that $\sigma \in W^?(\overline{\rho}^{\text{sp}})$.

Proof. By definition of τ , we have that $\tilde{w}(\tau)(\tilde{w}_h\tilde{w}_\lambda)^{-1}(0) = \tilde{w}(\overline{\rho}^{\text{sp}})\tilde{w}^{-1}(0)$. Note that the lowest alcove presentation of τ is $2h_{\eta_0}$ -generic by the depth assumption on σ . Then σ corresponds to the pair $(\tilde{w}_\lambda, \tilde{w}_h\tilde{w}_\lambda)$ in (2.5).

Suppose that $(\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 \tilde{w} \in \text{Adm}(e\eta_0)$. If we let $w_\lambda \in W$ be the image of \tilde{w}_λ , then we claim that $w_\lambda (\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 \tilde{w} \in \widetilde{W}^+$. Indeed since $\tilde{w}_\lambda \in \widetilde{W}_1^+$, $w_0 w_\lambda (\tilde{w}_h \tilde{w}_\lambda)^{-1}$ is an antidominant translation so that $w_0 w_\lambda (\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 \tilde{w} \cdot C_0$ is in the antidominant Weyl chamber. By [HH17, Corollary 4.4], we have that

$$(\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 \tilde{w} \leq t_{w_\lambda^{-1}(e\eta_0)} = (\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda.$$

Since these expressions are reduced by [LLHL19, Lemma 4.9], we conclude that $\tilde{w} \leq t_{(e-1)\eta_0} \tilde{w}_\lambda$ which implies that $\tilde{w} \uparrow t_{(e-1)\eta_0} \tilde{w}_\lambda$. We conclude from Proposition 2.4.2 that $\sigma \in W^?(\bar{\rho}^{\text{SP}})$. \square

Denote by $W^?(\bar{\rho}^{\text{SP}}, \tau)$ the intersection $W^?(\bar{\rho}^{\text{SP}}) \cap \text{JH}(\bar{\sigma}(\tau))$.

Proposition 2.4.6. *Suppose we fix compatible $(\max\{2, e\}h_{\eta_0})$ -generic and $2h_{\eta_0}$ -generic lowest alcove presentations of tame inertial types $\bar{\rho}^{\text{SP}}$ and τ over \mathbb{F} and E , respectively. Then the set $W^?(\bar{\rho}^{\text{SP}}, \tau)$ is exactly the set of weights in (2.5) such that $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$.*

Proof. Consider an element $\sigma \in \text{JH}(\bar{\sigma}(\tau))$. Let \tilde{w}_λ and \tilde{w}_2 be as in Proposition 2.4.1. By Proposition 2.4.2 and uniqueness of compatible lowest alcove presentations (see [LLHLMa, Lemma 2.2.3]), $\sigma \in W^?(\bar{\rho}^{\text{SP}})$ if and only if there exist $\tilde{s}_2 \in \widetilde{W}^+$ with $\tilde{s}_2 \uparrow t_{(e-1)\eta_0} \tilde{w}_\lambda$ such that

$$\tilde{w}(\tau) \tilde{w}_2^{-1}(0) = \tilde{w}(\bar{\rho}^{\text{SP}}) \tilde{s}_2^{-1}(0),$$

or equivalently, $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \in W \tilde{s}_2$.

We now show that there exists $\tilde{s}_2 \in \widetilde{W}^+$ with $\tilde{s}_2 \uparrow t_{(e-1)\eta_0} \tilde{w}_\lambda$ such that $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \in W \tilde{s}_2$ if and only if $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$. First suppose that such an \tilde{s}_2 exists. This implies that

$$\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0 \tilde{s}_2 \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda,$$

where the second inequality follows from the fact that $\tilde{s}_2 \leq t_{(e-1)\eta_0} \tilde{w}_\lambda$ by Wang's theorem. Conversely, if $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$, then using that $w_0(t_{(e-1)\eta_0} \tilde{w}_\lambda)$ is a reduced factorization, $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) = w \tilde{s}_2$ for some $\tilde{s}_2 \in \widetilde{W}^+$ with $\tilde{s}_2 \leq t_{(e-1)\eta_0} \tilde{w}_\lambda$ (or equivalently $\tilde{s}_2 \uparrow t_{(e-1)\eta_0} \tilde{w}_\lambda$ by Wang's theorem) and $w \in W$. \square

Corollary 2.4.7. *Suppose that tame inertial types $\bar{\rho}^{\text{SP}}$ and τ over \mathbb{F} and E have compatible $(\max\{2, e\}h_{\eta_0})$ -generic and $2h_{\eta_0}$ -generic lowest alcove presentations, respectively. If $W^?(\bar{\rho}^{\text{SP}}, \tau)$ is nonempty, then $\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \in \text{Adm}(e\eta_0)$.*

Proof. As in the statement of the corollary, we fix compatible lowest alcove presentations for $\bar{\rho}^{\text{SP}}$ and τ , respectively. If $W^?(\bar{\rho}^{\text{SP}}, \tau)$ is nonempty, by Proposition 2.4.6 we have that $\tilde{w}_2 \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$ for some $\tilde{w}_2 \in \widetilde{W}^+$ and $\tilde{w}_\lambda \in \widetilde{W}_1^+$ with $\tilde{w}_\lambda \uparrow \tilde{w}_h^{-1} \tilde{w}_2$. Then $\tilde{w}_2 \uparrow \tilde{w}_h \tilde{w}_\lambda$ by [LLHL19, Proposition 4.1.2] so that $\tilde{w}_2 \leq \tilde{w}_h \tilde{w}_\lambda$. Since $\tilde{w}_2^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$ and $(\tilde{w}_h \tilde{w}_\lambda)^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda$ are reduced expressions by [LLHL19, Lemma 4.1.9], we have that

$$\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq \tilde{w}_2^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda \leq \tilde{w}_\lambda^{-1} \tilde{w}_h^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}_\lambda = t_{w_\lambda^{-1}(e\eta_0)}.$$

\square

We now establish some results which will be used to prove modularity of certain Serre weights.

Proposition 2.4.8. *Suppose that $\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \in w^{-1} W_{a,\alpha} t_{e\eta_0} w \cap \text{Adm}(e\eta_0)$ for compatible $(\max\{2, e\}h_{\eta_0})$ -generic and $2h_{\eta_0}$ -generic lowest alcove presentations for tame inertial types $\bar{\rho}^{\text{SP}}$ and τ over \mathbb{F} and E , respectively.*

Then $W^?(\bar{\rho}^{\text{SP}}, \tau)$ equals

$$\begin{cases} \{F_{(\tilde{w}, \tilde{w}(\tau)\tilde{w}^{-1}\tilde{w}_h^{-1}(0))}\} & \text{if } \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) = t_{w^{-1}(e\eta_0)} \\ \{F_{(\widetilde{s_\alpha w}, \widetilde{w}(\tau)\widetilde{s_\alpha w}^{-1}\widetilde{w}_h^{-1}(0))}\} & \text{if } \tilde{w}(\bar{\rho}^{\text{SP}}, \tau) = t_{(s_\alpha w)^{-1}(e\eta_0)} \\ \{F_{(\tilde{w}, \tilde{w}(\tau)\tilde{w}^{-1}\tilde{w}_h^{-1}(0)), F_{(\widetilde{s_\alpha w}, \widetilde{w}(\tau)\widetilde{s_\alpha w}^{-1}\widetilde{w}_h^{-1}(0))}\} & \text{otherwise.} \end{cases}$$

Moreover, each weight appears as a Jordan–Hölder factor of $\bar{\sigma}(\tau)$ with multiplicity one.

Proof. Suppose that a weight σ of the form (2.5) is in $\text{JH}(\bar{\sigma}(\tau))$. Then by Proposition 2.4.6, $\sigma \in W^?(\bar{\rho}^{\text{SP}})$ if and only if $\tilde{w}_2\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \leq w_0t_{(e-1)\eta_0}\tilde{w}_\lambda$. By Proposition 2.2.8, $\tilde{w}_h^{-1}\tilde{w}_2$ and \tilde{w}_λ are both either \tilde{w} or $\widetilde{s_\alpha w}$. The last part of Proposition 2.2.8 implies the inclusion of $W^?(\bar{\rho}^{\text{SP}}, \tau)$ in the casewise defined sets.

On the other hand, using Proposition 2.2.6 and

$$(2.6) \quad t_{w^{-1}(e\eta_0-k\alpha)} = \tilde{w}^{-1}\tilde{w}_h^{-1}w_0t_{(e-1)\eta_0-k\alpha}\tilde{w} = \widetilde{s_\alpha w}^{-1}\tilde{w}_h^{-1}w_0t_{(e-1)\eta_0-(e-k)\alpha}\widetilde{s_\alpha w}$$

and

$$(2.7) \quad \tilde{w}^{-1}s_\alpha t_{e\eta_0-(k+1)\alpha}\tilde{w} = \tilde{w}^{-1}\tilde{w}_h^{-1}w_0s_\alpha t_{(e-1)\eta_0-k\alpha}\tilde{w} = \widetilde{s_\alpha w}^{-1}\tilde{w}_h^{-1}w_0s_\alpha t_{(e-1)\eta_0-(e-k-1)\alpha}\widetilde{s_\alpha w}.$$

we have that $\tilde{w}_h\tilde{w}(\bar{\rho}^{\text{SP}}, \tau)$ is either $w_0t_{(e-1)\eta_0-k\alpha}\tilde{w}$ or $w_0s_\alpha t_{(e-1)\eta_0-k\alpha}\tilde{w}$, which is less than or equal to $w_0t_{(e-1)\eta_0}\tilde{w}$ if $k \neq e$. This implies that $F_{(\tilde{w}, \tilde{w}(\tau)\tilde{w}^{-1}\tilde{w}_h^{-1}(0))} \in W^?(\bar{\rho}^{\text{SP}}, \tau)$ if $\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \neq t_{(s_\alpha w)^{-1}(e\eta_0)}$. Similarly, $\tilde{w}_h\widetilde{s_\alpha w}(\bar{\rho}^{\text{SP}}, \tau)$ is either $w_0t_{(e-1)\eta_0-(e-k)\alpha}\widetilde{s_\alpha w}$ or $w_0s_\alpha t_{(e-1)\eta_0-(e-k-1)\alpha}\widetilde{s_\alpha w}$, so that $F_{(\widetilde{s_\alpha w}, \widetilde{w}(\tau)\widetilde{s_\alpha w}^{-1}\widetilde{w}_h^{-1}(0))} \in W^?(\bar{\rho}^{\text{SP}}, \tau)$ if $\tilde{w}(\bar{\rho}^{\text{SP}}, \tau) \neq t_{w^{-1}(e\eta_0)}$. This gives the reverse inclusion. The multiplicity statement follows from that of Proposition 2.4.1. \square

Proposition 2.4.9. *Suppose we fix an $(e+2)h_{\eta_0}$ -generic lowest alcove presentation of a tame inertial type $\bar{\rho}^{\text{SP}}$ over \mathbb{F} . For $0 \leq k \leq e$, let τ_{2k} be the tame inertial type over E with compatible lowest alcove presentation such that $\tilde{w}(\tau) = \tilde{w}(\bar{\rho}^{\text{SP}})t_{w^{-1}(k\alpha-e\eta_0)}$. For $0 \leq k \leq e-1$, let τ_{2k+1} be the tame inertial type over E with compatible lowest alcove presentation such that $\tilde{w}(\tau) = \tilde{w}(\bar{\rho}^{\text{SP}})\tilde{w}^{-1}t_{k\alpha-e\eta_0}s_\alpha\tilde{w}$.*

For $0 \leq k \leq e-1$, let σ_{2k} be

$$F_{(\tilde{w}, \tilde{w}(\bar{\rho}^{\text{SP}})\tilde{w}^{-1}(k\alpha-(e-1)\eta_0))},$$

and let σ_{2k+1} be

$$F_{(\widetilde{s_\alpha w}, \widetilde{w}(\bar{\rho}^{\text{SP}})\widetilde{s_\alpha w}^{-1}((e-k-1)\alpha-(e-1)\eta_0))}.$$

Then for $0 \leq m \leq 2e$, $W^?(\bar{\rho}^{\text{SP}}, \tau_m) = \{\sigma_{m-1}, \sigma_m\}$ (where σ_ℓ should be omitted for $\ell = -1$ or $2e$). Moreover, σ_{m-1} and σ_m appear as Jordan–Hölder factors of $\bar{\sigma}(\tau_m)$ with multiplicity one.

Proof. This follows from Propositions 2.2.6 and 2.4.8 using (2.6) and (2.7) noting that τ_m is $2h_{\eta_0}$ -generic for all m . \square

2.4.1. The case of GL_2 : a comparison with Schein’s recipe. [Sch08] explicitly describes a set of Serre weights for a semisimple $\bar{\rho} : G_K \rightarrow \text{GL}_2(\mathbb{F})$ with K possibly ramified over \mathbb{Q}_p in terms of a “reflection operation” \mathcal{R}^δ similar to \mathcal{R} above. We compare this description in generic cases with the set $W^?(\bar{\rho})$ defined in §2.3.

Assume $e \leq p-1$ and let $\bar{\rho} : G_K \rightarrow \text{GL}_2(\mathbb{F})$ be semisimple. In [Sch08, Conjecture 1], a set of Serre weights is associated to $\bar{\rho}$, in terms of a reflection operation denoted as \mathcal{R}_p^δ in *loc. cit.* The superscript δ is an element in $\{0, \dots, e-1\}^{\mathcal{J}}$ and leads to the notion of δ -regular weight:

Definition 2.4.10 ([Sch08]). A Serre weight $F(\lambda)$ is δ -regular if $p-1-\langle \lambda, \alpha_j^\vee \rangle \in \{1, \dots, p\} + (2\delta_j - e+1)$ for all $j \in \mathcal{J}$. (Note that this definition does not depend of the lift of $\lambda \in X_1(\underline{T})/(p-\pi)X_0(\underline{T})$.)

A direct computation shows that if $\lambda \in \underline{C}_0$ is $(e-1)$ -deep then λ is δ -regular for any $\delta \in \{0, \dots, e-1\}^{\mathcal{J}}$, and moreover $\lambda + \nu \in \underline{C}_0$ for any weight ν appearing in $(W((1-e)w_0(\eta_0))_{/\mathbb{F}})|_{\underline{T}}$.

Let now $\lambda \in \underline{C}_0$ be $(e-1)$ -deep. A direct computation using the definition of $\mathcal{R}_{\mathfrak{p}}^{\delta}$ for δ -regular weights yields:

$$\mathcal{R}_{\mathfrak{p}}^{\delta}(F(\lambda)) = F(\tilde{w}_h \cdot (\lambda - (e-1)\eta_0 + \sum_{j \in \mathcal{J}} \delta_j \alpha_j))$$

and hence

$$\bigcup_{\delta \in \{0, \dots, e-1\}^{\mathcal{J}}} \mathcal{R}_{\mathfrak{p}}^{\delta}(F(\lambda)) = \mathcal{R}(\text{JH}(F(\lambda)) \otimes W((1-e)w_0\eta_0))$$

by the translation principle (cf. [LMS, Proposition 3.3]). From [DL21, Proposition 2.15] (or Proposition 2.4.1 above when $e \geq 2$), noting that for an e -generic Deligne–Lusztig representation R all the Serre weights $F \in \text{JH}(R)$ are $e-1$ -deep, we deduce

Proposition 2.4.11. *Let $\bar{\rho} : G_K \rightarrow \text{GL}_2(\mathbb{F})$ be semisimple and e -generic. Then the set of weights $W_{\mathfrak{p}}^{\delta}(\bar{\rho})$ defined in [Sch08, Conjecture 1(1)] coincides with the set $W^{\delta}(\bar{\rho})$ of Definition 2.3.1 above.*

3. BREUIL–KISIN MODULES

3.1. Moduli of Breuil–Kisin modules and local models. In this section, we introduce background on Breuil–Kisin modules with tame descent. We closely follow [LLHLMa, §5] making the necessary modifications to allow K/\mathbb{Q}_p to be ramified. We will generally admit proofs as the generalizations are straightforward. Throughout this section, we take $G = \mathrm{GL}_n$.

Let K/\mathbb{Q}_p be finite. We let K_0 be the maximal unramified subextension of K , with $f \stackrel{\mathrm{def}}{=} [K_0 : \mathbb{Q}_p]$ and $e \stackrel{\mathrm{def}}{=} [K : K_0]$. Let k denote the residue field of K , of cardinality p^f and which coincide with the residue field of K_0 . Let $W(k)$ be ring of Witt vectors of k , which is also the ring of integers of K_0 . We denote the arithmetic Frobenius automorphism on $W(k)$ by φ , which acts as raising to p -th power on the residue field.

We fix a uniformizer $\pi_K \in \overline{K}$ of K . Let $E(v) \in W(k)[v]$ be the minimal polynomial for π_K over K_0 , of degree e .

Let $\mathcal{J}_K = \mathrm{Hom}(K, E)$ and $\mathcal{J} = \mathrm{Hom}(k, \mathbb{F}) = \mathrm{Hom}(K_0, E)$. Recall that we have fixed an embedding $\sigma_0 : K_0 \hookrightarrow E$, hence an identification $\mathcal{J} \xrightarrow{\sim} \mathbb{Z}/f\mathbb{Z}$ given by $\sigma_j \stackrel{\mathrm{def}}{=} \sigma_0 \circ \varphi^{-j} \mapsto j$.

Let τ be a tame inertial type having a 1-generic lowest alcove presentation $(s, \mu) \in W^{\mathcal{J}} \times X^*(T)^{\mathcal{J}}$, which we now fix throughout this section. Recall from [LLHLMa, Example 2.4.1] that we have a combinatorial data attached to (s, μ) , in particular the element $s_{\tau} \in W$ (when $K = \mathbb{Q}_p$, this is the *niveau* of τ).

Let r be the order of s_{τ} . We write K' for the subfield of \overline{K} which is unramified of degree r over K , k' for its residue field, and K'_0 denote maximal unramified subextension of K' . Set $\mathcal{J}_{K'} = \mathrm{Hom}(K', E)$, $\mathcal{J}' \stackrel{\mathrm{def}}{=} \mathrm{Hom}(k', \mathbb{F})$. Let $f' \stackrel{\mathrm{def}}{=} fr$, $e' \stackrel{\mathrm{def}}{=} p^{f'} - 1$ and fix an embedding $\sigma'_0 : K'_0 \hookrightarrow E$ which extends $\sigma_0 : K_0 \hookrightarrow E$, so that the identification $\mathcal{J}' \cong \mathbb{Z}/f'\mathbb{Z}$ given by $\sigma_{j'} \stackrel{\mathrm{def}}{=} \sigma'_0 \circ \varphi^{-j'} \mapsto j'$ induces the natural surjection $\mathbb{Z}/f'\mathbb{Z} \twoheadrightarrow \mathbb{Z}/f\mathbb{Z}$ when considering the restriction of embedding from K'_0 to K_0 .

We fix an e' -root $\pi_{K'} \in \overline{K}$ of π_K and set $L' \stackrel{\mathrm{def}}{=} K'(\pi_{K'})$. Let $\Delta' \stackrel{\mathrm{def}}{=} \mathrm{Gal}(L'/K') \subset \Delta \stackrel{\mathrm{def}}{=} \mathrm{Gal}(L'/K)$. We set $\omega_{K'}(g) \stackrel{\mathrm{def}}{=} \frac{g(\pi_{K'})}{\pi_{K'}}$ for $g \in \Delta'$; then $\omega_{K'}$ does not depend on the choice of π' . Composing with $\sigma'_{j'} \in \mathcal{J}'$, we get a corresponding character $\omega_{K', \sigma'_{j'}} : \Delta' \rightarrow \mathcal{O}^{\times}$ which will also be seen as a character of $I_{K'} = I_K$. For $j' = 0$ we set $\omega_{j'} \stackrel{\mathrm{def}}{=} \omega_{K', \sigma'_{j'}}$.

Let R be an \mathcal{O} -algebra. Let $\mathfrak{S}_{L'} \stackrel{\mathrm{def}}{=} W(k')[[u']]$ and $\mathfrak{S}_{L', R} \stackrel{\mathrm{def}}{=} (W(k') \otimes_{\mathbb{Z}_p} R)[[u']]$. As usual, $\varphi : \mathfrak{S}_{L', R} \rightarrow \mathfrak{S}_{L', R}$ acts as Frobenius on $W(k')$, trivially on R , and sends u' to $(u')^p$. Note that for any $\mathfrak{S}_{L', R}$ -module \mathfrak{M} , we have the standard $R[[u']]$ -linear decomposition $\mathfrak{M} \cong \bigoplus_{j' \in \mathcal{J}'} \mathfrak{M}^{(j')}$, induced by the maps $W(k') \otimes_{\mathbb{Z}_p} R \rightarrow R$ defined by $x \otimes r \mapsto \sigma_{j'}(x)r$ for $j' \in \mathcal{J}'$.

We endow $\mathfrak{S}_{L', R}$ with an action of Δ as follows: for any g in Δ' , $g(u') = \omega_{K'}(g)u'$ and g acts trivially on the coefficients; if $\sigma^f \in \mathrm{Gal}(L'/K)$ is the lift of the p^f -Frobenius on $W(k')$ which fixes $\pi_{K'}$, then σ^f is a generator for $\mathrm{Gal}(K'/K)$, acting in natural way on $W(k')$ and trivially on both u' and R . Set $v = (u')^{e'}$, and define $\mathfrak{S}_R \stackrel{\mathrm{def}}{=} (\mathfrak{S}_{L', R})^{\Delta=1} = (W(k) \otimes_{\mathbb{Z}_p} R)[[v]]$. Note that $E(v) = E((u')^{e'})$ is the minimal polynomial for $\pi_{K'}$ over K_0 .

We will make use of the following group schemes over \mathcal{O} . For $j \in \mathcal{J}$ and for any \mathcal{O} -algebra R , define

$$\begin{aligned} \mathcal{L}\mathcal{G}^{(j)}(R) &\stackrel{\mathrm{def}}{=} \{A \in \mathrm{GL}_n(R[v]^{\wedge E_j}[\frac{1}{E_j}])\}; \\ L^+\mathcal{G}^{(j)}(R) &\stackrel{\mathrm{def}}{=} \{A \in \mathrm{GL}_n(R[v]^{\wedge E_j}), \text{ is upper triangular modulo } v\} \end{aligned}$$

where $E_j = \sigma_j(E(v)) \in \mathcal{O}[v]$, and \wedge_{E_j} stands for the E_j -adic completion. In particular if R is p -adically complete, this is the same as the v -adic completion of $R[v]$.

3.1.1. Breuil–Kisin modules with tame descent. Let R be a p -adically complete Noetherian \mathcal{O} -algebra. For any positive integer h , let $Y^{[0,h],\tau}(R)$ denote the groupoid of Breuil–Kisin module of rank n over $\mathfrak{S}_{L',R}$, height in $[0, h]$ and descent data of type τ (cf. [CL18, §3], [LLHLMa, Definitions 5.1.1 and 5.1.3]):

Definition 3.1.1. An object of $Y^{[0,h],\tau}(R)$ consists of

- a finitely generated projective $\mathfrak{S}_{L',R}$ -module \mathfrak{M} which is locally free of rank n ;
- an injective $\mathfrak{S}_{L',R}$ -linear map $\phi_{\mathfrak{M}} : \varphi^*(\mathfrak{M}) \rightarrow \mathfrak{M}$ whose cokernel is killed by $E(v)^h$;
- a semilinear action of Δ on \mathfrak{M} which commutes with $\phi_{\mathfrak{M}}$, and such that Zariski locally on R , for each $j' \in \mathcal{J}'$,

$$\mathfrak{M}^{(j')} \pmod{u'} \cong \tau^\vee \otimes_{\mathcal{O}} R$$

as Δ' -representations.

Morphisms are $\mathfrak{S}_{L',R}$ -linear maps respecting all the above structures.

We will often omit the additional data and just write $\mathfrak{M} \in Y^{[0,h],\tau}(R)$ in what follows. It is known that $Y^{[0,h],\tau}$ is a p -adic formal algebraic stack over $\mathrm{Spf} \mathcal{O}$ (see, for example, [CL18, Theorem 4.7]).

Recall that an eigenbasis of $\mathfrak{M} \in Y^{[0,h],\tau}(R)$ is a collection of bases $\beta^{(j')}$ for each $\mathfrak{M}^{(j')}$ for $j' \in \mathcal{J}'$ compatible with the descent datum (see [LLHLMa, Definition 5.1.6] for details). Given the lowest alcove presentation (s, μ) of τ , and element $\mathfrak{M} \in Y^{[0,h],\tau}(R)$ and an eigenbasis β of \mathfrak{M} , equation (5.4) in [LLHLMa] defines the matrix $A_{\mathfrak{M},\beta}^{(j')} \in \mathrm{Mat}_n(\mathfrak{S}_R)$ for each $j' \in \mathcal{J}'$. We refer the reader to *loc. cit.* for details rather than recall the excessive notation needed to make a precise definition. We will recall the properties we need as we go along.

First, the matrix $A_{\mathfrak{M},\beta}^{(j')}$ only depends on $j' \pmod{f}$. Abusing notation, we occasionally write $A_{\mathfrak{M},\beta}^{(j)}$ for $j \in \mathcal{J}$ with the obvious meaning. Because τ is 1-generic, the height condition is equivalent to $A_{\mathfrak{M},\beta}^{(j)}$ and $(E_j)^h (A_{\mathfrak{M},\beta}^{(j)})^{-1}$ both lying in $\mathrm{Mat}_n(R[[v]])$ and being upper triangular modulo v , for all $j \in \mathcal{J}$.

Definition 3.1.2. (1) For integers $a \leq b$, define

$$L^{[a,b]}\mathcal{G}^{(j)}(R) := \{A \in L\mathcal{G}^{(j)}(R) \mid E_j^{-a}A, E_j^bA^{-1} \in \mathrm{Mat}_n(R[[v]]) \text{ and upper triangular mod } v\}.$$

(2) Given a pair $(s, \mu) \in W^{\mathcal{J}} \times X^*(T)^{\mathcal{J}}$, we define the (s, μ) -twisted φ -conjugation action of $\prod_{j \in \mathcal{J}} L^+\mathcal{G}^{(j)}(R)$ on $\prod_{j \in \mathcal{J}} L^{[a,b]}\mathcal{G}^{(j)}(R)$ by

$$(3.1) \quad (I^{(j)}) \cdot (A^{(j)}) = I^{(j)} A^{(j)} (\mathrm{Ad}(s_j^{-1} v^{\mu_j + \eta_{0,j}})(\varphi(I^{(j-1)})^{-1})).$$

Remark 3.1.3. (1) The change of basis formula in [LLHLMa, Proposition 5.1.8] can be summarized as follows. For the fixed lowest alcove presentation $(s, \mu) \in W^{\mathcal{J}} \times X^*(T)^{\mathcal{J}}$ of τ , the set of eigenbases of \mathfrak{M} is a torsor for $\prod_{j \in \mathcal{J}} L^+\mathcal{G}^{(j)}(R)$, and given two eigenbases β and β' differing by $(I^{(j)})_{j \in \mathcal{J}} \in \prod_{j \in \mathcal{J}} L^+\mathcal{G}^{(j)}(R)$, the collections $(A_{\mathfrak{M},\beta}^{(j)})$ and $(A_{\mathfrak{M},\beta'}^{(j)})$ differ by (s, μ) -twisted φ -conjugation by $(I^{(j)})_{j \in \mathcal{J}}$.

(2) Since eigenbases exist locally, we have the presentation

$$Y_{\mathbb{F}}^{[0,h],\tau} \cong \left[\prod_{j \in \mathcal{J}} L^{[0,h]}\mathcal{G}^{(j)} /_{(s,\mu),\varphi} \prod_{j \in \mathcal{J}} L^+\mathcal{G}^{(j)} \right]$$

where the quotient is with respect to the twisted φ -conjugation (3.1).

- (3) Let $\tilde{w}^*(\tau) = s^{-1}t_{\mu+\eta}$. A key observation which we use frequently is that (s, μ) -twisted conjugation $\prod_{j \in \mathcal{J}} L^{[a,b]} \mathcal{G}^{(j)}$ is the same as usual φ -conjugation on the right translation $\prod_{j \in \mathcal{J}} L^{[a,b]} \mathcal{G}^{(j)} \tilde{w}^*(\tau)$.

We now recall some useful results mod p . We write $\mathcal{I} \stackrel{\text{def}}{=} L^+ \mathcal{G}_{\mathbb{F}}^{(j)}$, which is the usual Iwahori group scheme over Noetherian \mathbb{F} -algebras, in particular it is independent of the choice $j \in \mathcal{J}$. We also write $\mathcal{I}_1 \subset \mathcal{I}$ for pro- v Iwahori consisting of upper unipotent matrices mod p . Note that $E_j \equiv v^e \pmod{p}$.

Lemma 3.1.4 (Lemma 5.2.2 [LLHLMa]). *Let R be an \mathbb{F} -algebra and $(A_1^{(j)})_{j \in \mathcal{J}}, (A_2^{(j)})_{j \in \mathcal{J}} \in L^{[a,b]} \mathcal{G}^{(j)}(R)$. Let $\tilde{z} = s^{-1}t_{\mu+\eta} \in \widetilde{W}^{\vee, \mathcal{J}}$ where μ is $(e(b-a)+1)$ -deep in \underline{C}_0 and $s \in W^{\mathcal{J}}$. Then, there is a bijection between the following:*

- (1) *Tuples $(I^{(j)})_{j \in \mathcal{J}} \in \mathcal{I}_1(R)^{\mathcal{J}}$ such $A_2^{(j)} \tilde{z}_j = I^{(j)} A_1^{(j)} \tilde{z}_j \varphi(I^{(j-1)})^{-1}$ for all $j \in \mathcal{J}$;*
- (2) *Tuples $(X_j)_{j \in \mathcal{J}} \in \mathcal{I}_1(R)^{\mathcal{J}}$ such that $A_2^{(j)} = X_j A_1^{(j)}$ for all $j \in \mathcal{J}$.*

Remark 3.1.5. As in [LLHLMa, Corollary 5.2.3], Lemma 3.1.4 gives a presentation of $Y_{\mathbb{F}}^{[a,b], \tau}$ as quotient of $\prod_{j \in \mathcal{J}} \mathcal{I}_1 \backslash L^{[a,b]} \mathcal{G}_{\mathbb{F}}^{(j)}$ by (s, μ) -twisted conjugation by the torus $T_{\mathbb{F}}^{\vee, \mathcal{J}}$ when μ is $(e(a-b)+1)$ -deep.

Definition 3.1.6. Let \mathbb{F}'/\mathbb{F} be finite extension. The *shape* of a Breuil–Kisin module $\mathfrak{M} \in Y^{[0,h], \tau}(\mathbb{F}')$ with respect to τ is the element $\tilde{z} = (\tilde{z}_j)_{j \in \mathcal{J}} \in \widetilde{W}^{\vee, \mathcal{J}}$ such that for any eigenbasis β and any $j \in \mathcal{J}$, the matrix $A_{\mathfrak{M}, \beta}^{(j)}$ lies in $\mathcal{I}(\mathbb{F}') \tilde{z}_j \mathcal{I}(\mathbb{F}')$.

Proposition 3.1.7. *For each $\tilde{z} \in \widetilde{W}^{\vee, \mathcal{J}}$ such that $\tilde{z}_j \in L^{[0,h]} \mathcal{G}^{(j)}(\mathbb{F})$ for $j \in \mathcal{J}$, there is a locally closed substack $Y_{\mathbb{F}, \tilde{z}}^{[0,h], \tau} \stackrel{\text{def}}{=} \left[\prod_{j \in \mathcal{J}} \mathcal{I} \tilde{z}_j \mathcal{I} / (s, \mu), \varphi \prod_{j \in \mathcal{J}} \mathcal{I} \right] \subset Y_{\mathbb{F}}^{[0,h], \tau}$ whose \mathbb{F}' -points are the Breuil–Kisin modules of shape \tilde{z} . The closure of $Y_{\mathbb{F}, \tilde{z}}^{[0,h], \tau}$ is contained in the union of the strata $Y_{\mathbb{F}, \tilde{z}'}^{[0,h], \tau}$ such that $\tilde{z}' \leq \tilde{z}$ in the Bruhat order.*

Proof. This follows from

$$\begin{array}{ccc}
 Y_{\mathbb{F}}^{[0,h], \tau} & & \prod_{j \in \mathcal{J}} L^{[0,h]} \mathcal{G}_{\mathbb{F}}^{(j)} / \mathcal{I} \\
 & \searrow & \swarrow \\
 & \prod_{j \in \mathcal{J}} [\mathcal{I} \backslash L^{[0,h]} \mathcal{G}_{\mathbb{F}}^{(j)} / \mathcal{I}] &
 \end{array}$$

where the right arrow is an $\mathcal{I}^{\mathcal{J}}$ -torsor (cf. [CL18, Proposition 5.4]). \square

We define the cocharacter $\eta \stackrel{\text{def}}{=} ((n-1, \dots, 1, 0), \dots, (n-1, \dots, 1, 0)) \in X_*(T^{\vee})^{\text{Hom}(K, E)}$. There is closed p -adic formal algebraic stack $Y^{\leq \eta, \tau} \subset Y^{[0, n-1], \tau}$ defined in [CL18, Theorem 5.3] [LLHLMa, §5.3]. We recall the following result, deduced either from [CL18, Theorems 2.15 and 5.3] by reducing to principal series case or, when τ is n -generic and K/\mathbb{Q}_p is unramified, from [LLHLMa, Corollary 5.2.3 and Proposition 5.4.7].

Proposition 3.1.8. *The special fiber of $Y_{\mathbb{F}}^{\leq \eta, \tau}$ satisfies*

$$Y_{\mathbb{F}, \text{red}}^{\leq \eta, \tau} \subset \bigcup_{\tilde{z} \in \text{Adm}^{\vee}(e\eta_0)} Y_{\mathbb{F}, \tilde{z}}^{\leq \eta, \tau}.$$

Remark 3.1.9. In fact, the special fiber of $Y^{\leq \eta, \tau}$ is reduced and the inclusion in Proposition 3.1.8 is an equality. This is shown in the principal series case in [CL18]. The general case is best handled via the ramified generalization of [LLHLMa] which we will return to in the future. (See also Remark 4.2.3 and the discussion preceding it.)

3.1.2. Étale φ -modules. Let $\mathcal{O}_{\mathcal{E}, K}$ (resp. $\mathcal{O}_{\mathcal{E}, L'}$) be the p -adic completion of $(W(k)[[v]])[1/v]$ (resp. of $(W(k')[[u']])[1/u']$). It is endowed with a continuous Frobenius morphism φ extending the Frobenius on $W(k)$ (resp. on $W(k')$) and such that $\varphi(v) = v^p$ (resp. $\varphi(u') = (u')^p$). Let R be a p -adically complete Noetherian \mathcal{O} -algebra. We then have the groupoid $\Phi\text{-Mod}_K^{\text{ét}, n}(R)$ (resp. $\Phi\text{-Mod}_{dd, L'}^{\text{ét}, n}(R)$) of étale $(\varphi, \mathcal{O}_{\mathcal{E}, K} \widehat{\otimes}_{\mathbb{Z}_p} R)$ -modules (resp. étale $(\varphi, \mathcal{O}_{\mathcal{E}, L'} \widehat{\otimes}_{\mathbb{Z}_p} R)$ -modules with descent data from L' to K). Its objects are rank n projective modules \mathcal{M} over $\mathcal{O}_{\mathcal{E}, K} \widehat{\otimes}_{\mathbb{Z}_p} R$ (resp. $\mathcal{O}_{\mathcal{E}, L'} \widehat{\otimes}_{\mathbb{Z}_p} R$), endowed with a Frobenius semilinear endomorphism $\phi_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M}$ (resp. a Frobenius semilinear endomorphism $\phi_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M}$ commuting with the descent data) inducing an isomorphism on the pull-back: $\text{id} \otimes_{\varphi} \phi_{\mathcal{M}} : \varphi^*(\mathcal{M}) \xrightarrow{\sim} \mathcal{M}$. It is known that $\Phi\text{-Mod}_K^{\text{ét}, n}(R)$ and $\Phi\text{-Mod}_{dd, L'}^{\text{ét}, n}(R)$ form fppf stacks over $\text{Spf } \mathcal{O}$ (see [EGa, §3.1], [EGb, §5.2], [CEGS, §3.1] where they are denoted $\mathcal{R}_n, \mathcal{R}_{n, L'}^{dd}$ respectively). We use $\Phi\text{-Mod}_K^{\text{ét}}(R)$ (resp. $\Phi\text{-Mod}_{dd, L'}^{\text{ét}}(R)$) to denote the category of étale $(\varphi, \mathcal{O}_{\mathcal{E}, K} \widehat{\otimes}_{\mathbb{Z}_p} R)$ -modules (resp. $\mathcal{O}_{\mathcal{E}, L'} \widehat{\otimes}_{\mathbb{Z}_p} R$ -modules with descent from L' to K) of arbitrary finite rank.

Given $\mathfrak{M} \in Y^{[0, h], \tau}(R)$, $\mathfrak{M} \otimes_{\mathcal{S}_{L', R}} (\mathcal{O}_{\mathcal{E}, L'} \widehat{\otimes}_{\mathbb{Z}_p} R)$ is naturally an object $\Phi\text{-Mod}_{dd, L'}^{\text{ét}, n}(R)$, and we define an étale φ -module $\mathcal{M} \in \Phi\text{-Mod}_K^{\text{ét}, n}(R)$ by

$$\mathcal{M} \stackrel{\text{def}}{=} (\mathfrak{M} \otimes_{\mathcal{S}_{L', R}} (\mathcal{O}_{\mathcal{E}, L'} \widehat{\otimes}_{\mathbb{Z}_p} R))^{\Delta=1}$$

with the induced Frobenius. This construction defines a morphism of stacks $\varepsilon_{\tau} : Y^{[0, h], \tau} \rightarrow \Phi\text{-Mod}_K^{\text{ét}, n}$ which is representable by algebraic spaces, proper, and of finite presentation (see [LLHLMa, Proposition 5.4.1], which carries through in our ramified setting). Note that ε_{τ} is independent of any presentation of τ .

For any $(\mathcal{M}, \phi_{\mathcal{M}}) \in \Phi\text{-Mod}_K^{\text{ét}}(R)$, we decompose $\mathcal{M} = \bigoplus_{j \in \mathcal{J}} \mathcal{M}^{(j)}$ over the embeddings $\sigma_j : W(k) \rightarrow \mathcal{O}$, with induced maps $\phi_{\mathcal{M}}^{(j)} : \mathcal{M}^{(j-1)} \rightarrow \mathcal{M}^{(j)}$. We can define the map ε_{τ} explicitly in some cases:

Proposition 3.1.10. ([LLHLMa, Proposition 5.4.2]) *Let $\mathfrak{M} \in Y^{[0, h], \tau}(R)$ and set $\mathcal{M} = \varepsilon_{\tau}(\mathfrak{M})$. Let (s, μ) be the fixed lowest alcove presentation of τ . If β is an eigenbasis of \mathfrak{M} , then there exists a basis \mathfrak{f} (determined by β) for \mathcal{M} such that the matrix of $\phi_{\mathcal{M}}^{(j)}$ with respect to \mathfrak{f} is given by*

$$A_{\mathfrak{M}, \beta}^{(j)} s_j^{-1} v^{\mu_j + \eta_{0, j}} = A_{\mathfrak{M}, \beta}^{(j)} \tilde{w}^*(\tau)_j.$$

Finally, we recall that in generic situations the map ε_{τ} does not lose information:

Proposition 3.1.11. ([LLHLMa, Proposition 5.4.3]) *Assume τ is $(eh + 1)$ -generic. Then the map $\varepsilon_{\tau} : Y^{[0, h], \tau} \rightarrow \Phi\text{-Mod}_K^{\text{ét}, n}$ is a closed immersion.*

We briefly recall the relations between Breuil–Kisin modules and Galois representations. Recall from 1.5.2 the extension K_{∞}/K , and let $G_{K_{\infty}} \subset G_K$ denote the absolute Galois group of K_{∞} . We

have an anti-equivalence of categories established by the exact functor

$$\mathbb{V}_K^* : \Phi\text{-Mod}_K^{\acute{e}t, n}(R) \rightarrow \text{Rep}_R^n(G_{K_\infty})$$

defined through the theory of fields of norms (cf. [LLHLM18, §2.3 and §6.1] for details) and therefore a functor $T_{dd}^* : Y^{[0, h], \tau}(R) \rightarrow \text{Rep}_R^n(G_{K_\infty})$ defined as the composite of ε_τ followed by \mathbb{V}_K^* .

We finally recall from [LLHLMa, §5.5] the notion of *shape* of an n -dimensional \mathbb{F} -representation of G_K (or G_{K_∞}) with respect to τ .

Definition 3.1.12. Assume that τ is $(e(n-1)+1)$ -generic. Let $\bar{\rho}$ be an n -dimensional \mathbb{F} -representation of G_K or G_{K_∞} . If there exists $\overline{\mathfrak{M}} \in Y_{\mathbb{F}}^{\leq n, \tau}(\mathbb{F})$ such that $T_{dd}^*(\overline{\mathfrak{M}}) \cong \bar{\rho}|_{G_{K_\infty}}$ then we say that $\bar{\rho}$ is τ -admissible, and we define $\tilde{w}(\bar{\rho}, \tau) \in \text{Adm}(e\eta_0)$ to be the shape of $\overline{\mathfrak{M}}$ with respect to τ (Definition 3.1.6). This is well-defined by Proposition 3.1.11.

Proposition 3.1.13. ([LLHLMa, Proposition 5.5.7]) *Assume that the fixed lowest alcove presentation (s, μ) of τ is $(e(n-1)+1)$ -generic. Let $\bar{\rho}$ be a semisimple representation of G_K over \mathbb{F} . Then $\bar{\rho}$ is τ -admissible if and only if $\bar{\rho}|_{I_K}$ admits a lowest alcove presentation (w, ν) compatible with the lowest alcove presentation of τ such that $s^{-1}t_{\nu-\mu}w \in \text{Adm}(e\eta_0)$. Furthermore, if $\bar{\rho}$ is τ -admissible then $\tilde{w}(\bar{\rho}, \tau) = s^{-1}t_{\nu-\mu}w$.*

Proposition 3.1.14. *Assume that the fixed lowest alcove presentation (s, μ) of τ is $(eh+1)$ -generic. Let $\mathfrak{M} \in Y^{[0, h], \tau}(\mathbb{F})$ with shape $\tilde{z} \in \widetilde{W}^{\mathcal{J}}$. Then, there exists an eigenbasis β for \mathfrak{M} , unique up to scaling by $T(\mathbb{F})^{\mathcal{J}}$, such that*

$$A_{\mathfrak{M}, \beta}^{(j)} \in T(\mathbb{F})\tilde{z}_j N_{\tilde{z}_j}(\mathbb{F})$$

where $N_{\tilde{z}_j}$ is unipotent subgroup scheme of \mathcal{I} defined in [LLHLMa, Definition 4.2.9].

Proof. This follows from [LLHLMa, Proposition 4.2.13 and Corollary 5.2.3]. \square

3.2. Mod p monodromy. Let $\text{Fl} := \mathcal{I} \backslash LG_{\mathbb{F}}$ denote the affine flag variety over \mathbb{F} where $LG_{\mathbb{F}} = LG_{\mathbb{F}}^{(j)}$ for any j denotes the usual loop group. Given $\tilde{w} \in \widetilde{W}^{\vee}$, we write $S_{\mathbb{F}}^{\circ}(\tilde{w})$ for the affine open Schubert cell associated to \tilde{w} . Let $\mathbf{a} \in (\mathcal{O}^n)^{\mathcal{J}}$. Define a closed subfunctor of LG by

$$(3.2) \quad LG^{\nabla \mathbf{a}}(R) \stackrel{\text{def}}{=} \left\{ (A^{(j)}) \in LG(R) \mid v \frac{dA^{(j)}}{dv} (A^{(j)})^{-1} + A^{(j)} \text{Diag}(\mathbf{a}_j) (A^{(j)})^{-1} \in \frac{1}{v^e} \text{Lie } \mathcal{I}(R) \text{ for all } j \in \mathcal{J} \right\}.$$

This condition defines a closed sub-ind-scheme $\text{Fl}_{\mathcal{J}}^{\nabla \mathbf{a}} \subset \text{Fl}_{\mathcal{J}}^{\vee}$. For any subset $S \subset LG(R)$, we set $S^{\nabla \mathbf{a}} := S \cap LG^{\nabla \mathbf{a}}(R)$; similarly for any subscheme $X \subset \text{Fl}_{\mathcal{J}}^{\vee}$, set $X^{\nabla \mathbf{a}} \stackrel{\text{def}}{=} X \cap \text{Fl}_{\mathcal{J}}^{\nabla \mathbf{a}}$.

Following [LLHLMa, Definition 4.2.2], given an integer $m \geq 0$, we say that an element $(a_1, \dots, a_n) \in R^n$ is m -generic if $a_i - a_k \notin \{-m, -m+1, \dots, m-1, m\}$ for all $i \neq k$ (where $-m, -m+1, \dots, m-1, m$ are considered as elements of R via the canonical embedding $\mathbb{F}_p \hookrightarrow \mathbb{F}$ and the structural morphism $\mathbb{F} \hookrightarrow R$.)

Proposition 3.2.1. *Let h be a positive integer. Let $\tilde{w} \in \widetilde{W}^{\mathcal{J}}$ and $\mathbf{a} = (\mathbf{a}_j)_{j \in \mathcal{J}} \in (\mathcal{O}^n)^{\mathcal{J}}$. Assume that \tilde{w} is e -regular and h -small (see Definitions 2.1.5 and 2.1.7(3)) and that $\mathbf{a}_j \bmod \varpi \in \mathbb{F}^n$ is h -generic for all $j \in \mathcal{J}$. Then the intersection $S_{\mathbb{F}}^{\circ}(\tilde{w}^*) \cap \text{Fl}_{\mathcal{J}}^{\nabla \mathbf{a}}$ is an affine space of dimension $[K : \mathbb{Q}_p] \dim(B \backslash \text{GL}_n)_{\mathbb{F}}$.*

Proof. This is a direct generalization of [LLHLMa, Theorem 4.2.4] to the ramified setting. We only briefly outline the proof. It suffices to consider the case when $\#\mathcal{J} = 1$. By [LLHLMa, Proposition 4.2.13], there is an isomorphism $\tilde{w}^* N_{\tilde{w}^*} \cong S^{\circ}(\tilde{w}^*)$ where $N_{\tilde{w}^*}$ is a unipotent subgroup scheme of \mathcal{I}

isomorphic to an affine space of dimension $\ell(\tilde{w})$. As \tilde{w} is e -regular, for each α in the support of $N_{\tilde{w}^*}$, we have $N_{\tilde{w}^*, \alpha} = v^{\delta_{\alpha < 0}} f_\alpha$ where f_α is a polynomial of degree at least $e - 1$ (cf. [LLHLMa, Corollary 4.2.5], and note that, more precisely, the degree is $[\langle \tilde{w}(x), -\alpha^\vee \rangle] - \delta_{\alpha < 0}$, which is at least $e - 1$ by the e -regularity condition). Condition (3.2) does not impose any constraint on the coefficients of degree $\deg(f_\alpha), \deg(f_\alpha) - 1, \dots, \deg(f_\alpha) - (e - 1)$ of f_α , while the coefficients of degree strictly smaller than $\deg(f_\alpha) - (e - 1)$ are solved in terms of the coefficients of the polynomials $f_{\alpha'}$ with $\alpha' <_{\mathcal{C}} \alpha$ for a partial order $<_{\mathcal{C}}$ on Φ determined by \tilde{w} (cf. equation (4.6) in *loc. cit.*). Hence, $(\tilde{w}^* N_{\tilde{w}^*})^{\nabla_{\mathbf{a}}}$ is an affine space of dimension $e \dim(B \backslash \mathrm{GL}_n)_{\mathbb{F}}$. \square

Let $\tilde{z} = s^{-1} t_\mu \in \widetilde{W}^{\vee, \mathcal{J}}$ acting by right translation on $\mathrm{Fl}^{\mathcal{J}}$. Let $\mathbf{a} \in (\mathbb{Z}^n)^{\mathcal{J}}$ and assume that $\mathbf{a}_j \equiv s_j^{-1}(\mu_j) \pmod{p}$ for all $j \in \mathcal{J}$. An easy calculation shows that:

$$LG\tilde{z} \cap LG^{\nabla_0} = LG^{\nabla_{\mathbf{a}}}\tilde{z}, \quad \mathrm{Fl}^{\mathcal{J}}\tilde{z} \cap \mathrm{Fl}_{\mathcal{J}}^{\nabla_0} = \mathrm{Fl}_{\mathcal{J}}^{\nabla_{\mathbf{a}}}\tilde{z}$$

We can now state the main result of the section which is the ramified analogue of [LLHLMa, Proposition 4.3.4]:

Proposition 3.2.2. *Let $\tilde{w}, \tilde{w}' \in \widetilde{W}^{\mathcal{J}}$ be h -small, e -regular elements such that $\tilde{w}' \leq \tilde{w}$. Write $\tilde{w}^* = (\tilde{w}')^* \tilde{z}'$ and assume this is a reduced expression for \tilde{w}^* . Let $\tilde{z} \in \widetilde{W}^{\vee, \mathcal{J}}$ be $2h$ -generic. Then*

$$(\mathcal{I}(\mathbb{F})(\tilde{w}'_j)^* \mathcal{I}(\mathbb{F})\tilde{z}'_j \tilde{z}_j)^{\nabla_0} = (\mathcal{I}(\mathbb{F})\tilde{w}_j^* \mathcal{I}(\mathbb{F})\tilde{z}_j)^{\nabla_0}$$

for all $j \in \mathcal{J}$.

Proof. Again, the proof is very similar to the proof of [LLHLMa, Proposition 4.3.4], and we refer the reader to *loc. cit.* for further detail.

Since $(\tilde{w}')^* \tilde{z} = \tilde{w}^*$ is a reduced expression, there is an inclusion of the left side in the right side. Since both sides are invariant under $\mathcal{I}(\mathbb{F})$, we can descend to $\mathrm{Fl}_{\mathcal{J}}^{\nabla_0}$ and reduce to showing

$$(S^\circ(\tilde{w}')^* \tilde{z}' \tilde{z})^{\nabla_0} = (S^\circ(\tilde{w}^*) \tilde{z})^{\nabla_0}.$$

By the assumptions, both \tilde{z} and $\tilde{z}' \tilde{z}$ are h -generic and so by Proposition 3.2.1, both sides are affine spaces of the same dimension and so inclusion implies equality. \square

Definition 3.2.3. Assume that the lowest alcove presentation (s, μ) of τ is $(eh + 1)$ -generic. We say that $\mathfrak{M} \in Y^{[0, h], \tau}(\mathbb{F})$ satisfies the *mod p monodromy condition* if for any choice of eigenbasis β of \mathfrak{M} , the collection $(A_{\mathfrak{M}, \beta}^{(j)} \tilde{w}^*(\tau)_j)$ is in $LG^{\nabla_0}(\mathbb{F})^{\mathcal{J}}$.

3.3. Semicontinuity I. We fix a tame inertial type τ with a 1-generic lowest alcove presentation (s, μ) , as defined in §2.3. In this section, we show a semicontinuity result for the shape of a mod p Kisin module of type τ with respect to the shape of its semisimplification. This is preliminary to a more general semicontinuity result (Theorem 3.5.1 in section 3.5).

Proposition 3.3.1. *Let $\mathfrak{M} \in Y_{\mathbb{F}}^{[0, h], \tau}(\mathbb{F}')$ and set $\bar{\rho} := T_{dd}^*(\mathfrak{M})$ for any finite extension \mathbb{F}'/\mathbb{F} . There exists $\mathfrak{M}_0 \in Y_{\mathbb{F}}^{[0, h], \tau}(\mathbb{F}')$ such that*

$$T_{dd}^*(\mathfrak{M}_0) = \bar{\rho}^{\mathrm{ss}}.$$

Furthermore, the shape of \mathfrak{M}_0 with respect to τ is less than or equal to the shape of \mathfrak{M} with respect to τ in the Bruhat order on $\widetilde{W}^{\vee, \mathcal{J}}$.

Proof. By the closure relations for the stratum of the stack $Y_{\mathbb{F}}^{[0, h], \tau}$ (Proposition 3.1.7), it suffices to construct a map $\mathbb{A}_{\mathbb{F}}^1 \rightarrow Y_{\mathbb{F}}^{[0, h], \tau}$, sending x to \mathfrak{M}_x , such that

$$(1) \quad T_{dd}^*(\mathfrak{M}_0) \cong \bar{\rho}^{\mathrm{ss}}; \text{ and}$$

(2) for all $x \in \overline{\mathbb{F}}, x \neq 0, T_{dd}^*(\mathfrak{M}_x) \cong \overline{\rho}$.

The construction of the map proceeds as in the proof of [LLHLMa, Proposition 5.5.9]. Let α be the eigenbasis for \mathfrak{M} constructed in *loc. cit.* adapted to the filtration (\mathcal{M}_i) on the étale φ -module $\mathfrak{M}[1/u']$. Define the matrix $C^{(j)} \in G(\mathbb{F}'((u')))$ by the condition

$$\phi_{\mathfrak{M}}^{(j)}(\varphi^*(\alpha^{(j)})) = \alpha^{(j+1)} C^{(j)}.$$

By construction, $C^{(j)}$ lies in a parabolic subgroup $P(\mathbb{F}'((u'))) \subset G(\mathbb{F}'((u')))$ corresponding to the filtration (\mathcal{M}_i) . Let L denote the corresponding Levi subgroup which contains the diagonal torus T . Choose a dominant cocharacter λ such that L is the centralizer of λ .

For $x \neq 0$, define \mathfrak{M}_x to be the free Breuil–Kisin module of rank n with basis α_x such that Δ acts on α_x in the same way it acts on α and such that the Frobenius acts by $C_x^{(j)} = \lambda(x)C^{(j)}\lambda(x)^{-1}$ (with respect to α_x). Observe that the limit of $C_x^{(j)}$ as $x \rightarrow 0$ exists and lies in the Levi subgroup $L(\mathbb{F}'((u')))$. Thus, we can extend this to a family over $\mathbb{A}_{\mathbb{F}}^1$. It is easy to check property (1). For property (2), we note that for any $x \neq 0, C_x^{(j)}$ is the matrix for Frobenius with respect to the basis $(\alpha^{(j)} \cdot \lambda(x))$ and so \mathfrak{M}_x is isomorphic to \mathfrak{M}_1 . \square

Corollary 3.3.2. *Assume that τ is $(e(n-1)+1)$ -generic. If $\overline{\rho}$ is τ -admissible, then $\overline{\rho}^{\text{ss}}$ is τ -admissible and for all $j \in \mathcal{J}$,*

$$\tilde{w}(\overline{\rho}^{\text{ss}}, \tau)_j \leq \tilde{w}(\overline{\rho}, \tau)_j.$$

3.4. Specializations. Throughout this section we consider a continuous Galois representation $\overline{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$. We say that $\overline{\rho}$ is N -generic if the tame inertial \mathbb{F} -type $\overline{\rho}^{\text{ss}}|_{I_K}$ is N -generic (see §2.3.3). *All lowest alcove presentations for tame inertial types (over \mathbb{F} or over E) will always be compatible with a given lowest alcove presentation $\overline{\rho}^{\text{ss}}$.*

If $\overline{\rho}^{\text{sp}}$ is a tame inertial \mathbb{F} -type for K and τ is an inertial type over E with compatible lowest alcove presentation, then recall the combinatorially defined shape $\tilde{w}(\overline{\rho}^{\text{sp}}, \tau) = \tilde{w}(\tau)^{-1}\tilde{w}(\overline{\rho}^{\text{sp}}) \in \widetilde{W}^{\mathcal{J}}$ defined in §2.4.

Definition 3.4.1. Let $\overline{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be a continuous Galois representation.

- (1) A tame inertial \mathbb{F} -type $\overline{\rho}^{\text{sp}}$ for K over \mathbb{F} is a *specialization* of $\overline{\rho}$ if there exists an $(e(n-1)+1)$ -generic tame inertial type τ such that $\overline{\rho}$ is τ -admissible and $\tilde{w}(\overline{\rho}, \tau) = \tilde{w}(\overline{\rho}^{\text{sp}}, \tau)$. We say that τ *exhibits* the specialization.
- (2) A specialization $\overline{\rho}^{\text{sp}}$ is called an *extremal specialization* of $\overline{\rho}$ if there exists a τ exhibiting the specialization such that $\tilde{w}(\overline{\rho}^{\text{sp}}, \tau) = t_{w^{-1}(e\eta_0)}$ for some $w \in W^{\mathcal{J}}$ and if the unique $\mathfrak{M} \in Y^{\leq \eta, \tau}(\mathbb{F})$ such that $T_{dd}^*(\mathfrak{M}) \cong \overline{\rho}|_{G_{K_\infty}}$ satisfies the mod p monodromy condition (Definition 3.2.3).

Remark 3.4.2. By a version of [LLHLMa, Proposition 7.4.1] in the ramified setting (based on the analysis of the monodromy condition in characteristic 0, cf. Proposition 4.4.1 below), if τ is $(e+1)(n-1)$ -generic and $\overline{\rho}$ admits a potentially crystalline lift of type (τ, η) then the unique $\mathfrak{M} \in Y^{\leq \eta, \tau}(\mathbb{F})$ such that $T_{dd}^*(\mathfrak{M}) \cong \overline{\rho}|_{G_{K_\infty}}$ satisfies the mod p monodromy condition. Thus, the technical condition in Definition 3.4.1(2) could be replaced by the existence of a potentially crystalline lift.

Remark 3.4.3. Using the methods of [LLHLMa], it can be shown under suitable genericity conditions that all specializations are extremal when K/\mathbb{Q}_p is unramified. It is likely that the same is true in the ramified case but we do not attempt to prove it here.

Let $S(\overline{\rho})$ (resp. $S_{\text{ext}}(\overline{\rho})$) denote be the set of specializations (resp. extremal specializations) of $\overline{\rho}$.

Remark 3.4.4. The sets $S(\bar{\rho})$ and $S_{\text{ext}}(\bar{\rho})$ are finite because the set of $(e(n-1)+1)$ -generic types τ for which $\bar{\rho}^{\text{ss}}$ is τ -admissible is finite by Proposition 3.1.13 and the set of types for which $\bar{\rho}$ is τ -admissible is a subset of this set by Corollary 3.3.2. In Theorem 3.8.2, we show that $\bar{\rho}^{\text{ss}}|_{I_K}$ is an extremal specialization of $\bar{\rho}$ so $S_{\text{ext}}(\bar{\rho})$ is also non-empty.

Example 3.4.5. We have the following examples when $K = \mathbb{Q}_p$ and $n = 2, 3$.

- (1) Let $1 < a < p-1$ and assume that $\bar{\rho}|_{I_{\mathbb{Q}_p}}$ is of the form $\begin{pmatrix} \omega^a & * \\ 0 & 1 \end{pmatrix}$, where $* \neq 0$. Then we have two specializations, given by $\omega^a \oplus 1$ and $\omega_2^a \oplus \omega_2^{pa}$. A type which exhibits the specialization $\omega_2^a \oplus \omega_2^{pa}$ is $(\omega_2^{a+1} \oplus \omega_2^{p(a+1)}) \otimes \omega^{-1}$.
- (2) ([LLHLMb, Theorem 4.2.5]) Assume that $(a+1, b+1, c+1) \in \mathbb{Z}^3$ is 6-deep in alcove C_0 and that $\bar{\rho}|_{I_K}$ is of the form

$$\begin{pmatrix} \omega^a & *_{1} & * \\ 0 & \omega^b & *_{2} \\ 0 & 0 & \omega^c \end{pmatrix}$$

where $*_{1}, *_{2}$ denote non-split extensions. Then $\bar{\rho}$ has up to 6 specializations, namely $\omega^a \oplus \omega^b \oplus \omega^c$, $\omega_2^{a+pb} \oplus \omega_2^{b+pa} \oplus \omega^c$, $\omega_2^{a+pc} \oplus \omega^b \oplus \omega_2^{c+pa}$, $\omega^a \oplus \omega_2^{b+pc} \oplus \omega_2^{c+pb}$, $\omega_3^{a+pb+p^2c} \oplus \omega_3^{b+pc+p^2a} \oplus \omega_3^{c+pa+p^2b}$ and $\omega_3^{a+pc+p^2b} \oplus \omega_3^{b+pa+p^2c} \oplus \omega_3^{c+pb+p^2a}$. A type which exhibits the specialization $\bar{\rho}^{\text{ss}}$ is $\omega_2^{(a-1)+p(c-1)} \oplus \omega^{b-1} \oplus \omega_2^{(c-1)+p(a-1)}$. We have 4 specializations precisely when $\bar{\rho}$ has either a potentially crystalline lift of type $\omega_{a-1} \oplus \omega_2^{b+p(c-2)} \oplus \omega_2^{(c-2)+pb}$ (in which case the specializations $\omega_2^{a+pc} \oplus \omega^b \oplus \omega_2^{c+pa}$ and $\omega_3^{a+pc+p^2b} \oplus \omega_3^{b+pa+p^2c} \oplus \omega_3^{c+pb+p^2a}$ do not appear) or of type $\omega_2^{a+p(b-2)} \oplus \omega_2^{(b-2)+pa} \oplus \omega^{c-1}$ (in which case the specializations $\omega_2^{a+pc} \oplus \omega^b \oplus \omega_2^{c+pa}$ and $\omega_3^{a+pb+p^2c} \oplus \omega_3^{b+pc+p^2a} \oplus \omega_3^{c+pa+p^2b}$ do not appear).

3.5. Semicontinuity II. The following theorem generalizes Proposition 3.3.1.

Theorem 3.5.1. *Let $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be a $3e(n-1)$ -generic continuous Galois representation. Assume that $\bar{\rho}$ specializes to a tame inertial \mathbb{F} -type $\bar{\rho}^{\text{sp}}$ for K and that $\bar{\rho}$ is τ -admissible. For each $j \in \mathcal{J}$, we have the inequality*

$$\tilde{w}(\bar{\rho}^{\text{sp}}, \tau)_j \leq \tilde{w}(\bar{\rho}, \tau)_j.$$

We begin by stating two combinatorial lemmas which will be needed in the proof of Theorem 3.5.1.

Lemma 3.5.2. *Let τ and τ' be $(e(n-1)+1)$ -generic tame inertial types over E . Assume there exists a $\bar{\rho}$ which is both τ and τ' -admissible. Then, for any choice of lowest alcove presentation of $\bar{\rho}^{\text{ss}}$, τ and τ' admit lowest alcove presentations (s, μ) and (s', μ') , compatible with that of $\bar{\rho}^{\text{ss}}$, such that*

$$|\mu_{j,i} - \mu'_{j,i}| \leq e(n-1).$$

Proof. Since $\bar{\rho}$ is both τ and τ' -admissible, the same is true for $\bar{\rho}^{\text{ss}}$ by Proposition 3.3.2. Fixing a lowest alcove presentation of $\bar{\rho}^{\text{ss}}$, τ and τ' admit compatible presentation (s, μ) and (s', μ') respectively and we have $e\eta_0$ -admissible elements $\tilde{w}(\bar{\rho}^{\text{ss}}, \tau) = t_{\nu}w$ and $\tilde{w}(\bar{\rho}^{\text{ss}}, \tau') = t_{\nu'}w'$. Since ν and ν' are in the convex hull of $\underline{W}e\eta_0$ (cf. [HC02, Theorem 3.3]),

$$(3.3) \quad 0 \leq \nu_{j,i}, \nu'_{j,i} \leq e(n-1).$$

By Proposition 3.1.13, $\bar{\rho}^{\text{ss}}|_{I_K}$ has lowest alcove presentation $(sw, \mu + s(\nu)) = (s'w', \mu' + s'(\nu'))$. The result now follows from this equation and (3.3). \square

Lemma 3.5.3. *Let μ, μ' be dominant cocharacters which are $2e(n-1)$ -deep in alcove \underline{C}_0 , and assume that for all $j \in \mathcal{J}$, $1 \leq i \leq n$,*

$$|\mu_{j,i} - \mu'_{j,i}| \leq e(n-1).$$

Let $(B_j), (B'_j) \in \text{Mat}_n(\mathbb{F}[[v]])^{\mathcal{J}}$ such that for all j , $v^{e(n-1)}B_j^{-1} \in \text{Mat}_n(\mathbb{F}[[v]])$. Assume that for all $j \in \mathcal{J}$ there exists $C_j \in \text{GL}_n(\mathbb{F}((v)))$ such that

$$(3.4) \quad C_j B'_j v^{\mu'_j + \eta_{0,j}} = B_j v^{\mu_j + \eta_{0,j}} \varphi(C_{j-1}).$$

Then $C_j \in \mathcal{I}(\mathbb{F})$ for all $j \in \mathcal{J}$.

Proof. The technique is similar to the proof of [LLHLM18, Theorem 3.2] and [LLHLMa, Lemma 5.4.5]. We first show that for all $j \in \mathcal{J}$ we have $C_j \in \text{Mat}_n(\mathbb{F}[[v]])$. For all $j \in \mathcal{J}$, write $C_j = v^{-k_j} C_j^+$ with $k_j \in \mathbb{Z}$, $C_j^+ \in \text{Mat}_n(\mathbb{F}[[v]])$ and $C_j^+ \not\equiv 0$ modulo v . Rearranging equation (3.4), we can write:

$$(3.5) \quad v^{-pk_{j-1}} \text{Ad}(v^{\mu_j + \eta_{0,j}}) \varphi(C_{j-1}^+) = v^{-k_j} B_j^{-1} C_j^+ B'_j v^{\mu'_j - \mu_j}.$$

Since the RHS of (3.5) becomes integral after multiplying by $v^{k_j + e(n-1) + \max_{1 \leq i \leq n} |\mu'_{j,i} - \mu_{j,i}|}$, we get that

$$k_j + p - 1 > k_j + e(n-1) + \max_{1 \leq i \leq n} |\mu'_{j,i} - \mu_{j,i}| + \max_{\alpha \in \Phi} |\langle \mu_j, \alpha^\vee \rangle| \geq pk_{j-1}.$$

This shows that if $k = \max_{1 \leq j \leq f} k_j$, then $(p-1)k < p-1$, hence $k_j \leq 0$ for all $j \in \mathcal{J}$. Thus $C_j \in \text{Mat}_n(\mathbb{F}[[v]])$, and comparing determinants we see that $C_j \in \text{GL}_n(\mathbb{F}[[v]])$, for all $j \in \mathcal{J}$. (In particular $k_j = 0$ for all $j \in \mathcal{J}$.) Finally, we show that $C_j \in \mathcal{I}(\mathbb{F})$. If this were not the case, then for some $\alpha \in \Phi^-$ the entry corresponding to α in $\text{Ad}(v^{\mu_j + \eta_{0,j}}) \varphi(C_{j-1}^+)$ will have v -adic valuation $\langle \mu_j + \eta_{0,j}, \alpha^\vee \rangle$. Comparing the α entry in the equation

$$\text{Ad}(v^{\mu_j + \eta_{0,j}}) \varphi(C_{j-1}^+) = B_j^{-1} C_j^+ B'_j v^{\mu'_j - \mu_j}$$

then shows that

$$\langle \mu_j + \eta_{0,j}, \alpha^\vee \rangle \geq -e(n-1) - \max_{1 \leq i \leq n} |\mu'_{j,i} - \mu_{j,i}|$$

which contradicts the deepness assumption on μ . \square

Proof of Theorem 3.5.1. Let τ' be a type which exhibits the specialization to $\bar{\rho}^{\text{sp}}$. Let (s, μ) and (s', μ') be lowest alcove presentations of τ and τ' respectively compatible with a fixed choice of $3e(n-1)$ -generic lowest alcove presentation of $\bar{\rho}^{\text{ss}}$. Note that μ and μ' are $2e(n-1)$ -deep and satisfy the conclusion of Lemma 3.5.2. Let $\tilde{w} = \tilde{w}(\bar{\rho}, \tau)$ and $\tilde{w}' = \tilde{w}(\bar{\rho}, \tau')$.

The strategy is similar to the proof of Proposition 3.3.2. We will construct a morphism

$$(3.6) \quad \begin{aligned} \mathbb{A}_{\mathbb{F}}^1 &\rightarrow Y_{\mathbb{F}}^{\leq \eta, \tau} \\ x &\mapsto \mathfrak{M}_x \end{aligned}$$

which satisfies the following properties:

- (1) for all $x \neq 0$, the Breuil–Kisin module \mathfrak{M}_x has shape $\tilde{w}(\bar{\rho}, \tau)$;
- (2) $T_{dd}^*(\mathfrak{M}_1) \cong \bar{\rho}|_{G_{K_\infty}}$; and
- (3) $T_{dd}^*(\mathfrak{M}_0)|_{I_{K_\infty}} \cong \bar{\rho}^{\text{sp}}$.

Let $\mathfrak{M}' \in Y_{\mathbb{F}}^{\leq \eta, \tau'}(\mathbb{F})$ be the unique Breuil–Kisin module satisfying $T_{dd}^*(\mathfrak{M}') \cong \bar{\rho}|_{G_{K_\infty}}$. By Proposition 3.1.14, there is an eigenbasis β' for \mathfrak{M}' such that

$$A_{\mathfrak{M}', \beta'}^{(j)} = D_j(\tilde{w}'_j)^* U_j$$

where $D_j \in T(\mathbb{F})$ and $U_j \in N_{(\tilde{w}'_j)^*}(\mathbb{F}) \subset \mathcal{I}(\mathbb{F})$ is defined in [LLHLMa, Definition 4.2.9]. Since U_j is unipotent ([LLHLMa, Corollary 4.2.16]), there exists $s_j \in W$ such that

$$(3.7) \quad \text{Ad}(s_j(\eta_{0,j})(x)) \cdot U_j \in 1 + x \text{Mat}_n(\mathbb{F}[x][[v]]).$$

We define a map $\kappa' : (\mathbb{G}_m)_{\mathbb{F}} \rightarrow Y_{\mathbb{F}}^{\leq \eta, \tau'}$ by specifying Breuil–Kisin module $\widetilde{\mathfrak{M}}'$ over $\mathbb{F}[x^{\pm 1}]$ of type τ' and eigenbasis $\tilde{\beta}'$ such that

$$A_{\widetilde{\mathfrak{M}}', \tilde{\beta}'}^{(j)} = D_j(\tilde{w}'_j)^* \text{Ad}(s_j(\eta_{0,j})(x)) \cdot U_j$$

for all $x \neq 0$. By (3.7), this map extends to a map $\kappa' : \mathbb{A}_{\mathbb{F}}^1 \rightarrow Y_{\mathbb{F}}^{\leq \eta, \tau'}$.

The map κ' gives rise to a family $\widetilde{\mathcal{M}} \stackrel{\text{def}}{=} \varepsilon_{\tau'}(\widetilde{\mathfrak{M}}')$ of étale ϕ -modules over K parametrized by $\mathbb{A}_{\mathbb{F}}^1$. Over \mathbb{G}_m , by Proposition 3.1.10, $\widetilde{\mathcal{M}}$ admits a basis \mathfrak{f} such that

$$Q'_j \stackrel{\text{def}}{=} \text{Mat}_{\mathfrak{f}}(\phi_{\widetilde{\mathcal{M}}}^{(j)}) = D_j(\tilde{w}'_j)^* (\text{Ad}(s_j(\eta_{0,j})(x)) \cdot U_j) (s'_j)^{-1} v^{\mu'_j + \eta_{0,j}}.$$

For $x \in \mathbb{G}_m$, we write $Q'_{j,x} = \text{Mat}_{\mathfrak{f}_x}(\phi_{\widetilde{\mathcal{M}}_x}^{(j)})$ in what follows. By construction, $\mathbb{V}_K^*(\widetilde{\mathcal{M}}_1) \cong \bar{\rho}|_{G_{K_\infty}}$ and $\mathbb{V}_K^*(\widetilde{\mathcal{M}}_0)|_{I_{K_\infty}} \cong \bar{\rho}^{\text{sp}}$.

By assumption, \mathcal{M}_1 is the étale φ -module over K associated to the unique $\mathfrak{M} \in Y_{\mathbb{F}}^{\leq \eta, \tau}(\mathbb{F})$ satisfying $T_{dd}^*(\mathfrak{M}) \cong \bar{\rho}|_{G_{K_\infty}}$. Choose an eigenbasis β for \mathfrak{M} . By Proposition 3.1.10, there exists $(C_j) \in \text{GL}_n(\mathbb{F}((v)))^{\mathcal{J}}$ such that for all $j \in \mathcal{J}$

$$(3.8) \quad C^{(j+1)} Q'_{j,1} = A_{\mathfrak{M}, \beta}^{(j)} s_j^{-1} v^{\mu_j + \eta_{0,j}} \varphi(C^{(j)}).$$

Applying Lemma 3.5.3 with $B'_j = Q'_{j,1} v^{-\mu'_j - \eta_{0,j}}$ and $B_j = A_{\mathfrak{M}, \beta}^{(j)} s_j^{-1}$, we conclude that $C^{(j)} \in \mathcal{I}(\mathbb{F})$ for all $j \in \mathcal{J}$. Hence, by changing the eigenbasis of \mathfrak{M} if necessary, we can arrange that $Q'_{j,1} = A_{\mathfrak{M}, \beta}^{(j)} s_j^{-1} v^{\mu_j + \eta_{0,j}}$.

We now construct a map $\kappa : (\mathbb{G}_m)_{\mathbb{F}} \rightarrow Y_{\mathbb{F}}^{\leq \eta, \tau}$ by specifying a Breuil–Kisin module $\widetilde{\mathfrak{M}}$ over $\mathbb{F}[x^{\pm 1}]$ of type τ with eigenbasis $\tilde{\beta}$ such that

$$A_{\widetilde{\mathfrak{M}}, \tilde{\beta}}^{(j)} = Q'_j v^{-\mu_j - \eta_{0,j}} s_j = D_j(\tilde{w}'_j)^* (\text{Ad}(s_j(\eta_{0,j})(x)) \cdot U_j) (s'_j)^{-1} v^{\mu'_j - \mu_j} s_j.$$

To see that κ is well-defined, observe that

$$A_{\widetilde{\mathfrak{M}}_x, \tilde{\beta}_x}^{(j)} = t_1 A_{\widetilde{\mathfrak{M}}_1, \tilde{\beta}_1}^{(j)} t_2 = t_1 A_{\mathfrak{M}, \beta}^{(j)} t_2$$

for suitable (constant) diagonal matrices $t_1, t_2 \in T(\mathbb{F}')$ depending on $x \in (\mathbb{F}')^\times$. This also shows that κ satisfies property (1). The map κ satisfies property (2) by construction

The construction of κ shows that the fiber $\widetilde{\mathcal{M}}_x$ of the family $\widetilde{\mathcal{M}}$ of étale ϕ -modules over $\mathbb{A}_{\mathbb{F}}^1$ comes from a point of $Y_{\mathbb{F}}^{\leq \eta, \tau}$ for each $x \neq 0$. Since this is a closed condition and the map $Y_{\mathbb{F}}^{\leq \eta, \tau} \rightarrow \Phi\text{-Mod}_K^{\text{ét}, n}$ is proper (Proposition 3.1.11), it follows that κ extends to a map $\kappa : \mathbb{A}_{\mathbb{F}}^1 \rightarrow Y_{\mathbb{F}}^{\leq \eta, \tau}$, and property (3) holds for this extension. \square

The proof of Theorem 3.5.1 has the following useful consequence.

Corollary 3.5.4. *Suppose that τ and τ' are $2e(n-1)$ generic tame types with compatible lowest alcove presentations. Assume that $\varepsilon_\tau(\mathfrak{M}) \cong \varepsilon_{\tau'}(\mathfrak{M}')$ for objects $\mathfrak{M} \in Y^{\leq \eta, \tau}(\mathbb{F})$, $\mathfrak{M}' \in Y^{\leq \eta, \tau'}(\mathbb{F})$.*

Then \mathfrak{M} and \mathfrak{M}' admit eigenbases β and β' respectively such that

$$A_{\mathfrak{M},\beta}^{(j)} \tilde{w}^*(\tau) = A_{\mathfrak{M}',\beta'}^{(j)} \tilde{w}^*(\tau')$$

for all $j \in \mathcal{J}$.

3.6. Specialization pairs. In this subsection, we enhance the notion of specialization of $\bar{\rho}$ to a pair of specialization and a Serre weight. The pairs exhibit a nice combinatorial structure indexed by the Weyl group (see Definition 3.6.3).

Lemma 3.6.1. *Suppose that $\bar{\rho}^{\text{sp}}$ is an extremal specialization of $\bar{\rho}$ and that τ is a tame inertial type exhibiting this specialization and such that $\tilde{w}(\bar{\rho}, \tau) = t_{w^{-1}(e\eta_0)}$ for some $w \in W^{\mathcal{J}}$. Let $\tilde{w} \in \widetilde{W}_1^{+, \mathcal{J}}$ be an element (unique up to $X^0(\underline{T})$) whose image in $W^{\mathcal{J}}$ is w . Let τ_g be the tame inertial type with lowest alcove presentation compatible with τ such that $\tilde{w}(\bar{\rho}^{\text{sp}}, \tau_g)$ is the unique element in $\Omega^{\mathcal{J}} w_0 t_{(e-1)\eta_0} \tilde{w} \cap t_{e\eta_0} W_a^{\mathcal{J}}$. Assume that τ_g is $(e(n-1) + 1)$ -generic. Then $\bar{\rho}$ is τ_g -admissible and $\tilde{w}(\bar{\rho}, \tau_g) = \tilde{w}(\bar{\rho}^{\text{sp}}, \tau_g)$.*

Proof. Note that $t_{w^{-1}(e\eta_0)} = \tilde{w}(\tau)^{-1} \tilde{w}(\bar{\rho}^{\text{sp}}, \tau) = \tilde{w}(\bar{\rho}^{\text{sp}})$ and we can write $t_{w^{-1}(e\eta_0)} = \tilde{w}_2^{-1} w_0 t_{(e-1)\eta_0} \tilde{w}$ where $\tilde{w}_2 \in \widetilde{W}_1^{+, \mathcal{J}}$. Let $\delta \in \Omega^{\mathcal{J}}$ such that $\delta \tilde{w}_2 \in W_a^{\mathcal{J}}$. We define τ_g to be the unique tame inertial type such that $\tilde{w}(\tau_g) = \tilde{w}(\tau)(\delta \tilde{w}_2)^{-1}$. By definition τ_g is endowed with a compatible lowest alcove presentation which is $(e(n-1) + 1)$ -generic and $\tilde{w}(\bar{\rho}^{\text{sp}}, \tau_g)$ is as desired. It remains to show that $\tilde{w}(\bar{\rho}, \tau_g) = \tilde{w}(\bar{\rho}^{\text{sp}}, \tau_g)$.

Let $\mathfrak{M} \in Y^{\eta, \tau}(\mathbb{F})$ be unique Breuil–Kisin module such that $T_{dd}^*(\mathfrak{M}) \cong \bar{\rho}|_{G_{K_\infty}}$. By assumption, \mathfrak{M} has shape $t_{w^{-1}(e\eta_0)}$ and satisfies the mod p monodromy condition (Definition 3.2.3). Hence, for any choice of eigenbasis β , we have that $\iota_\tau(\mathfrak{M})$ is the étale ϕ -module with partial Frobenii given by $A_{\mathfrak{M},\beta}^{(j)} \tilde{w}^*(\tau)_j$ for $j \in \mathcal{J}$, where

$$A_{\mathfrak{M},\beta}^{(j)} \tilde{w}^*(\tau)_j \in (\mathcal{I}(\mathbb{F}) t_{w^{-1}(e\eta_{0,j})} \mathcal{I}(\mathbb{F}) \tilde{w}^*(\tau)_j)^{\nabla_0}.$$

Applying Proposition 3.2.2 with $\tilde{w} = t_{w^{-1}(e\eta_{0,j})}$, $\tilde{w}' = \delta w_0 t_{(e-1)\eta_{0,j}} \tilde{w}_{1,j}$ and $\tilde{z} = \tilde{w}^*(\tau)_j$, we have for all $j \in \mathcal{J}$:

$$A_{\mathfrak{M},\beta}^{(j)} \tilde{w}^*(\tau)_j \in \mathcal{I}(\mathbb{F}) (\delta w_0 t_{(e-1)\eta_{0,j}} \tilde{w}_{1,j})^* \mathcal{I}(\mathbb{F}) \tilde{w}^*(\tau_g)_j$$

(note that $\tilde{w} = t_{w^{-1}(e\eta_{0,j})}$, $\tilde{w}' = \delta w_0 t_{(e-1)\eta_{0,j}} \tilde{w}_{1,j}$ are e -regular and $n-1$ -small by Proposition 2.1.6). Hence there exists $\mathfrak{M}' \in Y^{\eta, \tau_g}(\mathbb{F})$ such that $\varepsilon_{\tau_g}(\mathfrak{M}') \cong \varepsilon_\tau(\mathfrak{M})$ and such that \mathfrak{M}' has shape $\delta w_0 t_{(e-1)\eta_0} \tilde{w}_1 = \tilde{w}(\bar{\rho}^{\text{sp}}, \tau_g)$. \square

Definition 3.6.2 (Specialization pairs). Suppose that $\bar{\rho}$, $\bar{\rho}^{\text{sp}}$, τ , and w are as in Lemma 3.6.1. Let $\tilde{w} \in \widetilde{W}_1^+$ be the unique element whose projection in W is w . Let σ be the Serre weight

$$(3.9) \quad F(\pi^{-1}(\tilde{w}) \cdot (\tilde{w}(\tau) \tilde{w}^{-1} \tilde{w}_h^{-1}(0) - \eta_0)) = F_{(\tilde{w}, \tilde{w}(\tau) \tilde{w}^{-1} \tilde{w}_h^{-1}(0))}.$$

Then we say that $\bar{\rho}$ specializes to the pair $(\sigma, \bar{\rho}^{\text{sp}})$ and that τ exhibits this specialization. Let $SP(\bar{\rho})$ be the set of pairs to which $\bar{\rho}$ specializes.

Note that if $\bar{\rho}^{\text{sp}}$ is $\max\{2, e\}(n-1)$ -generic and τ is $2(n-1)$ -generic then σ is the unique element in $W^?(\bar{\rho}^{\text{sp}}, \tau)$ by Proposition 2.4.8, and is the extremal weight corresponding to w (see Definition 2.4.3).

If $\bar{\rho}$ is $2e(n-1) + 1$ -generic, we have a natural map $SP(\bar{\rho}) \rightarrow S_{\text{ext}}(\bar{\rho})$ which is surjective and hence the set $SP(\bar{\rho})$ is finite. If $\zeta \in X^*(\underline{T})$ and some $\bar{\rho}^{\text{sp}} \in S_{\text{ext}}(\bar{\rho})$ has a ζ -compatible lowest alcove presentation, then every element of $S_{\text{ext}}(\bar{\rho})$ has a ζ -compatible lowest alcove presentation.

Definition 3.6.3. Assume that $\bar{\rho}$ is $(2e(n-1)+1)$ -generic. Let $\zeta \in X^*(T)$ and suppose that some (equivalently any) element of $S_{\text{ext}}(\bar{\rho})$ has a ζ -compatible lowest alcove presentation. We define a map

$$\theta_{\bar{\rho}}^{\zeta} : SP(\bar{\rho}) \rightarrow W^{\mathcal{J}}$$

as follows: If $(\sigma, \bar{\rho}^{\text{SP}})$ is in $SP(\bar{\rho})$, $(w_{\bar{\rho}^{\text{SP}}}, \mu_{\bar{\rho}^{\text{SP}}})$ is a ζ -compatible lowest alcove presentation of $\bar{\rho}^{\text{SP}}$, and σ is the extremal weight corresponding to w , we set $\theta_{\bar{\rho}}^{\zeta}(\sigma, \bar{\rho}^{\text{SP}}) = w_{\bar{\rho}^{\text{SP}}} w^{-1}$.

Proposition 3.6.4. *Assume that $\bar{\rho}$ is $3e(n-1)$ -generic. The map $\theta_{\bar{\rho}}^{\zeta}$ is injective. (Later in §5.4, we show map is bijective.)*

Proof. Suppose that $\theta_{\bar{\rho}}^{\zeta}(\sigma, \bar{\rho}^{\text{SP}}) = \theta_{\bar{\rho}}^{\zeta}(\sigma', \bar{\rho}'^{\text{SP}})$. Let τ and τ' be as in Lemma 3.6.1 exhibiting these specializations with $\tilde{w}(\bar{\rho}, \tau) = t_{w^{-1}(e\eta_0)}$ and $\tilde{w}(\bar{\rho}', \tau') = t_{(w')^{-1}(e\eta_0)}$, and let τ_g and τ'_g also be as in Lemma 3.6.1. Then σ and σ' are the extremal weights of $\bar{\rho}^{\text{SP}}$ and $\bar{\rho}'^{\text{SP}}$ corresponding to w and $w' \in W$, respectively. Let \tilde{w} and $\tilde{w}' \in \widetilde{W}_1^+$ be elements with images w and $w' \in W$, respectively. Then by Lemma 3.6.1, there exist $\delta, \delta' \in \Omega$ such that $\tilde{w}(\bar{\rho}, \tau_g) = \delta w_0 t_{(e-1)\eta_0} \tilde{w}$ and $\tilde{w}(\bar{\rho}', \tau'_g) = \delta' w_0 t_{(e-1)\eta_0} \tilde{w}'$. By Corollary 3.5.4 and the fact that $\tilde{w}^*(\tau'_g) \varphi(\mathcal{I}) \tilde{w}^*(\tau'_g)^{-1} \subset \mathcal{I}$,

$$\mathcal{I} \tilde{w}(\bar{\rho}, \tau_g)^* \mathcal{I} \tilde{w}^*(\tau_g) \cap \mathcal{I} \tilde{w}(\bar{\rho}', \tau'_g)^* \mathcal{I} \tilde{w}^*(\tau'_g) \neq \emptyset,$$

or equivalently by taking transposes,

$$(3.10) \quad \tilde{w}(\tau_g) \delta \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w} \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}} \cap \tilde{w}(\tau'_g) \delta' \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w}' \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}} \neq \emptyset,$$

where \mathcal{I}^{op} is the opposite Iwahori group scheme.

To simplify notation, let \tilde{s} and \tilde{s}' be $\tilde{w}(\tau_g) \delta$ and $\tilde{w}(\tau'_g) \delta'$, respectively. Then we have $\tilde{w}(\bar{\rho}^{\text{SP}}) = \tilde{s} w_0 t_{(e-1)\eta_0} \tilde{w}$ and $\tilde{w}(\bar{\rho}'^{\text{SP}}) = \tilde{s}' w_0 t_{(e-1)\eta_0} \tilde{w}'$. Let $s, s', w(\bar{\rho}^{\text{SP}})$, and $w(\bar{\rho}'^{\text{SP}}) \in W$ be the images of $\tilde{s}, \tilde{s}', \tilde{w}(\bar{\rho})$, and $\tilde{w}(\bar{\rho}'^{\text{SP}})$, respectively. The equality $w(\bar{\rho}^{\text{SP}}) w^{-1} = \theta_{\bar{\rho}}^{\zeta}(\sigma, \bar{\rho}^{\text{SP}}) = \theta_{\bar{\rho}}^{\zeta}(\sigma', \bar{\rho}'^{\text{SP}}) = w(\bar{\rho}'^{\text{SP}}) (w')^{-1}$ implies that $s = s'$.

The previous paragraph and (3.10) and imply that there exists $\nu \in X^*(T)$ such that

$$t_{\nu} \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w} \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}} \cap \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w}' \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}} \neq \emptyset.$$

Both $t_{\nu} \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w} \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}}$ and $\mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w}' \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}}$ are stable under the left action of T . There is a \mathbb{G}_m -subgroup which contracts $t_{\nu} \mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w} \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}}$ to $t_{\nu} w_0 t_{(e-1)\eta_0} \tilde{w}$. So $t_{\nu} w_0 t_{(e-1)\eta_0} \tilde{w}$ is in the closure of $\mathcal{I}^{\text{op}} w_0 t_{(e-1)\eta_0} \tilde{w}' \mathcal{I}^{\text{op}} / \mathcal{I}^{\text{op}}$, or equivalently $t_{\nu} w_0 t_{(e-1)\eta_0} \tilde{w} \leq w_0 t_{(e-1)\eta_0} \tilde{w}'$. Symmetrically, $t_{-\nu} w_0 t_{(e-1)\eta_0} \tilde{w}' \leq w_0 t_{(e-1)\eta_0} \tilde{w}$. Lemma 2.2.4 implies that $\tilde{w}' = t_{\nu} \tilde{w}$ and that $\nu \in X^0(T)$. In particular, we have $w = w'$ so that $\tau = \tau', \sigma \cong \sigma'$, and $\bar{\rho}^{\text{SP}} = \bar{\rho}'^{\text{SP}}$. \square

3.7. Extremal weights. In this section, we define extremal weights and use them to give a tameness criterion for Galois representations.

Definition 3.7.1 (Extremal weights). Let $\bar{\rho}$ be a $(2e(n-1)+1)$ -generic representation of G_K . Define $W_{\text{extr}}(\bar{\rho})$ to be the set of Serre weights σ such that there exists some $\bar{\rho}^{\text{SP}}$ so that $(\sigma, \bar{\rho}^{\text{SP}}) \in SP(\bar{\rho})$.

Proposition 3.7.2. *If $\bar{\rho}$ is semisimple and $(2e(n-1)+1)$ -generic, then $W_{\text{extr}}(\bar{\rho})$ agrees with the set $W_{\text{extr}}(\bar{\rho}|_I)$ from Definition 2.4.3.*

Proof. We first note that if $\bar{\rho}$ is semisimple, then it only specializes to $\bar{\rho}|_{I_K}$ by Proposition 3.1.13.

Fix now a $(2e(n-1)+1)$ -generic lowest alcove presentation of $\bar{\rho}|_{I_K}$. For each $w \in W^{\mathcal{J}}$, there is a unique type τ (with compatible lowest alcove presentation) such that $\tilde{w}(\bar{\rho}|_{I_K}, \tau) = \tilde{w}(\bar{\rho}, \tau) = t_{w^{-1}(e\eta_0)}$. Let $\tilde{w} \in \widetilde{W}_1^{+, \mathcal{J}}$ be the unique element whose projection in $W^{\mathcal{J}}$ is w . This type realizes the

specialization to the pair $\bar{\rho}|_{I_K}$ and $\sigma = F_{(\tilde{w}, \tilde{w}(\tau)\tilde{w}^{-1}\tilde{w}_h^{-1}(0))}$ (see (3.9)). Using that $\tilde{w}(\tau) = \tilde{w}(\bar{\rho}|_{I_K})t_{e\eta_0}$, we see that σ is the extremal weight of $\bar{\rho}|_{I_K}$ corresponding to w . \square

Proposition 3.7.3. *Assume that $\bar{\rho}$ is $3e(n-1)$ -generic. The following are equivalent:*

- (1) $\bar{\rho}$ is semisimple; and
- (2) $\#W_{\text{extr}}(\bar{\rho}) = \#W^{\mathcal{J}}$.

Proof. Proposition 3.7.2 gives (1) implies (2). Next, assume that $\#W_{\text{extr}}(\bar{\rho}) = \#W^{\mathcal{J}}$. By the injectivity of $\theta_{\bar{\rho}}$ (Proposition 3.6.4), $\bar{\rho}$ has a unique extremal specialization, call it $\bar{\rho}^{\text{sp}}$, and furthermore, $W_{\text{extr}}(\bar{\rho}) = W_{\text{extr}}(\bar{\rho}^{\text{sp}})$.

Let $w \in W^{\mathcal{J}}$. Let σ (resp. σ') be the extremal weight associated to w (resp. $w\eta_0$). We show that if $\bar{\rho}$ specializes to both $(\sigma, \bar{\rho}^{\text{sp}})$ and $(\sigma', \bar{\rho}^{\text{sp}})$ then $\bar{\rho}$ is semisimple and $\bar{\rho}|_{I_K} \cong \bar{\rho}^{\text{sp}}$. Let τ and τ' be the types realizing these specialization in shape $\tilde{z} = t_{w^{-1}(e\eta_0)}$ and $\tilde{z}' = t_{w_0w^{-1}(e\eta_0)}$ with corresponding Breuil–Kisin module \mathfrak{M} and \mathfrak{M}' .

By Proposition 3.1.14, there exists eigenbases β and β' respectively such that

$$A_{\mathfrak{M}, \beta}^{(j)} = D_j U_j \tilde{z}_j, \quad A_{\mathfrak{M}', \beta'}^{(j)} = D'_j U'_j \tilde{z}'_j$$

where $D_j, D'_j \in T(\mathbb{F})$, $U_j \in \tilde{z}_j N_{\tilde{z}_j}(\mathbb{F}) \tilde{z}_j^{-1}$, and $U'_j \in \tilde{z}'_j N_{\tilde{z}'_j}(\mathbb{F}) (\tilde{z}'_j)^{-1}$. By definition of $N_{\tilde{z}}$ ([LLHLMa, Definition 4.2.9]), we have

$$U_j, U'_j \in L^{--}\mathcal{G}_{\mathbb{F}}^{(j)}(\mathbb{F})$$

where $L^{--}\mathcal{G}_{\mathbb{F}}^{(j)}$ denotes the negative loop group for $L\mathcal{G}_{\mathbb{F}}^{(j)}$ (in particular, its \mathbb{F} points consist of matrices $A \in \text{GL}_n(\mathbb{F}[1/v])$ which are lower unipotent modulo $1/v$).

By Corollary 3.5.4, there exists $(I^{(j)}) \in \mathcal{I}(\mathbb{F})^{\mathcal{J}}$ such that

$$(3.11) \quad D_j U_j \tilde{z}_j \tilde{w}^*(\tau) = I^{(j)} D'_j U'_j \tilde{z}'_j \tilde{w}^*(\tau') (\varphi(I^{(j-1)}))^{-1}$$

By scaling β' by an element of $T(\mathbb{F})^{\mathcal{J}}$ if necessary, we can arrange that $(I^{(j)}) \in \mathcal{I}_1(\mathbb{F})^{\mathcal{J}}$. Since both τ and τ' realize the same specialization, $\tilde{z}_j \tilde{w}^*(\tau) = \tilde{z}'_j \tilde{w}^*(\tau') = \tilde{w}^*(\bar{\rho}^{\text{sp}})$ and so (3.11) becomes

$$(3.12) \quad D_j U_j \tilde{w}^*(\bar{\rho}^{\text{sp}}) = I^{(j)} D'_j U'_j \tilde{w}^*(\bar{\rho}^{\text{sp}}) (\varphi(I^{(j-1)}))^{-1}.$$

By Lemma 3.1.4, there exists $(X_j) \in \mathcal{I}_1(\mathbb{F})^{\mathcal{J}}$ such that $D_j U_j = X_j D'_j U'_j$ for all $j \in \mathcal{J}$. Thus, $U_j (U'_j)^{-1} \in \mathcal{I}(\mathbb{F}) \cap L^{--}\mathcal{G}_{\mathbb{F}}^{(j)}(\mathbb{F})$ and so $U_j = U'_j$. Finally, since $t_{w^{-1}(e\eta_0)}(A_0)$ and $t_{w_0w^{-1}(e\eta_0)}(A_0)$ are in opposite Weyl chambers, $\tilde{z}_j^{-1} U_j \tilde{z}_j$ and $(\tilde{z}'_j)^{-1} U'_j \tilde{z}'_j$ are in opposite unipotents by [LLHLMa, Corollary 4.2.15]. Thus, U_j and U'_j are the identity ($\tilde{z}_j, \tilde{z}'_j$ are both translations). Since $A_{\mathfrak{M}, \beta}^{(j)} = D_j \tilde{z}_j$ for all $j \in \mathcal{J}$, it follows that $T_{dd}^*(\mathfrak{M})$ is semisimple (see [LLHLMa, Proposition 5.5.2] for example). \square

3.8. Maximally ordinary weights. In this section, we show that the set $W_{\text{extr}}(\bar{\rho})$ contains the set of *maximally ordinary weights*. We further show that the set of maximally ordinary weights is nonempty, so that in particular, the set $W_{\text{extr}}(\bar{\rho})$ is nonempty. When $\bar{\rho}$ is an iterated extension of characters, the set of maximally ordinary weights is the set of *ordinary weights*.

Lemma 3.8.1. *Suppose that $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ is $(e(n-1) + 2)$ -generic and an extension of $\bar{\rho}_2$ by $\bar{\rho}_1$. For $i = 1$ and 2 , let n_i be the dimension of $\bar{\rho}_i$. Suppose that $\bar{\rho}_i$ has a potentially crystalline lift $\rho_i : G_K \rightarrow \text{GL}_n(\mathcal{O}_E)$ of tame inertial type τ_i and parallel Hodge–Tate weights $(n_1 + n_2 - 1, \dots, n_2)$ (resp. $(n_2 - 1, \dots, 0)$) if $i = 1$ (resp. $i = 2$). Then $\bar{\rho}$ has a lift ρ which is an extension of ρ_2 by ρ_1 and is potentially crystalline of type (τ, η) where $\tau = \tau_1 \oplus \tau_2$.*

Proof. Note that by genericity, both $\bar{\rho}$ and τ are at least 2-generic, in particular are cyclotomic-free ([LLHLMa, Lemma 7.2.9]). By genericity, $\text{Ext}_{G_K}^2(\rho_2, \rho_1)$ is zero. So the natural reduction map $\text{Ext}_{G_K}^1(\rho_2, \rho_1) \rightarrow \text{Ext}_{G_K}^1(\bar{\rho}_2, \bar{\rho}_1)$ is surjective. We conclude that there exists a lift $\rho : G_K \rightarrow \text{GL}_n(\mathcal{O}_E)$ of $\bar{\rho}$ which is an extension of ρ_2 by ρ_1 .

Let $\rho_{i,E}$ be $\rho_i \otimes_{\mathcal{O}_E} E$. Then the containment $H_g^1(G_K, \rho_{2,E}^\vee \otimes_E \rho_{1,E}) \subset H^1(G_K, \rho_{2,E}^\vee \otimes_E \rho_{1,E})$ is an equality for dimension reasons. Indeed, by the local Euler characteristic formula and Tate duality, we have that $h^1(G_K, \rho_{2,E}^\vee \otimes_E \rho_{1,E}) = \dim_E \rho_{2,E}^\vee \otimes_E \rho_{1,E}$. On the other hand, $h_g^1(G_K, \rho_{2,E}^\vee \otimes_E \rho_{1,E}) = \dim_E D_{\text{dR}}(\rho_{2,E}^\vee \otimes_E \rho_{1,E}) / D_{\text{dR}}(\rho_{2,E}^\vee \otimes_E \rho_{1,E})^+$. Since the Hodge–Tate weights of ρ_2 are strictly less than those of ρ_1 , this latter expression is $\dim_E \rho_{2,E}^\vee \otimes_E \rho_{1,E}$ as well. We conclude that ρ is an \mathcal{O}_E -lattice in a potentially semistable representation. Moreover, ρ has parallel Hodge–Tate weights η .

There is an exact sequence of smooth I_K -representations

$$0 \rightarrow D_{\text{pst}}(\rho_{1,E}) \rightarrow D_{\text{pst}}(\rho_E) \rightarrow D_{\text{pst}}(\rho_{2,E}) \rightarrow 0.$$

We conclude that $D_{\text{pst}}(\rho_E) \cong \tau = \tau_1 \oplus \tau_2$. Moreover, by genericity, $\text{Hom}_{I_K}(\tau_2, \tau_1(-1)) = 0$ and so ρ must be potentially crystalline. \square

Let $P^\vee \subset \text{GL}_n$ be a parabolic subgroup with Levi quotient M^\vee . Then $M^\vee \cong \prod_{i=1}^k M_i^\vee$ where $M_i^\vee \cong \text{GL}_{n_i}$ and $\sum_{i=1}^k n_i = n$. Let N_i be $\sum_{j=i+1}^k n_j$. We index these dimensions so that for all $1 \leq i \leq k$, P^\vee has a quotient P_i^\vee which is isomorphic to a parabolic subgroup of GL_{N_i} with Levi quotient $\prod_{j=i+1}^k M_j^\vee$. In other words, if P is block upper diagonal, then starting from the top left, the i -th block has size n_i .

Theorem 3.8.2. *Let $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be $(2e(n-1)+1)$ -generic. Suppose that $\bar{\rho}$ factors through $P^\vee(\mathbb{F})$ for a parabolic subgroup $P^\vee \subset \text{GL}_n$ as above. Let M^\vee , M_i^\vee , and N_i be as above. Suppose that the associated representations $\bar{\rho}_i : G_K \rightarrow M_i^\vee(\mathbb{F})$ are semisimple. For each i , let ρ_i be a potentially crystalline lift of type τ_i and parallel Hodge–Tate weights $(n_i + N_i - 1, \dots, N_i)$ where $\tilde{w}(\bar{\rho}_i(-N_i), \tau_i)$ is extremal. Then $\bar{\rho}$ has a potentially diagonalizable lift ρ (in the sense of [BLGGT14, §1.4]) of type (τ, η) where $\tau = \bigoplus_{i=1}^k \tau_i$. The corresponding specialization is $\bigoplus_{i=1}^k \bar{\rho}_i$. In particular, the semisimplification $\bigoplus_{i=1}^k \bar{\rho}_i$ is an extremal specialization of $\bar{\rho}$.*

Proof. By iterated application of Lemma 3.8.1, we obtain a potentially crystalline lift ρ of type (τ, η) , which is an iterated extension of potentially crystalline lifts (of type τ_i and parallel Hodge–Tate weights $(n_i + N_i - 1, \dots, N_i)$) of the representations $\bar{\rho}_i$. In particular, the semisimplification of ρ is $\bigoplus_{i=1}^k \rho_i$. Then by the argument of proof of [LLHL19, Corollary 3.4.11] (replacing the reference to Proposition 3.4.8 in *loc. cit.* with Proposition 4.2.2 below, and noting that the semisimple Kisin module produced as in *loc. cit.* has Hodge–Tate weights exactly η), after restriction to a finite index subgroup the semisimplification of ρ is a direct sum of characters. By [BLGGT14, Lemma 1.4.3(1)], ρ is potentially diagonalizable.

Since $\tilde{w}(\bar{\rho}_i(-N_i), \tau_i)$ is extremal for all $i = 1, \dots, k$, so is $\tilde{w}(\bigoplus_{i=1}^k \bar{\rho}_i, \tau)$ by an easy computation. Since the semisimplification of $\bar{\rho}$ is $\bigoplus_{i=1}^k \bar{\rho}_i$, we deduce that $\tilde{w}(\bar{\rho}, \tau)$ is this same shape by Proposition 3.3.2. Thus τ exhibits the specialization $\bigoplus_{i=1}^k \bar{\rho}_i$ of ρ . \square

Suppose that $\bar{\rho}$ is as in Theorem 3.8.2 with P^\vee , M^\vee , and N_i as before. Let $P \subset \text{GL}_n$ be the dual parabolic subgroup. Let U be the unipotent radical of P . For each i let $\sigma_i \in W_{\text{extr}}(\bar{\rho}_i)$. Let σ be the unique Serre weight such that

$$\sigma^U \cong \boxtimes_{i=1}^k \sigma_i(-N_i).$$

We call a Serre weight constructed in this way *maximally ordinary*. Let $W_{\text{mord}}(\bar{\rho})$ be the set of maximally ordinary Serre weights. Since we can always find P^\vee as in Theorem 3.8.2, $W_{\text{mord}}(\bar{\rho})$ is nonempty. If $\bar{\rho}$ is semisimple, then we can take P^\vee to be GL_n so that $W_{\text{mord}}(\bar{\rho}) = W_{\text{extr}}(\bar{\rho})$. Taking P^\vee to be a minimal parabolic when $\bar{\rho}^{\text{ss}}$ is a direct sum of characters, we see that ordinary weights are maximally ordinary.

Proposition 3.8.3. *There is an inclusion $W_{\text{mord}}(\bar{\rho}) \subset W_{\text{extr}}(\bar{\rho})$.*

Proof. Let $\bar{\rho}$ and $\bar{\rho}_i$ be as in Theorem 3.8.2. Suppose that $\sigma \in W_{\text{mord}}(\bar{\rho})$. For each i , let τ_i be the tame type such that $W^2(\bar{\rho}_i, \tau_i) = \{\sigma_i\}$. Then if we let τ be $\bigoplus_{i=1}^k \tau_i$, then τ exhibits a specialization of $\bar{\rho}$ to $\bar{\rho}^{\text{ss}}$. Moreover, one can check that $W^2(\bar{\rho}^{\text{ss}}, \tau) = \{\sigma\}$ so that $(\sigma, \bar{\rho}^{\text{ss}}) \in SP(\bar{\rho})$. This shows that $\sigma \in W_{\text{extr}}(\bar{\rho})$. \square

3.9. Connections to Emerton–Gee stacks. This section is a series of remarks explaining how the notions of extremal weight and specialization can be interpreted geometrically on the stack of mod p Galois representations \mathcal{X}_n introduced by Emerton–Gee [EGa]. When K/\mathbb{Q}_p is unramified, everything can be proved using the techniques of [LLHLMa]. The ramified case requires extending [LLHLMa] which will be the subject of future work.

First, we briefly recall what we need from [EGa]. In [EGa, Theorem 6.5.1], Emerton and Gee describe a parametrization of the irreducible components of the underlying reduced stack $\mathcal{X}_{n,\text{red}}$ of the moduli of (φ, Γ) -modules \mathcal{X}_n by Serre weights of $\text{GL}_n(\mathcal{O}_K)$. Let $\sigma = F(\kappa)$ be a Serre weight of $\text{GL}_n(\mathcal{O}_K)$ with $\kappa = (\kappa_j) \in X_1(T)^\mathcal{J}$. We use the normalization as in [LLHLMa] where $\mathcal{C}_\sigma \stackrel{\text{def}}{=} \mathcal{X}_{EG,n,\text{red}}^{\sigma^\vee \otimes \det^{n-1}}$.

If κ is 1-deep, then \mathcal{C}_σ is uniquely characterized by the fact that has a Zariski open subset consisting of $\bar{\rho}$ of the form

$$\bar{\rho} \cong \begin{pmatrix} \chi_1 & * & \cdots & * \\ 0 & \chi_2 & \cdots & * \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \chi_n \end{pmatrix}$$

where $\chi_i|_{I_K} = \bar{\varepsilon}^i \prod_{j \in \mathcal{J}} \bar{\omega}_{K,\sigma_j}^{(\kappa_j)_i}$ and $\bar{\rho}$ admits a unique G_K -stable flag.

Remark 3.9.1. (1) Let $\bar{\rho}^{\text{sp}}$ be a sufficiently generic tame \mathbb{F} -type. If σ is an extremal weight of $\bar{\rho}^{\text{sp}}$ as in Definition 2.4.3, then there is a Zariski open subset $\mathcal{C}_\sigma^{\bar{\rho}^{\text{sp}}} \subset \mathcal{C}_\sigma$ such that $\bar{\rho}$ specializes to the pair $(\sigma, \bar{\rho}^{\text{sp}})$ if and only if $\bar{\rho} \in \mathcal{C}_\sigma^{\bar{\rho}^{\text{sp}}}$. The Zariski open can be constructed via the generalization of the diagram in [LLHLMa, Theorem 7.4.2] to the ramified setting.
 (2) Let σ be a sufficiently generic Serre weight. Then there are $(n!)^\mathcal{J}$ sufficiently generic tame \mathbb{F} -types which have σ as an extremal for $\bar{\rho}$ corresponding to w weight. Thus, the union $\mathcal{C}_\sigma^{\text{extr}} = \bigcup \mathcal{C}_\sigma^{\bar{\rho}^{\text{sp}}}$ where $\bar{\rho}^{\text{sp}}$ ranges over all such types is a Zariski open subset of \mathcal{C}_σ consisting exactly of the $\bar{\rho}$ which have σ as an extremal weight. (One can check that $\mathcal{C}_\sigma^{\text{extr}} = \mathcal{C}_\sigma$ only when K/\mathbb{Q}_p is unramified and σ is Fontaine–Laffaille.)

Remark 3.9.2. As has been introduced in other settings ([LLHLMb]), there is a natural set of Serre weights that can be associated to an arbitrary $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$, the *geometric weights*,

$$W^g(\bar{\rho}) = \{\sigma \mid \bar{\rho} \in \mathcal{C}_\sigma(\mathbb{F})\}.$$

Remark 3.9.1 says that $W_{\text{extr}}(\bar{\rho}) \subset W^g(\bar{\rho})$. Generally speaking the set of geometric weights will be larger.

3.10. The extremal locus. In this section, we discuss the relationship between $W_{\text{extr}}(\bar{\rho})$ and $W^g(\bar{\rho})$ when K/\mathbb{Q}_p is unramified. This gives, in this setting, an alternative to the proof of the existence of extremal weights in §3.8. The main result of this section will also be used to construct global lifts in §5.5.

Let K/\mathbb{Q}_p be unramified. Let (\tilde{w}_1, ω) be a lowest alcove presentation for a Serre weight σ compatible with $\zeta \in X^*(Z)$. Recall from [LLHLMa, Definition 4.6.1] that C_σ^ζ is the closure of

$$(\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I}(t_\omega)^*)^{\nabla_0}$$

inside $\text{Fl}_{\mathcal{J}}^{\nabla_0}$ (see also §3.2). We define $C_{\sigma, \text{extr}}^\zeta$ to be the (Zariski) open subset

$$\cup_{w \in \underline{W}} (\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I}(t_\omega w)^*)^{\nabla_0} \subset C_\sigma^\zeta.$$

Assume that σ is $(3n - 1)$ -deep. Then by [LLHLMa, Remark 7.4.3(2)], we have a local model diagram for \mathcal{C}_σ (the irreducible component of $\mathcal{X}_{n, \text{red}}$ corresponding to σ , cf. §3.9) and C_σ^ζ . We then let $\mathcal{C}_{\sigma, \text{extr}} \subset \mathcal{C}_\sigma$ be the Zariski open set of \mathcal{C}_σ corresponding to $C_{\sigma, \text{extr}}^\zeta \subset C_\sigma^\zeta$. (The definition of $\mathcal{C}_{\sigma, \text{extr}}$ does not depend on the lowest alcove presentation of σ .)

Proposition 3.10.1. *Let $\bar{\rho}$ be $(2(n - 1) + 1)$ -generic. If σ is $(3n - 1)$ -deep and $\bar{\rho} \in \mathcal{C}_\sigma$, then $\bar{\rho} \in \mathcal{C}_{\sigma, \text{extr}}$ if and only if $\sigma \in W_{\text{extr}}(\bar{\rho})$.*

Proof. We fix a lowest alcove presentation (\tilde{w}_1, ω) for σ compatible with ζ . Let $x \in C_\sigma^\zeta$ correspond to $\bar{\rho} \in \mathcal{C}_\sigma$ in the local model diagram [LLHLMa, Theorem 7.4.2]. If $\sigma \in W_{\text{extr}}(\bar{\rho})$, then let τ be a tame inertial type exhibiting the extremal weight σ . Then

$$\begin{aligned} x &\in (\mathcal{I} \backslash \mathcal{I} t_{(w_1)^{-1}(\eta)}^* \mathcal{I} \tilde{w}(\tau)^*)^{\nabla_0} \\ &= (\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I}(\tilde{w}(\tau)(\tilde{w}_h \tilde{w}_1)^{-1})^*)^{\nabla_0} \\ &= (\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I}(t_\omega w(\tau)(w_0 w_1)^{-1})^*)^{\nabla_0} \\ &\subset C_{\sigma, \text{extr}}^\zeta \end{aligned}$$

where $\tilde{w}(\tau)$ is defined with respect to the lowest alcove presentation of τ compatible with ζ .

Conversely, suppose that $\bar{\rho} \in \mathcal{C}_{\sigma, \text{extr}}$. Let $w \in \underline{W}$ be such that $x \in (\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I}(t_\omega w)^*)^{\nabla_0}$. Then we let τ be such that $\tilde{w}(\tau) = t_\omega w \tilde{w}_h \tilde{w}_1$. The above calculation shows that τ exhibits σ as an extremal weight of $\bar{\rho}$. \square

Proposition 3.10.2. *Assume that σ is $(4n - 2)$ -deep. There is an inclusion*

$$\mathcal{C}_\sigma \subset \cup_{\sigma \text{ covers } \sigma'} \mathcal{C}_{\sigma', \text{extr}}.$$

Proof. We choose a $(4n - 2)$ -deep lowest alcove presentation (\tilde{w}_1, ω) of σ and will show that

$$C_\sigma^\zeta \subset \cup_{\sigma \text{ covers } \sigma'} C_{\sigma', \text{extr}}^\zeta.$$

Since the elements of \widetilde{W} less than or equal to $w_0 \tilde{w}_1$ are exactly those of the form $s\tilde{w}$ for some $s \in \underline{W}$ and $\tilde{w} \in \widetilde{W}^+$ with $\tilde{w} \uparrow \tilde{w}_1$ (see the proof of Lemma 2.2.2), [IM65, Proposition 2.8] gives

$$C_\sigma^\zeta \subset \overline{\mathcal{I} \backslash \mathcal{I}(w_0 \tilde{w}_1)^* \mathcal{I} t_\omega^*}^{\nabla_0} = \cup_{s \in \underline{W}} \cup_{\substack{\tilde{w} \in \widetilde{W} \\ \tilde{w} \uparrow \tilde{w}_1}} (\mathcal{I} \backslash \mathcal{I}(s\tilde{w})^* \mathcal{I} t_\omega^*)^{\nabla_0}.$$

We will show that $(\mathcal{I} \backslash \mathcal{I}(s\tilde{w})^* \mathcal{I} t_\omega^*)^{\nabla_0} \subset C_{\sigma', \text{extr}}^\zeta$ for some σ' which σ covers.

Since $(w_0s^{-1})s\tilde{w}$ is a reduced factorization by Lemma 2.2.1,

$$\begin{aligned} (\mathcal{I}\backslash\mathcal{I}(s\tilde{w})^*\mathcal{I}t_\omega^*)^{\nabla_0} &\subset (\mathcal{I}\backslash\mathcal{I}(s\tilde{w})^*\mathcal{I}(w_0s^{-1})^*\mathcal{I}((w_0s^{-1})^{-1})^*t_\omega^*)^{\nabla_0} \\ &= (\mathcal{I}\backslash\mathcal{I}(w_0\tilde{w})^*\mathcal{I}(t_\omega sw_0^{-1})^*)^{\nabla_0}. \end{aligned}$$

To further analyze this, let $\tilde{w} = t_\nu\tilde{w}'_1$ where $\nu \in X^*(\underline{I})$ is dominant and $\tilde{w}'_1 \in \widetilde{W}_1^+$. Then $t_{w_0(\nu)}w_0\tilde{w}'_1$ is a reduced expression for $w_0\tilde{w}$ by [LLHL19, Lemma 4.1.9], from which we deduce as before that

$$\begin{aligned} (\mathcal{I}\backslash\mathcal{I}(w_0\tilde{w}'_1)^*\mathcal{I}(t_{\omega+s(\nu)}sw_0^{-1})^*)^{\nabla_0} &= (\mathcal{I}\backslash\mathcal{I}(w_0\tilde{w}'_1)^*\mathcal{I}(t_\omega sw_0^{-1}t_{w_0(\nu)})^*)^{\nabla_0} \\ &\subset (\mathcal{I}\backslash\mathcal{I}(w_0\tilde{w})^*\mathcal{I}(t_\omega sw_0^{-1})^*)^{\nabla_0}. \end{aligned}$$

On the other hand, these are irreducible varieties of the same dimension by [LLHLMa, Theorem 4.2.4] and thus must be equal. Letting σ' be the Serre weight with lowest alcove presentation $(\tilde{w}'_1, \omega + s(\nu))$, we have $(\mathcal{I}\backslash\mathcal{I}(w_0\tilde{w}'_1)^*\mathcal{I}(t_{\omega+s(\nu)}sw_0^{-1})^*)^{\nabla_0} \subset C_{\sigma', \text{extr}}^\zeta$. (Note that σ' is $(3n-1)$ -deep, hence $\mathcal{C}_{\sigma', \text{extr}}$ is defined.) It suffices to show that σ covers σ' , or by [LLHLMa, Proposition 2.3.12(ii)] that $t_{W(\nu)}\tilde{w}'_1 \uparrow \tilde{w}_1$. However, we have $t_{\underline{W}(\nu)}\tilde{w}'_1 \uparrow \tilde{w} \uparrow \tilde{w}_1$ where the first inequality follows from [Jan03, II.6.5(3)]. \square

Proposition 3.10.3. *Let K/\mathbb{Q}_p be a finite unramified extension and $\bar{\rho} : G_K \rightarrow \text{GL}_n(\mathbb{F})$ be a Galois representation. Let τ be a $(5n-1)$ -generic tame inertial L -parameter. Then the following are equivalent.*

- (1) $R_{\bar{\rho}}^{\tau}$ is nonzero;
- (2) $\bar{\rho}$ is $4n$ -generic and $W^g(\bar{\rho}) \cap \text{JH}(\bar{\sigma}(\tau)) \neq \emptyset$; and
- (3) $\bar{\rho}$ is $4n$ -generic and $W_{\text{extr}}(\bar{\rho}) \cap \text{JH}(\bar{\sigma}(\tau)) \neq \emptyset$.

Proof. (1) and (2) are equivalent by [LLHLMa, Theorem 7.4.2(1)]. Since $W_{\text{extr}}(\bar{\rho}) \subset W^g(\bar{\rho})$, (3) implies (2). For the converse, suppose that $\bar{\rho} \in \mathcal{C}_\sigma$ for some $\sigma \in \text{JH}(\bar{\sigma}(\tau))$. Proposition 3.10.2 implies that $\bar{\rho} \in \mathcal{C}_{\sigma', \text{extr}}$ for some σ' which σ covers. Then $\sigma' \in W_{\text{extr}}(\bar{\rho})$ by Proposition 3.10.1 and $\sigma' \in \text{JH}(\bar{\sigma}(\tau))$ by the definition of covering. (Note that Propositions 3.10.1, 3.10.2 apply by the genericity assumption on τ .) \square

4. SOME POTENTIALLY CRYSTALLINE DEFORMATION RINGS

The aim of this section is to compute potentially crystalline deformation rings for a certain class of shapes, namely those related to the subgroup $W_{a,\alpha} \subseteq \widetilde{W}$ defined in 2.2. We follow the general procedure appearing in [LLHL19], improved in [LLHLMa].

4.1. The main result on Galois deformation rings. For a mod p Galois representation $\bar{\rho}$, we write $R_{\bar{\rho}}^{\eta,\tau}$ (resp. $R_{\bar{\rho}}^{\leq\eta,\tau}$) for the framed universal deformation ring of $\bar{\rho}$ of tame inertial type τ for I_K over E and parallel Hodge–Tate weights η (resp. $\leq\eta$). The main result is the following:

Theorem 4.1.1. *Let τ be a $\max\{(3n-7)e-(n-2), (2n-3)e\}$ -generic tame inertial type. Suppose that $\widetilde{w}(\bar{\rho}, \tau)$ is $\widetilde{w}^{-1}t_{e\eta_0}\widetilde{w}_\alpha\widetilde{w}$ for some $\widetilde{w} \in \widetilde{W}_1^+$, some $\alpha \in \Delta^{\mathcal{J}}$, and $\widetilde{w}_{\alpha_j} \in W_{a,\alpha_j}$.*

Then $R_{\bar{\rho}}^{\eta,\tau} = R_{\bar{\rho}}^{\leq\eta,\tau}$ is either zero or is a normal domain. Furthermore:

- *If \widetilde{w}_{α_j} is id or $t_{-e\alpha_j}$ for each j , $R_{\bar{\rho}}^{\eta,\tau}$ is formally smooth over \mathcal{O} .*
- *In general, $\text{Spec } \overline{R_{\bar{\rho}}^{\leq\eta,\tau}}$ is reduced with 2^m geometrically irreducible components of the same dimension, where $m = \#\{j \in \mathcal{J} \mid \widetilde{w}_{\alpha_j} \neq \text{id}, t_{-e\alpha_j}\}$.*

Remark 4.1.2. A key ingredient in our proof of Theorem 4.1.1 is the fact that the local model (in the sense of [LLHLMa]) of our Galois deformation ring has a Levi reduction property: namely, it is formally smooth over a similar local model attached to a Levi subgroup of GL_n . This turns out to be a general phenomenon whenever the shape $\widetilde{w}(\bar{\rho}, \tau)^*$ is suitably “decomposable”, which may be of independent interest. In the specific case of Theorem 4.1.1, the Levi subgroup we can reduce to is $\text{GL}_2 \times \text{GL}_1^{n-2}$, which is why we have very precise control on the relevant local models, and hence the Galois deformation rings.

4.2. Gauge bases and parabolic structures. For each $j \in \mathcal{J}$, we set $E_j = \sigma_j(E(v)) \in \mathcal{O}[v]$. Let R be an \mathcal{O} -algebra. We have the usual notion of degrees on $R[v]$, which is submultiplicative $\deg(ab) \leq \deg(a) + \deg(b)$, with equality if either a or b are monic (but not in general). The notion degree and being monic extends elements of $R[v, E_j^{-1}]$. The set of elements of degree ≤ 0 form a subring of $R[v, E_j^{-1}]_{\leq 0}$. This subring contains the set of elements $R[v, E_j^{-1}]_{\leq -1} = R[v, E_j^{-1}]_{< 0}$ of degree < 0 as an ideal, and another ideal given by $vR[v, E_j^{-1}]_{< 0}$. More generally, the set $R[v, E_j^{-1}]_{\leq d}$ of elements of degree $\leq d$ form an $R[v, E_j^{-1}]_{\leq 0}$ -module.

Concretely, the elements of $R[v, E_j^{-1}]_{\leq 0}$ are exactly those of the form $\frac{P}{E_j^m}$ with $P \in R[v]$ such that $\deg P \leq me$, with the extra condition $v \mid P$ (for some choice of fractions with m sufficiently large) for elements of $vR[v, E_j^{-1}]_{< 0}$, and the extra condition $\deg P < me$ for elements of $R[v, E_j^{-1}]_{< 0}$. Finally, note that for an element a represented by $\frac{P}{E_j^m}$ with $P(v) \in R[v]$, the \mathcal{O} -algebra generated by the coefficients of P is independent of the choice of representing fraction.

Let R be a Noetherian \mathcal{O} -algebra. We define

$$LG^{(j)}(R) \stackrel{\text{def}}{=} \{A \in \text{GL}_n(R[v]^{\wedge E_j}[\frac{1}{E_j}]), A \text{ is upper triangular mod } v\};$$

$$L^+ \mathcal{M}^{(j)}(R) \stackrel{\text{def}}{=} \{A \in \text{Mat}_n(R[v]^{\wedge E_j}), A \text{ is upper triangular mod } v\};$$

For $\widetilde{z} = zt_\nu \in \widetilde{W}^\vee$ such that $e \mid \|\nu\|$, define $\mathcal{U}(\widetilde{z})^{\det, \leq h}(R)$ to be the collection of $A \in LG^{(j)}(R)$ such that

- For $1 \leq i, k \leq n$,

$$A_{ik} = v^{\delta_{i>k}} \frac{P}{E_j^h}$$

with $P \in R[v]$ such that $\deg P \leq he + \nu_k - \delta_{i>k} - \delta_{i<z(k)}$. Furthermore, this is an equality when $i = z(k)$, in which case P is monic. In particular, $A \in \frac{1}{E_j^h} L^+ \mathcal{M}^{(j)}(R)$.

- $\det A = \det(z) E_j^{\frac{\|\nu\|}{e}}$.

If R is furthermore \mathcal{O} -flat, then for such A we have

$$A_{ik}^{-1} = v^{\delta_{i>k}} \frac{Q}{E_j^H}$$

with $Q \in R[v]$ and H sufficiently large, such that $\deg Q \leq He - \nu_i - \delta_{i>k} - \delta_{z(i)<k}$ (The condition that R is \mathcal{O} -flat is used to show that divisibility by v in $R[v, E_j^{-1}]$ is equivalent to evaluating to 0 at $v = 0$, and hence the numerators of all representing fractions have 0 constant terms).

For each $j \in \mathcal{J}$, we define $U^{[a,b]}(\tilde{z}_j) \subset \mathcal{U}^{\det, \leq -a}(\tilde{z}_j)$ to be the subfunctor consisting of A such that $E_j^b A^{-1} \in L^+ \mathcal{M}^{(j)}(R) \cap LG^{(j)}(R)$ and $E_j^{-a} A \in L^+ \mathcal{M}^{(j)}(R) \cap LG^{(j)}(R)$. This is clearly representable by a finite type affine \mathcal{O} -scheme, with a set of generators given by the coefficients of the entries of $A^{(j)}$. Note that this depends on j , a choice that is implicit in the symbol \tilde{z}_j .

If $\tilde{z} = (\tilde{z}_j)_j \in \widetilde{W}^{\vee, \mathcal{J}}$, we set $U^{[a,b]}(\tilde{z}) = \prod U^{[a,b]}(\tilde{z}_j)$. We have the following definition:

Definition 4.2.1. Let (R, \mathfrak{m}) be a complete local Noetherian \mathcal{O} -algebra and assume that $\mathfrak{M} \in Y^{[0, n-1], \tau}(R)$ such that $\mathfrak{M} \otimes_R R/\mathfrak{m}$ has shape \tilde{z} with respect to τ . An eigenbasis β for \mathfrak{M} is said to be a *gauge basis* if $A_{\mathfrak{M}, \beta}^{(j)} \in T^\vee(R) U^{[0, n-1]}(\tilde{z}_j)(R)$ for all $j \in \mathcal{J}$.

Proposition 4.2.2. *Assume that τ admits a $(e(n-1)+1)$ -deep lowest alcove presentation. Suppose R is a complete local Noetherian \mathcal{O} -algebra and let $\mathfrak{M} \in Y^{[0, n-1], \tau}(R)$ such that $\overline{\mathfrak{M}} \in Y^{[0, n-1], \tau}(\mathbb{F})$ has shape \tilde{z} with respect to τ . Then \mathfrak{M} has a gauge basis. Moreover the set of gauge basis for \mathfrak{M} is a torsor for the natural action of $T^{\vee, \mathcal{J}}(R)$.*

Proof. The proof of [LLHLMa, Proposition 5.2.7] generalizes verbatim by replacing the reference to Proposition 5.1.8 in *loc. cit.* by Remark 1 above, and noting that the statement of Lemma 5.1.10 in *loc. cit.* holds true in our setting. Note the proof in *loc. cit.* in fact proves a more general statement where R is only assumed to be merely p -adically complete. \square

Suppose we are given a gauge basis $\bar{\beta}$ for $\overline{\mathfrak{M}} \in Y^{[0, n-1], \tau}(\mathbb{F})$ with shape \tilde{z} and write

$$A_{\overline{\mathfrak{M}}, \bar{\beta}}^{(j)} = \overline{D}^{(j)} \overline{U}^{(j)}$$

where $\overline{D}^{(j)} \in T^\vee(\mathbb{F})$, $\overline{U}^{(j)} \in U^{[0, n-1]}(\tilde{z}_j)(\mathbb{F})$.

If R is a complete local Noetherian \mathcal{O} -algebra, and $\mathfrak{M} \in Y^{[0, n-1], \tau}(R)$ is such that $\mathfrak{M} \otimes_R \mathbb{F} \cong \overline{\mathfrak{M}}$, then the set of gauge basis for \mathfrak{M} lifting $\bar{\beta}$ is a torsor under the natural action of $\ker(T^{\vee, \mathcal{J}}(R) \rightarrow T^{\vee, \mathcal{J}}(\mathbb{F}))$. Thus, the functor representing deformations (\mathfrak{M}, β) of the pair $(\overline{\mathfrak{M}}, \bar{\beta})$ is representable by the completion of $T^\vee U^{[0, n-1]}(\tilde{z}_j)$ at the point corresponding to $(\overline{D}^{(j)} \overline{U}^{(j)})$, and it is formally smooth over the completion of $Y^{[0, n-1], \tau}$ at $\overline{\mathfrak{M}}$. The subfunctor classifying deformations (\mathfrak{M}, β) such that \mathfrak{M} furthermore belongs to $Y^{\leq \eta, \tau}$ correspond to the completion of the closed subscheme $T^\vee U(\tilde{z}, \leq \eta)$ of $T^\vee U^{[0, n-1]}(\tilde{z})$ characterized by:

- $T^\vee U(\tilde{z}, \leq \eta)$ is \mathcal{O} -flat and reduced.

- The elementary divisors of $(A^{(j)} \in T^\vee U(\tilde{z}, \leq \eta)(R) \subset \prod LG^{(j)}(R)$ are bounded by $E_j^{(n-1, \dots, 0)}$, i.e. for each $1 \leq k \leq n$, each $k \times k$ minors of $A^{(j)}$ (which belong to $R[v]$) are divisible by $E_j^{\frac{(k-1)k}{2}}$ (in $R[v]$).

Remark 4.2.3. Let

$$LG^{+, (j)}(R) \stackrel{\text{def}}{=} \{A \in \text{GL}_n(R[v]^{\wedge E_j}), A \text{ is upper triangular mod } v\}$$

a twisted positive loop group. Then $\text{Gr}_{\mathcal{G}}^{(j)} = LG^{+, (j)} \backslash LG^{(j)}$ is a twisted affine Grassmannian. Then the generic fiber $\text{Gr}_{\mathcal{G}, E}^{(j)} \cong (\text{Gr}_{\text{GL}_n, E})^e$ identifies with the product of e copies of the affine Grassmannian for the split group GL_n , while the special fiber $\text{Gr}_{\mathcal{G}, \mathbb{F}}^{(j)} \cong \text{Fl}$ identifies with the affine flag variety. The Pappas–Zhu local model $M_j(\leq \eta)$ for $\text{Res}_{\mathcal{O}_K \otimes W^{(k), \sigma_j} \mathcal{O} / \mathcal{O}} \text{GL}_n$ as defined in [Lev16] is the Zariski closure of the open Schubert variety for the cocharacter $(n-1, n-2, \dots, 1, 0)$ for each copy of $\text{Gr}_{\text{GL}_n, E}$. In this setup, the scheme $U(\tilde{z}_j, \leq \eta)$ identifies with an (possibly empty) open affine subscheme of $M_j(\leq \eta)$, cf. the discussion preceding [LLHLMa, Theorem 5.3.3]. In particular, if non-empty, $U(\tilde{z}_j, \leq \eta)$ has dimension $e \sum_{\beta > 0} \langle \eta_0, \beta^\vee \rangle = e \frac{(n-1)n(n+1)}{6}$.

The following Proposition shows that in certain cases, any element of $T^\vee U(\tilde{z}, \leq \eta)$ automatically acquires a parabolic structure. In Propositions 4.2.4 and 4.2.5, we work with fixed $j \in \mathcal{J}$ and drop the subscript for notational ease.

Proposition 4.2.4. *Let $w \in W^\vee$, $r + s = n$ and $\tilde{z} = zt_\nu = \begin{pmatrix} \tilde{z}_t & 0 \\ 0 & \tilde{z}_b \end{pmatrix} \in \widetilde{W}^\vee$ with block sizes r, s .*

Let $w = w_M w^M$ be the factorization so that w^M has minimal length and $w_M = (w_t, w_b) \in W(M) = W(\text{GL}_r) \times W(\text{GL}_s)$ (where M is the standard Levi for the partition $r + s = n$). Assume that

- $w_b^{-1} \tilde{z}_b w_b$ has elementary divisors bounded by $v^{e(s-1, \dots, 0)}$.
- $v^{-es} w_t^{-1} \tilde{z}_t w_t$ has elementary divisors bounded by $v^{e(r-1, \dots, 0)}$.

Suppose R is a \mathcal{O} -flat algebra and $A \in T^\vee U(w^{-1} \tilde{z} w, \leq \eta)(R)$. Then $A = Dw^{-1}Pw$ with $D \in T^\vee(R)$ and

$$P = \begin{pmatrix} M_t & 0 \\ X & M_b \end{pmatrix}$$

is parabolic with diagonal block sizes r, s , and furthermore:

- (1) $M_t \in E_j^s w_t U(t_{-se(1, \dots, 1)} w_t^{-1} \tilde{z}_t w_t) w_t^{-1}$ and has elementary divisors bounded by $E_j^{(n-1, \dots, s)}$.
- (2) $M_b \in w_b U(w_b^{-1} \tilde{z}_b w_b) w_b^{-1}$ and has elementary divisors bounded by $E_j^{(r-1, \dots, 0)}$.
- (3) $(X M_t^{-1})_{ik} \in v^{\delta_{w^{-1}(i) > w^{-1}(k)}} R[v, E_j^{-1}] \cap R[v, E_j^{-1}]_{\leq -\delta_{w^{-1}(i) < w^{-1}(k)}}$.
- (4) $(M_b^{-1} X)_{ik} \in v^{\delta_{w^{-1}(i) > w^{-1}(k)}} R[v] \cap R[v, E_j^{-1}]_{\leq \nu_k - \nu_i - \delta_{w^{-1}z(i) < w^{-1}z(k)}}$. (In the last two items, we interpret the indices to run over the rows and columns of X as a submatrix of P , i.e. $r+1 \leq i \leq n$, $1 \leq k \leq r$.)

Proof. We write $A = Dw^{-1}Pw$ so that $P \in wU(w^{-1} \tilde{z} w, \leq \eta)w^{-1}$. This means that P has entries in $R[v]$, with the degree bounds

$$P_{ik} \in v^{\delta_{w^{-1}(i) > w^{-1}(k)}} R[v] \cap R[v, E_j^{-1}]_{\leq \nu_k - \delta_{w^{-1}(i) < w^{-1}z(k)}}$$

and that the leading coefficient of $P_{iz(k)}$ are 1. We call the corresponding entry the pivot entries.

Write $P = \begin{pmatrix} M_t & Y \\ X & M_b \end{pmatrix}$. We first show that $Y = 0$. The degree bounds on P imply that when expanding $\det M_b$, there is a unique maximal degree term, which is given by the product of the top degree terms in the pivot entries in M_t (one can see this by noting that this is a combinatorial statement on the degree bounds which can be checked over rings S where $p = 0$, where it reduces to the fact that $M_b v^{-\nu_b} z_b^{-1}$ is conjugate to a matrix in with coefficients in $S[v^{-1}]$ which is upper triangular unipotent mod $v^{-1}S[v^{-1}]$). This shows $\det M_b = \det z_b E_j^{\frac{(s-1)s}{2}}$. Now

$$(Y M_b^{-1})_{ik} = \sum_l Y_{il} (M_b^{-1})_{lk}.$$

We observe

- $Y_{il} \in v^{\delta_{w^{-1}(i) > w^{-1}(l)}}$, $(M_b^{-1})_{lk} \in v^{\delta_{w^{-1}(l) > w^{-1}(k)}}$. Hence $(Y M_b^{-1})_{ik}$ is divisible by $v^{\delta_{w^{-1}(i) > w^{-1}(k)}}$ in $R[v, E_j^{-1}]$.
- $Y_{il} \in R[v, E_j^{-1}]_{\leq \nu_l - \delta_{w^{-1}(i) < w^{-1}(z(l))}}$, $(M_b^{-1})_{lk} \in R[v, E_j^{-1}]_{\leq -\nu_l - \delta_{w^{-1}(z(l)) < w^{-1}(k)}}$. Hence $(Y M_b^{-1})_{ik} \in R[v, E_j^{-1}]_{\leq -\delta_{w^{-1}(i) < w^{-1}(k)}}$.

However, the elementary divisor conditions together with the degree bounds imply that the minor formed by replacing one row of M_b with one row of Y belongs to $E_j^{\frac{(s-1)s}{2}} R$, hence Cramer's rule shows that the entries of $Y M_b^{-1}$ are in R . Since by the above, these entries also belong to $v^{\delta_{w^{-1}(i) < w^{-1}(k)}} R[v, E_j^{-1}] \cap R[v, E_j^{-1}]_{\leq -\delta_{w^{-1}(i) < w^{-1}(k)}}$, they must be all 0.

Thus, we see that P has the desired parabolic structure. The first two items immediately follow from the degree bounds on P and the elementary divisor conditions. The third and fourth items follow from the same argument used above in showing $Y = 0$. \square

By applying Proposition 4.2.4 to the universal case, we get

Proposition 4.2.5. *Assume the setting of Proposition 4.2.4. Let $R^{\text{univ}} = \mathcal{O}(U(w^{-1}\tilde{z}w, \leq \eta))$, so that the universal $A^{\text{univ}} \in U(\tilde{z}, \leq \eta)$ factors as*

$$A^{\text{univ}} = D^{\text{univ}} w^{-1} \begin{pmatrix} M_t^{\text{univ}} & 0 \\ X^{\text{univ}} & M_b^{\text{univ}} \end{pmatrix} w.$$

Then the map $A^{\text{univ}} \mapsto (\frac{1}{E_j} w_t^{-1} M_t^{\text{univ}} w_t, w_b^{-1} M_b w_b)$ exhibits $U(w^{-1}\tilde{z}w, \leq \eta)$ as an affine space over $U(t_{-se(1, \dots, 1)} w_t^{-1} \tilde{z}_t w_t, \leq (r-1, \dots, 0)) \times U(w_b^{-1} \tilde{z}_b w_b, \leq (s-1, \dots, 0))$, whose coordinates are the coefficients of the entries of $(M_b^{\text{univ}})^{-1} X^{\text{univ}}$ (which are subject to the degree bounds dictated by Proposition 4.2.4).

Proof. The fact that we get a map follows from Proposition 4.2.4, which clearly induces a closed immersion from $U(w^{-1}\tilde{z}w, \leq \eta)$ into the appropriate affine space over $U(t_{-se(1, \dots, 1)} w_t^{-1} \tilde{z}_t w_t, \leq (r-1, \dots, 0)) \times U(w_b^{-1} \tilde{z}_b w_b, \leq (s-1, \dots, 0))$. To see this injection is an isomorphism, observe that if we set Z to be a matrix subject to the degree bounds of Proposition 4.2.4(4) and whose coefficients are free variables, then

$$w^{-1} \begin{pmatrix} M_t^{\text{univ}} & 0 \\ M_b^{\text{univ}} Z & M_b^{\text{univ}} \end{pmatrix} w = w^{-1} \begin{pmatrix} M_t^{\text{univ}} & 0 \\ 0 & M_b^{\text{univ}} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ Z & 0 \end{pmatrix} w$$

satisfies the necessary elementary divisors and degree bounds characterizing $U(w^{-1}\tilde{z}w, \leq \eta)$. \square

4.3. Interlude: GL_2 Pappas–Zhu models. We specialize the previous section to $n = 2$. Thus, $M_j(t_{(1,0)})$ is a Pappas–Zhu local model for the Weil restricted group $\mathrm{Res}_{\mathcal{O}_K \otimes_{W(k), \sigma_j} \mathcal{O}/\mathcal{O}} \mathrm{GL}_2$, the (minuscule) cocharacter $(t_{(1,0)}, \dots, t_{(1,0)}) \in (\mathbb{Z}^2)^e$, and Iwahori level structure. The following summarizes the known geometric properties of $M_j(t_{(1,0)})$ (see Theorem A in [PR05] or Theorem 2.3.3 and 2.3.5 in [Lev16]):

Proposition 4.3.1. (1) $M_j(t_{(1,0)})_E \cong (\mathbb{P}_E^1)^e$.

(2) $M_j(t_{(1,0)})_{\mathbb{F}}$ is (geometrically) reduced, and identifies with the reduced union of $S(t_{(e,0)}) \cup S(t_{(0,e)})$ of $\mathrm{Fl} = \mathrm{Gr}_{\mathbb{F}}^{(j)}$. Each of its irreducible components are (geometrically) normal.

In particular, $M_j(t_{(1,0)})$ is a normal domain, whose special fiber has two irreducible components. Further more any $x \in S^0(t_{(e,0)}) \cup S^0(t_{(0,e)})$ belongs to the regular locus of $M_j(t_{(1,0)})$.

Note that the reducedness of the special fiber and the geometric normality of its irreducible component are preserved under taking products.

We note that the $e(1,0)$ -admissible elements are exactly $t_{(e-k,k)}$, $0 \leq k \leq e$ and $t_{(e-k,k)} s_\alpha$ with $0 < k \leq e$.

Corollary 4.3.2. Let \tilde{z}_j be $(e,0)$ -admissible. Then $U(\tilde{z}_j, \leq \eta)$ is a normal domain, and it is formally smooth over \mathcal{O} if $\tilde{z}_j \in \{t_{(e,0)}, t_{(0,e)}\}$. Otherwise, its special fiber has two (geometrically) normal irreducible components.

We deduce the following combinatorial property about the admissible set from our geometric considerations:

Corollary 4.3.3. Let $\tilde{w}_j = w_j^{-1} W_{a, \alpha_j} t_{e\eta_0} w_j \cap \mathrm{Adm}(e\eta_0)$ for some simple root α_j and $w_j \in W$. Then there are at most two $\sigma \in W$ such that $\tilde{w}_j \leq t_{\sigma^{-1}(e\eta_0)}$.

Proof. Set $\tilde{z}_j = \tilde{w}_j^*$. By Proposition 4.2.5, $U(\tilde{z}_j, \leq \eta)$ is an affine space over an affine scheme of the form as in Corollary 4.3.2. In particular, $U(\tilde{z}_j, \leq \eta)_{\mathbb{F}}$ has at most two irreducible components. On the other hand, this is an open neighborhood of \tilde{z}_j in the special fiber $M_j(\leq \eta)_{\mathbb{F}}$ of a Pappas–Zhu model. We conclude from the fact that $M_j(\leq \eta)_{\mathbb{F}} = \cup_{\sigma \in W} S(t_{\sigma(e\eta_0)})$. \square

4.4. Analysis of the monodromy condition. Suppose $\bar{\rho}$ admits a Breuil–Kisin module $\overline{\mathfrak{M}} \in Y^{[0, n-1], \tau}(\mathbb{F})$ of type τ , with shape \tilde{z} and a gauge basis $\bar{\beta}$. To analyze the potentially crystalline deformation ring $R_{\bar{\rho}}^{\leq \eta, \tau}$, we need to recall its relationship with the finite height deformation ring $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}}$, as in [LLHL19, §3,4] and [LLHLMa, §7.1]. One has a diagram (cf [LLHL19, Diagram (3.16)], [LLHLMa, Proposition 7.2.3])

$$(4.1) \quad \begin{array}{ccccc} & & \mathrm{Spf} R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \square, \nabla} & \xrightarrow{f.s.} & \mathrm{Spf} R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla} \\ & \nearrow \cong & \downarrow & & \downarrow \\ \mathrm{Spf} R_{\overline{\mathfrak{M}}, \bar{\rho}}^{\tau, \bar{\beta}, \square} & \hookrightarrow & \mathrm{Spf} R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \square} & \xrightarrow{f.s.} & \mathrm{Spf} R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}} \\ \downarrow f.s. & & & & \\ \mathrm{Spf} R_{\bar{\rho}}^{\leq \eta, \tau} & & & & \end{array}$$

where

- $R_{\bar{\rho}}^{\leq \eta, \tau}$ is the framed potentially crystalline deformation ring representing Galois deformations ρ with Hodge-Tate weights $\leq \eta$ and inertial type τ . Note that it is either zero, or is \mathcal{O} -flat, reduced and of Krull dimension $n^2 + 1 + \frac{n(n-1)}{2}[K : \mathbb{Q}_p]$.
- $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}$ represents deformations (\mathfrak{M}, β) of $(\bar{\mathfrak{M}}, \bar{\beta})$ where \mathfrak{M} belongs to $Y^{\leq \eta, \tau}$ and β is a gauge basis of \mathfrak{M} .
- $R_{\bar{\mathfrak{M}}, \bar{\rho}}^{\tau, \bar{\beta}, \square}$ represents potentially crystalline Galois deformations ρ of type $(\leq \eta, \tau)$, together with a gauge basis β of its (unique) Breuil–Kisin module \mathfrak{M} in $Y^{\leq \eta, \tau}$. It is formally smooth over $R^{\leq \eta, \tau}$ of relative dimension nf .
- $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}, \square}$ represents a deformation (\mathfrak{M}, β) of $(\bar{\mathfrak{M}}, \bar{\beta})$ as above together with a framing basis of the G_{K_∞} -representation associated to \mathfrak{M} . This is formally smooth over $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}$ of relative dimension n^2 .
- $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla}$ (resp. $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}, \square, \nabla}$) is the \mathcal{O} -flat reduced quotient of $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}$ (resp. $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}, \square}$) cut out by imposing the monodromy condition on the universal Breuil–Kisin module after inverting p .

We elaborate on the monodromy condition on the universal Breuil–Kisin module on $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}$. Recall that $E(v)$ is the Eisenstein polynomial of a chosen uniformizer of K over K_0 , and that $e' = p^{f'} - 1 = p^{f^r} - 1$. Recall from [LLHLMa, §7.1] the ring $\mathcal{O}^{\text{rig}} \stackrel{\text{def}}{=} \mathcal{O}_{K', R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}}^{\text{rig}}$, endowed with a canonical derivation

$N_{\nabla} = -u' \lambda \frac{d}{du'}$ (where $\lambda = \prod_{i=0}^{\infty} \frac{\varphi^i(E((u')^{e'}))}{E(0)}$ is constructed out of $E(v) = E((u')^{e'})$ instead of $v+p$), and the module $\mathfrak{M}^{\text{univ, rig}} \stackrel{\text{def}}{=} \mathfrak{M}^{\text{univ}} \otimes_{R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}} \mathcal{O}^{\text{rig}}$, such that $\mathfrak{M}^{\text{univ, rig}}[1/\lambda]$ is endowed with a canonical derivation $N_{\mathfrak{M}^{\text{univ, rig}}}$ over N_{∇} (cf. [LLHLMa, Proposition 7.1.3(1)]). Then the *monodromy condition* alluded to above is the condition that $N_{\mathfrak{M}^{\text{univ, rig}}}$ preserves $\mathfrak{M}^{\text{univ, rig}}$.

We now choose a lowest alcove presentation $\tau \cong \tau(s, \mu + \eta_0)$. Recall from §2.3.3 that attached to (s, μ) we have the data $s'_{\text{or}, j'} \in W$, $\mathbf{a}'^{(j')} \in \mathbb{Z}^n$. We write $A^{(j')}$ for the matrices constructed out of the universal Breuil–Kisin module and its universal gauge basis over $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}}$ or $R_{\bar{\mathfrak{M}}}^{\tau, \bar{\beta}, \square}$. We get the following control of the monodromy condition:

Proposition 4.4.1. *Assume $\tau(s, \mu + \eta_0)$ is an m -deep lowest alcove presentation of τ . If the monodromy condition holds, then for each j' , $0 \leq t < n - 2$ and π a root of E_j , the result of the operator $(\frac{d}{dv})^t|_{v=\pi}$ acting on*

$$\left(e'v \frac{d}{dv} A^{(j')} + [A^{(j')}, \text{Diag}((s'_{\text{or}, j'})^{-1}(\mathbf{a}'^{(j')}))] \right) (A^{(j')})^{-1} E_j^{n-1}$$

belongs to $p^{\frac{m+1-(n-2)e-t}{e}} R$.

Proof. This is a straightforward generalization of the computation in [LLHLMa, Proposition 7.1.10], with the following changes: h in *loc.cit.* becomes $n - 1$, occurrences of p (outside any evaluation at $v = -p$) becomes $E(0)$, occurrences of $(v + p)^h (A^{(j')})^{-1}$ becomes $E_{j'}^{n-1} (A^{(j')})^{-1}$, occurrences of $|_{v=-p}$ becomes $|_{v=\pi}$. Note that $E_j(0) \in p\mathcal{O}^\times$, and $E_{j'} = E_j$ depends only on $j \bmod f$. More specifically, the computation in *loc.cit.* expresses the monodromy condition as

$$\varphi(\lambda)^{n-1} \left(e'v \frac{d}{dv} A^{(j')} + [A^{(j')}, \text{Diag}((s'_{\text{or}, j'})^{-1}(\mathbf{a}'^{(j')}))] \right) (A^{(j')})^{-1} E_j^{n-1} + \text{Err}$$

has zeroes of order $n - 2$ along the roots of E_j , for an appropriate error term Err . It follows that the operator $(\frac{d}{dv})^t|_{v=\pi}$ annihilates this expression, for $0 \leq t < n - 2$ and π a root of E_j .

The error term Err has the form

$$\sum_{i=1}^{\infty} \varphi^{i+1}(\lambda)^{n-1} Z_i^{(j')}$$

with $Z_i^{(j')} \in \frac{1}{p^{i(n-2)}} v^{1+m\frac{p^i-1}{p-1}} \text{Mat}_n(R[[v]])$. We conclude from the analysis of the effect of $(\frac{d}{dv})^t|_{v=\pi}$ on the error term Err as in *loc.cit.* (except that we use the differential operator $\frac{d}{dv}$ as opposed to $v\frac{d}{dv}$), noting that in our current situation

- $(\frac{d}{dv})^t|_{v=\pi} \varphi^k(\lambda) \in p^{1+\frac{p-t}{e}} \mathcal{O} + p^{p-\frac{t}{e}} \mathcal{O}$ for any $t, k \geq 1$.
- If $F \in v^M \text{Mat}_n(R[[v]])$ then $(\frac{d}{dv})^t|_{v=\pi} F \in p^{\frac{M-t}{e}} \text{Mat}_n(R[[v]])$.

□

Lemma 4.4.2. *Let R be a p -flat \mathcal{O} -algebra. Let N, k be non-negative integers and $F \in R[v]$. Assume that $N < p$ and that $(\frac{d}{dv})^t|_{v=\pi}(F) \in p^k R$ for $0 \leq t < N$ and π is any root of E_j . Write $F = E_j^N q + r$ where $q, r \in R[v]$ such that $\deg r < Ne$ (this uniquely determines q, r) Then $r \in p^{k-(2N-1)(1-\frac{1}{e})}[v]$*

Proof. Our hypothesis implies $(\frac{d}{dv})^t|_{v=\pi}(r) \in p^k R$ for $t < N$ and $E_j(\pi) = 0$. We decompose $r = \sum_{t=0}^{N-1} E_j^t r_t$ with $\deg r_i < e$. Then the reduction to R/p^k of the coefficients of r_0 form an element in the kernel space of the Vandermonde matrix on the roots of E_j . It follows that $r_0 \in p^{k-\frac{e-1}{e}} R[v]$. For $t \geq 0$, Then $t!(E_j')^t(\pi)r_t(\pi)$ differs from $(\frac{d}{dv})^t|_{v=\pi}(r)$ by a polynomial in the coefficients of $r_{t'}$ for $t' < t$. This implies $r_t \in p^{k-(2t+1)(1-\frac{1}{e})} R[v]$ by induction on t . □

The following Lemma studies the effect of the approximation of the monodromy condition under the presence of a suitable parabolic structure:

Lemma 4.4.3. *Let R be a Noetherian \mathcal{O} -algebra, N, r, s non-negative integers such that $r + s = n$. Let $\kappa = \begin{pmatrix} \kappa_t & 0 \\ 0 & \kappa_b \end{pmatrix} \in X^*(T) \otimes \mathcal{O}$ viewed as a constant diagonal matrix, $w \in W^\vee$, and $\tilde{z} = zt_\nu = \begin{pmatrix} \tilde{z}_t & 0 \\ 0 & \tilde{z}_b \end{pmatrix}$ (with block sizes r, s). Suppose we are also given $P = \begin{pmatrix} A & 0 \\ C & D \end{pmatrix} \in \text{Mat}_n(R[[v]])$, a block lower triangular matrix corresponding to the partition $r + s = n$ satisfying*

$$(4.2) \quad (v\frac{d}{dv}P - [P, \kappa])P^{-1} \in \frac{1}{E_j} \text{Mat}_n(R[[v]])$$

Assume the following

- (1) $\begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \in wU(w^{-1}\tilde{z}w)w^{-1}$.
- (2) For $\beta \in \Phi$, the β -th entry of CA^{-1} (inserted inside Mat_n at the same position as C) belong to $v^{\delta_{w^{-1}(\beta) < 0}} R[[v, E_j^{-1}]]_{< 0}$.
- (3) $E_j^N P^{-1} \in \text{Mat}_n(R[[v]])$.
- (4) $\langle \tilde{z}(0) - z(\kappa), \beta^\vee \rangle + k \in \mathcal{O}^\times$ for all $k \in \{0, \dots, -Ne\}$.

Let \mathcal{O}_P be the \mathcal{O} -algebra generated by the coefficients of the entries of P , and $\mathcal{O}_{A,D}$ be the \mathcal{O} -algebra generated by the coefficients of the entries of A, D . Then \mathcal{O}_P is generated over $\mathcal{O}_{A,D}$ by at most e elements.

Proof. In this proof only, we abbreviate $\delta_\beta = \delta_{w^{-1}(\beta) < 0}$, to avoid cluttering notation.

Our hypothesis on $\begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix}$ implies

$$w^{-1} \begin{pmatrix} (v \frac{d}{dv} A - [A, \kappa_t]) A^{-1} & 0 \\ 0 & (v \frac{d}{dv} D - [D, \kappa_b]) D^{-1} \end{pmatrix} w \in \text{Mat}_n(R[v, E_j^{-1}]_{\leq 0}),$$

and whose entries above the diagonal are in $R[v, E_j^{-1}]_{< 0}$ and whose entries on and below the diagonal are in $vR[v, E_j^{-1}]_{< 0}$. Furthermore, modulo $R[v, E_j^{-1}]_{< 0}$, the diagonal part is exactly $\frac{vE'_j}{E_j} \text{Ad}(z) \nu_e + (1 - \text{Ad}(z))(\kappa)$.

Now

$$(v \frac{d}{dv} P - [P, \kappa]) P^{-1} = \begin{pmatrix} v \frac{d}{dv} A - [A, \kappa_t] & 0 \\ v \frac{d}{dv} C - C\kappa_t + \kappa_b C & v \frac{d}{dv} D - [D, \kappa_b] \end{pmatrix} \begin{pmatrix} A^{-1} & 0 \\ -D^{-1} C A^{-1} & D^{-1} \end{pmatrix}$$

Set $B = C A^{-1}$, then the bottom left block of the above expression is

$$\begin{aligned} & (v \frac{d}{dv} (BA) - BA\kappa_t + \kappa_b BA) A^{-1} - (v \frac{d}{dv} D - [D, \kappa_b]) D^{-1} B \\ &= v \frac{d}{dv} B - B\kappa_t + \kappa_b B + B(v \frac{d}{dv} A - [A, \kappa_t]) A^{-1} - (v \frac{d}{dv} D - [D, \kappa_b]) D^{-1} B \end{aligned}$$

We abbreviate $\nabla B = v \frac{d}{dv} B - B\kappa_t + \kappa_b B$, $\nabla A = (v \frac{d}{dv} A - [A, \kappa]) A^{-1}$ and $\nabla D = (v \frac{d}{dv} D - [D, \kappa_b]) D^{-1}$.

In what follows, we label the entries of various matrices of size smaller than $n \times n$ using roots/indices of the $n \times n$ matrix P , by interpreting such matrices as one of the non-trivial block of P corresponding to its size. We observe:

- $E_j \nabla(A) = A_{e-1} + \cdots + A_0$, where $A_{i,\beta} = v^{\delta_\beta + i} a_{i,\beta}$, $A_{i,ll} = v^{1+i} a_{i,ll}$ with $a_{i,\beta}, a_{i,ll} \in R$, for all β and l such that the relevant entry exists in A .
- $E_j \nabla(D) = D_{e-1} + \cdots + D_0$, where $D_{i,\beta} = v^{\delta_\beta + i} d_{i,\beta}$, $D_{i,ll} = v^{1+i} d_{i,ll}$ with $d_{i,\beta}, d_{i,ll} \in R$, for all β and l such that the relevant entry exists in D .
- The matrices A_{lead} , D_{lead} obtained by extracting the degree e coefficients of $E_j \nabla(A)$, $E_j \nabla(D)$ satisfy

$$w^{-1} \begin{pmatrix} A_{lead} & 0 \\ 0 & D_{lead} \end{pmatrix} w$$

is lower triangular, with diagonal entries $\text{Ad}(w^{-1})(z(\nu) + \kappa - z(\kappa))$.

- $B = \frac{1}{E_j^N} (B_0 + B_{-1} + \cdots)$ where $B_{i,\beta} = b_{i,\beta} v^{\delta_\beta + N e - 1 + i}$ with $b_{i,\beta} \in R$, and $b_{i,\beta} = 0$ if $i < -N e$.

Condition (4.2) means

$$\nabla B + B(\nabla A) - (\nabla D)B \in \frac{1}{E_j} \text{Mat}_n(R[v]).$$

Using $v \frac{d}{dv} (\frac{F}{E_j^N}) = -N \frac{vE'_j}{E_j^{N+1}} F + \frac{v}{E_j^N} \frac{dF}{dv}$, clearing denominators in the above expression yields

(4.3)

$$-N v E'_j (E_j^N B) + v E_j \frac{d}{dv} (E_j^N B) - E_j (E_j^N B) \kappa_t + E_j \kappa_b (E_j^N B) + (E_j^N B) (E_j \nabla A) - (E_j \nabla D) (E_j^N B) = E_j^N X$$

for some $X \in \text{Mat}_n(R[v])$.

The observations on the degree ranges of A_i, D_i, B_i show that for each relevant $\beta \in \Phi$, $X_\beta = v^{\delta_\beta} \sum_{i \geq 0} x_{i,\beta} v^i$ (recall that an element of $R[v][\frac{1}{E_j}]$ is divisible by v if and only if its evaluation at $v = 0$ is 0, a condition that makes sense because $R \subset R[\frac{1}{p}]$).

The degree $Ne + e - 1 + i + \delta_\beta$ part of the β -th entry of equation (4.3) reads

$$(4.4) \quad -Neb_{i,\beta} + (\delta_\beta + Ne - 1 + i)b_{i,\beta} + \langle \kappa, \beta^\vee \rangle b_{i,\beta} + O(> i, \beta) + \\ + \sum_{\beta = \beta' + \beta''} \sum_{k,l} b_{k,\beta'} a_{l,\beta''} + \sum_{\beta = \gamma' + \gamma''} \sum_{k',l'} d_{l',\gamma'} b_{k',\gamma''} = X_{e-1+i,\beta} + O(> e - 1 + i, \beta)$$

where

- The symbol $O(> i, \beta)$ (resp. $O(> e - 1 + i, \beta)$) stands for a polynomial with \mathcal{O} -coefficients in $b_{i',\beta}$ (resp. $X_{e-1+i',\beta}$) for $i' > i$.
- The decompositions $\beta = \beta' + \beta''$ runs over decompositions in Φ , with the added possibility that $\beta'' = 0$, in which case $a_{l,\beta''}$ is interpreted as the unique diagonal term $a_{l,tt}$ that contributes to the β -entry of the matrix product. A similar remark applies to $\beta = \gamma' + \gamma''$.
- The pairs k, l and k', l' are constrained by

$$Ne - 1 + k + l + \delta_{\beta'} + \delta_{\beta''} = Ne + e - 1 + i + \delta_\beta$$

$$Ne - 1 + k' + l' + \delta_{\gamma'} + \delta_{\gamma''} = Ne + e - 1 + i + \delta_\beta.$$

In particular, we learn that $k \geq i$ (resp. $k' \geq i$), with equality if and only if $l = e - 1$ and $\delta_\beta + 1 = \delta_{\beta'} + \delta_{\beta''}$ (resp. $l' = e - 1$ and $\delta_\beta + 1 = \delta_{\gamma'} + \delta_{\gamma''}$). Also observe that when $k = i$ the product $b_{k,\beta'} a_{l,\beta''}$ (resp. $d_{l',\gamma'} b_{k',\gamma''}$) is zero as soon as $\delta_{\beta''} = 0$ (resp. $\delta_{\gamma''} = 0$).

Let $\mathcal{O}_{A,D,B-top}$ be the \mathcal{O} -algebra generated by the coefficients of A, D and B_i for $i \geq 1 - e$. The above observation implies that $X_\beta v^{-\delta_\beta}$ has degree $\leq e - 1$, and each of its coefficients belong to $\mathcal{O}_{A,D,B-top}$.

We now show that the coefficient of each entry of B_i belongs to $\mathcal{O}_{A,D,B-top}$ by downward induction on i . The claim clearly holds for $i \geq 1 - e$. Suppose it holds up to $i + 1$. Let B_i^+ be the matrix given by $B_{i,\beta}^+ = \delta_\beta B_{i,\beta}$. It follows (using $\delta_{\beta'} + \delta_{\beta''} = 1 + \delta_\beta = 2$ if and only if $\delta_{\beta'} = \delta_{\beta''} = 1$) from the above facts that

$$iB_i^+ + (\kappa_b - D_{lead})B_i^+ - B_i^+(\kappa_t - A_{lead}) \in M_{s \times r}(\mathcal{O}_{A,D,B-top}[v])$$

As in Proposition 4.2.4, the element $w \in W(\mathrm{GL}_n)$ induces an element $(w_t, w_b) \in W(\mathrm{GL}_r) \times W(\mathrm{GL}_s)$. We then have $\mathrm{Ad}(w_t^{-1})(A_{lead}), \mathrm{Ad}(w_b^{-1})(D_{lead})$ are lower triangular. Thus Lemma 4.4.4 below applies, and shows $B_i^+ \in M_{s \times r}(\mathcal{O}_{A,D,B-top}[v])$. Now set $B_i^- = B_i - B_i^+$. Using what we just proved, we also get

$$(i - 1)B_i^- + (\kappa_b - D_{lead})B_i^- - B_i^-(\kappa_t - A_{lead}) \in M_{s \times r}(\mathcal{O}_{A,D,B-top}[v])$$

and the same argument shows $B_i^- \in M_{s \times r}(\mathcal{O}_{A,D,B-top}[v])$. This finishes the inductive step.

Finally, since $C = BA$, \mathcal{O}_P also belongs to $\mathcal{O}_{A,D,B-top}$. \square

Lemma 4.4.4. *Let R be a ring with a subring S , $r + s = n$, $w_1 \in W(\mathrm{GL}_r), w_2 \in W(\mathrm{GL}_s)$. Suppose we are given $A_1 \in M_r(S), A_2 \in M_r(S), B \in M_{s \times r}(R)$ such that*

- $\mathrm{Ad}(w_i^{-1})(A_i)$ is lower triangular for $i = 1, 2$.
- If s_1 is a diagonal entry of A_1 and s_2 is a diagonal entry of A_2 , then $s_1 - s_2 \in S^\times$.
- $BA_1 - A_2B \in M_{s \times r}(S)$.

Then $B \in M_{s \times r}(S)$.

Proof. Replacing B by $w_2 B w_1^{-1}$, we may assume $w_1 = 1$, $w_2 = 1$. In this case, looking at the (k, l) -th entry of $BA_1 - A_2 B$ shows that $(s_1 - s_2)B_{kl}$ belong to the subalgebra generated by S and $B_{k'l'}$ with $k' - l' < k - l$, where s_1, s_2 are suitable diagonal entries of A_1, A_2 . We conclude induction on $k - l$ that $B_{kl} \in S$. \square

Remark 4.4.5. Suppose that in the setting of Lemma 4.4.3, we don't have equation (4.2) exactly but only an approximately: for $0 \leq t < N$ and π a root of E_j , the operator $(\frac{d}{dv})^t(E_j^{N+1} \cdot)|_{v=\pi}$ hitting on the matrix in (4.2) belongs to $p^k \text{Mat}_n(R)$. Then the proof shows that the conclusion of Lemma 4.4.3 also holds approximately: there is an \mathcal{O} -subalgebra S of R generated over $\mathcal{O}_{A,D}$ by at most ers elements such that $\mathcal{O}_P \subset S + p^{k-(2N-1)(1-\frac{1}{e})}R$. This follows from Lemma 4.4.2.

4.5. Proof of Theorem 4.1.1.

Proof. We recall the setting of Theorem 4.1.1. We are given τ , a tame inertial type over E , together with a fixed lowest alcove presentation (s, μ) for it, such that μ is $\max\{(3n-7)(e-1) + 2n-6, (2n-3)e\}$ -deep. Furthermore, $\tilde{w}(\bar{\rho}, \tau) = (\tilde{w}_j^{-1} t_{e\eta} \tilde{w}_{\alpha_j} \tilde{w}_j)_j$ for some simple root α_j for each $j \in \mathcal{J}$.

We assume $R_{\bar{\rho}}^{\leq \eta, \tau} \neq 0$, otherwise there is nothing to prove. In particular we obtain $\overline{\mathfrak{M}} \in Y^{\leq \eta, \tau}(\mathbb{F})$ such that $T_{dd}^*(\overline{\mathfrak{M}}) \cong \bar{\rho}|_{I_K}$. Then $\overline{\mathfrak{M}}$ has shape $w^{-1} \tilde{z} w = \tilde{w}(\bar{\rho}, \tau)^*$.

We need to analyze $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla}$ in the context of diagram (4.1).

We first observe that for each j , \tilde{z}_j has a block diagonal structure

$$\tilde{z}_j = \begin{pmatrix} \tilde{z}_{j,t} & 0 & 0 \\ 0 & \tilde{z}_{\alpha_j} & 0 \\ 0 & 0 & \tilde{z}_{j,b} \end{pmatrix}$$

with sizes $r, 2, s$ where

- $\tilde{z}_{j,t} = t_{e(n-1, \dots, s+2)}$
- $\tilde{z}_{j,b} = t_{e(s-1, \dots, 0)}$.
- $v^{-es} \tilde{z}_{\alpha_j}$ has elementary divisors bounded by $v^{(e,0)}$.

In particular, we are in a position to repeatedly apply Proposition 4.2.4 to $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}}$ and each $A_{\overline{\mathfrak{M}}}^{(j)}$ for the universal Breuil–Kisin module $\overline{\mathfrak{M}}$. This gives

$$A_{\overline{\mathfrak{M}}}^{(j)} = D^{(j)} w_j^{-1} (P^{(j)}) w_j$$

where $P^{(j)}$ is block lower triangular, whose Levi blocks from top to bottom are $E_j^{n-1}, \dots, E_j^{s+2}, M_{\alpha_j}, E_j^{s-1}, \dots, 1$.

Furthermore, the entries of $D^{(j)}, P^{(j)}$ over all j topologically generate $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}}$. By Proposition 4.2.5 and Corollary 4.3.2, the data M_{α_j} identifies with the universal deformation of the point $x_j \in M_j(t_{(1,0)})(\mathbb{F})$ given by its reduction \overline{M}_{α_j} modulo the maximal ideal. Set $M_{\mathcal{J}}(t_{(1,0)}) = \prod_j M_j(t_{(1,0)})$, which contains $x = (x_j)$ as an \mathbb{F} -point. Thus $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}}$ acquires an $\widehat{\otimes}_{\mathcal{O}} \mathcal{O}_{M_{\mathcal{J}}(1,0), x}^{\wedge} = \mathcal{O}_{M_{\mathcal{J}}(t_{(1,0)}, x)}^{\wedge}$ -algebra structure, whose image coincide with the topological subalgebra generated by the coefficients of the entries of (M_{α_j}) for all possible j .

Repeated applications of the approximate version of Lemma 4.4.3 to $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla}$ as in Remark 4.4.5 (with the control of the monodromy condition obtained by combining Proposition 4.4.1 and Lemma 4.4.2) show that $R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla}$ is topologically generated over $\mathcal{O}_{M_{\mathcal{J}}(1,0), x}^{\wedge}$ by $fn + e \sum_j \dim N_{-\alpha_j} = fn + (\frac{n(n-1)}{2} - 1)[K : \mathbb{Q}_p]$ elements. But since we assumed $R^{\leq \eta, \tau} \neq 0$, $\dim R_{\overline{\mathfrak{M}}}^{\tau, \bar{\beta}, \nabla} = 1 + fn + \frac{n(n-1)}{2}[K : \mathbb{Q}_p] = \dim \mathcal{O}_{M_{\mathcal{J}}(1,0), x}^{\wedge} + fn + (\frac{n(n-1)}{2} - 1)[K : \mathbb{Q}_p]$. Since $\mathcal{O}_{M_{\mathcal{J}}(1,0), x}^{\wedge}$ is an integral domain (being the

completion of an excellent normal scheme), the equality of dimension can only happen if $R_{\overline{\mathfrak{M}}}^{\tau, \overline{\beta}, \nabla}$ is a power series ring over $\mathcal{O}_{M_{\mathcal{J}}(1,0),x}^{\wedge}$ in the correct number of variables. All the assertions of Theorem 4.1.1 now follows from properties of the $M_{\mathcal{J}}(t_{(1,0)})$ which follows from Corollary 4.3.2. \square

5. THE MAIN RESULTS

In this section, we prove our main results on the weight part of Serre's conjecture. We start with an axiomatic setup before defining the relevant spaces of automorphic forms in §5.5.

Recall from §2.3.6 that given an \mathbb{F} -valued L -homomorphism $\bar{\rho} : G_{\mathbb{Q}_p} \rightarrow {}^L G(\mathbb{F})$ (resp. a tame inertial L -parameter $\tau : I_{\mathbb{Q}_p} \rightarrow \underline{G}^\vee(E)$) we have a corresponding collection $(\bar{\rho}_v)_{v \in S_p}$ of continuous Galois representations $\bar{\rho}_v : G_{F_v^+} \rightarrow \mathrm{GL}_n(\mathbb{F})$ (resp. a corresponding collection $(\tau_v)_{v \in S_p}$ of tame inertial types $\tau_v : I_{F_v^+} \rightarrow \mathrm{GL}_n(E)$).

5.1. Weight elimination.

Theorem 5.1.1. *Let $\bar{\rho} : G_{\mathbb{Q}_p} \rightarrow {}^L G(\mathbb{F})$ be a $3e(n-1)$ -generic \mathbb{F} -valued L -homomorphism. Let $\bar{\rho}^{\mathrm{sp}}$ be a specialization of $\bar{\rho}$ with a compatible $\max\{2, e\}(n-1)$ -generic lowest alcove presentation. Assume that we have a set $W_{\mathrm{elim}}(\bar{\rho})$ of $3(n-1)$ -generic Serre weights satisfying the following local-global compatibility axiom:*

- (i) *for any tame inertial L -parameter τ , $\mathrm{JH}(\bar{\sigma}(\tau)) \cap W_{\mathrm{elim}}(\bar{\rho}) \neq \emptyset$ implies that $\bar{\rho}$ has a potentially crystalline lift of type (τ, η) .*

Then $W_{\mathrm{elim}}(\bar{\rho}) \subset W^?(\bar{\rho}^{\mathrm{sp}})$.

Proof. Suppose that $F(\lambda) \in W_{\mathrm{elim}}(\bar{\rho})$. Choose the tame inertial L -parameter τ with $F(\lambda) \in \mathrm{JH}(\bar{\sigma}(\tau))$ constructed in Proposition 2.4.5. By Theorem 3.5.1, $\tilde{w}(\bar{\rho}^{\mathrm{sp}}, \tau) \in \mathrm{Adm}(e\eta_0)$, and we conclude by Proposition 2.4.5. \square

Remark 5.1.2. If $e \geq 2$ the hypothesis on $\bar{\rho}^{\mathrm{sp}}$ follows from the hypothesis on $\bar{\rho}$.

5.2. **Patching functors.** We recall weak patching functors. Let

$$R_{\bar{\rho}} \stackrel{\mathrm{def}}{=} \widehat{\bigotimes}_{v \in S_p, \mathcal{O}} R_{\bar{\rho}_v}^{\square},$$

and let R^p be a nonzero complete local Noetherian equidimensional flat \mathcal{O} -algebra with residue field \mathbb{F} such that each irreducible component of $\mathrm{Spec} R^p$ and of $\mathrm{Spec} \bar{R}^p$ is geometrically irreducible. (The latter hypothesis can be guaranteed after passing to a finite extension of the coefficient field E .) We let $R_{\infty} \stackrel{\mathrm{def}}{=} R_{\bar{\rho}} \widehat{\otimes}_{\mathcal{O}} R^p$ and suppress the dependence on R^p below. We let $R_{\bar{\rho}}^{\eta, \tau}$ be

$$\widehat{\bigotimes}_{v \in S_p, \mathcal{O}} R_{\bar{\rho}_v}^{\eta_v, \tau_v}$$

and define $R_{\infty}(\tau) \stackrel{\mathrm{def}}{=} R_{\infty} \otimes_{R_{\bar{\rho}}} R_{\bar{\rho}}^{\eta, \tau}$. We write X_{∞} , $X_{\infty}(\tau)$, and $\bar{X}_{\infty}(\tau)$ for $\mathrm{Spec} R_{\infty}$, $\mathrm{Spec} R_{\infty}(\tau)$, and $\mathrm{Spec} \bar{R}_{\infty}(\tau)$ respectively, denote by $\mathrm{Mod}(X_{\infty})$ the category of coherent sheaves over X_{∞} , and let $\mathrm{Rep}_{\mathcal{O}}(\mathrm{GL}_n(\mathcal{O}_p))$ be the category of topological $\mathcal{O}[\mathrm{GL}_n(\mathcal{O}_p)]$ -modules which are finitely generated over \mathcal{O} . We say that an \bar{E} -point of $\mathrm{Spec} R_{\bar{\rho}}$ is *potentially diagonalizable* if for each $v \in S_p$, the corresponding Galois representation $G_{F_v^+} \rightarrow \mathrm{GL}_n(\bar{E})$ is potentially diagonalizable in the sense of [BLGGT14, §1.4]. We say that an \bar{E} -point of X_{∞} is potentially diagonalizable if its image in $\mathrm{Spec} R_{\bar{\rho}}$ is.

Definition 5.2.1. A *weak patching functor* for an L -homomorphism $\bar{\rho} : G_{\mathbb{Q}_p} \rightarrow {}^L G(\mathbb{F})$ is a nonzero covariant exact functor $M_{\infty} : \mathrm{Rep}_{\mathcal{O}}(\mathrm{GL}_n(\mathcal{O}_p)) \rightarrow \mathrm{Mod}(X_{\infty})$ satisfying the following: if τ is an inertial L -parameter and $\sigma^{\circ}(\tau)$ is an \mathcal{O} -lattice in $\sigma(\tau)$ then

- (1) $M_{\infty}(\sigma^{\circ}(\tau))$ is a maximal Cohen–Macaulay sheaf on $X_{\infty}(\tau)$;
- (2) for all $\sigma \in \mathrm{JH}(\bar{\sigma}^{\circ}(\tau))$, $M_{\infty}(\sigma)$ is a maximal Cohen–Macaulay sheaf on $\bar{X}_{\infty}(\tau)$ (or is 0); and

- (3) if there is an inertial L -parameter τ_0 such that $\text{Supp}M_\infty(\sigma(\tau_0)^\circ)$ contains a potentially diagonalizable \overline{E} -point, then for any inertial L -parameter τ , $\text{Supp}M_\infty(\sigma(\tau)^\circ)$ contains all potentially diagonalizable \overline{E} -points.

We say that a weak patching functor M_∞ is *minimal* if R^p is formally smooth over \mathcal{O} and whenever τ is an inertial L -parameter, $M_\infty(\sigma^\circ(\tau))[p^{-1}]$, which is locally free over (the regular scheme) $\text{Spec } R_\infty(\tau)[p^{-1}]$, has rank at most one on each connected component.

Definition 5.2.2. We say that a weak patching functor M_∞ is *potentially diagonalizable* if there exists τ_0 as in Definition 5.2.1(3).

5.3. Cycles from patching functors. We recall some notation from [EG14, §2.2]. Let \mathcal{X} be an equidimensional Noetherian scheme of dimension d . Let $\mathcal{Z}(\mathcal{X})$ be the free abelian group generated by integral subschemes of \mathcal{X} of *maximal dimension* d . If \mathcal{M} is a coherent sheaf on \mathcal{X} with finite-dimensional support, then we can define $Z(\mathcal{M}) \in \mathcal{Z}(\mathcal{X})$ to be $Z_d(\mathcal{M})$ which is defined as in *loc. cit.*

Now suppose that \mathcal{X} is a p -flat equidimensional Noetherian scheme over \mathcal{O} . Then $\mathcal{X}[p^{-1}] \stackrel{\text{def}}{=} \mathcal{X} \otimes_{\mathcal{O}} E$ and $\overline{\mathcal{X}} \stackrel{\text{def}}{=} \mathcal{X} \otimes_{\mathcal{O}} \mathbb{F}$ are equidimensional Noetherian schemes, and there is a natural reduction map $\text{red} : \mathcal{Z}(\mathcal{X}[p^{-1}]) \rightarrow \mathcal{Z}(\overline{\mathcal{X}})$. Moreover, if we let $\mathcal{M}[p^{-1}] \stackrel{\text{def}}{=} \mathcal{M} \otimes_{\mathcal{O}} E$ and $\overline{\mathcal{M}} \stackrel{\text{def}}{=} \mathcal{M} \otimes_{\mathcal{O}} \mathbb{F}$ be the corresponding sheaves on $\mathcal{X}[p^{-1}]$ and $\overline{\mathcal{X}}$, respectively, we have the following fact.

Proposition 5.3.1. *If \mathcal{M} is an \mathcal{O} -flat coherent sheaf over \mathcal{X} with finite-dimensional support, then $\text{red}(Z(\mathcal{M}[p^{-1}])) = Z(\overline{\mathcal{M}})$.*

We introduce notation for completed products of cycles. Suppose that R and S are equidimensional complete local Noetherian flat \mathcal{O} -algebras. If \overline{Z}_1 and \overline{Z}_2 are geometrically integral subschemes of $\text{Spec } R \otimes_{\mathcal{O}} \mathbb{F}$ and $\text{Spec } S \otimes_{\mathcal{O}} \mathbb{F}$ corresponding to prime ideals $\overline{\mathfrak{p}}$ and $\overline{\mathfrak{q}}$, respectively, then we denote by $\overline{Z}_1 \times \overline{Z}_2$ the subscheme

$$\text{Spec } (R \otimes_{\mathcal{O}} \mathbb{F}) / \overline{\mathfrak{p}} \widehat{\otimes}_{\mathbb{F}} (S \otimes_{\mathcal{O}} \mathbb{F}) / \overline{\mathfrak{q}} \subset \text{Spec } (R \otimes_{\mathcal{O}} \mathbb{F}) \widehat{\otimes}_{\mathbb{F}} (S \otimes_{\mathcal{O}} \mathbb{F})$$

which is geometrically integral by [BGHT11, Lemma 3.3(4)]. Similarly, if Z_1 and Z_2 are geometrically integral subschemes of $\text{Spec } R[p^{-1}]$ and $\text{Spec } S[p^{-1}]$ corresponding to prime ideals \mathfrak{p} and \mathfrak{q} , then we denote by $Z_1 \times Z_2$ the subscheme

$$\text{Spec } (R / (\mathfrak{p} \cap R) \widehat{\otimes}_{\mathcal{O}} S / (\mathfrak{q} \cap S)) [p^{-1}] \subset \text{Spec } R \widehat{\otimes}_{\mathcal{O}} S [p^{-1}]$$

which is geometrically integral by [BGHT11, Lemma 3.3(3)].

We now specialize to some schemes in our patching axioms. Let $\overline{\rho}$ be an L -homomorphism over \mathbb{F} . Fix a finite set \mathcal{T} of inertial L -parameters such that:

- (ii) for all $\tau \in \mathcal{T}$ the irreducible components of $\text{Spec } R_{\overline{\rho}}^\tau$ and $\text{Spec } \overline{R}_{\overline{\rho}}^\tau$ are geometrically integral.

Let $R_{\overline{\rho}}^{\mathcal{T}}$ be the reduced union $\cup_{\tau \in \mathcal{T}} \text{Spec } R_{\overline{\rho}}^\tau$. Let M_∞ be a weak patching functor for $\overline{\rho}$. We write $R_\infty(\mathcal{T})$ for $R_\infty \widehat{\otimes}_{R_{\overline{\rho}}^{\mathcal{T}}} R_{\overline{\rho}}^{\mathcal{T}} \cong R^p \widehat{\otimes}_{\mathcal{O}} R_{\overline{\rho}}^{\mathcal{T}}$ and $X_\infty(\mathcal{T})$ for $\text{Spec } R_\infty(\mathcal{T})$. Recall that by assumption, the irreducible components of $\text{Spec } R^p[p^{-1}]$ and $\text{Spec } \overline{R}^p$ are geometrically irreducible. Every irreducible cycle $Z \in \mathcal{Z}(\mathcal{X}_\infty(\mathcal{T})[p^{-1}])$ is of the form $Z^p \times Z_p$ for geometrically irreducible cycles $Z^p \in \mathcal{Z}(\text{Spec } R^p[p^{-1}])$ and $Z_p \in \mathcal{Z}(\text{Spec } R_{\overline{\rho}}^{\mathcal{T}})$ by [BGHT11, Lemma 3.3(5)]. Similarly, every irreducible cycle $\overline{Z} \in \mathcal{Z}(\overline{\mathcal{X}}_\infty(\mathcal{T}))$ is of the form $\overline{Z}^p \times \overline{Z}_p$ for geometrically irreducible cycles $\overline{Z}^p \in \mathcal{Z}(\text{Spec } \overline{R}^p)$ and $\overline{Z}_p \in \mathcal{Z}(\text{Spec } \overline{R}_{\overline{\rho}}^{\mathcal{T}})$ by [BGHT11, Lemma 3.3(6)].

Let $e : \mathcal{Z}(\mathrm{Spec} \overline{R}^p) \rightarrow \mathbb{Z}$ be the homomorphism that sends the cycle of an integral subscheme to 1. We define the maps

$$\begin{aligned} \mathrm{pr} : \mathcal{Z}(X_\infty(\mathcal{T})[p^{-1}]) &\rightarrow \mathcal{Z}(\mathrm{Spec} R_\rho^{\mathcal{T}}[p^{-1}]) \\ Z^p \times Z_p &\mapsto e(\mathrm{red}(Z^p))Z_p \end{aligned}$$

and

$$\begin{aligned} \overline{\mathrm{pr}} : \mathcal{Z}(\overline{X}_\infty(T)) &\rightarrow \mathcal{Z}(\mathrm{Spec} \overline{R}_\rho^{\mathcal{T}}) \\ \overline{Z}^p \times \overline{Z}_p &\mapsto \overline{Z}_p. \end{aligned}$$

We have that $\mathrm{red} \circ \mathrm{pr} = \overline{\mathrm{pr}} \circ \mathrm{red} : \mathcal{Z}(X_\infty(\mathcal{T})[p^{-1}]) \rightarrow \mathcal{Z}(\mathrm{Spec} \overline{R}_\rho^{\mathcal{T}})$ (using that $\mathrm{red}(Z^p \times Z_p) = \mathrm{red}(Z^p) \times \mathrm{red}(Z_p)$), from which we immediately obtain the following corollary.

Corollary 5.3.2. *If $\tau \in T$, then the image of the composition*

$$\mathcal{Z}(X_\infty(\tau)[p^{-1}]) \xrightarrow{\mathrm{red}} \mathcal{Z}(\overline{X}_\infty(\tau)) \xrightarrow{\overline{\mathrm{pr}}} \mathcal{Z}(\mathrm{Spec} \overline{R}_\rho^{\mathcal{T}})$$

is contained in $\mathrm{red}(\mathcal{Z}(\mathrm{Spec} R_\rho^{\mathcal{T}}[p^{-1}]))$.

Depending on context, we denote either $\mathrm{pr} \circ Z$ or $\overline{\mathrm{pr}} \circ Z$ by Z_p .

5.4. Weight elimination and modularity of extremal weights. Let $\overline{\rho}$ be a 1-generic L -homomorphism, with a lowest alcove presentation for it. Fix a weak patching functor M_∞ for $\overline{\rho}$. Let $W_{M_\infty}(\overline{\rho})$ be the set of $3(n-1)$ -generic Serre weights σ such that $M_\infty(\sigma)$ is nonzero.

Proposition 5.4.1. *The set $W_{M_\infty}(\overline{\rho})$ satisfies the condition (i) for $\overline{\rho}$.*

Proof. Suppose that $\sigma \in \mathrm{JH}(\overline{\sigma}(\tau)) \cap W_{M_\infty}(\overline{\rho})$ for a generic tame inertial L -parameter τ . Then $M_\infty(\sigma)$ is nonzero so that $M_\infty(\sigma(\tau)^\circ)$ is nonzero for any lattice $\sigma(\tau)^\circ \subset \sigma(\tau)$ by exactness. Definition 5.2.1(1) implies that $R_\infty(\tau)$, and so $R_\rho^{\mathcal{T}}$, is nonzero. \square

Definition 5.4.2. We say that a weak patching functor M_∞ for $\overline{\rho}$ is *extremal* if $W_{\mathrm{extr}}(\overline{\rho}) \cap W_{M_\infty}(\overline{\rho})$ is nonempty.

Theorem 5.4.3. *Let $\overline{\rho}$ be $6e(n-1)$ -generic. If a weak patching functor M_∞ is extremal, then $W_{\mathrm{extr}}(\overline{\rho}) \subset W_{M_\infty}(\overline{\rho})$, and moreover, the map $\theta_{\overline{\rho}} : SP(\overline{\rho}) \rightarrow W^{\mathcal{J}}$ is a bijection.*

The proof of Theorem 5.4.3 requires the following two results.

Lemma 5.4.4. *Assume that $\overline{\rho}$ is $(2e + \max\{2, e\})(n-1)$ -generic. Suppose that M_∞ is a weak patching functor for $\overline{\rho}$, $(\sigma, \overline{\rho}^{\mathrm{SP}}) \in SP(\overline{\rho})$, and $\sigma \in W_{M_\infty}(\overline{\rho})$. Assume that $\overline{\rho}^{\mathrm{SP}}$ has a compatible $4e(n-1)$ -generic lowest alcove presentation. Suppose that σ is the extremal weight of $\overline{\rho}^{\mathrm{SP}}$ corresponding to $w \in W$. Let α be a simple root.*

Using $\overline{\rho}^{\mathrm{SP}}$, w , and α , we define as in Proposition 2.4.9 (using the above $4e(n-1)$ -generic lowest alcove presentation) σ_m and τ_m for $0 \leq m \leq 2e-1$ and $0 \leq m \leq 2e$, respectively, so that $\sigma_0 = \sigma$. Then there exists $0 \leq k \leq 2e-1$ such that $\sigma_m \in W_{M_\infty}(\overline{\rho})$ if and only if $m \leq k$. Moreover, τ_{k+1} exhibits a specialization of $\overline{\rho}$ to $(\sigma_k, \overline{\rho}'^{\mathrm{SP}})$ for some \mathbb{F} -valued tame inertial L -parameter $\overline{\rho}'^{\mathrm{SP}}$.

Proof. Fix \mathcal{O} -lattices $\sigma(\tau_m)^\circ \subset \sigma(\tau_m)$ for $0 \leq m \leq 2e$ (the choices will not affect the argument below). Let $0 \leq k \leq 2e-1$ be such that $\sigma_m \in W_{M_\infty}(\overline{\rho})$ for $0 \leq m \leq k$ and either $\sigma_{k+1} \notin W_{M_\infty}(\overline{\rho})$ or $k = 2e-1$. That $\sigma_m \in W_{M_\infty}(\overline{\rho})$ implies that $M_\infty(\sigma(\tau_{m+1})^\circ)$ is nonzero. Therefore $R_\rho^{\mathcal{T}_{m+1}}$ is nonzero for $0 \leq m \leq k$. We will first show that $\tilde{w}(\overline{\rho}, \tau_{m+1}) \in w^{-1}t_{e\eta_0}W_{a,\alpha}w \cap \mathrm{Adm}^\vee(e\eta_0)$ for $0 \leq m \leq k$.

Fix m with $0 \leq m \leq k$. Suppose that τ_{m+1} exhibits the specialization to the \mathbb{F} -valued tame inertial L -parameter $\bar{\rho}'^{\text{SP}}$ i.e. that $\tilde{w}(\bar{\rho}, \tau_{m+1}) = \tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_{m+1})$. Since $\sigma_m \in W_{M_\infty}(\bar{\rho}) \subset W^?(\bar{\rho}'^{\text{SP}})$ by Theorem 5.1.1 and Proposition 5.4.1, Proposition 2.4.6 implies that $\tilde{w}_h \tilde{s} \tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_{m+1}) \leq w_0 t_{(e-1)\eta_0} \tilde{s}$ for $s \in \{w, s_\alpha w\}$. (Note that Theorem 5.1.1 applies to $\bar{\rho}$ and $\bar{\rho}'^{\text{SP}}$, by the genericity assumption on $\bar{\rho}$.) This implies that $\tilde{w}(\bar{\rho}, \tau_{m+1}) = \tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_{m+1}) \leq t_{s^{-1}(e\eta_0)}$. Combining this with the fact that $\tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_{m+1}) \leq \tilde{w}(\bar{\rho}, \tau_{m+1})$ by Theorem 3.5.1, we have that $\tilde{w}(\bar{\rho}, \tau_{m+1}) \in w^{-1} t_{e\eta_0} W_{a,\alpha} w \cap \text{Adm}^\vee(e\eta_0)$ by Proposition 2.2.9.

Now Proposition 2.4.9 applied to $\bar{\rho}^{\text{SP}}$, Theorem 5.1.1, and Proposition 5.4.1 imply that

$$W_{M_\infty}(\bar{\rho}) \cap \text{JH}(\bar{\sigma}(\tau_{m+1})^\circ) \subset W^?(\bar{\rho}^{\text{SP}}) \cap \text{JH}(\bar{\sigma}(\tau_{m+1})^\circ) = \{\sigma_m, \sigma_{m+1}\}$$

(or $\{\sigma_m\}$ if $m = 2e - 1$). We now use notation from §5.3 with $\mathcal{T} = \{\tau_m \mid 0 \leq m \leq 2e - 1\}$. The set of types \mathcal{T} satisfies condition (ii) by the genericity assumption on $\bar{\rho}^{\text{SP}}$ and Theorem 4.1.1. We continue to fix m with $0 \leq m \leq k$. Since σ_m and σ_{m+1} appear as Jordan–Hölder factors of $\bar{\sigma}(\tau_{m+1})^\circ$ with multiplicity one, exactness of M_∞ gives

$$(5.1) \quad Z_{\mathfrak{p}}(M_\infty(\bar{\sigma}(\tau_{m+1})^\circ)) = Z_{\mathfrak{p}}(M_\infty(\sigma_m)) + Z_{\mathfrak{p}}(M_\infty(\sigma_{m+1})),$$

for $0 \leq m \leq k - 1$ and $Z_{\mathfrak{p}}(M_\infty(\bar{\sigma}(\tau_{k+1})^\circ)) = Z_{\mathfrak{p}}(M_\infty(\sigma_k))$. We will use (5.1) and the previous paragraph to show that $\tilde{w}(\bar{\rho}, \tau_{k+1})$ is $t_{s^{-1}(\eta_0)}$ for some $s \in \{w, s_\alpha w\}$.

Let us call a cycle *balanced* if it is a multiple of the sum of two distinct integral subschemes and *unbalanced* if it is supported on at most two integral subschemes with distinct multiplicities. In particular, an unbalanced cycle is nonzero. For $0 \leq m \leq k - 1$, $\tilde{w}(\bar{\rho}, \tau_{m+1}) \notin \{t_{w^{-1}(\eta_0)}, t_{(s_\alpha w)^{-1}(\eta_0)}\}$ since otherwise

$$2 = \#W_{M_\infty}(\bar{\rho}) \cap \text{JH}(\bar{\sigma}(\tau_{m+1})^\circ) \leq \#W^?(\bar{\rho}'^{\text{SP}}) \cap \text{JH}(\bar{\sigma}(\tau_{m+1})^\circ) = 1$$

by Proposition 5.4.1. Then $Z_{\mathfrak{p}}(M_\infty(\bar{\sigma}(\tau_{m+1})^\circ))$ is balanced by Corollary 5.3.2 since $R_{\bar{\rho}}^{\tau_{m+1}}[p^{-1}]$ is geometrically irreducible and $Z(\text{Spec } \bar{R}_{\bar{\rho}}^{\tau_{m+1}})$ is balanced (see Theorem 4.1.1). By (5.1), $Z_{\mathfrak{p}}(M_\infty(\sigma_m))$ is balanced (resp. unbalanced) if and only if $Z_{\mathfrak{p}}(M_\infty(\sigma_{m+1}))$ is balanced (resp. unbalanced) for $0 \leq m \leq k - 1$. Since $Z_{\mathfrak{p}}(M_\infty(\sigma_0)) = Z_{\mathfrak{p}}(M_\infty(\bar{\sigma}(\tau_0)^\circ))$ is unbalanced as $R_{\bar{\rho}}^{\tau_0}$ is formally smooth over \mathcal{O} , we conclude that $Z_{\mathfrak{p}}(M_\infty(\sigma_k)) = Z_{\mathfrak{p}}(M_\infty(\bar{\sigma}(\tau_{k+1})^\circ))$ is unbalanced. We conclude from the argument above that $\tilde{w}(\bar{\rho}, \tau_{k+1})$ is $t_{s^{-1}(\eta_0)}$ for some $s \in \{w, s_\alpha w\}$. In particular, τ_{k+1} exhibits a specialization of $\bar{\rho}$ to $(\sigma_k, \bar{\rho}'^{\text{SP}})$ (not necessarily the same $\bar{\rho}'^{\text{SP}}$ from the first paragraph).

By the definition of σ_k and using that σ_k is an extremal weight of $\bar{\rho}'^{\text{SP}}$, we see that

$$\tilde{w}(\bar{\rho}, \tau_{k+1}) = \tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_{k+1}) = \begin{cases} t_{w^{-1}(e\eta_0)} & \text{if } k \text{ is even} \\ t_{(s_\alpha w)^{-1}(e\eta_0)} & \text{if } k \text{ is odd.} \end{cases}$$

Then a computation shows that

$$\tilde{w}(\bar{\rho}'^{\text{SP}}) = \begin{cases} \tilde{w}(\bar{\rho}^{\text{SP}}) \tilde{w}^{-1} t_{(\frac{k}{2}-e)\alpha} s_\alpha \tilde{w} & \text{if } k \text{ is even} \\ \tilde{w}(\bar{\rho}^{\text{SP}}) \tilde{w}^{-1} t_{(\frac{k+1}{2}-e)\alpha} \tilde{w} & \text{if } k \text{ is odd.} \end{cases}$$

Note that $\tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_m) \in w^{-1} W_{a,\alpha} t_{e\eta_0} w$ for all $0 \leq m \leq 2e$. Another computation shows that if $m > k + 1$, then $\tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_m)$ is not listed in Proposition 2.2.6. This implies that $\tilde{w}(\bar{\rho}'^{\text{SP}}, \tau_m) \notin \text{Adm}(e\eta_0)$ for $m > k + 1$. Corollary 2.4.7 implies that $W^?(\bar{\rho}'^{\text{SP}}, \tau_m) = \emptyset$ for $m > k + 1$. In particular, $\sigma_m \notin W^?(\bar{\rho}'^{\text{SP}})$ for $m > k$. Theorem 5.1.1 and Proposition 5.4.1 imply that $\sigma_m \notin W_{M_\infty}(\bar{\rho})$ for $m > k$. \square

Corollary 5.4.5. *Let $(\sigma, \bar{\rho}^{\text{SP}}) \in SP(\bar{\rho})$ and $\sigma \in W_{M_\infty}(\bar{\rho})$ be as in Lemma 5.4.4. Let α be a simple root. Then there exists $(\sigma', \bar{\rho}'^{\text{SP}})$ such that $\theta_{\bar{\rho}}(\sigma', \bar{\rho}'^{\text{SP}}) = \theta_{\bar{\rho}}(\sigma, \bar{\rho}^{\text{SP}})s_\alpha$. Moreover, if $\sigma \in W_{M_\infty}(\bar{\rho})$, then $\sigma' \in W_{M_\infty}(\bar{\rho})$ as well.*

Proof. Let $\sigma, \sigma_k, \bar{\rho}^{\text{SP}}$, and $\bar{\rho}'^{\text{SP}}$ be as in Lemma 5.4.4. Let σ' be σ_k . Then $\sigma' \in W_{M_\infty}(\bar{\rho})$. It suffices to show that $\theta_{\bar{\rho}}(\sigma', \bar{\rho}'^{\text{SP}}) = \theta_{\bar{\rho}}(\sigma, \bar{\rho}^{\text{SP}})s_\alpha$. We have that

$$w(\bar{\rho}'^{\text{SP}}) = \begin{cases} w(\bar{\rho}^{\text{SP}})w^{-1}s_\alpha w & \text{if } k \text{ is even} \\ w(\bar{\rho}^{\text{SP}}) & \text{if } k \text{ is odd} \end{cases}$$

and σ_k is the extremal weight of $\bar{\rho}'^{\text{SP}}$ corresponding to

$$\begin{cases} w & \text{if } k \text{ is even.} \\ s_\alpha w & \text{if } k \text{ is odd.} \end{cases}$$

We conclude that $\theta_{\bar{\rho}}(\sigma', \bar{\rho}'^{\text{SP}}) = w(\bar{\rho})w^{-1}s_\alpha = \theta_{\bar{\rho}}(\sigma, \bar{\rho}^{\text{SP}})s_\alpha$. \square

Proof of Theorem 5.4.3. Suppose that $\sigma \in W_{M_\infty}(\bar{\rho})$ and that $(\sigma, \bar{\rho}^{\text{SP}}) \in SP(\bar{\rho})$. Then using Corollary 5.4.5 and the fact that simple reflections generate W , we see that for each $w \in W$, there is $(\sigma_w, \bar{\rho}_w^{\text{SP}}) \in SP(\bar{\rho})$ such that $\theta_{\bar{\rho}}(\sigma_w, \bar{\rho}_w^{\text{SP}}) = w$ and $\sigma_w \in W_{M_\infty}(\bar{\rho})$. This first implies that the map $\theta_{\bar{\rho}}$ is surjective and hence an isomorphism by Proposition 3.6.4. It also implies that $W_{\text{extr}}(\bar{\rho}) \subset W_{M_\infty}(\bar{\rho})$. \square

Theorem 5.4.6. *Let $\bar{\rho}$ be $6e(n-1)$ -generic and let M_∞ be a weak patching functor for $\bar{\rho}$. The following are equivalent.*

- (1) M_∞ is extremal.
- (2) $W_{\text{extr}}(\bar{\rho}) \subset W_{M_\infty}(\bar{\rho})$.
- (3) M_∞ is potentially diagonalizable.

Proof. (1) implies (2) by Theorem 5.4.3. We next show that (2) implies (3). Let σ be in $W_{\text{mord}}(\bar{\rho})$ so that τ exhibits the specialization pair $(\sigma, \bar{\rho}^{\text{SS}}) \in SP(\bar{\rho})$ as in the proof of Proposition 3.8.3. Then $M_\infty(\sigma(\tau)^\circ)$ is nonzero since $M_\infty(\sigma)$ is. Since $R_{\bar{\rho}}^T$ is a domain and $\bar{\rho}$ has a potentially diagonalizable lift of type (τ, η) by Theorem 3.8.2, M_∞ is potentially diagonalizable.

Finally, we show that (3) implies (1). Again, let σ be in $W_{\text{mord}}(\bar{\rho})$ so that τ exhibits the specialization pair $(\sigma, \bar{\rho}^{\text{SS}}) \in SP(\bar{\rho})$ as in the proof of Proposition 3.8.3. Then since M_∞ is potentially diagonalizable and $\bar{\rho}$ has a potentially diagonalizable lift of type (τ, η) as before, $M_\infty(\sigma(\tau)^\circ)$ is nonzero. Since $W_{M_\infty}(\bar{\rho}) \subset W^?(\bar{\rho}^{\text{SS}})$ by Theorem 5.1.1 and $W^?(\bar{\rho}^{\text{SS}}, \tau) = \{\sigma\}$ as in the proof of Proposition 3.8.3, $M_\infty(\sigma)$ is nonzero. Thus $W_{\text{mord}}(\bar{\rho}) \cap W_{M_\infty}(\bar{\rho})$ is nonempty. The result now follows from Proposition 3.8.3. \square

Remark 5.4.7. Theorem 5.4.6 generalizes [LLHL19, Theorem 4.3.8] to the nonsemisimple case in an abstract setting. Moreover, the above proof (and §5.5) gives a different proof of this theorem. (Specifically, the order of implications proved is reversed.) Indeed, we do not know whether every extremal lift is potentially diagonalizable when $\bar{\rho}$ is wildly ramified.

Corollary 5.4.8. *Suppose that \mathcal{O}_p is étale over \mathbb{Z}_p , i.e., F_p^+ is a product of unramified extensions of \mathbb{Q}_p . Let $\bar{\rho}$ be an L -homomorphism over \mathbb{F} . Suppose that M_∞ is a weak patching functor for $\bar{\rho}$ satisfying the equivalent conditions of Theorem 5.4.6. (In particular, $\bar{\rho}$ is $7(n-1)$ -generic.) If τ is an n -generic tame inertial L -parameter, then $R_{\bar{\rho}}^T$ is nonzero if and only if $M_\infty(\sigma^\circ(\tau))$ is nonzero for any \mathcal{O} -lattice $\sigma^\circ(\tau) \subset \sigma(\tau)$.*

Proof. If $M_\infty(\sigma^\circ(\tau))$ is nonzero, then $R_\infty(\tau)$, and thus R_ρ^τ , is nonzero. Conversely, if R_ρ^τ is nonzero, then $W_{\text{extr}}(\bar{\rho}) \cap \text{JH}(\bar{\sigma}(\tau)) \neq \emptyset$ by Proposition 3.10.3. Theorem 5.4.6(2) and exactness of M_∞ imply that $M_\infty(\sigma^\circ(\tau))$ is nonzero. \square

The following freeness result follows from our previous results and the Diamond–Fujiwara trick.

Theorem 5.4.9. *Let M_∞ be a minimal weak patching functor for $\bar{\rho}$. Suppose that the equivalent conditions of Theorem 5.4.6 hold for M_∞ and that $\sigma \in W_{\text{extr}}(\bar{\rho})$. Then $M_\infty(\sigma)$ is free of rank 1 over its support (which is formally smooth over \mathbb{F}).*

Proof. There exists a generic tame inertial L -parameter τ which exhibits the specialization $(\sigma, \bar{\rho}^{\text{SP}}) \in SP(\bar{\rho})$ for some \mathbb{F} -valued inertial L -parameter $\bar{\rho}^{\text{SP}}$. By Lemma 3.6.1, we can assume without loss of generality that $\tilde{w}(\bar{\rho}, \tau) = t_{w^{-1}(e\eta_0)}$ for some $w \in W$. By Theorem 4.1.1, R_ρ^τ is formally smooth over \mathcal{O} , so that $R_\infty(\tau)$ is as well. Since for any \mathcal{O} -lattice $\sigma^\circ(\tau) \subset \sigma(\tau)$, $M_\infty(\sigma^\circ(\tau))$ is nonzero, finitely generated, and maximally Cohen–Macaulay over $R_\infty(\tau)$, it must be free over $\bar{R}_\infty(\tau)$ by Serre’s theorem on finiteness of projective dimension and the Auslander–Buchsbaum formula. Since the generic rank is at most 1, its rank must be 1. \square

5.5. Global results. In this section, we discuss algebraic automorphic forms on certain definite unitary groups to which the Taylor–Wiles patching construction can be applied to obtain patching functors as in §5.2. This gives a context to which results in the previous section can be applied.

5.5.1. Algebraic automorphic forms on some definite unitary groups. Let F^+/\mathbb{Q} be a totally real field not equal to \mathbb{Q} , and let $F \subset \bar{F}^+$ be a CM extension of F^+ . We say that a finite place of F^+ is *split* (resp. ramified or inert) if it splits (resp. ramifies or is inert) in F . We say that a place of F is *split* (resp. ramified or inert) if its restriction to F^+ is split (resp. ramified or inert) in F .

Let $G_{/F^+}$ be a reductive group which is an outer form of GL_n such that

- $G_{/F}$ is an inner form of GL_n ;
- $G_{/F^+}(F_v^+) \cong U_n(\mathbb{R})$ for all $v|\infty$; and
- $G_{/F^+}$ is quasisplit at all inert and ramified finite places.

By [EGH13, §7.1], G admits a reductive model \mathcal{G} over $\mathcal{O}_{F^+}[1/N]$, for some $N \in \mathbb{N}$, and an isomorphism

$$(5.2) \quad \iota : \mathcal{G}_{/\mathcal{O}_F[1/N]} \xrightarrow{\iota} \text{GL}_n/\mathcal{O}_F[1/N]$$

which specializes to $\iota_w : \mathcal{G}(\mathcal{O}_{F_v^+}) \xrightarrow{\sim} \mathcal{G}(\mathcal{O}_{F_w}) \xrightarrow{\iota} \text{GL}_n(\mathcal{O}_{F_w})$ for all split finite places w in F prime to N where v is $w|_{F^+}$ here. For each split place v of F^+ , we choose a place \tilde{v} of F dividing v . For a split v prime to N , let ι_v be the composition of $\iota_{\tilde{v}}$ and the canonical isomorphism $\text{GL}_n(\mathcal{O}_{F_{\tilde{v}}}) \cong \text{GL}_n(\mathcal{O}_{F_v^+})$ (suppressing the dependence on the choice of \tilde{v}).

Let S_p be the set of all places in F^+ dividing p . Suppose from now on that all places in S_p are split. If $U = U_p U^{\infty,p} \leq G(\mathbb{A}_{F^+,p}^\infty) \times G(\mathbb{A}_{F^+}^{\infty,p})$ is a compact open subgroup and W is a finite \mathcal{O} -module endowed with a continuous action of U_Σ for some finite set Σ of finite places of F^+ , then we define the space of algebraic automorphic forms on G of level U and coefficients in W to be the (finite) \mathcal{O} -module

$$(5.3) \quad S(U, W) \stackrel{\text{def}}{=} \{ f : G(F^+) \backslash G(\mathbb{A}_{F^+}^\infty) \rightarrow W \mid f(gu) = u_\Sigma^{-1} f(g) \ \forall g \in G(\mathbb{A}_{F^+}^\infty), u \in U \}.$$

We recall that the level U is said to be *sufficiently small* if for all $t \in G(\mathbb{A}_{F^+}^\infty)$, the order of the finite group $t^{-1}G(F^+)t \cap U$ is prime to p . If U is sufficiently small, then $S(U, -)$ defines an exact functor from finite \mathcal{O} -modules with a continuous U_p -action to finite \mathcal{O} -modules. From now on we assume that U is sufficiently small.

For a finite place v of F^+ prime to N , we say that U is *unramified* at v if one has a decomposition $U = \mathcal{G}(\mathcal{O}_{F_v^+})U^v$. Let S be a finite set of finite places in F^+ containing all places dividing pN , Σ , and all places at which U is *not* unramified.

Let \mathcal{P}_S be the set of split finite places w of F such that $w|_{F^+} \notin S$. For any subset $\mathcal{P} \subseteq \mathcal{P}_S$ of finite complement that is closed under complex conjugation, we write $\mathbb{T}_{\mathcal{P}} \stackrel{\text{def}}{=} \mathcal{O}[T_w^{(i)}, w \in \mathcal{P}, 0 \leq i \leq n]$ for the universal Hecke algebra on \mathcal{P} . The space of algebraic automorphic forms $S(U, W)$ is endowed with an action of $\mathbb{T}_{\mathcal{P}}$, where $T_w^{(i)}$ acts by the usual double coset operator

$$\iota_w^{-1} \left[\text{GL}_n(\mathcal{O}_{F_w}) \begin{pmatrix} \varpi_w \text{Id}_i & 0 \\ 0 & \text{Id}_{n-i} \end{pmatrix} \text{GL}_n(\mathcal{O}_{F_w}) \right].$$

Let $\mathbb{T}_{\mathcal{P}}(U, W)$ be the image of $\mathbb{T}_{\mathcal{P}}$ in $\text{End}_{\mathcal{O}}(S(U, W))$ —it is a finite flat \mathcal{O} -algebra and in particular a complete semilocal ring. Enlarging E if necessary, we assume that the residue fields are identified with \mathbb{F} . If Q is the (finite) set $\{w|_{F^+} : w \in \mathcal{P}_S \setminus \mathcal{P}\}$, then we also denote $\mathbb{T}_{\mathcal{P}}(U, W)$ by $\mathbb{T}^Q(U, W)$. For a maximal ideal $\mathfrak{m} \subset \mathbb{T}^Q(U, W)$, there is a semisimple Galois representation $\bar{r} \stackrel{\text{def}}{=} \bar{r}_{\mathfrak{m}} : G_{F^+, S} \rightarrow \mathcal{G}_n(\mathbb{F})$, where \mathcal{G}_n is the group scheme over \mathbb{Z} defined in [CHT08, §2.1], uniquely determined by the equation

$$(5.4) \quad \det(1 - \bar{r}_{\mathfrak{m}}|_{G_F}(\text{Frob}_w)X) = \sum_{j=0}^n (-1)^j (\mathbf{N}_{F/\mathbb{Q}}(w))^{(j)} (T_w^{(j)} \bmod \mathfrak{m}) X^j.$$

Definition 5.5.1. We say that such a Galois representation $\bar{r} : G_{F^+, S} \rightarrow \mathcal{G}_n(\mathbb{F})$ is *automorphic* of level U and coefficients W if \bar{r} satisfies (5.4) for a finite subset $Q \subset \mathcal{P}_S$ closed under complex conjugation and a maximal ideal $\mathfrak{m} \subset \mathbb{T}^Q(U, W)$. In this case, we say that \mathfrak{m} is the maximal ideal (of $\mathbb{T}^Q(U, W)$ or $\mathbb{T}_{\mathcal{P}}$) corresponding to \bar{r} .

We say that \bar{r} is *automorphic* if \bar{r} is automorphic of some level U and some coefficients W .

We now suppose that $\bar{r}_{\mathfrak{m}}$ is absolutely irreducible. Let $\alpha : \mathbb{T}_{\mathcal{P}} \twoheadrightarrow \mathbb{T}^Q(U, W)_{\mathfrak{m}}$ be the natural quotient map. Then there is a Galois representation $r_{\mathfrak{m}} \stackrel{\text{def}}{=} r(U, W)_{\mathfrak{m}} : G_{F^+, S} \rightarrow \mathcal{G}_n(\mathbb{T}^Q(U, W)_{\mathfrak{m}})$ determined by the equations

$$\det(1 - r(U, W)_{\mathfrak{m}}|_{G_F}(\text{Frob}_w)X) = \sum_{j=0}^n (-1)^j (\mathbf{N}_{F/\mathbb{Q}}(w))^{(j)} \alpha(T_w^{(j)}) X^j$$

for all $w \in \mathcal{P}$.

For each $v \in S_p$, there is an isomorphism $\iota_v : G_{/F_v^+} \cong G_{/F_v} \cong \text{GL}_{d_v}(D_{\tilde{v}/F_v})$ for some $d_v \in \mathbb{N}$ and some central division algebra $D_{\tilde{v}}$ over $F_{\tilde{v}}$ where $\text{GL}_{d_v}(D_{\tilde{v}/F_v})(R) \stackrel{\text{def}}{=} \text{GL}_{d_v}(D_{\tilde{v}} \otimes_{F_{\tilde{v}}} R)$. We now let U_v be $\iota_v^{-1}(\text{GL}_{d_v}(\mathcal{O}_{D_{\tilde{v}}}))$ and U_p be $\prod_{v \in S_p} U_v = \iota_p^{-1}(\prod_{v \in S_p} \text{GL}_{d_v}(\mathcal{O}_{D_{\tilde{v}}}))$ where ι_p denotes $\prod_{v \in S_p} \iota_v$.

Definition 5.5.2. Suppose that $U^{\infty, p}$ is such that $U = U_p U^{\infty, p}$ is a sufficiently small compact open subgroup of $G(\mathbb{A}_{F^+}^{\infty})$ and let σ be an irreducible representation of $\prod_{v \in S_p} \text{GL}_{d_v}(\mathcal{O}_{D_{\tilde{v}}})$ over \mathbb{F} .

We say that \bar{r} is *automorphic of weight* σ and level U if \bar{r} is automorphic of level U and coefficients $\sigma^{\vee} \circ \iota_p$, where σ^{\vee} denotes the \mathbb{F} -dual of σ . We say that \bar{r} is automorphic of weight σ or σ is a *modular (Serre) weight* for \bar{r} if \bar{r} is automorphic of weight σ and some level U .

Let $W(\bar{r})$ be the set of modular Serre weights of \bar{r} .

For each v , we fix an embedding $\overline{F}^+ \hookrightarrow \overline{F}_v^+$ such that the restriction $F \hookrightarrow \overline{F}_v^+$ induces the place \tilde{v} . Let \bar{r}_v be the restriction of \bar{r} to $G_{F_v^+} \cong G_{F_{\tilde{v}}}$, and let \bar{r}_p be the L -homomorphism over \mathbb{F} corresponding to the collection $(\bar{r}_v)_{v \in S_p}$. One expects that $W(\bar{r})$ depends only on \bar{r}_p .

5.5.2. *Minimal level.* We now introduce a space of modular forms at *minimal level*. Suppose that F/F^+ , G , and \bar{r} are as before. Assume moreover that F/F^+ is unramified at all finite places and that \bar{r} is ramified only at split places.

We begin with some notation and terminology. If v is a split place of F^+ , then we define the minimally ramified type τ_v at v (with respect to \bar{r}) to be the inertial type obtained from the restriction to inertia of any minimally ramified lift of $\bar{r}|_{G_{F_v^+}}$ in the sense of [CHT08, Definition 2.4.14]).

Let v_1 be a split place of F^+ away from p such that

- v_1 does not split completely in $F(\zeta_p)$; and
- $\bar{r}|_{G_{F_{v_1}^+}}$ is unramified and $\bar{r}(\text{Frob}_{F_{v_1}^+})$ has distinct eigenvalues, no two of which have ratio equal to $(\mathbf{N}v_1)^{\pm 1}$.

(It is possible to find such a v_1 if $\bar{r}(G_F)$ contains $\text{GL}_n(\mathbb{F})$ with $\#\mathbb{F} > 3n$, see [CEG⁺16, §2.3].)

Let $U \subset G(\mathbb{A}_{F^+}^\infty)$ be the compact open subgroup $\prod_v U_v$ where U_v is

- $\iota_{\bar{v}}^{-1}(\text{GL}_n(\mathcal{O}_{F_{\bar{v}}}))$ if v is a split place of F^+ not equal to v_1 ;
- the preimage of the upper triangular matrices under the composition

$$G(\mathcal{O}_{F_{v_1}^+}) \xrightarrow{\iota_{\bar{v}_1}} \text{GL}_n(\mathcal{O}_{F_{\bar{v}_1}}) \rightarrow \text{GL}_n(k_{\bar{v}_1})$$

if $v = v_1$; and

- hyperspecial if v is an inert place.

Then the compact open subgroup U is sufficiently small.

Let Σ be the set of places of F^+ away from p where \bar{r} ramifies. Recall that S is a finite set of places of F^+ containing all places dividing pN , Σ , and v_1 . For any subset $\mathcal{P} \subseteq \mathcal{P}_S$ of finite complement that is closed under complex conjugation, we write $\mathbb{T}'_{\mathcal{P}} \stackrel{\text{def}}{=} \mathbb{T}_{\mathcal{P}}[T_{\bar{v}_1}^{(i)}, 0 \leq i \leq n]$ where $\mathbb{T}_{\mathcal{P}}$ is the universal Hecke algebra on \mathcal{P} as before. For a U_p -module V , $\mathbb{T}'_{\mathcal{P}}$ acts on the space

$$S(U, (\otimes_{v \in \Sigma} \sigma(\tau_v^\vee)^\circ \circ \iota_v) \otimes V)$$

where the action of $T_{\bar{v}_1}^{(i)}$ is by the double coset operator $U_{v_1} \iota_{\bar{v}_1}^{-1} \begin{pmatrix} \varpi_w \text{Id}_i & 0 \\ 0 & \text{Id}_{n-i} \end{pmatrix} U_{v_1}$.

Choose an ordering $\delta_1, \dots, \delta_n$ of the distinct eigenvalues of $\bar{r}(\text{Frob}_{\bar{v}_1})$ and let \mathfrak{m}' be the maximal ideal of $\mathbb{T}'_{\mathcal{P}}$ generated by $\mathfrak{m} \subset \mathbb{T}_{\mathcal{P}}$ and the elements $T_{\bar{v}_1}^{(i)} - (\mathbf{N}v_1)^{i(1-i)/2}(\delta_1 \cdots \delta_i)$. Then the space $S(U, (\otimes_{v \in \Sigma} \sigma(\tau_v^\vee)^\circ \circ \iota_v) \otimes V)_{\mathfrak{m}'}$ is nonzero.

5.5.3. *G quasisplit at p .* With G as in §5.5.1, we furthermore suppose in this section that $G_{/F_v^+}$ is quasisplit for all $v \in S_p$, i.e., $G_{/F_v^+} \cong \text{GL}_{n/F_v^+}$.

Definition 5.5.3. We say that \bar{r} is *potentially diagonalizably automorphic* if there is a U , W , Q , and a homomorphism $\lambda : \mathbb{T}^Q(U, W)_{\mathfrak{m}} \rightarrow \overline{\mathbb{Q}}_p$ such that if $r_\lambda : G_{F^+} \rightarrow \mathcal{G}(\overline{\mathbb{Q}}_p)$ is the attached semisimple Galois representation characterized by the equation

$$(5.5) \quad \det(1 - r_\lambda|_{G_F}(\text{Frob}_w)X) = \sum_{j=0}^n (-1)^j (\mathbf{N}_{F/\mathbb{Q}}(w)) \binom{j}{2} \lambda(T_w^{(j)}) X^j,$$

then $r_{\lambda,v}$ is potentially diagonalizable for all $v \in S_p$.

Lemma 5.5.4. *Let U_p be as in §5.5.1 and suppose that $U = U_p U^{\infty,p} \subset G(\mathbb{A}_{F^+}^\infty)$ is a sufficiently small compact open subgroup. Let Σ be a finite set of finite places of F^+ away from p . Let W be a finite $\mathcal{O}[U_\Sigma]$ -module.*

Then there is a patching functor M_∞ such that for any finite \mathbb{F} -module V with a continuous $\prod_{v \in S_p} \mathrm{GL}_n(\mathcal{O}_{F_v^+})$ -action,

$$(5.6) \quad M_\infty(V)/\mathfrak{m}_\infty \cong S(U, W \otimes_{\mathcal{O}} V^\vee \circ \iota_p)[\mathfrak{m}]^\vee,$$

where $\mathfrak{m}_\infty \subset R_\infty$ denotes the maximal ideal. In particular, $M_\infty(V)$ is nonzero if and only if $S(U, W \otimes_{\mathcal{O}} V^\vee \circ \iota_p)_{\mathfrak{m}}$ is nonzero.

If \bar{r} is potentially diagonalizably automorphic, then there is an M_∞ as above which is moreover potentially diagonalizable.

Suppose now that F/F^+ is unramified at all finite places and that \bar{r} is ramified only at split places. Let U and \mathfrak{m}' be as in §5.5.2. If W is $\otimes_{v \in \Sigma} \sigma(\tau_v^\vee)^\circ \circ \iota_v$ where τ_v is the minimally ramified type with respect to \bar{r} and $\sigma(\tau_v^\vee)^\circ \subset \sigma(\tau_v^\vee)$ is an \mathcal{O} -lattice, then there is a minimal patching functor M_∞ such that for any finite \mathbb{F} -module V as before,

$$(5.7) \quad M_\infty(V)/\mathfrak{m}_\infty \cong S(U, W \otimes_{\mathcal{O}} V^\vee \circ \iota_p)[\mathfrak{m}']^\vee.$$

If \bar{r} is potentially diagonalizably automorphic, then this minimal M_∞ can be taken to be potentially diagonalizable.

Proof. Except for Definition 5.2.1(3) and the minimality, this follows from the proof of [LLHLMa, Lemma A.1.1] using that \mathfrak{m}_∞ is the preimage of \mathfrak{m} in *loc. cit.* under the map $R_\infty \rightarrow R_\infty/\mathfrak{a}_\infty$.

Suppose the existence of τ_0 as in Definition 5.2.1(3). Then by the above, \bar{r} is potentially diagonalizably automorphic. Let τ be an inertial L -parameter and x be a potentially diagonalizable \bar{E} -point of $\mathrm{Spec} R_\infty(\tau)$. There is an \bar{E} -point y of $\mathrm{Spec} R_\infty(\tau)/\mathfrak{a}_\infty$ which is on the same irreducible component of $\mathrm{Spec} R_\infty(\tau)$ as x by [Paš16, Lemma 3.9]. For any \mathcal{O} -lattice $\sigma(\tau)^\circ \subset \sigma(\tau)$, $M_\infty(\sigma(\tau)^\circ)/\mathfrak{a}_\infty$, and thus $M_\infty(\sigma(\tau)^\circ)$, is supported at y by [LLHL19, Theorem 4.3.1] and the properties of $\sigma(\tau)$ (see §2.3.4). Since $M_\infty(\sigma(\tau)^\circ)$ is a maximal Cohen–Macaulay $R_\infty(\tau)$ -module, it is supported at x as well.

The construction of M_∞ in the minimal level case is as in [Le18, §4] ($n = 3$ and p is assumed to be split, but the modifications are simple). \square

Theorem 5.5.5 (Modularity of extremal weights). *Let $\bar{r} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be an automorphic representation such that*

- $\bar{r}|_{G_{F(\zeta_p)}}$ is adequate; and
- \bar{r}_p is $6e(n-1)$ -generic (in particular $p \nmid 2n$).

Then the following are equivalent:

- (1) if $W_{\mathrm{extr}}(\bar{r}_p) \cap W(\bar{r}_p) \neq \emptyset$;
- (2) $W_{\mathrm{extr}}(\bar{r}_p) \subset W(\bar{r}_p)$; and
- (3) \bar{r} is potentially diagonalizably automorphic.

Proof. Using Lemma 5.5.4 with $U' = U$, the result follows from Theorem 5.4.6. \square

Theorem 5.5.6 (Automorphic tameness criterion). *Let $\sigma_w, \sigma_{w_0w} \in W_{\mathrm{extr}}(\bar{r}_p^{\mathrm{ss}})$ be the extremal weights of \bar{r}_p^{ss} corresponding to w and $w_0w \in W$, respectively. Suppose that $\sigma_w \in W(\bar{r}_p)$. Then the following are equivalent:*

- (1) $\sigma_{w_0w} \in W(\bar{r}_p)$; and
- (2) $\bar{r}_p = \bar{r}_p^{\mathrm{ss}}$.

Proof. Use Theorem 5.5.5 and Proposition 3.7.3. \square

Corollary 5.5.7. *Suppose that $F(\lambda) \in W(\bar{r}_p)$ for $\lambda \in \mathcal{C}_0$ (in particular, $F(\lambda) \in W_{\text{extr}}(\bar{r}_p)$). Let $(F(\lambda), \bar{r}_p^{\text{sp}}) \in SP(\bar{r}_p)$ any lift of $F(\lambda) \in W_{\text{extr}}(\bar{r}_p)$. Then \bar{r}_p is semisimple if and only if*

$$F\left(\left(t_\eta w_0 w \theta_{\bar{r}_p}^\zeta((F(\lambda), \bar{r}_p^{\text{sp}})) w_0^{-1}\right) \cdot (e(w_0(\eta) - \eta)) + w_0 \cdot (\lambda - \eta)\right) \in W(\bar{r}_p)$$

where $w \in \underline{W}$ is such that $F(\lambda)$ is the obvious weight of \bar{r}_p^{sp} corresponding to w .

Theorem 5.5.8 (mod p multiplicity one). *Suppose that F/F^+ is unramified at all finite places, G is quasisplit at all finite places, and that if $\bar{r}|_{G_{F_v^+}}$ is ramified for a finite place v of F^+ , then v splits in F . Let U be as in §5.5.2.*

Let Σ be the set of finite places of F^+ away from p at which \bar{r} is ramified. For each $v \in \Sigma$, let τ_v be the minimally ramified inertial type corresponding to $\bar{r}|_{G_{F_v^+}} : G_{F_v^+} \rightarrow \text{GL}_n(\mathbb{F})$. If \bar{r} satisfies the equivalent conditions of Theorem 5.5.5, then for each $\sigma \in W_{\text{extr}}(\bar{r}_p)$,

$$(5.8) \quad S(U, \otimes_{v \in \Sigma} \sigma^\circ(\tau_v^\vee) \circ \iota_\Sigma \otimes_{\mathcal{O}} \sigma \circ \iota_p)[\mathfrak{m}]$$

is one-dimensional over \mathbb{F} .

Proof. This follows from (5.7) and Theorem 5.4.9. \square

Remark 5.5.9. Using Theorem 5.5.8, one can recover the main results of [Enn18] (with stronger genericity assumptions) which assert a multiplicity one statement for the *ordinary part* of (5.8).

We require the following “change of type” result.

Theorem 5.5.10. *Let F^+ be a totally real field and $F \subset \bar{F}^+$ a CM extension where every place of F^+ dividing p splits in F . Suppose further that $\zeta_p \notin F$. For each place v of F^+ dividing p , choose an embedding $\bar{F}^+ \hookrightarrow \bar{F}_v^+$.*

Let $\bar{r} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be a Galois representation such that $\bar{r}(G_{F(\zeta_p)})$ is adequate and there is a RACSDC automorphic representation Π of $\text{GL}_n(\mathbb{A}_F)$ such that

- $\bar{r}|_{G_F} \cong \bar{r}_{p,\iota}(\Pi)$; and
- for each $v|p$, $r_{p,\iota}(\Pi)|_{G_{F_v^+}}$ is potentially diagonalizable.

Let Δ be a finite set of places in F away from p which split in F such that if $w \in \Delta$, then Π_w is supercuspidal. For each place v of F^+ dividing p , suppose that \bar{r}_v admits a potentially diagonalizable lift which is potentially crystalline of type $(\lambda_v + \eta_v, \tau_v)$.

Then there exists a RACSDC automorphic representation π of $\text{GL}_n(\mathbb{A}_F)$ such that

- $\bar{r}|_{G_F} \cong \bar{r}_{p,\iota}(\pi)$;
- for each $v|p$, $r_{p,\iota}(\pi)|_{G_{F_v^+}}$ is potentially diagonalizable and potentially crystalline of type $(\lambda_v + \eta_v, \tau_v)$; and
- for each $w \in \Delta$, π_w is supercuspidal.

Proof. This follows from [LLHL19, Theorem 4.3.1], which is based on [BGG18, Theorem 3.1.3], except for the assertion of supercuspidality. However, [BGG18, Theorem 3.1.3] with S chosen to contain S_p and $\Delta^+ \stackrel{\text{def}}{=} \{w|_{F^+} \mid w \in \Delta\}$ guarantees that one can choose π so that $r_{p,\iota}(\Pi)|_{G_{F_w}} \sim r_{p,\iota}(\pi)|_{G_{F_w}}$ for each $w \in \Delta$. In particular, the irreducibility of $\text{WD}(r_{p,\iota}(\Pi)|_{G_{F_w}})|_{W_{F_w}}$ implies the irreducibility of $\text{WD}(r_{p,\iota}(\pi)|_{G_{F_w}})|_{W_{F_w}}$, which implies the desired assertion. \square

Corollary 5.5.11. *Let F^+ be a totally real field and $F \subset \bar{F}^+$ a CM extension where every finite place of F^+ is unramified in F and every place dividing p splits. Suppose further that p is unramified*

in F^+ and that $\zeta_p \notin F$. For each place v of F^+ dividing p , choose an embedding $\overline{F}^+ \hookrightarrow \overline{F}_v^+$. Fix a set Δ of split places in F away from p .

Let $\bar{r} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be a Galois representation such that $\bar{r}(G_{F(\zeta_p)})$ is adequate, $\bar{r}|_{G_{F_v^+}}$ is $(6n-2)$ -generic for all $v|p$, and there is a RACSDC automorphic representation Π of $\mathrm{GL}_n(\mathbb{A}_F)$ such that

- $\bar{r}|_{G_F} \cong \bar{r}_{p,\iota}(\Pi)$;
- for each $v|p$, $r_{p,\iota}(\Pi)|_{G_{F_v^+}}$ is potentially diagonalizable; and
- for each $w \in \Delta$, Π_w is supercuspidal.

For each place v of F^+ dividing p , let τ_v be a tame inertial type. Then the following are equivalent:

- (1) $R_{\bar{r}_v}^{\tau_v}$ is nonzero for all places v of F^+ dividing p ;
- (2) there is a RACSDC automorphic representation π of $\mathrm{GL}_n(\mathbb{A}_F)$ such that
 - $\bar{r}|_{G_F} \cong \bar{r}_{p,\iota}(\pi)$;
 - $r_{p,\iota}(\pi)|_{G_{F_v^+}}$ is potentially crystalline of type (η_v, τ_v) for all $v|p$; and
 - for each $w \in \Delta$, π_w is supercuspidal.

Proof. (2) immediately implies (1). We now assume (1) and show the converse. (1) in particular implies that τ is $(5n-1)$ -generic, so that Proposition 3.10.3 applies. Indeed, τ is $(5n-4)$ -generic by [Enn19, Proposition 7]. Then τ is in fact $(5n-1)$ -generic by [LLHL19, Theorem 3.2.1]. Let Δ^+ be the set $\{w|_{F^+} \mid w \in \Delta\}$. Recall from the proof of Theorem 5.5.5 that for each $v \in S_p$, \bar{r}_v admits a potentially diagonalizable lift of type (η_v, τ'_v) for some tame inertial type τ'_v . Let π be the RACSDC automorphic representation of $\mathrm{GL}_n(\mathbb{A}_F)$ guaranteed by Theorem 5.5.10. [Lab11, Theorem 5.4] implies that for some supercuspidal inertial types $(\tau'_v)_{v \in \Delta^+}$,

$$S(U, \otimes_{v \in S_p} \sigma^\circ(\tau'_v{}^N) \circ \iota_p \otimes \otimes_{v \in \Delta^+} \sigma^\circ(\tau'_v{}^N) \circ \iota_{\Delta^+})_{\mathfrak{m}} \neq 0$$

where $\sigma^\circ(\tau'_v{}^N) \subset \sigma(\tau'_v{}^N)$ is a $\mathrm{GL}_n(\mathcal{O}_{F_v^+})$ -stable \mathcal{O} -lattice for each $v \in S_p \cup \Delta^+$. Let M_∞ be the potentially diagonalizable patching functor guaranteed by Lemma 5.5.4 with $W \stackrel{\mathrm{def}}{=} \otimes_{v \in \Delta^+} \sigma^\circ(\tau'_v{}^N) \circ \iota_{\Delta^+}$.

Theorem 5.4.6 implies that $W_{\mathrm{extr}}(\bar{r}_p) \subset W_{M_\infty}(\bar{r}_p)$. Properties of M_∞ from Lemma 5.5.4 imply that

$$S(U, \otimes_{v \in S_p} (\sigma^\vee \circ \iota_p) \otimes \otimes_{v \in \Delta^+} (\sigma^\circ(\tau'_v{}^N) \circ \iota_{\Delta^+}))_{\mathfrak{m}} \neq 0$$

for any $\sigma \in W_{\mathrm{extr}}(\bar{r}_p)$. Exactness of $S(U, -)_{\mathfrak{m}}$ and Proposition 3.10.3 imply that

$$S(U, \otimes_{v \in S_p} (\sigma^\circ(\tau'_v{}^N) \circ \iota_p) \otimes \otimes_{v \in \Delta^+} (\sigma^\circ(\tau'_v{}^N) \circ \iota_{\Delta^+}))_{\mathfrak{m}} \neq 0.$$

We conclude with an application of [Lab11, Corollaire 5.3]. □

5.5.4. *G anisotropic mod center at p.* With G as in §5.5.1, we furthermore suppose in this section that for all $v \in S_p$, $G_{/F_v^+}$ is anisotropic modulo center, i.e., we have an isomorphism $\iota_v : G_{/F_v^+} \xrightarrow{\sim} D_{\bar{v}/F_{\bar{v}}}^\times$. We first recall the set of irreducible $\mathcal{O}_{D_{\bar{v}}}^\times$ -representations over \mathbb{F} (or *Serre weights*).

Let $\mathfrak{m}_{D_{\bar{v}}} \subset \mathcal{O}_{D_{\bar{v}}}$ denote the maximal ideal. Then $k_{D_{\bar{v}}} \stackrel{\mathrm{def}}{=} \mathcal{O}_{D_{\bar{v}}}/\mathfrak{m}_{D_{\bar{v}}}$ is a degree n field extension of the residue field $k_{\bar{v}}$ of $F_{\bar{v}}$. We say that a character of $\mathcal{O}_{D_{\bar{v}}}^\times$ is *tame* if it factors through $k_{D_{\bar{v}}}^\times$.

Since $1 + \mathfrak{m}_{D_{\bar{v}}}$ is a pro- p group (under multiplication), it acts trivially on any irreducible $\mathcal{O}_{D_{\bar{v}}}^\times$ -representation over \mathbb{F} . Thus any irreducible $\mathcal{O}_{D_{\bar{v}}}^\times$ -representation over \mathbb{F} is a tame \mathbb{F} -character. Moreover, the \mathcal{O} -Teichmüller lift gives a bijection between irreducible $\mathcal{O}_{D_{\bar{v}}}^\times$ -representations over \mathbb{F} and tame \mathcal{O} -valued characters of $\mathcal{O}_{D_{\bar{v}}}^\times$.

Given a tame character $\chi_v : \mathcal{O}_{D_{\tilde{v}}}^\times \rightarrow k_{D_{\tilde{v}}}^\times \rightarrow \mathcal{O}^\times$, we define a tame inertial type $\tau(\chi_v)$ as follows. Let $K_{\tilde{v}}$ be $W(k_{D_{\tilde{v}}})[p^{-1}] \otimes_{W_{k_{\tilde{v}}}[p^{-1}]} F_{\tilde{v}}$ and choose an $F_{\tilde{v}}$ -linear embedding of $K_{\tilde{v}} \hookrightarrow \overline{F}_v^+$. We also denote by χ_v the character $\mathcal{O}_{K_{\tilde{v}}}^\times \rightarrow k_{D_{\tilde{v}}}^\times \xrightarrow{\chi_v} \mathcal{O}^\times$. Then we let $\tau(\chi_v)$ be $\text{Ind}_{W_{K_{\tilde{v}}}}^{W_{F_v^+}} (\tilde{\chi}_v \circ \text{Art}_{K_{\tilde{v}}}^{-1})|_{I_{F_v^+}}$ for an extension $\tilde{\chi}_v : K_{\tilde{v}}^\times \rightarrow \mathcal{O}^\times$ of $\chi_v|_{\mathcal{O}_{K_{\tilde{v}}}^\times}$. The tame inertial type $\tau(\chi_v)$ does not depend on the choice of embedding $K_{\tilde{v}} \hookrightarrow \overline{F}_v^+$ or extension $\tilde{\chi}_v$.

Lemma 5.5.12. *Let $\lambda : \mathbb{T}^Q(U, W) \rightarrow \overline{\mathbb{Q}}_p$ be a homomorphism and $r_\lambda : G_{F^+} \rightarrow \mathcal{G}(\overline{\mathbb{Q}}_p)$ be the attached semisimple Galois representation characterized by (5.5).*

Let $\chi = \otimes_{v \in S_p} \chi_v : \prod_{v \in S_p} \mathcal{O}_{D_{\tilde{v}}}^\times \rightarrow \mathcal{O}^\times$ be a tame character. If $\tau(\chi_v)$ is a regular tame inertial type for all $v \in S_p$, then the following are equivalent:

- (1) *for each $v \in S_p$, $r_\lambda|_{G_{F_v^+}}$ is potentially crystalline of type $(\eta_v, \tau(\chi_v))$; and*
- (2) *$S(U, \chi^\vee \circ \iota_p)_{\ker(\lambda)} \neq 0$.*

If $\tau(\chi_v)$ is not regular and $S(U, \chi^\vee \circ \iota_p)_{\ker(\lambda)} \neq 0$, then $r_\lambda|_{G_{F_v^+}}$ is potentially semistable of type (η_v, τ_v) with τ_v not regular.

Proof. Let π be the automorphic representation of $G(\mathbb{A}_{F^+})$ corresponding to λ . First suppose that $\tau(\chi_v)$ is regular for all $v \in S_p$. We will show that (1) and (2) are equivalent to

$$(5.9) \quad \text{rec}_{F_{\tilde{v}}}(\text{JL}(\pi_v))|_{I_{F_{\tilde{v}}}} \cong \tau(\chi_v)$$

for all $v \in S_p$.

Choosing a subring of $F_{\tilde{v}, n} \subset D_{\tilde{v}}$ which is a degree n unramified field extension of $F_{\tilde{v}}$ for each $v \in S_p$, (2) is equivalent to the fact that for each $v \in S_p$, π_v is isomorphic to $\text{Ind}_{F_{\tilde{v}, n}^\times(1+m_{D_{\tilde{v}}})}^{D_{\tilde{v}}^\times} \tilde{\chi}_v$ for some character $\tilde{\chi}_v : F_{\tilde{v}, n}^\times \rightarrow E^\times$ extending χ_v (see [BH11, §1.5]). This is in turn equivalent to (5.9) for all $v \in S_p$ by the main result of [BH11].

Let Π be the automorphic representation of $\text{GL}_n(\mathbb{A}_F)$ in [HKV20, Proposition 6.5.1]. Fixing $v \in S_p$, $|\text{LJ}_{G(F_{\tilde{v}})}|_{\Pi_{\tilde{v}}} \cong \pi_v$ so that $\Pi_{\tilde{v}} \cong \text{JL}(\pi_v)$ (see [Bad08, §3]). Then

$$(5.10) \quad \text{WD}(r_\lambda|_{G_{F_v^+}})^{F\text{-ss}}|_{W_{F_v^+}} \cong \text{WD}(r_{p, \iota}(\Pi)|_{G_{F_{\tilde{v}}}})^{F\text{-ss}}|_{W_{F_{\tilde{v}}}} \cong \text{rec}_{F_{\tilde{v}}}(\text{JL}(\pi_v) \otimes |\det|^{\frac{1-n}{2}})|_{W_{F_{\tilde{v}}}}$$

by [HKV20, Lemma 6.2.2]. Since $r_\lambda|_{G_{F_v^+}}$ is potentially semistable of weight η_v by [BLGGT14, Theorem 2.1.1], we conclude that (1) is also equivalent to (5.9) for all $v \in S_p$.

Now suppose that $\tau(\chi_v)$ is not regular for some $v \in S_p$ and that $S(U, \chi^\vee \circ \iota_p)_{\ker(\lambda)} \neq 0$. As before $r_\lambda|_{G_{F_v^+}}$ is potentially semistable of type (η_v, τ_v) for some inertial type τ_v . We will show that τ_v is tame and is not regular. Let Π be as above. Then as before, $\Pi_{\tilde{v}} \cong \text{JL}(\pi_v)$ so that (5.10) holds. Since $\text{rec}_{F_{\tilde{v}}}(\text{JL}(\pi_v) \otimes |\det|^{\frac{1-n}{2}})|_{I_{F_{\tilde{v}}}}$ is tame and is not regular by [HKV20, Proposition 6.2.3], we conclude that τ_v is tame and is not regular. \square

Theorem 5.5.13. *Let F^+ be a totally real field and $F \subset \overline{F}^+$ a CM extension where every finite place of F^+ is unramified in F and every place dividing p splits. Suppose further that p is unramified in F^+ and that $\zeta_p \notin F$.*

Let $\bar{r} : G_{F^+} \rightarrow \mathcal{G}(\mathbb{F})$ be an automorphic Galois representation such that $\bar{r}(G_{F(\zeta_p)})$ is adequate, $\bar{r}|_{G_{F_v^+}}$ is $(6n-2)$ -generic for all $v|p$, and there is a RACSDC automorphic representation Π of $\text{GL}_n(\mathbb{A}_F)$ such that

- $\bar{r}|_{G_F} \cong \bar{r}_{p, \iota}(\Pi)$;

- for each $v|p$, $r_{p,\iota}(\Pi)|_{G_{F_v^+}}$ is potentially diagonalizable; and
- for each finite place w of F for which G/F_w is not quasisplit, Π_w is supercuspidal.

Let $\chi : D_v^\times \rightarrow E^\times$ be a character. Then $\bar{\chi} \in W(\bar{r})$ if and only if $\bar{r}|_{G_{F_v^+}}$ has a potentially crystalline lift of type $\tau(\chi_v)$ for every $v \in S_p$.

Proof. Suppose that $\bar{\chi} \in W(\bar{r})$. Then $S(U, \chi^\vee \circ \iota_p)_\mathfrak{m}$ is a nonzero finite free \mathcal{O} -module so that there exists λ as in Lemma 5.5.12 such that $S(U, \chi^\vee \circ \iota_p)_{\ker(\lambda)} \neq 0$. We first claim that $\tau(\chi_v)$ is regular for every $v \in S_p$. If $\tau(\chi_v)$ is not regular for some $v \in S_p$, then Lemma 5.5.12 implies that $\bar{r}|_{G_{F_v^+}}$ has a potentially semistable lift of type (η_v, τ_v) for some tame inertial type τ_v which is not regular. This leads to a contradiction since $\bar{r}|_{G_{F_v^+}}$ has no such lift by [Enn19, Proposition 7]. Now since $\tau(\chi_v)$ is regular for every $v \in S_p$, the existence of desired local lifts follows from Lemma 5.5.12.

Suppose now that $\bar{r}|_{G_{F_v^+}}$ has a potentially crystalline lift of type $\tau(\chi_v)$ for every $v \in S_p$. Then let π be as in Corollary 5.5.11. (As in the proof of Corollary 5.5.11, $\tau(\chi_v)$ is $(5n - 1)$ -generic for all $v \in S_p$ and thus cuspidal.) Let π' be the base change cuspidal automorphic representation of $G(\mathbb{A}_F)$ guaranteed by [HKV20, Proposition 6.5.2]. Then $r_{p,\iota}(\pi) \cong r_{p,\iota}(\pi')$ so that in particular $\bar{r}_{p,\iota}(\pi') \cong \bar{r}$ and $r_{p,\iota}(\pi')|_{G_{F_v^+}}$ is potentially crystalline of type (η_v, τ_v) for each $v \in S_p$. Taking λ in Lemma 5.5.12 corresponding to π' , we have that $S(U, \chi^\vee \circ \iota_p)_{\ker(\lambda)}$, and thus $S(U, \bar{\chi}^\vee \circ \iota_p)_\mathfrak{m}$, is nonzero. \square

Remark 5.5.14. In the setting of Theorem 5.5.13, Proposition 3.10.3 implies that for $v|p$, $\bar{r}|_{G_{F_v^+}}$ has a potentially crystalline lift of type $\tau(\chi_v)$ if and only if $W^g(\bar{r}|_{G_{F_v^+}}) \cap \text{JH}(\bar{\sigma}(\tau(\chi_v)))$ is nonempty. Since $\bar{\sigma}(\tau(\chi_v)) = \text{JL}_p(\bar{\chi}_v)$ with JL_p as defined in [Dota, §5], Theorem 5.5.13 implies that $\chi \in W(\bar{r}_p)$ if and only if $\text{JL}_p(\bar{\chi}_v) \cap W^g(\bar{r}|_{G_{F_v^+}})$ is nonempty for all $v|p$. For general G as in §5.5.1, we would expect that $\sigma \in W(\bar{r}_p)$ if and only if $\text{JL}_p(\sigma_v) \cap W^{\text{BM}}(\bar{r}|_{G_{F_v^+}})$ is nonempty for all $v|p$ when the set $W^{\text{BM}}(\bar{r}|_{G_{F_v^+}})$ is defined (see [LLHLMa, §1.6, 8]), but we use $W^g(\bar{r}|_{G_{F_v^+}})$ here because it has been defined in greater generality.

REFERENCES

- [Bad08] Alexandru Ioan Badulescu, *Global Jacquet-Langlands correspondence, multiplicity one and classification of automorphic representations. With an appendix by Neven Grbac.*, Invent. Math. **172** (2008), no. 2, 383–438 (English).
- [BDJ10] Kevin Buzzard, Fred Diamond, and Frazer Jarvis, *On Serre’s conjecture for mod ℓ Galois representations over totally real fields*, Duke Math. J. **155** (2010), no. 1, 105–161. MR 2730374 (2012k:11067)
- [BGG18] Thomas Barnet-Lamb, Toby Gee, and David Geraghty, *Serre weights for $U(n)$* , J. Reine Angew. Math. **735** (2018), 199–224 (English).
- [BGHT11] Tom Barnet-Lamb, David Geraghty, Michael Harris, and Richard Taylor, *A family of Calabi-Yau varieties and potential automorphy. II.*, Publ. Res. Inst. Math. Sci. **47** (2011), no. 1, 29–98 (English).
- [BH11] Colin J. Bushnell and Guy Henniart, *Explicit functorial correspondences for level zero representations of p -adic linear groups*, J. Number Theory **131** (2011), no. 2, 309–331 (English).
- [BHH⁺] Christophe Breuil, Florian Herzig, Yongquan Hu, Stefano Morra, and Benjamin Schraen, *Conjectures and results on modular representations of $GL_n(K)$ for a p -adic field K* , <https://arxiv.org/pdf/2102.06188.pdf>, preprint (2021).
- [BLGGT14] Thomas Barnet-Lamb, Toby Gee, David Geraghty, and Richard Taylor, *Potential automorphy and change of weight*, Ann. of Math. (2) **179** (2014), no. 2, 501–609. MR 3152941
- [BP13] C. Breuil and V. Paskunas, *Towards a modulo p langlands correspondence for GL_2* , vol. 2016, Memoirs of Amer. Math. Soc., 2013.
- [CDT99] Brian Conrad, Fred Diamond, and Richard Taylor, *Modularity of certain potentially Barsotti-Tate Galois representations*, J. Amer. Math. Soc. **12** (1999), no. 2, 521–567. MR 1639612 (99i:11037)
- [CEG⁺16] Ana Caraiani, Matthew Emerton, Toby Gee, David Geraghty, Vytautas Paškūnas, and Sug Woo Shin, *Patching and the p -adic local Langlands correspondence*, Camb. J. Math. **4** (2016), no. 2, 197–287. MR 3529394
- [CEGM17] Frank Calegari, Matthew Emerton, Toby Gee, and Lambros Mavrides, *Explicit Serre weights for two-dimensional Galois representations*, Compos. Math. **153** (2017), no. 9, 1893–1907 (English).
- [CEGS] Ana Caraiani, Matthew Emerton, Toby Gee, and David Savitt, *Moduli stacks of two-dimensional Galois representations*, Proceedings of the International Colloquium on Arithmetic Geometry, TIFR Mumbai, Jan. 6–10, 2020, to appear.
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci. (2008), no. 108, 1–181, With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras. MR 2470687 (2010j:11082)
- [CL18] Ana Caraiani and Brandon Levin, *Kisin modules with descent data and parahoric local models*, Ann. Sci. Éc. Norm. Supér. (4) **51** (2018), no. 1, 181–213. MR 3764041
- [CV92] Robert F. Coleman and José Felipe Voloch, *Companion forms and Kodaira-Spencer theory*, Invent. Math. **110** (1992), no. 2, 263–281 (English).
- [DDR16] Lassina Dembélé, Fred Diamond, and David P. Roberts, *Serre weights and wild ramification in two-dimensional Galois representations*, Forum Math. Sigma **4** (2016), 49 (English), Id/No e33.
- [DL21] Andrea Dotto and Daniel Le, *Diagrams in the mod p cohomology of Shimura curves*, Compos. Math. **157** (2021), no. 8, 1653–1723 (English).
- [Dota] Andrea Dotto, *Breuil-Mézard conjectures for central division algebras*, Algebra and Number Theory, to appear.
- [Dotb] ———, *Serre weights for $GL_m(D)$* , in preparation, (2022).
- [Edi92] Bas Edixhoven, *The weight in Serre’s conjectures on modular forms*, Invent. Math. **109** (1992), no. 3, 563–594 (English).
- [EGa] Matthew Emerton and Toby Gee, *Moduli stacks of étale (φ, Γ) -modules and the existence of crystalline lifts*, Annals of Mathematical Studies, to appear (2022).
- [EGb] ———, *‘Scheme theoretic images’ of morphisms of stacks*, Algebraic Geometry, to appear.
- [EG14] ———, *A geometric perspective on the Breuil-Mézard conjecture*, J. Inst. Math. Jussieu **13** (2014), no. 1, 183–223. MR 3134019
- [EGH13] Matthew Emerton, Toby Gee, and Florian Herzig, *Weight cycling and Serre-type conjectures for unitary groups*, Duke Math. J. **162** (2013), no. 9, 1649–1722. MR 3079258
- [Enn18] John Enns, *Multiplicities in the ordinary part of mod p cohomology for $GL_n(\mathbb{Q}_p)$* , 2018.
- [Enn19] ———, *On weight elimination for $GL_n(\mathbb{Q}_{p^f})$* , Math. Res. Lett. **26** (2019), no. 1, 53–66. MR 3963975

- [Gee11] T. Gee, *On the weights of mod p Hilbert modular forms*, *Invent. Math.* **184** (2011), no. 1, 1–46. MR 2782251 (2012g:11088)
- [GG12] Toby Gee and David Geraghty, *Companion forms for unitary and symplectic groups*, *Duke Math. J.* **161** (2012), no. 2, 247–303. MR 2876931
- [GHS18] Toby Gee, Florian Herzig, and David Savitt, *General Serre weight conjectures*, *J. Eur. Math. Soc. (JEMS)* **20** (2018), no. 12, 2859–2949. MR 3871496
- [GK14] Toby Gee and Mark Kisin, *The Breuil-Mézard conjecture for potentially Barsotti-Tate representations*, *Forum Math. Pi* **2** (2014), e1, 56. MR 3292675
- [GLS14] Toby Gee, Tong Liu, and David Savitt, *The Buzzard-Diamond-Jarvis conjecture for unitary groups*, *J. Amer. Math. Soc.* **27** (2014), no. 2, 389–435. MR 3164985
- [GLS15] ———, *The weight part of Serre’s conjecture for $GL(2)$* , *Forum Math. Pi* **3** (2015), e2, 52. MR 3324938
- [Gro90] Benedict H. Gross, *A tameness criterion for Galois representations associated to modular forms (mod p)*, *Duke Math. J.* **61** (1990), no. 2, 445–517 (English).
- [GS11] Toby Gee and David Savitt, *Serre weights for quaternion algebras*, *Compos. Math.* **147** (2011), no. 4, 1059–1086. MR 2822861 (2012h:11070)
- [HC02] Thomas J. Haines and Ngô Bao Châu, *Alcoves associated to special fibers of local models*, *Amer. J. Math.* **124** (2002), no. 6, 1125–1152. MR 1939783
- [Her09] Florian Herzig, *The weight in a Serre-type conjecture for tame n -dimensional Galois representations*, *Duke Math. J.* **149** (2009), no. 1, 37–116. MR 2541127 (2010f:11083)
- [HH17] Thomas J. Haines and Xuhua He, *Vertexwise criteria for admissibility of alcoves*, *Amer. J. Math.* **139** (2017), no. 3, 769–784. MR 3650232
- [HKV20] Florian Herzig, Karol Koziol, and Marie-France Vignéras, *On the existence of admissible supersingular representations of p -adic reductive groups*, *Forum Math. Sigma* **8** (2020), 73 (English), Id/No e2.
- [HN14] Xuhua He and Sian Nie, *Minimal length elements of extended affine Weyl groups.*, *Compos. Math.* **150** (2014), no. 11, 1903–1927 (English).
- [IM65] N. Iwahori and H. Matsumoto, *On some Bruhat decomposition and the structure of the Hecke rings of p -adic Chevalley groups*, *Inst. Hautes Études Sci. Publ. Math.* (1965), no. 25, 5–48. MR 185016
- [Jan03] Jens Carsten Jantzen, *Representations of algebraic groups*, second ed., *Mathematical Surveys and Monographs*, vol. 107, American Mathematical Society, Providence, RI, 2003. MR 2015057 (2004h:20061)
- [KR00] R. Kottwitz and M. Rapoport, *Minuscule alcoves for GL_n and GSp_{2n}* , *Manuscripta Math.* **102** (2000), no. 4, 403–428. MR 1785323
- [KW09] Chandrashekar Khare and Jean-Pierre Wintenberger, *Serre’s modularity conjecture. I*, *Invent. Math.* **178** (2009), no. 3, 485–504 (English).
- [Lab11] J.-P. Labesse, *Changement de base CM et séries discrètes*, *On the stabilization of the trace formula, Stab. Trace Formula Shimura Var. Arith. Appl.*, vol. 1, Int. Press, Somerville, MA, 2011, pp. 429–470. MR 2856380
- [Le18] Daniel Le, *Lattices in the cohomology of $U(3)$ arithmetic manifolds*, *Math. Ann.* **372** (2018), no. 1-2, 55–89. MR 3856806
- [Lev16] Brandon Levin, *Local models for Weil-restricted groups*, *Compos. Math.* **152** (2016), no. 12, 2563–2601 (English).
- [LLHL19] Daniel Le, Bao Viet Le Hung, and Brandon Levin, *Weight elimination in Serre-type conjectures*, *Duke Math. J.* **168** (2019), no. 13, 2433–2506. MR 4007598
- [LLHLMa] Daniel Le, Bao Viet Le Hung, Brandon Levin, and Stefano Morra, *Local models for Galois deformation rings and applications*, <https://arxiv.org/pdf/2007.05398.pdf>, preprint (2020).
- [LLHLMb] ———, *Serre weights for three-dimensional wildly ramified Galois representations*, <https://arxiv.org/pdf/2202.03303.pdf>, preprint (2021).
- [LLHLM18] ———, *Potentially crystalline deformation rings and Serre weight conjectures: shapes and shadows*, *Invent. Math.* **212** (2018), no. 1, 1–107. MR 3773788
- [LLHM⁺] Daniel Le, Bao Viet Le Hung, Stefano Morra, Chol Park, and Zicheng Qian, *Moduli of Fontaine-Laffaille representations and a mod- p compatibility result*, <https://arxiv.org/pdf/2109.02720.pdf>, preprint (2021).
- [LMS] D. Le, S. Morra, and B. Schraen, *Multiplicity one at full congruence level*, *Journal de l’Institut de Mathématiques de Jussieu*, to appear.
- [New14] James Newton, *Serre weights and Shimura curves*, *Proc. Lond. Math. Soc. (3)* **108** (2014), no. 6, 1471–1500. MR 3218316

- [Paš16] Vytautas Paškūnas, *On 2-dimensional 2-adic Galois representations of local and global fields*, Algebra Number Theory **10** (2016), no. 6, 1301–1358. MR 3544298
- [PR05] G. Pappas and M. Rapoport, *Local models in the ramified case. II: Splitting models*, Duke Math. J. **127** (2005), no. 2, 193–250 (English).
- [Sch08] Michael M. Schein, *Weights in Serre's conjecture for Hilbert modular forms: the ramified case*, Israel J. Math. **166** (2008), 369–391. MR 2430440 (2009e:11090)
- [Ser87] Jean-Pierre Serre, *Sur les représentations modulaires de degré 2 de $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$* , Duke Math. J. **54** (1987), no. 1, 179–230. MR 885783 (88g:11022)
- [Wan87] Jian Pan Wang, *Partial orderings on affine Weyl groups*, J. East China Norm. Univ. Natur. Sci. Ed. (1987), no. 4, 15–25. MR 980127

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAFAYETTE, IN 47907-2067

Email address: ledt@purdue.edu

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, 2033 SHERIDAN ROAD, EVANSTON, ILLINOIS 60208, USA

Email address: lhvietbao@googlemail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, 617 N SANTA RITA AVENUE, TUCSON, ARIZONA 85721, USA

Email address: bwlevin@math.arizona.edu

LABORATOIRE D'ANALYSE, GÉOMÉTRIE, ALGÈBRE, UNIVERSITÉ PARIS 13 ET PARIS 8, 99 AVE. J-B. CLÉMENT, 93430 VILLETANEUSE, FRANCE

Email address: morra@math.univ-paris13.fr