



Extension of Alon's and Friedman's conjectures to Schottky surfaces

Michael Magee, Frédéric Naud

► To cite this version:

Michael Magee, Frédéric Naud. Extension of Alon's and Friedman's conjectures to Schottky surfaces. 2022. hal-03841740

HAL Id: hal-03841740

<https://hal.science/hal-03841740>

Preprint submitted on 7 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Extension of Alon's and Friedman's conjectures to Schottky surfaces

Michael Magee and Frédéric Naud

June 11, 2021

Abstract

Let $X = \Lambda \backslash \mathbb{H}$ be a Schottky surface, that is, a conformally compact hyperbolic surface of infinite area. Let δ denote the Hausdorff dimension of the limit set of Λ .

We prove that for any compact subset $\mathcal{K} \subset \{s : \operatorname{Re}(s) > \frac{\delta}{2}\}$, if one picks a random degree n cover X_n of X uniformly at random, then with probability tending to one as $n \rightarrow \infty$, there are no resonances of X_n in \mathcal{K} other than those already belonging to X (and with the same multiplicity). This result is conjectured to be the optimal one for bounded frequency resonances and is analogous to both Alon's and Friedman's conjectures for random graphs, which are now theorems due to Friedman and Bordenave-Collins, respectively.

Contents

1	Introduction	2
2	Preliminaries	5
2.1	Boundary coding of Schottky groups and Bowen's formula	5
2.2	Random covering spaces, permutations, and representations	7
2.3	Transfer operator and function spaces	7
2.4	Deterministic a priori bounds	9
3	Random permutations and random graphs	14
3.1	Symmetric random permutations	14
3.2	Graphs and paths	15
3.3	Tangles	15
4	Non-backtracking operator and path decomposition	16
4.1	Conjugation between transfer operator and non-backtracking operator	16
4.2	Bordenave's path decomposition	18
5	High trace method	21
5.1	Norm of $\underline{B}(s)^{(\ell)}$	21
5.2	Norm of $R_k(s)^{(\ell)}$	27

1 Introduction

Let X be an infinite area surface equipped with a metric of constant curvature -1 , with a finitely generated non-abelian fundamental group and without cusps. Such a surface is called a *conformally compact hyperbolic surface*.

This paper addresses the question of whether typical such surfaces have (almost) optimal spectral gaps. Here, a spectral gap refers either to a gap in the spectrum of the Laplace-Beltrami operator or, more ambitiously, to a region where there are no resonances of the meromorphically continued resolvent. We give the background to this spectral theory now; for general background on the spectral theory of infinite area hyperbolic surfaces the reader should see [Bor16].

One can obtain X as a quotient $X = \Lambda \backslash \mathbb{H}$ where \mathbb{H} is the hyperbolic upper half plane and Λ is a discrete, finitely generated, non-abelian free subgroup of $\mathrm{PSL}_2(\mathbf{R})$, and, in fact, by a result of Button [But98] there are generators $\gamma_1, \dots, \gamma_d$ of Λ for some $d \geq 2$ which form a Schottky system in the sense of §§2.1. The orbit of Λ on any fixed point of \mathbb{H} accumulates on $\partial\mathbb{H}$ in a set called the *limit set* of Λ ; we write $\delta = \delta(\Lambda)$ for the Hausdorff dimension of the limit set. The assumption that Λ is non-abelian and X is infinite area implies $\delta \in (0, 1)$.

Let Δ_X denote the Laplace-Beltrami operator on $L^2(X)$. The spectrum of Δ_X in the range $[\frac{1}{4}, \infty)$ is always continuous with no embedded eigenvalues, and the spectrum is always discrete below $\frac{1}{4}$, by work of Lax-Phillips [LP81]. If $\delta > \frac{1}{2}$ then the bottom of the spectrum occurs at $\delta(1 - \delta)$ by a result of Patterson [Pat76]. Therefore a naive definition of spectral gap in this case is $\lambda_1 - \delta(1 - \delta)$ where λ_1 is the minimum element of the spectrum other than $\delta(1 - \delta)$, including $\delta(1 - \delta)$ itself if it occurs with multiplicity larger than one. However, if $\delta \leq \frac{1}{2}$ then the spectrum is precisely $[\frac{1}{4}, \infty)$ and this notion of spectral gap has no meaning. Moreover, even if $\delta > \frac{1}{2}$, it does not give the strongest possible information as we explain now.

The resolvent operator

$$R_X(s) \stackrel{\text{def}}{=} (\Delta_X - s(1 - s))^{-1} : C_0^\infty(X) \rightarrow C^\infty(X)$$

has meromorphic continuation from $\mathrm{Re}(s) > \frac{1}{2}$ to the entire complex plane [MM87]. The poles of this family of operators are called *resonances* of X . The multiplicity of a resonance s is given by $\mathrm{rank} \left(\int_\gamma R_X(s) ds \right)$ where γ is an anticlockwise oriented circle enclosing s and no other resonance of X . Resonances s with $\mathrm{Re}(s) > \frac{1}{2}$ yield eigenvalues $s(1 - s)$ with the same multiplicity. We write \mathcal{R}_X for the multiset of resonances of X , including multiplicities.

Now it is apparent that beyond L^2 spectral gaps, we can ask for resonance-free regions of the complex plane. We summarize what is known in this regard assuming that X is connected (otherwise there is no spectral gap in any sense). In the half-plane $\{\mathrm{Re}(s) > \frac{1}{2}\}$, there are finitely many resonances that all lie on the real line, the right-most one being at $s = \delta$. If $\delta \leq \frac{1}{2}$, then we know from [Nau05] that there exists $\epsilon(\Lambda) > 0$ such that in the half plane

$\{\operatorname{Re}(s) > \delta - \epsilon\}$, the only resonance is at $s = \delta$. Therefore the spectral gap is then defined as the maximal size of this $\epsilon(\Lambda)$. It was shown in [JNS19] that this gap can be arbitrarily small.

A conjecture of Jakobson and Naud [JN12, Conj. 2] predicts (among other things) that for any $\epsilon > 0$ there are infinitely many resonances with $\operatorname{Re}(s) > \frac{\delta}{2} - \epsilon$ and hence one does not expect to obtain spectral gap larger than $\frac{\delta}{2}$. It is a pressing question as to whether ‘typical’ surfaces have a spectral gap close to this optimal value; this question has famous analogs in graph theory that we will turn to shortly.

In [MN20] we introduced a model of random conformally compact hyperbolic manifolds based on random covering spaces. For any $n \in \mathbf{N}$ the collection of degree n Riemannian covering spaces of X is a finite set and hence we can pick one of these covering spaces uniformly at random. Note that if X' covers X , then any resonance of X is a resonance of X' , with at least as large multiplicity. The main theorem of this paper is the following.

Theorem 1.1. *Let X_n denote a uniformly random degree n covering space of X . For any compact set $\mathcal{K} \subset \{s : \operatorname{Re}(s) > \frac{\delta}{2}\}$, with probability tending to one as $n \rightarrow \infty$*

$$\mathcal{R}_{X_n} \cap \mathcal{K} = \mathcal{R}_X \cap \mathcal{K}.$$

This theorem says that for any $\epsilon > 0$, the resonance set of random X_n in the region $\{s : \operatorname{Re}(s) > \frac{\delta}{2} + \epsilon\}$ is almost surely optimal *provided we restrict to bounded frequency (imaginary part)*. A weaker version of Theorem 1.1 with $\frac{\delta}{2}$ replaced by $\frac{3\delta}{4}$ was proved in [MN20, Thm. 1.1]

Theorem 1.1 together with the main result of [BMM17] implies the following corollary on L^2 eigenvalues.

Corollary 1.2. *Assume that $\delta > \frac{1}{2}$, and that the base surface¹ X has Euler characteristic -1 . Let X_n denote a uniformly random degree n covering space of X . With probability tending to one as $n \rightarrow \infty$ the only eigenvalue of Δ_{X_n} is $\delta(1 - \delta)$.*

We now explain the analogy with random graphs; this is also discussed in detail in [MN20, Introduction]. A celebrated conjecture of Alon [Alo86] predicted that for any $\epsilon > 0$, a uniformly random d -regular graph on n vertices, with probability tending to one as $n \rightarrow \infty$, has no eigenvalues of its adjacency operator larger than $2\sqrt{d-1} + \epsilon$, other than d . The relevance of the value $2\sqrt{d-1}$ is both that it is the spectral radius of the adjacency operator on the universal cover of a d -regular graph (a d -regular tree), and also, a result of Alon-Boppana [Nil91] states that any sequence of d -regular graphs on n vertices has second largest eigenvalue of their adjacency operators at least $2\sqrt{d-1} - o(1)$ as $n \rightarrow \infty$. Hence the value $2\sqrt{d-1}$ is analogous to $\frac{\delta}{2}$ here. It is interesting however that even though the Alon-Boppana bound is not very hard to prove, the analog of this fact in the current setting is still only a conjecture. Alon’s conjecture was proved by Friedman in [Fri08] (see also [Bor] for a new proof and [Pud15] for a proof of only slightly weaker result).

¹In this case X is either a pair of pants with three funnels or a torus with one funnel.

Friedman conjectured in [Fri03] that a variant of Alon’s conjecture should hold for random degree n covering spaces² of a fixed finite base graph provided

- One replaces $2\sqrt{d-1}$ by the spectral radius of the adjacency operator on the universal cover of the graph, and
- One allow eigenvalues of the adjacency operator that already belonged to the base graph (as one must).

Friedman’s conjecture was proved in a breakthrough work of Bordenave and Collins [BC19]. In fact Bordenave-Collins proved a vast generalization of Friedman’s conjecture where one twists a random Hecke operator, formed from random permutations, by arbitrary fixed finite dimensional matrices, assuming the matrices satisfy a symmetry condition that forces the resulting operator to be self-adjoint. The work of Bordenave-Collins is a vital ingredient of the current work.

The first result on spectral gap of random hyperbolic surfaces is due to Brooks and Makover [BM04] who prove that for a combinatorial model of random closed hyperbolic surface, depending on a parameter n that influences the genus (non-deterministically), that there exists a constant $C > 0$ such that with probability tending to one as $n \rightarrow \infty$, the first non-zero eigenvalue (λ_1) of the Laplacian satisfies $\lambda_1 \geq C$. Mirzakhani proved in [Mir13] that for Weil-Petersson random closed hyperbolic surfaces of genus g , with probability tending to one as $g \rightarrow \infty$ one has $\lambda_1 \geq 0.0024$. In [MN20] the authors of the current paper proved that Theorem 1.1 holds with $\frac{\delta}{2}$ replaced by $\frac{3\delta}{4}$.

Returning to closed surfaces, by building on [MP20], the authors and Puder proved in [MNP20] that for a uniformly random degree n cover of a fixed closed hyperbolic surface, there are for any $\epsilon > 0$, with probability tending to one as $n \rightarrow \infty$, no new eigenvalues of the covering space below $\frac{3}{16} - \epsilon$. This result was adapted to Weil-Petersson random surfaces independently by Wu and Xue [WX21] and Lipnowski and Wright [LW21]; here the corresponding statement is that there are no eigenvalues between 0 and $\frac{3}{16} - \epsilon$. These ‘ $\frac{3}{16}$ ’ results are, when it comes *solely* to L^2 eigenvalues, at the strength of the result of [MN20] giving resonance-free regions in terms of $\frac{3\delta}{4}$; for compact surfaces $\delta = 1$ and the eigenvalue is written $\lambda = s(1-s)$ ($\frac{3}{4}(1-\frac{3}{4}) = \frac{3}{16}$). On the other hand, closed surfaces involve additional difficulties due to their non-free fundamental groups.

There are other related works on Weil-Petersson random surfaces that do not imply spectral gaps, but instead offer some spectral delocalization results [Mon20, GLMST21].

Uniform spectral gap for (deterministic) covering spaces of infinite area hyperbolic surfaces has also been of interest in number theoretic settings; see [BGS11, OW16, MOW17, Gam02] for a selection of results; the single quantitative result here is by Gamburd [Gam02]. Much of the motivation of these spectral gap results came from the ‘thin groups’ research program; see Sarnak’s article [Sar14] for an overview.

Another closely related concept is that of *essential spectral gap*; referring to a half-plane where only finitely many resonances appear. Two important results here are due to Bourgain

²In graph-theoretic literature, these are called n -lifts.

and Dyatlov. Let X be conformally compact as above. The first result, proved in [BD17], says that there is $\epsilon > 0$, *depending only on* δ , such that there are only finitely many resonances in $\{s : \operatorname{Re}(s) > \delta - \epsilon\}$. This result is relevant if $\delta \leq 1/2$. On the other hand, it is proved in [BD18] that there are only finitely many resonances in $\{s : \operatorname{Re}(s) > \frac{1}{2} - \eta\}$ for some $\eta = \eta(\Lambda) > 0$, this result being relevant if $\delta > \frac{1}{2}$. A conjecture of Jakobson-Naud [JN12] says that the optimal essential spectral gap corresponds to finitely many resonances in $\{\operatorname{Re}(s) > \frac{\delta}{2} + \varepsilon\}$ for any $\varepsilon > 0$. We do not know yet if our probabilistic techniques can be used to address high frequency problems (i.e. resonances with large imaginary parts) and have not attempted to do so in the present paper, however the present paper says that in some sense this conjecture holds in the bounded frequency, large cover regime. For a broader perspective on resonances of hyperbolic surfaces than we are able to offer here, the reader can see Zworski's survey article [Zwo17].

Acknowledgments

We thank Charles Bordenave, Benoît Collins, and Doron Puder for helpful conversations related to this work.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 949143).

2 Preliminaries

2.1 Boundary coding of Schottky groups and Bowen's formula

In this section, we set some notations (which are essentially similar to [MN20] and [BD17] with some slight changes).

Let $d \geq 2$ and $\mathcal{I} = \{1, \dots, 2d\}$. If $i \in \mathcal{I}$, then we write $\bar{i} \stackrel{\text{def}}{=} i + d \bmod 2d$ so $i \in \mathcal{I}$. For each $i \in \mathcal{I}$ we are given an *open* disc D_i in \mathbf{C} with center in \mathbf{R} . The closures of the discs D_i for $i \in \mathcal{I}$ are assumed to be disjoint from one another. We let $I_i \stackrel{\text{def}}{=} D_i \cap \mathbf{R}$, an open interval. We write $\mathbf{D} \stackrel{\text{def}}{=} \cup_{i \in \mathcal{I}} D_i$ for the union of the discs.

We consider the usual action of $\operatorname{SL}_2(\mathbf{R})$ by Möbius transformations on the extended complex plane $\hat{\mathbf{C}} = \mathbf{C} \cup \{\infty\}$. We are given for each $i \in \mathcal{I}$ a matrix $\gamma_i \in \operatorname{SL}_2(\mathbf{R})$ with the mapping property

$$\gamma_i \left(\hat{\mathbf{C}} \setminus D_{\bar{i}} \right) = \overline{D_i}, \quad \gamma_{\bar{i}} = \gamma_i^{-1}.$$

We recall that we denote $\Lambda \stackrel{\text{def}}{=} \langle \gamma_i : i \in \mathcal{I} \rangle$ for the group generated by the γ_i . Since the discs D_i are disjoint, Klein's Ping-Pong Lemma shows that Λ is a *free subgroup* of $\operatorname{SL}_2(\mathbf{R})$. The converse is actually true in dimension 2: every conformally compact hyperbolic surface X can be uniformized by a Schottky group Λ so that $X = \Lambda \backslash \mathbb{H}$, see [But98]. The elements of Λ can be encoded by words in the alphabet \mathcal{I} as follows. A *word* is a finite sequence

$$\mathbf{i} = (i_1, \dots, i_n), \quad n \in \mathbf{N} \cup \{0\}$$

such that $i_j \neq \overline{i_{j+1}}$ for $j = 1, \dots, n-1$. We say that n is the *length* of \mathbf{i} and denote this by $|\mathbf{i}| = n$. We write \mathcal{W} for the collection of all words, \mathcal{W}_N for the words of length N , and $\mathcal{W}_{\geq N}$ for the words of length $\geq N$. We write \emptyset for the empty word and write $\mathcal{W}^\circ = \mathcal{W} - \{\emptyset\}$. For $\mathbf{i} = (i_1, \dots, i_n)$, $\mathbf{j} = (j_1, \dots, j_m) \in \mathcal{W}$ we write

- $\mathbf{i}' \stackrel{\text{def}}{=} (i_1, \dots, i_{n-1})$ if $\mathbf{i} = (i_1, \dots, i_n)$ and $n \geq 1$.
- $\hat{\mathbf{i}} \stackrel{\text{def}}{=} (i_2, \dots, i_n)$ if $\mathbf{i} = (i_1, \dots, i_n)$ and $n \geq 1$.
- $\mathbf{i} \rightarrow \mathbf{j}$ if either of \mathbf{i} or \mathbf{j} is empty, or else $i_n \neq \overline{j_1}$, in which case $(i_1, \dots, i_n, j_1, \dots, j_m)$ is in \mathcal{W}° and we write \mathbf{ij} for this concatenation.

If $\mathbf{i} = (i_1, \dots, i_n) \in \mathcal{W}$ then we associate to \mathbf{i} the group element $\gamma_{\mathbf{i}} \stackrel{\text{def}}{=} \gamma_{i_1} \dots \gamma_{i_n}$; here $\gamma_\emptyset = \text{id}$. The map $\mathbf{i} \in \mathcal{W} \mapsto \gamma_{\mathbf{i}} \in \Gamma$ is a one-to-one encoding of Γ . We write $\bar{\mathbf{i}} \stackrel{\text{def}}{=} (\overline{i_n}, \dots, \overline{i_1})$ and call this the *mirror word* of \mathbf{i} . Note that $\gamma_{\bar{\mathbf{i}}} = \gamma_{\mathbf{i}}^{-1}$. If $\mathbf{i} = (\mathbf{i}_1, \dots, \mathbf{i}_n) \in \mathcal{W}^\circ$ we set

$$D_{\mathbf{i}} = \gamma_{\mathbf{i}'}(D_{i_n}), \quad I_{\mathbf{i}} = \gamma_{\mathbf{i}'}(I_{i_n})$$

and write $|I_{\mathbf{i}}|$ for the length of the open interval $I_{\mathbf{i}}$. We view this set up as fixed henceforth, so all constants will depend on Λ .

The *Bowen-Series map* $T : \mathbf{D} \rightarrow \hat{\mathbf{C}}$ is given by

$$T|_{D_{\mathbf{i}}} = \gamma_{\bar{\mathbf{i}}}^{-1} = \gamma_{\bar{\mathbf{i}}}.$$

The Bowen-Series map is eventually expanding [Bor16, Prop. 15.5]; this will be made explicit below so we do not give the general definition now. The limit set $K = K(\Lambda)$ of Λ , defined in the Introduction, coincides with the non-wandering set of T :

$$K(\Lambda) \stackrel{\text{def}}{=} \bigcap_{n=1}^{\infty} T^{-n}(\mathbf{D}).$$

The limit set K is a compact T -invariant subset of \mathbf{R} . Given a Hölder continuous map $\varphi : K \rightarrow \mathbf{R}$, the *topological pressure* $P(\varphi)$ can be defined through the variational formula:

$$P(\varphi) \stackrel{\text{def}}{=} \sup_{\mu} \left(h_{\mu}(T) + \int_{\Lambda} \varphi d\mu \right),$$

where the supremum is taken over all T -invariant probability measures on $K(\Lambda)$, and $h_{\mu}(T)$ stands for the measure-theoretic entropy. A famous result of Bowen [Bow79] says that the map $\mathbf{R} \rightarrow \mathbf{R}$,

$$r \mapsto P(-r \log |T'|)$$

is convex³, strictly decreasing and vanishes exactly at $r = \delta \stackrel{\text{def}}{=} \delta(\Lambda)$, the Hausdorff dimension of the limit set $K(\Lambda)$. In addition, it is not difficult to see from the variational formula that

³Convexity follows obviously from the variational formula above.

$P(-r \log |T'|)$ tends to $-\infty$ as $r \rightarrow +\infty$. For simplicity, we will use the notation $P(r)$ in place of $P(-r \log |T'|)$. The pressure functional plays an important role in the sequel.

2.2 Random covering spaces, permutations, and representations

Let S_n denote the group of permutations of $[n] \stackrel{\text{def}}{=} \{1, \dots, n\}$. Our random covering spaces of X are parameterized by

$$\phi \in \text{Hom}(\Lambda, S_n).$$

Of course, choosing ϕ is the same as choosing

$$\sigma = (\sigma_1, \dots, \sigma_d) \in S_n^d$$

where $\sigma_i \stackrel{\text{def}}{=} \phi(\gamma_i)$. We view σ and ϕ as coupled in this way in the rest of the paper.

Given $\phi \in \text{Hom}(\Lambda, S_n)$, we construct a covering space of X as follows. Let Λ act on $\mathbb{H} \times [n]$ by

$$\gamma(z, i) = (\gamma.z, \phi(\gamma)[i])$$

where the action on the first factor is by Möbius transformation. The quotient

$$X_\phi \stackrel{\text{def}}{=} \Lambda \backslash_\phi (\mathbb{H} \times [n])$$

is a degree n cover of X . Choosing ϕ uniformly at random in $\text{Hom}(\Lambda, S_n)$, the resulting X_ϕ is a uniformly random degree n Riemannian cover of X .

Let $V_n \stackrel{\text{def}}{=} \ell^2([n])$. Given $\phi \in \text{Hom}(\Lambda, S_n)$ as above, we let $\rho_n : \Lambda \rightarrow \text{End}(V_n)$ be the representation obtained by composing ϕ with the standard permutation representation

$$S_n \rightarrow \text{End}(V_n).$$

We let ρ_n^0 denote the restriction of ρ_n to the space V_n^0 of functions on $[n]$ that are orthogonal to constants. The parameters ϕ, ρ_n , and ρ_n^0 (as well as σ) are now coupled for the rest of the paper.

2.3 Transfer operator and function spaces

Let V be a finite dimensional complex Hilbert space. We consider the Bergman space $\mathcal{H}(\mathbf{D}; V)$ that is the space of V -valued holomorphic functions on \mathbf{D} with finite norm with respect to the given inner product

$$\langle f, g \rangle \stackrel{\text{def}}{=} \int_{\mathbf{D}} \langle f(x), g(x) \rangle_V dm(x).$$

Here dm is Lebesgue measure on \mathbf{D} . This splits as an orthogonal direct sum

$$\mathcal{H}(\mathbf{D}; V) = \bigoplus_{i \in \mathcal{I}} \mathcal{H}(D_i; V).$$

If $\{\mathbf{e}_k\}_{k=1}^\infty$ is any orthonormal basis of $\mathcal{H}(D_i; \mathbf{C})$, and $x_1, x_2 \in D_i$, then the sum

$$\sum_{k=1}^{\infty} \mathbf{e}_k(x_1) \overline{\mathbf{e}_k(x_2)} \stackrel{\text{def}}{=} B_{D_i}(x_1, x_2)$$

converges uniformly on every compact subset of D_i and is called the *Bergman kernel of D_i* . It is given by the explicit formula (cf. [Bor16, pg. 378])

$$B_{D_i}(x_1, x_2) = \frac{r_i^2}{\pi [r_i^2 - (\bar{x}_2 - c_i)(x_1 - c_i)]^2} \quad (2.1)$$

where r_i, c_i are the radius and center of D_i . We let

$$B_{\mathbf{D}}(z, w) \stackrel{\text{def}}{=} \sum_{i \in \mathcal{I}} \mathbf{1}\{z, w \in D_i\} B_{D_i}(z, w).$$

An important fact that will be used several times in this paper is the reproducing property of the Bergman kernel: if $f \in \mathcal{H}(\mathbf{D}, V)$, then we have for all $z \in \mathbf{D}$,

$$f(z) = \int_{\mathbf{D}} B_{\mathbf{D}}(z, w) f(w) dm(w),$$

in particular for all compact subsets $K \subset \mathbf{D}$, the above explicit formulas for the kernels show that there exists $C_K > 0$ such that

$$\sup_{z \in K} \|f(z)\|_V \leq C_K \|f\|_{\mathcal{H}(\mathbf{D}, V)}.$$

Throughout the sequel, $\rho : \Lambda \rightarrow \mathbf{U}(V)$ will denote a finite dimensional unitary representation of the Schottky group Λ . The transfer operator acting on $\mathcal{H}(\mathbf{D}, V)$ is defined as follows:

$$\mathcal{L}_{s, \rho}[f](x) f \stackrel{\text{def}}{=} \sum_{\substack{i \in \mathcal{I} \\ i \rightarrow j}} \gamma'_i(x)^s \rho(\gamma_i^{-1}) f(\gamma_i(x)) \quad x \in D_j, j \in \mathcal{I}.$$

Here $(\gamma'_i(x))^s$ is understood as $(\gamma'_i(x))^s := e^{-s\tau(\gamma_i(x))}$, where $\tau(z)$ is the analytic continuation to $\cup_{|\mathbf{i}|=2} D_{\mathbf{i}}$ of $\tau(x) = \log |T'(x)|$, and where T is, as defined above, the Bowen-Series map. In addition, we can assume that τ is also continuous on $\cup_{|\mathbf{i}|=2} \overline{D_{\mathbf{i}}}$.

The following result appears in [Bor16, Lemma 15.7] in the case $V = \mathbf{C}$, and easily extends to the case of finite dimensional V .

Lemma 2.1. *The operator $\mathcal{L}_{s, \rho}$ is a trace class operator on the Hilbert space $\mathcal{H}(\mathbf{D}, V)$.*

Our interest in considering these operators stems from setting $\rho = \rho_n^0$ from §§2.2. The following fact follows from [MN20, Thm 2.2. and Prop. 4.4(2)].

Proposition 2.2. *If s is a resonance of X_ϕ that appears with greater multiplicity than in X ,*

for example, by not being a resonance of X , then

$$\det(1 - \mathcal{L}_{s, \rho_n^0}) = 0$$

and hence 1 is an eigenvalue of $\mathcal{L}_{s, \rho_n^0}$. (The determinant above is a Fredholm determinant.)

2.4 Deterministic a priori bounds

We will now introduce a useful notation from [MN20]. In the rest of the paper, for any $\mathbf{i} \in \mathcal{W}^\circ$, we define

$$\Upsilon_{\mathbf{i}} \stackrel{\text{def}}{=} |I_{\mathbf{i}}|,$$

that is the length of the interval $I_{\mathbf{i}}$ associated to the word \mathbf{i} . For all $\mathbf{i} \in \mathcal{W}_{\geq 2}$, we have obviously

$$\Upsilon_{\mathbf{i}} \leq \Upsilon_{\mathbf{i}'}, \quad (2.2)$$

since $I_{\mathbf{i}} \subset I_{\mathbf{i}'}$. Therefore there exists trivially $c = c(\Lambda) > 0$ such that for any $\mathbf{i} \in \mathcal{W}^\circ$,

$$0 < \Upsilon_{\mathbf{i}} \leq c. \quad (2.3)$$

Lemma 2.3. *There exists a constant $K = K(\Lambda) > 1$ such that the following bounds hold.*

(Rough multiplicativity) *For all $\mathbf{i}, \mathbf{j} \in \mathcal{W}^\circ$ with $\mathbf{i} \rightarrow \mathbf{j}$, we have*

$$K^{-1} \Upsilon_{\mathbf{i}} \Upsilon_{\mathbf{j}} \leq \Upsilon_{\mathbf{ij}} \leq K \Upsilon_{\mathbf{i}} \Upsilon_{\mathbf{j}}. \quad (2.4)$$

(Mirror estimate) *For all $\mathbf{j} \in \mathcal{W}^\circ$, we have*

$$K^{-1} \Upsilon_{\bar{\mathbf{j}}} \leq \Upsilon_{\mathbf{j}} \leq K \Upsilon_{\bar{\mathbf{j}}}. \quad (2.5)$$

(Derivatives) *For all $x \in D_{i_{|\mathbf{i}|}}$, we have*

$$K^{-1} \Upsilon_{\mathbf{i}} \leq |\gamma'_{\mathbf{i}'}(x)| \leq K \Upsilon_{\mathbf{i}}. \quad (2.6)$$

(Exponential Bound) *There is a constant $D = D(\Gamma) > 1$ such that*

$$\Upsilon_{\mathbf{i}} \leq K D^{-|\mathbf{i}|}. \quad (2.7)$$

The proof of these bounds follow from [MN20, Lemmas 3.1, 3.2, 3.4, 3.5] and follows previous estimates from [BD17]. The following result is [MN20, Lemma 3.10].

Lemma 2.4. *For all $r_1, Q \in \mathbf{R}$ such that $0 \leq r_1 < Q$ there is a constant $C = C(r_1, Q) > 0$ such that for all $N \in \mathbf{N}_0$ and $r \in [r_1, Q]$ we have*

$$\sum_{\mathbf{i} \in \mathcal{W}_N} \Upsilon_{\mathbf{i}}^r \leq C \exp(NP(r_1)). \quad (2.8)$$

Lemma 2.5. *Suppose that $\mathcal{K} \subset \mathbf{C}$ is compact. There is a constant $C = C(\mathcal{K}) > 0$ such that for all $\ell \in \mathbf{N}$ and $s, s_0 \in \mathcal{K}$*

$$\|\mathcal{L}_{s, \rho_n^0}^\ell - \mathcal{L}_{s_0, \rho_n^0}^\ell\| \leq |s - s_0| C^\ell,$$

where $\|\cdot\|$ denotes the operator norm on $\mathcal{H}(\mathbf{D}, V_n^0)$.

Proof. Let $f \in \mathcal{H}(\mathbf{D}, V_n^0)$. We write

$$\begin{aligned} \|\mathcal{L}_{s, \rho_n^0}^\ell(f) - \mathcal{L}_{s_0, \rho_n^0}^\ell(f)\|^2 &= \int_{\mathbf{D}} \|\mathcal{L}_{s, \rho_n^0}^\ell(f) - \mathcal{L}_{s_0, \rho_n^0}^\ell(f)\|_{V_n^0}^2 dm \\ &= \sum_{j=1}^{2d} \int_{D_j} \left\| \sum_{\mathbf{i} \rightarrow j, |\mathbf{i}|=\ell} \left(e^{-s\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} - e^{-s_0\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} \right) \rho_n^0(\gamma_{\mathbf{i}}^{-1}) f(\gamma_{\mathbf{i}}z) \right\|_{V_n^0}^2 dm(z), \end{aligned}$$

where we have set

$$\tau^{(\ell)}(\gamma_{\mathbf{i}}z) \stackrel{\text{def}}{=} \tau(\gamma_{i_1 \dots i_\ell} z) + \tau(\gamma_{i_2 \dots i_\ell} z) + \dots + \tau(\gamma_{i_\ell} z).$$

Notice that there exists $M > 0$ independent of ℓ such that for all z, \mathbf{i} above we have $|\tau^{(\ell)}(\gamma_{\mathbf{i}}z)| \leq \ell M$. We will then use the following basic bound, valid for all $z_1, z_2 \in \mathbf{C}$:

$$|e^{z_1} - e^{z_2}| \leq |z_1 - z_2| e^{\max\{\text{Re}(z_1), \text{Re}(z_2)\}},$$

which yields

$$\left| e^{-s\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} - e^{-s_0\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} \right| \leq M\ell |s - s_0| e^{\ell M \max\{|s_0|, |s|\}}.$$

Using unitarity of the representation ρ_n^0 and triangle inequality leads to the very crude bound

$$\left\| \sum_{\mathbf{i} \rightarrow j, |\mathbf{i}|=\ell} \left(e^{-s\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} - e^{-s_0\tau^{(\ell)}(\gamma_{\mathbf{i}}z)} \right) \rho_n^0(\gamma_{\mathbf{i}}^{-1}) f(\gamma_{\mathbf{i}}z) \right\|_{V_n^0} \leq M\ell |s - s_0| e^{\ell M \max\{|s_0|, |s|\}} \# \mathcal{W}_\ell \left(\sup_{\mathbf{i} \rightarrow j} \sup_{z \in D_j} \|f(\gamma_{\mathbf{i}}z)\|_{V_n^0} \right),$$

which then gives

$$\|\mathcal{L}_{s, \rho_n^0}^\ell(f) - \mathcal{L}_{s_0, \rho_n^0}^\ell(f)\|^2 \leq m(\mathbf{D}) (M\ell |s - s_0|)^2 e^{2\ell M \max\{|s_0|, |s|\}} (\# \mathcal{W}_\ell)^2 \left(\sup_{\mathbf{i} \rightarrow j} \sup_{z \in D_j} \|f(\gamma_{\mathbf{i}}z)\|_{V_n^0}^2 \right).$$

By the reproducing property of Bergman's kernel, and since all $\gamma_{\mathbf{i}}$ map uniformly each D_j in a compact subset of \mathbf{D} , we deduce that there exists a constant $C > 0$ uniform in ℓ such that

$$\sup_{\mathbf{i} \rightarrow j} \sup_{z \in D_j} \|f(\gamma_{\mathbf{i}}z)\|_{V_n^0}^2 \leq C \|f\|_{\mathcal{H}(\mathbf{D}, V_n^0)}^2.$$

The proof is now done if we use the fact that s, s_0 are in a compact set. \square

We then define the following operators $A_i(s) : \mathcal{H}(\mathbf{D}, \mathbf{C}) \rightarrow \mathcal{H}(\mathbf{D}, \mathbf{C})$ by

$$A_i(s)[f](x) \stackrel{\text{def}}{=} \mathbf{1}\{x \notin D_i\} e^{-s\tau(\gamma_{\bar{i}}(x))} f(\gamma_{\bar{i}}(x)). \quad (2.9)$$

Being a composition operator which maps admissible discs D_i compactly into \mathbf{D} , it is easy to see that each operator $A_i(s)$ is trace class on $\mathcal{H}(\mathbf{D}; \mathbf{C})$. If $\mathbf{i} = (i_1, i_2, \dots, i_\ell)$ is an admissible word, we can then define the following composition

$$A(s)_{\mathbf{i}} \stackrel{\text{def}}{=} A_{i_1}(s) A_{i_2}(s) \dots A_{i_\ell}(s)[f] = \mathbf{1}\{x \notin D_{i_1}\} e^{-s\tau^{(\ell)}(\gamma_{\bar{\mathbf{i}}}x)} f \circ \gamma_{\bar{\mathbf{i}}}(x),$$

where we have set for all admissible word \mathbf{j} of length ℓ , $\tau^{(\ell)}(\gamma_{\mathbf{j}}z) = \tau(\gamma_{j_1 \dots j_\ell} z) + \tau(\gamma_{j_2 \dots j_\ell} z) + \dots + \tau(\gamma_{j_\ell} z)$. It will be useful in the following to have a kernel formula for the action of the adjoint operator $A(s)_{\mathbf{i}}^*$.

Lemma 2.6. *Using the above notations, we have for all $f \in \mathcal{H}(\mathbf{D}, \mathbf{C})$,*

$$A(s)_{\mathbf{i}}^*[f](x) = \int_{\mathbf{D} \setminus D_{i_1}} e^{-\overline{s\tau^{(\ell)}(\gamma_{\bar{\mathbf{i}}}z)}} B_{\mathbf{D}}(x, \gamma_{\bar{\mathbf{i}}}z) f(z) dm(z).$$

Proof. Using the reproducing kernel property, we have for all $f, g \in \mathcal{H}(\mathbf{D}, \mathbf{C})$,

$$\langle A(s)_{\mathbf{i}} f, g \rangle = \int_{\mathbf{D}} \mathbf{1}\{z \notin D_{i_1}\} e^{-s\tau^{(\ell)}(\gamma_{\bar{\mathbf{i}}}z)} \int_{\mathbf{D}} B_D(\gamma_{\bar{\mathbf{i}}}z, w) f(w) dm(w) \overline{g(z)} dm(z).$$

We observe that since there exists a compact set $K \subset \mathbf{D}$ such that for all $z \notin D_{i_1}$, $\gamma_{\bar{\mathbf{i}}}(z) \in K$, we have (by using the explicit formula for the Bergman kernel)

$$\sup_{z, w \in \mathbf{D}} |B_D(\gamma_{\bar{\mathbf{i}}}z, w) \mathbf{1}\{z \notin D_{i_1}\}| < \infty.$$

Therefore we have

$$\int_{\mathbf{D}} \int_{\mathbf{D}} \mathbf{1}\{z \notin D_{i_1}\} \left| e^{-s\tau^{(\ell)}(\gamma_{\bar{\mathbf{i}}}z)} \right| |B(\gamma_{\bar{\mathbf{i}}}z, w) f(w) g(z)| dm(z) dm(w) < \infty,$$

and we can use Fubini's theorem to write

$$\langle A(s)_{\mathbf{i}} f, g \rangle = \int_{\mathbf{D}} f(w) \overline{\int_{\mathbf{D}} \mathbf{1}\{z \notin D_{i_1}\} e^{-\overline{s\tau^{(\ell)}(\gamma_{\bar{\mathbf{i}}}z)}} B_{\mathbf{D}}(w, \gamma_{\bar{\mathbf{i}}}z) g(z) dm(z) dm(w)},$$

where we have used the fact that $B_{\mathbf{D}}(z, w) = \overline{B_{\mathbf{D}}(w, z)}$ and the proof is done. \square

Notice that we have $A_i^*(s) \neq A_{\bar{i}}(s)$, which is a source of non-selfadjointness issues and (besides the obvious infinite dimensional setting) one of the main differences with [BC19], where a symmetry hypothesis is assumed. See §§4.1 for a more detailed discussion of these differences to [BC19].

In this paper, we will have to deal with integral operators acting on $\mathcal{H}(\mathbf{D}, \mathbf{C})$ (or their vector valued extensions). We need a simple criterion to compute their traces, which is as

follows.

Lemma 2.7. *Let $T_{\mathcal{K}} : \mathcal{H}(\mathbf{D}, \mathbf{C}) \rightarrow \mathcal{H}(\mathbf{D}, \mathbf{C})$ be an integral operator whose kernel $\mathcal{K}(z, w)$ satisfies the following properties: assume that $\mathcal{K} \in C^0(\overline{\mathbf{D} \times \mathbf{D}})$, and that for all $w \in \mathbf{D}$, the map $z \mapsto \mathcal{K}(z, w)$ is holomorphic and extends to a C^2 function on the boundary $\partial \mathbf{D}$. Then, if $T_{\mathcal{K}}$ is trace class, we have the identity*

$$\mathrm{tr}(T_{\mathcal{K}}) = \int_{\mathbf{D}} \mathcal{K}(z, z) dm(z).$$

Proof. It is enough to work in the unit disc \mathbb{D} to simplify. Let $(e_p)_{\{p \in \mathbf{N}\}}$ be the Hilbert basis of $\mathcal{H}(\mathbb{D}, \mathbf{C})$ given explicitly by

$$e_p(z) = \sqrt{\frac{p+1}{\pi}} z^p.$$

Because $T_{\mathcal{K}}$ is assumed to be trace class, we have

$$\mathrm{tr}(T_{\mathcal{K}}) = \sum_{p=0}^{\infty} \langle T_{\mathcal{K}} e_p, e_p \rangle = \sum_p \int_{\mathbb{D}} \int_{\mathbb{D}} \mathcal{K}(z, w) e_p(w) dm(w) \overline{e_p(z)} dm(z),$$

which by Fubini can be rewritten as

$$\mathrm{tr}(T_{\mathcal{K}}) = \sum_p \int_{\mathbb{D}} \int_{\mathbb{D}} \mathcal{K}(z, w) \overline{e_p(z)} dm(z) e_p(w) dm(w).$$

Since $z \mapsto \mathcal{K}(z, w)$ is holomorphic on the unit disc, we know that we have the convergent Taylor expansion

$$\mathcal{K}(z, w) = \sum_{p=0}^{\infty} a_p(w) z^p,$$

where for all $0 < r < 1$, we can write

$$a_p(w) = \frac{1}{2\pi} \int_0^{2\pi} \mathcal{K}(re^{i\theta}, w) (re^{i\theta})^{-p} d\theta.$$

because $z \mapsto \mathcal{K}(z, w)$ is actually continuous up to the boundary we can write

$$a_p(w) = \frac{1}{2\pi} \int_0^{2\pi} \mathcal{K}(e^{i\theta}, w) (e^{i\theta})^{-p} d\theta,$$

and use the C^2 regularity to integrate by parts two times which yields uniformly for all $w \in \overline{\mathbb{D}}$,

$$|a_p(w)| = O(p^{-2}).$$

We now notice that

$$\int_{\mathbb{D}} \mathcal{K}(z, w) \overline{e_p(z)} dm(z) = a_p(w) \sqrt{\frac{\pi}{p+1}},$$

which shows that uniformly for all $w \in \overline{\mathbb{D}}$,

$$\left| \int_{\mathbb{D}} \mathcal{K}(z, w) \overline{e_p(z)} dm(z) \right| |e_p(w)| = O(p^{-2}),$$

and the series

$$\sum_p \int_{\mathbb{D}} \mathcal{K}(z, w) \overline{e_p(z)} dm(z) e_p(w)$$

are therefore uniformly convergent (with limit $\mathcal{K}(w, w)$) on $\overline{\mathbb{D}}$, which allows to write

$$\text{tr}(T_{\mathcal{K}}) = \int_{\mathbb{D}} \left(\sum_{p=0}^{\infty} \int_{\mathbb{D}} \mathcal{K}(z, w) \overline{e_p(z)} dm(z) e_p(w) \right) dm(w) = \int_{\mathbb{D}} \mathcal{K}(w, w) dm(w),$$

and the proof is done. \square

Lemma 2.8. *Let \mathcal{K} be any compact subset of \mathbf{C} . There is $C = C(\mathcal{K}) > 0$ such that for any $s = r + it \in \mathcal{K}$, all $m > 0$ and all $2m$ -tuples of admissible sequences $\mathbf{i}^1, \dots, \mathbf{i}^m, \mathbf{j}^1, \dots, \mathbf{j}^m$, we have*

$$\left| \text{tr} \left(A(s)_{\mathbf{i}^1} A(s)_{\mathbf{j}^1}^* A(s)_{\mathbf{i}^2} A(s)_{\mathbf{j}^2}^* \cdots A(s)_{\mathbf{i}^m} A(s)_{\mathbf{j}^m}^* \right) \right| \leq C^m \prod_{k=1}^m \Upsilon_{\mathbf{i}^k}^r \Upsilon_{\mathbf{j}^k}^r.$$

Proof. Given an admissible word \mathbf{i} , we denote by $\mathcal{K}_{\mathbf{i}}(z, w)$ the integral kernel of $A(s)_{\mathbf{i}}$ which is given by

$$\mathcal{K}_{\mathbf{i}}(z, w) = \mathbf{1}\{z \notin D_{i_1}\} e^{-s\tau(|\mathbf{i}|)(\gamma_{\bar{\mathbf{i}}} z)} B_{\mathbf{D}}(\gamma_{\bar{\mathbf{i}}} z, w).$$

Similarly we will denote by $\mathcal{K}_{\mathbf{i}}^*(z, w)$ the integral kernel of $A(s)_{\mathbf{i}}^*$, which is

$$\mathcal{K}_{\mathbf{i}}^*(z, w) = \mathbf{1}\{w \notin D_{i_1}\} e^{-\overline{s\tau(|\mathbf{i}|)(\gamma_{\bar{\mathbf{i}}} w)}} B_{\mathbf{D}}(z, \gamma_{\bar{\mathbf{i}}} w).$$

An important observation is due to the fact that each $\gamma_{\bar{\mathbf{i}}}$ maps $\mathbf{D} \setminus D_{i_1}$ uniformly and compactly into \mathbf{D} , one can use the explicit formula for the Bergman kernel to prove that there exists $C_0 > 0$ such that for all admissible \mathbf{i} ,

$$\sup_{z, w \in \mathbf{D}} |\mathbf{1}\{z \notin D_{i_1}\} B(\gamma_{\bar{\mathbf{i}}} z, w)| \leq C_0 \text{ and } \sup_{z, w \in \mathbf{D}} |\mathbf{1}\{w \notin D_{i_1}\} B(z, \gamma_{\bar{\mathbf{i}}} w)| \leq C_0.$$

Because each kernel $\mathcal{K}_{\mathbf{i}}(z, w)$, $\mathcal{K}_{\mathbf{j}}^*(z, w)$ is bounded on $\mathbf{D} \times \mathbf{D}$ (by the above observation), we can use Fubini's theorem to compute the kernel of

$$A(s)_{\mathbf{i}^1} A(s)_{\mathbf{j}^1}^* A(s)_{\mathbf{i}^2} A(s)_{\mathbf{j}^2}^* \cdots A(s)_{\mathbf{i}^m} A(s)_{\mathbf{j}^m}^*,$$

which is given by

$$\mathcal{K}_{\mathbf{i}^1, \mathbf{j}^1, \dots, \mathbf{i}^m, \mathbf{j}^m}(z, w) = \int_{\mathbf{D}^{2m-1}} \mathcal{K}_{\mathbf{i}^1}(z, w_1) \mathcal{K}_{\mathbf{j}^1}^*(w_1, w_2) \mathcal{K}_{\mathbf{i}^2}(w_2, w_3) \cdots \mathcal{K}_{\mathbf{j}^m}^*(w_{2m-1}, w) dm(w_1) \cdots dm(w_{2m-1}).$$

We can then check that $z \mapsto \mathcal{K}_{\mathbf{i}^1, \mathbf{j}^1, \dots, \mathbf{i}^m, \mathbf{j}^m}(z, w)$ is holomorphic with an analytic extension to $\partial \mathbf{D}$. Being a product of trace class operators, $A(s)_{\mathbf{i}^1} A(s)_{\mathbf{j}^1}^* A(s)_{\mathbf{i}^2} A(s)_{\mathbf{j}^2}^* \cdots A(s)_{\mathbf{i}^m} A(s)_{\mathbf{j}^m}^*$ is therefore trace class and we can use Lemma 2.7 to obtain (again rearranging the order of integration by Fubini)

$$\begin{aligned} & \operatorname{tr} \left(A(s)_{\mathbf{i}^1} A(s)_{\mathbf{j}^1}^* A(s)_{\mathbf{i}^2} A(s)_{\mathbf{j}^2}^* \cdots A(s)_{\mathbf{i}^m} A(s)_{\mathbf{j}^m}^* \right) \\ &= \int_{\mathbf{D}^{2m}} \mathcal{K}_{\mathbf{i}^1}(w_{2m}, w_1) \mathcal{K}_{\mathbf{j}^1}^*(w_1, w_2) \cdots \mathcal{K}_{\mathbf{i}^m}(w_{2m-2}, w_{2m-1}) \mathcal{K}_{\mathbf{j}^m}^*(w_{2m-1}, w_{2m}) dm(w_1) \cdots dm(w_{2m}). \end{aligned}$$

We now recall that for all $z \in \mathbf{D} \setminus D_{i_1}$,

$$e^{-s\tau^{(\mathbf{i}^1)}(\gamma_{\bar{\mathbf{i}}^1} z)} = (\gamma'_{\bar{\mathbf{i}}^1}(z))^s = e^{s \log \gamma'_{\bar{\mathbf{i}}^1}(z)},$$

where $\log(w) = \log|w| + i \arg(w)$ is the principal branch of the complex logarithm on $\mathbf{C} \setminus (-\infty, 0]$. Using (2.6), this leads to the bound for $s \in \mathcal{K}$

$$\left| (\gamma'_{\bar{\mathbf{i}}^1}(z))^s \right| \leq e^{\operatorname{Re}(s \log \gamma'_{\bar{\mathbf{i}}^1}(z))} \leq C_1(\mathcal{K}) \Upsilon_{\bar{\mathbf{i}}^1}^{\operatorname{Re}(s)},$$

where $C_1(\mathcal{K}) > 0$. Using the mirror estimate (2.5), we end up with (up to a slight change of constant)

$$\left| (\gamma'_{\bar{\mathbf{i}}^1}(z))^s \right| \leq C_2(\mathcal{K}) \Upsilon_{\bar{\mathbf{i}}^1}^{\operatorname{Re}(s)}.$$

We have therefore if $\operatorname{Re}(s) = r$

$$\left| \operatorname{tr} \left(A(s)_{\mathbf{i}^1} A(s)_{\mathbf{j}^1}^* A(s)_{\mathbf{i}^2} A(s)_{\mathbf{j}^2}^* \cdots A(s)_{\mathbf{i}^m} A(s)_{\mathbf{j}^m}^* \right) \right| \leq C_0^{2m} (C_2(\mathcal{K}))^{2m} (\operatorname{Vol}(\mathbf{D}))^{2m} \prod_{k=1}^m \Upsilon_{\mathbf{i}^k}^r \Upsilon_{\mathbf{j}^k}^r,$$

which concludes the proof. \square

3 Random permutations and random graphs

3.1 Symmetric random permutations

We write $[n] \stackrel{\text{def}}{=} \{1, \dots, n\}$ and we write S_n for the symmetric group of permutations of $[n]$. We view $d \geq 2$ as a fixed integer as in §§2.1. As before, for $i \in [2d]$ let $\bar{i} \stackrel{\text{def}}{=} i + d \bmod 2d$ so $\bar{i} \in [2d]$. We say that a tuple in $(S_n)^{2d}$ is *symmetric* if $\sigma_{\bar{i}} = \sigma_i^{-1}$. To obtain symmetric random permutations we choose $\sigma_1, \dots, \sigma_d$ independently and uniformly in S_n . (Of course, choosing these plays the same role as in §§2.2 and hence is the same thing as choosing our random cover X_ϕ .)

We let $\sigma_{\bar{i}} = \sigma_i^{-1}$ to extend the indices of σ_i to $[2d]$ and we write S_i for the matrix representing σ_i as a 0-1 matrix. We also let

$$\underline{S}_i \stackrel{\text{def}}{=} S_i - \frac{1}{n} \mathbf{1} \otimes \mathbf{1},$$

this is just the matrix of σ_i acting in V_n^0 .

3.2 Graphs and paths

First following [BC19, Def. 6] we define the category of graphs that we work with.

Definition 3.1. A colored edge is an equivalence class of element (x, i, y) of $[n] \times [d] \times [n]$ with respect to the equivalence relation generated by

$$(x, i, y) \sim (y, \bar{i}, x). \quad (3.1)$$

A colored graph is a pair consisting of a vertex set $V \subset [n]$ and a set of colored edges E as above. It is clear that any colored graph has underlying multigraph obtained by forgetting colors. When we refer to paths, cycles, etc, in colored graphs, they are just paths and cycles in the underlying multigraph.

For symmetric permutations $(\sigma_1, \dots, \sigma_{2d})$ in $(S_n)^{2d}$, G^σ is the colored graph with vertex set $[n]$ and colored edges given by $[x, i, y]$ such that $\sigma_i(x) = y$, modulo the equivalence relation on colored edges.

We also introduce a way to talk about paths and the graphs they trace out.

Definition 3.2. The set E is the collection of pairs $e = (x, i)$ with $x \in [n]$ and $i \in [2d]$. Each of these be thought of as a half-edge emanating from x and labeled by i . A *path* of length k is a sequence in E^k . For $k \in \mathbf{N}$ let $\gamma = (\gamma_1, \dots, \gamma_k) \in E^k$ be a path, and write each $\gamma_t = (x_t, i_t) \in [n] \times [2d]$.

- Let $V_\gamma = \{x_t : 1 \leq t \leq k\}$ and $E_\gamma = \{[x_t, i_t, x_{t+1}] : 1 \leq t \leq k-1\}$. We denote by G_γ the colored graph with vertex set V_γ and colored edges E_γ modulo the equivalence relation (3.1).
- A path is non-backtracking if for each $1 \leq t \leq k-1$ as above, $i_t \neq \bar{i}_{t+1}$. We write Γ^k for the subset of non-backtracking paths in E^k . For $e, f \in E$ we write $\Gamma_{ef}^k \subset \Gamma^k$ for the non-backtracking paths with $\gamma_1 = e, \gamma_k = f$.

Remark 3.3. Note that in Definition 3.2 the paths, etc have nothing a priori to do with G^σ . Rather, their role is as objects that could *potentially* appear in G^σ .

Definition 3.4. Given a non-backtracking path $\gamma = ((x_1, i_1), \dots, (x_k, i_k)) \in \Gamma^k$, the *color sequence* of the path is

$$\mathbf{i}(\gamma) \stackrel{\text{def}}{=} (i_1, \dots, i_k).$$

3.3 Tangles

One important notion in the works of Bordenave-Collins [BC19], originating in Friedman [Fri08], is *tangles*. These relate to events of small probability that make large, argument-destroying, contributions to the expected value of traces. The following definition is [BC19, Def. 19]

Definition 3.5. Let H be a colored graph. H is *tangle-free* if it contains at most one cycle. For any vertex x of H , let $(H, x)_\ell$ denote the subgraph of H containing all colored edges and vertices that belong to a path of length at most ℓ beginning at x . H is ℓ -tangle-free if for any vertex x , $(H, x)_\ell$ is tangle-free, and H is ℓ -tangled otherwise. We say that a path $\gamma \in E^k$ is tangle-free if G_γ is tangle-free. We write F^k (resp. F_{ef}^k) for the collection of tangle-free non-backtracking paths in Γ^k (resp. Γ_{ef}^k).

The following lemma appearing in [BC19, Lemma 23] gives the probability that G^σ is ℓ -tangled when (σ_i) are symmetric random permutations in $(S_n)^{2d}$.

Lemma 3.6. *There is a constant $c > 0$ such that for (σ_i) symmetric random permutations in $(S_n)^{2d}$ and $1 \leq \ell \leq \sqrt{n}$, the probability that G^σ is ℓ -tangled is at most*

$$c \frac{\ell^3 (2d-1)^{4\ell}}{n}.$$

4 Non-backtracking operator and path decomposition

4.1 Conjugation between transfer operator and non-backtracking operator

The purpose of this section is to show how the transfer operator, after a unitary conjugation, is roughly of the form as the non-backtracking operator of [BC19]. This conjugation is not essential for the argument and merely makes it easier to read the current work alongside [BC19].

Recall from §2.3 that our transfer operator

$$\mathcal{L}_{s, \rho_n}[f](x) \stackrel{\text{def}}{=} \sum_{i \in \mathcal{A}} e^{-s\tau(\gamma_i(x))} \rho_n(\gamma_i)^{-1} f(\gamma_i(x))$$

acts on $\mathcal{H}(\mathbf{D}, V_n)$ and the restriction of this operator to $\mathcal{H}(\mathbf{D}, V_n^0)$ coincides with $\mathcal{L}_{s, \rho_n^0}$; we aim to prove for certain values of s that $\mathcal{L}_{s, \rho_n^0}$ has no eigenvalue 1 (cf. Proposition 2.2).

Let

$$K_0 \stackrel{\text{def}}{=} \mathcal{H}(\mathbf{D}) \otimes V_n^0 \subset \mathcal{H}(\mathbf{D}) \otimes \ell^2([n]).$$

We let S_i denote the 0-1 matrix of $\sigma_i = \phi(\gamma_i)$ as in §2.2 (recall also that ρ_n and ϕ are coupled). For each element $e = (x, i) \in E$ we consider the subspace

$$\mathcal{H}_e \stackrel{\text{def}}{=} \mathcal{H}(D_{\bar{i}}) \otimes \mathbf{C}x \subset \mathcal{H}(\mathbf{D}) \otimes \ell^2([n]).$$

(The reason \bar{i} appears here is to make things line up with Bordenave-Collins machinery). In $\ell^2([n])$, we write y for the indicator function of y (this corresponds to identification of $\ell^2([n])$ with formal complex linear combinations of $[n]$). This gives

$$\mathcal{H}(\mathbf{D}) \otimes \ell^2([n]) = \bigoplus_{e \in E} \mathcal{H}_e.$$

Write \mathbf{p}_e for the orthogonal projection onto \mathcal{H}_e . For any endomorphism M of $\mathcal{H}(\mathbf{D}) \otimes \ell^2([n])$ and for each $e, f \in E$ we write

$$M_{ef} \stackrel{\text{def}}{=} \mathbf{p}_e M \mathbf{p}_f.$$

Clearly M is determined by the values M_{ef} . Let $\mathcal{E}_{xy} : \ell^2([n]) \rightarrow \ell^2([n])$ be the map with

$$(\mathcal{E}_{xy})(y') = \delta_{yy'} x$$

and $\delta_{yy'}$ is the Kronecker delta.

Recall the definition of operators $A_j(s)$ from (2.9). For $e = (x, i)$ and $f = (y, j)$ we define

$$B(s)_{ef} \stackrel{\text{def}}{=} \mathbf{1}\{i \neq \bar{j}\} (S_i)_{xy} \mathbf{p}_e [A_j(s) \otimes \mathcal{E}_{xy}] \mathbf{p}_f; \quad (4.1)$$

this defines $B(s) \in \text{End}(\mathcal{H}(\mathbf{D}) \otimes \ell^2([n]))$. The operator $B(s)$ is in a similar form to the operator B defined by Bordenave and Collins in [BC19, (12)] (cf. also the display equation following [BC19, (12)]). However our $B(s)$ and Bordenave-Collins' B differ in the following ways:

- our $\mathcal{H}(\mathbf{D})$ is the analog of $\mathbf{C}^r \otimes \ell^2([2d])$, however, $\mathcal{H}(\mathbf{D})$ does not split as a tensor product in this way.
- Also, our $A_j(s)$ is slightly more general (in the Bordenave-Collins setting, it is equivalent to considering $a_j \in \text{End}(\mathbf{C}^r \otimes \ell^2([2d]))$ rather than $a_j \in \text{End}(\mathbf{C}^r)$).
- each $A_j(s)$ acts on a infinite dimensional Hilbert space $\mathcal{H}(\mathbf{D})$.
- **most critically**, the symmetry condition $A_i = A_i^*$ that Bordenave-Collins assume does not hold here.

These differences, and *especially the lack of the symmetry condition*, have to be worked around in the sequel.

Proposition 4.1. *There is a unitary operator $U : \mathcal{H}(\mathbf{D}, V_n) \rightarrow \mathcal{H}(\mathbf{D}) \otimes \ell^2([n])$ such that*

1. *U conjugates the holomorphic family of trace class operators \mathcal{L}_{s, ρ_n} to the family of operators $B(s)$.*
2. *U conjugates the holomorphic family of trace class operators $\mathcal{L}_{s, \rho_n^0}$ to the family of operators $B(s)|_{K_0}$.*

As such, $B(s)$ and $B(s)|_{K_0}$ are holomorphic families of trace class operators and $\|\mathcal{L}_{s, \rho_n^0}^\ell\| = \|B(s)^\ell|_{K_0}\|$ for all $s \in \mathbf{C}$ and $\ell \in \mathbf{N}$.

Proof. First of all, by the obvious identification

$$\mathcal{H}(\mathbf{D}, V_n) \cong \mathcal{H}(\mathbf{D}) \otimes \ell^2([n])$$

the operator \mathcal{L}_{s,ρ_n} is conjugated to $b(s) \in \text{End}(\mathcal{H}(\mathbf{D}) \otimes \ell^2([n]))$,

$$b(s) \stackrel{\text{def}}{=} \sum_i A_i(s) \otimes \sigma_i.$$

Consider the unitary operator $Q \in \text{End}(\mathcal{H}(\mathbf{D}) \otimes \ell^2([n]))$ defined for $e = (x, i)$ and $f = (y, j)$ by

$$Q_{ef} \stackrel{\text{def}}{=} \mathfrak{p}_e \left(\text{Id}_{\mathcal{H}(D_{\bar{j}})} \otimes \sigma_j^{-1} \right) \mathfrak{p}_f.$$

Then for $F \otimes y \in \mathcal{H}_f$, $f = (y, j)$, $F \in \mathcal{H}(D_{\bar{j}})$, using $A_i(s)F = 0$ unless $i = j$ we obtain

$$\begin{aligned} [Q^{-1}b(s)Q]_{ef}[F \otimes y] &= \mathfrak{p}_e Q b(s) [F \otimes \sigma_j^{-1}(y)] \\ &= \mathfrak{p}_e Q [(A_j(s)F) \otimes y] \\ &= \mathfrak{p}_e (A_j(s)F \otimes \sigma_i^{-1}(y)) \\ &= \mathbf{1}\{i \neq \bar{j}\} \mathbf{1}\{\sigma_i(x) = y\} \mathfrak{p}_e (A_j(s)F \otimes x) \\ &= B(s)_{ef}[F \otimes y] \end{aligned}$$

proving that $Q^{-1}b(s)Q = B(s)$. It is also not hard to see that $B(s)$ leaves K_0 invariant and Q also conjugates the restriction $B(s)|_{K_0}$ to \mathcal{L}_{s,ρ_n^0} . Finally, noting that the conjugation between \mathcal{L}_{s,ρ_n} and $B(s)$ does not depend on s completes the proof. \square

4.2 Bordenave's path decomposition

Here we perform a path decomposition method that originates in work of Bordenave [Bor]. Recall from Definition 3.5 that $F^{\ell+1}$ is the collection of tangle-free paths in $\Gamma^{\ell+1}$. For $1 \leq k \leq \ell$ we write $F_k^{\ell+1}$ for the collection of paths γ in $\Gamma^{\ell+1}$ such that the first k half-edges of γ (in Γ^k) form a tangle-free path and the last $\ell - k + 1$ half-edges (in $\Gamma^{\ell-k+1}$) also form a tangle-free path.

Recall the definition of $\mathbf{i}(\gamma)$ from Definition 3.4 and the notation $\hat{\mathbf{i}} = (i_2, \dots, i_n)$ from §§2.1. As in [BC19, pg. 834] we obtain with $e = (x, i)$ and $f = (y, j)$

$$B(s)_{ef}^\ell = \mathfrak{p}_e \left(\sum_{\gamma \in \Gamma_{ef}^{\ell+1}} (A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \quad (4.2)$$

For $\ell \in \mathbf{N}$ we define as in [BC19, (32), (34)]

$$\begin{aligned} \underline{B}(s)^{(\ell)} &\stackrel{\text{def}}{=} \sum_{\gamma=((x_1, i_1), \dots, (x_{\ell+1}, i_{\ell+1})) \in F^{\ell+1}} \mathfrak{p}_{(x_1, i_1)}[A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{x_1 x_{\ell+1}}] \mathfrak{p}_{(x_{\ell+1}, i_{\ell+1})} \left(\prod_{t=1}^{\ell} (\underline{S}_{i_t})_{x_t x_{t+1}} \right), \\ R_k(s)^{(\ell)} &\stackrel{\text{def}}{=} \sum_{\gamma=((x_1, i_1), \dots, (x_{\ell+1}, i_{\ell+1})) \in F_k^{\ell+1} \setminus F^{\ell+1}} \mathfrak{p}_{(x_1, i_1)}[A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{x_1 x_{\ell+1}}] \mathfrak{p}_{(x_{\ell+1}, i_{\ell+1})} \left(\prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \right) \left(\prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right). \end{aligned}$$

Remark 4.2. Since each $A(s)_i$ is trace class (see §§2.4), both $\underline{B}(s)^{(\ell)}$ and $R_k(s)^{(\ell)}$ are trace class operators for any $\ell \in \mathbf{N}$.

The following is proved by Bordenave and Collins in the course of proving [BC19, Lemma 20] (see the last display equation before [BC19, Lemma 20]).

Lemma 4.3. *Let $\ell \geq 1$ be an integer and suppose G^σ is ℓ -tangle free. Then for all $F \in K_0$,*

$$B(s)^\ell F = \underline{B}(s)^{(\ell)} - \frac{1}{n} R_k(s)^{(\ell)} F. \quad (4.3)$$

Hence

$$\|B(s)^\ell|_{K_0}\| \leq \|\underline{B}(s)^{(\ell)}\| + \frac{1}{n} \sum_{k=1}^{\ell} \|R_k(s)^{(\ell)}\|.$$

Proof. It clearly suffices to prove the first statement. This is similar to [BC19, pgs. 836-837] but there are subtle differences so we give the details for completeness. Let $e = (x, i)$ and $f = (y, j)$ in E . Beginning with the expression (4.2) for $B(s)_{ef}^\ell$, on the event that G^σ is ℓ -tangle free we can write

$$\begin{aligned} B(s)_{ef}^\ell &= \mathfrak{p}_e \left(\sum_{\gamma \in \Gamma_{ef}^{\ell+1}} (A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{\ell} (\underline{S}_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \\ &= \mathfrak{p}_e \left(\sum_{\gamma \in F_{ef}^{\ell+1}} (A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{\ell} (\underline{S}_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \end{aligned} \quad (4.4)$$

$$+ \frac{1}{n} \sum_{k=1}^{\ell} \mathfrak{p}_e \left(\sum_{\gamma \in F_{ef}^{\ell+1}} (A(s)_{\hat{\mathbf{i}}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f, \quad (4.5)$$

where the last equality used

$$\prod_{t=1}^{\ell} (S_{i_t})_{x_t x_{t+1}} = \prod_{t=1}^{\ell} (\underline{S}_{i_t})_{x_t x_{t+1}} + \frac{1}{n} \sum_{k=1}^{\ell} \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}}.$$

The operator on line (4.4) is $\underline{B}(s)_{ef}^{(\ell)}$ so it suffices to describe the operator on line (4.5). We write

$$\begin{aligned} & \mathfrak{p}_e \left(\sum_{\gamma \in F_{ef}^{\ell+1}} (A(s)_{\mathfrak{i}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \\ &= \mathfrak{p}_e \left(\sum_{\gamma \in F_{k,ef}^{\ell+1}} (A(s)_{\mathfrak{i}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \end{aligned} \quad (4.6)$$

$$- \mathfrak{p}_e \left(\sum_{\gamma \in F_{k,ef}^{\ell+1} \setminus F_{ef}^{\ell+1}} (A(s)_{\mathfrak{i}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_e. \quad (4.7)$$

The terms in line (4.7) give the required contributions to $-\frac{1}{n}R_k(s)^{(\ell)}$ in (4.3) so it suffices to show that the operator M defined by

$$M_{ef} \stackrel{\text{def}}{=} (4.6)$$

kills every element of K_0 .

Let $J : \mathcal{H}(\mathbf{D}) \otimes \ell^2([n]) \rightarrow \mathcal{H}(\mathbf{D}) \otimes \ell^2([n])$ denote the operator defined for $e = (x, i)$, $f = (y, j)$

$$J_{ef} = \mathbf{1}\{i \neq \bar{j}\} \mathfrak{p}_e[A_j(s) \otimes \mathcal{E}_{xy}] \mathfrak{p}_f$$

(contrast this to (4.1)). Note that if $F = F_0 \otimes v \in K_0$, with $F_0 \in \mathcal{H}(D_{\bar{j}})$ and $v = (v_1, \dots, v_n) \in V_n^0$, $\sum_{y \in [n]} v_y = 0$, then

$$\begin{aligned} J[F_0 \otimes v] &= \sum_{\substack{e=(x,i) \\ i \neq \bar{j}}} \sum_{y \in [n]} \mathfrak{p}_e[A_j(s) \otimes \mathcal{E}_{xy}] \mathfrak{p}_{(y,j)}(F_0 \otimes v) \\ &= \sum_{\substack{e=(x,i) \\ i \neq \bar{j}}} \sum_{y \in [n]} v_y \mathfrak{p}_e[A_j(s) \otimes \mathcal{E}_{xy}](F_0 \otimes y) \\ &= \sum_{\substack{e=(x,i) \\ i \neq \bar{j}}} [(A_j(s)F_0) |_{D_{\bar{i}} \otimes x}] \sum_{y \in [n]} v_y = 0. \end{aligned}$$

This implies $K_0 \subset \ker J$.

Since we still assume G^σ is ℓ -tangle-free, and hence k -tangle free for all $k \in [\ell]$, we have

$$\begin{aligned} M_{ef} &= \mathfrak{p}_e \left(\sum_{\gamma \in F_{k,ef}^{\ell+1}} (A(s)_{\mathfrak{i}(\gamma)} \otimes \mathcal{E}_{xy}) \prod_{t=1}^{k-1} (\underline{S}_{i_t})_{x_t x_{t+1}} \prod_{t=k+1}^{\ell} (S_{i_t})_{x_t x_{t+1}} \right) \mathfrak{p}_f \\ &= (\tilde{M}JB(s)^{\ell-k})_{ef} \end{aligned}$$

for some operator \tilde{M} . Since $B(s)$ preserves K_0 , this implies K_0 is in the kernel of M as required. \square

Lemma 4.3 tells us that bounds on both $\|\underline{B}(s)^{(\ell)}\|$ and $\|R_k(s)^{(\ell)}\|$ that hold with high probability can be coupled with the fact that G^σ is ℓ -tangle-free with high-probability (for ℓ in a suitable range) to establish bounds for $\|B(s)^\ell|_{K_0}\|$ that hold with high probability. A high probability bound for $\|\underline{B}(s)^{(\ell)}\|$ is the subject of §5.1 and high probability bounds for $\|R_k(s)^{(\ell)}\|$ are the subject of §5.2.

5 High trace method

5.1 Norm of $\underline{B}(s)^{(\ell)}$

The main result of this §5.1 is the following.

Proposition 5.1. *Let \mathcal{K} be a compact subset of $\{s : \operatorname{Re}(s) > \frac{\delta}{2}\}$. There is $c = c(\mathcal{K}) > 1$, $\rho = \rho(\mathcal{K}) < 1$ such that for all $s \in \mathcal{K}$, for all $1 \leq \ell \leq \log n$ the event*

$$\|\underline{B}(s)^{(\ell)}\| \leq (\log n)^{20} \rho^\ell$$

holds with probability at least $1 - c \exp\left(-\frac{\ell \log n}{c \log \log n}\right)$.

Here we follow the structure of [BC19, §§4.4.1] up to Proposition 5.6, when we make our departure. The method used here to bound $\|\underline{B}(s)^{(\ell)}\|$ is the ‘high-trace’ method originating in work of Füredi and Komlós [FK81].

For $\ell, m \in \mathbf{N}$ we let $W_{\ell, m}$ denote the set of

$$\gamma = (\gamma_1, \dots, \gamma_{2m}) \in (F^{\ell+1})^{2m}$$

such that for $j = 1, \dots, m$, the first half-edge (i.e. $\gamma_{2j,1} = (x_{2j,1}, i_{2j,1})$) of γ_{2j} is the same as the first half-edge of γ_{2j+1} , and the last half-edge of γ_{2j-1} is the same as the last half-edge of γ_{2j} . Here we let $\gamma_{2m+1} \stackrel{\text{def}}{=} \gamma_1$.

Following [BC19, (36)] we get

$$\begin{aligned} \|\underline{B}(s)^{(\ell)}\|^{2m} &\leq \operatorname{tr} \left(\left(\underline{B}(s)^{(\ell)} (\underline{B}(s)^{(\ell)})^* \right)^m \right) \\ &= \sum_{\substack{\gamma = (\gamma_1, \dots, \gamma_{2m}) \in W_{\ell, m} \\ \gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))}} \left(\operatorname{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \right) \\ &\quad \cdot \prod_{j=1}^{2m} \prod_{t=1}^{\ell} (\underline{S}_{i_{j,t}})_{x_{j,t} x_{j,t+1}}. \end{aligned}$$

Here we used that all $A(s)_{\mathbf{i}}$ and indeed also $\underline{B}(s)^{(\ell)}$ are trace class operators (cf. Remark 4.2).

Therefore

$$\mathbb{E}[\|B(s)^{(\ell)}\|^{2m}] \leq \sum_{\substack{\gamma=(\gamma_1,\dots,\gamma_{2m}) \in W_{\ell,m} \\ \gamma_j=((x_{j,1},i_{j,1}),\dots,(x_{j,\ell+1},i_{j,\ell+1}))}} \left| \text{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \right| \cdot \left| \mathbb{E} \left[\prod_{j=1}^{2m} \prod_{t=1}^{\ell} (S_{i_{j,t}})_{x_{j,t}x_{j,t+1}} \right] \right|. \quad (5.1)$$

Both the expectation and the absolute value of the trace in the right hand side of the above inequality will be estimated in terms of topological properties of γ , among other things. Recall that in Definition 3.2 we defined for a path $\gamma \in E^k$ a colored graph G_γ that is roughly speaking the image of the path γ . Now, for $\gamma = (\gamma_1, \dots, \gamma_{2m}) \in W_{\ell,m}$ we define G_γ to be the union of the graphs G_{γ_i} . In other words, the vertices of G_γ are the elements of $[n]$ visited by γ and the edges are just the ones followed by γ with their corresponding colors. We say that a colored edge of G_γ has *multiplicity one* if it is traversed exactly once by γ (in either direction, so an edge traversed in both directions is **not** multiplicity one).

Notice that for $\gamma \in W_{\ell,m}$, G_γ is always connected. We let $W_{\ell,m}(v, e)$ denote the collection of $\gamma \in W_{\ell,m}$ such that G_γ has v vertices and e edges; by the previous remark $W_{\ell,m}(v, e)$ is empty unless $e - v + 1 \geq 0$.

We say two elements γ and γ' of $W_{\ell,m}$ are *isomorphic* if γ' is obtained from γ by changing the labels (in $[n]$) of the vertices of G_γ and changing the colors (in $[2d]$) assigned to edges in G_γ ; these changes to G_γ clearly induce a well-defined change of γ . The reader can see [BC19, pp. 841-842] for a more formal definition.

Bordenave-Collins prove, by associating to each isomorphism class in $W_{\ell,m}$ a canonical element, and cleverly counting these canonical elements, the following result [BC19, Lemma 25].

Lemma 5.2. *The number of isomorphism classes in $W_{\ell,m}(v, e)$ is*

$$\leq (4d\ell m)^{6m \cdot \text{rank} + 10m}$$

where⁴ $\text{rank} \stackrel{\text{def}}{=} e - v + 1$.

Remark 5.3. It is in this lemma that the tangle-free hypothesis is crucially used.

Bordenave-Collins also prove the following random matrix result in [BC19, Lemma 27].

Lemma 5.4. *There is a constant $c > 0$ such that if $2\ell m \leq \sqrt{n}$ and $\gamma = (\gamma_1, \dots, \gamma_{2m}) \in W_{\ell,m}(v, e)$, with $\gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))$ for $j \in [2m]$, with G_γ having e_1 edges of multiplicity 1, then*

$$\left| \mathbb{E} \left[\prod_{j=1}^{2m} \prod_{t=1}^{\ell} (S_{i_{j,t}})_{x_{j,t}x_{j,t+1}} \right] \right| \leq c^{m+\text{rank}} \left(\frac{1}{n} \right)^e \left(\frac{6\ell m}{\sqrt{n}} \right)^{\max(e_1 - 4 \cdot \text{rank} - 4m, 0)},$$

⁴We choose to use rank rather than what Bordenave-Collins call χ , because of potential confusion with Euler characteristic.

where $\text{rank} = e - v + 1$.

Remark 5.5. The quantity $\left| \mathbb{E} \left[\prod_{j=1}^{2m} \prod_{t=1}^{\ell} (\underline{S}_{i_{j,t}})_{x_{j,t}x_{j,t+1}} \right] \right|$ is constant on isomorphism classes in $W_{\ell,m}$.

Now grouping terms in (5.1) into isomorphism classes and using the previous lemmas (this will be done formally later), the only estimate left is the following one. This is a main point of departure from Bordenave-Collins and replaces [BC19, Lemma 26].

Proposition 5.6. *For any compact subset $\mathcal{K} \subset \{s : \text{Re}(s) > \frac{\delta}{2}\}$, there is $\rho = \rho(\mathcal{K}) < 1$ such that for all $s \in \mathcal{K}$, all $m > 0$ and $\gamma^0 \in W_{\ell,m}(v, e)$*

$$\sum_{\substack{\gamma \text{ isomorphic to } \gamma^0 \\ \gamma = (\gamma_1, \dots, \gamma_{2m})}} \left| \text{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \right| \leq C^{m+\text{rank}+e_1} n^v \rho^{2\ell m}.$$

where e_1 is the number of edges of G_{γ^0} of multiplicity one and $\text{rank} = e - v + 1$.

Proof. Write $s = r + it$. First, we use Lemma 2.8 to get

$$\begin{aligned} & \sum_{\substack{\gamma \text{ isomorphic to } \gamma^0 \\ \gamma = (\gamma_1, \dots, \gamma_{2m})}} \left| \text{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \right| \\ & \leq C^m \sum_{\substack{\gamma \text{ isomorphic to } \gamma^0 \\ \gamma = (\gamma_1, \dots, \gamma_{2m})}} \Upsilon_{\mathbf{i}(\gamma_1)}^r \Upsilon_{\mathbf{i}(\gamma_2)}^r \cdots \Upsilon_{\mathbf{i}(\gamma_{2m})}^r \\ & \leq C_1^m \sum_{\substack{\gamma \text{ isomorphic to } \gamma^0 \\ \gamma = (\gamma_1, \dots, \gamma_{2m})}} \Upsilon_{\mathbf{i}(\gamma_1)'}^r \Upsilon_{\mathbf{i}(\gamma_2)'}^r \cdots \Upsilon_{\mathbf{i}(\gamma_{2m})'}^r \end{aligned} \tag{5.2}$$

for some $C_1(\mathcal{K}) > 0$, where the last inequality used (2.2), $\Upsilon_i \geq c(\Gamma) > 0$ for $i \in [2d]$, and (2.4) to deduce

$$\Upsilon_{\mathbf{i}} \leq c^{-1} \Upsilon_{\mathbf{i}} \Upsilon_{i_1} \leq c^{-1} K \Upsilon_{\mathbf{i}} \leq c^{-1} K \Upsilon_{\mathbf{i}'}.$$

Here we view γ^0 as fixed; let

$$\gamma^0 = (\gamma_1^0, \dots, \gamma_{2m}^0), \quad \gamma_j^0 = ((x_{j,1}^0, i_{j,1}^0), \dots, (x_{j,\ell+1}^0, i_{j,\ell+1}^0)).$$

We build a graph \tilde{G} as follows. Let \tilde{V} denote the vertices of G_{γ^0} that are either of degree ≥ 3 or of the form $x_{j,1}^0$ or $x_{j,\ell+1}^0$ for $j \in [2m]$. Now, every vertex of G_{γ^0} that is not in \tilde{V} has degree 2 and for each of these vertices, we remove it by merging the two adjacent edges into one edge; this can be done sequentially. The resulting graph on \tilde{V} is called \tilde{G} and has edge set that we call \tilde{E} . Every edge of \tilde{E} is labeled by a sequence of edges of G_{γ^0} . We write e' for the number of vertices of \tilde{G} and v' for the number of vertices. Recall that $\text{rank} \stackrel{\text{def}}{=} e - v + 1$.

The following (in)equalities appear in [BC19, proof of Lemma 26] and are either obvious

or not hard to check directly:

$$e = v + \text{rank} - 1, \quad (5.3)$$

$$e' = v' + \text{rank} - 1,$$

$$e' \leq 3 \cdot \text{rank} + 4m - 3. \quad (5.4)$$

Choosing γ isomorphic to γ^0 amounts to the following choices:

- Choosing distinct numbers in n for the vertices of G_{γ^0} , and, independently,
- Choosing ‘allowable’ labels in $[2d]$ for the edges of E_{γ^0} .

There are clearly $(n)(n-1)\cdots(n-v+1) \leq n^v$ of the first kind of choice. We will later incorporate this into our final estimate. Now we count the second kind of choice.

The reason for manipulating (5.2) into its current form is that the sequences

$$\mathbf{i}(\gamma_1)', \mathbf{i}(\gamma_2)', \dots, \mathbf{i}(\gamma_{2m})' \quad (5.5)$$

appearing therein are exactly the sequence of edge colors read by walking in G_γ according to the respective γ_j . Note that by assumption, the underlying graph of G_γ (forgetting edge colors) is the same as that of G_{γ_0} and hence has underlying topological graph \tilde{G} . For each edge of \tilde{G} , the number of times it is *covered* by γ is the number of times γ traverses it in either direction. (Note that γ and γ_0 cover each edge of \tilde{G} the same number of times, and in the same orders and directions.)

We now partition **separately** each $\mathbf{i}(\gamma_j)'$ in (5.2) into subsequences of three⁵ mutually exclusive types:

- 0) a subsequence that arises from γ traversing an edge of \tilde{G} that is only ever covered once by γ_0 and hence also γ .
- 1) a subsequence that arises from γ traversing an edge of \tilde{G} for the first or second time (in any direction) that is covered at least two times by γ_0 and hence also γ .
- 2) a subsequence that arises from γ traversing a **sequence** of edges of \tilde{G} where each edge is traversed for at least the third time (in either direction).

(Here subsequences are repeated according to their multiplicity.) There are at most $2e'$ subsequences of type **0** or type **1** and hence also at most $2e' + 4m$ subsequences of type **2** and at most $4e' + 4m$ subsequences in total.

We use the multiplicative estimate (2.4) for Υ to obtain

$$\Upsilon_{\mathbf{i}(\gamma_1)'}^r \Upsilon_{\mathbf{i}(\gamma_2)'}^r \cdots \Upsilon_{\mathbf{i}(\gamma_{2m})'}^r \leq K^{(8e'+8m)r} \left(\prod_{\text{type } \mathbf{0} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r \right) \left(\prod_{\text{type } \mathbf{1} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r \right) \left(\prod_{\text{type } \mathbf{2} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r \right). \quad (5.6)$$

⁵In Bordenave-Collins type **0** and type **1** are considered as one type but here it is useful to separate these out.

We deal with these factors in turn.

Write $\{\mathbf{U}_i\}$ for the type **0** subsequences of γ . Their total length is clearly e_1 and there are at most e' of them.

The subsequences of type **1** come in pairs \mathbf{u}, \mathbf{v} of the form either

$$\mathbf{u} = \mathbf{v}, \quad \text{or} \quad \mathbf{u} = \bar{\mathbf{v}}.$$

In either case the contribution of this pair to $\left(\prod_{\text{type } \mathbf{1} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r\right)$, by the mirror estimate (2.5), is

$$\leq K^r \Upsilon_{\mathbf{u}}^{2r}.$$

Let $\{\mathbf{u}_i\}$ denote a collection of representative subsequences of type **1**, one for each pair of type **1**. The total number of these pairs is $\leq e'$. The total length of the \mathbf{u}_i is $e - e_1$. Therefore

$$\prod_{\text{type } \mathbf{1} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r \leq \prod_i (K^r \Upsilon_{\mathbf{u}_i}^{2r}) = K^{e'r} \prod_i \Upsilon_{\mathbf{u}_i}^{2r}. \quad (5.7)$$

Finally we deal with the type **2** subsequences. Recalling that there are at most $2e' + 4m$ of these and their total length is $2(\ell m - e) + e_1$ as in [BC19, (44)], by using (2.7) we obtain

$$\prod_{\text{type } \mathbf{2} \text{ } \mathbf{u}} \Upsilon_{\mathbf{u}}^r \leq \prod_{\text{type } \mathbf{2} \text{ } \mathbf{u}} K^r D^{-r|\mathbf{u}|} \leq K^{r(2e'+4m)} D^{-r(2(\ell m - e) + e_1)}. \quad (5.8)$$

Now, to choose γ isomorphic to γ_0 , we need to choose the vertex labels in $[n]$ (of which there are $\leq n^v$ choices) and then

- Choose the sequence of edge colors in each type **0** subsequence \mathbf{U}_i
- Choose the sequence of edge colors in each type **1** subsequence \mathbf{u}_i
- (This completes the choice of γ because type **2** subsequences are determined by the previous choices.)

Now combining (5.2), (5.5), (5.6), (5.7), (5.8) we obtain

$$\begin{aligned} & \sum_{\substack{\gamma \text{ isomorphic to } \gamma^0 \\ \gamma = (\gamma_1, \dots, \gamma_{2m})}} \left| \text{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \right| \\ & \leq n^v C_1^m K^{(11e'+12m)r} D^{-r(2(\ell m - e) + e_1)} \left(\prod_i \Upsilon_{\mathbf{U}_i}^r \right) \left(\prod_i \Upsilon_{\mathbf{u}_i}^{2r} \right). \end{aligned} \quad (5.9)$$

Now using the bound in terms of the pressure from Lemma 2.4 we obtain that

$$\begin{aligned} (5.9) & \leq n^v C_1^m K^{(11e'+12m)r} D^{-r(2(\ell m - e) + e_1)} C^{2e'} \left(\prod_i \exp(|\mathbf{U}_i| P(r_1)) \right) \left(\prod_i \exp(|\mathbf{u}_i| P(2r_1)) \right) \\ & = n^v C_1^m K^{(11e'+12m)r} D^{-r(2(\ell m - e) + e_1)} C^{2e'} \exp(e_1 P(r_1)) \exp((e - e_1) P(2r_1)) \end{aligned}$$

where

$$r_1 \stackrel{\text{def}}{=} \min\{ \operatorname{Re}(s) : s \in \mathcal{K} \} > \frac{\delta}{2}.$$

We conclude by using (5.3) and (5.4) to obtain for some consolidated constant $c > 1$ depending on all parameters

$$(5.9) \leq c^{m+\operatorname{rank}+e_1} n^v D^{-r_1 2(\ell m - v)} \exp(vP(2r_1)).$$

The key point here is that $P(2r_1) < 0$.

If $v \leq \frac{\ell m}{2}$ we get the stated result with $\rho = D^{-\frac{r_1}{2}}$. Otherwise $v > \frac{\ell m}{2}$ and we get the result with $\rho = \exp\left(\frac{P(2r_1)}{4}\right) < 1$. In any case we get the result with $\rho < 1$ equal to the maximum of these two values. This concludes the proof. \square

Proof of Proposition 5.1. Given Lemmas 5.2 and 5.4 and Proposition 5.6 the proof is very similar to [BC19, proof of Prop. 24] but we give the details here for completeness.

For $n \geq 3$ let

$$m = \left\lfloor \frac{\log n}{13 \log \log n} \right\rfloor.$$

We partition all possible isomorphism classes of paths γ^0 in $W_{\ell, m}$ according to the number v of vertices and number e of edges of G_{γ^0} and also the number e_1 of edges of G_{γ^0} that are multiplicity one. We write $\operatorname{rank} \stackrel{\text{def}}{=} e - v + 1$. We have the inequalities

$$2(e - \ell m) \leq e_1 \leq e \tag{5.10}$$

by [BC19, Lemma 27] and can assume $e - v + 1 \geq 0$ (or else $W_{\ell, m}(v, e)$ is empty).

The contribution to (5.1) from a given isomorphism class with parameters v, e, e_1 is

$$\leq n(cC)^{m+\operatorname{rank}} \left(\frac{1}{n}\right)^{\operatorname{rank}} \left(\frac{6\ell m}{\sqrt{n}}\right)^{\max(e_1 - 4 \cdot \operatorname{rank} - 4m, 0)} C^{e_1} \rho^{2\ell m} \tag{5.11}$$

by Lemma 5.4 and Proposition 5.6.

If $e_1 - 4 \cdot \operatorname{rank} - 4m \leq 0$ then $C^{e_1} \leq (C^4)^{\operatorname{rank}+m}$ and therefore

$$(5.11) \leq n(cC^5)^{m+\operatorname{rank}} \left(\frac{1}{n}\right)^{\operatorname{rank}} \left(\frac{6\ell m}{\sqrt{n}}\right)^0 \rho^{2\ell m}.$$

If alternatively $e_1 - 4 \cdot \operatorname{rank} - 4m \geq 0$ then by (5.10) we get

$$\begin{aligned} (5.11) &\leq n(cC^5)^{m+\operatorname{rank}} \left(\frac{1}{n}\right)^{\operatorname{rank}} \left(\frac{6C\ell m}{\sqrt{n}}\right)^{e_1 - 4 \cdot \operatorname{rank} - 4m} \rho^{2\ell m} \\ &\leq n(cC^5)^{m+\operatorname{rank}} \left(\frac{1}{n}\right)^{\operatorname{rank}} \left(\frac{6C\ell m}{\sqrt{n}}\right)^{2e - 2\ell m - 4 \cdot \operatorname{rank} - 4m} \rho^{2\ell m} \\ &= n(cC^5)^{m+\operatorname{rank}} \left(\frac{1}{n}\right)^{\operatorname{rank}} \left(\frac{6C\ell m}{\sqrt{n}}\right)^{2v - 2(\ell+2)m - 2 \cdot \operatorname{rank}} \rho^{2\ell m}. \end{aligned}$$

In any case we get that the contribution to (5.1) from a single isomorphism class is

$$\leq n(cC^5)^{m+\text{rank}} \left(\frac{1}{n}\right)^{\text{rank}} \left(\frac{6C\ell m}{\sqrt{n}}\right)^{\max(2v-2(\ell+2)m-2\cdot\text{rank},0)} \rho^{2\ell m}.$$

Now, using Lemma 5.2 and summing over v and rank we get

$$\begin{aligned} & \mathbb{E}[\|\underline{B}(s)^{(\ell)}\|^{2m}] \\ & \leq n\rho^{2\ell m} \sum_{v=1}^{\infty} \sum_{\text{rank}=0}^{\infty} (4d\ell m)^{6m\cdot\text{rank}+10m} (cC^5)^{m+\text{rank}} \left(\frac{1}{n}\right)^{\text{rank}} \left(\frac{6C\ell m}{\sqrt{n}}\right)^{\max(2v-2(\ell+2)m-2\cdot\text{rank},0)}. \end{aligned}$$

Now a series of elementary bounds involving geometric series that can be found in [BC19, proof of Prop. 24] leads to

$$\mathbb{E}[\|\underline{B}(s)^{(\ell)}\|^{2m}] \leq n\rho^{2\ell m} (c'\ell m)^{10m}$$

for some new constant $c' = c'(\mathcal{K}) > 0$. Let $\rho' > \rho$ be a value in $(\rho, 1)$.

Applying Markov's inequality now gives

$$\begin{aligned} \text{Prob}[\|\underline{B}(s)^{(\ell)}\| \geq (\log n)^{20}(\rho')^{\ell}] & \leq \frac{n(c'\ell m)^{10m}}{(\log n)^{40m}} \left(\frac{\rho}{\rho'}\right)^{2\ell m} \\ & \leq \frac{n(c' \log n)^{20m}}{(\log n)^{40m}} \left(\frac{\rho}{\rho'}\right)^{2\ell m} \leq C' \left(\frac{\rho}{\rho'}\right)^{2\ell m} \end{aligned}$$

as required. \square

5.2 Norm of $R_k(s)^{(\ell)}$

Proposition 5.7. *Let \mathcal{K} be a compact subset of $\{s : \text{Re}(s) > \frac{\delta}{2}\}$. There is $c = c(\mathcal{K}) > 0$ and $\rho_1 = \rho_1(\mathcal{K}) > 1$ such that for any $s \in \mathcal{K}$, for all $1 \leq k \leq \ell \leq \log n$ the event*

$$\|R_k(s)^{(\ell)}\| \leq (\log n)^{40} \rho_1^{\ell}$$

holds with probability at least $1 - c \exp\left(-\frac{\ell \log n}{c \log \log n}\right)$.

First we follow [BC19, §§4.4.2] and define⁶ $\hat{W}_{\ell,m,k}$ to be the set of $\gamma = (\gamma_1, \dots, \gamma_{2m})$ satisfying the same conditions as the elements of $W_{\ell,m}$ **except here**, each γ_i is required to be in $F_k^{\ell+1} \setminus F^{\ell+1}$, recalling the definition of $F_k^{\ell+1}$ and $F^{\ell+1}$ from Definition 3.5.

⁶Here there is a typo in [BC19]; we checked with the authors that they meant to define $\hat{W}_{\ell,m,k}$ (which Bordenave-Collins call $\hat{W}_{\ell,m}$) using $F_k^{\ell+1} \setminus F^{\ell+1}$ (not $F_k^{\ell+1}$ as written there).

We then obtain as in [BC19, pg 851]

$$\begin{aligned}
\|R_k(s)^{(\ell)}\|^{2m} &\leq \text{tr} \left(\left(R_k(s)^{(\ell)} (R_k(s)^{(\ell)})^* \right)^m \right) \\
&= \sum_{\substack{\gamma=(\gamma_1, \dots, \gamma_{2m}) \in \hat{W}_{\ell, m, k} \\ \gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))}} \text{tr} \left[A(s)_{\mathbf{i}(\gamma_1)} A(s)_{\mathbf{i}(\gamma_2)}^* \cdots A(s)_{\mathbf{i}(\gamma_{2m-1})} A(s)_{\mathbf{i}(\gamma_{2m})}^* \right] \\
&\quad \cdot \prod_{j=1}^{2m} \left(\prod_{t=1}^{k-1} (\underline{S}_{i_{j,t}})_{x_{j,t}, x_{j,t+1}} \right) \left(\prod_{t=k+1}^{\ell} (S_{i_{j,t}})_{x_{j,t}, x_{j,t+1}} \right). \tag{5.12}
\end{aligned}$$

Isomorphism classes of elements of $\hat{W}_{\ell, m, k}$ are defined the same way as for isomorphism classes in $W_{\ell, m}$.

For $\gamma \in \hat{W}_{\ell, m, k}$ we define \hat{G}_γ to be the following graph. If we write

$$\gamma = (\gamma_1, \dots, \gamma_{2m}), \quad \gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))$$

then the vertex set of \hat{G}_γ is defined to be the set of all $x_{j,t}$ with $j \in [2m], t \in [\ell+1]$ and the colored edges

$$\{[x_{j,t}, i_{j,t}, x_{j,t+1}] : 1 \leq t \leq k-1, k+1 \leq t \leq \ell\}$$

modulo the equivalence relation (3.1). Notice this is a sub-colored-graph of G_γ with restriction on the edges corresponding to the matrix product in (5.12).

For $\gamma \in \hat{W}_{\ell, m, k}$ it is possible for \hat{G}_γ to be disconnected but, as in [BC19, (53)], \hat{G}_γ cannot have more vertices than edges; this is actually the purpose of ensuring each γ_i is tangled. We have from [BC19, Lemma 29] the following result.

Lemma 5.8. *Let $\hat{W}_{\ell, m, k}(v, e)$ denote the elements of $\hat{W}_{\ell, m, k}$ such that \hat{G}_γ has v vertices and e edges. The number of isomorphism classes in $\hat{W}_{\ell, m, k}(v, e)$ is*

$$\leq (2d\ell m)^{12m \cdot \text{rank} + 20m}$$

where as before, $\text{rank} \stackrel{\text{def}}{=} e - v + 1 \geq 1$ if $\hat{W}_{\ell, m, k}$ is non-empty.

We also have the following results from [BC19, pg. 853], the first of which is trivial.

Lemma 5.9. *The number of elements of any isomorphism class in $\hat{W}_{\ell, m, k}(v, e)$ is at most*

$$n^v d^e.$$

Lemma 5.10. *There is a constant $c > 0$ such that if $2\ell m \leq \sqrt{n}$ and $\gamma = (\gamma_1, \dots, \gamma_{2m}) \in \hat{W}_{\ell, m, k}(v, e)$, with $\gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))$ for $j \in [2m]$, then*

$$\left| \mathbb{E} \left[\prod_{j=1}^{2m} \left(\prod_{t=1}^{k-1} (\underline{S}_{i_{j,t}})_{x_{j,t}, x_{j,t+1}} \right) \left(\prod_{t=k+1}^{\ell} (S_{i_{j,t}})_{x_{j,t}, x_{j,t+1}} \right) \right] \right| \leq c \left(\frac{9}{n} \right)^e.$$

Proof of Proposition 5.7. Suppose $n \geq 3$ and define

$$m = \left\lfloor \frac{\log n}{25 \log \log n} \right\rfloor.$$

Taking the expected value of (5.12), and using Lemma 2.8 and the fact that $\Upsilon_{\mathbf{i}} \leq C$ for any \mathbf{i} we obtain for $C = C(\Lambda) > 0$

$$\begin{aligned} & \mathbb{E} \left[\|R_k(s)^{(\ell)}\|^{2m} \right] \\ & \leq C^m \sum_{\substack{\gamma=(\gamma_1, \dots, \gamma_{2m}) \in \hat{W}_{\ell, m, k} \\ \gamma_j = ((x_{j,1}, i_{j,1}), \dots, (x_{j,\ell+1}, i_{j,\ell+1}))}} \left| \mathbb{E} \left[\prod_{j=1}^{2m} \left(\prod_{t=1}^{k-1} (S_{i_{j,t}})_{x_{j,t} x_{j,t+1}} \right) \left(\prod_{t=k+1}^{\ell} (S_{i_{j,t}})_{x_{j,t} x_{j,t+1}} \right) \right] \right|. \end{aligned}$$

Now noting that the product of matrix coefficients above is constant on isomorphism classes in $\hat{W}_{\ell, m, k}(v, e)$, and using Lemmas 5.8, 5.9, and 5.10, we obtain

$$\begin{aligned} \mathbb{E} \left[\|R_k(s)^{(\ell)}\|^{2m} \right] & \leq c C^m \sum_{v=1}^{2\ell m} \sum_{e=v}^{\infty} (2d\ell m)^{12m \cdot (e-v+1) + 20m} \left(\frac{9}{n} \right)^e n^v d^e \\ & = (C' d\ell m)^{32m} \sum_{v=1}^{2\ell m} \left(\frac{n}{(2d\ell m)^{12m}} \right)^v \sum_{e=v}^{\infty} \left(\frac{9d}{n} (2d\ell m)^{12m} \right)^e. \end{aligned}$$

The inner geometric series is convergent by our constraint on ℓ, m so we obtain

$$\begin{aligned} \mathbb{E} \left[\|R_k(s)^{(\ell)}\|^{2m} \right] & \leq (C' d\ell m)^{32m} \sum_{v=1}^{2\ell m} \left(\frac{n}{(2d\ell m)^{12m}} \frac{9d}{n} (2d\ell m)^{12m} \right)^v \\ & = (C' d\ell m)^{32m} \sum_{v=1}^{2\ell m} (9d)^v \\ & \leq (C'' d\ell m)^{32m} (9d)^{2\ell m}. \end{aligned}$$

Using Markov's inequality yields

$$\begin{aligned} \text{Prob}[\|R_k(s)^{(\ell)}\| \geq (\log n)^{40} (10d)^{\ell}] & \leq \frac{(C'' d\ell m)^{32m}}{(\log n)^{80m}} \left(\frac{9}{10} \right)^{2\ell m} \\ & \leq \frac{(c(\log n))^{64m}}{(\log n)^{80m}} \left(\frac{9}{10} \right)^{2\ell m} \\ & \leq c' \left(\frac{9}{10} \right)^{2\ell m}. \end{aligned}$$

This directly implies the result. □

6 Proof of Theorem 1.1

It is sufficient to prove Theorem 1.1 when \mathcal{K} is a rectangle in $\{s : \operatorname{Re}(s) > \frac{\delta}{2}\}$ (by covering \mathcal{K} with finitely many rectangles). So suppose \mathcal{K} is such a rectangle. Let ρ and ρ_1 be the constants provided by Propositions 5.1 and 5.7 for this \mathcal{K} . Let C_0 be the constant provided by Lemma 2.5 for this \mathcal{K} .

We let

$$\beta = \min \left(\frac{1}{8 \log(2d-1)}, \log \left(\frac{\rho_1}{\rho} \right)^{-1} \right), \quad (6.1)$$

$$\alpha = 2\beta \log C_0. \quad (6.2)$$

A δ -net of \mathcal{K} is a finite subset of \mathcal{K} such that any point of \mathcal{K} is within Euclidean distance δ of some point of the δ -net. Now it is easy to see that for each n we can choose an $n^{-\alpha}$ -net \mathcal{N}_n of \mathcal{K} with

$$|\mathcal{N}_n| \leq Cn^{2\alpha}$$

for some $C = C(\mathcal{K}, \alpha) > 0$. We want all the following events to hold simultaneously for

$$\ell = \lfloor \beta \log n \rfloor.$$

A G_σ is ℓ -tangle-free.

B $\|\underline{B}(s_i)^{(\ell)}\| \leq (\log n)^{20} \rho^\ell$ for $\rho < 1$ as in Proposition 5.1 for all $s_i \in \mathcal{N}_n$.

C $\|R_k(s_i)^{(\ell)}\| \leq (\log n)^{40} \rho_1^\ell$ for $\rho_1 > 0$ as in Proposition 5.2 for all $s_i \in \mathcal{N}_n$ and all $1 \leq k \leq \ell$.

By Lemma 3.6 and Propositions 5.1 and 5.2, these events all hold with probability at least

$$1 - c \frac{\ell^3 (2d-1)^{4\ell}}{n} - cCn^{2\alpha} \exp \left(-\frac{\ell \log n}{c \log \log n} \right)$$

for some $c > 0$. The last term above tends to zero for any fixed α, β and the second term tends to zero by our choice of β in (6.1). So with probability tending to one as $n \rightarrow \infty$, all three families **A**, **B**, **C** of events above hold.

Now, if **A**, **B**, **C** hold then by Lemma 4.3

$$\begin{aligned} \|B(s_i)^\ell|_{K_0}\| &\leq \|\underline{B}(s_i)^{(\ell)}\| + \frac{1}{n} \sum_{k=1}^{\ell} \|R_k(s_i)^{(\ell)}\| \\ &\leq (\log n)^{20} \rho^\ell + \frac{\ell}{n} (\log n)^{40} \rho_1^\ell \\ &\leq (1 + \beta) (\log n)^{41} \rho^\ell \end{aligned}$$

for all $s_i \in \mathcal{N}_n$, where the last inequality used (6.1). Therefore by Proposition 4.1 we have

$$\|\mathcal{L}_{s_i, \rho_n^0}^\ell\| \leq (1 + \beta) (\log n)^{41} \rho^\ell \quad (6.3)$$

for all $s_i \in \mathcal{N}_n$.

Now by Lemma 2.5 there is a constant $C_0 = C_0(K) > 0$ such that for all $s \in \mathcal{K}$, if s_i is a point in \mathcal{N}_n for distance $\leq n^{-\alpha}$ from s then

$$\|\mathcal{L}_{s, \rho_n^0}^\ell - \mathcal{L}_{s_0, \rho_n^0}^\ell\| \leq |s - s_0| C_0^\ell \leq n^{-\alpha} C_0^\ell \leq n^{(\beta \log C_0 - \alpha)} < n^{-\frac{\alpha}{2}} \quad (6.4)$$

by our choice of α in (6.2).

Combining (6.3) and (6.4) we obtain that with probability tending to one as $n \rightarrow \infty$, for all $s \in \mathcal{K}$

$$\|\mathcal{L}_{s, \rho_n^0}^\ell\| < 1.$$

This immediately implies that $\mathcal{L}_{s, \rho_n^0}$ does not have 1 as an eigenvalue, and hence by Proposition 2.2 there are no new resonances of the random cover X_ϕ in \mathcal{K} . \square

References

- [Alo86] N. Alon. Eigenvalues and expanders. *Combinatorica*, 6(2):83–96, 1986. Theory of computing (Singer Island, Fla., 1984). 3
- [BC19] C. Bordenave and B. Collins. Eigenvalues of random lifts and polynomials of random permutation matrices. *Ann. of Math. (2)*, 190(3):811–875, 2019. 4, 11, 15, 16, 17, 18, 19, 21, 22, 23, 25, 26, 27, 28
- [BD17] J. Bourgain and S. Dyatlov. Fourier dimension and spectral gaps for hyperbolic surfaces. *Geom. Funct. Anal.*, 27(4):744–771, 2017. 4, 5, 9
- [BD18] J. Bourgain and S. Dyatlov. Spectral gaps without the pressure condition. *Ann. of Math. (2)*, 187(3):825–867, 2018. 4
- [BGS11] J. Bourgain, A. Gamburd, and P. Sarnak. Generalization of Selberg’s $\frac{3}{16}$ theorem and affine sieve. *Acta Math.*, 207(2):255–290, 2011. 4
- [BM04] R. Brooks and E. Makover. Random construction of Riemann surfaces. *J. Differential Geom.*, 68(1):121–157, 2004. 4
- [BMM17] W. Ballmann, H. Matthiesen, and S. Mondal. Small eigenvalues of surfaces of finite type. *Compos. Math.*, 153(8):1747–1768, 2017. 3
- [Bor] C. Bordenave. A new proof of Friedman’s second eigenvalue theorem and its extension to random lifts. *Annales scientifiques de l’Ecole normale supérieure*. to appear, available at arxiv:1502.04482. 3, 18
- [Bor16] D. Borthwick. *Spectral theory of infinite-area hyperbolic surfaces*, volume 318 of *Progress in Mathematics*. Birkhäuser/Springer, [Cham], second edition, 2016. 2, 6, 8

- [Bow79] R. Bowen. Hausdorff dimension of quasicircles. *Inst. Hautes Études Sci. Publ. Math.*, (50):11–25, 1979. [6](#)
- [But98] J. Button. All Fuchsian Schottky groups are classical Schottky groups. In *The Epstein birthday schrift*, volume 1 of *Geom. Topol. Monogr.*, pages 117–125. Geom. Topol. Publ., Coventry, 1998. [2](#), [5](#)
- [FK81] Z. Füredi and J. Komlós. The eigenvalues of random symmetric matrices. *Combinatorica*, 1(3):233–241, 1981. [21](#)
- [Fri03] J. Friedman. Relative expanders or weakly relatively Ramanujan graphs. *Duke Math. J.*, 118(1):19–35, 2003. [3](#)
- [Fri08] J. Friedman. A proof of Alon’s second eigenvalue conjecture and related problems. *Mem. Amer. Math. Soc.*, 195(910):viii+100, 2008. [3](#), [15](#)
- [Gam02] A. Gamburd. On the spectral gap for infinite index “congruence” subgroups of $SL_2(\mathbf{Z})$. *Israel J. Math.*, 127:157–200, 2002. [4](#)
- [GLMST21] C. Gilmore, E. Le Masson, T. Sahlsten, and J. Thomas. Short geodesic loops and L^p norms of eigenfunctions on large genus random surfaces. *Geom. Funct. Anal.*, 31(1):62–110, 2021. [4](#)
- [JN12] D. Jakobson and F. Naud. On the critical line of convex co-compact hyperbolic surfaces. *Geom. Funct. Anal.*, 22(2):352–368, 2012. [3](#), [5](#)
- [JNS19] D. Jakobson, F. Naud, and L. Soares. Large covers and sharp resonances of hyperbolic surfaces. *To appear, Ann. Institut Fourier.*, 2019. [3](#)
- [LP81] P. D. Lax and R. S. Phillips. The asymptotic distribution of lattice points in Euclidean and non-Euclidean spaces. In *Functional analysis and approximation (Oberwolfach, 1980)*, volume 60 of *Internat. Ser. Numer. Math.*, pages 373–383. Birkhäuser, Basel-Boston, Mass., 1981. [2](#)
- [LW21] M. Lipnowski and A. Wright. Towards optimal spectral gaps in large genus, 2021. Preprint, arXiv:2103.07496. [4](#)
- [Mir13] M. Mirzakhani. Growth of Weil-Petersson volumes and random hyperbolic surfaces of large genus. *J. Differential Geom.*, 94(2):267–300, 2013. [4](#)
- [MM87] R. R. Mazzeo and R. B. Melrose. Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature. *J. Funct. Anal.*, 75(2):260–310, 1987. [2](#)
- [MN20] M. Magee and F. Naud. Explicit spectral gaps for random covers of Riemann surfaces. *Publ. Math. Inst. Hautes Études Sci.*, 132:137–179, 2020. [3](#), [4](#), [5](#), [8](#), [9](#)

- [MNP20] M. Magee, F. Naud, and D. Puder. A random cover of a compact hyperbolic surface has relative spectral gap $\frac{3}{16} - \varepsilon$, 2020. Preprint, arXiv:2003.10911. [4](#)
- [Mon20] L. Monk. Benjamini-Schramm convergence and spectrum of random hyperbolic surfaces of high genus. *Analysis & PDE*, to appear, available at arXiv:arXiv:2002.00869, page arXiv:2002.00869, February 2020. [4](#)
- [MOW17] M. Magee, H. Oh, and D. Winter. Uniform congruence counting for Schottky semigroups in $\mathrm{SL}_2(\mathbf{Z})$, with appendix by J. Bourgain, A. Kontorovich, and M. Magee. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 01 2017. [4](#)
- [MP20] M. Magee and D. Puder. The asymptotic statistics of random covering surfaces, 2020. Preprint, arXiv:2003.05892v1. [4](#)
- [Nau05] F. Naud. Expanding maps on Cantor sets and analytic continuation of zeta functions. *Ann. Sci. École Norm. Sup. (4)*, 38(1):116–153, 2005. [2](#)
- [Nil91] A. Nilli. On the second eigenvalue of a graph. *Discrete Math.*, 91(2):207–210, 1991. [3](#)
- [OW16] H. Oh and D. Winter. Uniform exponential mixing and resonance free regions for convex cocompact congruence subgroups of $\mathrm{SL}_2(\mathbf{Z})$. *J. Amer. Math. Soc.*, 29(4):1069–1115, 2016. [4](#)
- [Pat76] S. J. Patterson. The limit set of a Fuchsian group. *Acta Math.*, 136(3-4):241–273, 1976. [2](#)
- [Pud15] D. Puder. Expansion of random graphs: new proofs, new results. *Invent. Math.*, 201(3):845–908, 2015. [3](#)
- [Sar14] P. Sarnak. Notes on thin matrix groups. In *Thin groups and superstrong approximation*, volume 61 of *Math. Sci. Res. Inst. Publ.*, pages 343–362. Cambridge Univ. Press, Cambridge, 2014. [4](#)
- [WX21] Y. Wu and Y. Xue. Random hyperbolic surfaces of large genus have first eigenvalues greater than $\frac{3}{16} - \epsilon$, 2021. Preprint, arXiv:2102.05581. [4](#)
- [Zwo17] M. Zworski. Mathematical study of scattering resonances. *Bull. Math. Sci.*, 7(1):1–85, 2017. [5](#)

Michael Magee,
Department of Mathematical Sciences,
Durham University,
Lower Mountjoy, DH1 3LE Durham,
United Kingdom
`michael.r.magee@durham.ac.uk`

Frédéric Naud,
Institut Mathématique de Jussieu,
Sorbonne Université,
4 place Jussieu, 75252 Paris Cedex 05
France
`frederic.naud@imj-prg.fr`