



Physiological changes induced by sodium chloride stress in *Aphanizomenon gracile*, *Cylindrospermopsis raciborskii* and *Dolichospermum* sp.

Emilie Houliez, Enora Briand, Florent Malo, Georges-Augustin Rovillon,
Fabienne Hervé, Elise Robert, Laetitia Marchand, Agata Zykwinska,
Amandine M.N. Caruana

► To cite this version:

Emilie Houliez, Enora Briand, Florent Malo, Georges-Augustin Rovillon, Fabienne Hervé, et al.. Physiological changes induced by sodium chloride stress in *Aphanizomenon gracile*, *Cylindrospermopsis raciborskii* and *Dolichospermum* sp.. Harmful Algae, 2021, 103, pp.102028. 10.1016/j.hal.2021.102028 . hal-03842083

HAL Id: hal-03842083

<https://hal.science/hal-03842083>

Submitted on 7 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Physiological changes induced by sodium chloride stress in *Aphanizomenon gracile*, *Cylindrospermopsis raciborskii* and *Dolichospermum* sp.

Houliez Emilie ^{1,*}, Briand Enora ¹, Malo Florent ¹, Rovillon Georges-Augustin ¹, Hervé Fabienne ¹, Robert Elise ¹, Marchand Laetitia ², Zykawska Agata ², Caruana Amandine ¹

¹ IFREMER, Phycotoxins Laboratory, F-44311 Nantes, France

² IFREMER, EM3B Laboratory, F-44311 Nantes, France

* Corresponding author : Emilie Houliez, email address : emilie.houliez@outlook.fr

Abstract :

Due to anthropogenic activities, associated with climate change, many freshwater ecosystems are expected to experience an increase in salinity. This phenomenon is predicted to favor the development and expansion of freshwater cyanobacteria towards brackish waters due to their transfer along the estuarine freshwater-marine continuum. Since freshwater cyanobacteria are known to produce toxins, this represents a serious threat for animal and human health. Saxitoxins (STXs) are classified among the most powerful cyanotoxins. It becomes thus critical to evaluate the capacity of cyanobacteria producing STXs to face variations in salinity and to better understand the physiological consequences of sodium chloride (NaCl) exposure, in particular on their toxicity. Laboratory experiments were conducted on three filamentous cyanobacteria species isolated from brackish (*Dolichospermum* sp.) and fresh waters (*Aphanizomenon gracile* and *Cylindrospermopsis raciborskii*) to determine how salinity variations affect their growth, photosynthetic activity, pigment composition, production of reactive oxygen species (ROS), synthesis of compatible solutes and STXs intracellular quotas. Salinity tolerance was found to be species-specific. *Dolichospermum* sp. was more resistant to salinity variations than *A. gracile* and *C. raciborskii*. NaCl variations reduced growth in all species. In *A. gracile*, carotenoids content was dose-dependently reduced by NaCl. By contrast, in *C. raciborskii* and *Dolichospermum* sp., variations in carotenoids content did not show obvious relationships with NaCl concentration. While in *Dolichospermum* sp. phycocyanin and phycoerythrin increased within the first 24 h exposure to NaCl, in both *A. gracile* and *C. raciborskii*, these pigments decreased proportionally to NaCl concentration. Low changes in salinity did not impact STXs production in *A. gracile* and *C. raciborskii* while higher increase in salinity could modify the toxin profile and content of *C. raciborskii* (intracellular STX decreased while dc-GTX2 increased). In estuaries, *A. gracile* and *C. raciborskii* would not be able to survive beyond the oligohaline area (i.e. salinity > 5). Conversely, in part due to its ability to accumulate compatible solutes, *Dolichospermum* sp. has the potential to face consequent salinity variations and to survive in the polyhaline area (at least up to salinity = 24).

Highlights

► In all species, sodium chloride exposure resulted in reduced growth rates. ► The responses of photosynthetic activity, pigment composition, ROS production as well as the capacity to accumulate compatible solutes were species-specific. ► In part due to its ability to accumulate compatible solutes, *Dolichospermum* sp. was the most tolerant species. ► Low sodium chloride concentrations did not impact STXs production in *Aphanizomenon gracile* and *Cylindrospermopsis raciborskii* while higher concentrations modified the toxin profile and content of *C. raciborskii* (intracellular STX decreased while dc-GTX2 increased).

Keywords : Cyanobacteria, Saxitoxins, Paralytic shellfish toxins, Harmful algal bloom, Sodium chloride, Cyanotoxins

1 Introduction

In the last decades, cyanobacteria blooms have considerably increased in frequency, magnitude and duration in many countries and ecosystems (O'Neil et al., 2012; Paerl et al., 2001; Paerl and Huisman, 2009). This phenomenon is becoming a [worldwide](#) public health and environmental concern (Burford et al., 2020). Indeed, some strains of cyanobacteria produce toxic metabolites (cyanotoxins) that are responsible for serious human and animal health problems and can even lead to mortalities in some cases. Mass development of cyanobacteria also strongly impacts ecosystems functioning and human activities. Indeed, it increases turbidity of water and accumulates as thick scums and mats that can lead to anoxia when they decomposed. A variety of odor and taste compounds are also produced making affected waters unsuitable for drinking and recreational purposes (Carmichael, 2001; Dittmann et al., 2012; Paerl et al., 2001). Saxitoxins (STXs) are among the most powerful cyanotoxins with a reported 50% mouse lethal dose of 5-10 $\mu\text{g kg}^{-1}$ (Munday, 2014; Schantz et al., 1975). STXs are a group of 57 alkaloids also known as paralytic shellfish toxins (PSTs) due to their association with paralytic shellfish poisoning (PSP) which occurs after ingestion of shellfish contaminated with PSTs (Wiese et al., 2010). STXs are neurotoxins that selectively block voltage-gated sodium channels present in neuronal cell membranes and prolong the gating of potassium channels in heart muscle cells (Pearson et al., 2010). PSP symptoms includes perspiration, vomiting and cardiovascular failure due to respiratory muscles paralysis and heart malfunctioning (Llewellyn, 2006; Wiese et al., 2010). So far, STXs synthesis has been described in nine genera of filamentous cyanobacteria mainly found in freshwaters: *Dolichospermum* (formerly known as *Anabaena*), *Aphanizomenon*, *Cuspidothrix*, *Cylindrospermopsis*, *Lyngbya*, *Planktothrix*, *Raphidiopsis*, *Scytonema* and *Phormidium* (Cirés and Ballot, 2016; Pearson et al., 2016). Water contamination with STXs

produced by these genera is a worldwide phenomenon detected in Asia, North and South America, Europe and Oceania. In addition, the frequency and distribution of STXs-containing cyanobacteria blooms appears to be on the rise (Pearson et al., 2016). In the last 10 years, STXs-producing cyanobacteria (STXs-cyanobacteria) have, for example, been detected for the first time in Arctic (Kleinteich et al., 2013), New Zealand (Smith et al., 2011), Canada (Lajeunesse et al., 2012) and several locations in Europe (Ballot et al., 2010; Jančula et al., 2014; Rapala et al., 2005; Teneva et al., 2010; Wörmer et al., 2011) with some of those blooms reaching high STXs concentrations (such as 25-1000 μg PSTs L^{-1} (Rapala et al., 2005; Wörmer et al., 2011) [knowing that in Europe, the acute reference dose for STXs has been fixed at 0.5 \$\mu\text{g}\$ STX equivalents \$\text{kg}^{-1}\$ of body weight \(Paredes et al., 2011\)](#)). Recently, climate change has been linked to the extension and intensification of harmful cyanobacterial blooms (Paerl and Huisman, 2009; Paerl and Paul, 2012; Slim et al., 2014). Even though the main influence of climate change is global warming with its direct influence on water temperature, climate change is also likely to induce salinity fluctuations in brackish and fresh waters due to sea level rising [\(through the penetration of seawater into estuaries over a higher distance and increasing submersions\)](#) and modifications in frequency, intensity and duration of precipitations and droughts (Li et al., 2016; Paerl and Paul, 2012). This, associated with the increasing anthropogenic salinization of inland waters, may strongly modify phytoplankton community structure and blooms occurrence (Williams, 2001). It is thought that salinity fluctuations would be favorable for cyanobacteria. Indeed, one of the impacts of salinization is increased vertical density stratification (Paerl and Huisman, 2009 and citations therein). Such conditions would facilitate buoyancy of some cyanobacteria because they have gas vesicles in cells that enable them to float to the water surface and in this way, they outcompete other algae for available light and nutrients (Antunes et al., 2015; Cirés and Ballot, 2016; Li et al., 2016). [Additionally, the sparse information on the response of harmful](#)

bloom forming cyanobacteria to salinity seems to indicate that some genera or strains are rather tolerant to salinity changes, sometimes more than their counterpart eukaryotic freshwater phytoplankton species (Brutemark et al., 2015; Moisander et al., 2002; Paerl and Paul, 2012; Preece et al., 2017). The salt tolerance of these cyanobacteria would be one of the causes of the increasing number of harmful cyanobacteria blooms in brackish waters (Paerl and Paul, 2012; Soto-Liebe et al., 2013). This salt tolerance also spotlights estuaries as potential vectors for increasing spread of harmful cyanobacteria through possible transfers along the freshwater-marine continuum (Bormans et al., 2019). Several estuarine localities have already experienced harmful cyanobacteria appearance and sometimes bloom formation in response to salinity fluctuations (John and Kemp, 2006; Preece et al., 2017; Robson and Hamilton, 2013; Ross et al., 2019). Such an increased expansion of harmful cyanobacteria blooms exposes other aquatic organisms and human users of these waters to elevated risks of contamination with cyanotoxins (Paerl and Paul, 2012). Especially since field studies have suggested causal relationships between variations in salinity (or conductivity) and increases in water STXs concentration (Brentano et al., 2016 and references therein). Despite the public health concern and economic importance of such toxic cyanobacteria occurrences, relatively few studies have actually focused on the salt tolerance of STXs-cyanobacteria and the resulting effects on STXs production. This contrasts with the significant amount of publications dealing with STXs production in toxic dinoflagellates (e.g. Aguilera-Belmonte et al., 2013 and references therein) or salt effects on other cyanotoxins such as microcystins (e.g. Georges des Aulnois et al., 2020; Georges des Aulnois et al., 2019; Melero-Jiménez et al., 2020 and references therein). Additionally, some of the STXs-cyanobacteria species remain poorly studied. To our knowledge, this is the case of *Aphanizomenon gracile* that has not been previously used as a model species to study effects of salinity variations on its physiology and toxin production. Some physiological processes potentially involved in salt stress acclimation,

such as photosynthesis, ROS production and accumulation of compatible solutes, have also received little attention in STXs-cyanobacteria. Even in *Cylindrospermopsis raciborskii*, one of the most-studied STXs-cyanobacteria, in which salt stress effects on growth and STX production have been previously described (Carneiro et al., 2013; Ongley et al., 2016; Pomati et al., 2004b). It becomes thus crucial to acquire more knowledge about the physiological response of STXs-cyanobacteria to salt exposure to fully understand their invasive potential. This study investigates impacts of increasing salt concentrations on survival, growth, pigments composition, photosynthetic activity, production of reactive oxygen species and intracellular compatible solutes and STXs contents of STXs-cyanobacteria. The objectives were: 1) to evaluate the capacity of each species to face salinity variations, 2) to identify the physiological processes impacted by salt stress and, as far as possible, their role in salt acclimation, 3) to determine whether the different species use similar strategies to deal with salinity variations and 4) to understand how salinity variations may alter the toxicity (toxin content and profile) of these species. *The physiological responses of two freshwater cyanobacteria (A. gracile and C. raciborskii) grown under controlled conditions were investigated and compared to the responses of a brackish water strain (Dolichospermum sp.). The Dolichospermum sp. strain was used as a model of cyanobacteria already used to face high salinity exposure in its natural environment.*

2 Materials and methods

2.1 Cyanobacteria strains and growth conditions

Two freshwater STX-producing strains, *Aphanizomenon gracile* (PMC627.10, isolated from Champs-sur-Marne in France, Ledreux et al., 2010) and *Cylindrospermopsis raciborskii* (PMC00.01, isolated from Jucazinho in Brazil, Bernard et al., 2003) were obtained from the microalgae and cyanobacteria culture collection of the National Museum of Natural History

(Paris, France). The brackish water non-toxin-producing strain, *Dolichospermum* sp. (CCY9401, isolated from Arcachon in France, salinity = 9) was obtained from the culture collection of the Royal Netherlands Institute of Sea Research (Yerseke, Netherlands). None of these strains has been found to produce detectable amount of anatoxins, homoanatoxins, cylindrospermopsin, desoxycylindrospermopsin, microcystins or nodularins. These three strains were maintained at 25°C under illumination of 40 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (PAR) provided by cool-white fluorescent tubes (Osram lumilux cool daylight L18w/865) with a 12:12 h light/dark cycle. Although they were diazotrophic species, the three strains were maintained in nitrogen replete culture media. *A. gracile* and *C. raciborskii* were maintained in BG11 medium (Allen, 1968; Allen and Stanier, 1968; Rippka et al., 1979). *Dolichospermum* sp. was maintained in BG11 medium in which 9 g L⁻¹ NaCl were added because this strain was isolated from brackish waters.

2.2 Salt stress experiment

Cultures in exponential growth phase were used as inoculum for new batch cultures in media containing different amount of NaCl for 4 days. *A. gracile* and *C. raciborskii* were exposed to BG11 (control conditions), BG11 + 2 g L⁻¹ NaCl (i.e. 34 mM NaCl), BG11 + 9 g L⁻¹ NaCl (i.e. 154 mM NaCl) and BG11 + 15 g L⁻¹ NaCl (i.e. 257 mM NaCl). In an estuary, these salinity variations would correspond to a transfer from freshwaters to the oligohaline area (from BG11 to BG11 + 34 mM NaCl) and from freshwaters to the mesohaline area (from BG11 to BG11 + 154 mM NaCl and from BG11 to BG11 + 257 mM NaCl). *Dolichospermum* sp. was exposed to two media in common with *A. gracile* and *C. raciborskii*: BG11 + 154 mM NaCl (which corresponds here to control conditions) and BG11 + 257 mM NaCl. It was also exposed to BG11 + 18 g L⁻¹ NaCl (i.e. 308 mM NaCl) and BG11 + 24 g L⁻¹ NaCl (i.e. 411 mM NaCl): this corresponds respectively to an increase of 154 and 257 mM NaCl in

comparison to control conditions as it was the case for *A. gracile* and *C. raciborskii*. Each condition was performed in triplicates. Sampling in aseptic conditions was done after different exposure times (0, 1, 4, 6, 24, 48 and 72h). For each treatment, cyanobacteria growth was measured by following chlorophyll a (chl a) content (see below for its measurement) and growth rate (μ) was calculated following Wood et al. (2005).

2.3 Photosynthetic activity and mean chl a-specific absorption coefficient (\bar{a}_{phy}^*)

Photosynthetic activity was measured by Pulse Amplitude Modulated (PAM) fluorometry using a PHYTO-PAM fluorometer PHYTO-ED (Heinz Walz GmbH, Effeltrich, Germany). For a detailed description of the PHYTO-PAM, see Kolbowski and Schreiber (1995) and Schreiber (1998). Before measurement, samples were dark acclimated for 15 minutes at sampling temperature. Fluorescence levels F_0 and F_m were respectively measured before and after a saturating pulse (400 ms to 4000 $\mu\text{mol photons quanta m}^{-2} \text{s}^{-1}$) to obtain the maximum quantum yield (F_v/F_m):

$$F_v/F_m = \frac{(F_m - F_0)}{F_m}$$

Rapid Light curves were then constructed by exposing the sample for 10 s to 21 sequential increasing light steps with irradiance levels ranging from 1 to 2064 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$. Actinic light was provided by 16 LEDs peaking at 655 nm. At each light step, the fluorescence levels before and after applying a saturating pulse were measured with four pulse-modulated measuring lights peaking at 470, 520, 645 and 665 nm. At each light step and for each measuring light, the effective quantum yield of photosystem II (ϕPSII also noted $\Delta F/F'_m$) was calculated according to Genty et al. (1989):

$$\phi\text{PSII} = \frac{\Delta F}{F'_m} = \frac{(F'_m - F)}{F'_m}$$

where F is the fluorescence level of the light-acclimated sample just prior to the saturating pulse and F'_m is the maximum fluorescence emitted by the light-acclimated sample after the saturating pulse.

ϕ PSII measured with the measuring light peaking at 645 nm was then selected to calculate the absolute electron transport rate (ETR) since it targets phycocyanin and was the most appropriate wavelength to our cyanobacteria species. ETR was calculated as:

$$\text{ETR} = \phi\text{PSII} \times E \times 0.36 \times \bar{a}^*_{\text{phy}}$$

where E ($\mu\text{mol photons m}^{-2} \text{ s}^{-1}$) is the actinic irradiance, 0.36 is the fraction of absorbed quanta to PSII ($f\text{AQ}_{\text{PSII}}$) in cyanobacteria (Johnsen and Sakshaug, 2007) and \bar{a}^*_{phy} ($\text{m}^2 (\text{mg chl a})^{-1}$) is the spectrally averaged (400-700 nm) chlorophyll a-specific absorption coefficient (see below for its measurement). The obtained ETR vs. E curves were fitted using the model of Eilers and Peeters (1988) to estimate the maximal light utilization efficiency (α) which corresponds to the initial slope of the curve and, maximum electron transport rate (ETR_m) which is the asymptote of the curve. Curve fitting was carried out using the downhill simplex method of the Nelder-Mead model (1965) and standard deviation of each parameter was estimated by an asymptotic method. All fittings were tested by analyses of variance ($P < 0.001$), residues being tested for normality and homogeneity of variance, and parameters significance by Student's t test ($P < 0.05$). All curve fitting processes and associated statistics were coded under MATLAB R2010b.

At each light step, non-photochemical quenching of fluorescence (NPQ) was calculated from the fluorescence levels measured with the measuring light peaking at 645 nm using the equation proposed by Serôdio et al. (2005):

$$\text{NPQ} = \frac{(F'_{\text{mm}} - F'_m)}{F'_m}$$

where F'_{mm} is the maximum F'_m value, higher than F_m , that is measured under low actinic irradiance. Then, NPQ_{max} that corresponds to the highest NPQ value measured during ETR vs E curve was extracted.

To measure the mean chl a-specific absorption coefficient (\bar{a}^*_{phy}), cells were harvested from 25 mL of culture by centrifugation (4000 g, for 15 min at 25°C) and the pellet obtained was resuspended in 1 mL of Milli-Q water. The optical density (OD) spectrum was then measured between 350 and 850 nm with 1 nm-increment using a spectrophotometer UV-Vis 6705 (Jenway). A cuvette filled with Milli-Q water was used as a reference (blank). The OD between 750 and 800 nm was used to correct for scattering. The absorption coefficient of particles ($ap(\lambda)$, in m^{-1}) was calculated according to the equation of Mitchell et al. (2003):

$$ap(\lambda) = \frac{2.303}{l} [OD_{sample}(\lambda) - OD_{blank}(\lambda) - OD_{null}(\lambda)]$$

where l is the cuvette pathlength, $OD_{sample}(\lambda)$ is the sample absorbance, $OD_{blank}(\lambda)$ is the reference absorbance and $OD_{null}(\lambda)$ is the OD between 750 and 800 nm used to correct for scattering. The depigmented particle absorption coefficient ($ad(\lambda)$, in m^{-1}) was obtained by bleaching the sample with a solution of sodium hypochlorite (0.1% active chlorine). The spectral absorption coefficient for cyanobacteria pigments ($a_{phy}(\lambda)$) was then calculated by subtracting $ad(\lambda)$ to $ap(\lambda)$. $a_{phy}(\lambda)$ was normalized to the chlorophyll a concentration to obtain the chl a-specific absorption ($a^*_{phy}(\lambda)$, in $m^2 (mg \text{ chl a})^{-1}$). Finally, the mean absorption coefficient between 400 and 700 nm (\bar{a}^*_{phy} , in $m^2 (mg \text{ chl a})^{-1}$) was calculated.

2.4 Chlorophyll a, carotenoids, phycoerythrin and phycocyanin content

Chlorophyll a (chl a) and carotenoids concentrations were measured by filtering known volume of sample through 25 mm GF/F glass-fiber filter. The filters were stored at – 80°C. Chl a and carotenoids were subsequently dark-extracted in 90% acetone, clarified by

centrifugation and quantified spectrophotometrically. Chl a and carotenoids concentrations were respectively calculated according to the equations of UNESCO (1966) and Strickland and Parsons (1972).

Phycoerythrin and phycocyanin contents were measured using in vivo fluorescence excitation spectra. Prior to the measurements, samples were treated with 50 μ M (final concentration) of 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU, Sigma Aldrich) and illuminated with a strong light to ensure the complete saturation of PSII and avoid variable fluorescence (Duysens, 1972). Spectra were recorded using a microplate reader infinite 200 (Tecan). Excitation light was provided at 5 nm bandwidth from 400 to 700 nm with emission recorded at 730 nm. Phycoerythrin and phycocyanin contents were determined by using respectively the fluorescence intensity measured after excitation at 565-575 and 645-655 nm (Seppälä, 2009). All spectra were normalized at the chl a peak (425 nm) for comparison.

2.5 Production of reactive oxygen species (ROS)

Intracellular ROS production was assayed by measuring the oxidation of the 2'-7'-dichlorohydrofluorescein diacetate (DCFH-DA, Sigma Aldrich) according to the protocol of Rajneesh et al. (2017) (steps 3-7) with small modifications. Cells were harvested from 5 mL of culture by centrifugation (4000 g for 15 min at 25°C) to obtain cell pellet and supernatant. Supernatant was discarded. Cell pellet was resuspended in 1 mL of phosphate buffer (50 mM) and 2.5 μ L of 2 mM DCFH-DA solubilized in ethanol was added. Two blanks were also prepared: 1) a cell pellet harvested from a 5 mL of sample centrifuged and resuspended in 1 mL phosphate buffer to correct for pigments fluorescence in the absence of DCFH-DA and 2) 1 mL of buffer with 2.5 μ L of 2 mM DCFH-DA to correct for background fluorescence from potential auto-oxidation of DCFH-DA. Samples and blanks were incubated at room temperature in the dark for 1h on a shaking table (15 rpm). OD₇₅₀ and fluorescence of samples

and blanks were measured using a microplate reader infinite 200 (Tecan). For fluorescence measurements, excitation light was provided at 485 nm and emission was registered between 515 and 600 nm to target fluorescence from 2',7'-dichlorofluorescein (DCF). The relative ROS production was calculated by averaging the fluorescence intensity measured between 525 and 535 nm after blanks correction and was normalized by OD₇₅₀.

2.6 Extraction and analysis of saxitoxins (STX)

Cyanobacteria were harvested from 25 mL of culture by centrifugation (4000 g for 15 min at 25°C) to obtain cell pellet and supernatant. Cell pellet was resuspended for toxins extraction in 1 mL of 0.1 N acetic acid. Cells were lysed using 0.25 g of glass beads with a ball mills (MM400, Retsch GmbH, Germany) for 30 min at 30 Hz. Samples were centrifuged at 15000 g for 15 min at 4°C. The supernatant was then filtrated through a Nanosep MF centrifugal devices 0.2 µm (4000 g for 15 min at 4°C).

Analysis of STXs was conducted by HPLC-FLD with a protocol adapted from the method of Van de Riet et al. (2009) which uses a reversed-phase LC with post-column oxidation. Samples were analyzed on a Zorbax bonus RP column (4.6 mm x 150 mm, 3.5 µm) using a mobile phase composed of two solvents: one containing 11 mM heptane sulfonate and 5.5 mM phosphoric acid (H₃PO₄) and the second containing 11 mM heptane sulfonate, 16.5 mM H₃PO₄ and 11.5% acetonitrile. The post-column oxidant was a mixture of 100 mM H₃PO₄ and 5 mM periodic acid. 0.75 M of nitric acid was used as a post-column acid. Fluorescence was monitored using an excitation light at 330 nm and emission at 390 nm. Six standard mixtures containing known increasing concentrations of toxins were prepared using solutions of gonyautoxin 1, 2, 3, 4 and 5 (respectively GTX1, GTX2, GTX3, GTX4 and GTX5), decarbamoylogonyautoxin 2 and 3 (dc-GTX2 and dc-GTX3), neosaxitoxin (Neo-STX), decarbamoylsaxitoxin (dc-STX) and saxitoxin (STX) purchased from the Institute for

Marine Biosciences (National Research Council, Halifax, NS, Canada). During HPLC measurements, each run of samples was preceded and followed by these standards mixtures and blanks. Toxins in samples were identified by comparison with the known retention time of each standard toxin (supplementary materials Fig. S1 and S2). Toxins in samples were quantified by comparison of their integrated area to a calibration curve built, for each standard toxin, from integrated areas and known concentrations in mixtures (supplementary materials Fig. S3). Intracellular toxin contents were normalized by chlorophyll a and concentrations were expressed per chlorophyll unit. However to be sure that observed variations did not reflect changes in chl content, toxins were also normalized by OD₇₅₀.

2.7 Extraction and analysis of DMSP, glycine betaine, methionine and proline

Cyanobacteria were harvested from 15 mL of culture by centrifugation (4000 g for 15 min at 25°C) to obtain cell pellet and supernatant. Cell pellet was resuspended for DMSP, glycine betaine (GB), methionine and proline extraction in 1 mL of 100% methanol (Honeywell). Cells were lysed using 0.25 g of glass beads with a ball mills (MM400, Retsch GmbH, Germany) for 20 min at 30 Hz. Samples were centrifuged at 10000 g for 2 min at 4°C. The supernatant was then filtrated through a Nanosep MF centrifugal devices 0.2 µm (10000 g for 2 min at 4°C).

Intracellular contents in DMSP, GB, methionine and proline were measured by LC-MS/MS on a LC System (UFLC XR, Shimadzu) coupled to a triple quadrupole mass spectrometer (4000 QTrap, ABSciex). A Hypersil GOLD HILIC column (150 x 3.0 mm, 3 µm, ThermoScientific) with a suited guard column was used to perform chromatography using a protocol adapted from Curson et al. (2018). The binary gradient was composed of water/acetonitrile (90:10; V/V) containing 4.5 mM ammonium formate (A) and water/acetonitrile (5:95, V/V) containing 5mM ammonium formate (B). Injection volume was

5 μL and flow rate equaled 0.4 mL min^{-1} . Column temperature was 30°C and that of sample 4°C . Elution gradient was 90% B (0-1 min), 90-45% B (1-8 min), 45% B (8-12 min) and 90% B (12-15 min). A positive ionization mode and MRM with two transitions per compounds was used. Compounds were quantified by comparison with 5-point calibration curves built with standards (Sigma Aldrich) dissolved in 100% methanol at concentrations ranging from 50 nM to 5000 nM. Measurements and data processing were carried out using the Analyst 1.6.3 (ABSciex) software. Intracellular contents in DMSP, GB, methionine and proline were normalized by chlorophyll a and concentrations were expressed per chlorophyll unit. However to be sure that observed variations did not reflect changes in chl content, they were also normalized by OD_{750} .

2.8 Extraction and analysis of mono and disaccharides

Cyanobacteria were harvested from 15 mL of culture by centrifugation (4000 g for 15 min at 25°C) to obtain cell pellets and supernatants. Cell pellets were resuspended in 1 mL of absolute ethanol and incubated for 3h at 65°C . Samples were then filtered through a Nanosep MF centrifugal devices $0.2\text{ }\mu\text{m}$ (8000 g for 1 min at 4°C) to remove salt. Ethanol was evaporated from the salt-free extracts under a stream of nitrogen at 65°C . Dried pellets were suspended in Milli-Q water and sugar contents were analyzed by gas chromatography (GC-FID, Agilent Technologies 6890N) following Adams et al. (1999). Four osidic residues were investigated: sucrose, fructose, glucose and D-trehalose (Sigma Aldrich). Intracellular contents in sucrose, fructose, glucose and D-trehalose were normalized by chlorophyll a and concentrations were expressed per chlorophyll unit. However to be sure that observed variations did not reflect changes in chl content, they were also normalized by OD_{750} .

2.9 Statistical analysis

For each parameter, differences between control and other culture conditions were tested using two-way ANOVA followed by post-hoc multiple comparisons using the Holm-Šidák method (Holm, 1979). Normality and equal variance were tested using respectively the Shapiro-Wilk (Royston, 1982) and Brown-Forsythe (Brown and Forsythe, 1974) tests. All analyses were performed using SigmaPlot 14.0.

3 Results

3.1 Growth and survival

Growth and survival time of *A. gracile* (Fig. 1A) and *C. raciborskii* (Fig. 1B) were inversely proportional to NaCl concentration in culture medium. In BG11, growth of *A. gracile* started after a 48h lag phase ($\mu=0.12 \text{ day}^{-1}$). In BG11 + 34 mM NaCl, the cell density of *A. gracile* slightly increased during the 72h of experiment and remained close to the BG11 lag phase ($\mu=0.07 \text{ day}^{-1}$). In BG11 + 9 g. L⁻¹ NaCl, the cell density of *A. gracile* moderately decreased after 24h exposure to NaCl and remained stable until the end of the experiment. In BG11 + 257 mM NaCl, the quantity of chl a dropped in the first 6h after the salt treatment and *A. gracile* only survived 24h. *C. raciborskii* only grew in BG11 and BG11 + 34 mM NaCl with a lower growth in BG11 + 34 mM NaCl ($\mu=0.03 \text{ day}^{-1}$) than in BG11 ($\mu=0.06 \text{ day}^{-1}$). In BG11 + 154 mM NaCl, *C. raciborskii* survived until the end of the experiment but the cell density declined over time as indicated by the drop in chl a concentration. In BG11 + 257 mM NaCl, *C. raciborskii* only survived 24h. *Dolichospermum* sp. survived until the end of the experiment in all culture media (Fig. 1C). There was no difference in *Dolichospermum* sp. growth in BG11 + 154 mM NaCl ($\mu=0.17 \text{ day}^{-1}$) and BG11 + 257 mM NaCl ($\mu=0.17 \text{ day}^{-1}$). Growth in BG11 + 308 mM NaCl and BG11 + 411 mM NaCl was lower than in BG11 + 154

mM NaCl and inversely proportional to NaCl amount ($\mu=0.10 \text{ day}^{-1}$ in BG11 + 308 mM NaCl and 0.08 day^{-1} in BG11 + 411 mM NaCl). In BG11 + 411 mM NaCl, the lag phase was longer than in other media.

3.2 Pigments

In all species, carotenoids concentration changed significantly over time and with NaCl treatments ($p<0.05$). There was a significant interaction between these factors ($p<0.001$). In *A. gracile*, carotenoids concentration differed from control conditions (BG11) only from 4 to 6h of NaCl exposure when carotenoids concentration became lower in NaCl containing media than in BG11 (Fig. 2A). In *C. raciborskii*, carotenoids concentration in BG11 + 34 mM NaCl changed in two steps (Fig. 2B). It increased after 4h exposure and recovered a level not significantly different from control conditions (BG11) after 6h exposure. Then, it increased again after 24h exposure and remained stable until the end of the experiment. In BG11 + 154 mM NaCl, carotenoids concentration was not significantly different from control conditions and remained relatively stable during the 72h of experiment. In BG11 + 257 mM NaCl, carotenoids concentration decreased from 24h exposure. In *Dolichospermum* sp., carotenoids concentration at $t=0$ was around 2 times lower than in *A. gracile* and *C. raciborskii*. In BG11 + 257 mM NaCl was not significantly different from control conditions (BG11 + 154 mM NaCl) (Fig. 2C). In BG11 + 308 mM NaCl, carotenoids concentration decreased from 48 to 72h exposure. In BG11 + 411 mM NaCl, carotenoids concentration increased after 4h exposure. Then, it recovered a level not significantly different from control conditions and remained relatively stable until the end of the experiment.

Phycocyanin (PC) was the dominant phycobiliprotein in *A. gracile* and *C. raciborskii* while phycoerythrin (PE) dominated in *Dolichospermum* sp. (Fig. 2). In all species, PC and PE content varied significantly over time and with NaCl treatments ($p<0.001$). There was a

significant interaction ($p < 0.001$) between these factors. In *A. gracile*, exposure to NaCl decreased PE and PC contents proportionally to NaCl concentration in culture media from 24 to 48h exposure and from 6 to 48h exposure respectively (Fig. 2D, G). After 72h exposure, in media in which *A. gracile* survived PC and PE content did not seem to be further impacted by NaCl. In *C. raciborskii*, variation in PC and PE content in response to NaCl exposure was biphasic (Fig. 2E, H). After 1h exposure, PC and PE content were higher in NaCl containing media than in BG11. Then, from 4 to 24h exposure, there was no significant difference between BG11 and NaCl treatments. After 24h exposure, PC content in BG11 remained stable while it became inversely proportional to NaCl amount in the other media. PE content followed the same pattern as PC but after 48h exposure. In *Dolichospermum* sp., PC and PE content showed similar patterns of variation (Fig. 2F, I). In control conditions (BG11 + 154 mM NaCl), these contents remained stable. By contrast, from 1 to 24h exposure in other media, PC and PE content was proportional to the NaCl amount. After 48h exposure, PC and PE recovered levels not significantly different from control conditions.

3.3 Photosynthetic parameters

The maximum quantum yield (F_v/F_m), measured in the diverse NaCl treatments, ranged from 0.59 to 0.20 in *A. gracile* (Fig. 3A), 0.61 to 0.04 in *C. raciborskii* (Fig. 3B) and 0.52 to 0.23 in *Dolichospermum* sp. (Fig. 3C). In all species, F_v/F_m changed significantly over time and with NaCl treatments ($p < 0.001$). There was a significant interaction ($p < 0.001$) between these factors. When *A. gracile* was exposed to BG11, BG11 + 34 mM NaCl and BG11 + 154 mM NaCl, F_v/F_m stayed high with few differences between treatments. By contrast BG11 + 257 mM NaCl resulted in cell death after 48h exposure and F_v/F_m decrease drastically (from 0.52 to 0.26) during the last day of life (24h). When *C. raciborskii* was exposed to BG11 + 34 mM NaCl and BG11 + 154 mM NaCl, F_v/F_m increased proportionally to the amount of NaCl

during the first 24h exposure. After 48h in BG11 + 34 mM NaCl, F_v/F_m recovered values not significantly different from control conditions. Inversely, in BG11 + 154 mM NaCl, F_v/F_m dropped from 0.57 to 0.17 and reached values close to zero after 72h exposure. BG11 + 257 mM NaCl resulted in cell death after 48h exposure. F_v/F_m started to decrease after 1h exposure, remained stable during the first 6h and dropped from 0.37 to 0.19 during the last day of life (24h). In *Dolichospermum* sp., F_v/F_m increased over time in all media. During the first 4h exposure, F_v/F_m was not affected by NaCl and only decreased proportionally to the amount of NaCl between 6 and 48h exposure. In BG11 + 257 mM NaCl and BG11 + 308 mM NaCl, F_v/F_m recovered after 72h exposure while, in BG11 + 411 mM NaCl, F_v/F_m stayed lower than in control conditions until the end of the experiment.

In all species, the maximal light utilization efficiency (α) and maximum electron transport rate (ETR_m) varied conjointly. α ranged from 0.0006 to 0.0045 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1})^{-1}$ in *A. gracile* (Fig. 3D), 0.0003 to 0.0038 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1})^{-1}$ in *C. raciborskii* (Fig. 3E) and 0.0003 to 0.0025 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1})^{-1}$ in *Dolichospermum* sp. (Fig. 3F). ETR_m ranged from 0.32 to 3.35 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ in *A. gracile* (Fig. 3G), 0.11 to 2.57 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ in *C. raciborskii* (Fig. 3H) and 0.22 to 1.84 $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ in *Dolichospermum* sp. (Fig. 3I). In *A. gracile* and *C. raciborskii*, α and ETR_m changed significantly over time and with NaCl treatments ($p < 0.001$). There was a significant interaction ($p < 0.001$) between these factors. In *A. gracile*, α and ETR_m increased in response to salt stress after 4-6h exposure. In BG11 + 34 mM NaCl, this increase did not persist as α and ETR_m recovered values not significantly different from the control after 24h exposure. In BG11 + 154 mM NaCl, α recovered after 48h while ETR_m recovered after 72h. In BG11 + 257 mM NaCl, α and ETR_m did not recover and decreased after 24h exposure. In *C. raciborskii* exposed to BG11 + 34 mM NaCl, α and ETR_m were not significantly different from control conditions. While α and ETR_m decreased in BG11 during

the first 4-6 hours, in BG11 + 154 mM NaCl, α and ETR_m stayed high and reached their highest values after 24h exposure. Then, they decreased until the end of the experiment. In BG11 + 257 mM NaCl, α did not change over time and was not significantly different from control conditions. ETR_m was significantly lower than in control conditions only after 24h exposure. In *Dolichospermum* sp., α and ETR_m changed significantly over time ($p < 0.001$) but there was no significant difference between NaCl treatments.

In *A. gracile* and *C. raciborskii*, maximum non-photochemical quenching of fluorescence (NPQ_{max}) changed significantly over time and with NaCl treatments ($p < 0.001$). There was a significant interaction ($p < 0.001$) between these factors. In *A. gracile*, NPQ_{max} in BG11 + 34 mM NaCl was not significantly different from NPQ_{max} in BG11 (Fig. 3J). In BG11 + 154 mM NaCl, NPQ_{max} increased significantly after 48h exposure and stayed high until the end of the experiment. In BG11 + 257 mM NaCl, NPQ_{max} increased significantly after 24h exposure. In *C. raciborskii*, NPQ_{max} in BG11 + 34 mM NaCl and BG11 + 257 mM NaCl was not significantly different from NPQ_{max} in control conditions (Fig. 3K). In BG11 + 154 mM NaCl, NPQ_{max} increased significantly only after 24h exposure. In *Dolichospermum* sp., NPQ_{max} did not change significantly over time and with NaCl treatments (Fig. 3L).

3.4 Reactive oxygen species (ROS)

In all species, ROS production (Fig. 4) changed significantly over time and with NaCl treatments ($p < 0.05$). There was a significant interaction ($p < 0.001$) between these factors. In *A. gracile*, ROS production in BG11 + 34 mM NaCl was not significantly different from control conditions (Fig. 4A). In BG11 + 154 mM NaCl and BG11 + 257 mM NaCl, ROS production of *A. gracile* was proportional to NaCl amount. In BG11 + 154 mM NaCl, ROS production was biphasic. It started after 6h exposure, remained stable between 6 and 48h exposure and increased after 72h. In BG11 + 257 mM NaCl, ROS production occurred between 4 and 6h

exposure. Then, ROS production declined. In *C. raciborskii*, ROS production in BG11 + 34 mM NaCl was not significantly different from control conditions except after 24h exposure where it was higher than in BG11 (Fig. 4B). In BG11 + 154 mM NaCl and BG11 + 257 mM NaCl, ROS production was biphasic. In BG11 + 154 mM NaCl, a first increase in ROS production occurred after 1h exposure. Then, ROS declined after 4h exposure and a second phase of ROS production took place from 6 to 24h exposure. After, 48h exposure, ROS decreased and became lower than in BG11. In BG11 + 257 mM NaCl, ROS production increased after 1h exposure and remained stable between 1 and 6h exposure. A second peak of ROS production occurred after 24h exposure. In *Dolichospermum* sp., ROS production in BG11 + 257 mM NaCl was not significantly different from control conditions (BG11 + 154 mM NaCl) (Fig. 4C). In BG11 + 308 mM NaCl and BG11 + 411 mM NaCl, ROS production was proportional to NaCl amount. In BG11 + 308 mM NaCl, ROS production started after 1h exposure and reached its maximum after 4h exposure. It remained stable between 4 and 24h exposure and recovered its basal level after 48h exposure. In BG11 + 411 mM NaCl, ROS production started after 1h exposure and increased with a maximum reached after 24h exposure. Then, ROS production decreased to a level not significantly different from control conditions and remained stable until the end of the experiment.

3.5 Toxins

In *A. gracile* samples, two toxin analogs, neosaxitoxin (Neo-STX) and saxitoxin (STX), were detected (supplementary materials Fig. S4). The total amount of paralytic shellfish toxins (PSTs) (Fig. 5A) as well as Neo-STX (Fig. 5B) and STX (Fig. 5C) concentrations varied significantly over time and with NaCl treatments ($p < 0.05$). There was a significant interaction between these factors ($p < 0.001$). Neo-STX and STX showed similar variations and thus changes in PSTs concentrations were the result of changes in both toxins. PSTs,

Neo-STX and STX concentrations in BG11 + 34 mM NaCl and BG11 + 257 mM NaCl were not significantly different from control conditions in BG11. In BG11 + 154 mM NaCl, PSTs, Neo-STX and STX concentrations were significantly higher than in control conditions and increased with exposure time to NaCl. At t=0, the relative proportions of Neo-STX and STX were respectively 75% and 25% and Neo-STX stayed the dominant analog in all treatments (Fig. 6).

In *C. raciborskii*, 3 toxins were detected: decarbamoylogonyautoxin 2 (dc-GTX2), Neo-STX and STX (supplementary materials Fig. S5). An unidentified peak appearing between GTX4 and GTX1 standards was also detected during toxin analyses. The quantity of PSTs produced by *C. raciborskii* was 5-6 times lower than in *A. gracile*. The total amount of PSTs (Fig. 5B) and concentrations of Neo-STX (Fig. 5D), STX (Fig. 5F), and dc-GTX2 (Fig. 5G) changed significantly over time and with NaCl treatments ($p < 0.05$). There was a significant interaction between these factors ($p < 0.001$). PSTs, Neo-STX, STX and dc-GTX2 concentrations in BG11 + 34 mM NaCl were not significantly different from control conditions in BG11. In BG11 + 154 mM NaCl, PSTs and Neo-STX contents both decreased with exposure time to NaCl. While in control conditions STX concentration increased from 6h, in BG11 + 154 mM NaCl STX content stayed low. By contrast, dc-GTX2 became significantly higher than in control conditions after 6h exposure and then gradually decreased from 24h exposure. From 48h exposure, dc-GTX2 was the only analog detected in this culture medium. In BG11 + 257 mM NaCl, PSTs, Neo-STX, STX and dc-GTX2 concentrations decreased with exposure time to NaCl. At t=0 in BG11, the relative proportions of Neo-STX, STX and dc-GTX2 were respectively 74%, 8% and 18% (Fig. 6). Neo-STX stayed the dominant analog in all treatments during the first 24h exposure. In BG11 + 34 mM NaCl and BG11 + 257 mM NaCl, the relative proportions of Neo-STX, STX and dc-GTX2 followed the same temporal variations than in control conditions in BG11. By contrast, in BG11 + 154 mM

NaCl, dc-GTX2 proportion increased with time exposure to NaCl while those of Neo-STX and STX decreased Fig. 6G). Similar trends of variation were obtained when toxins were normalized by OD₇₅₀ (supplementary materials Fig. S6 and S7). This indicates that observed variations were the results of toxins contents variations and not the results of changes in intracellular chl a concentrations. No toxin was detected in *Dolichospermum* sp. (supplementary materials Fig. S8).

3.6 Methionine

In all species, methionine content varied significantly over time and with NaCl treatments ($p < 0.05$). There was a significant interaction ($p < 0.001$) between these factors. In *A. gracile* (Fig. 7A) and *C. raciborskii* (Fig. 7B), methionine content in BG11 + 34 mM NaCl was not significantly different from control conditions in BG11. In *A. gracile*, methionine content was proportional to NaCl concentration from 6 to 48h exposure. In BG11 + 154 mM NaCl, methionine concentration increased strongly after 72h exposure. In *C. raciborskii*, methionine content was inversely proportional to NaCl concentration from 6 to 24h exposure. After 48h exposure, methionine content in BG11 + 154 mM NaCl increased and methionine became proportional to NaCl concentration. After 72h exposure, methionine content was again inversely proportional to NaCl concentration. In *Dolichospermum* sp. (Fig. 7C), from 6 to 24h exposure, only the methionine content in BG11 + 411 mM NaCl was significantly different from control conditions in BG11 + 154 mM NaCl. After 48h exposure, only the methionine content in BG11 + 257 mM NaCl was significantly lower than in control conditions. After 72h exposure, methionine contents in BG11 + 257 mM NaCl and BG11 + 411 mM NaCl were both significantly lower than in control conditions while in BG11 + 308 mM NaCl, methionine content increased.

3.7 DMSP, glycine betaine, proline, sucrose, fructose, glucose and D-trehalose

In all species, DMSP and glycine betaine concentrations were below the detection limit (50 nM) in all treatments. In *A. gracile* (Fig. 7D) and *C. raciborskii* (Fig. 7E), proline content changed significantly over time and with NaCl treatments ($p < 0.05$). There was a significant interaction ($p < 0.001$) between these factors. By contrast, in *Dolichospermum* sp. (Fig. 7F), NaCl treatments did not significantly modify proline content and the maximum level of proline produced was threefold lower than in *A. gracile* and *C. raciborskii*. In *A. gracile*, proline content increased dose-dependently with NaCl after 24h exposure. After 48h exposure, proline content in BG11 + 154 mM NaCl recovered a level not significantly different from control conditions while it increased in BG11 + 34 mM NaCl. After 72h exposure, proline content was again proportional to NaCl concentration. In *C. raciborskii*, proline content decreased dose-dependently with NaCl and was inversely proportional to NaCl concentration after 24h exposure. After 48h exposure, proline content in BG11 + 34 mM NaCl recovered a level not significantly different from control conditions while it increased in BG11 + 154 mM NaCl. After 72h exposure, proline content in BG11 + 154 mM NaCl decreased but it was still significantly higher than in control conditions. In *A. gracile* and *C. raciborskii*, sucrose, fructose, glucose and D-trehalose concentrations were below the detection limit ($8 \mu\text{g L}^{-1}$) in all treatments. In *Dolichospermum* sp., sucrose concentration was below the detection limit in BG11 + 154 mM NaCl and BG11 + 257 mM NaCl. By contrast, in BG11 + 308 mM NaCl and BG11 + 411 mM NaCl sucrose was detected from 6h exposure (Fig. 8). Sucrose concentration was higher in BG11 + 411 mM NaCl than in BG11 + 308 mM NaCl. In both media, sucrose concentration remained relatively stable until the end of the experiment. The other mono and disaccharides (fructose, glucose and D-trehalose) were below the detection limit in all treatments. Normalization by OD₇₅₀ confirmed that observed variations were the results of changes in intracellular methionine, proline and sucrose in

response to NaCl treatments and not related to modifications in intracellular chl a concentrations (supplementary materials Fig. S9 and S10).

4 Discussion

4.1 Growth and salt tolerance

A. gracile, *C. raciborskii* and *Dolichospermum* sp. presented distinct tolerance and survival capacities to salinity stress. *C. raciborskii* seemed to be the less tolerant as it only survived 24h when it was exposed to the highest salinity and the cell density declined over time after exposure to BG11 + 154 mM NaCl. By opposition, the brackish water species (*Dolichospermum* sp.) was the most tolerant. It survived and grew in all tested salinities. In all species, NaCl dose-dependently slowed down growth. However in *A. gracile* and *C. raciborskii*, growth seemed to be inhibited when $[\text{NaCl}] \geq 154 \text{ mM}$ while in *Dolichospermum* sp., growth at high salinities was reduced but not inhibited. The slow-down or inhibition of growth by NaCl has been frequently observed in several species of cyanobacteria (Carneiro et al., 2013; Chen et al., 2015; Georges des Aulnois et al., 2019; Hagemann, 2011; Pomati et al., 2004b; Swapnil et al., 2017; Tonk et al., 2007). This happens because high NaCl concentrations lower the water potential in the culture medium to an extent that makes it difficult for cyanobacteria to obtain and retain water inside their cells. Turgor, the driving force for growth, is consequently reduced or lost and growth is affected (Hagemann, 2011). Growth reduction may also results from the extra energy expenditure due to Na^+/H^+ antiporters activity and implementation of the "salt-out" strategy (Carneiro et al., 2013; Hagemann, 2011). In the present study, intensity of growth slow-down (or inhibition) also depended on cyanobacteria capacity to maintain their photosynthetic activity under salt stress conditions (see section 4.3. for more details). The higher salt tolerance of *Dolichospermum* sp. (*Anabaena* sp.) in comparison to *C. raciborskii* is consistent with the Tonk et al. (2007)'s data

compilation on the salt tolerance of cyanobacteria as well as the previously reported *C. raciborskii* preference for low salinity conditions with optimal growth in freshwaters to oligohaline conditions (Antunes et al., 2015).

4.2 Pigments

Carotenoids content was differently impacted by NaCl treatments in the freshwater (*A. gracile*, *C. raciborskii*) and brackish water (*Dolichospermum* sp.) species. While in *A. gracile*, NaCl reduced carotenoids content from 4 to 6h exposure, in *C. raciborskii* and *Dolichospermum* sp., variations in carotenoids content differed among salinity treatments without any obvious relationships with NaCl concentration. In cyanobacteria, carotenoids are photosynthetic pigments described under salt stress as having a protective role on chlorophyll as well as antioxidant effects on reactive oxygen species (ROS) (Bhargava and Srivastava, 2013). However, in cyanobacteria, effects of NaCl stress on carotenoids content do not seem to be universal. In freshwater cyanobacteria, some authors found a decrease in carotenoids content with increasing NaCl concentration (Chen et al., 2015) while others observed an increase (Kumar et al., 2015; Schubert et al., 1993). A production of carotenoids to protect cells against ROS could not be confirmed in the present study since carotenoids trends did not correspond to ROS variations. For instance, in *Dolichospermum* sp. cultures exposed to 411 mM NaCl, there was an increase in ROS production from 4 to 24h while on the same period carotenoids content decreased from 4 to 6h and then remained relatively stable until 48h. Besides, although ROS variations were biphasic in *A. gracile* and *C. raciborskii*, these species presented completely different trends in their carotenoids content.

While in *Dolichospermum* sp. PC and PE contents increased within the first 24h exposure to NaCl, in both *A. gracile* and *C. raciborskii*, PC and PE contents decreased proportionally to NaCl concentration but with different timing. The decrease of these

phycobiliproteins in response to salinity stress has been described in a freshwater strain of *Microcystis aeruginosa* (Chen et al., 2015) and several species of freshwater filamentous cyanobacteria (Kumar et al., 2015; Lu and Vonshak, 2002). In cyanobacteria, phycobiliproteins constitute phycobilisomes that serve as the primary light-harvesting antenna for photosystems II. Under stress conditions, the constituents of phycobilisomes are often degraded. This degradation could provide the cell with amino acids used for the synthesis of proteins required for the acclimation processes (Grossman et al., 1993). Modifications in PC and PE contents can also have important consequences on cyanobacteria photosynthetic capacities because this can modify energy transfers from phycobilisomes to photosystems II (Schubert et al., 1993; Sudhir and Murthy, 2004; Zhang et al., 2010).

4.3 Photosynthesis

PSII activity was differently impacted by NaCl treatments in *A. gracile*, *C. raciborskii* and *Dolichospermum* sp. In *A. gracile* and *C. raciborskii*, partly due to phycobilisomes modifications, the treatment with the highest NaCl concentration (BG11 + 257 mM NaCl) strongly impacted PSII functioning. The maximum quantum yield (F_v/F_m) and maximum electron transport rate (ETR_m) were reduced and the maximum non-photochemical quenching of fluorescence (NPQ_{max}) increased in *A. gracile*. These results suggest that the stress level generated by 257 mM NaCl was sufficiently high to down regulate the photosynthetic activity and to stimulate photoprotective mechanisms (NPQ). None of these parameters recovered during the experiment leading to cell death. In the other treatments, F_v/F_m was not impacted. The maximal light utilization efficiency (α) and ETR_m increased but recovered with a recovery time inversely proportional to NaCl concentration. NPQ_{max} increased only in BG11 + 154 mM NaCl. Hence, exposure to 34 and 154 mM NaCl reversibly modified photosynthetic activity and slowed down growth. In *C. raciborskii*, F_v/F_m first increased

proportionally to the amount of NaCl during the first 24h exposure. Then, F_v/F_m recovered in BG11 + 34 mM NaCl while in BG11 + 154 mM NaCl, it strongly decreased. α and ETR_m were not impacted in BG11 + 34 mM NaCl. In BG11 + 154 mM NaCl, they decreased after 24h exposure and did not recover. NPQ_{max} increased simultaneously to α and ETR_m decrease in BG11 + 154 mM NaCl. While exposure to 34 mM NaCl did not severely impacted photosynthetic activity and slightly reduced *C. raciborskii* growth, 154 mM NaCl treatment irreversibly inhibited photosynthesis (both the capacity to absorb light and transport electrons) and led to cell death. In the brackish water species (*Dolichospermum* sp.), NaCl had a weak effect on PSII activity. Growth was thus slowed down but maintained in all salt treatments. α , ETR_m and NPQ_{max} were not affected. Only F_v/F_m decreased proportionally to the amount of NaCl and recovered in BG11 + 257 mM NaCl and BG11 + 308 mM NaCl but not in BG11 + 411 mM NaCl.

Like in the present study, different effects of salinity stress on cyanobacteria PSII activity have been reported in literature. Although it has been found that PSII of a freshwater strain of *Synechocystis* sp. (PCC6803) were not impacted by a salt stress of 550 mM NaCl (i.e. 32.14 g L⁻¹ NaCl) (Jeanjean et al., 1993), the majority of studies on freshwater cyanobacteria observed a decrease of PSII activity under salinity stress (e.g. Chen et al., 2015; Gong et al., 2008; Lu and Vonshak, 2002; Ross et al., 2019). In accordance with the present study, when salinity concentration was sufficiently high to induce a stress, a decreased F_v/F_m was observed in *Spirulina platensis* (Gong et al., 2008; Lu et al., 1999) and *M. aeruginosa* (Georges des Aulnois et al., 2020; Ross et al., 2019). In line with the *A. gracile* and *C. raciborskii* results under high salinity concentrations, a decreased ETR_m under salinity stress has also been found in a freshwater strain of *M. aeruginosa* (Chen et al., 2015).

The underlying mechanisms of the salt-stress reduction in PSII activity, and photosynthesis in general, are not completely resolved in cyanobacteria. In the freshwater

cyanobacteria: *Synechocystis* sp. and *S. platensis*, it has been suggested that the decreased PSII activity under salt stress is associated with a transition from state 1 to state 2 i.e. a modification in the distribution and transfer of excitation energy from phycobilisomes to PSII and photosystem I (PSI) (Gong et al., 2008; Schubert et al., 1993; Schubert and Hagemann, 1990). In *S. platensis*, Gong et al. (2008) found that salt stress resulted in an increase of state 2 stability at the donor side and this was associated with a dissociation of the PSII extrinsic protein (PsbO) from the thylakoid membrane. The transition to state 2 involves the functional decoupling of phycobilisomes from PSII and their reassociation with PSI. Light energy transfer to PSII is thus reduced in favor of PSI (Joshua and Mullineaux, 2004) and the resulting F_v/F_m decreases (Ogawa et al., 2017). PSII is responsible for the production of NADPH, whereas PSI is involved in ATP generation (Bhargava and Srivastava, 2013). This transition to state 2 and the resulting boost in PSI activity would thus increase the relative amount of ATP produced by photosynthesis. This ATP may be used to fuel P-type ATPases and to respond to the over demand of ATP for Na^+ extrusion and restoration of cellular homeostasis (Bhargava and Srivastava, 2013; Hagemann, 2011). The increase in NPQ_{max} observed in *A. gracile* and *C. raciborskii* is consistent with the occurrence of this state transition because it has been shown in many studies that non photochemical quenching in cyanobacteria largely reflects changes in the PSII fluorescence yield as a result of the state transition mechanism (Lu and Vonshak, 2002 and references therein). In addition to state transition, the decreased F_v/F_m may be due to PSII reaction centers inactivation and inhibition of electron transport at both donor and acceptor sides (Gong et al., 2008; Lu et al., 1999; Lu and Vonshak, 2002). In *S. platensis*, it has been shown that salt stress induced a F_v/F_m decrease associated with a modification of the plastoquinone Q_B niche at the acceptor side (i.e. one of the PSII electron transporters) (Gong et al., 2008).

4.4 Reactive oxygen species (ROS)

ROS production was observed in the three tested cyanobacteria species when salinity was sufficiently high to induce a stress. However, the ROS accumulation timing and sequence differed among the three species. In the freshwater species (*A. gracile* and *C. raciborskii*), ROS accumulation was biphasic. In *A. gracile*, the time at which the first phase started was dependent on NaCl concentration. The higher the NaCl concentration, the sooner ROS accumulated. This trend was not observed in *C. raciborskii* in which ROS accumulation started after 1h exposure at any NaCl concentration. A biphasic ROS accumulation induced by salinity has been previously reported in the freshwater cyanobacteria *Anabaena fertilissima* (Swapnil et al., 2017). According to Swapnil et al. (2017), the first phase would be produced by NADPH oxidases located in the plasma membrane as well as cell wall peroxidases. These ROS would act as signal molecules to regulate ion homeostasis. The second phase would result from altered photosynthetic activity (Swapnil et al., 2017). Altered photosynthetic activity caused by salt stress may generate electrons in excess, which associated with changes at the electron acceptor sides, reduces O₂ and forms ROS (Bhargava and Srivastava, 2013; Swapnil and Rai, 2018). In the same way, a salinity induced increase in respiration may over reduce the ubiquinone pool and lead to electron leakage from complex I and III to O₂ which generates ROS (Swapnil and Rai, 2018; Swapnil et al., 2017). Results from the present study are insufficient to confirm that the first ROS accumulation in *A. gracile* and *C. raciborskii* acted as signal molecules because no measurement of intracellular ionic concentration or NADPH oxidases and cell peroxidases activity has been made. However, the results seem to confirm that ROS production was related to photosynthetic activity. In *A. gracile*, timing of variation in ROS accumulation corresponded to changes in α and ETR_m, while in *C. raciborskii*, it corresponded in part to changes in F_v/F_m and ETR_m. In *Dolichospermum* sp., ROS accumulation was not biphasic and did not seem to be related to PSII activity because it

started before any visible variation in F_v/F_m , α and ETR_m . Additional measurements of PSI activity and respiration would be required to better understand the processes behind this ROS production. The buildup of ROS, such as singlet oxygen (1O_2), superoxide anion (O_2^-), hydrogen peroxide (H_2O_2) and hydroxyl radical (OH^\cdot), may be problematic for prokaryotes such as cyanobacteria because these compounds have the potential to damage cell constituents. It has been shown that ROS production is related to several cellular toxic processes including damages to proteins, membrane lipid peroxidation, enzyme inactivation and DNA breakage (Latifi et al., 2009). ROS accumulation can also lead to the degradation of cellular membranes and consequent toxin release (Ross et al., 2006). This could not be confirmed in the present study because timing of ROS accumulation did not match with the changes observed in toxin content.

4.5 Toxins

Changes in cyanobacteria toxin content in response to environmental conditions are an important consideration when estimating the risk of animal and human exposure. In the present study, toxin content of *A. gracile* did not seem to be affected in a dose dependent-manner by salinity variations, since no significant changes in PSTs, Neo-STX and STX concentrations and proportions has been observed at the lowest as well as the highest NaCl concentrations tested. Only exposure to BG11+ 154 mM NaCl induced a significant increase in toxin content associated with an increase in ROS, methionine and proline contents after 72h exposure. There was also no apparent relationship between growth rate and toxin content. To our knowledge, *A. gracile* has not been previously used as a model species to study salinity effect on the production of STX and its analogs. Light, temperature and nutrients are the main abiotic factors that have been investigated in *Aphanizomenon* spp. (see Cirés and Ballot, 2016 for a review). The absence of an evident relationship with growth rate is consistent with

previous studies, since to date, no clear relationship between toxin content and growth rate has been found in *Aphanizomenon spp.* (Casero et al., 2014; Cirés and Ballot, 2016; Preußel et al., 2009). Toxin production by *C. raciborskii* seemed to be more sensitive to salinity variations. A decreased PSTs concentration was observed rapidly after exposure to the two highest NaCl levels tested. The sensitivity of STX content to NaCl stress in *C. raciborskii* is in agreement with previous studies [on the Brazilian freshwater *C. raciborskii* strain T3](#) (Carneiro et al., 2013; Ongley et al., 2016; Pomati et al., 2004b). However, results from the different studies are not completely consistent, since some authors found that NaCl stress resulted in intracellular STX accumulation in a dose-dependent manner (Carneiro et al., 2013; Pomati et al., 2004b) while, others observed a decrease (Ongley et al., 2016).

Although it is well known that STX, and its analogs, perturb voltage-gated Na⁺ channels in vertebrate cells, the stimuli inducing (or repressing) their production within cyanobacteria cells as well as their metabolic role for the producing organisms remain unclear (D'Agostino et al., 2016; Ongley et al., 2016; Soto-Liebe et al., 2012). The fact that STX seems to respond to salinity variations, more precisely intracellular sodium levels, led to think that STX biosynthesis might be influenced by the protective mechanisms that cyanobacteria use to maintain homeostasis or that PSTs themselves could interact with ion transport (Pomati et al., 2004b; Soto-Liebe et al., 2012). [In the Brazilian freshwater *Raphidiopsis brookii* strain D9](#), exposure to NaCl has been shown to stimulate the transport of STX outside of the cell likely through an active transport involving the multidrug and toxic compound extrusion (MATE) family of transporters (Soto-Liebe et al., 2012). This toxin transport would act as a protective mechanism to face salinity variations. However, by examining expression of genes involved in STX biosynthesis (*sxtA*) along with that of MATE transporters in *R. brookii* D9 and *C. raciborskii* T3, Ongley et al. (2016) found that both species showed opposing transcriptional responses to the same NaCl stress. In the present study, extracellular level of

STX has not been monitored and it could not be confirmed whether any extracellular transport of STX occurred in response to NaCl exposure. *Dolichospermum* sp. did not produce any toxins under the studied conditions, and yet was the most salt tolerant strain. This suggests the involvement of other metabolites to ensure the protection against salinity stress.

In *C. raciborskii*, results from the present study showed shifts in STX and dc-GTX2 proportions after exposure to 154 mM NaCl. Few studies have investigated how the different PSTs analogs produced by a single cyanobacteria species may vary in response to NaCl stress exposure. In *C. raciborskii* T3, Carneiro et al (2013) observed that NaCl modified the ratio of STX to Neo-STX. The relative proportion of STX increased but Neo-STX remained the major analog. The physiological reason of this shift remained unexplained. However, in *R. brookii* D9, Soto-Liebe et al. (2012) indicated that in the presence of the ionophore monensin that removes the Na⁺ gradient between the cytoplasm of cyanobacteria cells and culture medium, GTX2/3 levels were not impacted while, after 1h exposure, intracellular levels of STX decreased. These authors suggested that STX synthesis is sensitive to the Na⁺ gradient or that high Na⁺ concentrations may have affected enzymes involved in STX biosynthesis (Soto-Liebe et al., 2012). This could explain why STX, which is the precursor of both Neo-STX and GTX2/3 (D'Agostino et al., 2014; Soto-Liebe et al., 2010), can be differently affected by NaCl exposure than Neo-STX and GTX2/3. In the case of STX and dc-GTX2, the situation is different because STX is not the direct precursor of dc-GTX2 (D'Agostino et al., 2014). The suggestions of Soto-Liebe et al. (2012) remain, nonetheless, plausible. Distinct degradation rates have also been suggested to explain imbalance among PSTs analogs (Pomati et al., 2004a). Further research is, however, needed to better understand the underlying processes responsible for these shifts in STX and dc-GTX2 as well as why similar modifications have not been observed after exposure to 257 mM NaCl. Since toxicity of PSTs analogs is different (Wiese et al., 2010), the shifts in the relative proportions of PSTs analogs observed in *C.*

raciborskii in response to NaCl exposure means that salinity has the potential to modify the toxicity of this species. This is an important consideration when evaluating the risk of toxicity for animals and human beings.

Ongley et al. (2016)'s results, associated with those from the present study, showed that cyanobacteria with PSTs analogs in common can differently respond to the same salinity stress. This means that the physiological response involved in PSTs biosynthesis under NaCl stress is species-specific.

4.6 *Methionine and compatible solutes*

Methionine is essential in all organisms as it is both a proteinogenic amino acid and a component of the cofactor S-adenosyl methionine (SAM) (Ferla and Patrick, 2014). In cyanobacteria, PSTs would be biosynthesized via a pathway requiring 3 precursors: arginine, acetate (via acetyl-coenzyme A) and methionine methyl (via SAM) (Kellmann et al., 2008). In the present study, variations in methionine content in response to NaCl exposure were species-specific. The smallest variations were observed in *Dolichospermum* sp. Methionine content in *A. gracile* and *C. raciborskii* showed opposite variations during the first 24h of NaCl exposure. In *A. gracile*, methionine content increased in a dose dependent-manner with NaCl concentration. By contrast, it decreased in *C. raciborskii*. A potential link between the available methionine pool and PSTs content was difficult to establish with the present results. The hypothesis that, in comparison to *C. raciborskii*, toxin content in *A. gracile* was less sensitive to NaCl because methionine content increased in response to NaCl exposure whereas it decreased in *C. raciborskii* is plausible. However, this does not explain why, in *C. raciborskii*, STX level remained high in the cells presented the lowest methionine content (i.e. in BG11 + 257 mM NaCl) while in BG11 + 154 mM NaCl, STX concentration was low with more methionine available. Beside their role in toxin biosynthesis, methionine enters in

proteins composition. Variations in methionine content under salt stress, in toxic as well as non-toxic cyanobacteria, may thus result from protein synthesis alteration and anabolic processes. In stressed cyanobacteria cells, it has been shown that protein synthesis is usually modified leading to reduced synthesis of most normal cell proteins while synthesis of a set of special stress proteins is induced or enhanced (Fulda et al., 1999). Production of free methionine takes place in two possible ways: synthesis via homocysteine and release from protein degradation by proteases (Gröne and Kirst, 1992).

Under salt exposure, it has been shown that cyanobacteria may accumulate compatible solutes to compensate for differences in water potential between their intracellular medium and the surrounding environment (Hagemann, 2011; Klahn and Hagemann, 2011). These compatible solutes are of different chemical nature including carbohydrates, polyols, organosulfur compounds, amino acids and their derivatives (Oren, 2007). In the present study, the presence and potential accumulation of 7 compounds (DMSP, glycine betaine, proline, sucrose, fructose, glucose and D-trehalose) previously described as compatible solutes in microalgae and cyanobacteria was investigated. None of the three cyanobacteria species presented a detectable accumulation of DMSP, glycine betaine, fructose, glucose or D-trehalose in response to NaCl exposure. DMSP has been described as a constitutive compatible solute (Stefels, 2000). It has been found to increase with salinity in several microalgae species and has been detected in several cyanobacteria genera including *Anabaena* species and *Aphanizomenon flos-aquae* (Oren, 2007; Steinke et al., 2018). However, the measured concentrations in cyanobacteria were low and it is still unclear to what extent cyanobacteria rely on DMSP for osmotic balance (Oren, 2007). Considering the environment from which the three cyanobacteria species were isolated (fresh- and brackish waters), it is not surprising that they did not accumulate glycine betaine. Indeed, among cyanobacteria, a correlation has been found between their final salt tolerance and major compatible solute they

accumulate. Glycine betaine is usually synthesized by halophytic strains able to grow in saturated salt concentrations (Hagemann, 2011 and references therein). Moderately halotolerant (generally marine) strains accumulate glucosylglycerol as major compatible solute and sometimes glucosylglycerate as secondary solute. Freshwater strains with low halotolerance mainly synthesize sucrose and/or trehalose. Among the three species investigated, only *Dolichospermum* sp. isolated from brackish waters was found to accumulate sucrose in response to the two highest salinity levels tested. This agrees with the list of compatible solutes found in *Dolichospermum* (*Anabaena*) species (supplementary table in Hagemann, 2011). This is also in line with the sucrose accumulation observed in a particularly salt-tolerant *M. aeruginosa* strain, isolated from brackish waters (Georges des Aulnois et al., 2019). The protective role of sucrose during salt stress was further confirmed in the same *M. aeruginosa* strain by the over-expression of genes involved in its synthesis in response to salinity increase (Georges des Aulnois et al., 2020).

The presence of proline was found in all three cyanobacteria. In *Dolichospermum* sp., proline did not seem to be used as a compatible solute because NaCl exposure did not significantly modify its intracellular concentration. *A. gracile* and *C. raciborskii* showed different patterns in proline variations. In *A. gracile*, proline accumulated proportionally to NaCl concentration at two different times: 24h and 72h. In *C. raciborskii*, after a first phase during which proline content was inversely proportional to NaCl concentration, proline accumulated in a dose-dependent manner after 48h of NaCl exposure. However, with the present results, it is difficult to confirm with certainty the role of proline as a compatible solute in *A. gracile* and *C. raciborskii* because unlike most other compatible solutes, proline is used in primary metabolism (Hagemann, 2011). In addition, in plants, proline has been recognized as a multifunctional amino acid (Szabados and Saviouré, 2010). It has been proposed to act as a compatible solute, to be a ROS scavenger, to function as a molecular

chaperone stabilizing the structure of proteins and to act as a signaling molecule to modulate mitochondrial functions (Ben Rejeb et al., 2014; Szabados and Savouré, 2010; Verbruggen and Hermans, 2008). In *A. gracile* and *C. raciborskii*, ROS decreased when proline accumulated in cells respectively after 24 and 48h exposure to NaCl. However, it remains difficult to interpret this as a cause and effect relationships confirming that proline acted as a ROS scavenger. In *A. gracile* and *C. raciborskii*, proline trends were similar to variations in methionine content. This suggests that proline content may also have been influenced by protein synthesis alteration and anabolic processes. There are extensive reports on salt stress induced proline accumulation in higher plants (see Mansour and Ali, 2017 for a review). By contrast, in cyanobacteria, not much work has been done to understand the role of proline in salinity stress tolerance. The action of proline as a potential compatible solute has been shown in *Nostoc muscorum* (Singh et al., 1996). A mutant strain, over accumulating proline due to their deficiency in proline oxidase activity, has been found to be protected against the lethal effects of 150 mol m⁻³ NaCl. The culture medium supplementation with proline, and its subsequent uptake, offered a similar protection to the parent strain. Intracellular proline content has also been shown to naturally accumulate in response to NaCl exposure in [an halophilic *Synechococcus* sp. strain](#) (Fulda et al., 1999), [cyanobacteria isolated from rice fields](#) (*Cylindrospermum* sp. (Chris et al., 2006) and *Anabaena variabilis* (Syiem and Nongrum, 2011)) and two strains of *M. aeruginosa* ([one isolated from a freshwater lake and the other from a brackish water reservoir](#)) (Georges des Aulnois et al., 2019). In *Synechococcus* sp., proline accumulation preceded the synthesis of glycine betaine (Fulda et al., 1999).

5 Conclusions

The brackish (*Dolichospermum* sp.) and freshwater (*A. gracile*, *C. raciborskii*) species presented different tolerance to salinity stress. In estuaries, *A. gracile* and *C. raciborskii* would not be able to survive beyond the oligohaline area while *Dolichospermum* sp. seems to be able to colonize a larger zone and to potentially survive up to the polyhaline area. In all species, NaCl exposure resulted in reduced growth rates. By contrast, the responses of photosynthetic activity, pigment composition, ROS production as well as the capacity to accumulate compatible solutes were species-specific. *A. gracile* and *C. raciborskii* responded differently to salinity stress even though they were isolated from similar systems (freshwaters). This illustrates the diversity of mechanisms operated by cyanobacteria to face salinity variations. Low sodium chloride concentrations did not impact STXs production in *A. gracile* and *C. raciborskii* while, higher concentrations modified the toxin profile and content of *C. raciborskii* (intracellular STX decreased while dc-GTX2 increased). Since toxicity of PSTs analogs is different, these variations in toxin profile imply that salinity variations have the potential to modify the toxicity of this species. This is an important fact to consider when assessing the risk of animal and human exposure to cyanobacterial PSTs.

Acknowledgements

This study was funded by IFREMER and the Atlantic Microalgae project (AMI) supported by the Région Pays de la Loire.

References

Adams, M.A., Chen, Z., Landman, P., Colmer, T.D., 1999. Simultaneous Determination by Capillary Gas Chromatography of Organic Acids, Sugars, and Sugar Alcohols in Plant Tissue Extracts as Their Trimethylsilyl Derivatives. *Analytical Biochemistry* 266(1), 77-84.

- 924 Aguilera-Belmonte, A., Inostroza, I., Carrillo, K.S., Franco, J.M., Riobó, P., Gómez, P.I.,
 925 2013. The combined effect of salinity and temperature on the growth and toxin content of four
 926 Chilean strains of *Alexandrium catenella* (Whedon and Kofoid) Balech 1985 (Dinophyceae)
 927 isolated from an outbreak occurring in southern Chile in 2009. *Harmful Algae* 23, 55-59.
- 928 Allen, M.M., 1968. Simple conditions for growth of unicellular blue-green algae on plates.
 929 *Journal of Phycology* 4, 1-4.
- 930 Allen, M.M., Stanier, R.Y., 1968. Growth and division of some unicellular blue-green algae.
 931 *Journal of General Microbiology* 51, 199-202.
- 932 Antunes, J.T., Leão, P.N., Vasconcelos, V.M., 2015. *Cylindrospermopsis raciborskii*: review
 933 of the distribution, phylogeography, and ecophysiology of a global invasive species. *Frontiers*
 934 *in Microbiology* 6(473), article 473
- 935 Ballot, A., Fastner, J., Wiedner, C., 2010. Paralytic Shellfish Poisoning Toxin-Producing
 936 Cyanobacterium *Aphanizomenon gracile* in Northeast Germany. *Applied and Environmental*
 937 *Microbiology* 76, 1173-1180.
- 938 Ben Rejeb, K., Abdelly, C., Savouré, A., 2014. How reactive oxygen species and proline face
 939 stress together. *Plant Physiology and Biochemistry* 80, 278-284.
- 940 Bernard, C., Harvey, M., Briand, J.F., Biré, R., Krys, S., Fontaine, J.J., 2003. Toxicological
 941 comparison of diverse *Cylindrospermopsis raciborskii* strains: Evidence of liver damage
 942 caused by a French *C. raciborskii* strain. *Environmental toxicology* 18(3), 176-186.
- 943 Bhargava, P., Srivastava, A.K., 2013. Salt toxicity and survival strategies of cyanobacteria, In:
 944 Srivastava, A.K., Rai, A.N., Neilan, B.A. (Eds.), *Stress biology of cyanobacteria: molecular*
 945 *mechanisms to cellular responses*. CRC Press Boca Raton, FL, USA, pp. 171-188.
- 946 Bormans, M., Amzil, Z., Mineaud, E., Brient, L., Savar, V., Robert, E., Lance, E., 2019.
 947 Demonstrated transfer of cyanobacteria and cyanotoxins along a freshwater-marine
 948 continuum in France. *Harmful Algae* 87, 101639.
- 949 Brentano, D.M., Giehl, E.L.H., Petrucio, M.M., 2016. Abiotic variables affect STX
 950 concentration in a meso-oligotrophic subtropical coastal lake dominated by
 951 *Cylindrospermopsis raciborskii* (Cyanophyceae). *Harmful Algae* 56, 22-28.
- 952 Brown, M.B., Forsythe, A.B., 1974. Robust tests for equality of variances. *Journal of*
 953 *American Statistical Association* 69, 364-367.
- 954 Brutemark, A., Vandellannoote, A., Engström-Öst, J., Suikkanen, S., 2015. A Less Saline
 955 Baltic Sea Promotes Cyanobacterial Growth, Hampers Intracellular Microcystin Production,
 956 and Leads to Strain-Specific Differences in Allelopathy. *PLOS ONE* 10(6), e0128904.
- 957 Burford, M.A., Carey, C.C., Hamilton, D.P., Huisman, J., Paerl, H.W., Wood, S.A., Wulff,
 958 A., 2020. Perspective: Advancing the research agenda for improving understanding of
 959 cyanobacteria in a future of global change. *Harmful Algae* 91, 101601.
- 960 Carmichael, W.W., 2001. Health effects of toxin-producing cyanobacteria: "the cyanoHABs".
 961 *Human and ecological risk assessment: an international journal* 7, 1393-1407.
- 962 Carneiro, R., Pacheco, A., de Oliveira e Azevedo, S., 2013. Growth and Saxitoxin Production
 963 by *Cylindrospermopsis raciborskii* (Cyanobacteria) Correlate with Water Hardness. *Marine*
 964 *Drugs* 11(8), 2949.
- 965 Casero, M.C., Ballot, A., Agha, R., Quesada, A., Cirés, S., 2014. Characterization of saxitoxin
 966 production and release and phylogeny of sxt genes in paralytic shellfish poisoning toxin-
 967 producing *Aphanizomenon gracile*. *Harmful Algae* 37, 28-37.
- 968 Chen, L., Mao, F.J., Kirumba, G.C., Jiang, C., Manefield, M., He, Y.L., 2015. Changes in
 969 metabolites, antioxidant system, and gene expression in *Microcystis aeruginosa* under sodium
 970 chloride stress. *Ecotoxicology and Environmental Safety* 122, 126-135.
- 971 Chris, A., Zeeshan, M., Abraham, G., Prasad, S.M., 2006. Proline accumulation in
 972 *Cylindrospermum* sp. *Environmental and Experimental Botany* 57(1), 154-159.

- 973 Cirés, S., Ballot, A., 2016. A review of the phylogeny, ecology and toxin production of
 974 bloom-forming *Aphanizomenon spp.* and related species within the Nostocales
 975 (cyanobacteria). *Harmful Algae* 54, 21-43.
- 976 Curson, A.R.J., Williams, B.T., Pinchbeck, B.J., Sims, L.P., Martínez, A.B., Rivera, P.P.L.,
 977 Kumaresan, D., Mercadé, E., Spurgin, L.G., Carrión, O., Moxon, S., Cattolico, R.A.,
 978 Kuzhiumparambil, U., Guagliardo, P., Clode, P.L., Raina, J.-B., Todd, J.D., 2018. DSYB
 979 catalyses the key step of dimethylsulfoniopropionate biosynthesis in many phytoplankton.
 980 *Nature Microbiology* 3(4), 430-439.
- 981 D'Agostino, P.M., Moffitt, M., Neilan, B., 2014. Current knowledge of paralytic shellfish
 982 toxin biosynthesis, molecular detection and evolution, Active compounds from microalgae,
 983 pp. 251-280.
- 984 D'Agostino, P.M., Song, X.M., Neilan, B.A., Moffitt, M.C., 2016. Proteogenomics of a
 985 saxitoxin-producing and non-toxic strain of *Anabaena circinalis* (cyanobacteria) in response
 986 to extracellular NaCl and phosphate depletion. *Environmental Microbiology* 18(2), 461-476.
- 987 Dittmann, E., Fewer, D.P., Neilan, B.A., 2012. Cyanobacterial toxins: biosynthetic routes and
 988 evolutionary roots. *FEMS Microbiology Reviews* 37, 23-43.
- 989 Duysens, L.N., 1972. 3-(3,4-Dichlorophenyl)-1,1-dimethylurea (DCMU) inhibition of system
 990 II and light-induced regulatory changes in energy transfer efficiency. *Biophysical journal* 12,
 991 858-863.
- 992 Eilers, P.H.C., Peeters, J.C.H., 1988. A model for the relationship between light intensity and
 993 the rate of photosynthesis in phytoplankton. *Ecological modelling* 42, 199-215.
- 994 Ferla, M.P., Patrick, W.M., 2014. Bacterial methionine biosynthesis. *Microbiology Society*,
 995 pp. 1571-1584.
- 996 Fulda, S., Huckauf, J., Schoor, A., Hagemann, M., 1999. Analysis of Stress Responses in the
 997 Cyanobacterial Strains *Synechococcus sp.* PCC 7942, *Synechocystis sp.* PCC 6803, and
 998 *Synechococcus sp.* PCC 7418: Osmolyte Accumulation and Stress Protein Synthesis. *Journal*
 999 *of Plant Physiology* 154(2), 240-249.
- 1000 Genty, B., Briantais, J.M., Baker, N.R., 1989. The relationship between the quantum yield of
 1001 photosynthetic electron transport and quenching of fluorescence *Biochemical and Biophysical*
 1002 *Acta* 990, 87-92.
- 1003 Georges des Aulnois, M., Réveillon, D., Robert, E., Caruana, A., Briand, E., Guljamow, A.,
 1004 Dittmann, E., Amzil, Z., Bormans, M., 2020. Salt Shock Responses of Microcystis Revealed
 1005 through Physiological, Transcript, and Metabolomic Analyses. *Toxins* 12(3), 192.
- 1006 Georges des Aulnois, M., Roux, P., Caruana, A., Réveillon, D., Briand, E., Hervé, F., Savar,
 1007 V., Bormans, M., Amzil, Z., 2019. Physiological and Metabolic Responses of Freshwater and
 1008 Brackish-Water Strains of *Microcystis aeruginosa* Acclimated to a Salinity Gradient: Insight
 1009 into Salt Tolerance. *Applied and Environmental Microbiology* 85(21), e01614-01619.
- 1010 Gong, H., Tang, Y., Wang, J., Wen, X., Zhang, L., Lu, C., 2008. Characterization of
 1011 photosystem II in salt-stressed cyanobacterial *Spirulina platensis* cells. *Biochimica et*
 1012 *Biophysica Acta (BBA) - Bioenergetics* 1777(6), 488-495.
- 1013 Gröne, T., Kirst, G.O., 1992. The effect of nitrogen deficiency, methionine and inhibitors of
 1014 methionine metabolism on the DMSP contents of *Tetraselmis subcordiformis* (Stein). *Marine*
 1015 *Biology* 112(3), 497-503.
- 1016 Grossman, A.R., Schaefer, M.R., Chiang, G.G., Collier, J.L., 1993. The phycobilisome, a
 1017 light-harvesting complex responsive to environmental conditions, *Microbiological reviews*,
 1018 pp. 725-749.
- 1019 Hagemann, M., 2011. Molecular biology of cyanobacterial salt acclimation. *FEMS*
 1020 *Microbiology Reviews* 35(1), 87-123.
- 1021 Holm, S., 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal*
 1022 *of Statistics* 6, 65-70.

- 1023 Jančula, D., Straková, L., Sadilek, J., Maršálek, B., Babica, P., 2014. Survey of cyanobacterial
1024 toxins in Czech water reservoirs - the first observation of neurotoxic saxitoxins.
1025 Environmental Science and Pollution Research 21, 8006-2015.
- 1026 Jeanjean, R., Matthijs, H.C.P., Onana, B., Havaux, M., Joset, F., 1993. Exposure of the
1027 Cyanobacterium *Synechocystis* PCC6803 to Salt Stress Induces Concerted Changes in
1028 Respiration and Photosynthesis. Plant and Cell Physiology 34(7), 1073-1079.
- 1029 John, J., Kemp, A., 2006. Cyanobacterial Blooms in the Wetlands of the Perth
1030 region, Taxonomy and Distribution: an Overview. Journal of the Royal Society of Western
1031 Australia 89, 51-26.
- 1032 Johnsen, G., Sakshaug, E., 2007. Biooptical characteristics of PSII and PSI in 33 species (13
1033 pigment groups) of marine phytoplankton, and the relevance for pulse-amplitude-modulated
1034 and fast-repetition-rate fluorometry. Journal of Phycology 43, 1236-1251.
- 1035 Joshua, S., Mullineaux, C.W., 2004. Phycobilisome Diffusion Is Required for Light-State
1036 Transitions in Cyanobacteria. Bioenergetics and photosynthesis 135(4), 2112-2119.
- 1037 Kellmann, R., Mihali, T.K., Jeon, Y.J., Pickford, R., Pomati, F., Neilan, B.A., 2008.
1038 Biosynthetic intermediate analysis and functional homology reveal a saxitoxin gene cluster in
1039 cyanobacteria. Applied and environmental microbiology 74(13), 4044-4053.
- 1040 Klahn, S., Hagemann, M., 2011. Compatible solute biosynthesis in cyanobacteria.
1041 Environmental Microbiology 13(3), 551-562.
- 1042 Kleinteich, J., Wood, S.A., Puddick, J., Schleheck, D., Küpper, F.C., Dietrich, D., 2013.
1043 Potent toxins in Arctic environments - presence of saxitoxins and an unusual microcystin
1044 variant in Arctic freshwater ecosystems. Chemico-biological interactions 206, 423-431.
- 1045 Kolbowski, J., Schreiber, U., 1995. Computer-controlled phytoplankton analyser based on 4-
1046 wavelengths PAM chlorophyll fluorometer. In: Mathis P. (ed) Photosynthesis: from Light to
1047 Biosphere (V). Kluwer Academic Publishers, Dordrecht. pp 825-828.
- 1048 Kumar, J., Singh, V.P., Prasad, S.M., 2015. NaCl-induced physiological and biochemical
1049 changes in two cyanobacteria *Nostoc muscorum* and *Phormidium foveolarum* acclimatized to
1050 different photosynthetically active radiation. Journal of Photochemistry and Photobiology B-
1051 Biology 151, 221-232.
- 1052 Lajeunesse, A., Segura, P.A., Gélinas, M., Hudon, C., Thomas, K., Quilliam, M.A., Gagnon,
1053 C., 2012. Detection and confirmation of saxitoxin analogues in freshwater benthic *Lyngbya*
1054 *wollei* algae collected in the St. Lawrence River (Canada) by liquid chromatography–tandem
1055 mass spectrometry. Journal of chromatography A 1219, 93-103.
- 1056 Latifi, A., Ruiz, M., Zhang, C.-C., 2009. Oxidative stress in cyanobacteria. FEMS
1057 Microbiology Reviews 33(2), 258-278.
- 1058 Ledreux, A., Thomazeau, S., Catherine, A., Duval, C., Yepremian, C., Marie, A., Bernard, C.,
1059 2010. Evidence for saxitoxins production by the cyanobacterium *Aphanizomenon gracile* in a
1060 French recreational water body. Harmful Algae 10(1), 88-97.
- 1061 Li, X.C., Dreher, T.W., Li, R.H., 2016. An overview of diversity, occurrence, genetics and
1062 toxin production of bloom-forming *Dolichospermum* (*Anabaena*) species. Harmful Algae 54,
1063 54-68.
- 1064 Llewellyn, L.E., 2006. Saxitoxin, a toxic marine natural product that targets a multitude of
1065 receptors. Natural Product Reports 23(2), 200-222.
- 1066 Lu, C., Torzillo, G., Vonshak, A., 1999. Kinetic response of photosystem II photochemistry in
1067 the cyanobacterium *Spirulina platensis* to high salinity is characterized by two distinct phases.
1068 Functional Plant Biology 26(3), 283-292.
- 1069 Lu, C., Vonshak, A., 2002. Effects of salinity stress on photosystem II function in
1070 cyanobacterial *Spirulina platensis* cells. Physiologia Plantarum 114(3), 405-413.
- 1071 Mansour, M.M.F., Ali, E.F., 2017. Evaluation of proline functions in saline conditions.
1072 Phytochemistry 140, 52-68.

- 1073 Melero-Jiménez, I.J., Martín-Clemente, E., García-Sánchez, M.J., Bañares-España, E., Flores-
1074 Moya, A., 2020. The limit of resistance to salinity in the freshwater cyanobacterium
1075 *Microcystis aeruginosa* is modulated by the rate of salinity increase. *Ecology and Evolution*
1076 00, 1-11.
- 1077 Mitchell, B.G., Kahru, M., Wieland, J., Stramska, M., 2003. Determination of spectral
1078 absorption coefficient of particles, dissolved material and phytoplankton for discrete water
1079 samples. In: Mueller J.L., Fargion G.S., McClain CR (eds) *Ocean optics protocols for*
1080 *satellite ocean color sensor validation, Revision 4, vol IV. NASA Technical Memorandum*
1081 2003-211621. NASA GFSC, Greenbelt, Maryland, pp 39-64.
- 1082 Moisander, P.H., McClinton, E., Paerl, H.W., 2002. Salinity effects on growth, photosynthetic
1083 parameters, and nitrogenase activity in estuarine planktonic cyanobacteria. *Microbial Ecology*
1084 43(4), 432-442.
- 1085 Munday, R., 2014. Toxicology of seafood toxins: A critical review, In: Botana, L. (Ed.),
1086 *Seafood and freshwater toxins: pharmacology, physiology, and detection. CRC press: Boca*
1087 *Raton, FL, USA, pp. 197-290.*
- 1088 Nelder, J.A., Mead, R., 1965. A simplex method for function minimization. *The Computer*
1089 *Journal* 7, 308-313.
- 1090 O'Neil, J.M., Davis, T.W., Burford, M.A., Gobler, C.J., 2012. The rise of harmful
1091 cyanobacteria blooms: the potential roles of eutrophication and climate change. *Harmful*
1092 *Algae* 14, 313-334.
- 1093 Ogawa, T., Misumi, M., Sonoike, K., 2017. Estimation of photosynthesis in cyanobacteria by
1094 pulse-amplitude modulation chlorophyll fluorescence: problems and solutions. *Photosynthesis*
1095 *Research* 133(1-3), 63-73.
- 1096 Ongley, S.E., Pengelly, J.J.L., Neilan, B.A., 2016. Elevated Na⁺ and pH influence the
1097 production and transport of saxitoxin in the cyanobacteria *Anabaena circinalis* AWQC131C
1098 and *Cylindrospermopsis raciborskii* T3. *Environmental Microbiology* 18(2), 427-438.
- 1099 Oren, A., 2007. Diversity of organic osmotic compounds and osmotic adaptation in
1100 cyanobacteria and algae, *Algae and Cyanobacteria in Extreme Environments. Springer, pp.*
1101 *639-655.*
- 1102 Paerl, H.W., Fulton, R.S., Moisander, P.H., Dyble, J., 2001. Harmful freshwater algal blooms,
1103 with an emphasis on cyanobacteria. *The scientific world* 1, 76-113.
- 1104 Paerl, H.W., Huisman, J., 2009. Climate change: a catalyst for global expansion of harmful
1105 cyanobacterial blooms. *Environmental Microbiology Reports* 1, 27-37.
- 1106 Paerl, H.W., Paul, V.J., 2012. Climate change: links to global expansion of harmful
1107 cyanobacteria. *Water research* 46, 1349-1363.
- 1108 Paredes, I., Rietjens, I.M.C.M., Vieites, J.M., Cabado, A.G., 2011. Update of risk assessments
1109 of main marine biotoxins in the European Union. *Toxicon* 58(4), 336-354.
- 1110 Pearson, L., Mihali, T., Moffitt, M., Kellmann, R., Neilan, B., 2010. On the chemistry,
1111 toxicology and genetics of the cyanobacterial toxins, microcystins, nodularin, saxitoxin and
1112 cylindrospermopsin. *Marine Drugs* 8, 1650-1680.
- 1113 Pearson, L.A., Dittmann, E., Mazmouz, R., Ongley, S.E., D'Agostino, P.M., Neilan, B.A.,
1114 2016. The genetics, biosynthesis and regulation of toxic specialized metabolites of
1115 cyanobacteria. *Harmful Algae* 54, 98-111.
- 1116 Pomati, F., Moffitt, M.C., Cavaliere, R., Neilan, B.A., 2004a. Evidence for differences in the
1117 metabolism of saxitoxin and C1+2 toxins in the freshwater cyanobacterium
1118 *Cylindrospermopsis raciborskii* T3. *Biochimica Et Biophysica Acta-General Subjects*
1119 1674(1), 60-67.
- 1120 Pomati, F., Rossetti, C., Manarolla, G., Burns, B.P., Neilan, B.A., 2004b. Interactions
1121 between intracellular Na⁺ levels and saxitoxin production in *Cylindrospermopsis raciborskii*
1122 T3. *Microbiology-Sgm* 150, 455-461.

- 1123 Preece, E.P., Hardy, F.J., Moore, B.C., Bryan, M., 2017. A review of microcystin detections
1124 in Estuarine and Marine waters: Environmental implications and human health risk. *Harmful*
1125 *Algae* 61, 31-45.
- 1126 Preußel, K., Wessel, G., Fastner, J., Chorus, I., 2009. Response of cylindrospermopsin
1127 production and release in *Aphanizomenon flos-aquae* (Cyanobacteria) to varying light and
1128 temperature conditions. *Harmful Algae* 8(5), 645-650.
- 1129 Rajneesh, Pathak, j., Chatterjee, A., Singh, S.P., Sinha, R.P., 2017. Detection of reactive
1130 oxygen species (ROS) in cyanobacteria using the oxidant-sensing probe 2'-7'-
1131 Dichlorodihydrofluorescein diacetate (DCFH-DA). *Bio-protocol* 7, BioProtoc.2545.
- 1132 Rapala, J., Robertson, A., Negri, A.P., Berg, K.A., Tuomi, P., Lyra, C., Erkomaa, K., Lahti,
1133 K., Hoppu, K., Lepistö, L., 2005. First report of saxitoxin in Finnish lakes and possible
1134 associated effects on human health. *Environmental toxicology* 20, 331-340.
- 1135 Rippka, R., Deruelles, J., Waterbury, J.B., Herdman, M., Stanier, R.Y., 1979. Generic
1136 assignments, strain histories and properties of pure cultures of cyanobacteria. *Journal of*
1137 *General Microbiology* 111, 1-61.
- 1138 Robson, B.J., Hamilton, D.P., 2013. Summer flow event induces a cyanobacterial bloom in a
1139 seasonal Western Australian estuary. *Marine and Freshwater Research* 54, 139-151.
- 1140 Ross, C., Santiago-Vázquez, L., Paul, V., 2006. Toxin release in response to oxidative stress
1141 and programmed cell death in the cyanobacterium *Microcystis aeruginosa*. *Aquatic*
1142 *Toxicology* 78(1), 66-73.
- 1143 Ross, C., Warhurst, B.C., Brown, A., Huff, C., Ochrietor, J.D., 2019. Mesohaline conditions
1144 represent the threshold for oxidative stress, cell death and toxin release in the cyanobacterium
1145 *Microcystis aeruginosa*. *Aquatic Toxicology* 206, 203-211.
- 1146 Royston, J.P., 1982. An extension of Shapiro and Wilk's W test for normality to large
1147 samples. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 31, 115-124.
- 1148 Schantz, R., Ghazarossian, V., Schnoes, H.K., Strong, F., Springer, J., Pezzanite, J.O., Clardy,
1149 J., 1975. Structure of saxitoxin. *Journal of American Chemical Society* 97, 1238-1239.
- 1150 Schreiber, U., 1998. Chlorophyll fluorescence: New instruments for special applications. In:
1151 Garab G. (ed) *Photosynthesis: mechanisms and effects*, vol. 5. Kluwer Academic Publishers,
1152 Dordrecht. pp 4253-4258.
- 1153 Schubert, H., Fulda, S., Hagemann, M., 1993. Effects of Adaptation to Different Salt
1154 Concentrations on Photosynthesis and Pigmentation of the Cyanobacterium *Synechocystis sp.*
1155 PCC 6803. *Journal of Plant Physiology* 142(3), 291-295.
- 1156 Schubert, H., Hagemann, M., 1990. Salt effects on 77K fluorescence and photosynthesis in
1157 the cyanobacterium *Synechocystis sp.* PCC 6803. *FEMS Microbiology Letters* 71(1-2), 169-
1158 172.
- 1159 Seppälä, J., 2009. Fluorescence properties of Baltic sea phytoplankton. *Monographs of the*
1160 *Boreal environment research* (34). Edita Prima Ltd, Helsinki, pp 83.
- 1161 Seródio, J., Cruz, S., Vieira, S., Brotas, V., 2005. Non-photochemical quenching of
1162 fluorescence and operation of the xanthophyll cycle in estuarine microphytobenthos. *Journal*
1163 *of experimental marine biology and ecology* 326, 157-169.
- 1164 Singh, A.K., Chakravarthy, D., Singh, T.P.K., Singh, H.N., 1996. Evidence for a role for L-
1165 proline as a salinity protectant in the cyanobacterium *Nostoc muscorum*. *Plant Cell and*
1166 *Environment* 19(4), 490-494.
- 1167 Slim, K., Fadel, A., Atoui, A., Lemaire, B.J., Vinçon-Leite, B., Tassin, B., 2014. Global
1168 warming as a driving factor for cyanobacterial blooms in Lake Karaoun, Lebanon.
1169 *Desalination and Water Treatment* 52, 2094-2101.
- 1170 Smith, F.M.J., Wood, S.A., van Ginkel, R., Broady, P.A., Gaw, S., 2011. First report of
1171 saxitoxin production by a species of the freshwater benthic cyanobacterium, *Scytonema*
1172 *Agardh*. *Toxicon* 57, 566-573.

- 1173 Soto-Liebe, K., Lopez-Cortes, X.A., Fuentes-Valdes, J.J., Stucken, K., Gonzalez-Nilo, F.,
 1174 Vasquez, M., 2013. In Silico Analysis of Putative Paralytic Shellfish Poisoning Toxins Export
 1175 Proteins in Cyanobacteria. *Plos One* 8(2).
- 1176 Soto-Liebe, K., Mendez, M.A., Fuenzalida, L., Krock, B., Cembella, A., Vasquez, M., 2012.
 1177 PSP toxin release from the cyanobacterium *Raphidiopsis brookii* D9 (Nostocales) can be
 1178 induced by sodium and potassium ions. *Toxicon* 60(7), 1324-1334.
- 1179 Soto-Liebe, K., Murillo, A.A., Krock, B., Stucken, K., Fuentes-Valdes, J.J., Trefault, N.,
 1180 Cembella, A., Vasquez, M., 2010. Reassessment of the toxin profile of *Cylindrospermopsis*
 1181 *raciborskii* T3 and function of putative sulfotransferases in synthesis of sulfated and
 1182 sulfonated PSP toxins. *Toxicon* 56(8), 1350-1361.
- 1183 Stefels, J., 2000. Physiological aspects of the production and conversion of DMSP in marine
 1184 algae and higher plants. *Journal of Sea Research* 43(3), 183-197.
- 1185 Steinke, M., Hodapp, B., Subhan, R., Bell, T.G., Martin-Creuzburg, D., 2018. Flux of the
 1186 biogenic volatiles isoprene and dimethyl sulfide from an oligotrophic lake. *Scientific Reports*
 1187 8(1), 630.
- 1188 Strickland, I.D.H., Parsons, T.R., 1972. A practical handbook of seawater analysis. 2nd Ed.
 1189 Fisheries Research Board of Canada, Ottawa. pp: 310.
- 1190 Sudhir, P., Murthy, S.D.S., 2004. Effects of salt stress on basic processes of photosynthesis.
 1191 *Photosynthetica* 42(4), 481-486.
- 1192 Swapnil, P., Rai, A.K., 2018. Physiological responses to salt stress of salt-adapted and
 1193 directly salt (NaCl and NaCl+Na₂SO₄ mixture)-stressed cyanobacterium *Anabaena*
 1194 *fertilissima*. *Protoplasma* 255(3), 963-976.
- 1195 Swapnil, P., Yadav, A.K., Srivastav, S., Sharma, N.K., Srikrishna, S., Rai, A.K., 2017.
 1196 Biphasic ROS accumulation and programmed cell death in a cyanobacterium exposed to
 1197 salinity (NaCl and Na₂SO₄). *Algal Research-Biomass Biofuels and Bioproducts* 23, 88-95.
- 1198 Syiem, M.B., Nongrum, N.A., 2011. Increase in intracellular proline content in *Anabaena*
 1199 *variabilis* during stress conditions. *Journal of Applied and Natural Science* 3(1), 119-123.
- 1200 Szabados, L., Savouré, A., 2010. Proline: a multifunctional amino acid. *Trends in Plant*
 1201 *Science* 15(2), 89-97.
- 1202 Teneva, I., Mladenov, R., Belkinova, D., Dimitrova-Dyulgerova, I., Dzhabazov, B., 2010.
 1203 Phytoplankton community of the drinking water supply reservoir Borovitsa (South Bulgaria)
 1204 with an emphasis on cyanotoxins and water quality. *Central European Journal of Biology* 5,
 1205 231-239.
- 1206 Tonk, L., Bosch, K., Visser, P., Huisman, J., 2007. Salt tolerance of the harmful
 1207 cyanobacterium *Microcystis aeruginosa*. *Aquatic Microbial Ecology* 46(2), 117-123.
- 1208 UNESCO, 1966. Determination of photosynthetic pigments in seawater. Report of SCOR-
 1209 UNESCO Working group 17. Monographs on Oceanographic Methodology 1, 1-69.
- 1210 Van de Riet, J.M., Gibbs, R.S., Chou, F.W., Muggah, P.M., Rourke, W.A., Burns, G., 2009.
 1211 Liquid chromatographic post-column oxidation method for analysis of paralytic shellfish
 1212 toxins in mussels, clams, scallops, and oysters: single-laboratory validation. *Journal of AOAC*
 1213 *International* 92, 1690-1704.
- 1214 Verbruggen, N., Hermans, C., 2008. Proline accumulation in plants: a review. *Amino Acids*
 1215 35(4), 753-759.
- 1216 Wiese, M., D'Agostino, P.M., Mihali, T.K., Moffitt, M.C., Neilan, B.A., 2010. Neurotoxic
 1217 Alkaloids: Saxitoxin and Its Analogs. *Marine Drugs* 8(7), 2185-2211.
- 1218 Williams, W.D., 2001. Anthropogenic salinisation of inland waters. *Hydrobiologia* 466, 329-
 1219 337.
- 1220 Wood, A.M., Everroad, R.C., Wingard, L.M., 2005. Measuring growth rates in microalgal
 1221 cultures, In: Andersen, R.A. (Ed.), *Algal Culturing techniques*. Elsevier Academic Press
 1222 Burlington, MA, pp. 269-286.

- 1223 Wörmer, L., Cirés, S., Agha, R., Verdugo, M., de Hoyos, C., Quesada, A., 2011. First
1224 detection of cyanobacterial PSP (paralytic shellfish poisoning) toxins in Spanish freshwaters.
1225 Toxicon 57, 918-921.
- 1226 Zhang, T., Gong, H., Wen, X., Lu, C., 2010. Salt stress induces a decrease in excitation
1227 energy transfer from phycobilisomes to photosystem II but an increase to photosystem I in the
1228 cyanobacterium *Spirulina platensis*. Journal of Plant Physiology 167(12), 951-958.
1229
1230

Figures captions

Fig. 1: Growth curves of A) *Aphanizomenon gracile* and B) *Cylindrospermopsis raciborskii* in BG11, BG11 + 34 mM NaCl, BG11 + 154 mM NaCl and BG11 + 257 mM NaCl. C) Growth curves of *Dolichospermum* sp. in BG11 + 154 mM NaCl, BG11 + 257 mM NaCl, BG11 + 308 mM NaCl, and BG11 + 411 mM NaCl. Data are average of triplicate cultures and error bars represent standard deviation

Fig. 2: Carotenoids content (in μg ($\mu\text{g chl a}^{-1}$)) (A, B, C), phycoerythrine content (in relative units) (D, E, F) and phycocyanin content (in relative units) (G, H, I) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D, G, J), *Cylindrospermopsis raciborskii* (B, E, H, K) and *Dolichospermum* sp. (C, F, I, L). Data are average of triplicate cultures and error bars represent standard deviation

Fig. 3: Maximum quantum yield (F_v/F_m) (A, B, C), maximal light utilization efficiency (α in $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$) $^{-1}$) (D, E, F), maximum electron transport rate (ETR_m in $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$) (G, H, I) and maximum non-photochemical quenching (NPQ_{max}) (J, K, L) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D, G), *Cylindrospermopsis raciborskii* (B, E, H) and *Dolichospermum* sp. (C, F, I). Data are average of triplicate cultures and error bars represent standard deviation

Fig. 4: ROS production measured as 2',7'-dichlorofluorescein (DCF) fluorescence level (in relative units) over time and along a salinity gradient in *Aphanizomenon gracile* (A), *Cylindrospermopsis raciborskii* (B) and *Dolichospermum* sp. (C). Results are represented as relative changes from the respective control conditions. Data are average of triplicate cultures and error bars represent standard deviation

Fig. 5: Intracellular quota of paralytic shellfish toxins (PSTs) (A, B), neo-saxitoxin (C, D), saxitoxin (E, F) and decarbamoylogonyautoxin 2 (G) over time and along a salinity gradient in *Aphanizomenon gracile* (A, C, E) and *Cylindrospermopsis raciborskii* (B, D, F, G). Data are average of triplicate cultures and error bars represent standard deviation. Please note the different y-axis limits

Fig. 6: Intracellular proportions of paralytic shellfish toxins analogs in *Aphanizomenon gracile* (A, B, C, D) and *Cylindrospermopsis raciborskii* (E, F, G, H) exposed to BG11 (A, E), BG11 + 34 mM NaCl (B, F), BG11 + 154 mM NaCl (C, G) and BG11 + 257 mM NaCl (D, H)

Fig. 7: Intracellular quota of methionine (A, B, C) and proline (D, E, F) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D) and *Cylindrospermopsis raciborskii* (B, E) and *Dolichospermum* sp. (C, F). Data are average of triplicate cultures and error bars represent standard deviation

Fig. 8: Intracellular sucrose accumulation in *Dolichospermum* sp. after a salt shock




Fig. 1: Growth curves of A) *Aphanizomenon gracile* and B) *Cylandropermopsis raciborskii* in BG11, BG11 + 34 mM NaCl, BG11 + 154 mM NaCl and BG11 + 257 mM NaCl. C) Growth curves of *Dolichospermum* sp. in BG11 + 154 mM NaCl, BG11 + 257 mM NaCl, BG11 + 308 mM NaCl, and BG11 + 411 mM NaCl. Data are average of triplicate cultures and error bars represent standard deviation




Fig. 2: Carotenoids content (in $\mu\text{g } (\mu\text{g chl a})^{-1}$) (A, B, C), phycoerythrin content (in relative units) (D, E, F) and phycocyanin content (in relative units) (G, H, I) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D, G), *Cylindrospermopsis raciborskii* (B, E, H) and *Dolichospermum* sp. (C, F, I). Data are average of triplicate cultures and error bars represent standard deviation




Fig. 3: Maximum quantum yield (F_v/F_m) (A, B, C), maximal light utilization efficiency (α in $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$ ($\mu\text{mol photons m}^{-2} \text{s}^{-1}$) $^{-1}$) (D, E, F), maximum electron transport rate (ETR_m in $\mu\text{mol e}^- \text{mg chl a}^{-1} \text{s}^{-1}$) (G, H, I) and maximum non-photochemical quenching (NPQ_{max}) (J, K, L) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D, G, J), *Cylindrospermopsis raciborskii* (B, E, H, K) and *Dolichospermum sp.* (C, F, I, L). Data are average of triplicate cultures and error bars represent standard deviation




Fig. 4: ROS production measured as 2',7'-dichlorofluorescein (DCF) fluorescence level (in relative units) over time and along a salinity gradient in *Aphanizomenon gracile* (A), *Cylindrospermopsis raciborskii* (B) and *Dolichospermum sp.* (C). Results are represented as relative changes from the respective control conditions. Data are average of triplicate cultures and error bars represent standard deviation




Fig. 5: Intracellular quota of paralytic shellfish toxins (PSTs) (A, B), neo-saxitoxin (C, D), saxitoxin (E, F) and decarbamoylogonyautoxin 2 (G) over time and along a salinity gradient in *Aphanizomenon gracile* (A, C, E) and *Cylindrospermopsis raciborskii* (B, D, F, G). Data are average of triplicate cultures and error bars represent standard deviation. Please note the different y-axis limits




Fig. 6: Intracellular proportions of paralytic shellfish toxins analogs in *Aphanizomenon gracile* (A, B, C, D) and *Cylindrospermopsis raciborskii* (E, F, G, H) exposed to BG11 (A, E), BG11 + 34 mM NaCl (B, F), BG11 + 154 mM NaCl (C, G) and BG11 + 257 mM NaCl (D, H)




Fig. 7: Intracellular quota of methionine (A, B, C) and proline (D, E, F) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D) and *Cylindrospermopsis raciborskii* (B, E) and *Dolichospermum* sp. (C, F). Data are average of triplicate cultures and error bars represent standard deviation




Fig. 8: Intracellular sucrose accumulation in *Dolichospermum* sp. after a salt shock

Supplementary materials

Physiological changes induced by sodium chloride stress in *Aphanizomenon gracile*, *Cylindrospermopsis raciborskii* and *Dolichospermum* sp.

Emilie Houliez, Enora Briand, Florent Malo, Georges-Augustin Rovillon, Fabienne Hervé, Elise Robert, Laetitia Marchand, Agata Zykwinska, Amandine Caruana




Fig. S1: High performance liquid chromatogram showing the retention time of paralytic shellfish toxins (PSTs) standards: gonyautoxin (GTX4), gonyautoxin (GTX1), decarbamoylogonyautoxin (dc-GTX3), gonyautoxin 5 (GTX5), decarbamoylogonyautoxin 2 (dc-GTX2), gonyautoxin 3 (GTX3), gonyautoxin 2 (GTX2), neosaxitoxin (NEO STX), decarbamoylsaxitoxin (dc-STX) and saxitoxin (STX)




Fig. S2: High performance liquid chromatogram of a blank




Fig. S3: A) Neosaxitoxin calibration curve. B) Saxitoxin calibration curve. C) Decarbamoylogonyautoxin 2 (dc-GTX2) calibration curve




Fig. S4: High performance liquid chromatogram of a sample of *Aphanizomenon gracile* (PMC627.10) culture




Fig. S5: High performance liquid chromatogram of a sample of *Cylindrospermopsis raciborskii* (PMC00.01) culture




Fig. S6: Intracellular quota of paralytic shellfish toxins (PSTs) (A, B), neo-saxitoxin (C, D), saxitoxin (E, F) and decarbamoylogonyautoxin 2 (G) over time and along a salinity gradient in *Aphanizomenon gracile* (A, C, E) and *Cylindrospermopsis raciborskii* (B, D, F, G). Data are average of triplicate cultures and error bars represent standard deviation. All toxins values were normalized by OD₇₅₀. Please note the different y-axis limits




Fig. S7: Intracellular proportions of paralytic shellfish toxins analogs in *Aphanizomenon gracile* (A, B, C, D) and *Cylindrospermopsis raciborskii* (E, F, G, H) exposed to BG11 (A, E), BG11 + 34 mM NaCl (B, F), BG11 + 154 mM NaCl (C, G) and BG11 + 257 mM NaCl (D, H). All toxins concentrations were normalized by OD₇₅₀ before the calculation of proportions




Fig. S8: High performance liquid chromatogram of a sample of *Dolichospermum* sp. (CCY9401) culture




Fig. S9: Intracellular quota of methionine (A, B, C) and proline (D, E, F) over time and along a salinity gradient in *Aphanizomenon gracile* (A, D) and *Cylindrospermopsis raciborskii* (B, E) and *Dolichospermum sp.* (C, F). Data are average of triplicate cultures and error bars represent standard deviation. All concentrations were normalized by OD₇₅₀







Fig. S10: Intracellular sucrose accumulation in *Dolichospermum* sp. after a salt shock when concentrations were normalized by OD_{750}


Figure1

[Click here to access/download;Figure;figure1.TIF](#)















Figure6







We declare that the work presented in this manuscript has not been published previously and is not under consideration for publication elsewhere. All authors read the manuscript, approved its content and agreed with its publication. If accepted, this work will not be published in the same form, in English or in any other language, without the written consent of the copyright-holder.

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: