

# State-Specific Configuration Interaction for Excited States

Fábris Kossoski, Pierre-François Loos

## ▶ To cite this version:

Fábris Kossoski, Pierre-François Loos. State-Specific Configuration Interaction for Excited States. Journal of Chemical Theory and Computation, 2023, 19 (8), pp.2258-2269. 10.1021/acs.jctc.3c00057 . hal-03843259

## HAL Id: hal-03843259 https://hal.science/hal-03843259

Submitted on 9 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

### State-Specific Configuration Interaction for Excited States

Fábris Kossoski<sup>1, a)</sup> and Pierre-Francois Loos<sup>1, b)</sup>

Laboratoire de Chimie et Physique Quantiques (UMR 5626), Université de Toulouse, CNRS, UPS, France

Abstract: We introduce and benchmark a new systematically improvable route for excited-state calculations, statespecific configuration interaction ( $\Delta$ CI). Starting with a reference built from optimized configuration state functions, separate CI calculations are performed for each targeted state (hence state-specific orbitals and determinants). Accounting for single and double excitations produces the  $\Delta$ CISD model, which can be improved with second-order Epstein-Nesbet perturbation theory ( $\Delta$ CISD+EN2) or a posteriori Davidson corrections ( $\Delta$ CISD+Q). These models were gauged against a vast and diverse set of 294 reference excitation energies. We have found that  $\Delta$ CI is significantly more accurate than standard ground-state-based CI, whereas close performances were found between  $\Delta$ CISD and EOM-CC2, and between  $\Delta$ CISD+EN2 and EOM-CCSD. For larger systems,  $\Delta$ CISD+Q delivers more accurate results than EOM-CC2 and EOM-CCSD. The  $\Delta$ CI route can handle challenging multireference problems, singly- and doubly-excited states, from closed- and open-shell species, with overall comparable accuracy, and thus represents a promising alternative to more established methodologies.



Most molecular electronic structure methods rely on different descriptions for ground and excited states. The ground state is described first, at a given level of theory, providing a baseline for later accessing the excited states, which in turn makes use of another approach or a distinct formalism altogether. For example, Kohn-Sham (KS) density-functional theory (DFT) is a ground-state method,<sup>1-3</sup> whereas the excited states are obtained later with a linear response treatment of time-dependent density-functional theory (TDDFT).4-7 Similarly, the coupled-cluster  $(CC)^{8-11}$  equations are solved for the ground state, whereas a diagonalization of the similaritytransformed Hamiltonian is implied in excited-state calculations based on the equation-of-motion (EOM)<sup>12,13</sup> or linearresponse<sup>14,15</sup> formalisms. Within configuration interaction (CI) methods.<sup>16</sup> the underlying formalism is the same for ground and excited states, but typical implementations also rely on a special treatment for the ground state, given the use of groundstate Hartree-Fock (HF) orbitals and the fact that the truncated CI space is spanned by excitations from the ground-state HF determinant. Apart from truly multireference approaches,<sup>17,18</sup> excited state methods entail a formal distinction between the targeted excited states and the ground state.

It is thus important to devise methods that minimize this unbalance as much as possible, aiming at a more unified description of ground and excited states, while keeping a modest computational cost. This also means a more balanced description among the excited states, and here we highlight the case of singly- and doubly-excited states, which differ by the number of excited electrons during the electronic transition. Most excited-state methodologies either fail to properly describe doubly-excited states or require higher-order excitations to be accounted for.<sup>19</sup> In this sense, a methodology that offers a comparable accuracy for singly- and doubly-excited states would be equally desirable.

In light of these motivations, there has been an ever-growing interest in so-called state-specific methods. The general principle is to employ a single formalism, approaching each state of interest independently, and without resorting to any prior knowledge about the other states. The first and probably the most well-known state-specific method is  $\Delta$ SCF,<sup>20,21</sup> where excited states are described by a single determinant and represent higher-lying solutions of the HF or KS equations. By optimizing the orbitals for a non-Aufbau determinant,  $\Delta$ SCF attempts to recover relaxation effects already at a mean-field level. There is a growing body of evidence showing that DFTbased  $\triangle$ SCF usually outperforms TD-DFT,<sup>22–32</sup> most notably for doubly-excited and charge transfer states.<sup>25,26</sup> However,  $\Delta$ SCF still represents a major limitation to open-shell singlet states, because of strong spin-contamination associated with the single-determinant ansatz. Restricted open-shell Kohn-Sham (ROKS)<sup>22,33</sup> offers one way around this problem, by optimizing the orbitals for a Lagrangian that considers both the mixed-spin determinant and the triplet determinant with spin projection  $m_s = 1$ . In wave-function-based methods, excitedstate mean field (ESMF) theory<sup>27-29</sup> has been proposed as a state-specific alternative to excited states. In the ESMF approach, excited-state orbitals are optimized for a CI with single excitations (CIS) ansatz,<sup>16</sup> and energies can be further

a)Electronic mail: fkossoski@irsamc.ups-tlse.fr

<sup>&</sup>lt;sup>b)</sup>Electronic mail: loos@irsamc.ups-tlse.fr

corrected with second-order Møller-Plesset (MP2) perturbation theory.<sup>27,34</sup> An extension of ESMF to DFT has also been proposed.<sup>35</sup> Variants of CC methods that directly target excited states have also been actively pursued.<sup>36–39</sup>

An important practical question for all the aforementioned methods concerns the optimization of orbitals for excited states, which typically appear as saddle point solutions in the orbital parameter space,  $^{39-42}$  therefore being more difficult to obtain than ground-state solutions.<sup>43–45</sup> In this sense, specialized algorithms for obtaining excited-state orbitals have been proposed and developed by several groups.<sup>23–25,30,31</sup> Related methods that aim at describing multiple states within the same theoretical framework, though usually in a state-averaged fashion, include complete active space self-consistent field,<sup>46</sup> ensemble DFT,<sup>47–56</sup> and multi-state TDDFT.<sup>57–61</sup>

Here we propose state-specific CI as a novel route for excited-state calculations. First, the orbitals are optimized for a minimal set of configuration state functions (CSFs), as illustrated in Fig. 1, which provides a state-specific reference. By running separate calculations for the ground state and for a targeted excited state, excitation energies can be obtained as the energy difference between the individual total energies. We label this approach  $\Delta$ CSF, in close parallel to the  $\Delta$ SCF method. By being a proper eigenstate of the total spin operator,  $\Delta$ CSF cures the spin-contamination problem of  $\Delta$ SCF, thus leading to truly state-specific orbitals and an improved reference, particularly for singlet excited states. Being a mean-field method [with an  $O(N^5)$  computational cost associated with the integral transformation, where N is the number of basis functions],  $\Delta$ CSF is intended to provide a balanced set of reference wave functions for a subsequent, more accurate calculation. At this second stage, which we label  $\Delta CI$ , correlation effects are captured by performing separate CI calculations for each state. When accounting for all single and double excitations, we obtain the  $\Delta$ CISD model, which is now expected to provide decent excitation energies with an  $O(N^6)$  computational scaling. Notice that, because we perform all singles and doubles with respect to each reference determinant, the maximum excitation degree is potentially higher than two (except of course for a one-determinant reference). This also applies to higher-order CI calculations. In this way, each state is described as much as possible in a state-specific way, with a different set of orbitals as well as determinants.

We can further compute the renormalized second-order Epstein-Nesbet (EN2) perturbation correction<sup>62</sup> from the determinants left outside the truncated CISD space of each calculation, giving rise to the  $\Delta$ CISD+EN2 model. The EN2 perturbative correction involves a single loop over external determinants that are connected to the internal determinants via at most double excitations, thus entailing an overall  $O(N^8)$  scaling associated with the number of quadruply-excited determinants. Despite this unfavorable scaling, the corresponding prefactor of the EN perturbative correction is rather small, making such calculations still affordable. Alternatively, we could calculate one of the several types of a posteriori Davidson corrections<sup>17,63,64</sup> in a state-specific fashion, leading to a  $\Delta$ CISD+Q approach. We recall that computing Davidson corrections is virtually free, such that  $\Delta$ CISD+Q presents the



FIG. 1. Types of configuration state functions (CSFs) employed as a reference for different classes of excited states in our  $\Delta$ CSF and  $\Delta$ CISD approaches.

same computational cost and  $O(N^6)$  scaling as  $\Delta$ CISD.<sup>65</sup>

The remaining question is how to build an appropriate reference for each state of interest. Our general guideline is to select the smallest set of CSFs that provides a qualitatively correct description of the state of interest, as shown in Fig. 1. Here we adopted the spin-restricted formalism. The HF determinant is the obvious choice for the ground state of closed-shell singlets. For singly-excited states of closed-shell systems, we chose either one or two CSFs, depending on each particular excited state. For most cases, a single CSF associated with two unpaired electrons should be enough. Some excited states, however, display a strong multireference character, like those of N<sub>2</sub>, CO<sub>2</sub>, and acetylene, thus requiring two CSFs. For genuine doubly-excited states where a pair of opposite-spin electrons are promoted from the same occupied to the same virtual orbital, we selected a single determinant associated with the corresponding double excitation. For ground and excited states of open-shell doublets, a single-determinant restricted open-shell HF reference is adopted as well.

Strictly speaking, our  $\Delta$ CISD approach can be seen as a type of multireference CI (MRCI), though with two key differences with respect to typical realizations of MRCI.<sup>17</sup> First, it relies on a minimal set of CSFs as the reference space, whereas in the conventional form of MRCI the reference is built from a much larger complete active space. This means that the CI space becomes more amenable in the former approach, enabling calculations for larger systems. The second important difference is that the reference in  $\Delta$ CISD is state-specific, which is expected to favor the overall fitness of the orbitals when compared with state-averaged orbitals of standard MRCI (whenever excited states are involved).  $\Delta CISD$  also resembles the ESMF theory<sup>27–29</sup> of Neuscamman and coworkers in their underlying motivation: a state-specific mean-field-like starting point, subject to a subsequent treatment of correlation effects. In  $\Delta$ CISD, however, the starting point is much more

compact and arguably closer to a mean-field description than the CIS-like ansatz of ESMF. This makes the CI expansion up to double excitations feasible in our approach, though not in ESMF, which in turn resorts to generalized MP2 to describe correlation.<sup>27,34</sup> This  $\Delta$ CSF ansatz has already been suggested as a more compact alternative to the ESMF one,<sup>28</sup> but again in the spirit of recovering correlation at the MP2 level, whereas we propose a state-specific CISD expansion, that could be followed by Davidson or EN2 perturbative corrections.

Our state-specific CI approach was implemented in QUANTUM PACKAGE,<sup>62</sup> whose determinant-driven framework provides a very convenient platform for including arbitrary sets of determinants in the CI expansion. In this way, we can easily select only the determinants that are connected to the reference determinants according to a given criterion provided by the user. On top of that, the state-specific implementation further profits from the *configuration interaction using a perturbative* selection made iteratively (CIPSI) algorithm<sup>66–69</sup> implemented in QUANTUM PACKAGE, which allows for a large reduction of the CI space without loss of accuracy. At each iteration of the CIPSI algorithm, the CI energies are obtained with the Davidson iterative algorithm,<sup>70</sup> which is ended when the EN2 perturbation correction computed in the truncated CI space lies below 0.01 m $E_{\rm h}$ .<sup>69</sup> Our state-specific CI implementation can be employed for different selection criteria for the excited determinants, based for example on the seniority number,<sup>71</sup> the hierarchy parameter,<sup>72</sup> or the excitation degree. Here, we considered the more traditional excitation-based CI. After the CI calculation, we computed the renormalized EN2 perturbation correction<sup>62</sup> from the determinants left outside the truncated CI space, which is relatively cheap because of the semi-stochastic nature of our algorithm.<sup>73</sup> We also evaluate the seven variants of Davidson corrections discussed in Ref. 17. To get statespecific orbitals, we first run a CIS calculation and obtained the natural transition orbitals (NTOs),<sup>74</sup> which proved to be more suitable guess orbitals than the canonical HF orbitals. The dominant hole and particle NTOs are taken as the singlyoccupied orbitals, and for pronounced multireference states, the second most important pair of NTOs was also considered (as illustrated in Fig. 1). The orbital optimization was performed with the Newton-Raphson method, also implemented in quantum package.<sup>39,41</sup>

Having our state-specific approaches presented, our main goal here is to assess their performance in describing electronic excited states. For that, we calculated vertical excitation energies for an extensive set of 294 electronic transitions, for systems, states, and geometries provided in the QUEST database.<sup>75</sup> We considered small-<sup>76,77</sup> and mediumsized<sup>78,79</sup> organic compounds, radicals and "exotic" systems,<sup>80</sup> and doubly-excited states.<sup>77-79</sup> The set of excited states comprises closed-shell (singlets and triplets) and open-shell (doublets) systems, ranging from one up to six non-hydrogen atoms, and of various characters (valence and Rydberg states, singlyand doubly-excited states). We employed the aug-cc-pVDZ basis set for systems having up to three non-hydrogen atoms, and the 6-31+G(d) basis set for the larger ones. We compared the excitation energies obtained with our state-specific approaches against more established alternatives, such as CIS,<sup>81</sup> CIS with

perturbative doubles [CIS(D)],<sup>82,83</sup> CC with singles and doubles (CCSD)<sup>13,84–86</sup> and the second-order approximate CC with singles and doubles (CC2),<sup>87,88</sup> the latter two understood as EOM-CC. The excitation energies obtained with the different methodologies were gauged against very accurate reference values, of high-order CC or extrapolated full CI quality.<sup>76–80</sup> The complete set of reference methods and energies are provided in the Supporting Information.

Our first important result is that the combination of the Newton-Raphson method starting with NTOs proved to be quite reliable in converging the  $\Delta$ CSF ansatz to excited-state solutions. In most cases, the orbitals are optimized with relatively few iterations (typically less than 10), and to the correct targeted state. A second-order method such as Newton-Raphson is required if the targeted solution is a saddle point in the orbital rotation landscape, which is expected to be the case for excited states.<sup>42,89</sup> At convergence, the number of negative eigenvalues of the orbital Hessian matrix, i.e., the saddle point order, can provide further insights about the topology of the solutions for a given CSF ansatz. For the closed-shell systems, we found that the lowest-lying solution (global minimum) obtained with the open-shell CSF is always an excited state since it cannot properly describe the closed-shell ground state. In turn, higher-lying excited states tend to appear as saddle points, with increasing order as one goes up in energy, even though this behavior is not very systematic. It was not uncommon, for example, to encounter two different excited states as local minima or sharing the same saddle point order. For some systems, we searched for symmetry-broken solutions of excited states by rotating the orbitals along the direction associated with a negative eigenvalue of the orbital Hessian, but this procedure leads to solutions representing a different state. We did not explore this exhaustively though, and we cannot rule out the existence of symmetry-broken excited-state solutions. Importantly, state-specific solutions could be found for different types of states, including singly- and doubly-excited states, for closed-shell singlets and open-shell doublets, and for the first as well as higher-lying states of a given point group symmetry. The full list of saddle point orders is shown in the Supporting Information.

Before diving into the main discussion, we mention that the excited-state reference could be based on single-determinant  $\Delta$ SCF orbitals, rather than the  $\Delta$ CSF orbitals we have adopted. However, the former method is heavily spin-contaminated, being an exact mixture of singlet and triplet, whereas the latter method targets one spin multiplicity at a time. In this way, the excitation energies obtained with  $\Delta$ CSF appear above (for singlets) and below (for triplets) the single energy obtained with  $\Delta$ SCF, overall improving the comparison with the reference values. In turn, we compared  $\Delta$ SCF and  $\Delta$ CSF excited-state orbitals for  $\Delta$ CISD calculations, and found overall little differences in the excitation energies. Still, we think  $\Delta$ CSF is preferable because it delivers truly state-specific orbitals, whereas  $\Delta$ SCF produces the same orbitals for the singlet and triplet states, and is thus less state-specific.

The state-specific  $\Delta$ CI approach offers a well-defined route towards full CI by increasing the excitation degree, in analogy with standard ground-state-based CI methods. We explored

both routes by calculating 16 excitation energies for small systems, by considering up to quadruple excitations (full set of results are available in the Supporting Information). Even though this is a small set for obtaining significant statistics, it is enough to showcase the main trends when comparing statespecific and ground-state-based CI methods. The mean signed error (MSE) and mean absolute error (MAE) are shown in Table I. The convergence for standard CI is quite slow, with CISD largely overestimating the excitation energies, CISDT leading to more decent results, which are improved at the CISDTQ level. In turn, we found that  $\Delta CI$  displays much more accurate results and accelerated convergence than their ground-statebased counterparts. Already at the  $\Delta$ CISD level, the accuracy is far superior to that of standard CISD, being comparable to CISDT. Going one step further ( $\Delta$ CISDT) does not lead to a visible improvement, whereas the state-specific quadruple excitations of  $\Delta$ CISDTQ recover much of the remaining correlation energy, hence the motivation to investigate various flavors of Davidson correction. We found that the MAEs of  $\Delta$ CISD are comparable to that of CISDT, which can be understood from the observation that the doubly-excited determinants accessed from the excited-state reference can only be achieved via triple excitations from the ground-state reference. The comparison between state-specific and ground-state-based CI for a given excitation degree ( $\Delta$ CISD against CISD and  $\Delta$ CISDTQ against CISDTQ) shows that the MAEs are reduced by one order of magnitude in the former route, when compared with the latter. No gain is observed from CISDT to  $\Delta$ CISDT, though.

TABLE I. Mean Signed Error (MSE) and Mean Absolute Error (MAE), in Units of eV, with Respect to Reference Theoretical Values, for the Set of 16 Excitation Energies Listed in the Supporting Information.

method	MSE	MAE
CISD	+3.91	3.91
CISDT	+0.07	0.17
CISDTQ	+0.13	0.13
ΔCISD	-0.14	0.18
∆CISDT	-0.20	0.20
∆CISDTQ	-0.02	0.02
$\Delta \text{CISD}+\text{EN2}$	-0.00	0.03
$\Delta CISD+PC$	-0.10	0.14
CIS(D)	-0.03	0.35
CC2	-0.05	0.32
CCSD	+0.01	0.08
CC3	+0.01	0.03
CCSDT	-0.01	0.02

We now start the discussion on how well our state-specific CI approaches compare with more established methods by presenting in Fig. 2 the distribution of errors and statistical measures associated with a set of 237 singly-excited states of closed-shell systems. At the  $\Delta$ CSF level, the excitation energies are systematically underestimated, thus a substantially negative MSE. This happens because the CSF reference for the excited states (typically containing two determinants) already recovers some correlation, whereas the one-determinant HF reference of the ground state does not. The MAE of the  $\Delta$ CSF

approach (0.62 eV) is comparable to that of CIS (0.65 eV). The overall similar performance of these two methods is somewhat expected since the orbital relaxation that takes place in the state-specific CSF is partially described via the single excitations of CIS. Moving to the  $\Delta$ CISD level, we find that correlation effects are described in a reasonably balanced way for ground and excited states. The MAE is significantly reduced (0.18 eV) with respect to that of  $\Delta$ CSF, being smaller than in CIS(D) (0.21 eV) and comparable to CC2 (0.17 eV). The absolute MSE also decreases, but remains negative, whereas the other CI- or CC-based methods present positive MSEs. This shows that there is still some bias toward a better description of excited states at the  $\Delta$ CISD level, probably reflecting the two-determinant reference (compared to one-determinant for the ground state).

The perturbative correction introduced at the  $\Delta CISD + EN2$ approach reduces the statistical errors even more, showing the same MAE as CCSD (0.06 eV). At times, however, the EN2 correction leads to erroneous results due to the presence of intruder states, which typically occurs for some higher-lying states of a given symmetry.<sup>90</sup> Instead of relying on perturbation theory to correct the CISD total energies, we can resort to one of Davidson corrections.<sup>17</sup> Even though this correction is not as accurate as the EN2 perturbative energy, more often than not it improves upon  $\Delta$ CISD, and with virtually no additional computational cost.<sup>91</sup> We found that all seven  $\Delta CISD+Q$  variants provide MAEs in the 0.10 eV to 0.12 eV range. Since the Pople correction<sup>64</sup> is arguably the most well-behaved one, with fewer outlier excitation energies, we only present here the statistical measures associated with this correction, labeled  $\Delta$ CISD+PC from hereon, while the full set of results can be found in the Supporting Information.

We also surveyed the performance of our state-specific methods for 10 genuine doubly-excited states<sup>19</sup> and 47 excited states of open-shell doublets (doublet-doublet transitions),<sup>80</sup> both sets were extracted from the QUEST database.<sup>75</sup> The statistical measures are shown in Table S1 of the Supporting Information, together with those of singly-excited states of closed-shell systems. The important finding in this comparison is that the state-specific methods provide reasonably similar MAEs for the three types of excited states. For instance,  $\Delta$ CISD has MAEs of 0.18 eV for singly-excited states of closed-shell singlets, 0.17 eV for doublet-doublet transition and 0.16 eV for doubly-excited states. This contrasts with the case of more familiar methods, which cannot describe doubly-excited states unless higher-order excitations are included.<sup>19</sup> We notice that the MSE of  $\Delta$ CSF is more negative for singly-excited states of closed-shell molecules (-0.55 eV) than for doubly-excited states (-0.20 eV), being closer to zero for doublet-doublet transitions (0.07 eV), which reflects the one-determinant reference adopted for both the excited and ground states in the latter cases. This difference does not translate into comparatively smaller errors in the correlated results though.

For the doubly-excited states, we further compare in Table II the performance of state-specific CI against higher-order CC methods. The accuracy of the  $\Delta$ CSF mean-field model is superior to CC3 and approaches CCSDT, which highlights the importance of orbital relaxation for doubly-excited states.



FIG. 2. Distribution of Errors in Excitation Energies with Respect to Reference Theoretical Values and Corresponding Mean Signed Error (MSE) and Mean Absolute Error (MAE), for Various Excited-State Methodologies.

 $\Delta$ CISD is significantly more accurate than CCSDT, whereas the perturbative and the Davidson corrections bring a small improvement.

TABLE II. Mean Signed Error (MSE) and Mean Absolute Error (MAE), in Units of eV, with Respect to Reference Theoretical Values, for the Set of 10 Doubly-Excited States Listed in the Supporting Information.

Method	MSE	MAE
ΔCSF	-0.20	0.46
ΔCISD	+0.03	0.16
$\Delta CISD+EN2$	-0.05	0.13
$\Delta \text{CISD+PC}$	+0.02	0.13
CC3	+0.85	0.85
CCSDT	+0.38	0.38
CCSDTQ	+0.03	0.03

The performance of our state-specific methods can also be assessed for specific types of excited states, say for  $\pi\pi^*$  transitions or for systems of a given size. This is shown in Table III, which compares the MAEs across different categories. Many trends can be identified, but here we highlight the most notorious and interesting ones, starting with spin multiplicity. While the  $\Delta$ CISD results are comparable for singlets and triplets, we found that the perturbative correction has a more pronounced effect for the triplets, bringing the MAE down to 0.06 eV, the same as obtained with CCSD. To some extent, the worse performance of the EN2 correction for the singlets stems from intruder states (most noticeably the ground state when it shares the same point group symmetry as the targeted excited state). We also found that the Davidson corrections bring a somewhat larger improvement for triplets than for singlets, with some flavors having MAEs of 0.07 eV for the triplets, being essentially as accurate as CCSD (MAE of 0.06 eV). For the

doublet-doublet transitions, the EN2 and +Q corrections are as helpful as they are for the triplets (see Table S1 of the Supporting Information).

Regarding the character of the excitations, we found that  $\Delta$ CISD is considerably better for Rydberg (MAE of 0.12 eV) than for valence (MAE of 0.21 eV) excited states. Interestingly, the difference at the  $\Delta$ CSF level is proportionally smaller (0.65 eV against 0.56 eV), suggesting that the  $\Delta$ CISD difference actually stems from the description of correlation. In turn, the EN2 correction has a larger impact on valence excitations, making little difference for the Rydberg states, such that the  $\Delta$ CISD+EN2 results become comparable for both types of excitation. Additional trends can be observed when dividing the valence excitations into  $n\pi^*$ ,  $\pi\pi^*$ , or  $\sigma\pi^*$ , and the Rydberg excitations as taking place from n or  $\pi$  orbitals. Our state-specific methods are typically more accurate for  $n\pi^*$  excitations than for  $\pi\pi^*$  excitations.  $\Delta$ CISD+EN2, for example, is as accurate as CCSD for  $n\pi^*$  transitions, with corresponding MAEs of 0.06 eV. We also found that the less common  $\sigma\pi^*$  excitations are much better described across all methods than the more typical  $n\pi^*$  and  $\pi\pi^*$  transitions. For this type of state,  $\Delta$ CISD+EN2 is the best performing method, with MAEs as small as 0.03 eV. When separating the Rydberg states by the character of the hole orbital, n or  $\pi$ , additional interesting features can be seen. Except for CCSD, all the other methods considered here provide more accurate results for the Rydberg excitations involving the  $\pi$  orbitals. Not only that, but the MAEs are quite small and comparable across all methods (except for  $\triangle$ CSF and CIS), ranging from 0.06 eV to 0.11 eV. Surprisingly, CIS is much more accurate for  $\pi$  Rydberg (MAE of 0.29 eV) than for n Rydberg (MAE of 1.17 eV) excitations.

The third and most important line of comparison concerns the system size. Under this criterion, we divided the excited states into three groups, small, medium, and large, depending on the number of non-hydrogen atoms (see Table III). We

	# States	∆CSF	ΔCISD	$\Delta CISD+EN2$	$\Delta CISD+PC$	CIS	CIS(D)	CC2	CCSD
All states	237	0.62	0.18	0.08	0.10	0.65	0.21	0.17	0.08
Singlet	127	0.56	0.17	0.10	0.12	0.68	0.22	0.19	0.10
Triplet	110	0.69	0.18	0.06	0.09	0.61	0.19	0.15	0.06
Valence	155	0.65	0.21	0.08	0.10	0.61	0.19	0.14	0.08
Rydberg	82	0.56	0.12	0.10	0.11	0.72	0.24	0.22	0.08
$n\pi^*$	56	0.60	0.16	0.06	0.07	0.54	0.10	0.09	0.06
$\pi\pi^*$	80	0.75	0.26	0.10	0.12	0.74	0.27	0.20	0.10
$\sigma\pi^*$	18	0.39	0.13	0.03	0.06	0.26	0.11	0.07	0.04
n Rydberg	40	0.61	0.12	0.12	0.14	1.17	0.37	0.39	0.07
$\pi$ Rydberg	38	0.46	0.10	0.08	0.09	0.29	0.11	0.06	0.10
1-2 non-H atoms	69	0.83	0.18	0.06	0.12	0.71	0.24	0.23	0.06
3-4 non-H atoms	122	0.57	0.18	0.09	0.10	0.70	0.20	0.16	0.08
5-6 non-H atoms	46	0.41	0.15	0.12	0.07	0.43	0.18	0.12	0.10

TABLE III. Mean Absolute Error, in Units of eV, with Respect to Reference Theoretical Values, for Different Types of Singly-Excited States of Closed-Shell Systems Listed in the Supporting Information.

found that  $\Delta$ CSF becomes more accurate as the system size increases, which we assign to a diminishing effect of the onevs two-determinant imbalance discussed above. As the system size increases, the correlation energy recovered by the two determinants of the excited states is expected to become smaller in comparison to the total correlation energy, thus alleviating this imbalance. In contrast, CISD should provide less accurate total energies for larger systems, due to its well-known lack of size-consistency. This issue would be expected to reflect on excitation energies to some degree, which are not absolute but relative energies. However, a more balanced reference provided by  $\Delta$ CSF might compensate for the lack of sizeconsistency when larger systems are targeted. Indeed,  $\Delta CISD$ presents comparable MAEs across the three sets of system size (0.15 eV to 0.18 eV). In contrast,  $\Delta$ CISD+Q and  $\Delta$ CISD+EN2 seem to go opposite ways: the former becomes more accurate and the latter less as a function of system size. Similarly, CC2 becomes more accurate and CCSD loses accuracy as the system size increases,  $^{78,92}$  to the point where the theoretically more approximate CC2 becomes the favored methodological choice. Our results thus show that state-specific methods represent a competitive alternative for larger systems. For molecules containing five or six non-hydrogen atoms,  $\Delta$ CISD+EN2 becomes practically as accurate as CCSD, with MAEs in the 0.10 eV to 0.12 eV range. The  $\Delta$ CISD+Q models turn out to be the most accurate choice for systems of this size, with MAEs ranging from 0.07 eV to 0.09 eV (see Table S5 of the Supporting Information).

Butadiene, glyoxal, the carbon dimer  $(C_2)$  and trimer  $(C_3)$  are particularly interesting and challenging systems that deserve a dedicated discussion. The excitation energies for these molecules can be found in the Supporting Information.

The dark  $2^1A_g$  excited state of butadiene is a notoriously famous example, having received much attention (see Ref. 19 and references therein). Prior studies had assigned it as a doubly-excited state, due to important contributions from doubly-excited determinants.<sup>93,94</sup> More recently, though, it has been re-assigned as a singly-excited state,<sup>95</sup> meaning that the doubly-excited determinants actually represent strong orbital relaxation effects. Here, our state-specific results support this interpretation, since one CSF associated with a single excitation provided reasonable excitation energies, whereas attempts to employ a doubly-excited reference produced much higher-lying solutions. Already at the  $\Delta$ CSF level, we obtained an excitation energy (7.18 eV) comparable to the much more expensive CCSD (7.20 eV), though still overestimating the CCSDTQ reference value of 6.56 eV.78 This result demonstrates the ability of  $\Delta$ CSF to capture orbital relaxation effects, at only a mean-field cost, which in contrast needs at least double excitations in EOM-CC. Inclusion of correlation at the  $\Delta$ CISD level brings the excitation energy down to 6.93 eV. An important question in butadiene concerns the energy gap between the  $2^{1}A_{g}$  dark state and the lower-lying  $1^{1}B_{u}$  bright state, whose correct ordering has only recently been settled.<sup>96</sup> Having the CCSDTQ reference value of 0.15 eV for the energy gap,<sup>78</sup> we observe that EOM-CC methods considerably overestimate it (0.94 eV in CC2 and 0.65 eV in CCSD), whereas the state-specific methods deliver improved results (0.65 eV in  $\Delta$ CSF and 0.39 eV in  $\Delta$ CISD).

Another challenging system is glyoxal, which presents excited states of genuine multireference character.<sup>97</sup> While the first pair of NTOs has a dominant weight, the second pair is non-negligible. In this sense, most of the first excited states of glyoxal lie in between the cases of most singly-excited states (that can be qualitatively described with one CSF) and those that need two CSFs. Being an intermediate case, here we performed  $\Delta$ CISD calculations with references containing one or two CSFs, for the first two singlet states and first four triplet states. With one CSF only,  $\Delta$ CSF typically overestimates the reference excitation energies, with the corresponding  $\Delta$ CISD improving the overall comparison. For this set of six excited states, the associated MAEs are 1.14 eV for  $\Delta$ CSF and 0.65 eVfor  $\Delta$ CISD when using a single CSF as reference. Despite the improvement brought at the CISD level, this is still limited by the lack of an actual multiconfigurational reference for these states. When two CSFs are employed as the reference for the excited states, the MAEs are reduced to 0.58 eV  $(\Delta CSF)$  and 0.22 eV ( $\Delta CISD$ ), which can be further decreased to 0.08 eV with  $\Delta$ CISD+PC. We thus recommend augmenting the excited-state reference whenever it displays at least some

multireference character, and the weight of the first pairs of NTOs could serve as an easy proxy for this.

Finally, we comment on the lowest-lying  $1^1\Delta_g$  and higherlying  $2^{1}\Sigma_{\rho}^{+}$  doubly-excited states of C<sub>2</sub> and C<sub>3</sub>, which would require at least CCSDTQ quality calculations to become accurate to within 0.1 eV.<sup>19</sup> C<sub>2</sub> displays a strong multireference ground state, and thus we employed two CSFs as the reference, the closed-shell HF and the determinant associated with the  $(\sigma_{2s}^*)^2 \rightarrow (\sigma_{2p_2})^2$  transition. For its doubly-excited states, we employed the two CSFs needed to describe both doublyexcited states, generated from the HF determinant through the  $(\pi_{2p_x})^2 \rightarrow (\sigma_{2p_z})^2$  and  $(\pi_{2p_y})^2 \rightarrow (\sigma_{2p_z})^2$  excitations,  $\pi_{2p_x}$  and  $\pi_{2p_y}$  being degenerate orbitals. In C<sub>3</sub>, the multireference character of the ground state is weaker, and thus we adopted a single HF determinant as reference. In turn, four CSFs are needed for its doubly-excited states, built from the HF determinant by performing  $(\sigma_g)^2 \to (\pi_{2p_x}^*)^2$ ,  $(\sigma_g)^2 \to (\pi_{2p_y}^*)^2$ ,  $(\sigma_u)^2 \to (\pi_{2p_x}^*)^2$ , and  $(\sigma_u)^2 \to (\pi_{2p_y}^*)^2$  transitions, where  $\pi_{2p_x}^*$  and  $\pi_{2p_y}^*$  are degenerate orbitals. We therefore re-assign the doubly-excited states of C<sub>3</sub> as  $(\sigma)^2 \rightarrow (\pi^*)^2$ , which had been first assigned as  $(\pi)^2 \to (\sigma^*)^2$ .<sup>19</sup> Notice that, for both systems, what differentiates  $1^1\Delta_g$  and  $2^1\Sigma_g^+$  is essentially the phase between the two CSFs differing by the occupation of the degenerate orbitals  $(\pi \text{ in } C_2, \pi^* \text{ in } C_3)$ . Thus, the higher-lying state orbitals were obtained by optimizing for the second CI root associated with the reference (two CSFs in  $C_2$ , four in  $C_3$ ). We found that  $\Delta$ CISD is more accurate than CCSDT for C<sub>2</sub>, and even more accurate than CCSDTQ for C<sub>3</sub>.

To summarize, here we have presented a novel route to perform excited-state calculations, based on a state-specific CI approach. The orbitals are optimized for a targeted state with a minimal set of CSFs, serving as the reference wave function for the CI calculations, which can be further corrected with Epstein-Nesbet perturbation theory or with a posteriori Davidson corrections. We have surveyed these methods against more established alternatives by computing excitation energies for a vast class of molecules and types of excitations from the QUEST database. State-specific CI was found to be substantially more accurate than the standard CI methods based on a ground-state reference. Importantly, it delivers reliable results across different types of excited states, most notably when comparing singly- and doubly-excited states, and can easily handle ground and excited states of multireference nature. The overall accuracy of  $\Delta$ CISD rivals that of CC2 (MAEs of 0.17 eV to 0.18 eV), whereas  $\Delta$ CISD+EN2 is comparable to CCSD (MAEs of 0.08 eV), with  $\Delta$ CISD+Q lying in-between (MAE of 0.10 eV to 0.12 eV). For larger systems,  $\Delta CISD+Q$ leads to more accurate results (MAE of 0.07 eV to 0.09 eV) than CC2 and CCSD (MAEs of 0.10 eV to 0.12 eV).

There are many exciting possibilities to be pursued from this work. One is to develop analogous state-specific coupledcluster methods. In light of the huge improvement we have observed when going from ground-state-based to state-specific CI, we expect a similar gain when comparing EOM-CC to statespecific CC methods where tailored CSFs are employed as the reference wave function.<sup>98–100</sup> One could also develop statespecific implementations of seniority-based<sup>71</sup> and hierarchybased<sup>72</sup> CI for excited states. It would be important to assess the performance of our state-specific approaches to even larger systems, which would require switching from a determinantdriven to an integral-driven implementation. Although evaluating nonorthogonal matrix elements is more challenging than their orthogonal analogs, the calculation of static properties such as dipole moments and oscillator strengths is possible thanks to the recent generalized extension of the nonorthogonal Wick's theorem proposed by Burton.<sup>101,102</sup> Yet another exciting possibility is to move from a state-specific to a stateaveraged reference, while contemplating only a small set of important determinants for describing a given set of states. This would be expected to solve some of the issues encountered here when two states of the same symmetry are strongly coupled, and to provide an even more balanced description of correlation between ground and excited states.

#### ACKNOWLEDGMENTS

This work was performed using HPC resources from CALMIP (Toulouse) under allocation 2021-18005. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 863481).

### SUPPORTING INFORMATION AVAILABLE

Additional statistical measures for different sets of excited states and for all flavors of  $\Delta CISD+Q$  models. Separate tables gathering the excitation energies for butadiene, glyoxal,  $C_2$ , and  $C_3$ . For the full set of 294 excited states, total energies and excitation energies obtained at the  $\Delta$ CSF,  $\Delta$ CISD,  $\Delta CISD+EN2$ , and seven variants of  $\Delta CISD+Q$  models, excitation energies computed with CIS, CIS(D), CC2, and CCSD, number of determinants in the reference, saddle point order associated with the  $\Delta$ CSF solutions, the reference excitation energies and corresponding method, and additional statistical measures. For a subset of 16 excited states, total energies and excitation energies obtained at the  $\Delta$ CISDT,  $\Delta$ CISDTQ and ground-state-based CISDT and CISDTQ levels of theory. For the subset of 10 doubly-excited states, excitation energies obtained at the CC3, CCSDT, CC4, and CCSDTQ levels of theory.

<sup>1</sup>Hohenberg, P.; Kohn, W. Inhomogeneous electron gas. *Phys. Rev.* **1964**, *136*, B864–B871.

- <sup>2</sup>Kohn, W.; Sham, L. J. Self-consistent equations including exchange and correlation effects. *Phys. Rev.* **1965**, *140*, A1133–A1138.
- <sup>3</sup>Parr, R. G.; Yang, W. *Density-functional theory of atoms and molecules*; Oxford: Clarendon Press, 1989.
- <sup>4</sup>Runge, E.; Gross, E. K. U. Density-Functional Theory for Time-Dependent Systems. *Phys. Rev. Lett.* **1984**, *52*, 997–1000.
- <sup>5</sup>Burke, K.; Werschnik, J.; Gross, E. K. U. Time-dependent density functional theory: Past, present, and future. *J. Chem. Phys.* 2005, *123*, 062206.
  <sup>6</sup>Casida, M.; Huix-Rotllant, M. Progress in Time-Dependent Density-Functional Theory. *Annu. Rev. Phys. Chem.* 2012, *63*, 287–323.
- <sup>7</sup>Huix-Rotllant, M.; Ferré, N.; Barbatti, M. Quantum Chemistry and Dynamics of Excited States; John Wiley & Sons, Ltd, 2020; Chapter 2, pp 13–46.

- <sup>8</sup>Čížek, J. On the Correlation Problem in Atomic and Molecular Systems. Calculation of Wavefunction Components in Ursell-Type Expansion Using Quantum-Field Theoretical Methods. J. Chem. Phys. **1966**, 45, 4256–4266.
- <sup>9</sup>Crawford, T. D.; Schaefer, H. F. *Reviews in Computational Chemistry*; John Wiley & Sons, Ltd, 2000; pp 33–136.
- <sup>10</sup>Bartlett, R. J.; Musiał, M. Coupled-Cluster Theory in Quantum Chemistry. *Rev. Mod. Phys.* **2007**, *79*, 291–352.
- <sup>11</sup>Shavitt, I.; Bartlett, R. J. Many-Body Methods in Chemistry and Physics: MBPT and Coupled-Cluster Theory; Cambridge Molecular Science; Cambridge University Press: Cambridge, 2009.
- <sup>12</sup>Rowe, D. J. Equations-of-Motion Method and the Extended Shell Model. *Rev. Mod. Phys.* **1968**, *40*, 153–166.
- <sup>13</sup>Stanton, J. F.; Bartlett, R. J. The equation of motion coupled-cluster method. A systematic biorthogonal approach to molecular excitation energies, transition probabilities, and excited state properties. *J. Chem. Phys.* **1993**, *98*, 7029–7039.
- <sup>14</sup>Monkhorst, H. J. Calculation of properties with the coupled-cluster method. *Int. J. Quantum Chem.* **1977**, *12*, 421–432.
- <sup>15</sup>Koch, H.; Kobayashi, R.; Sanchez de Merás, A.; Jorgensen, P. Calculation of size-intensive transition moments from the coupled cluster singles and doubles linear response function. J. Chem. Phys. **1994**, 100, 4393–4400.
- <sup>16</sup>Szabo, A.; Ostlund, N. S. *Modern quantum chemistry*; McGraw-Hill: New York, 1989.
- <sup>17</sup>Szalay, P. G.; Müller, T.; Gidofalvi, G.; Lischka, H.; Shepard, R. Multiconfiguration self-consistent field and multireference configuration interaction methods and applications. *Chem. Rev.* **2012**, *112*, 108–81.
- <sup>18</sup>Lischka, H.; Nachtigallová, D.; Aquino, A. J. A.; Szalay, P. G.; Plasser, F.; Machado, F. B. C.; Barbatti, M. Multireference Approaches for Excited States of Molecules. *Chem. Rev.s* **2018**, *118*, 7293–7361.
- <sup>19</sup>Loos, P. F.; Boggio-Pasqua, M.; Scemama, A.; Caffarel, M.; Jacquemin, D. Reference Energies for Double Excitations. *J. Chem. Theory Comput.* **2019**, *15*, 1939–1956.
- <sup>20</sup>Ziegler, T.; Rauk, A.; Baerends, E. J. On the calculation of multiplet energies by the Hartree-Fock-Slater method. *Theor. Chim. Acta* **1977**, *43*, 261–271.
- <sup>21</sup>Kowalczyk, T.; Yost, S. R.; Voorhis, T. V. Assessment of the ΔSCF density functional theory approach for electronic excitations in organic dyes. J. Chem. Phys. **2011**, 134, 054128.
- <sup>22</sup>Kowalczyk, T.; Tsuchimochi, T.; Chen, P.-T.; Top, L.; Van Voorhis, T. Excitation energies and Stokes shifts from a restricted open-shell Kohn-Sham approach. J. Chem. Phys. **2013**, *138*, 164101.
- <sup>23</sup>Gilbert, A. T.; Besley, N. A.; Gill, P. M. Self-consistent field calculations of excited states using the maximum overlap method (MOM). *J. Phys. Chem.* A 2008, 112, 13164–13171.
- <sup>24</sup>Barca, G. M.; Gilbert, A. T.; Gill, P. M. Simple Models for Difficult Electronic Excitations. J. Chem. Theory Comput. **2018**, 14, 1501–1509.
- <sup>25</sup>Hait, D.; Head-Gordon, M. Excited State Orbital Optimization via Minimizing the Square of the Gradient: General Approach and Application to Singly and Doubly Excited States via Density Functional Theory. *J. Chem. Theory Comput.* **2020**, *16*, 1699–1710.
- <sup>26</sup>Hait, D.; Head-Gordon, M. Orbital Optimized Density Functional Theory for Electronic Excited States. J. Phys. Chem. Lett. **2021**, 12, 4517–4529.
- <sup>27</sup>Shea, J. A. R.; Neuscamman, E. Communication: A mean field platform for excited state quantum chemistry. J. Chem. Phys. 2018, 149, 081101.
- <sup>28</sup>Shea, J. A. R.; Gwin, E.; Neuscamman, E. A Generalized Variational Principle with Applications to Excited State Mean Field Theory. *J. Chem. Theory Comput.* **2020**, *16*, 1526–1540.
- <sup>29</sup>Hardikar, T. S.; Neuscamman, E. A self-consistent field formulation of excited state mean field theory. J. Chem. Phys. **2020**, 153, 164108.
- <sup>30</sup>Levi, G.; Ivanov, A. V.; Jónsson, H. Variational Density Functional Calculations of Excited States via Direct Optimization. *J. Chem. Theory Comput.* 2020,
- <sup>31</sup>Carter-Fenk, K.; Herbert, J. M. State-Targeted Energy Projection: A Simple and Robust Approach to Orbital Relaxation of Non-Aufbau Self-Consistent Field Solutions. J. Chem. Theory Comput. 2020, 16, 5067–5082.
- <sup>32</sup>Toffoli, D.; Quarin, M.; Fronzoni, G.; Stener, M. Accurate Vertical Excitation Energies of BODIPY/Aza-BODIPY Derivatives from Excited-State Mean-Field Calculations. *The J. Phys. Chem. A* **2022**, *126*, 7137–7146.
- <sup>33</sup>Filatov, M.; Shaik, S. A spin-restricted ensemble-referenced Kohn–Sham method and its application to diradicaloid situations. *Chem. Phys. Lett.*

**1999**, *304*, 429–437.

- <sup>34</sup>Clune, R.; Shea, J. A. R.; Neuscamman, E. N5-Scaling Excited-State-Specific Perturbation Theory. J. Chem. Theory Comput. 2020, 16, 6132– 6141.
- <sup>35</sup>Zhao, L.; Neuscamman, E. Density Functional Extension to Excited-State Mean-Field Theory. J. Chem. Theory Comput. 2020, 16, 164–178.
- <sup>36</sup>Piecuch, P.; Kowalski, K. Computational Chemistry: Reviews of Current Trends; Computational Chemistry: Reviews of Current Trends; WORLD SCIENTIFIC, 2000; Vol. Volume 5; pp 1–104.
- <sup>37</sup>Mayhall, N. J.; Raghavachari, K. Multiple Solutions to the Single-Reference CCSD Equations for NiH. J. Chem. Theory Comput. **2010**, *6*, 2714–2720.
- <sup>38</sup>Lee, J.; Small, D. W.; Head-Gordon, M. Excited States via Coupled Cluster Theory without Equation-of-Motion Methods: Seeking Higher Roots with Application to Doubly Excited States and Double Core Hole States. *J. Chem. Phys.* **2019**, *151*, 214103.
- <sup>39</sup>Kossoski, F.; Marie, A.; Scemama, A.; Caffarel, M.; Loos, P.-F. Excited States from State-Specific Orbital-Optimized Pair Coupled Cluster. J. Chem. Theory Comput. 2021, 17, 4756.
- <sup>40</sup>Marie, A.; Kossoski, F.; Loos, P.-F. Variational coupled cluster for ground and excited states. J. Chem. Phys. **2021**, 155, 104105.
- <sup>41</sup>Damour, Y.; Véril, M.; Kossoski, F.; Caffarel, M.; Jacquemin, D.; Scemama, A.; Loos, P.-F. Accurate full configuration interaction correlation energy estimates for five- and six-membered rings. *J. Chem. Phys.* **2021**, *155*, 134104.
- <sup>42</sup>Burton, H. G. Energy Landscape of State-Specific Electronic Structure Theory. J. Chem. Theory Comput. 2022, 18, 1512–1526.
- <sup>43</sup>Cuzzocrea, A.; Scemama, A.; Briels, W. J.; Moroni, S.; Filippi, C. Variational Principles in Quantum Monte Carlo: The Troubled Story of Variance Minimization. J. Chem. Theory Comput. **2020**, *16*, 4203–4212.
- <sup>44</sup>Otis, L.; Craig, I. M.; Neuscamman, E. A hybrid approach to excited-state-specific variational Monte Carlo and doubly excited states. *J. Chem. Phys.* **2020**, *153*, 234105.
- <sup>45</sup>Shepard, S.; Panadés-Barrueta, R. L.; Moroni, S.; Scemama, A.; Filippi, C. Double Excitation Energies from Quantum Monte Carlo Using State-Specific Energy Optimization. J. Chem. Theory Comput. 0, 0, null, PMID: 36314602.
- <sup>46</sup>Roos, B. O.; Taylor, P. R.; Sigbahn, P. E. A complete active space SCF method (CASSCF) using a density matrix formulated super-CI approach. *Chem. Phys.* **1980**, *48*, 157–173.
- <sup>47</sup>Cernatic, F.; Senjean, B.; Robert, V.; Fromager, E. Ensemble Density Functional Theory of Neutral and Charged Excitations. *Top. Curr. Chem.* **2022**, *380*, 1–80.
- <sup>48</sup>Gould, T.; Hashimi, Z.; Kronik, L.; Dale, S. G. Single Excitation Energies Obtained from the Ensemble "HOMO–LUMO Gap": Exact Results and Approximations. J. Phys. Chem. Lett. **2022**, 13, 2452–2458.
- <sup>49</sup>Gould, T.; Kronik, L.; Pittalis, S. Double excitations in molecules from ensemble density functionals: Theory and approximations. *Phys. Rev. A* **2021**, *104*, 022803.
- <sup>50</sup>Marut, C.; Senjean, B.; Fromager, E.; Loos, P.-F. Weight dependence of local exchange–correlation functionals in ensemble density-functional theory: double excitations in two-electron systems. *Faraday Discuss.* **2020**, *224*, 402.
- <sup>51</sup>Loos, P.-F.; Fromager, E. A weight-dependent local correlation densityfunctional approximation for ensembles. *J. Chem. Phys.* **2020**, *152*, 214101.
- <sup>52</sup>Deur, K.; Fromager, E. Ground and excited energy levels can be extracted exactly from a single ensemble density-functional theory calculation. *J. Chem. Phys.* **2019**, *150*, 094106.
- <sup>53</sup>Gould, T.; Kronik, L.; Pittalis, S. Charge Transfer Excitations from Exact and Approximate Ensemble Kohn-Sham Theory. J. Chem. Phys. 2018, 148, 174101.
- <sup>54</sup>Deur, K.; Mazouin, L.; Fromager, E. Exact Ensemble Density Functional Theory for Excited States in a Model System: Investigating the Weight Dependence of the Correlation Energy. *Phys. Rev. B* 2017, *95*, 035120.
- <sup>55</sup>Filatov, M. In *Density-Functional Methods for Excited States*; Ferré, N., Filatov, M., Huix-Rotllant, M., Eds.; Springer International Publishing: Cham, 2016; pp 97–124.
- <sup>56</sup>Senjean, B.; Knecht, S.; Jensen, H. J. A.; Fromager, E. Linear Interpolation Method in Ensemble Kohn-Sham and Range-Separated Density-Functional Approximations for Excited States. *Phys. Rev. A* **2015**, *92*, 012518.

- <sup>57</sup>Gao, J.; Grofe, A.; Ren, H.; Bao, P. Beyond Kohn–Sham Approximation: Hybrid Multistate Wave Function and Density Functional Theory. J. Phys. Chem. Lett. 2016, 7, 5143–5149.
- <sup>58</sup>Lu, Y.; Gao, J. Multistate Density Functional Theory of Excited States. J. Phys. Chem. Lett. **2022**, 13, 7762–7769.
- <sup>59</sup>Ren, H.; Provorse, M. R.; Bao, P.; Qu, Z.; Gao, J. Multistate Density Functional Theory for Effective Diabatic Electronic Coupling. J. Phys. Chem. Lett. 2016, 7, 2286–2293.
- <sup>60</sup>Horbatenko, Y.; Lee, S.; Filatov, M.; Choi, C. H. Performance Analysis and Optimization of Mixed-Reference Spin-Flip Time-Dependent Density Functional Theory (MRSF-TDDFT) for Vertical Excitation Energies and Singlet–Triplet Energy Gaps. J. Phys. Chem. A **2019**, *123*, 7991–8000.
- <sup>61</sup>Horbatenko, Y.; Sadiq, S.; Lee, S.; Filatov, M.; Choi, C. H. Mixed-Reference Spin-Flip Time-Dependent Density Functional Theory (MRSF-TDDFT) as a Simple yet Accurate Method for Diradicals and Diradicaloids. *J. Chem. Theory Comput.* **2021**, *17*, 848–859.
- <sup>62</sup>Garniron, Y. et al. Quantum Package 2.0: a open-source determinant-driven suite of programs. J. Chem. Theory Comput. 2019, 15, 3591.
- <sup>63</sup>Langhoff, S. R.; Davidson, E. R. Configuration interaction calculations on the nitrogen molecule. *Int. J. Quantum Chem.* **1974**, *8*, 61–72.
- <sup>64</sup>Pople, J. A.; Seeger, R.; Krishnan, R. Variational configuration interaction methods and comparison with perturbation theory. *Int. J. Quantum Chem.* **1977**, *12*, 149–163.
- <sup>65</sup>It is worth noting that the ΔCSF, ΔCISD, and ΔCISD+Q excitation energies are invariant under rotations within the doubly-occupied and virtual blocks, although the ΔCISD+EN2 energies are not, which is well-known for the EN2 perturbative correction.<sup>62</sup>.
- <sup>66</sup>Huron, B.; Malrieu, J. P.; Rancurel, P. Iterative perturbation calculations of ground and excited state energies from multiconfigurational zeroth-order wavefunctions. J. Chem. Phys. **1973**, 58, 5745–5759.
- <sup>67</sup>Giner, E.; Scemama, A.; Caffarel, M. Using perturbatively selected configuration interaction in quantum Monte Carlo calculations. *Can. J. Chem.* **2013**, *91*, 879–885.
- <sup>68</sup>Giner, E.; Scemama, A.; Caffarel, M. Fixed-node diffusion Monte Carlo potential energy curve of the fluorine molecule F2 using selected configuration interaction trial wavefunctions. *J. Chem. Phys.* **2015**, *142*, 044115.
- <sup>69</sup>Garniron, Y.; Scemama, A.; Giner, E.; Caffarel, M.; Loos, P. F. Selected Configuration Interaction Dressed by Perturbation. *J. Chem. Phys.* 2018, *149*, 064103.
- <sup>70</sup>Davidson, E. R. The iterative calculation of a few of the lowest eigenvalues and corresponding eigenvectors of large real-symmetric matrices. *J. Comput. Phys.* **1975**, *17*, 87–94.
- <sup>71</sup>Bytautas, L.; Henderson, T. M.; Jiménez-Hoyos, C. A.; Ellis, J. K.; Scuseria, G. E. Seniority and Orbital Symmetry as Tools for Establishing a Full Configuration Interaction Hierarchy. J. Chem. Phys. 2011, 135, 044119.
- <sup>72</sup>Kossoski, F.; Damour, Y.; Loos, P.-F. Hierarchy Configuration Interaction: Combining Seniority Number and Excitation Degree. J. Phys. Chem. Lett. 2022, 13, 4342–4349.
- <sup>73</sup>Garniron, Y.; Scemama, A.; Loos, P.-F.; Caffarel, M. Hybrid stochasticdeterministic calculation of the second-order perturbative contribution of multireference perturbation theory. J. Chem. Phys. 2017, 147, 034101.
- <sup>74</sup>Martin, R. L. Natural transition orbitals. J. Chem. Phys. 2003, 118, 4775– 4777.
- <sup>75</sup>Véril, M.; Scemama, A.; Caffarel, M.; Lipparini, F.; Boggio-Pasqua, M.; Jacquemin, D.; Loos, P.-F. QUESTDB: a database of highly-accurate excitation energies for the electronic structure community. *WIREs Comput. Mol. Sci.* 2021, e1517.
- <sup>76</sup>Loos, P. F.; Scemama, A.; Blondel, A.; Garniron, Y.; Caffarel, M.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly Accurate Reference Energies and Benchmarks. J. Chem. Theory Comput. 2018, 14, 4360–4379.
- <sup>77</sup>Loos, P.-F.; Matthews, D. A.; Lipparini, F.; Jacquemin, D. How accurate are EOM-CC4 vertical excitation energies? J. Chem. Phys. **2021**, 154, 221103.
- <sup>78</sup>Loos, P. F.; Lipparini, F.; Boggio-Pasqua, M.; Scemama, A.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly Accurate Energies and Benchmarks for Medium Sized Molecules. *J. Chem. Theory Comput.* **2020**, *16*, 1711–1741.
- <sup>79</sup>Loos, P.-F.; Lipparini, F.; Matthews, D. A.; Blondel, A.; Jacquemin, D. A Mountaineering Strategy to Excited States: Revising Reference Values with EOM-CC4. J. Chem. Theory Comput. **2022**, 18, 4418–4427.

- <sup>80</sup>Loos, P.-F.; Scemama, A.; Boggio-Pasqua, M.; Jacquemin, D. Mountaineering Strategy to Excited States: Highly Accurate Energies and Benchmarks for Exotic Molecules and Radicals. *J. Chem. Theory Comput.* **2020**, *16*, 3720–3736.
- <sup>81</sup>Del Bene, J. E.; Ditchfield, R.; Pople, J. A. Self-Consistent Molecular Orbital Methods. X. Molecular Orbital Studies of Excited States with Minimal and Extended Basis Sets. J. Chem. Phys. **1971**, 55, 2236.
- <sup>82</sup>Head-Gordon, M.; Rico, R. J.; Oumi, M.; Lee, T. J. A doubles correction to electronic excited states from configuration interaction in the space of single substitutions. *Chem. Phys. Lett.* **1994**, *219*, 21–29.
- <sup>83</sup>Head-Gordon, M.; Maurice, D.; Oumi, M. A perturbative correction to restricted open shell configuration interaction with single substitutions for excited states of radicals. *Chem. Phys. Lett.* **1995**, *246*, 114–121.
- <sup>84</sup>Purvis III, G. P.; Bartlett, R. J. A Full Coupled-Cluster Singles and Doubles Model: The Inclusion of Disconnected Triples. *J. Chem. Phys.* **1982**, *76*, 1910–1918.
- <sup>85</sup>Scuseria, G. E.; Scheiner, A. C.; Lee, T. J.; Rice, J. E.; Schaefer, H. F. The Closed-shell Coupled Cluster Single and Double Excitation (CCSD) Model for the Description of Electron Correlation. A Comparison with Configuration Interaction (CISD) Results. J. Chem. Phys. **1987**, 86, 2881– 2890.
- <sup>86</sup>Koch, H.; Jensen, H. J. A.; Jorgensen, P.; Helgaker, T. Excitation Energies from the Coupled Cluster Singles and Doubles Linear Response Function (CCSDLR). Applications to Be, CH<sup>+</sup>, CO, and H<sub>2</sub>O. *J. Chem. Phys.* **1990**, *93*, 3345–3350.
- <sup>87</sup>Christiansen, O.; Koch, H.; Jørgensen, P. The Second-Order Approximate Coupled Cluster Singles and Doubles Model CC2. *Chem. Phys. Lett.* **1995**, 243, 409–418.
- <sup>88</sup>Hättig, C.; Weigend, F. CC2 excitation energy calculations on large molecules using the resolution of the identity approximation. *J. Chem. Phys.* **2000**, *113*, 5154–5161.
- <sup>89</sup>Burton, H. G. A.; Wales, D. J. Energy Landscapes for Electronic Structure. J. Chem. Theory Comput. **2021**, 17, 151–169.
- <sup>90</sup>We have discarded 10 out of 294 problematic cases when evaluating the statistics of the ΔCISD+EN2 results.
- <sup>91</sup>For the  $\Delta$ CISD+Q statistics, we discarded 12 out of 294 data points where ||c|| < 0.9, where *c* gathers the coefficients of the reference determinants in the CI expansion.
- <sup>92</sup>Loos, P.-F.; Jacquemin, D. A Mountaineering Strategy to Excited States: Highly Accurate Energies and Benchmarks for Bicyclic Systems. *The J. Phys. Chem. A* **2021**, *125*, 10174–10188.
- <sup>93</sup>Serrano-Andrés, L.; Merchán, M.; Nebot-Gil, I.; Lindh, R.; Roos, B. O. Towards an accurate molecular orbital theory for excited states: Ethene, butadiene, and hexatriene. J. Chem. Phys. **1993**, *98*, 3151–3162.
- <sup>94</sup>Starcke, J. H.; Wormit, M.; Schirmer, J.; Dreuw, A. How much double excitation character do the lowest excited states of linear polyenes have? *Chem. Phys.* **2006**, *329*, 39–49, Electron Correlation and Multimode Dynamics in Molecules.
- <sup>95</sup>Barca, G. M.; Gilbert, A. T.; Gill, P. M. Excitation Number: Characterizing Multiply Excited States. J. Chem. Theory Comput. 2018, 14, 9–13.
- <sup>96</sup>Watson, M. A.; Chan, G. K.-L. Excited States of Butadiene to Chemical Accuracy: Reconciling Theory and Experiment. J. Chem. Theory Comput. 2012, 8, 4013–4018.
- <sup>97</sup>Hollauer, E.; Nascimento, M. A. C. A CASSCF description of the nπ<sup>\*</sup> singlet and triplet electronic excited states of the trans-glyoxal molecule. *Chem. Phys. Lett.* **1991**, *181*, 463–466.
- <sup>98</sup>Piecuch, P.; Paldus, J. Orthogonally spin-adapted multi-reference Hilbert space coupled-cluster formalism: diagrammatic formulation. *Theoret. Chim. Acta* 1992, 83.
- <sup>99</sup>Piecuch, P.; Paldus, J. Orthogonally spin-adapted state-universal coupledcluster formalism: Implementation of the complete two-reference theory including cubic and quartic coupling terms. J. Chem. Phys. **1994**, 101, 5875–5890.
- <sup>100</sup>Lutz, J. J.; Nooijen, M.; Perera, A.; Bartlett, R. J. Reference dependence of the two-determinant coupled-cluster method for triplet and open-shell singlet states of biradical molecules. *J. Chem. Phys.* **2018**, *148*, 164102.
- <sup>101</sup>Burton, H. G. A. Generalized nonorthogonal matrix elements: Unifying Wick's theorem and the Slater–Condon rules. J. Chem. Phys. 2021, 154, 144109.

<sup>102</sup>Burton, H. G. A. Generalized nonorthogonal matrix elements for arbitrary excitations. 2022; https://arxiv.org/abs/2208.10208.