



HAL
open science

Hormones and Vestibular Disorders: The Quest for Biomarkers

Rhizlane El Khiati, Brahim Tighilet, Stephane Besnard, Christian Chabbert

► **To cite this version:**

Rhizlane El Khiati, Brahim Tighilet, Stephane Besnard, Christian Chabbert. Hormones and Vestibular Disorders: The Quest for Biomarkers. *Brain Sciences*, 2022, 12 (5), pp.592. 10.3390/brain-sci12050592. hal-03843271

HAL Id: hal-03843271

<https://hal.science/hal-03843271v1>

Submitted on 28 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Hormones and Vestibular Disorders: The Quest for Biomarkers

Rhizlane El Khiati ^{1,2}, Brahim Tighilet ^{1,2}, Stephane Besnard ^{1,2} and Christian Chabbert ^{1,2,*} 

- ¹ Team Pathophysiology and Therapy of Vestibular Disorders, Laboratory of Cognitive Neurosciences, UMR7291, Aix Marseille University-CNRS, CEDEX 07, 13007 Marseille, France; ghizlaineelkhiati@gmail.com (R.E.K.); brahim.thigilet@univ-amu.fr (B.T.); stephane.besnard@unicaen.fr (S.B.)
- ² Research Group on Vestibular Pathophysiology, Unit GDR2074 CNRS, Aix Marseille University-CNRS, CEDEX 07, 13284 Marseille, France
- * Correspondence: christian.chabbert@univ-amu.fr

Abstract: The vestibular system exerts control over various functions through neural pathways that are not yet fully mapped. Functional dysregulations or tissue lesions at different levels of the peripheral and the central vestibular networks can alter these different functions, causing a wide variety of symptoms, ranging from posturo-locomotor alterations to psychiatric syndromes such as PPPD, including the deregulation of the main biological functions. These different symptoms differ by their expression kinetics (they each appear and regress with their own kinetics) by the targets affected (muscles, organs, and brain areas) and by the sensitivity specific to each individual. Vestibular pathologies thus cover a mosaic of distinct effects, and they involve various effectors—which constitute the many markers of their different types and stages. It is therefore crucial, to predict the onset of a vertigo syndrome, to follow its temporal course, or to monitor the impact of therapeutic approaches, and to have specific and reliable biomarkers. Hormonal variations are among the possible sources of biomarkers for neurotology. We know that specific hormonal profiles can promote the appearance of vestibular disorders. We also know that the expression of vertigo syndrome is accompanied by measurable hormonal variations. The link between endocrine deregulation and vestibular alterations therefore no longer needs to be proven. However, there are still few data on their precise correlations with the vertigo syndrome. This study was undertaken with the aim to deliver an extensive review of the hormonal alterations linked to vestibular disorders. A review of the literature covering the last two decades was carried out using the MEDLINE and COCHRANE databases in order to identify studies associating the terms vestibular system or vestibular pathologies and hormones. Bibliographic data provides several outcomes in terms of therapeutic innovation in the diagnosis and therapeutic follow-up of vestibular pathologies.



Citation: El Khiati, R.; Tighilet, B.; Besnard, S.; Chabbert, C. Hormones and Vestibular Disorders: The Quest for Biomarkers. *Brain Sci.* **2022**, *12*, 592. <https://doi.org/10.3390/brainsci12050592>

Academic Editors: Paola Perin, Alessandra Fioretti, Luigi Califano and Elouise Koops

Received: 21 March 2022

Accepted: 29 April 2022

Published: 2 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: biomarkers; dizziness; hormones; Ménière's disease; vestibular neuritis

1. Vestibular Impairment: A Mosaic of Functional Alterations and as Many Potential Biomarkers

The vestibular system impacts different functions through different anatomical pathways (Figure 1). The vestibulo-spinal pathway, the first listed following the work of Flourens on the pigeon in 1830, ensures through descending bundles of motoneurons the control of the trunk and the leg muscles, allowing real time reactions to any disturbance of our equilibrium through adapted postural and locomotor responses to avoid falling. Thirty years later, Vulpian was the first to use the word nystagmus, soon followed by neurologists who established the existence of the vestibulo-ocular reflex. This reflex through the continuous control of the oculomotor muscles when moving the head, makes it possible to fix the gaze on a chosen target (an interlocutor, the sidewalk on which we walk, the staircase which we descend). Barany used it as a basis for the establishment of the caloric test of the vestibule function (rewarded with a Nobel Prize in 1914) [1].




Figure 1. Biomarkers at the different levels of vestibular sensory information integration. VSP: Vestibulo Spinal Pathway; VOP: Vestibulo Ocular Pathway. SVN: Superior Vestibular Nuclei; MVN: Medial Vestibular Nuclei; LVN: Lateral Vestibular Nuclei; IVN: Inferior Vestibular Nuclei: Courtesy Dr S Besnard.

It is only from the 1990s that we started to obtain anatomical and functional evidence of pathways originating from the brainstem vestibular nuclei and projecting towards the bulbar centers and the hypothalamic–pituitary complex. Yates is one of the pioneers who documented the functional pathways between vegetative brainstem centers and vestibular nuclei neurovegetative regulation [2,3]. Thanks to their work on animal models placed in modified gravity, Fuller and Besnard paved the way for the demonstration of vestibular control over the regulation of chronobiological functions, such as sleep, thermoregulation, food intake, and bone renewal [4–9].

We had to wait until 2010 to see an accumulation of experimental confirmations of a direct link between vestibule and cognition, with experimental validation of the role of the vestibule in memory and spatial navigation as well as in bodily self-awareness [10,11]. Finally, the vestibulo–cortical pathway currently deepened with anatomical and functional evidence of a strong link between the vestibule and emotions, with the recent inclusion in the International Classification of Vestibular Disorders (ICVD) of a new syndrome, PPPD, combining postural and perceptual deficits and psychiatric alterations [12].

Functional dysregulations or tissue lesions on the peripheral and the central vestibular networks impact these different functions and result in a wide variety of symptoms ranging from posturo-locomotor alterations to psychiatric syndromes such as PPPD (see other articles in this special issue). These different symptoms differ in their kinetics of expression (they each appear and regress with their own kinetics) in the organs they affect (muscles, organs, and areas of the brain) and sensitivity specific to each individual. These different symptoms thus cover a mosaic of effects and effectors which constitute as many biomarkers, characterizing the different types and stages of vestibular pathologies. It is therefore crucial to predict the onset of a vertigo syndrome, to follow its temporal course, or to monitor the impact of therapeutic approaches, and to have specific and reliable biomarkers.

2. Vestibular Disorders and Hormonal Dysregulation: An Obvious but Still Poorly Documented Link

A number of observations reveal clear links between the endocrine and the vestibular systems. Biological or histological analyses carried out in mammals have shown that different hormones are present in vestibular tissues. Hormones such as aldosterone, testosterone,

progesterone, or estradiol have been found in the epithelial cells of the endolymphatic sac in mammals [13–15]. Vasopressin, through its action on transepithelial exchanges between perilymph and endolymph, is believed to be a key player in the generation of hydrops associated with Ménière's disease [16–20]. It has also been suggested as an inductor of motion sickness [21–23]. Furthermore, stimulation of the vestibular pathways has been shown to activate the central secretory nuclei, which in turn triggers endocrine variations [24–26]. Thus, stimulation of the vestibular pathways induces responses from the paraventricular nucleus (PVN) and the supraoptic nucleus [22,24,26,27]. The latter secrete vasopressin. Thyrotropin-releasing hormone (TRH) neurons in the paraventricular hypothalamic nuclei (PVN) also play a critical role in the function of the hypothalamic–pituitary–thyroid axis, as they stimulate TSH secretion [26]. Steroids are known to be directly involved in both the facilitative and deleterious effects of stress on vestibular compensation [28]. Vestibular stimulation also activates the sympathetic system [29] and the hypothalamic–pituitary axis [30], with modulation of CRH, ACTH, LHRH, and LH secretions [31]. The vestibular system sends projections to the suprachiasmatic nuclei (SCN) and the raphe [32] involved in the secretion of melatonin [26,33] and serotonin [34] as well as a projection bidirectional with orexinergic neurons [35,36]. The vestibular organs receive feedback from melatonin neurons [37]. Furthermore, receptors for different hormones such as vasopressin, cortisol, adrenaline, insulin, or sex hormones (testosterone, progesterone, and estrogen) [14,16,38–42] are expressed in the inner ear sensory organs and all along the nervous tract that projects to the brainstem vestibular nuclei [40]. The conditions for a modulating action of these hormones on the vestibular sensors and at the level of the vestibular sensory information integration zones are thus met. However, the molecular mechanisms supporting these actions, as well as their consequences on the vestibular function remain to be determined.

In order to determine whether the different phases of the vertigo syndrome are correlated with variations in the blood level of hormones, several clinical teams have compared blood samples taken at the time of the acute attack of peripheral vestibulopathies such as Ménière's disease or BPPV and at a distance from them. These studies have demonstrated that vasopressin [31,43], cortisol [44,45], or parathormone [46] levels are significantly increased at the time of an acute vertigo attack in patients with Ménière's disease and BPPV, respectively. These first data suggest that blood variations in circulating hormones could represent biomarkers of the different phases of the vertigo syndrome, with the potential opportunity to discriminate between the different types of vestibular pathologies.

Several epidemiological studies reveal that certain physiological states displaying particular hormonal profiles are especially exposed to vestibular disorders. This is the case in diabetes [47–53] and in thyroid pathologies [54–56]. Benign paroxysmal positional vertigo (BPPV) is predominant in postmenopausal women [57–67], while vestibular migraines may be linked to hormonal alterations [68,69]. If we can multiply the examples of comorbidity between specific hormonal profiles and vestibular disorders, the precise role of circulating hormones as triggers, accompanying factors, or reactive mechanisms to vestibular disorders remains to be determined. Together, these structural, functional, and epidemiological observations suggest a clear link between the hormonal and the vestibular spheres.

3. Hormonal Dysregulations as Biomarkers of the Different Types and Stages of Vestibular Disorders

The principle of a biomarker is to constitute a specific signature of a particular physiological or pathological state. To be considered as a biomarker, a single hormonal variation, or coordinated hormonal variations must occur only at a precise stage, or for a specific type of vestibular pathology. Such an event must be reproduced identically in the same individual, when the same physiological or pathological conditions are repeated; and, of course, it must be interindividually transposable. Taking the specific characteristics of each type of vestibular disorder, such as the etiology, neural circuits involved, organs and modulated cerebral zones, and time course, it is possible to conceive that a pattern of hormonal variations could represent a reproducible signature comparable to a biomarker.

By definition, a biomarker must be measurable, quantifiable in a reliable way, and with standards in terms of methodology and norms. Hormone assays are routinely used in our disease areas to monitor the blood concentration of sex, thyroid, and adrenal hormones. Most biomarkers routinely measured in clinics for diagnosis or follow-up come from blood samples. Chemoluminescence immunoassay techniques are generally used as they are suitable for routine use on analyzers. Some hormones such as estradiol can still be measured by radioimmunology because they are more sensitive, especially in children. Hormone dosage is not a common practice.

4. Conclusions

Hormone dosage is not a common practice in the current diagnosis of vestibular patients. However, it could become a standard of care whether suitable protocols are established according to individual particularities, conditions of access to the patient, and the assay methods available in hospitals. Multicenter, double-blind clinical studies are now urgently needed to identify the different types of hormones, whose plasma or salivary levels may be altered during the different types of vestibular pathologies. Establishing controlled sampling and dosage protocols will open up new avenues for the development of new diagnostic tools and the therapeutic monitoring of vestibular pathologies.

Author Contributions: Conceptualization, R.E.K., S.B. and C.C.; methodology, R.E.K. and C.C.; data curation, R.E.K.; writing—original draft preparation, R.E.K. and C.C.; writing—review and editing, R.E.K., C.C., B.T. and S.B.; supervision, S.B. and C.C. All authors have read and agreed to the published version of the manuscript.

Funding: REK was laureate of the 2021 award of the Research Group on Vestibular Pathophysiology.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lopez, C.; Blanke, O. Nobel Prize centenary: Robert Bárány and the vestibular system. *Curr. Biol.* **2014**, *24*, R1026–R1028. [[CrossRef](#)]
2. Yates, B.J.; Bronstein, A.M. The effects of vestibular system lesions on autonomic regulation: Observations, mechanisms, and clinical implications. *J. Vestib. Res.* **2005**, *15*, 119–129. [[CrossRef](#)] [[PubMed](#)]
3. Yates, B.J.; Bolton, P.S.; Macefield, V.G. Vestibulo-Sympathetic Responses. *Compr. Physiol.* **2014**, *4*, 851–887. [[CrossRef](#)] [[PubMed](#)]
4. Fuller, P.M.; Jones, T.A.; Jones, S.M.; Fuller, C.A. Neurovestibular modulation of circadian and homeostatic regulation: Vestibulo-hypothalamic connection? *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 15723–15728. [[CrossRef](#)]
5. Martin, T.; Mauvieux, B.; Bulla, J.; Quarck, G.; Davenne, D.; Denise, P.; Philoxène, B.; Besnard, S. Vestibular loss disrupts daily rhythm in rats. *J. Appl. Physiol.* **2015**, *118*, 310–318. [[CrossRef](#)] [[PubMed](#)]
6. Kompotis, K.; Hubbard, J.; Emmenegger, Y.; Perrault, A.; Mühlethaler, M.; Schwartz, S.; Bayer, L.; Franken, P. Rocking Promotes Sleep in Mice through Rhythmic Stimulation of the Vestibular System. *Curr. Biol.* **2019**, *29*, 392–401.e4. [[CrossRef](#)] [[PubMed](#)]
7. Besnard, S.; Tighilet, B.; Chabbert, C.; Hitier, M.; Toulouse, J.; Le Gall, A.; Machado, M.-L.; Smith, P.F. The balance of sleep: Role of the vestibular sensory system. *Sleep Med. Rev.* **2018**, *42*, 220–228. [[CrossRef](#)]
8. Martin, T.; Dauvilliers, Y.; Koumar, O.-C.; Bouet, V.; Freret, T.; Besnard, S.; Dauphin, F.; Bessot, N. Dual orexin receptor antagonist induces changes in core body temperature in rats after exercise. *Sci. Rep.* **2019**, *9*, 18432, Erratum in. *Sci Rep.* **2020**, *10*, 6030. [[CrossRef](#)]
9. Van Sluijs, R.; Wilhelm, E.; Rondei, Q.; Omlin, X.; Crivelli, F.; Straumann, D.; Jäger, L.; Riener, R.; Achermann, P. Gentle rocking movements during sleep in the elderly. *J. Sleep Res.* **2020**, *29*, e12989. [[CrossRef](#)]
10. Ehitier, M.; Ebesnard, S.; Smith, P.F. Vestibular pathways involved in cognition. *Front. Integr. Neurosci.* **2014**, *8*, 59. [[CrossRef](#)]
11. Lopez, C. The vestibular system: Balancing more than just the body. *Curr. Opin. Neurol.* **2016**, *29*, 74–83. [[CrossRef](#)]
12. Staab, J.P.; Eckhardt-Henn, A.; Horii, A.; Jacob, R.; Strupp, M.; Brandt, T.; Bronstein, A. Diagnostic criteria for persistent postural-perceptual dizziness (PPPD): Consensus document of the committee for the Classification of Vestibular Disorders of the Bárány Society. *J. Vestib. Res.* **2017**, *27*, 191–208. [[CrossRef](#)]
13. Kumagami, H. Sex Hormones in the Kidney and Endolymphatic Sac: An Immunohistological Study. *Acta OtoLaryngol.* **1994**, *114*, 48–51. [[CrossRef](#)]

14. Grassi, S.; Frondaroli, A.; Di Mauro, M.; Pettorossi, V. Influence of testosterone on synaptic transmission in the rat medial vestibular nuclei: Estrogenic and androgenic effects. *J. Neurosci.* **2010**, *171*, 666–676. [[CrossRef](#)]
15. Motohashi, R.; Takumida, M.; Shimizu, A.; Konomi, U.; Fujita, K.; Hirakawa, K.; Suzuki, M.; Anniko, M. Effects of age and sex on the expression of estrogen receptor α and beta; in the mouse inner ear. *Acta OtoLaryngol.* **2010**, *130*, 204–214. [[CrossRef](#)] [[PubMed](#)]
16. Horii, A.; Koike, K.; Uno, A.; Uno, Y.; Kubo, T. Vestibular modulation of plasma vasopressin levels in rats. *Brain Res.* **2001**, *914*, 179–184. [[CrossRef](#)]
17. Horii, A.; Kitahara, T.; Uno, A.; Kondoh, K.; Morihana, T.; Okumura, S.; Nakagawa, A.; Mitani, K.; Masumura, C.; Kubo, T. Vestibular function and vasopressin. *Acta Oto-Laryngol.* **2004**, *124*, 50–53. [[CrossRef](#)]
18. Takeda, T.; Takeda, S.; Kitano, H.; Okada, T.; Kakigi, A. Endolymphatic hydrops induced by chronic administration of vasopressin. *Hearth Res.* **2000**, *140*, 1–6. [[CrossRef](#)]
19. Katagiri, Y.; Takumida, M.; Hirakawa, K.; Anniko, M. Long-term administration of vasopressin can cause Ménière's disease in mice. *Acta OtoLaryngol.* **2014**, *134*, 990–1004. [[CrossRef](#)] [[PubMed](#)]
20. Kim, M.; Kim, K.S. Vestibular Function Change in a Vasopressin-Induced Hydrops Model. *Otol. Neurotol.* **2017**, *38*, e495–e500. [[CrossRef](#)] [[PubMed](#)]
21. Li, X.; Jiang, Z.-L.; Wang, G.-H.; Fan, J.-W. Plasma Vasopressin, an Etiologic Factor of Motion Sickness in Rat and Human? *J. Neuroendocr.* **2005**, *81*, 351–359. [[CrossRef](#)]
22. Asmar, M.-H.; Gaboury, L.; Saliba, I. Ménière's Disease Pathophysiology: Endolymphatic Sac Immunohistochemical Study of Aquaporin-2, V2R Vasopressin Receptor, NKCC2, and TRPV4. *Otolaryngol. Neck Surg.* **2018**, *158*, 721–728. [[CrossRef](#)] [[PubMed](#)]
23. Xu, L.-H.; Yang, Y.; Liu, H.-X.; Xiao, S.-F.; Qiu, W.-X.; Wang, J.-X.; Zhao, C.-C.; Gui, Y.-H.; Liu, G.-Z.; Peng, B.; et al. Inner Ear Arginine Vasopressin-Vasopressin Receptor 2-Aquaporin 2 Signaling Pathway Is Involved in the Induction of Motion Sickness. *J. Pharmacol. Exp. Ther.* **2020**, *373*, 248–260. [[CrossRef](#)]
24. Liu, F.; Inokuchi, A.; Komiyama, S. Neuronal responses to vestibular stimulation in the guinea pig hypothalamic paraventricular nucleus. *Eur. Arch. OtoRhinoLaryngol.* **1997**, *254*, 95–100. [[CrossRef](#)] [[PubMed](#)]
25. Markia, B.; Kovács-Palkovits, M. Projections from the vestibular nuclei to the hypothalamic paraventricular nucleus: Morphological evidence for the existence of a vestibular stress pathway in the rat brain. *Brain Struct. Func.* **2008**, *213*, 239–245. [[CrossRef](#)]
26. Sailesh, K.S.; Mukkadan, J.K. Vestibular modulation of endocrine secretions—A review. *Intern. J. Res. Health Sci.* **2014**, *2*, 68–78.
27. Palkovits, M.; De Jong, W.; De Wied, D. Hypothalamic Control of Aldosterone Production in Sodium-Deficient Rats. *Neuroendocrinology* **1974**, *14*, 297–309. [[CrossRef](#)]
28. Tighilet, B.; Manrique, C.; Lacour, M. Stress axis plasticity during vestibular compensation in the adult cat. *J. Neurosci.* **2009**, *160*, 716–730. [[CrossRef](#)]
29. Ray, C.A.; Monahan, K.D. Vestibulo sympathetic reflex in humans neural interactions between cardio vascular reflexes. *Clin. Exp. Pharmacol. Physiol.* **2002**, *29*, 98–102. [[CrossRef](#)]
30. Samuel, R.; Denmeade, M.D.; John, T.; Isaacs, P.D. Overview of Regulation of Systemic Androgen Levels. In *Holland-Frei Cancer Medicine*, 6th ed.; BC Decker: Hamilton, ON, USA, 2014.
31. Aoki, M.; Asai, M.; Nishihori, T.; Mizuta, K.; Ito, Y.; Ando, K. The Relevance of an Elevation in the Plasma Vasopressin Levels to the Pathogenesis of Meniere's Attack. *J. Neuroendocr.* **2007**, *19*, 901–906. [[CrossRef](#)]
32. Cavdar, S.; Onat, F.; Aker, R.; Sehirli, U.; San, T.; Yananli, H.R. The afferent connections of the posterior hypothalamic nucleus in the rat using horseradish peroxidase. *J. Anat.* **2001**, *198*, 463e72. [[CrossRef](#)]
33. Dubocovich, M.L. Melatonin receptors: Role on sleep and circadian rhythm regulation. *Sleep Med.* **2007**, *8* (Suppl. 3), 34–42. [[CrossRef](#)]
34. Mickle, W.A.; Ades, H.W. Rostral Projection Pathway of the Vestibular System. *Am. J. Physiol. Content* **1954**, *176*, 243–246. [[CrossRef](#)]
35. Zhang, J.; Li, B.; Yu, L.; He, Y.-C.; Li, H.-Z.; Zhu, J.-N.; Wang, J.-J. A Role for Orexin in Central Vestibular Motor Control. *J. Neuron* **2011**, *69*, 793–804. [[CrossRef](#)]
36. Yu, L.; Zhang, X.-Y.; Chen, Z.-P.; Zhuang, Q.-X.; Zhu, J.-N.; Wang, J.-J. Orexin excites rat inferior vestibular nuclear neurons via co-activation of OX1 and OX2 receptors. *J. Neural Transm.* **2014**, *122*, 747–755. [[CrossRef](#)] [[PubMed](#)]
37. Takumida, M.; Anniko, M. Localization of melatonin and its receptors (melatonin 1a and 1b receptors) in the mouse inner ear. *Acta OtoLaryngol.* **2019**, *139*, 948–952. [[CrossRef](#)]
38. Wangemann, P.; Liu, J.; Shimozone, M.; Scofield, M.A. Beta1-adrenergic receptors but not beta2-adrenergic or vasopressin receptors regulate K⁺ secretion in vestibular dark cells of the inner ear. *J. Membr. Biol.* **1999**, *170*, 67–77. [[CrossRef](#)] [[PubMed](#)]
39. Stenberg, A.E.; Wang, H.; Sahlin, L.; Hultcrantz, M. Mapping of estrogen receptors in the inner ear of mouse and rat. *Hear. Res.* **1999**, *136*, 29–34. [[CrossRef](#)]
40. Seemungal, B.M.; Gresty, M.A.; Bronstein, A.M. The endocrine system, vertigo and balance. *Curr. Opin. Neurol.* **2001**, *14*, 27–34. [[CrossRef](#)]
41. Patel, A.K.; Alexander, T.H.; Andalibi, A.; Ryan, A.F.; Doherty, J.K. Vestibular schwannoma quantitative polymerase chain reaction expression of estrogen and progesterone receptors. *Laryngoscope* **2008**, *118*, 1458–1463. [[CrossRef](#)]
42. Kawao, N.; Takafuji, Y.; Ishida, M.; Okumoto, K.; Morita, H.; Muratani, M.; Kaji, H. Roles of the vestibular system in obesity and impaired glucose metabolism in high-fat diet-fed mice. *PLoS ONE* **2020**, *15*, e0228685. [[CrossRef](#)]

43. Aoki, M.; Hayashi, H.; Kuze, B.; Mizuta, K.; Ito, Y. The association of the plasma vasopressin level during attacks with a prognosis of Meniere's disease. *Intern. J. Audiol.* **2010**, *49*, 1–6. [[CrossRef](#)]
44. Dagilas, A.; Kimiskidis, V.; Aggelopoulou, M.; Kapaki, E.; Fiteli, C.; Libitaki, G.; Papagiannopoulos, S.; Kazis, D.; Kazis, A.; Aidonis, A. Changes in Blood Neurotransmitter and Steroid Levels During Evoked Vertigo. *Otol. Neurotol.* **2005**, *26*, 476–480. [[CrossRef](#)]
45. Cozma, S.; Ghiciuc, C.M.; Damian, L.; Pasquali, V.; Saponaro, A.; Lupusoru, E.C.; Patacchioli, F.R.; Dima-Cozma, L.C. Distinct activation of the sympathetic adreno-medullary system and hypothalamus-pituitary-adrenal axis following the caloric vestibular test in healthy subjects. *PLoS ONE* **2018**, *13*, e0193963. [[CrossRef](#)]
46. Kahraman, S.S.; Ozcan, O.; Arli, C.; Ustun, I.; Erduran, R.; Akoglu, E.; Gokce, C. Calcium Homeostasis During Attack and Remission in Patients with Idiopathic Benign Paroxysmal Positional Vertigo. *Otol. Neurotol.* **2016**, *37*, 1388–1392. [[CrossRef](#)]
47. Rybak, L.P. Metabolic disorders of the vestibular system. *Otolaryngol. Head Neck Surg.* **1995**, *112*, 128–132. [[CrossRef](#)]
48. Degerman, E.; Rauch, U.; Lindberg, S.; Caye-Thomasen, P.; Hultgårdh, A.; Magnusson, M. Expression of insulin signalling components in the sensory epithelium of the human saccule. *Cell Tissue Res.* **2013**, *352*, 469–478. [[CrossRef](#)]
49. Doroszewska, G.; Kaźmierczak, H. Hyperinsulinemia in vertigo, tinnitus and hearing loss. *Otolaryngol. Polska* **2002**, *56*, 57–62.
50. Gawron, W.; Pospiech, L.; Orendorz-Fraczkowska, K.; Noczynska, A. Are there any disturbances in vestibular organ of children and young adults with Type I diabetes? *Diabetologia* **2002**, *45*, 728–734. [[CrossRef](#)]
51. Webster, G.; Sens, P.M.; Salmito, M.C.; Cavalcante, J.D.R.; dos Santos, P.R.B.; da Silva, A.L.M.; de Souza, É.C.F. Hyperinsulinemia and hyperglycemia: Risk factors for recurrence of benign paroxysmal positional vertigo. *Braz. J. Otorhinolaryngol.* **2015**, *81*, 347–351. [[CrossRef](#)] [[PubMed](#)]
52. Bittar, R.S.M.; Santos, M.D.; Mezzalira, R. Glucose metabolism disorders and vestibular manifestations: Evaluation through computerized dynamic posturography. *Braz. J. Otorhinolaryngol.* **2016**, *82*, 372–376. [[CrossRef](#)] [[PubMed](#)]
53. Gioacchini, F.M.; Albera, R.; Re, M.; Scarpa, A.; Cassandro, C.; Cassandro, E. Hyperglycemia and diabetes mellitus are related to vestibular organs dysfunction: Truth or suggestion? A literature review. *Acta Diabetol.* **2018**, *55*, 1201–1207. [[CrossRef](#)]
54. Lin, W.-L.; Chen, C.-Y.; Hsu, T.-Y.; Chen, W.-K.; Lin, C.-L.; Chen, H.-C. Hypothyroidism is an independent risk factor for Menière's disease: A Population-Based Cohort Study. *Medicine* **2019**, *98*, e15166. [[CrossRef](#)] [[PubMed](#)]
55. Miśkiewicz-Orczyk, K.A.; Lisowska, G.; Kajdaniuk, D.; Wojtulek, M. Can Hashimoto's thyroiditis cause vertigo? Czy choroba Hashimoto może być przyczyną zawrotów głowy? *Endokrynol. Polska* **2020**, *71*, 76–86. [[CrossRef](#)]
56. Choi, H.; Song, Y.; Wee, J.; Min, C.; Yoo, D.; Kim, S. Analyses of the Relation Between BPPV and Thyroid Diseases: A Nested Case-Control Study. *Diagnostics* **2021**, *11*, 329. [[CrossRef](#)] [[PubMed](#)]
57. Vibert, D.; Kompis, M.; Häusler, R. Benign Paroxysmal Positional Vertigo in Older Women May be Related to Osteoporosis and Osteopenia. *Ann. Otol. Rhinol. Laryngol.* **2003**, *112*, 885–889. [[CrossRef](#)] [[PubMed](#)]
58. Ogun, O.A.; Büki, B.; Cohn, E.S.; Janky, K.L.; Lundberg, Y.W. Menopause and benign paroxysmal positional vertigo. *Menopause* **2014**, *21*, 886–889. [[CrossRef](#)] [[PubMed](#)]
59. Liu, D.-H.; Kuo, C.-H.; Wang, C.-T.; Chiu, C.-C.; Chen, T.-J.; Hwang, D.-K.; Kao, C.-L. Age-Related Increases in Benign Paroxysmal Positional Vertigo Are Reversed in Women Taking Estrogen Replacement Therapy: A Population-Based Study in Taiwan. *Front. Aging Neurosci.* **2017**, *9*, 404. [[CrossRef](#)] [[PubMed](#)]
60. Wang, S.F.; Zhang, L.; Li, G.H.; Zhang, W.W.; Wang, Y.P.; Geng, B. The change of female progesterone level and blood calcium concentration in perimenopausal women with benign paroxysmal positional vertigo. *Chin. J. Otorhinolaryngol. Head Neck Surg.* **2017**, *52*, 287–290. [[CrossRef](#)]
61. Zhang, Y.M.; Yang, Z.D.; Li, W.X.; Shi, C.; Yu, Y.F. The relationship between the recurrence of benign paroxysmal positional vertigo and the level of bone mineral as well as estrogen in postmenopausal women. *Chin. J. Otorhinolaryngol. Head Neck Surg.* **2017**, *52*, 881–884. [[CrossRef](#)]
62. Yang, L.; Xu, Y.; Zhang, Y.; Vijayakumar, S.; Jones, S.M.; Lundberg, Y.W. Mechanism Underlying the Effects of Estrogen Deficiency on Otoconia. *J. Assoc. Res. Otolaryngol.* **2018**, *19*, 353–362. [[CrossRef](#)] [[PubMed](#)]
63. Chen, Y.G.; Chen, J.; Cao, X.L.; Li, Z.X.; Jiang, H.Q. Relationship between the level of estrogen, calcium and phosphorus concentration in serum with benign paroxysmal positional vertigo. *J. Clin. Otorhinolaryngol. Head Neck Surg.* **2019**, *33*, 497–500. [[CrossRef](#)]
64. Feng, M.Y.; Zhuang, J.H.; Gu, H.H.; Tian, Q.; Zhang, Z.H. Changes of serum E2 and Otolin-1 levels in postmenopausal women with BPPV. *J. Clin. Otorhinolaryngol. Head Neck Surg.* **2019**, *33*, 1138–1141.
65. Feng, M.Y.; Gu, H.H.; Tian, Q.; Yang, H.L.; Zhuang, J.H. Molecular Mediators of Estrogen Reduction-induced Otolith Shedding. *Curr. Med. Sci.* **2021**, *41*, 667–672. [[CrossRef](#)]
66. Guerra, J.; Devesa, J. Causes and treatment of idiopathic benign paroxysmal positional vertigo based on endocrinological and other metabolic factors. *J. Otol.* **2020**, *15*, 155–160. [[CrossRef](#)] [[PubMed](#)]
67. Jeong, S.-H. Benign Paroxysmal Positional Vertigo Risk Factors Unique to Perimenopausal Women. *Front. Neurol.* **2020**, *11*, 589605. [[CrossRef](#)]
68. Park, J.; Viirre, E. Vestibular migraine may be an important cause of dizziness/vertigo in perimenopausal period. *Med. Hypotheses* **2010**, *75*, 409–414. [[CrossRef](#)]
69. Tang, B.; Yu, X.; Jiang, W.; Zhang, C.; Zhan, T.; He, Y. Clinical significance of serum sex hormones in postmenopausal women with vestibular migraine: Potential role of estradiol. *J. Int. Med. Res.* **2021**, *49*, 03000605211016379. [[CrossRef](#)]