



**HAL**  
open science

## Tracking a killer shrimp: *Dikerogammarus villosus* invasion dynamics across Europe

Ismael Soto, Ross N Cuthbert, Danish A Ahmed, Antonín Kouba, Sami Domisch, Jaime R G Marquez, Ayah Beidas, Giuseppe Amatulli, Jens Kiesel, Longzhu Q Shen, et al.

► **To cite this version:**

Ismael Soto, Ross N Cuthbert, Danish A Ahmed, Antonín Kouba, Sami Domisch, et al.. Tracking a killer shrimp: *Dikerogammarus villosus* invasion dynamics across Europe. *Diversity and Distributions*, 2023, 29 (1), pp.157-172. 10.1111/ddi.13649 . hal-03843377

**HAL Id: hal-03843377**

**<https://hal.science/hal-03843377>**

Submitted on 8 Nov 2022

























**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Tracking a killer shrimp: *Dikerogammarus villosus* invasion dynamics across Europe

Ismael Soto<sup>1</sup>  | Ross N. Cuthbert<sup>2,3</sup>  | Danish A. Ahmed<sup>4</sup>  | Antonín Kouba<sup>1</sup>  | Sami Domisch<sup>5</sup>  | Jaime R. G. Marquez<sup>5</sup> | Ayah Beidas<sup>4</sup> | Giuseppe Amatulli<sup>6</sup>  | Jens Kiesel<sup>7,8</sup>  | Longzhu Q. Shen<sup>5,9</sup>  | Margarita Florencio<sup>10,11</sup>  | Herlander Lima<sup>12</sup> | Elizabeta Briski<sup>2</sup>  | Florian Altermatt<sup>13,14</sup>  | Gaït Archambaud-Suard<sup>15</sup>  | Peter Borza<sup>16</sup> | Zoltan Csabal<sup>17,18</sup>  | Thibault Datry<sup>19</sup> | Mathieu Floury<sup>20</sup>  | Maxence Forcellini<sup>19</sup>  | Jean-François Fruget<sup>21</sup> | Patrick Leitner<sup>22</sup>  | Marie-Hélène Lizée<sup>15</sup> | Anthony Maire<sup>23</sup>  | Anthony Ricciardi<sup>24</sup>  | Ralf B. Schäfer<sup>25</sup>  | Rachel Stubbington<sup>26</sup>  | Gea H. Van der Lee<sup>27</sup> | Gábor Várbbíró<sup>28</sup>  | Ralf C. M. Verdonschot<sup>27</sup>  | Peter Haase<sup>8,29</sup>  | Phillip J. Haubrock<sup>1,29</sup> 

<sup>1</sup>Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, University of South Bohemia in České Budějovice, Vodňany, Czech Republic

<sup>2</sup>GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Kiel, Germany

<sup>3</sup>School of Biological Sciences, Queen's University Belfast, Belfast, UK

<sup>4</sup>Center for Applied Mathematics and Bioinformatics (CAMB), Department of Mathematics and Natural Sciences, Gulf University for Science and Technology, Hawally, Kuwait

<sup>5</sup>Department of Community and Ecosystem Ecology, Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

<sup>6</sup>School of the Environment, Yale University, New Haven, Connecticut, USA

<sup>7</sup>Department of Hydrology and Water Resources Management, Institute for Natural Resource Conservation, Christian-Albrechts-University Kiel, Kiel, Germany

<sup>8</sup>Faculty of Biology, University of Duisburg-Essen, Essen, Germany

<sup>9</sup>Institute for Green Science, Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

<sup>10</sup>Inland-Water Ecosystems Team (I-WET), Departamento de Ecología, Edificio de Biología, Facultad de Ciencias, Universidad Autónoma de Madrid, Madrid, Spain

<sup>11</sup>Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid, Madrid, Spain

<sup>12</sup>GloCEE – Global Change Ecology & Evolution Group, Department of Life Sciences, University of Alcalá, Alcalá de Henares, Spain

<sup>13</sup>Department of Aquatic Ecology, Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

<sup>14</sup>Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zürich, Switzerland

<sup>15</sup>INRAE, UMR RECOVER, Aix Marseille Univ., Centre d'Aix-en-Provence, Aix-en-Provence Cedex 5, France

<sup>16</sup>Centre for Ecological Research, Institute of Aquatic Ecology, Budapest, Hungary

<sup>17</sup>Department of Hydrobiology, University of Pécs, Pécs, Hungary

<sup>18</sup>Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic

<sup>19</sup>RiverLY Research Unit, National Research Institute for Agriculture, Food and Environment (INRAE), Villeurbanne, France

<sup>20</sup>UMR 5023 LEHNA, Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, Villeurbanne, France

<sup>21</sup>ARALEP, Ecologie des Eaux Douces, Villeurbanne Cedex, France

<sup>22</sup>Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Life Sciences, Vienna, Austria

<sup>23</sup>EDF R&D, Laboratoire National d'Hydraulique et Environnement (LNHE), Chatou Cedex, France

<sup>24</sup>Redpath Museum and Bieler School of Environment, McGill University, Montreal, Quebec, Canada

<sup>25</sup>Institute for Environmental Sciences, University of Koblenz Landau, Landau, Germany

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Diversity and Distributions* published by John Wiley & Sons Ltd.

<sup>26</sup>School of Science & Technology, Nottingham Trent University, Nottingham, UK

<sup>27</sup>Wageningen Environmental Research, Wageningen University and Research, Wageningen, The Netherlands

<sup>28</sup>Department of Tisza River Research, Centre for Ecological Research, Institute of Aquatic Ecology, Debrecen, Hungary

<sup>29</sup>Department of River Ecology and Conservation, Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany

### Correspondence

Ismael Soto, Faculty of Fisheries and Protection of Waters, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, University of South Bohemia in České Budějovice, Vodňany, Czech Republic.

Email: [isma-sa@hotmail.com](mailto:isma-sa@hotmail.com)

### Funding information

Kuwait Foundation for the Advancement of Sciences (KFAS), Grant/Award Number: PR1914SM-01; Gulf University for Science and Technology (GUST); Leverhulme Trust Early Career Fellowship, Grant/Award Number: ECF-2021-001; EU Horizon 2020, Grant/Award Number: 871128; German Federal Ministry of Education and Research, Grant/Award Number: 033W034A

Editor: April Blakeslee

### Abstract

**Aim:** Invasive alien species are a growing problem worldwide due to their ecological, economic and human health impacts. The “killer shrimp” *Dikerogammarus villosus* is a notorious invasive alien amphipod from the Ponto-Caspian region that has invaded many fresh and brackish waters across Europe. Understandings of large-scale population dynamics of highly impactful invaders such as *D. villosus* are lacking, inhibiting predictions of impact and efficient timing of management strategies. Hence, our aim was to assess trends and dynamics of *D. villosus* as well as its impacts in freshwater rivers and streams.

**Location:** Europe.

**Methods:** We analysed 96 European time series between 1994 and 2019 and identified trends in the relative abundance (i.e. dominance %) of *D. villosus* in invaded time series, as well as a set of site-specific characteristics to identify drivers and determinants of population changes and invasion dynamics using meta-regression modelling. We also looked at the spread over space and time to estimate the invasion speed (km/year) of *D. villosus* in Europe. We investigated the impact of *D. villosus* abundance on recipient community metrics (i.e. abundance, taxa richness, temporal turnover, Shannon diversity and Pielou evenness) using generalized linear models.

**Results:** Population trends varied across the time series. Nevertheless, community dominance of *D. villosus* increased over time across all time series. The frequency of occurrences (used as a proxy for invader spread) was well described by a Pareto distribution, whereby we estimated a lag phase (i.e. the time between introduction and spatial expansion) of approximately 28 years, followed by a gradual increase before new occurrences declined rapidly in the long term. *D. villosus* population change was associated with decreased taxa richness, community turnover and Shannon diversity.

**Main Conclusion:** Our results show that *D. villosus* is well-established in European waters and its abundance significantly alters ecological communities. However, the multidecadal lag phase prior to observed spatial expansion suggests that initial introductions by *D. villosus* are cryptic, thus signalling the need for more effective early detection methods.

### KEYWORDS

biological invasion, crustacean, freshwater ecosystem, invasive alien species, long-term monitoring, time-series

## 1 | INTRODUCTION

Humans have translocated thousands of invasive alien species beyond their native ranges (Seebens et al., 2021). Their establishment and spread have been recognized as a leading cause of biodiversity loss and a growing socio-economic burden worldwide (Bellard et al., 2016; Diagne et al., 2021). It is therefore imperative to understand and predict large-scale invasion patterns, identify future

invaders (i.e. introduced species that become invasive; Fournier et al., 2019; Pyšek et al., 2020) and assess their impacts (Essl et al., 2020). With the ongoing global increase in invasion rates (Seebens et al., 2021) and a growing threat to ecosystems and economies (Pyšek et al., 2020), there is an urgent need to characterize population dynamics at large scales to inform effective detection, management actions and monitoring (Cuthbert, Kotronaki, Carlton, et al., 2022; Seebens et al., 2021).

The form of population dynamics may differ among invaders, invasion pathways, biogeographical regions, abiotic or biotic gradients and spatio-temporal scales (Arim et al., 2006; Haubrock et al., 2020). Characterization of population dynamics of invasive alien species at a broad spatio-temporal scale has been limited by insufficient long-term data. However, advances in availability in long-term biodiversity monitoring (Dornelas et al., 2018; Mirtl et al., 2018) and data analysis approaches for biodiversity time series (Bowler et al., 2017; Dornelas et al., 2014; Pilotto et al., 2020) have been made. These and other initiatives have collated large datasets for diverse taxonomic groups and examined biodiversity and ecosystem function trends at regional and global scales (Seebens et al., 2019, 2020). Large-scale, community-level data can also be used to examine distribution and abundance patterns of invasive alien species, as well as their potential effects on ecological communities over time (Dornelas et al., 2014; Haubrock et al., 2020).

Invaders may take considerable time before becoming established, detected and disruptive (Crooks, 2005; Spear et al., 2021), and such time lags are very difficult to predict (Coutts et al., 2018). Furthermore, the invaders may not remain disruptive or could become less so, owing to population declines (e.g. reflecting boom-and-bust cycles and/or community adjustment; Strayer et al., 2017). Finally, the type and magnitude of ecological impacts an invader causes are dependent, in part, on its abundance (Sofaer et al., 2018; Yokomizo et al., 2009), and thus can change through time according to the invader's population dynamics. Given that the long-term population dynamics of even widespread conspicuous invaders are often poorly characterized (Strayer et al., 2006), this gap challenges our ability to (i) predict in which situations the invader's impacts on invaded ecosystems will be maximal and (ii) decide if/when management should intervene. However, species having well-documented invasion histories within large contiguous regions offer temporally highly resolved data whose collation and analysis could reveal essential information to guide risk assessment and management prioritization.

One such species is the "killer shrimp" *Dikergammarus villosus*, an invasive alien freshwater amphipod listed among the "100 worst" invaders in Europe (Nentwig et al., 2018). The confirmed native range of *D. villosus* is the northern margins of the Ponto-Caspian region (i.e. Black Sea, Caspian Sea and Azov Sea; Dedyu, 1980; Mordukhai-Boltovskoi, 1960). This voracious predator has spread rapidly through European inland waterways and the Baltic Sea, aided by canalization and anthropogenic vectors (Cuthbert et al., 2020; De Ventura et al., 2017), and also poses an invasion risk to the North American Great Lakes (Kramer et al., 2017). Facilitated by the known influence of streams on the spread of *D. villosus*, it has substantial impacts on biodiversity in invaded regions, causing marked declines in native macroinvertebrates via predation and competition (Dick et al., 2002; Dick & Platvoet, 2000). It also potentially impacts egg and embryonic stages of large, ecologically important crustaceans, fish and amphibians (Roje et al., 2021; Taylor & Dunn, 2017; Warren

et al., 2021), causing disruptions across multiple trophic levels (i.e. trophic cascades; Van Riel, Van der Velde, & Bij de Vaate, 2006). Its displacement of native species (MacNeil et al., 2011; MacNeil & Platvoet, 2013) also facilitates the establishment of other invasive alien species (Bollache et al., 2004; Leuven et al., 2009). Notably, it is the only invasive amphipod with documented monetary costs (Kouba et al., 2021). Yet, information on this invader's effects on native species assemblages over time is still scarce.

*Dikergammarus villosus* invasion success has been promoted by its suitability for transport by anthropogenic vectors (e.g. ships; Anderson et al., 2014), wide thermal and salinity tolerances (Cuthbert et al., 2020), aggressive competitive behaviours (Kobak et al., 2016), effective anti-predator strategies (Rolla, Consuegra, & de Leaniz, 2020) and high growth rates and fecundity (Holdich & Pöckl, 2007; Pöckl, 2009), as one female can carry nearly 200 eggs (Pöckl, 2009). Accordingly, this invader's potential to successfully establish after introduction – from even a single gravid female – into a novel environment is substantial (Devin et al., 2004; Lockwood et al., 2005). Reflecting its impacts on recipient ecosystems (e.g. extirpation of native species and changes in biotic indices; Kouba et al., 2021; MacNeil et al., 2013), *D. villosus* has been the focus of various management actions, although measures for management post-establishment are undeveloped, and prevention of initial introduction has been advocated (Bradbeer et al., 2020; Wood et al., 2021). Moreover, understanding the population dynamics of this species remains limited at large spatial scales regarding temporal and climatic gradients (e.g. time since invasion; variability in temperature and precipitation), hindering quantification of impact and population trends that could inform management strategies. For example, earlier invasions, such as those detected in Hungary (Bij de Vaate et al., 2002: in 1926; Huber et al., 2015 in 1970; Figure 1), might already be regressing from their peak abundance to a stable, intermediate level or exhibiting "boom-bust" dynamics (Rolla, Consuegra, Hall, & Garcia de Leaniz, 2020; Strayer et al., 2017) and therefore be expanding to new areas more slowly than populations with a more recent invasion history (Seebens et al., 2018). More recent invasions may be at an early stage of spread, potentially preceding future exponential population increases. In addition, invasion success and exerted impacts are intertwined with anthropogenic stressors such as hydromorphological alterations and dam construction, which can influence spread rates (Colautti et al., 2006; MacNeil & Platvoet, 2013).

To characterize the population dynamics of *D. villosus* across time series, we collated European riverine macroinvertebrate bio-monitoring data containing this species. We hypothesized that the (i) dominance of *D. villosus* populations is increasing within time series, whereas the number of new occurrences is declining in the long-term, indicating a deceleration in the invasion; (ii) recent invasions exhibit more rapid population growth, whereas populations originating from earlier invasions are stable or declining; (iii) population dynamics of *D. villosus* are mediated by site-specific climatic and spatial characteristics, with populations affected by warming and the presence of anthropogenic barriers such as dams. Lastly, we

hypothesized (iv) changes in recipient communities to be associated with *D. villosus* population dynamics.

## 2 | METHODS

### 2.1 | Data compilation

To investigate the population dynamics of *D. villosus* and the response of recipient communities across Europe, we considered 1816 time series (Peter Haase, unpublished data) reporting the abundance of macroinvertebrate taxa in streams and rivers across 22 European countries. Macroinvertebrates were sampled using different methods and protocols (see Table S1) among time series, but were consistent within each time series. Each time series comprised macroinvertebrate assemblages collected at a single site in multiple years. We initially selected all 132 time series with *D. villosus* populations and then excluded 36 time series that contained  $\leq 2$  sampled years, retaining 96 time series (Figure 1), of which most are from large European rivers (e.g. Danube or Rhine) across six European countries (i.e. Austria, France, Germany, Hungary, Netherlands and Switzerland). For time series in which samples were not collected in all years, we coded missing years as “not available” (NA). Time series spanned a mean  $\pm$  SD of  $10.1 \pm 3.4$  years and contained  $7.7 \pm 3.6$  sampling years between 1994 and 2019.

### 2.2 | Trend identification and meta-regression modelling

To synthesize and describe the directionality and the trends in the number of sampled *D. villosus* individuals, we used a meta-regression modelling approach, which synthesizes the slopes of

individual regressions (as in our case from different time series). For this, we used the “rma.mv” function of the R package *metafor* (Viechtbauer, 2010), using the time-series Mann–Kendall trend test (*S*-statistics) and respective variance as the effect size. The Mann–Kendall trend test is a non-parametric test to evaluate a monotonic increase or decrease in trends. In particular, we used the trends calculated by modified Mann–Kendall trend tests with variance correction to account for temporal auto-correlation (Hamed & Rao, 1998; Maire et al., 2019). To correct the spatial autocorrelation between time series, we used a random effect model, specifying the geographic coordinates as a random effect according to a Gaussian correlation structure (Cressie, 1993; Maire et al., 2019). This approach uses a regression model and enables comparability of time series by analysing the individual time series abundance trends (i.e. its slope) rather than the raw abundances. These models account for the variance of each individual temporal trend and treat each population as an individual spatial unit associated with scale and sampling protocol (Viechtbauer et al., 2015). Lastly, based on two data clusters, we classified the time series according to their first year of sampling into two groups and visually inspect if there are differences in population growth among groups. For this, we selected 2003 as a middle point between earlier (i.e. the first year of the time series before 2003) and later (i.e. the first year of the time series after 2003) invasions.

Although raw abundances are not comparable directly across time series due to differences in sampling methods, we approximated the temporal trend in *D. villosus* relative abundance (%) by averaging abundance records across all samples from a given year, as a proxy of dominance of the species to avoid introducing a bias from the comparison of different sampling methods. We considered only those years in which at least 15 assemblages were sampled and thus omitted 1994, 2018 and 2019 (5–7 sampled assemblages). Once removed, the mean number of assemblages sampled was  $27.3 \pm 6.8$ ,




FIGURE 1 Map summarizing the native and invaded range of *Dikerogammarus villosus* in Europe, showing populations reported by the Global Biodiversity Information Facility (GBIF, 2021; grey triangles) and our time series (white triangles). Years within countries indicate the year of the first introduction according to the sTWIST database of first records (Seebens et al., 2018). The invasion pathways by which *D. villosus* has spread (i.e. Rhine, Danube, Volga and Dnieper rivers) are inferred from Bij de Vaate et al. (2002).



and maximum 42 in the year 2007. We tested a linear model for the dominance of *D. villosus* over time using least squares regression and thus estimated the proportion of abundance (%) at the time the assemblage was sampled.

### 2.3 | Effects of site-specific characteristics on *D. villosus*

To investigate the spatial and climatic drivers of *D. villosus* trends, we used site-specific characteristics of each time series. Climatic regions (i.e. Köppen-Geiger climate zone) were extracted from Beck et al. (2018). Biogeographical regions were defined following the European Environment Agency classification map and estimated visually using the site-specific coordinates (EEA, 2021). We further classified the type of ecosystem of each time series based on Strahler order (i.e. stream <8 or large rivers ≥8) to evaluate differences in the degree of invasions between both. We obtained site-specific runoff data, expressed as the annual Q (mm), from the TerraClimate dataset at 4-km spatial resolution (Abatzoglou et al., 2018). We extracted the elevation of each site from the MERIT Hydro digital elevation model (Yamazaki et al., 2019) at 90-m spatial resolution and used the Hydrography90m (Amatulli et al., 2022) stream network, catchments and sub-catchments (catchments between network nodes) as underlying spatial units. For each site, we computed the stream slope using the `r.stream.slope` function. We extracted land cover data from the European Space Agency Climate Change Initiative (ESA CCI) Land Cover time series v2.0.7 dataset, at 300-m spatial resolution (ESA, 2017), as the percentage cover of a given land cover category within the sub-catchment. We used the Global Reservoir and Dam Database (GRanD) v1.3 to identify dams along the river network. We then measured the distance to the nearest dam to investigate the effect of instream barriers. We extracted mean daily temperature and total daily precipitation data from a gridded European scale observation-based dataset (spatial resolution: 0.1°; Cornes et al., 2018) and calculated the average annual temperature and precipitation for each sampled assemblage in each sampling year (Pilotto et al., 2020), as well as their respective *S*-statistics (i.e. the Mann-Kendall trend test statistic, see above), as indicators of climatic changes. Precipitation in particular can be used as a proxy for river discharge (Higashino & Stefan, 2019) and was included as it affects the availability of water and nutrients as well as the habitat suitability for many species (Gallardo et al., 2012; Shi et al., 2010). We then calculated the mean maximum and minimum temperature and precipitation and each respective *S*-statistic. *Dikerogammarus villosus* slopes (to investigate factors determining *D. villosus*' rate of change over time) and relative abundances (as a proxy of *D. villosus*' dominance in invaded ecosystems over time) were analysed as a function of these spatial, temporal, and site-specific characteristics, to identify significant drivers of temporal trends (see Table S2). We used generalized linear models (GLMs) via the MASS R package (Ripley et al., 2013). We used a Gaussian distribution for continuous data and quasibinomial distribution (to account for high variance of

the dataset) for proportion data (i.e. relative abundances) after visually inspecting their respective residual distributions for normality through histograms.

To identify the best model, we first tested for collinearity among the numerical variables using the variance inflation factor (VIF) for continuous predictors using the "corvif" function (Zuur et al., 2009). We selected  $VIF > 5$  as the threshold, and those variables with high VIF values for each model were assessed for their ecological relevance based on expert knowledge (Table S3; Dorman et al., 2013; Zuur et al., 2009). Regarding the four categorical variables (country, biogeographical region, Köppen-Geiger classification and ecosystem type), we used chi-square tests to investigate the collinearity, and retained only the biogeographical region (Table S4). We considered each model and the respective predictors, using expert-based opinion to determine if the inclusion predictors would make sense from an ecological perspective (Table S5). Hence, the model consisted of a single response variable (i.e. the Mann-Kendall trend test slopes of *D. villosus* abundances or the relative proportion of *D. villosus*) and site-specific characteristics (see above; Table S2). Following the alphabetical order, the Alpine region was used as a reference factor (i.e. as intercept), and therefore, we do not infer any results about this region.

### 2.4 | Modelling occurrence frequency and invasion speed

We combined the first occurrence of *D. villosus* in each time series in our data with those occurrences (as coordinates and year of records) in the Global Biodiversity Information Facility database (GBIF, 2021). Eight GBIF occurrences were removed due to insufficient information (e.g. no recorded year), resulting in 400 records.

We excluded those sample years from the dataset of 400 records that reported relatively high occurrence frequencies, that is, any number of occurrences that was greater than  $Q_3 + 3 \times IQR$ , where  $Q_3$  is the upper quartile and IQR is the interquartile range of the dataset. A single outlier was found with 120 occurrences in the year 2009, and thus, it was removed. We modelled the remaining 280 occurrences to represent invader spread, using a logistic distribution and a two-tailed Pareto distribution. A key difference between these distributions is the decay rate at the end tails: the logistic distribution decays exponentially fast (thin tails), and for the Pareto distribution, the decay is much slower according to an inverse power law (fat tails; Nolan, 2020). Moreover, Pareto distributions with distinct parameters were considered (i.e. two-tailed) for the early and late phases of the invasion. Both distributions were fitted against the occurrence data using the non-linear regression tool `lsqcurvefit` in Matlab. The better-fitting model was determined based on lower number of parameters and higher  $R^2$  value (see Note S2). Further, we estimated the frequency of occurrences  $f_0$  at the time of first sampling ( $t = 0$ ) the time of introduction  $t_{intro}$ , and the duration of the lag phase  $t_{lag}$  that is, the period before *D. villosus* was observed in additional assemblages, evaluated at 10% of the largest recorded occurrence frequency  $f^*$ .

As a complementary analysis, we computed the distance between the locations of every sampled site from the first invaded site using site location data (GPS coordinates recorded as latitude and longitude) over the years 1994–2021. An estimate for the invasion speed (km/year) was obtained by computing the mean distance (i.e. total distance averaged over the number of occurrences per year) over time and modelled using a linear equation (Bagnara et al., 2022). We also calculated the differences between the first record of *D. villosus* in our data and sTWIST database (the most comprehensive source of first records of alien species, integrating several databases and merging them into a single database; Seebens et al., 2018).

## 2.5 | Impacts of *D. villosus* on community metrics

To assess the effect of *D. villosus* abundance on recipient communities (proxied by the *S*-statistic of *D. villosus* trends), we computed five common metrics for each community and year within each time series: total abundance (i.e. individuals), taxon richness (i.e. the total number of taxa), temporal turnover (i.e. the proportion of species either gained or lost over time relative to the total number of species observed; Carvalho et al., 2012), the Shannon diversity index (Shannon & Weaver, 1949) and Pielou's evenness (Pielou, 1966). The metrics were calculated considering all species in the community except *D. villosus*, potentially including both native and other non-native species. For evenness, we followed the formula:  $H/\ln(S)$  (where  $H$  is the Shannon Index and  $S$  is the taxon richness of a community). Metrics were calculated using the “diversity” function in the R package *vegan* (Oksanen et al., 2013) and the “turnover” function in the R package *codyn* (Hallett et al., 2016). In analysing these metrics as response variables in meta-regression models, we included the middle point of each time series (see above) in addition to the rate change of *D. villosus* to infer the effects of temporal variability in changing temporal trends (i.e. slopes) over time and evaluate the change of sampled individuals of *D. villosus* individuals to test its associated effect on community metrics.

All models used restricted maximum likelihood estimation (REML). We also quantified the proportion of variance in the model not attributed to sampling error by using the  $I^2$  statistic. In addition, we evaluated the results of the meta-regression by a graphical representation (i.e. forest plots) using the “forest” function of the R package *metafor* (Figure S2; Viechtbauer, 2010). To inspect potential biases that may alter the results, we checked the symmetry of the data using funnel plots and statistically evaluated this symmetry using the Egger's test (Egger et al., 1997; see Figure S3; Table S6).


All analyses were carried out in R v.4.1.3. (R Core Team, 2022). The reproducible R script is available with the manuscript and lists all R packages that were used.

## 3 | RESULTS

### 3.1 | Trends of *D. villosus* across Europe

Across the sampling sites in Europe included in this study, the abundance trend of *D. villosus* increased in 49 locations, decreased in 44 locations and has not changed in 3 locations (Figure 2a). Our analysis therefore suggests that the total number of *D. villosus* individuals in the study region experienced no overall significant change in its rising trajectory between 1994 and 2019s (*S*-statistics = 4.74; CI: -9.32, 18.82,  $p = .50$ ; Table 1; Figure 2a), albeit expressing low heterogeneity ( $I^2 = 3.62\%$ ). In addition, we did not find differences in the population growth between earlier (for which  $t = 0$  was before 2003) and later time series (i.e.  $t = 0$  after 2003; Figure S1).

Averaged across all time series, the overall proportion of *D. villosus* was well described by a linear model ( $r = .45$ ), suggesting a steep rise in relative abundance over time. An average of 8.66% of *D. villosus* was recorded per sampled assemblage at the first time point. On average across all time series, the rate of increase in relative abundance (i.e. dominance) increased by 0.31% per year for each sampled assemblage (Figure 2b). In addition, our first records for *D. villosus* were on average



**FIGURE 2** Changes in trends (slopes) of *Dikerogammarus villosus* in individual time series (*S*-statistics ± confidence intervals): red represents negative trends, blue positive trends, and grey indicates no significant change over time (a). Relative abundance of *D. villosus* in sampled sites at the European level. Proportions were averaged over the number of sampled assemblages each year from 1995 ( $t = 0$ ) to 2017 ( $t = 22$ ; black dots) (b).

10.8 years later than those referenced in the sTWIST database. This difference was reduced to 2.8 years after excluding time series from Austria and Switzerland ( $n = 2$ ), for which the difference between both databases was 14 years (Seebens et al., 2018).

### 3.2 | Effect of site-specific characteristics on *D. villosus* trend

We did not identify a change in individuals of *D. villosus* sampled over time, despite a positive tendency (i.e. change in the  $S$ -statistics; GLM:  $0.59 \pm 0.45$ ;  $p = .19$ ; Figure 3a; Table S7). The rate of change in the trend of *D. villosus* increased significantly across the Mediterranean biogeographic region ( $p < .05$ ). The average minimum temperature and the rate change of the maximum temperature had positive effects on the rate of change of *D. villosus* individuals ( $p < .05$ ), while the distance to the next barrier had negative effects ( $p < .05$ ; Figure 3b; Table S7). Regarding the relative abundance of *D. villosus* over time, we identified a significant increase ( $p < .05$ ; Figure 3c; Table S7). This increase was shown in all biogeographic regions (relative to Alpine), as well as with increasing distance to the next barrier and the elevation of the stream ( $p < .01$ ; Figure 3d; Table S7). The relative abundance decreased in streams relative to large rivers ( $p < .01$ ; Figure 3d; Table S7).

### 3.3 | Modelling occurrence frequency and invasion speed at the European level

The mean  $\pm$  SD for the number of *D. villosus* occurrences was  $13.33 \pm 19.07$ , with the maximum recorded occurrence frequency

$f^* = 78$  in the year 2008 ( $t^* = 14$ ). The occurrence frequency of *D. villosus* was best described by a two-tailed Pareto distribution ( $R^2 = .90$  for  $t \leq t^*$  and  $R^2 = .98$  for  $t \geq t^*$  see Figure 4a), which fits better than the logistic distribution ( $R^2 = .86$ ; see Note S2). Moreover, the Pareto distribution depends on fewer parameters. On considering an occurrence frequency of 1 (i.e. first invaded site), we predicted the time of introduction  $t_{\text{intro}} = -20.67$  years prior to the first sampling event. The estimated number of occurrences at the time of first sampling was  $f_0 = 3.37$ . The duration of the lag phase was 27.64 years after  $t_{\text{intro}}$ . Beyond  $t^*$ , occurrence frequency rapidly declined, reaching low levels 45–50 years after  $t_{\text{intro}}$ , indicating low levels of spread for *D. villosus*.

We estimated the invasion speed of *D. villosus* at the time of first sampling as 80.27 km/year, with deceleration at a rate of 2.83 km/year<sup>2</sup>, eventually reaching minimum speed at time  $t = 28.34$  years (see Figure 4b). The estimation of null speed corresponds to a cessation in the frequency of occurrences at approximately the same time (49.03 years after  $t_{\text{intro}}$ ; Figure 4a).


### 3.4 | Impact of *D. villosus* on community metrics trends

We did not find a significant trend over time for community metrics (Table 1). The rate of change in the number of *D. villosus* individuals sampled over time (i.e. its slope) had a significant negative effect on trends in taxon richness, temporal turnover and Shannon diversity (Figure 5; Table 1). We did not find a significant effect of the number of sampled *D. villosus* individuals on total community abundance and Pielou's evenness trends (Table 1; Figure 5).

TABLE 1 Meta-regression results according to time and *Dikerogammarus villosus* abundance for the following response variables: *D. villosus* abundance (a), community abundance (b), richness (c), turnover (d), diversity (e) and evenness (f) of recipient community.

Response variable	Predictor	Estimate	Standard error	p-Value	Confidence interval (lower)	Confidence interval (upper)
(a) <i>D. villosus</i> abundance $I^2 = 3.62\%$	Intercept	4.74	7.18	.50	-9.32	18.82
(b) Abundance $I^2 = 21.86\%$	Intercept	-15.70	196.46	.93	-400.76	369.34
	Middle point of time series	0.13	0.13	.32	-0.13	0.39
	Change in <i>D. villosus</i> abundance	0.14	0.08	.10	-0.03	0.31
(c) Richness $I^2 = 68.96\%$	Intercept	-327.49	237.33	.16	-792.66	137.67
	Middle point of time series	0.11	0.20	.57	-0.28	0.52
	Change in <i>D. villosus</i> abundance	-0.24	0.09	.01	-0.43	-0.05
(d) Turnover $I^2 = 4.55\%$	Intercept	47.70	187.37	.75	-319.53	414.94
	Middle point of time series	-0.03	0.09	.75	-0.21	0.15
	Change in <i>D. villosus</i> abundance	-0.17	0.08	.03	-0.32	-0.01
(e) Diversity Shannon $I^2 = 21.80\%$	Intercept	74.45	188.24	.69	-294.50	443.41
	Middle point of time series	<0.01	0.15	.95	-0.30	0.32
	Change in <i>D. villosus</i> abundance	-0.22	0.08	.01	-0.39	-0.04
(f) Evenness Pielou $I^2 = 14.42\%$	Intercept	148.23	190.31	.43	-224.78	521.25
	Middle point of time series	-0.03	0.12	.78	-0.27	0.20
	Change in <i>D. villosus</i> abundance	-0.15	0.08	.08	-0.33	0.02






**FIGURE 3** Trend of *Dikerogammarus villosus* abundance over time  $\pm$  standard error (SE, blue shaded area) (a). Effect  $\pm$  SE of the predictors included in the *D. villosus* model (b). Relative abundance of *D. villosus* over time  $\pm$  standard error (SE, blue shaded area) (c). Effect  $\pm$  SE of the predictors included in the model (d). Solid trend lines are significant and dashed lines not significant. Blue dot: positive effect; red dot: negative effect; filled dots: significant effects; empty dots: non-significant effects.

## 4 | DISCUSSION

### 4.1 | Overview

We characterized the population dynamics of one of the most notorious invasive alien species in Europe, *D. villosus*, and its effects on freshwater macroinvertebrate community metrics across available European time series. Contrary to our first hypothesis, we detected no significant trend in the number of *D. villosus* individuals sampled, although its dominance in invaded ecosystems increased over time. Contrary to our second hypothesis, the growth of earlier and more recently invading populations was comparable. Supporting our third hypothesis, *D. villosus* populations were influenced by site-specific climatic and spatial characteristics. Finally, supporting our fourth hypothesis, *D. villosus* was negatively associated with trends in macroinvertebrate community taxon richness, temporal turnover and Shannon diversity. These results highlight the need towards proactive management actions to contain *D. villosus* as well as to better understand the potential synergistic effects of stressors (Ricciardi et al., 2021).

Contrary to our first hypothesis, our meta-regression models identified no trend in the *D. villosus* population across time series. This lack of identifiable patterns could (i) reflect the complex population dynamics of invasive alien species at large spatial scales, (ii) climatic variability across European countries and biogeographical regions or (iii) genetic differentiation across invaded sites, ultimately leading to differing trends (Arim et al., 2006; Haubrock et al., 2022). For example, in France, all temporal trends (i.e. S-Statistics) were positive, whereas most trends were negative in Hungary. The sustained dominance of *D. villosus* in recipient ecosystems increased over time, which could reflect its ability to rapidly reach high population densities in combination with its capacity to predate, eliminate and replace native and alien species (Dick & Platvoet, 2000; Nentwig et al., 2018; Warren et al., 2021). In addition, the difference between the first record of *D. villosus* in sTWIST and our database was  $\sim 10.8$  years. Yet, after excluding Switzerland and Austria—which had a difference of  $\sim 14$  years between both databases likely due to the scarce time series from that country ( $n = 2$ )—the difference among both databases shrank to  $\sim 2.8$  years only, underlining the accuracy of our data.



**FIGURE 4** Occurrence frequency of *D. villosus* between 1994 ( $t = 0$ ) and 2021 ( $t = 27$ ) (circle markers) (a). Pareto distributions were fitted with  $R^2 = .90$  for  $t \leq t^*$ ,  $RMSE = 6.7$  and  $R^2 = .98$  for  $t > t^*$ ,  $RMSE = 3.6$ , where the maximum occurrence frequency  $f^* = 78$  was recorded at  $t^* = 14$  years. Estimated model parameters for the left-hand tail are  $s_1 = 1.81$ ,  $\mu_1 = 1.45$ , which were used to predict the: time of introduction  $t_{\text{intro}} = -20.67$  years, number of occurrences at the time of first sampling ( $t = 0$ )  $f_0 = 3.37$ , duration of the lag phase  $t_{\text{lag}} = 27.64$  years (after  $t_{\text{intro}}$ ) with occurrence frequency  $f_{\text{lag}} = 7.8$  (fixed at 10% of  $f^*$ ). Model parameters for the right-hand tail are  $s_2 = 1.08 \times 10^4$ ,  $\mu_2 = 7.51 \times 10^3$ , see Note S2 for mathematical details related to the Pareto distributions, and how these key points were determined from estimated model parameters. Linear model  $v = -2.83t + 80.27$  fitted using least squares regression against estimated values of annual invasion speed with  $R = -0.70$ . The estimated speed at the time the first site was sampled is  $v_0 = 80.27$  km/year, and acceleration  $v_1 = -2.83$  km/year<sup>2</sup> (b).

In addition, contrary to our fourth hypothesis, we observed no difference in the growth rates of earlier and more recently invading populations, suggesting that the time since invasion does not influence *D. villosus* population dynamics. In addition, time series representing earlier and more recent invasions originated from different biogeographical regions: earlier ones were mostly from Atlantic and Continental regions, and more recent ones from Pannonian and Alpine regions.


#### 4.2 | Occurrence frequency and invasion speed of *D. villosus*

We estimated that the introduction of *D. villosus* occurred on average 21 years before the first sampling event suggesting current monitoring of European streams is insufficient for early detection of invasive alien species. Following their introduction, such species often have low abundance during an initial establishment phase before increasing or becoming detected (Crooks et al., 1999), although lag phases are rarely measured in freshwater systems (but see Karatayev et al., 2011). Here, after a considerably long lag phase of 28 years from the time of introduction, *D. villosus* then only took another seven years to reach peak abundance. A lengthy lag period could reflect non-mutually exclusive phenomena including, *inter alia*, limits on the organism's reproductive rate in the early phase of exponential growth (e.g. Allee effects); (ii) multiple failed introductions prior to colonization success; (iii) genotypic selection of locally adapted organisms; and (iv) sudden population growth triggered by disturbance, environmental stochasticity or interspecific interactions (Crooks et al., 1999;

Crooks, 2005; Sakai et al., 2001; Spear et al., 2021). Although reporting efforts may be increased over time, this lag phase may explain the rapid decrease in new occurrences 35 years after an assemblage was first invaded (Ricciardi, 2013; Rouget et al., 2016). Nevertheless, records extracted from GBIF have to be taken with caution, as the taxonomic validity cannot always be ensured, simultaneously suggesting that many observations may be missing (Nekola et al., 2019; Shirey et al., 2019). Predicting future trends in the abundance of *D. villosus* is hampered by context dependencies, which may cause sudden shifts in population dynamics at different temporal scales, for example reflecting boom-bust dynamics (Strayer et al., 2017).

#### 4.3 | The influence of site-specific characteristics on *D. villosus*

Understanding how site-specific characteristics influence invasive populations can enable the identification of factors facilitating and limiting their spread. Supporting our third hypothesis, *D. villosus* populations were influenced by site-specific abiotic characteristics, in particular, elevation, distance to the next barrier and climatic variables (average minimum temperature and the trend of maximum temperature). Focusing first on the rate of change of *D. villosus* trends, the distance to the next barrier had a negative effect on *D. villosus* populations. The reservoirs created by the construction of barriers such as dams are a hotspot for the introduction of invasive alien species, due to, for example, recreational fishing activity (Anderson et al., 2014), with *D. villosus* able to survive for up to three and a half days out of water attached to ropes and other equipment



**FIGURE 5** Effect of general temporal trend (circles) and the number of sampled individuals of *Dikerogammarus villosus* (*D. villosus* shape) on the rate at which trends in community metrics (total abundance, richness, temporal turnover, Shannon diversity and Pielou's evenness) changed across all time series. Empty dots represent no significant effects  $\pm$  standard error (bars). Hollow shapes represent no significant effects of *D. villosus*, while the filled shapes represent significant effects. Red represents negative effects of *D. villosus* and negative trends in community metrics, while blue represents positive effects and positive trends of community metrics.

(Bacela-Spychalska et al., 2013). Furthermore, dams can be used as refuges and "stepping stones" for further spread, but also limit the spread upstream. Therefore, invasive alien species can accumulate near these barriers (Rahel, 2013). For invasive alien species, climatic variables – in particular temperature – can be the most important environmental variables determining the survival, reproduction and establishment in recipient ecosystems (Müller & Baur, 2011). Temperature is also well-known for its influence on life cycle characteristics such as fecundity in *D. villosus* (Pöckl et al., 2003). In congruence with Kobak et al. (2017), *D. villosus* preferred warm water and exhibited a stronger tendency to select extreme temperatures.

The dominance of *D. villosus* increased over time. This increase can be explained by propagule/colonization pressures, such as by the species exploiting increasing anthropogenic invasion corridors such as canals (Maclsaac et al., 2001; Lockwood et al., 2005). Reduced abiotic and biotic resistance resulting from degradation of ecosystems directly or indirectly by humans could also promote invasion (Hufbauer et al., 2012). The more rapid increase in *D. villosus* population growth at higher elevations nevertheless contradicts other studies that show lower elevation as high bioclimatic suitability for *D. villosus* (Gallardo et al., 2012), but suggests the species is invading higher elevated regions as a potential response to the ongoing climate change (Pauchard et al., 2016). Lastly, the dominance of *D.*

*villosus* was lower in streams in comparison to the major European rivers. This result may be partially explained by fast water flow or lower temperature in streams (Allan & Castillo, 2007; Grabowski et al., 2009), but also larger rivers having a greater level of confluence with smaller streams, functioning as shipping canals and therefore being prone to higher invasion rates. These results could be also affected by differences in sampling effort and the sampling methods among time series, which could have created biases and delays in detecting *D. villosus* at local and regional scales.

#### 4.4 | Impact of *D. villosus* on community metrics

Understandings of how invasive alien species impact the recipient ecosystems and the potential synergistic effects of anthropogenic stressors (abiotic and biotic factors) have become priorities in invasion science (Ricciardi et al., 2021). Supporting our fourth hypothesis, we identified a negative relationship between temporal changes in the number of sampled *D. villosus* individuals and in three metrics representing macroinvertebrate communities: taxon richness, temporal turnover and Shannon diversity. The negative impacts of *D. villosus* on invaded ecosystems are well-documented and include the depredation of a wide range of macroinvertebrates (e.g. chironomids, leeches, isopods and juvenile crayfish; Buřič et al., 2009; Dick et al., 2002; Platvoet et al., 2009), including via "wasteful" killing (Dick et al., 2002). Invasive amphipods have also been shown to display lower levels of omnivory than native species (Cuthbert, Kotronaki, Hütt, et al., 2022). This predatory capacity can reduce or replace functionally equivalent species via intraguild predation (e.g. native *Gammarus duebeni* by alien *G. tigrinus*; Rewicz et al., 2014). *Dikerogammarus villosus* also has negative effects on ecosystem functioning, including alteration of habitat structure, leaf litter decomposition and energy flows through food webs, potentially causing large-scale trophic cascades (Koester et al., 2016; MacNeil et al., 2011; Piscart et al., 2011; Van Riel et al., 2006). These impacts can create vacant niches that increase community susceptibility to other invasions and exacerbate the collective impacts of invasive alien species (Boets et al., 2010, 2011).

However, we stress that changes in community trends cannot be attributed exclusively to the effect of *D. villosus*, as correlation does not indicate causation, and were likely also altered by other anthropogenic and natural stressors not included in our models, including changes in water quality, disturbance events or even other invasive alien species (Didham et al., 2005; Haubrock et al., 2020; Pilotto et al., 2020). The combination of invasive alien species and other human impacts can promote the local extirpation of native species, reducing community diversity and driving biotic homogenization (Dormann et al., 2007; Ekroos et al., 2010; McKinney, 2004; McKinney & Lockwood, 1999), but can also promote or prevent invasions and/or increases in invader populations (Simberloff & Von Holle, 1999; Beaury et al., 2020).

## 4.5 | Implications and conclusions

The scope of our study was limited by the time series represented in our database. Invasive alien species, such as *D. villosus*, can affect food webs through either bottom-up or top-down regulation, potentially triggering trophic cascades that cause major disturbances in invaded ecosystems (Van Riel et al., 2006). The greater effects of *D. villosus* following steeper increases in its abundance suggest that the main effects on invaded communities are driven by high population densities, which overwhelm the ecological resistance of the recipient community. Surprisingly, the data from GBIF and our time series do not overlap in some cases, highlighting areas for future abundance survey efforts. *Dikerogammarus villosus* was first recorded in Italy in 1992 (Seebens et al., 2018), Belgium in 1998 (Seebens et al., 2018), France at the beginning of the 2000s (Devin et al., 2001) and the United Kingdom in 2010 (Bacela-Spychalska et al., 2013; Seebens et al., 2018), but *D. villosus* was not present in any time series from these countries, perhaps because our data are restricted to lotic systems. Our study was therefore limited by our focus on rivers and streams. *Dikerogammarus villosus* also occurs in lentic freshwaters including lakes, ponds, and brackish waters, and has marked impacts on these ecosystems (Bacela-Spychalska et al., 2013; Bollache et al., 2004; Minchin et al., 2019). Its occurrence in other countries could also be underestimated, resulting in relatively few time series, for example the Netherlands or Switzerland (Altermatt et al., 2014; Bij de Vaate & Klink, 1995). As such, further research would be needed to comprehensively characterize *D. villosus* impacts across all European freshwaters. Notwithstanding these data gaps, our results allow broad-scale inference of impacts using standardized time series across a range of invaded lotic European freshwaters.

Overall, our results show that *D. villosus* is well-established across the vast majority of Europe. Considering that *D. villosus* has invaded many freshwater and brackish ecosystems beyond those sites covered by the time series we considered in our analyses, it remains impossible to conclude anything about the capacity of *D. villosus* to expand further in Europe. Nonetheless, our documentation of *D. villosus* highlights the need for greater effort to reduce delays in the detection of invasive alien species to implement management techniques in an early stage of invasion, when such methods can be more effective (Ahmed et al., 2022; Lodge et al., 2016) and less expensive (Cuthbert, Diagne, et al., 2022; Fantle-Lepczyk et al., 2022; Hulme et al., 2009). These measures are especially important in those regions currently uninvaded, such as the North American Great Lakes that can act as “stepping stones” to assist further spread across the continent. Our use of long-term, large-scale time series also emphasizes the importance of long-term data in ecology (Crooks et al., 1999). Further long-term studies are necessary to increase our understanding of the population dynamics of *D. villosus* and other aquatic invaders across the breadth of ecosystems, and the context-dependencies that differentiate such dynamics, to provide better management information for stakeholders and governments.

## ACKNOWLEDGEMENTS

We acknowledge the contribution of Dr. rer. nat. Hanno Seebens, who helped with the extrapolation model of the occurrences of *Dikerogammarus villosus*. We also thank the Yale Centre for Research Computing for guidance and use of the research computing infrastructure. D.A.A. is funded by the Kuwait Foundation for the Advancement of Sciences (KFAS) (PR1914SM-01) and the Gulf University for Science and Technology (GUST) internal seed funds (Case no. 234597 & 253536). R.N.C. acknowledges funding from a Leverhulme Trust Early Career Fellowship (ECF-2021-001). P.J.H. and P.H. received funding from the EU Horizon 2020 project eLTER PLUS (Grant Agreement No. 871128). S.D. acknowledges funding by the Leibniz Competition (J45/2018) and support by the German Federal Ministry of Education and Research (BMBF; 033W034A). Open Access funding enabled and organized by Projekt DEAL.

## CONFLICT OF INTEREST

The authors declare no competing interests.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in a GitHub repository at <https://github.com/IsmaSA/Dikerogammarus-villosus-population-dynamics.git>.

## PEER REVIEW

The peer review history for this article is available at <https://publons.com/publon/10.1111/ddi.13649>.

## ORCID

Ismael Soto  <https://orcid.org/0000-0002-7288-6336>

Ross N. Cuthbert  <https://orcid.org/0000-0003-2770-254X>

Danish A. Ahmed  <https://orcid.org/0000-0002-2490-1546>

Antonín Kouba  <https://orcid.org/0000-0001-8118-8612>

Sami Domisch  <https://orcid.org/0000-0002-8127-9335>

Giuseppe Amatulli  <https://orcid.org/0000-0002-8341-2830>

Jens Kiesel  <https://orcid.org/0000-0002-4371-6434>

Longzhu Q. Shen  <https://orcid.org/0000-0001-5629-3007>

Margarita Florencio  <https://orcid.org/0000-0002-6688-7770>

Elizabeta Briski  <https://orcid.org/0000-0003-1896-3860>

Florian Altermatt  <https://orcid.org/0000-0002-4831-6958>

Gaït Archambaud-Suard  <https://orcid.org/0000-0001-9493-2279>

Zoltan Csabai  <https://orcid.org/0000-0003-1700-2574>

Mathieu Floury  <https://orcid.org/0000-0002-4952-5807>

Maxence Forcellini  <https://orcid.org/0000-0003-4921-2189>

Patrick Leitner  <https://orcid.org/0000-0001-8122-4265>

Anthony Maire  <https://orcid.org/0000-0003-0920-773X>

Anthony Ricciardi  <https://orcid.org/0000-0003-1492-0054>

Ralf B. Schärer  <https://orcid.org/0000-0003-3510-1701>

Rachel Stubbington  <https://orcid.org/0000-0001-8475-5109>

Gábor Várbiro  <https://orcid.org/0000-0001-5907-3472>

Ralf C. M. Verdonshot  <https://orcid.org/0000-0002-0977-5975>

Peter Haase  <https://orcid.org/0000-0002-9340-0438>

Phillip J. Haubrock  <https://orcid.org/0000-0003-2154-4341>



## REFERENCES

- Abatzoglou, J. T., Dobrowski, S. Z., Parks, S. A., & Hegewisch, K. C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958–2015. *Scientific Data*, 5(1), 1–12. [https://doi.org/10.1038/sdata.2017.191\(2018\)](https://doi.org/10.1038/sdata.2017.191(2018))
- Ahmed, D. A., Hudgins, E. J., Cuthbert, R. N., Kourantidou, M., Diagne, C., Haubrock, P. J., Leung, B., Liu, C., Leroy, B., Petrovskii, S., Beidas, A., & Courchamp, F. (2022). Managing biological invasions: The cost of inaction. *Biological Invasions*, 1–20, 1947–1948. <https://doi.org/10.1007/s10530-022-02799-2>
- Allan, J. D., & Castillo, M. M. (2007). The foundations of stream ecology. In *Stream Ecology: Structure and function of running waters* (pp. 359–372). Springer.
- Altermatt, F., Alther, R., Fišer, C., Jokela, J., Konec, M., Kury, D., Mächler, E., Stucki, P., & Westram, A. M. (2014). Diversity and distribution of freshwater amphipod species in Switzerland (Crustacea: Amphipoda). *PLoS One*, 9(10), e110328. <https://doi.org/10.1371/journal.pone.0110328>
- Amatulli, G., Garcia Marquez, J., Sethi, T., Kiesel, J., Grigoropoulou, A., Üblacker, M., Shen, L., & Domisch, S. (2022). Hydrography90m: A new high-resolution global hydrographic dataset. *Earth System Science Data Discussions*, 2022, 1–43. <https://doi.org/10.5194/essd-2022-9>
- Anderson, L. G., White, P. C., Stebbing, P. D., Stentiford, G. D., & Dunn, A. M. (2014). Biosecurity and vector behaviour: Evaluating the potential threat posed by anglers and canoeists as pathways for the spread of invasive non-native species and pathogens. *PLoS One*, 9(4), e92788. <https://doi.org/10.1371/journal.pone.0092788>
- Arim, M., Abades, S. R., Neill, P. E., Lima, M., & Marquet, P. A. (2006). Spread dynamics of invasive species. *Proceedings of the National Academy of Sciences of the United States of America*, 103(2), 374–378. <https://doi.org/10.1073/pnas.0504272102>
- Bacela-Spychalska, K., Grabowski, M., Rewicz, T., Konopacka, A., & Wattier, R. (2013). The 'killer shrimp' *Dikerogammarus villosus* (Crustacea, Amphipoda) invading Alpine lakes: Overland transport by recreational boats and scuba-diving gear as potential entry vectors? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23(4), 606–618. <https://doi.org/10.1002/aqc.2329>
- Bagnara, M., Nowak, L., Boehmer, H. J., Schöll, F., Schurr, F. M., & Seebens, H. (2022). Simulating the spread and establishment of alien species along aquatic and terrestrial transport networks—a multi-pathway and high-resolution approach. *Journal of Applied Ecology*, 59, 1769–1780. <https://doi.org/10.1111/1365-2664.14184>
- Beaury, E. M., Finn, J. T., Corbin, J. D., Barr, V., & Bradley, B. A. (2020). Biotic resistance to invasion is ubiquitous across ecosystems of the United States. *Ecology Letters*, 23(3), 476–482. <https://doi.org/10.1111/ele.13446>
- Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., & Wood, E. F. (2018). Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5(1), 1–12. <https://doi.org/10.1038/sdata.2018.214>
- Bellard, C., Cassey, P., & Blackburn, T. M. (2016). Alien species as a driver of recent extinctions. *Biology Letters*, 12(2), 20150623. <https://doi.org/10.1098/rsbl.2015.0623>
- Bij de Vaate, A., Jazdzewski, K., Ketelaars, H. A., Gollasch, S., & Van der Velde, G. (2002). Geographical patterns in range extension of Ponto-Caspian macroinvertebrate species in Europe. *Canadian Journal of Fisheries and Aquatic Sciences*, 59(7), 1159–1174. <https://doi.org/10.1139/f02-098>
- Bij de Vaate, A., & Klink, A. G. (1995). *Dikerogammarus villosus* Sowinsky (Crustacea: Gammaridae) a new immigrant in the Dutch part of the Lower Rhine. *Lauterbornia*, 20, 51–54.
- Boets, P., Lock, K., & Goethals, P. L. (2011). Using long-term monitoring to investigate the changes in species composition in the harbour of Ghent (Belgium). *Hydrobiologia*, 663(1), 155–166. <https://doi.org/10.1007/s10750-010-0567-2>
- Boets, P., Lock, K., Messiaen, M., & Goethals, P. L. (2010). Combining data-driven methods and lab studies to analyse the ecology of *Dikerogammarus villosus*. *Ecological Informatics*, 5(2), 133–139. <https://doi.org/10.1016/j.ecoinf.2009.12.005>
- Bollache, L., Devin, S., Wattier, R., Chovet, M., Beisel, J. N., Moreteau, J. C., & Rigaud, T. (2004). Rapid range extension of the Ponto-Caspian amphipod *Dikerogammarus villosus* in France: Potential consequences. *Hydrobiologia*, 160(1), 57–66. <https://doi.org/10.1127/0003-9136/2004/0160-0057>
- Bowler, D. E., Hof, C., Haase, P., Kröncke, I., Schweiger, O., Adrian, R., Baert, L., Bauer, H. G., Blick, T., Brooker, R. W., Dekoninck, W., Domisch, S., Eckmann, R., Hendrickx, F., Hickler, T., Klotz, S., Kraberg, A., Kühn, I., Matesanz, S., ... Böhning-Gaese, K. (2017). Cross-realm assessment of climate change impacts on species' abundance trends. *Nature Ecology & Evolution*, 1(3), 1–7. <https://doi.org/10.1038/s41559-016-0067>
- Bradbeer, S. J., Coughlan, N. E., Cuthbert, R. N., Crane, K., Dick, J. T., Caffrey, J. M., Lucy, F. E., Renals, T., Davis, E., Warren, D. A., Pile, B., Quinn, C., & Dunn, A. M. (2020). The effectiveness of disinfectant and steam exposure treatments to prevent the spread of the highly invasive killer shrimp, *Dikerogammarus villosus*. *Scientific Reports*, 10(1), 1–7. <https://doi.org/10.1038/s41598-020-58058-8>
- Burič, M., Kočí, L., Petrušek, A., Kouba, A., & Kozák, P. (2009). Invaders eating invaders: Potential trophic interactions between the amphipod *Dikerogammarus villosus* and juvenile crayfish *Orconectes limosus*. *Knowledge and Management of Aquatic Ecosystems*, 5, 394–395. <https://doi.org/10.1051/kmae/2009015>
- Carvalho, J. C., Cardoso, P., & Gomes, P. (2012). Determining the relative roles of species replacement and species richness differences in generating beta-diversity patterns. *Global Ecology and Biogeography*, 21, 760–771. <https://doi.org/10.1111/j.1466-8238.2011.00694.x>
- Colautti, R. I., Grigorovich, I. A., & MacIsaac, H. J. (2006). Propagule pressure: A null model for biological invasions. *Biological Invasions*, 8(5), 1023–1037. <https://doi.org/10.1007/s10530-006-9007-7>
- Cornes, R. C., van der Schrier, G., van den Besselaar, E. J., & Jones, P. D. (2018). An ensemble version of the E-OBS temperature and precipitation data sets. *Journal of Geophysical Research: Atmospheres*, 123(17), 9391–9409. <https://doi.org/10.1029/2017JD028200>
- Coutts, S. R., Helmstedt, K. J., & Bennett, J. R. (2018). Invasion lags: The stories we tell ourselves and our inability to infer process from pattern. *Diversity and Distributions*, 24(2), 244–251. <https://doi.org/10.1111/ddi.12669>
- Cressie, N. A. C. (1993). *Statistics for spatial data*. J. Wiley & Sons.
- Crooks, J. A. (2005). Lag times and exotic species: The ecology and management of biological invasions in slow-motion. *Ecoscience*, 12(3), 316–329. <https://doi.org/10.2980/i1195-6860-12-3-316.1>
- Crooks, J. A., Soulé, M. E., & Sandlund, O. T. (1999). Lag times in population explosions of invasive species: Causes and implications. *Invasive Species and Biodiversity Management*, 24, 103–125.
- Cuthbert, R. N., Diagne, C., Hudgins, E. J., Turbelin, A., Ahmed, D. A., Albert, C., Bodey, T. W., Briski, E., Essl, F., Haubrock, P. J., Gozlan, R. E., Kirichenko, N., Kourantidou, M., Kramer, A. M., & Courchamp, F. (2022). Biological invasion costs reveal insufficient proactive management worldwide. *Science of the Total Environment*, 819, 153404. <https://doi.org/10.1016/j.scitotenv.2022.153404>
- Cuthbert, R. N., Kotronaki, S. G., Carlton, J. T., Ruiz, G. M., Fofonoff, P., & Briski, E. (2022). Aquatic invasion patterns across the North Atlantic. *Global Change Biology*, 28, 1376–1387. <https://doi.org/10.1111/gcb.16016>
- Cuthbert, R. N., Kotronaki, S. G., Dick, J. T., & Briski, E. (2020). Salinity tolerance and geographical origin predict global alien amphipod



- invasions. *Biology Letters*, 16(9), 20200354. <https://doi.org/10.1098/rsbl.2020.0354>
- Cuthbert, R. N., Kotronaki, S. G., Hütt, J. C., Renk, E., Warlo, N., & Briski, E. (2022). Do alternative resources dampen functional responses of native but not alien gammarids? *Ecology and Evolution*, 12(9).
- De Ventura, L., Weissert, N., Tobias, R., Kopp, K., & Jokela, J. (2017). Identifying target factors for interventions to increase boat cleaning in order to prevent spread of invasive species. *Management of Biological Invasions*, 8(1), 71–84. <https://doi.org/10.3391/mbi.2017.8.1.07>
- Dedyu, I. I. (1980). *Amphipods of fresh and salt waters of the South-West part of the USSR*. Shtiintsa Publishers 220 pp.
- Devin, S., Beisel, J. N., Bachmann, V., & Moreteau, J. C. (2001). *Dikerogammarus villosus* (Amphipoda: Gammaridae): Another invasive species newly established in the Moselle River and French hydrosystems. *Annales de Limnologie*, 37(1), 21–27. <https://doi.org/10.1051/limn/2001001>
- Devin, S., Piscart, C., Beisel, J. N., & Moreteau, J. C. (2004). Life history traits of the invader *Dikerogammarus villosus* (Crustacea: Amphipoda) in the Moselle River, France. *International Review of Hydrobiology*, 89(1), 21–34. <https://doi.org/10.1002/iroh.200310667>
- Diagne, C., Leroy, B., Vaissière, A. C., Gozlan, R. E., Roiz, D., Jarić, I., Salles, J.-M., Bradshaw, C. J. A., & Courchamp, F. (2021). High and rising economic costs of biological invasions worldwide. *Nature*, 592(7855), 571–576. <https://doi.org/10.1038/s41586-022-05100-6>
- Dick, J. T., & Platvoet, D. (2000). Invading predatory crustacean *Dikerogammarus villosus* eliminates both native and exotic species. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 267(1447), 977–983.
- Dick, J. T., Platvoet, D., & Kelly, D. W. (2002). Predatory impact of the freshwater invader *Dikerogammarus villosus* (Crustacea: Amphipoda). *Canadian Journal of Fisheries and Aquatic Sciences*, 59(6), 1078–1084. <https://doi.org/10.1139/f02-074>
- Didham, R. K., Tylianakis, J. M., Hutchison, M. A., Ewers, R. M., & Gemmill, N. J. (2005). Are invasive species the drivers of ecological change? *Trends in Ecology & Evolution*, 20(9), 470–474. <https://doi.org/10.1016/j.tree.2005.07.006>
- Dorman, M., Svoray, T., Perevolotsky, A., & Sarris, D. (2013). Forest performance during two consecutive drought periods: Diverging long-term trends and short-term responses along a climatic gradient. *Forest Ecology and Management*, 310, 1–9.
- Dormann, C. F., Schweiger, O., Augenstein, I., Bailey, D., Billeter, R., de Blust, G., DeFilippi, R., Frenzel, M., Hendrickx, F., Herzog, F., Klotz, S., Liira, J., Maelfait, J.-P., Schmidt, T., Speelmans, M., Van Wingerden, W. K. R. E., & Zobel, M. (2007). Effects of landscape structure and land-use intensity on similarity of plant and animal communities. *Global Ecology and Biogeography*, 16, 774–787. <https://doi.org/10.1111/j.1466-8238.2007.00344.x>
- Dornelas, M., Antao, L. H., Moyes, F., Bates, A. E., Magurran, A. E., Adam, D., Akhmetzhanova, A. A., Appeltans, W., Arcos, J. M., Arnold, H., Ayyappan, N., Badihi, G., Baird, A. H., Barbosa, M., Barreto, T. E., Bässler, C., Bellgrove, A., Belmaker, J., Benedetti-Cecchi, L., & Murphy, G. (2018). BioTIME: A database of biodiversity time series for the Anthropocene. *Global Ecology and Biogeography*, 27(7), 760–786. <https://doi.org/10.1111/geb.12729>
- Dornelas, M., Gotelli, N. J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., & Magurran, A. E. (2014). Assemblage time series reveal biodiversity change but not systematic loss. *Science*, 344(6181), 296–299. <https://doi.org/10.1126/science.1248484>
- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315(7109), 629–634.
- Ekroos, J., Heliölä, J., & Kuussaari, M. (2010). Homogenisation of lepidopteran communities in intensively cultivated agricultural landscapes. *Journal of Applied Ecology*, 47, 459–467. <https://doi.org/10.1111/j.1365-2664.2009.01767.x>
- Essl, F., Lenzer, B., Bacher, S., Bailey, S., Capinha, C., Daehler, C., Dullinger, S., Genovesi, P., Hui, C., Hulme, P. E., Jeschke, J. M., Katsanevakis, S., Kühn, I., Leung, B., Liebhold, A., Liu, C., MacIsaac, H. J., Meyerson, L. A., Nuñez, M. A., ... Roura-Pascual, N. (2020). Drivers of future alien species impacts: An expert-based assessment. *Global Change Biology*, 26(9), 4880–4893. <https://doi.org/10.1111/gcb.15199>
- European Environment Agency (EEA). (2021). <https://www.eea.europa.eu/data-and-maps/indicators/european-precipitation-2/assessment>
- European Space Agency, ESA. (2017). *Land cover CCI product user guide version 2*. Technical report. [maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2\\_2.0.pdf](https://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-Ph2-PUGv2_2.0.pdf)
- Fantle-Lepczyk, J. E., Haubrock, P. J., Kramer, A. M., Cuthbert, R. N., Turbelin, A. J., Crystal-Ornelas, R., Diagne, C., & Courchamp, F. (2022). Economic costs of biological invasions in the United States. *Science of the Total Environment*, 806, 151318. <https://doi.org/10.1016/j.scitotenv.2021.151318>
- Fournier, A., Penone, C., Pennino, M. G., & Courchamp, F. (2019). Predicting future invaders and future invasions. *Proceedings of the National Academy of Sciences of the United States of America*, 116(16), 7905–7910. <https://doi.org/10.1073/pnas.1803456116>
- Gallardo, B., Errea, M. P., & Aldridge, D. C. (2012). Application of bioclimatic models coupled with network analysis for risk assessment of the killer shrimp, *Dikerogammarus villosus*, in Great Britain. *Biological Invasions*, 14(6), 1265–1278. <https://doi.org/10.1007/s10530-011-0154-0>
- GBIF.org. (2021). GBIF occurrence download. <https://doi.org/10.15468/dl.t2er6w>
- Grabowski, M., Bacela, K., Konopacka, A., & Jazdzewski, K. (2009). Salinity-related distribution of alien amphipods in rivers provides refugia for native species. *Biological Invasions*, 11(9), 2107–2117.
- Hallett, L. M., Jones, S. K., MacDonald, A. A. M., Jones, M. B., Flynn, D. F., Ripplinger, J., Slaughter, P., Gries, C., & Collins, S. L. (2016). codyn: An R package of community dynamics metrics. *Methods in Ecology and Evolution*, 7(10), 1146–1151.
- Hamed, K. H., & Rao, A. R. (1998). A modified Mann-Kendall trend test for autocorrelated data. *Journal of Hydrology*, 204(1–4), 182–196. [https://doi.org/10.1016/S0022-1694\(97\)00125-X](https://doi.org/10.1016/S0022-1694(97)00125-X)
- Haubrock, P., Ahmed, D. A., Cuthbert, R. N., Stubbington, R., Domisch, S., Marquez, J. R. G., Beidas, A., Amatulli, G., Kiesel, J., Shen, L. Q., Soto, I., Angeler, D. G., Bonada, N., Cañedo-Argüelles, M., Csabai, Z., Datry, T., de Eyto, E., Dohet, A., Drohan, E., & Haase, P. (2022). Invasion impacts and dynamics of a European-wide introduced species. *Global Change Biology*, 28, 4620–4632. <https://doi.org/10.1111/gcb.16207>
- Haubrock, P. J., Pilotto, F., Innocenti, G., Cianfanelli, S., & Haase, P. (2020). Two centuries for an almost complete community turnover from native to non-native species in a riverine ecosystem. *Global Change Biology*, 27(3), 606–623. <https://doi.org/10.1111/gcb.15442>
- Higashino, M., & Stefan, H. G. (2019). Variability and change of precipitation and flood discharge in a Japanese river basin. *Journal of Hydrology: Regional Studies*, 21, 68–79.
- Holdich, D. M., & Pöckl, M. (2007). Invasive crustaceans in European inland waters. In *Biological invaders in inland waters: Profiles, distribution, and threats* (pp. 29–75). Springer.
- Huber, T., Leitner, P., Remund, N., Graf, W., Paunović, M., Borza, P., Csányi, B., Szekeres, J., & Szekeres, J. (2015). Longitudinal distributional patterns of Peracarida (Crustacea, Malacostraca) in the River Danube. *Fundamental and Applied Limnology*, 187(2), 113–126. <https://doi.org/10.1127/fal/2015/0769>

- Hufbauer, R. A., Facon, B., Ravigné, V., Turgeon, J., Foucaud, J., Lee, C. E., Rey, O., & Estoup, A. (2012). Anthropogenically induced adaptation to invade (AIAI): Contemporary adaptation to human-altered habitats within the native range can promote invasions. *Evolutionary Applications*, 5(1), 89–101. <https://doi.org/10.1111/j.1752-4571.2011.00211.x>
- Hulme, P. E., Pyšek, P., Nentwig, W., & Vilà, M. (2009). Will threat of biological invasions unite the European Union. *Science*, 324(5923), 40–41. <https://doi.org/10.1126/science.1171111>
- Karatayev, A. Y., Burlakova, L. E., Mastitsky, S. E., Padilla, D. K., & Mills, E. L. (2011). Contrasting rates of spread of two congeners, *Dreissena polymorpha* and *Dreissena rostriformis bugensis*, at different spatial scales. *Journal of Shellfish Research*, 30(3), 923–931.
- Kobak, J., Jermacz, Ł., Marcińczyk, J., Bartoszyńska, E., Rutkowska, D., & Pawłowska, K. (2017). Abiotic factors affecting habitat selection by two invasive gammarids *Dikerogammarus villosus* and *Pontogammarus robustoides*. *Hydrobiologia*, 797(1), 247–263. <https://doi.org/10.1007/s10750-017-3185-4>
- Kobak, J., Rachalewski, M., & Bącela-Spychalska, K. (2016). Conquerors or exiles? Impact of interference competition among invasive Ponto-Caspian gammarideans on their dispersal rates. *Biological Invasions*, 18(7), 1953–1965. <https://doi.org/10.1007/s10530-016-1140-3>
- Koester, M., Bayer, B., & Gergs, R. (2016). Is *Dikerogammarus villosus* (Crustacea, Gammaridae) a 'killer shrimp' in the River Rhine system? *Hydrobiologia*, 768(1), 299–313. <https://doi.org/10.1007/s10750-015-2558-9>
- Kouba, A., Oficialdegui, F. J., Cuthbert, R. N., Kourantidou, M., South, J., Tricario, E., Leroy, B., Gozlan, R., Courchamp, F., & Haubrock, P. J. (2021). Identifying economic costs and knowledge gaps of invasive aquatic crustaceans. *Science of the Total Environment*, 813, 152325. <https://doi.org/10.1016/j.scitotenv.2021.152325>
- Kramer, A. M., Annis, G., Wittmann, M. E., Chadderton, W. L., Rutherford, E. S., Lodge, D. M., Mason, L., Beletsky, D., Riseng, C., & Drake, J. M. (2017). Suitability of Laurentian Great Lakes for invasive species based on global species distribution models and local habitat. *Ecosphere*, 8(7), e01883. <https://doi.org/10.1002/ecs2.1883>
- Leuven, R. S., van der Velde, G., Baijens, I., Snijders, J., van der Zwart, C., Lenders, H. R., & bij de Vaate, A. (2009). The river Rhine: A global highway for dispersal of aquatic invasive species. *Biological Invasions*, 11(9), 1989–2008. <https://doi.org/10.1007/s10530-009-9491-7>
- Lockwood, J. L., Cassey, P., & Blackburn, T. M. (2005). The role of propagule pressure in explaining species invasion. *Trends in Ecology Evolution*, 20, 223–228. <https://doi.org/10.1016/j.tree.2005.02.004>
- Lodge, D. M., Simonin, P. W., Burgiel, S. W., Keller, R. P., Bossenbroek, J. M., Jerde, C. L., Kramer, A. M., Rutherford, E. S., Barnes, M. A., Wittmann, M. E., Chadderton, W. L., Apriesnig, J. L., Beletsky, D., Cooke, R. M., Drake, J. M., Egan, S. P., Finnoff, D. C., Gantz, C. A., Grey, E. K., ... Zhang, H. (2016). Risk analysis and bioeconomics of invasive species to inform policy and management. *Annual Review of Environment and Resources*, 41, 453–488. <https://doi.org/10.1146/annurev-environ-110615-085532>
- MacIsaac, H. J., Grigorovich, I. A., & Ricciardi, A. (2001). Reassessment of species invasions concepts: The Great Lakes basin as a model. *Biological Invasions*, 3(4), 405–416. <https://doi.org/10.1023/A:1015854606465>
- MacNeil, C., Boets, P., Lock, K., & Goethals, P. L. (2013). Potential effects of the invasive 'killer shrimp' (*Dikerogammarus villosus*) on macroinvertebrate assemblages and biomonitoring indices. *Freshwater Biology*, 58(1), 171–182. <https://doi.org/10.1111/fwb.12048>
- MacNeil, C., Dick, J. T., Platvoet, D., & Briffa, M. (2011). Direct and indirect effects of species displacements: An invading freshwater amphipod can disrupt leaf-litter processing and shredder efficiency. *Journal of the North American Benthological Society*, 30(1), 38–48.
- MacNeil, C., & Platvoet, D. (2013). Could artificial structures such as fish passes facilitate the establishment and spread of the 'killer shrimp' *Dikerogammarus villosus* (Crustacea: Amphipoda) in river systems? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23(5), 667–677. <https://doi.org/10.1002/aqc.2337>
- Maire, A., Thierry, E., Viechtbauer, W., & Daufresne, M. (2019). Poleward shift in large-river fish communities detected with a novel meta-analysis framework. *Freshwater Biology*, 64(6), 1143–1156. <https://doi.org/10.1111/fwb.13291>
- McKinney, M. L. (2004). Do exotics homogenize or differentiate communities? Roles of sampling and exotic species richness. *Biological Invasions*, 6, 495–504. <https://doi.org/10.1023/B:BINV.0000041562.31023.42>
- McKinney, M. L., & Lockwood, J. L. (1999). Biotic homogenisation: A few winners replacing many losers in the next mass extinction. *Trends in Ecology & Evolution*, 14(11), 450–453. [https://doi.org/10.1016/S0169-5347\(99\)01679-1](https://doi.org/10.1016/S0169-5347(99)01679-1)
- Minchin, J. D., Arbačiauskas, K., Daunys, D., Ezhova, E., Grudule, N., Kotta, J., Molchanova, N., Olenin, S., Višinskienė, G., & Strake, S. (2019). Rapid expansion and facilitating factors of the Ponto-Caspian invader *Dikerogammarus villosus* within the eastern Baltic Sea. *Aquatic Invasions*, 14(2), 165–181. <https://doi.org/10.3391/ai.2019.14.2.02>
- Mirtl, M., Borer, E. T., Djukic, I., Forsius, M., Haubold, H., Hugo, W., Jourdan, J., Lindenmayer, D., McDowell, W. H., Muraoka, H., Orenstein, D. E., Pauw, J. C., Peterseil, J., Shibata, H., Wohner, C., Yu, X., & Haase, P. (2018). Genesis, goals and achievements of long-term ecological research at the global scale: A critical review ofILTER and future directions. *Science of the Total Environment*, 626, 1439–1462, 2435–2446. <https://doi.org/10.1016/j.scitotenv.2017.12.001>
- Mordukhai-Boltovskoi, F. D. (1960). *Caspian fauna in the Azov and Black Sea Basin*. Izdatelstvo Akademii Nauk SSSR.
- Müller, O., & Baur, B. (2011). Survival of the invasive clam *Corbicula fluminea* (Müller) in response to winter water temperature. *Malacologia*, 53(2), 367–371. <https://doi.org/10.4002/040.053.0207>
- Nekola, J. C., Hutchins, B. T., Schofield, A., Najev, B., & Perez, K. E. (2019). Caveat consumptor notitia museo: Let the museum data user beware. *Global Ecology and Biogeography*, 28(12), 1722–1734. <https://doi.org/10.1111/geb.12995>
- Nentwig, W., Bacher, S., Kumschick, S., Pyšek, P., & Vilà, M. (2018). More than "100 worst" alien species in Europe. *Biological Invasions*, 20(6), 1611–1621. <https://doi.org/10.1007/s10530-017-1651-6>
- Nolan, J. P. (2020). *Univariate stable distributions*. Springer.
- Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R. B., Simpson, L. G., & Oksanen, M. J. (2013). *Package 'vegan'*. Community ecology package, version, 2(9), 1–295.
- Pauchard, A., Milbau, A., Albiñ, A., Alexander, J., Burgess, T., Daehler, C., Englund, G., Essl, F., Evengård, B., Greenwood, G. B., Haider, S., Lenoir, J., McDougall, K., Muths, E., Nuñez, M. A., Olofsson, J., Pellissier, L., Rabitsch, W., Rew, L. J., ... Kueffer, C. (2016). Non-native and native organisms moving into high elevation and high latitude ecosystems in an era of climate change: New challenges for ecology and conservation. *Biological Invasions*, 18(2), 345–353. <https://doi.org/10.1007/s10530-015-1025-x>
- Pielou, E. C. (1966). The measurement of diversity in different types of biological collections. *Journal of Theoretical Biology*, 13, 131–144.
- Pilotto, F., Kühn, I., Adrian, R., Alber, R., Alignier, A., Andrews, C., Bäck, J., Barbaro, L., Beaumont, D., Beenaerts, N., Benham, S., Boukal, D. S., Bretagnolle, V., Camatti, E., Canullo, R., Cardoso, P. G., Ens, B. J., Everaert, G., Evtimova, V., ... Haase, P. (2020). Meta-analysis of multidecadal biodiversity trends in Europe. *Nature Communications*, 11(1), 1–11. <https://doi.org/10.1038/s41467-020-17171-y>
- Piscart, C., Kefford, B. J., & Beisel, J. N. (2011). Are salinity tolerances of non-native macroinvertebrates in France an indicator of potential for their translocation in a new area? *Limnologia*, 41(2), 107–112. <https://doi.org/10.1016/j.limno.2010.09.002>

- Platvoet, D., Dick, J. T., MacNeil, C., van Riel, M. C., & van der Velde, G. (2009). Invader–invader interactions in relation to environmental heterogeneity leads to zonation of two invasive amphipods, *Dikerogammarus villosus* (Sowinsky) and *Gammarus tigrinus* Sexton: Amphipod pilot species project (AMPIS) report 6. *Biological Invasions*, 11(9), 2085–2093.
- Pöckl, M. (2009). Success of the invasive Ponto-Caspian amphipod *Dikerogammarus villosus* by life history traits and reproductive capacity. *Biological Invasions*, 11(9), 2021–2041. <https://doi.org/10.1007/s10530-009-9485-5>
- Pöckl, M., Webb, B. W., & Sutcliffe, D. W. (2003). Life history and reproductive capacity of *Gammarus fossarum* and *G. roeseli* (Crustacea: Amphipoda) under naturally fluctuating water temperatures: A simulation study. *Freshwater Biology*, 48(1), 53–66. <https://doi.org/10.1046/j.1365-2427.2003.00967.x>
- Pyšek, P., Hulme, P. E., Simberloff, D., Bacher, S., Blackburn, T. M., Carlton, J. T., Dawson, W., Essl, F., Foxcroft, L. C., Genovesi, P., Jeschke, J. M., Kühn, I., Liebhold, A. M., Mandrak, N. E., Meyerson, L. A., Pauchard, A., Pergl, J., Roy, H. E., Seebens, H., ... Richardson, D. M. (2020). Scientists' warning on invasive alien species. *Biological Reviews*, 95(6), 1511–1534. <https://doi.org/10.1111/brv.12627>
- R Core Team. (2022). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rahel, F. J. (2013). Intentional fragmentation as a management strategy in aquatic systems. *Bioscience*, 63, 363–373. <https://doi.org/10.1525/bio.2013.63.5.9>
- Rewicz, T., Grabowski, M., MacNeil, C., & Bacela-Spychalska, K. (2014). The profile of a 'perfect' invader—the case of killer shrimp, *Dikerogammarus villosus*. *Aquatic Invasions*, 9(3), 267–288. <https://doi.org/10.3391/ai.2014.9.3.04>
- Ricciardi, A. (2013). Invasive species. In *Ecological systems* (pp. 161–178). Springer.
- Ricciardi, A., Iacarella, J. C., Aldridge, D. C., Blackburn, T. M., Carlton, J. T., Catford, J. A., Dick, J. T. A., Hulme, P. E., Jeschke, J. M., Liebhold, A. M., Lockwood, J., Maclsaac, H., Meyerson, L., Pyšek, P., Richardson, D., Ruiz, G., Simberloff, D., Vilà, M., & Wardle, D. A. (2021). Four priority areas to advance invasion science in the face of rapid environmental change. *Environmental Reviews*, 29(2), 119–141. <https://doi.org/10.1139/er-2020-0088>
- Ripley, B., Venables, B., Bates, D. M., Hornik, K., Gebhardt, A., Firth, D., & Ripley, M. B. (2013). Package 'mass'. *Cran r*, 538, 113–120.
- Roje, S., Švagrová, K., Veselý, L., Sentis, A., Kouba, A., & Buřič, M. (2021). Pilferer, murderer of innocents or prey? The potential impact of killer shrimp (*Dikerogammarus villosus*) on crayfish. *Aquatic Sciences*, 83(1), 1–12. <https://doi.org/10.1007/s00027-020-00762-8>
- Rolla, M., Consuegra, S., & de Leaniz, C. G. (2020). Predator recognition and anti-predatory behaviour in a recent aquatic invader, the killer shrimp (*Dikerogammarus villosus*). *Aquatic Invasions*, 15(3), 482–496. <https://doi.org/10.3391/ai.2020.15.3.08>
- Rolla, M., Consuegra, S., Hall, D. J., & Garcia de Leaniz, C. (2020). Seasonal and spatial variation in growth and abundance of zebra mussel (*Dreissena polymorpha*) in a recently invaded artificial lake: Implications for management. *Frontiers in Ecology and Evolution*, 8, 159. <https://doi.org/10.3389/fevo.2020.00159>
- Rouget, M., Robertson, M. P., Wilson, J. R., Hui, C., Essl, F., Renteria, J. L., & Richardson, D. M. (2016). Invasion debt—quantifying future biological invasions. *Diversity and Distributions*, 22(4), 445–456. <https://doi.org/10.1111/ddi.12408>
- Sakai, A. K., Allendorf, F. W., Holt, J. S., Lodge, D. M., Molofsky, J., With, K. A., Baughman, S., Cabin, R. J., Cohen, J. E., Ellstrand, N. C., McCauley, D. E., O'Neil, P., Parker, I. M., Thompson, J. N., & Weller, S. G. (2001). The population biology of invasive species. *Annual Review of Ecology and Systematics*, 32(1), 305–332.
- Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., Pagad, S., Pyšek, P., van Kleunen, M., Winter, M., Ansong, M., Arianoutsou, M., Bacher, S., Blasius, B., Brocknerhoff, E. G., Brundu, G., Capinha, C., Causton, C. E., Celesti-Grappow, L., ... Essl, F. (2018). Global rise in emerging alien species results from increased accessibility of new source pools. *Proceedings of the National Academy of Sciences of the United States of America*, 115(10), E2264–E2273. <https://doi.org/10.1073/pnas.1719429115>
- Seebens, H., Blackburn, T. M., Hulme, P. E., van Kleunen, M., Liebhold, A. M., Orlova-Bienkowskaja, M., Pyšek, P., Schindler, S., & Essl, F. (2021). Around the world in 500 years: Inter-regional spread of alien species over recent centuries. *Global Ecology and Biogeography*, 30, 1621–1632. <https://doi.org/10.1111/geb.13325>
- Seebens, H., Briski, E., Ghabooli, S., Shiganova, T., Maclsaac, H. J., & Blasius, B. (2019). Non-native species spread in a complex network: The interaction of global transport and local population dynamics determines invasion success. *Proceedings of the Royal Society B: Biological Sciences*, 286(1901), 20190036. <https://doi.org/10.1098/rspb.2019.0036>
- Seebens, H., Clarke, D. A., Groom, Q., Wilson, J. R., Garcia-Berthou, E., Kühn, I., Roigé, M., Pagad, S., Essl, F., Vicente, J., Winter, M., & McGeoch, M. (2020). A workflow for standardising and integrating alien species distribution data.
- Shannon, C. E., & Weaver, W. (1949). The mathematical theory of communication. *The Bell System Technical Journal*, 27.
- Shi, J., Luo, Y. Q., Zhou, F., & He, P. (2010). The relationship between invasive alien species and main climatic zones. *Biodiversity and Conservation*, 19(9), 2485–2500. <https://doi.org/10.1007/s10531-010-9855-4>
- Shirey, V., Seppälä, S., Branco, V. V., & Cardoso, P. (2019). Current GBIF occurrence data demonstrates both promise and limitations for potential red listing of spiders. *Biodiversity Data Journal*, 7, e47369. <https://doi.org/10.3897/BDJ.7.e47369>
- Simberloff, D., & Von Holle, B. (1999). Positive interactions of nonindigenous species: Invasional meltdown? *Biological Invasions*, 1(1), 21–32. <https://doi.org/10.1023/A:1010086329619>
- Sofaer, H. R., Jarnevich, C. S., & Pearse, I. S. (2018). The relationship between invader abundance and impact. *Ecosphere*, 9(9), e02415. <https://doi.org/10.1002/ecs2.2415>
- Spear, M. J., Walsh, J. R., Ricciardi, A., & Zanden, M. (2021). The invasion ecology of sleeper populations: Prevalence, persistence, and abrupt shifts. *Bioscience*, 71(4), 357–369. <https://doi.org/10.1093/biosci/biaa168>
- Strayer, D. L., D'Antonio, C. M., Essl, F., Fowler, M. S., Geist, J., Hilt, S., Jarić, I., Jöhnk, K., Jones, C. G., Lambin, X., Latzka, A. W., Pergl, J., Pyšek, P., Robertson, P., von Schmalensee, M., Steffansson, R. A., Wright, J., & Jeschke, J. M. (2017). Boom-bust dynamics in biological invasions: Towards an improved application of the concept. *Ecology Letters*, 20(10), 1337–1350. <https://doi.org/10.1111/ele.12822>
- Strayer, D. L., Eviner, V. T., Jeschke, J. M., & Pace, M. L. (2006). Understanding the long-term effects of species invasions. *Trends in Ecology & Evolution*, 21(11), 645–651. <https://doi.org/10.1016/j.tree.2006.07.007>
- Taylor, N. G., & Dunn, A. M. (2017). Size matters: Predation of fish eggs and larvae by native and invasive amphipods. *Biological Invasions*, 19(1), 89–107. <https://doi.org/10.1007/s10530-016-1265-4>
- Van Riel, M. C., Van der Velde, G., & Bij de Vaate, A. (2006). To conquer and persist: Colonisation and population development of the Ponto-Caspian amphipods *Dikerogammarus villosus* and *Chelicorophium curvispinum* on bare stone substrate in the main channel of the River Rhine. *Archiv für Hydrobiologie*, 166(1), 23–40. <https://doi.org/10.1127/0003-9136/2006/0166-0023>
- Van Riel, M. C., Van der Velde, G., Rajagopal, S., Marguillier, S., Dehairs, F., & Bij de Vaate, A. (2006). Trophic relationships in the Rhine food web during invasion and after establishment of the Ponto-Caspian invader *Dikerogammarus villosus*. *Hydrobiologia*, 565(1), 39–58. <https://doi.org/10.1007/s10750-005-1904-8>

- Viechtbauer, W. (2010). Conducting meta-analyses in R with the *metafor* package. *Journal of Statistical Software*, 36(3), 1–48.
- Viechtbauer, W., Smits, L., Kotz, D., Budé, L., Spigt, M., Serroyen, J., & Crutzen, R. (2015). A simple formula for the calculation of sample size in pilot studies. *Journal of Clinical Epidemiology*, 68(11), 1375–1379. <https://doi.org/10.1016/j.jclinepi.2015.04.014>
- Warren, D. A., Bradbeer, S. J., & Dunn, A. M. (2021). Superior predatory ability and abundance predicts potential ecological impact towards early-stage anurans by invasive 'Killer Shrimp' (*Dikerogammarus villosus*). *Scientific Reports*, 11(1), 1–12. <https://doi.org/10.1038/s41598-021-82630-5>
- Wood, L., Smith, E., Bojko, J., & Stebbing, P. (2021). Options for the control of *Dikerogammarus villosus* (killer shrimp) and other invasive amphipods: Invasive Amphipod Control. *Management of Biological Invasions*, 12, 662–684.
- Yamazaki, D., Ikeshima, D., Sosa, J., Bates, P. D., Allen, G. H., & Pavelsky, T. M. (2019). MERIT Hydro: A high-resolution global hydrography map based on latest topography dataset. *Water Resources Research*, 55(6), 5053–5073. <https://doi.org/10.1029/2019WR024873>
- Yokomizo, H., Possingham, H. P., Thomas, M. B., & Buckley, Y. M. (2009). Managing the impact of invasive species: The value of knowing the density–impact curve. *Ecological Applications*, 19(2), 376–386. <https://doi.org/10.1890/08-0442.1>
- Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A., & Smith, G. M. (2009). *Mixed effects models and extensions in ecology with R* (Vol. 574). Springer.

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Soto, I., Cuthbert, R. N., Ahmed, D. A., Kouba, A., Domisch, S., Marquez, J. R. G., Beidas, A., Amatulli, G., Kiesel, J., Shen, L. Q., Florencio, M., Lima, H., Briski, E., Altermatt, F., Archambaud-Suard, G., Borza, P., Csabai, Z., Datry, T., Floury, M. ... Haubrock, P. J. (2022). Tracking a killer shrimp: *Dikerogammarus villosus* invasion dynamics across Europe. *Diversity and Distributions*, 00, 1–16. <https://doi.org/10.1111/ddi.13649>

## BIOSKETCH

**Ismael Soto** is a PhD student at the faculty of fisheries and protection of waters at the University of South Bohemia (Czech Republic). His works is focused on the long-term trends and impacts of invasive alien species in European inland waters.

Author contributions: IS: Methodology, Formal analysis, Data Curation, Writing – Original Draft, Writing – review & editing. RC, DA: Conceptualization, Writing – Original Draft, Writing – review & editing. AK, SD, JM, AB, GA, JK, LS, MF, HL, EB, FA, GA, PB, ZC, TD, MF, MF, JF, PL, ML, AM, AR, RS, RS, GV, GV, RV: Resources, Writing – Review & Editing. PH: Funding acquisition and Project, Writing – Review & Editing Administration. PJH: Conceptualization, Supervision, Writing – original draft; Writing – review & editing.