



HAL
open science

Steady Kähler-Ricci solitons on crepant resolutions of finite quotients of C

Olivier Biquard, Heather Macbeth

► **To cite this version:**

Olivier Biquard, Heather Macbeth. Steady Kähler-Ricci solitons on crepant resolutions of finite quotients of C . Journal of the London Mathematical Society, 2024, 109, pp.e12833. 10.1112/jlms.12833. hal-03844675

HAL Id: hal-03844675

<https://hal.science/hal-03844675>

Submitted on 22 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Steady Kähler-Ricci solitons on crepant resolutions of finite quotients of \mathbb{C}^n

Olivier Biquard and Heather Macbeth

Abstract

We prove the existence of steady Kähler-Ricci solitons on equivariant crepant resolutions of \mathbb{C}^n/G , where G is a finite subgroup of $SU(n)$ acting freely on \mathbb{C}^n .

1 Introduction

1.1 Overview

The ‘fixed points’ of the Ricci flow are *steady Ricci solitons*. Such objects, natural generalizations of Ricci-flat metrics, are pairs (g, X) of a Riemannian metric and a vector field, satisfying the elliptic partial differential equation

$$\text{Ric}(g) + \frac{1}{2}\mathcal{L}_X g = 0.$$

They arise in the study of the Ricci flow as models of singularities [DdP07, GZ08, DS10, AIK15], and as backward limits of ancient solutions. They are also critical points, in a suitable sense [Has11], of the Perelman \mathcal{F} -functional, and thus can be considered canonical among all such pairs (g, X) .

Steady Ricci solitons which are not Ricci-flat must be noncompact [Ive93]. The few known examples include several which are Kähler, with holomorphic vector field: Hamilton’s *cigar soliton* [Ham88] on \mathbb{C} , H.-D. Cao’s generalizations [Cao96] on \mathbb{C}^n and $K_{\mathbb{C}P^{n-1}}$, and further generalizations by Dancer-M. Wang [DW11] and B. Yang [Yan12]. Non-Kähler examples include the well-known constructions of Bryant and of Ivey. All these examples – and, we believe, all known examples – are highly symmetric, and the soliton metric is given either explicitly or by solving an ODE.

In this article we use PDE methods to construct new steady Kähler-Ricci solitons (M, ω, X) , in all complex dimensions $n \geq 2$ (real dimensions $2n \geq 4$), of infinitely many topological types in dimensions 2 and 3 at least. Like all steady Kähler-Ricci solitons, they have first Chern class

$$c_1(M) = [\text{Ric}(\omega)] = [-\frac{1}{2}\mathcal{L}_X \omega] = 0.$$

Our construction proceeds by taking Joyce’s well-known family [Joy00] of Ricci-flat Kähler metrics on *crepant resolutions* of orbifolds \mathbb{C}^n/G (which automatically have $c_1(M) = 0$), and modifying their metrics near infinity by *gluing* them to a G -quotient of Cao’s soliton on \mathbb{C}^n . A precise statement is as follows:

This material is based upon work supported by the National Science Foundation under Grant No. DMS-1440140 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2016 semester.

Theorem 1.1. *Let G be a finite subgroup of $SU(n)$, which acts freely on $\mathbb{C}^n \setminus \{0\}$. Let M_G be a crepant resolution of \mathbb{C}^n/G , which is equivariant with respect to the action of \mathbb{C}^* . Let ϵ be a Kähler class on M_G which contains an asymptotically locally Euclidean Kähler metric.*

Then for all ϵ sufficiently small, there exists a steady Kähler-Ricci soliton on M_G in the cohomology class ϵ , whose drift vector field is the extension to M_G of the radial vector field $-2r\partial/\partial r$ (this extension exists by the equivariance of the resolution).

Moreover, outside a compact set these solitons have the form $\omega_0 + \frac{i}{2}\partial\bar{\partial}\Psi$, where ω_0 is Cao's steady Kähler-Ricci soliton metric on \mathbb{C}^n (descending to \mathbb{C}^n/G), and, for any k and any $\lambda < n$,

$$|\nabla^k \Psi|_{\omega_0} = O(e^{-\lambda t}),$$

where t is the distance to a fixed point in M_G .

Since crepant resolutions have trivial 1-cohomology, the soliton has to be a gradient solution, that is the drift vector field is the gradient of a function (the soliton potential): given the asymptotics obtained in the theorem, this function is asymptotic to the soliton potential of the model Cao soliton given in Section 2.

Such gluing constructions have been performed before in other geometric settings. Early uses of the method include Taubes' construction [Tau92] of anti-self-dual metrics, and Kapouleas' [Kap90, Kap91] of minimal surfaces. More recently the method has been used [Ngu09, Ngu13] to construct self-translators, the mean curvature flow analogue of steady Ricci solitons. Joyce himself [Joy00] used his Calabi-Yau's as building blocks for gluing constructions of manifolds of exceptional holonomy.

In the Kähler setting, Biquard-Minerbe [BM11] used gluing methods to construct noncompact Calabi-Yau manifolds of complex dimension 2, of several different asymptotic behaviours. Gluing techniques have also been used by various authors to construct constant scalar curvature and extremal Kähler metrics, see for example [RS05, AP06, APS11, Szé12, BR15].

We also prove a general uniqueness statement. We show that a steady Kähler-Ricci soliton is unique for its cohomology class, asymptotics (as previously mentioned, nontrivial steady solitons are noncompact), and drift vector field, in the following sense:

Proposition 1.2. *Let (M, ω, X) be a steady Kähler-Ricci soliton. Let $\omega + \frac{i}{2}\partial\bar{\partial}\Psi$ be another steady Kähler-Ricci soliton for the same drift vector field X , such that at infinity we have*

1. $\Psi \rightarrow 0$;
2. $X\Psi \rightarrow 0$;
3. $\partial\bar{\partial}\Psi \rightarrow 0$ (by comparison with ω).

Then $\Psi = 0$.

1.2 Related work

The form of Theorem 1.1 suggests generalization in two directions, and after this article was posted to the arXiv, Conlon-Deruelle [CD20] succeeded in carrying out both these extensions.

First, the orbifolds \mathbb{C}^n/G are only the simplest examples of Calabi-Yau cones. It turns out that equivariant crepant resolutions of arbitrary Calabi-Yau cones admit steady Kähler-Ricci solitons modeled on the Cao soliton, with a subtlety in the asymptotics of the soliton metric depending on

whether the Kähler class is or is not compactly supported (in the case of resolutions of \mathbb{C}^n/G all Kähler classes are compactly supported).

Secondly, a steady Kähler-Ricci soliton should exist on the manifolds M_G , associated to the radial vector field X , in every Kähler class, not just small Kähler classes as constructed in this article. Rather than using gluing machinery, the argument in [CD20] solves the PDE by a continuity method, with the C^0 estimate relying on, among other things, a weighted energy estimate and an adapted Nash-Moser iteration, as well as ideas from pluripotential theory. The alternate construction presented here remains useful in providing the asymptotics of the construction as the Kähler class tends to zero.

1.3 Outline

In Sections 2–5, we first review the construction of the Cao soliton, and then introduce the required analytical tools to handle the analysis in weighted Hölder spaces on the Cao soliton, with an exponential weight at infinity. Finally, for finite quotients of the Cao soliton, we combine this analysis with the weighted analysis at the singular point of the quotient.

In Section 6 we review the material needed on crepant resolutions of \mathbb{C}^n/G , and on the canonical Kähler, Ricci-flat, asymptotically locally Euclidean metrics on these spaces. We also discuss some specific examples of equivariant crepant resolutions.

The proof of uniqueness (Proposition 1.2) is given in Section 7.

Finally, in Sections 8–13, we proceed to the gluing construction, using in particular a blowup analysis (Section 11) to bound the inverse of the linearization of the problem in suitable weighted spaces. Theorem 1.1 is proved in Section 13.

2 $U(n)$ -invariant Kähler metrics and the soliton equation

We begin by recalling Cao’s construction [Cao96] of a steady soliton on \mathbb{C}^n and a one-parameter family of steady solitons on the bundle $\mathcal{O}(-n)$ over $\mathbb{C}P^{n-1}$. We habitually identify the complement of the zero section of $\mathcal{O}(-n)$ with $(\mathbb{C}^n \setminus \{0\})/\mathbb{Z}_n$, to which it is biholomorphic.

All Cao’s solitons are invariant under the natural action of $U(n)$ on finite quotients of \mathbb{C}^n : therefore we start by reviewing this ansatz, see [FIK03].

From Section 4 onwards, only the soliton on \mathbb{C}^n will be used. The family of solitons on $\mathcal{O}(-n)$ is reviewed because it is the motivating example for the solitons on desingularizations of $(\mathbb{C}^n \setminus \{0\})/G$ constructed in the rest of this article (see Remark 3.2).

Let $\Phi : \mathbb{R} \rightarrow \mathbb{R}$, and denote its derivative by $\varphi(t) = \Phi_t(t)$. Throughout this article we use interchangeably the variables $z \in \mathbb{C}^n \setminus \{0\}$, $r \in \mathbb{R}^+$, $t \in \mathbb{R}$, with

$$|z|^2 = r^2 = e^t.$$

We introduce real symmetric 2-tensors on $\mathbb{C}^n \setminus \{0\}$:

$$\begin{aligned} g_{FS} &= g_{S^{2n-1}} - \eta^2, \\ g_{cyl} &= \frac{dr^2}{r^2} + \eta^2 = \frac{1}{4}dt^2 + \eta^2 \end{aligned}$$

where $g_{S^{2n-1}}$ is the round sphere metric, the 1-form η is the connection 1-form on the Hopf bundle $S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$, and g_{FS} is the pull-back of the standard Fubini-Study metric of $\mathbb{C}P^{n-1}$ with holomorphic sectional curvature -4 . We denote by ω_{FS} and ω_{cyl} the associated (1,1)-forms.

The Cao $U(n)$ -invariant Kähler forms on $\mathbb{C}^n \setminus \{0\}$ will be of the form

$$\omega = \frac{i}{2} \partial \bar{\partial} \Phi(t) = \varphi(t) \omega_{FS} + \varphi_t(t) \omega_{cyl}. \quad (2.1)$$

This formula defines a Kähler metric outside the origin if and only if both φ and φ_t are everywhere positive. We henceforth assume that they are. The Ricci form is then

$$\rho(\omega) = i \partial \bar{\partial} \log \frac{\omega^n}{\text{vol}_{\mathbb{C}^n}} = i \partial \bar{\partial} \log (\varphi^{n-1} \varphi_t e^{-nt}) = i \partial \bar{\partial} (\log(\varphi^{n-1} \varphi_t) - nt). \quad (2.2)$$

Fix $\mu \in \mathbb{R}$. Let X be the radial vector field $-2\mu r \partial / \partial r = -4\mu \partial / \partial t$. The pair (ω, X) is a steady Ricci soliton if

$$\text{Ric}(g) + \frac{1}{2} \mathcal{L}_X g = 0,$$

which by (2.2) can be rewritten

$$\varphi^{n-1} \varphi_t e^{\mu \varphi} = cst. e^{nt}.$$

By a translation on t (that is, an homothety of \mathbb{C}^n), one can take the constant to be 1, and the final equation is therefore

$$\varphi^{n-1} \varphi_t e^{\mu \varphi} = e^{nt}. \quad (2.3)$$

Such Ricci solitons are *gradient* Ricci solitons: with the gradient function $f(z) = -\mu \varphi(t)$,

$$\text{Ric}(g_\Phi) + (\nabla^{g_\Phi})^2 f = 0.$$

By multiplying φ by a constant, one can restrict to the case $\mu = 1$, which we will henceforth do.

The equation (2.3) is solved in the following way. Let $F(s)$ be the degree- $(n-1)$ polynomial $F(s) = \sum_{r=0}^{n-1} (-1)^{n-r-1} \frac{(n-1)!}{r!} s^r$. Then one can check that

$$\frac{d}{ds} [F(s)e^s] = s^{n-1} e^s,$$

and for all $s > s_0 \geq 0$,

$$0 < F(s)e^s - F(s_0)e^{s_0} < s^{n-1} e^s. \quad (2.4)$$

It follows that the soliton equation (2.3) can be rewritten as

$$F(\varphi) e^\varphi = \frac{e^{nt}}{n} + cst.$$

We can therefore define a family of solitons parametrized by a nonnegative real number $a \geq 0$ by taking $\varphi_a = \varphi$ to be implicitly defined by the equation

$$F(\varphi(t)) e^{\varphi(t)} = \frac{e^{nt}}{n} + F(a) e^a. \quad (2.5)$$

From (2.4) the function φ is well defined, with $\varphi(t) > a$ for all t ; again from (2.4), one has $\varphi^{n-1} e^\varphi > \frac{e^{nt}}{n}$ so equation (2.3) implies that

$$0 < \varphi_t < n. \quad (2.6)$$

Thus $\Phi := \int \varphi dt$ is the Kähler potential of a steady Ricci soliton with associated holomorphic vector field $X = -2r \partial / \partial r = -4 \partial / \partial t$.

Example 2.1 (Hamilton's cigar soliton). In dimension $n = 1$, the equation (2.5) reduces to $e^{\varphi_a(t)} = e^t + e^a$, and then $(\varphi_a)_t = (1 + e^a e^{-t})^{-1}$ by (2.3). The associated Kähler-Ricci soliton on \mathbb{C} is $(\varphi_a)_t g_{cyl}$, so the dependence in a is just by translation, and we get a unique soliton (Hamilton's cigar soliton),

$$(\varphi_0)_t g_{cyl} = \frac{dx^2 + dy^2}{r^2 + 1}.$$

3 Asymptotics of the Cao potentials

In this section, we establish the asymptotics of the functions $\varphi_a(t)$ constructed in the previous section as $t \rightarrow -\infty$ and as $t \rightarrow \infty$. The former is necessary to determine the topological structure of the metric completion of the solitons. The latter determines the asymptotics at infinity of the Cao solitons.

Lemma 3.1. *As $t \rightarrow -\infty$,*

$$\varphi_a(t) = \begin{cases} a + \frac{1}{n}a^{1-n}e^{-a}e^{nt} + O(e^{2nt}), & a > 0. \\ e^t - \frac{1}{n+1}e^{2t} + O(e^{3t}), & a = 0. \end{cases}$$

Proof. Easy calculation starting from (2.5). □

Since $e^t = r^2$, when $a = 0$, the metric associated to $\varphi = \varphi_0$ on $\mathbb{C}^n \setminus \{0\}$ extends smoothly across the origin (φ_0 has a smooth development in powers of e^t). This is Cao's steady soliton (unique up to translation and rescaling) on \mathbb{C}^n .

When $a > 0$, then $\varphi_t(t) = a^{1-n}e^{-a}e^{nt} + O(e^{2nt})$ with $e^{nt} = r^{2n}$ (and again there is a smooth development in powers of e^{nt}), so formula (2.1) says that the metric associated to $\varphi = \varphi_a$ on the quotient $(\mathbb{C}^n \setminus \{0\})/\mathbb{Z}_n \cong \mathcal{O}(-n) \setminus \mathbb{C}P^{n-1}$ extends smoothly across the zero section $\mathbb{C}P^{n-1}$ to give a smooth metric on $\mathcal{O}(-n)$. This is Cao's family of steady solitons (a one-parameter family, up to translation and rescaling) on $\mathcal{O}(-n)$.

Remark 3.2. As $a \rightarrow 0$, the solitons on $\mathcal{O}(-n)$ converge to the Cao soliton on $\mathbb{C}^n/\mathbb{Z}_n$ with an orbifold singularity at the origin. This is an explicit example of our construction in this paper. It is also interesting to see what 'bubble' occurs in this limit: an easy calculation gives

$$\frac{\varphi_a(t + \log a)}{a} \rightarrow (1 + e^{nt})^{\frac{1}{n}}$$

(the translation in t corresponds to homotheties in \mathbb{C}^n). The limit is the well-known Kähler Ricci-flat metric on the total space of $\mathcal{O}(-n)$ constructed by Calabi [Cal79] (for $n = 2$ this is the Eguchi-Hanson metric).

We now study the behaviour at infinity. The following lemma is again a consequence of equation (2.5). For further terms in the expansion, down to an error of $O\left(\frac{(\log t)^2}{t^2}\right)$, see [CD20, Proposition 3.1].

Lemma 3.3. *As $t \rightarrow \infty$,*

$$\begin{aligned} \varphi(t) &= nt - (n-1)\log t - n\log n + O\left(\frac{\log t}{t}\right). \\ \varphi_t(t) &= n - \frac{n-1}{t} + O\left(\frac{\log t}{t^2}\right). \end{aligned}$$

□

Proof. Let $\varphi_0(t) := nt - (n-1)\log t - n\log n$. For a fixed real number A , consider the function

$$\left\{ \frac{nF\left(\varphi_0(t) + A\frac{\log t}{t}\right) \exp\left(\varphi_0(t) + A\frac{\log t}{t}\right)}{e^{nt}} - 1 - \frac{nF(a)e^a}{e^{nt}} \right\} \frac{t}{\log t}.$$

This function has limit

$$A - \frac{(n-1)^2}{n}$$

as $t \rightarrow \infty$. Therefore, for A sufficiently large and positive, say A_2 , this function is positive for all t sufficiently large, and for A sufficiently small, say A_1 , this function is negative for all t sufficiently large. That is, for all t sufficiently large,

$$F\left(\varphi_0(t) + A_1 \frac{\log t}{t}\right) e^{(\varphi_0(t) + A_1 \frac{\log t}{t})} < \frac{e^{nt}}{n} + F(a)e^a < F\left(\varphi_0(t) + A_2 \frac{\log t}{t}\right) e^{(\varphi_0(t) + A_2 \frac{\log t}{t})}.$$

Since $\frac{e^{nt}}{n} + F(a)e^a = F(\varphi(t))$ (by (2.5)), by the monotonicity of the function $F(s)e^s$ we conclude that for all t sufficiently large

$$\varphi_0(t) + A_1 \frac{\log t}{t} < \varphi(t) < \varphi_0(t) + A_2 \frac{\log t}{t}.$$

This establishes the first statement. For the second, substitute this expansion into equation (2.3). \square

Therefore the metric has the asymptotic behaviour when $t \rightarrow +\infty$

$$g \sim n\left(\frac{1}{4}dt^2 + \eta^2 + tg_{FS}\right) \quad (3.1)$$

and in particular the volume of the ball of radius t is of order t^n , that is half-dimensional; also the injectivity radius is bounded below. More generally:

Lemma 3.4. *The Cao solitons have injectivity radius bounded below and curvature and all their covariant derivatives bounded.*

Proof. It remains to be proved that the Riemannian curvature and its derivatives are bounded. Given the explicit form of the metric, this reduces to the fact that all derivatives of φ are bounded, which is obvious (at infinity $\varphi_t \rightarrow n$ and the higher derivatives go to zero). \square

4 Analysis on the Cao soliton

For the rest of this article, we need only the soliton obtained for $a = 0$; that is, the Cao soliton on \mathbb{C}^n with Kähler form $\omega_0 = \varphi\omega_{FS} + \varphi_t\omega_{cyl}$. In this section we do some analysis on this soliton and on manifolds asymptotic to it.

We first work generally on a Riemannian manifold (M, g) with a single end isometric to the portion $\{T_0 \leq t\}$ of the Cao soliton (\mathbb{C}, ω_0) , or to its quotient by some freely-acting $G \subseteq SU(n)$. Fix $\delta > 0$ and define a weight function

$$w(t) = e^{\delta\varphi}.$$

So $w \geq 1$ and at infinity $w(t) \sim e^{\delta nt}$. We use the Hölder weighted spaces $C_\delta^{k,\alpha}(M) = w^{-1}C^{k,\alpha}(M)$, where $C^{k,\alpha}(M)$ are the usual Hölder spaces.

Proposition 4.1 (Schauder estimate for Cao asymptotics). *Let a be a symmetric positive-definite bivector field, and b a vector field, whose (unweighted) C^α norms are finite. Suppose moreover that for some real $\lambda > 0$, the uniform global bound $a \geq \lambda g^{-1}$ holds. Denote by L the second-order elliptic operator*

$$Lu = \langle a, \nabla^2 u \rangle + \langle b, du \rangle.$$

Then there exists a constant C , such that for all t_0, t_1 with $T_0 + 1 \leq t_0 + 1 \leq t_1 < \infty$, for all u on $\{t < t_1\} \subseteq M$,

$$\|u\|_{C_\delta^{2,\alpha}(\{t < t_0\})} \leq C \left[\|u\|_{C_\delta^0(\{t < t_1\})} + \|Lu\|_{C_\delta^\alpha(\{t < t_1\})} \right].$$

Proof. By Lemma 3.4, there exist $s > 0$ and $Q > 0$ such that for all $x \in M$, there exist (harmonic) co-ordinates on the ball $B_x(s)$ in which the metric tensor g is $C^{1,\alpha}$ controlled by Q . See, for example, [Heb96, Section 1.2]. Therefore on balls of radii $\leq s$, the Schauder norms with respect to the metric g are uniformly equivalent to the Euclidean Schauder norms.

Moreover, by assumption, the coefficients a, b of the operator are uniformly controlled on balls $B_x(s)$. Therefore the Euclidean Schauder estimate [GT01, Theorem 6.2] also holds, with uniform constant, on balls $B_x(s)$ with Riemannian Schauder norm.

The weight function w_δ has the property that, for some uniform constant C , for all x and all $y \in B_x(s)$, $C^{-1}w(x) \leq w(y) \leq Cw(x)$. We may therefore combine the uniform Schauder estimates on balls $B_x(s)$ to give the desired global weighted estimate. \square

We now specialize to the Cao manifold (\mathbb{C}^n, ω_0) itself, with the weight function $w(t) = e^{\delta\varphi}$ itself. Denote by Δ the Laplace-Beltrami operator of ω_0 , and by X (as in Section 2) the radial vector field $-2r\partial/\partial r = -4\partial/\partial t$. The linearization of the soliton equation on a Kähler potential is the operator $\Delta - X$. We have the following C^0 estimates for this operator:

Lemma 4.2. *Let $0 < \delta < 1$.*

1. *For $t_0 \in \mathbb{R}$, and u on $\{t \leq t_0\} \subseteq \mathbb{C}^n$,*

$$\sup_{t \leq t_0} w|u| \leq \max \left[\sup_{t=t_0} w|u|, \frac{1}{4\delta(1-\delta)n} \sup_{t \leq t_0} w|(\Delta - X)u| \right]. \quad (4.1)$$

2. *For $r_0 \in [1, \infty]$, and u on $\{1 \leq r \leq r_0\} \subseteq \mathbb{C}^n$ such that $u|_{\{r=r_0\}} = (if\ r_0 < \infty)$ or $u \in C_\delta^2(\{1 \leq r\})$ (if $r_0 = \infty$),*

$$\sup_{1 \leq r \leq r_0} w|u| \leq \max \left[\sup_{r=1} w|u|, \frac{1}{4\delta(1-\delta)n} \sup_{1 \leq r \leq r_0} w|(\Delta - X)u| \right]. \quad (4.2)$$

Proof. The Laplacian of a radial function is

$$\Delta f = \frac{4}{\varphi^{n-1}\varphi_t} \frac{\partial}{\partial t} \left(\varphi^{n-1} \frac{\partial f}{\partial t} \right).$$

Therefore, using equation (2.3) and $X = -4\frac{\partial}{\partial t}$,

$$(\Delta - X)e^{-\delta\varphi} = -\frac{4\delta}{\varphi^{n-1}\varphi_t} \frac{\partial}{\partial t} (e^{-\delta\varphi} \varphi^{n-1} \varphi_t) + 4\delta\varphi_t$$

$$\begin{aligned}
&= -\frac{4\delta}{e^{nt-\varphi}} \frac{\partial}{\partial t} (e^{nt-(1+\delta)\varphi}) + 4\delta\varphi_t \\
&= -4\delta(n - \delta\varphi_t)e^{-\delta\varphi}.
\end{aligned}$$

Since $0 < \varphi_t < n$ by (2.6), one deduces

$$4\delta(1 - \delta)n < -e^{\delta\varphi}(\Delta - X)e^{-\delta\varphi} < 4\delta n.$$

This estimate enables the use of a barrier argument: if $(\Delta - X)f = g$, then write $f = e^{-\delta\varphi}F$; then

$$(\Delta - X)F + Fe^{\delta\varphi}(\Delta - X)e^{-\delta\varphi} - 2\delta\langle d\varphi, dF \rangle = e^{\delta\varphi}g. \quad (4.3)$$

On a compact domain $t \leq t_0$, the maximum principle applied to (4.3) implies

$$\sup_{\{t \leq t_0\}} e^{\delta\varphi}|f| = \sup_{\{t \leq t_0\}} |F| \leq \max \left[\sup_{\{t=t_0\}} |F| + \frac{1}{4\delta(1 - \delta)n} \sup_{\{t \leq t_0\}} e^{\delta\varphi}|g| \right].$$

This is (4.1). Similarly, if $r_0 < \infty$, then on the compact domain $1 \leq r \leq r_0$, if $u|_{\{r=r_0\}} = 0$, then we obtain (4.2).

Finally, if $r_0 = \infty$, we may select a sequence of compactly-supported functions χ_k , with $0 \leq \chi_k \leq 1$, with $\chi_k \rightarrow 1$ pointwise as $k \rightarrow \infty$, such that for all $u \in C_\delta^2(\{1 \leq r\})$,

$$\lim_{k \rightarrow \infty} \|(\Delta - X)(\chi_k u)\|_{C_\delta^0(\{1 \leq r\})} = \|(\Delta - X)u\|_{C_\delta^0(\{1 \leq r\})}.$$

(Indeed, we may take $\chi_k(t) = \chi(t/k)$, for some fixed cutoff function χ .) We can then extract the noncompact ($r_0 = \infty$) case of (4.2) from the sequence of compact results. \square

Theorem 4.3 (Drift Laplacian on the Cao soliton). *Let $0 < \delta < 1$. Then the operator*

$$\Delta - X : C_\delta^{2,\alpha}(\mathbb{C}^n) \rightarrow C_\delta^\alpha(\mathbb{C}^n)$$

is an isomorphism.

Proof. To prove surjectivity: Let $g \in C_\delta^\alpha(\mathbb{C}^n)$. For each of a sequence of domains $\{t \leq t_i\}$, with $t_i \rightarrow \infty$, let f_i be the solution to the Dirichlet problem $(\Delta - X)f_i = g$, $f_i|_{\{t=t_i\}} = 0$. Such a solution exists since $\Delta - X$ has no zero-th order term. By the estimate (4.1),

$$\sup_{t \leq t_i} w|f_i| \leq \frac{1}{4\delta(1 - \delta)n} \sup_{t \leq t_i} w|(\Delta - X)f_i| = \frac{1}{4\delta(1 - \delta)n} \sup_{t \leq t_i} w|g|.$$

So, by the Schauder estimate Proposition 4.1, for all t_0 and all i such that $t_i \geq t_0 + 1$,

$$\|f_i\|_{C_\delta^{2,\alpha}(\{t \leq t_0\})} \leq C \left[\|f_i\|_{C_\delta^0(\{t \leq t_i\})} + \|(\Delta - X)f_i\|_{C_\delta^\alpha(\{t \leq t_i\})} \right] \leq C \|g\|_{C_\delta^\alpha(\{t \leq t_i\})},$$

where the constant C is independent both of t_0 and of i .

By a diagonal argument, we can extract a subsequence, also denoted f_i , which C^2 -converges on each compact subset. Let f denote the limit. On each compact subset $\{t \leq t_0\}$, we have that

$$\|f\|_{C_\delta^{2,\alpha}(\{t \leq t_0\})} \leq \limsup_{i \rightarrow \infty} \|f_i\|_{C_\delta^{2,\alpha}(\{t \leq t_0\})} \leq C \|g\|_{C_\delta^\alpha}.$$

So $\|f\|_{C_\delta^{2,\alpha}} \leq c\|g\|_{C_\delta^g}$. This proves the surjectivity of $\Delta - X$.

To prove injectivity: suppose that $(\Delta - X)f = 0$ with $f = O(e^{-\delta\varphi})$. Let δ' be such that $0 < \delta' < \delta$. Applying the estimate (4.1) with the parameter δ' ,

$$\sup_{t \leq t_0} e^{\delta'\varphi} |f| \leq \sup_{t=t_0} e^{\delta'\varphi} |f| = e^{(\delta' - \delta)\varphi(t_0)} \sup_{t=t_0} e^{\delta\varphi} |f|.$$

The right hand side tends to 0 as $t_0 \rightarrow \infty$, so $f = 0$. \square

5 Analysis on finite quotients of the Cao soliton

For the rest of this article, fix a finite subgroup $G \subset SU(n)$ acting freely on $\mathbb{C}^n \setminus \{0\}$. We now get an orbifold soliton $(C_G = (\mathbb{C}^n \setminus \{0\})/G, \omega_0)$ with an isolated singularity at the origin.

It is possible to construct weighted orbifold Hölder spaces on C_G , consisting of the functions on C_G whose lifts to $\mathbb{C}^n \setminus \{0\}$ extend to functions in the spaces $C_\delta^{k,\gamma}(\mathbb{C}^n)$ of the previous section. By Theorem 4.3, the operator $\Delta - X$ is an isomorphism between appropriate such weighted-at-infinity orbifold Hölder spaces. However, this result is not sufficient for our gluing procedure.

We need to consider Hölder spaces which are also nontrivially weighted at the origin, with weight functions polynomial in the distance to the origin. We take as the weight function the positive radial function

$$w_0(t) := \begin{cases} e^{\gamma t/2}, & t < 0, \\ e^{\delta\varphi(t)}, & t > 0, \end{cases}$$

(we can smooth this function at $t = 0$ but this is not really needed). So $w_0(t) \sim e^{\delta nt}$ at infinity (the weights of the previous section) and $w_0(t) \sim r^\gamma$ at the origin. If it is necessary to specify the parameters, we will use the notation $w_{0,\gamma,\delta}$ for w_0 .

Choose a positive function $\sigma : C_G \rightarrow \mathbb{R}$ such that at each point x , the injectivity radius at x is greater than $\sigma(x)$. We can take $\sigma = \sigma(t)$ to be a radial function, with

$$\sigma(t) = \begin{cases} cst. e^{t/2} = cst. r, & t \ll 0, \\ cst, & t \gg 0. \end{cases} \quad (5.1)$$

We can now define the weighted Hölder space $C_{\gamma,\delta}^{k,\alpha}(C_G)$ by the norm

$$\|f\|_{C_{\gamma,\delta}^{k,\alpha}} = \sum_{i=0}^k \sup \sigma(x)^i w_0(x) |\nabla^i f| + \sup \sigma(x)^{k+\alpha} w_0(x) \sup_{y \in B_x(\sigma(x))} \frac{|P_{y,x} \nabla^k f(y) - \nabla^k f(x)|}{d(x,y)^\alpha}, \quad (5.2)$$

where $P_{y,x}$ is the parallel transport along the unique minimizing geodesic from y to x . (We use the Riemannian metric ω_0 for all geometric aspects of this construction: $\nabla, |\cdot|, d(\cdot, \cdot), P_{x,y}$.) This weighted Hölder norm is equivalent to the previous section's norm on \mathbb{C}^n , for functions supported on $\{t > 0\}$, and to the norm classically used on cones for functions supported on $\{t < 0\}$.

Before we do this, we note, for future reference, some (essentially classical) facts about $\Delta - X$ when considered as an operator on a bounded portion $\{t \leq t_0\}$ of C_G . Let $C_\gamma^{k,\alpha}(t \leq t_0)$ denote the Hölder space of functions on that portion, with norm the analogue of the formula (5.2) restricting to $x, y \in \{t \leq t_0\}$. Then let $\hat{C}_\gamma^{2,\alpha}(t \leq t_0)$ denotes the set of functions $f \in C_\gamma^{2,\alpha}(t \leq t_0)$ with (Dirichlet) boundary condition $f|_{t=t_0} = 0$.

Lemma 5.1. *For each $\gamma \in (0, 2n - 2)$, the operator*

$$\Delta - X : \hat{C}_{\gamma}^{2,\alpha}(t \leq t_0) \longrightarrow C_{\gamma+2}^{\alpha}(t \leq t_0)$$

is an isomorphism.

Proof. We note that the only critical weights of Δ at the origin in the interval $[0, 2n - 2]$ are 0 (corresponding to the constants) and $2n - 2$ (corresponding to the Green function). This implies that the Laplacian Δ has index zero as a map between the spaces $\hat{C}_{\gamma}^{2,\alpha}(t \leq t_0)$ and $C_{\gamma+2}^{\alpha}(t \leq t_0)$ (its adjoint is itself with the adjoint weight $\gamma' = 2n - 2 - \gamma \in (0, 2n - 2)$, so the operator and its adjoint have the same kernel).

Since $X = -2r \frac{\partial}{\partial r}$ is a compact operator between the spaces in question, this means that $\Delta - X$ has index zero.

To show the injectivity of the operator, observe that the calculation of the critical weights implies that a solution of $(\Delta - X)f = 0$ with $f \in \hat{C}_{\gamma}^{2,\alpha}(t \leq t_0)$ and $0 < \gamma < 2n - 2$ must actually be bounded and have a limit at the origin. By local elliptic regularity it must be smooth (in the orbifold sense of lifting to a smooth function on the G -prequotient $B = B_0(\exp(t_0/2)) \subseteq \mathbb{C}^n$). So we have a solution of $(\Delta - X)f = 0$ in the space $C^{2,\alpha}(B)$, with Dirichlet boundary conditions, so $f = 0$.

Therefore the operator is an isomorphism. \square

We can now give the doubly-weighted isomorphism.

Theorem 5.2 (Drift Laplacian on finite quotients of the Cao soliton). *Let $0 < \delta < 1$ and $0 < \gamma < 2n - 2$. Then the operator*

$$\Delta - X : C_{\gamma,\delta}^{2,\alpha}(C_G) \rightarrow C_{\gamma+2,\delta}^{\alpha}(C_G)$$

is an isomorphism.

Proof. This theorem will be a consequence of Theorem 4.3 (which deals with the analysis at infinity) and of the local theory for polynomial weights in cone points (as in Lemma 5.1).

To show the injectivity of the operator, we follow the same argument as in Lemma 5.1. A solution of $(\Delta - X)f = 0$ with $f \in C_{\gamma,\delta}^{2,\alpha}$ and $0 < \gamma < 2n - 2$ must actually be bounded and have a limit at the origin, and so lift to a solution of $(\Delta - X)f = 0$ in the space $C_{\delta}^{2,\alpha}$. We then find $f = 0$ by Theorem 4.3.

To show the surjectivity of the operator, take $g \in C_{\gamma+2}^{\alpha}(C_G)$. By Lemma 5.1 we find $f_0 \in C_{\gamma}^{2,\alpha}(\{t \leq 0\})$ defined for $t \leq 0$, such that $(\Delta - X)f_0 = g$. Take χ a cutoff function so that $\chi(t) = 0$ for $t \geq 0$ and $\chi(t) = 1$ for $t \leq -1$. Then, using the isomorphism of Theorem 4.3, and the fact that $g - (\Delta - X)(\chi f_0)$ has support included in $\{t \geq -1\}$, we can find $f_1 \in C_{\delta}^{2,\alpha}(C_G) \subseteq C_{\gamma,\delta}^{2,\alpha}(C_G)$ such that

$$(\Delta - X)f_1 = g - (\Delta - X)(\chi f_0).$$

Then $f = \chi f_0 + f_1 \in C_{\gamma,\delta}^{2,\alpha}(C_G)$ is the required solution to $(\Delta - X)f = g$. \square

We also note the appropriate Schauder estimate on these doubly-weighted spaces:

Proposition 5.3 (Schauder estimate on the Cao soliton quotient). *There exists a constant C , such that for all $t_0 \leq \infty$, for all u on $\{t \leq t_0\}$, with also $u|_{\{t=t_0\}} = 0$ if $t_0 < \infty$,*

$$\|u\|_{C_{\gamma,\delta}^{2,\alpha}(\{t < t_0\})} \leq C \left[\|u\|_{C_{\gamma,\delta}^0(\{t < t_0\})} + \|(\Delta - X)u\|_{C_{\gamma+2,\delta}^{\alpha}(\{t < t_0\})} \right].$$

Proof. The proof is similar to that of Proposition 4.1. In this situation we have that there exists $Q > 0$ such that for all $x \in M$, there exist (harmonic) co-ordinates on the ball $B_x(\sigma(x))$ in which the metric tensor g is $C^{1,\alpha}$ controlled by Q : the varying radius $\sigma(x)$ (defined in (5.1)) is needed because the injectivity radius is small near the orbifold point at the origin.

Therefore on balls of radii $\sigma(x)$, the *weighted* Schauder norms with respect to the metric g are uniformly equivalent to the *weighted* Euclidean Schauder norms. So the weighted Euclidean Schauder *boundary* estimate [GT01, Lemma 6.4] also holds, with uniform constant, on balls $B_x(\sigma(x))$ with weighted Riemannian Schauder norm.

Again, the weight functions $w_{\gamma,\delta}$ have the property that, for some uniform constant C , for all x and all $y \in B_x(\sigma(x))$, $C^{-1}w(x) \leq w(y) \leq Cw(x)$. We may therefore combine the uniform Schauder estimates on balls $B_x(\sigma(x))$ to give the desired global weighted estimate; the weight $w_{\gamma+2,\delta}$ appears as the product $\sigma^2 w_{\gamma,\delta}$. \square

6 Joyce metrics on crepant resolutions

We continue to fix a finite subgroup G of $SU(n)$ acting freely on $\mathbb{C}^n \setminus \{0\}$, as in Section 5. For the rest of this article we also fix an equivariant crepant resolution $\pi : J_G \rightarrow \mathbb{C}^n/G$ of the quotient. In this context, *equivariant* means that the \mathbb{C}^* action on \mathbb{C}^n/G extends to an action on J_G . Not all such quotients \mathbb{C}^n/G admit equivariant crepant resolutions, or even crepant resolutions. (By the argument of [CDS19, proof of Lemma 2.13], the existence of an *unique* crepant resolution implies that that resolution is equivariant.) We note some examples which do admit equivariant crepant resolutions:

Example 6.1. In dimension $n = 2$, finite subgroups $G \subseteq SU(2)$ which act freely are classified by Dynkin diagrams, falling into two infinite classes A_k, D_k together with some exceptional examples. The associated orbifolds \mathbb{C}^2/G are the *Kleinian* or *Du Val singularities*, and there is a unique *minimal* resolution of each such singularity, which is equivariant and crepant.

Example 6.2. In dimension $n = 3$, there are some ten infinite classes of such subgroups G . There exists a canonical equivariant crepant resolution of \mathbb{C}^3/G , the *G-orbit Hilbert scheme* of \mathbb{C}^3 , whose existence was conjectured by Nakamura [Nak01] and proved by Bridgeland-King-Reid [BKR01]. There may also exist other crepant resolutions (perhaps not equivariant).

Example 6.3. Toric geometry [DHZ01, CR04] provides numerous examples, in arbitrary dimension, of orbifolds with toric, and therefore equivariant, crepant resolutions.

The equivariance property means that the radial vector field $X = -2R \frac{\partial}{\partial R}$ on \mathbb{C}^n/G extends to J_G , and we will continue to denote this extension by X ; the vector field JX generates the action of $S^1 \subset \mathbb{C}^*$.

We will use the notation R for the radius function on J_G , to distinguish it from the homothetic radius r on the Cao soliton.

As previously for the Cao soliton, we introduce Hölder weighted spaces adapted to the geometry of the manifolds J_G . Consider a positive locally bounded weight w_γ^- , which for $R \geq 2$ satisfies

$$w_\gamma^- = R^\gamma.$$

If there is no ambiguity, we will denote w_γ^- by w^- .

Take any reference metric on J_G which coincides with the flat metric of \mathbb{C}^n/G near infinity. There is a positive function σ such that at each point $\sigma(x)$ is smaller than the injectivity radius at x , and near infinity $\sigma(x) = cst. R$. So w_γ^- and σ^γ are comparable (w_γ^-/σ^γ and σ^γ/w_γ^- are bounded). Then the analogue of formula (5.2), with this reference metric, this σ and this weight w_γ^- , defines a weighted Hölder space $C_\gamma^{k,\alpha}(J_G)$.

We recall the Ricci-flat asymptotically locally Euclidean (ALE) Kähler metrics on J_G , implicit in [TY91] and constructed directly in [Joy00]. These metrics are asymptotic to the flat metric on \mathbb{C}^n/G , up to a term in the weighted space $C_{2n}^\infty(J_G)$.

Theorem 6.4 (Joyce). *Let $\mathfrak{k} \in H^2(J_G, \mathbb{R})$ be a Kähler class on J_G containing an ALE Kähler metric. Then \mathfrak{k} contains a unique ALE Ricci-flat metric ω^- . Moreover, near infinity one has $\omega^- = \frac{i}{2}\partial\bar{\partial}\Phi^-$, with*

$$\Phi^- = R^2 + AR^{2-2n} + \psi, \quad \psi \in C_\gamma^\infty(J_G) \text{ for all } \gamma < 2n - 1.$$

Finally, if J_G is an equivariant resolution, the metric ω^- is S^1 -invariant.

The last statement follows from the uniqueness, since the S^1 action preserves the ALE behaviour at infinity.

Standard analysis on weighted spaces gives immediately:

Proposition 6.5. *For any $\gamma \in (0, 2n - 2)$, the Laplacian is an isomorphism*

$$\Delta : C_\gamma^{2,\alpha}(J_G) \rightarrow C_{\gamma+2}^\alpha(J_G).$$

Proposition 6.6 (Schauder estimate on the Joyce manifold). *There exists a constant C , such that for all u on J_G ,*

$$\|u\|_{C_\gamma^{2,\alpha}(J_G)} \leq C \left[\|u\|_{C_\gamma^0(J_G)} + \|\Delta u\|_{C_{\gamma+2}^\alpha(J_G)} \right].$$

Since $H^1(J_G, \mathbb{R}) = 0$, the S^1 -action generated by JX has a moment map μ^- , that is a function such that

$$d\mu^- = \iota_{JX}\omega^-. \tag{6.1}$$

It follows that

$$\mathcal{L}_X\omega^- = d\iota_X\omega^- = -dJd\mu = -2i\partial\bar{\partial}\mu^-.$$

Since near infinity $\omega^- = \frac{i}{2}\partial\bar{\partial}\Phi^-$, we observe that $d\mu^- = \iota_{JX}\frac{i}{2}\partial\bar{\partial}\Phi^- = -\frac{1}{4}d\Phi^-$. Therefore, up to a constant,

$$\mu^- = -\frac{1}{4}X\Phi^-.$$

7 Generalities on steady Kähler-Ricci solitons

In this section, we digress to prove some general facts about steady Kähler-Ricci solitons, culminating in the uniqueness theorem stated in the Introduction. We work on an arbitrary noncompact complex manifold M .

Lemma 7.1. *1. Suppose that the Kähler forms ω and $\omega + \frac{i}{2}\partial\bar{\partial}\Psi$ are both steady solitons with drift vector field X . Then*

$$0 = i\partial\bar{\partial} \left[-\log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} + \frac{1}{4}X\Psi \right].$$

2. Suppose that ω is Kähler, that μ is such that $\frac{1}{2}\mathcal{L}_X\omega = -i\partial\bar{\partial}\mu$, and that Ω is a global holomorphic n -form. Let $f = -\log \frac{\omega^n}{\Omega \wedge \bar{\Omega}} - \mu$. Then $\omega + \frac{i}{2}\partial\bar{\partial}\Psi$ is a steady soliton with drift vector field X , if and only if

$$0 = i\partial\bar{\partial} \left[-\log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} + \frac{1}{4}X\Psi + f \right].$$

Proof. For the first part, we calculate

$$\begin{aligned} 0 &= \text{Ric}(\omega + \frac{i}{2}\partial\bar{\partial}\Psi) + \frac{1}{2}\mathcal{L}_X(\omega + \frac{i}{2}\partial\bar{\partial}\Psi) - \text{Ric}(\omega) - \frac{1}{2}\mathcal{L}_X(\omega) \\ &= -i\partial\bar{\partial} \log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} + i\partial\bar{\partial}(\frac{1}{4}X\Psi). \end{aligned}$$

For the second,

$$\begin{aligned} \text{Ric}(\omega + \frac{i}{2}\partial\bar{\partial}\Psi) + \frac{1}{2}\mathcal{L}_X(\omega + \frac{i}{2}\partial\bar{\partial}\Psi) &= -i\partial\bar{\partial} \log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\Omega \wedge \bar{\Omega}} - i\partial\bar{\partial}(\mu - \frac{1}{4}X\Psi) \\ &= i\partial\bar{\partial} \left[-\log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} + \frac{1}{4}X\Psi + f \right]. \end{aligned} \quad (7.1)$$

□

We can now prove the uniqueness of the Kähler-Ricci soliton in its Kähler class, as stated in the Introduction. This relies on the well-known strong maximum principle of Hopf [GT01, Theorem 3.5].

Proof of Proposition 1.2. Suppose that the Kähler forms ω and $\omega + \frac{i}{2}\partial\bar{\partial}\Psi$ are both steady solitons, with Ψ , $X\Psi$ and $\partial\bar{\partial}\Psi$ going to 0 at infinity (the last in comparison with ω). Then

$$\log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} - \frac{1}{4}X\Psi$$

tends to 0 at infinity, and by Lemma 7.1

$$\Delta_\omega \left[-\log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} + \frac{1}{4}X\Psi \right] = 0.$$

By the strong maximum principle, Δ_ω -harmonic functions on M either do not attain their extrema, or are constant. So, since the function tends to 0 at infinity, it vanishes.

By the convexity of the logarithmic function,

$$0 = \log \frac{(\omega + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega^n} - \frac{1}{4}X\Psi \leq \frac{1}{4}(\Delta_\omega - X)\Psi.$$

Again by the strong maximum principle, $(\Delta_\omega - X)$ -subharmonic functions either do not attain their maximum, or are constant. Since $\Psi \rightarrow 0$ at infinity, $\Psi \leq 0$ everywhere.

Inverting the roles of ω and $\omega + \frac{i}{2}\partial\bar{\partial}\Psi$ gives the other inequality $\Psi \geq 0$, so $\Psi = 0$. □

8 The approximate solution and the Monge-Ampère equation

We now begin the gluing construction. For each small $\epsilon > 0$, define a diffeomorphism $p_\epsilon(z) = \epsilon^{-1}z$ of \mathbb{C}^n/G , which also extends as a diffeomorphism of the equivariant resolution J_G . Despite the fact that it is the same complex manifold, we wish to distinguish the desingularization of the Cao manifold, denoted M_G , from the ALE space J_G used for this desingularization. In this way, we see p_ϵ as a ‘blowup map’

$$p_\epsilon : M_G \longrightarrow J_G.$$

We will always consider J_G equipped with the Joyce metric ω^- (from Section 6), whereas we want to construct on M_G a Kähler-Ricci soliton desingularizing the Cao soliton ω_0 on $C_G = (\mathbb{C}^n \setminus \{0\})/G$ (from Section 3). Let Φ_0 (on C_G) and Φ_- (on the complement of a compact subset of J_G) be as in those sections, so that $\omega_0 = \frac{i}{2}\partial\bar{\partial}\Phi_0$ and $\omega_- = \frac{i}{2}\partial\bar{\partial}\Phi_-$.

For each small $\epsilon > 0$, we define a reference Kähler metric ω_ϵ on M_G in the following way. Let χ be a cutoff function such that $\chi(r) = 0$ for $r \leq 1$ and $\chi(r) = 1$ for $r \geq 2$, and $\chi_\lambda(r) = \chi(r/\lambda)$. The gluing procedure will occur around the radius

$$r_\epsilon = \epsilon^{\frac{n}{n+1}}.$$

We construct ω_ϵ in the following way:

- on $r \geq 2r_\epsilon$, let $\omega_\epsilon = \omega_0$;
- on $r \leq r_\epsilon$, let $\omega_\epsilon = \epsilon^2 p_\epsilon^* \omega_-$;
- on $r_\epsilon \leq r \leq 2r_\epsilon$, let $\omega_\epsilon = \frac{i}{2}\partial\bar{\partial}\Phi_\epsilon$, where

$$\Phi_\epsilon = \chi_{r_\epsilon}(r)\Phi_0 + (1 - \chi_{r_\epsilon}(r))\epsilon^2 p_\epsilon^* \Phi_-. \quad (8.1)$$

This is well-defined since in the intermediate region, ω_ϵ coincides with ω_0 near $r = 2r_\epsilon$ and with $\epsilon^2 p_\epsilon^* \omega_-$ near $r = r_\epsilon$. It is of course S^1 -invariant. Observe that the function Φ_ϵ defined in (8.1) is actually naturally defined on a larger region, say $r \geq \epsilon$ (corresponding to $R \geq 1$ in J_G), and that Φ_ϵ is a potential for ω_ϵ on this larger region.

To see that the forms ω_ϵ are indeed Kähler for ϵ sufficiently small, note that we have $\Phi_0 - r^2 = O(r^4)$ and $\Phi_-(R) - R^2 = O(R^{2-2n})$, with control on the derivatives: $\nabla^k(\Phi_-(R) - R^2) = O(R^{2-2n-k})$. It follows that, for $r_\epsilon \leq r \leq 2r_\epsilon$, one has

$$\begin{aligned} r^2 - \epsilon^2 p_\epsilon^* \Phi_- &= O(\epsilon^{2n} r^{2-2n}) = O(r_\epsilon^4), \\ \nabla^2(r^2 - \epsilon^2 p_\epsilon^* \Phi_-) &= O(\epsilon^{2n} r^{-2n}) = O(r_\epsilon^2). \end{aligned}$$

Therefore on $r_\epsilon \leq r \leq 2r_\epsilon$ we have

$$\omega_\epsilon - \frac{i}{2}\partial\bar{\partial}r^2 = O(r_\epsilon^2),$$

so for ϵ small enough ω_ϵ is positive, so is a Kähler form everywhere.

Since we also have bounds on all derivatives of Φ_- , we more generally get on the same transition region $r_\epsilon \leq r \leq 2r_\epsilon$ the estimates

$$\nabla^k(\Phi_\epsilon - r^2) = O(r_\epsilon^{4-k}), \quad \nabla^k(\omega_\epsilon - \frac{i}{2}\partial\bar{\partial}r^2) = O(r_\epsilon^{2-k}). \quad (8.2)$$

As in Section 6, there is a global moment map $\mu_\epsilon : M_G \rightarrow \mathbb{R}$, such that $d\mu_\epsilon = \iota_{JX}\omega_\epsilon$. We can fix the constant by deciding that, for $r \geq \epsilon$ one has

$$\mu_\epsilon = -\frac{1}{4}X\Phi_\epsilon = \begin{cases} -\frac{1}{4}X\Phi_0, & r \geq 2r_\epsilon; \\ -\frac{1}{4}X[\epsilon^2 p_\epsilon^* \Phi^-] = \epsilon^2 p_\epsilon^* \mu^-, & \epsilon \leq r \leq r_\epsilon. \end{cases} \quad (8.3)$$

The latter implies that, in fact, $\mu_\epsilon = \epsilon^2 p_\epsilon^* \mu^-$ throughout the region $r \leq r_\epsilon$. One again has $\mathcal{L}_X \omega_\epsilon = -2i\partial\bar{\partial}\mu_\epsilon$.

Let Ω be the global holomorphic n -form on M_G which agrees, on $C_G = \mathbb{C}^n/G$, with the Euclidean n -form. (Such an n -form exists since the resolution $\pi : M_G \rightarrow C_G$ is crepant.) Let

$$f_\epsilon = -\log \frac{\omega_\epsilon^n}{\Omega \wedge \bar{\Omega}} - \mu_\epsilon. \quad (8.4)$$

and observe that by (2.3), f_ϵ vanishes on $r \geq 2r_\epsilon$. Motivated by Lemma 7.1, we now introduce the operator

$$T_\epsilon(\Psi) := \frac{(\omega_\epsilon + \frac{i}{2}\partial\bar{\partial}\Psi)^n}{\omega_\epsilon^n} - e^{\frac{1}{4}X\Psi + f_\epsilon}. \quad (8.5)$$

For the rest of the article we study the operators T_ϵ .

9 Glued function spaces

We define weight functions $w_\epsilon = w_{\epsilon,\gamma,\delta}$ depending on the parameters $\gamma \in (0, 2n-2)$ and $\delta \in (0, 1)$ (we will add the dependence in the parameters, for example $w_{\epsilon,\gamma}$ to specify the dependence in γ , only in case of ambiguity) in the following way:

- for $r \geq r_\epsilon$ the weight w_ϵ coincides with the weight defined on the Cao soliton: $w_\epsilon = w_0$; recall that $w_0(r) = r^\gamma$ near $r = 0$;
- for $r \leq 2r_\epsilon$ the weight w_ϵ coincides up to a constant with the weight for the Joyce metric: $w_\epsilon = \epsilon^\gamma p_\epsilon^* w^-$; since for R large one has $w^-(R) = R^\gamma$, this choices ensures that the two definitions coincide for $r_\epsilon \leq r \leq 2r_\epsilon$.

We define a weighted Hölder space $C_{\epsilon,\gamma,\delta}^{k,\alpha}(M_G)$, “gluing the Hölder spaces of the Cao manifold and of the Joyce manifold,” by the analogue of formula (5.2) for the glued metrics ω_ϵ , glued weights $w_{\epsilon,\gamma,\delta}$, and similarly glued σ_ϵ . For example, the C^0 norm is

$$\|f\|_{C_{\epsilon,\gamma,\delta}^0} = \sup w_{\epsilon,\gamma,\delta} |f|.$$

A perhaps-clarifying remark is that, for cutoff functions χ_λ as in the previous section,

$$\|f\|_{C_{\epsilon,\gamma,\delta}^{k,\alpha}} \sim \|\chi_{r_\epsilon}(r)f\|_{C_{\gamma,\delta}^{k,\alpha}(C_G)} + \epsilon^\gamma \|(p_\epsilon)_*(1 - \chi_{r_\epsilon}(r))f\|_{C_\gamma^{k,\alpha}(J_G)}.$$

(The factor ϵ^γ is exactly the factor required to make the two norms be equivalent on the transition region $r_\epsilon \leq r \leq 2r_\epsilon$.)

Proposition 9.1 (Uniform Schauder estimate). *There exists a constant C , such that for all $\epsilon > 0$ sufficiently small, and all u on M_G ,*

$$\|u\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}(M_G)} \leq C \left[\|u\|_{C_{\epsilon,\gamma,\delta}^0(M_G)} + \|(\Delta_\epsilon - X)u\|_{C_{\epsilon,\gamma+2,\delta}^{0,\alpha}(M_G)} \right].$$

Proof. As for Propositions 4.1 and 5.3. □

10 Error term

We want to control f_ϵ defined by (8.4). We know that f_ϵ vanishes on $r \geq 2r_\epsilon$, so the needed estimate is only on the Joyce manifold. We prove:

Proposition 10.1. *For all $\gamma \geq -4$, δ , and k , there exists C such that for ϵ sufficiently small,*

$$\|f_\epsilon\|_{C_{\epsilon, \gamma+2, \delta}^k} \leq Cr_\epsilon^{4+\gamma}.$$

Corollary 10.2. *Under the same hypotheses, one has*

$$\|T_\epsilon(0)\|_{C_{\epsilon, \gamma+2, \delta}^k} = \|1 - e^{f_\epsilon}\|_{C_{\epsilon, \gamma+2, \delta}^k} \leq Cr_\epsilon^{4+\gamma}.$$

Proof of Corollary 10.2. Observe that the bound in Proposition 10.1 reads as

$$\epsilon^{\gamma+2} \sup_{R \leq 2\epsilon^{-1}r_\epsilon} w_{\gamma+2+k}^- |\nabla^k f_\epsilon|_{\omega^-} \leq C_k \epsilon^{\frac{n}{n+1}(4+\gamma)}$$

that is

$$\sup |\nabla^k f_\epsilon|_{\omega^-} \leq C_k (w_{\gamma+2+k}^-)^{-1} \epsilon^{\frac{2n-2-\gamma}{n+1}}$$

for ϵ small enough. Expanding $\nabla^k(e^{f_\epsilon} - 1)$ in terms of the derivatives of f_ϵ now gives the same bounds on $e^{f_\epsilon} - 1$. \square

Proof of Proposition 10.1. Since f_ϵ vanishes for $r \geq 2r_\epsilon$, we only need to bound f_ϵ for $r \leq 2r_\epsilon$. We will bound independently the two terms $\log \frac{\omega_\epsilon^n}{\Omega \wedge \bar{\Omega}}$ and μ_ϵ .

The first term is nonzero only on the transition region $r_\epsilon \leq r \leq 2r_\epsilon$. From the bounds (8.2), we get that on this region

$$\nabla^k \left(\frac{\omega_\epsilon^n}{\Omega \wedge \bar{\Omega}} \right) = O(r_\epsilon^{2-k}),$$

and so

$$\sup_{r_\epsilon \leq r \leq 2r_\epsilon} r^{2+\gamma+k} |\nabla^k \log \frac{\omega_\epsilon^n}{\Omega \wedge \bar{\Omega}}| \leq c_k r_\epsilon^{4+\gamma}$$

which is exactly the required bound.

By (8.3), the second term μ_ϵ coincides with $-\frac{1}{4}X\Psi_\epsilon$ on the region where the potential Ψ_ϵ exists, that is $r \geq \epsilon$; and with $\epsilon^2 p_\epsilon^* \mu^-$ on $r \leq r_\epsilon$. From the estimate (8.2) we see that on the transition region $r_\epsilon \leq r \leq 2r_\epsilon$ we have

$$\nabla^k \mu_\epsilon = O(r_\epsilon^{2-k}),$$

and so

$$\sup_{r_\epsilon \leq r \leq 2r_\epsilon} r^{2+\gamma+k} |\nabla^k \mu_\epsilon| \leq c_k r_\epsilon^{4+\gamma},$$

the desired estimate. Finally, on $r \leq r_\epsilon$ it is easier to write the required estimate in terms of norms on the manifold J_G :

$$\begin{aligned} \|\mu_\epsilon\|_{C_{\epsilon, \gamma+2, \delta}^k(\{r \leq r_\epsilon\})} &= \epsilon^{\gamma+2} \|(p_\epsilon^{-1})^* \mu_\epsilon\|_{C_{\gamma+2}^k(\{R \leq \epsilon^{-1}r_\epsilon\})} \\ &= \epsilon^{\gamma+4} \|\mu^-\|_{C_{\gamma+2}^k(\{R \leq \epsilon^{-1}r_\epsilon\})} \\ &\leq Cr_\epsilon^{4+\gamma} \end{aligned}$$

so long as $\gamma \geq -4$. \square

Remark 10.3. The choice of r_ϵ was made to get an optimal bound on f_ϵ : the two errors coming from cutting off the Cao soliton, and from gluing the Joyce manifold (which is Ricci-flat, not a soliton), are of the same order around r_ϵ . For example, for $n = 2$ we have $r_\epsilon = \epsilon^{\frac{2}{3}}$. A different choice (such as $r_\epsilon = \epsilon^{\frac{1}{2}}$, which means taking a bigger r_ϵ) would give a larger error term; it would then be necessary to correct the Joyce metric as in [BM11, § 2.4] before gluing, but we do not need to do this with our choice of r_ϵ .

11 First-order term: Blowup argument

The linearization of the operator T_ϵ defined in (8.5) is $\frac{1}{4}(\Delta_\epsilon - e^{f_\epsilon} X)$. We first study the simpler operator $\Delta_\epsilon - X$, then, in Corollary 11.4, use the control on e^{f_ϵ} from Corollary 10.2 to extend the results to $\Delta_\epsilon - e^{f_\epsilon} X$.

Lemma 11.1. *For each γ, δ , and each $\epsilon > 0$, the operator*

$$\Delta_\epsilon - X : C_{\epsilon, \gamma, \delta}^{2, \alpha}(M_G) \rightarrow C_{\epsilon, \gamma+2, \delta}^\alpha(M_G)$$

is continuous.

Proof. We have that $(\Delta_\epsilon - X)u = \langle \omega_\epsilon^{-1}, \nabla^2 u \rangle + \langle -X, du \rangle$, where $\langle \cdot, \cdot \rangle$ denotes a full trace. Since both ω_ϵ^{-1} and X are bounded with respect to the metric ω_ϵ , the operator is continuous. \square

Proposition 11.2. *For each $0 < \delta < 1$, each $0 < \gamma < 2n - 2$, and each $0 < \alpha < 1$, there exists a constant c , such that for all ϵ sufficiently small, for all $r_0 \in [1, \infty]$, and for all $u \in C_{\epsilon, \gamma, \delta}^2(\{r \leq r_0\})$, with the Dirichlet boundary condition $u|_{r=r_0} = 0$ if $r_0 < \infty$,*

$$\|u\|_{C_{\epsilon, \gamma, \delta}^0(r \leq r_0)} \leq c \|(\Delta_\epsilon - X)u\|_{C_{\epsilon, \gamma+2, \delta}^0(r \leq r_0)}.$$

Proof. We prove this by contradiction. Suppose this false; then there exist a sequence $\epsilon_i \rightarrow 0$, a sequence $r_i \in [1, \infty]$, and a sequence of functions u_i defined on $\{r \leq r_i\}$, satisfying $u_i|_{r=r_i} = 0$, with

$$\|(\Delta - X)u_i\|_{C_{\epsilon_i, \gamma+2, \delta}^0} \rightarrow 0, \tag{11.1}$$

but for all i ,

$$\|u_i\|_{C_{\epsilon_i, \gamma, \delta}^0} = 1.$$

The first observation is that by (4.2), together with the fact that for $r \geq 1$ the weight w used for the Cao metric coincides with the weights $w_{\epsilon, \gamma}$, $w_{\epsilon, \gamma+2}$ used to define the norms $C_{\epsilon, \gamma, \delta}^0$ and $C_{\epsilon, \gamma+2, \delta}^0$,

$$\sup_{1 \leq r \leq r_i} w_{\epsilon, \gamma} |u_i| \leq \max \left[\sup_{r=1} w_{\epsilon, \gamma} |u_i|, \frac{1}{4\delta(1-\delta)n} \sup_{1 \leq r \leq r_i} w_{\epsilon, \gamma+2} |(\Delta - X)u_i| \right].$$

Since by hypothesis $\sup w_{\epsilon, \gamma+2} |(\Delta - X)u_i| \rightarrow 0$, we see that there exist points z_i such that:

$$\begin{aligned} |z_i| &\leq 1 \\ w_{\epsilon, \gamma}(z_i) |u_i(z_i)| &= 1. \end{aligned}$$

We will now use a blowup argument at z_i . Up to extracting a subsequence, we can suppose that $r_i \rightarrow r_\infty \in [1, \infty]$. We distinguish three cases:

Case 1: $0 < \inf_i |z_i|$.

We can suppose $z_i \rightarrow z_\infty$. From local elliptic regularity (including at the boundary $r = r_i$ if it is finite), as in [GT01, Theorems 8.32-3], we have local $C^{1,\alpha'}$ bounds on u_i , so we can extract a subsequence $C^{1,\alpha}$ -converging on each compact subset of the domain $\{r < r_\infty\}$ of the Cao manifold C_G . Let u_∞ be its limit; since the weights w_ϵ converge to the weight w_0 for C_G , we get

$$\begin{aligned} (\Delta - X)u_\infty &= 0 \text{ (and } u_\infty \text{ is smooth),} \\ \sup w_0|u_\infty| &= w_0(z_\infty)|u_\infty(z_\infty)| = 1, \\ u_\infty|_{\{r=r_\infty\}} &= 0 \text{ if } r_\infty < \infty. \end{aligned}$$

Proposition 5.3, the Schauder estimate on the Cao manifold C_G , gives that

$$\|u_\infty\|_{C_{\gamma,\delta}^{2,\alpha}(\{r \leq r_\infty\})} \leq C \left[\|u_\infty\|_{C_{\gamma,\delta}^0(\{r \leq r_\infty\})} + \|(\Delta - X)u_\infty\|_{C_{\gamma+2,\delta}^\alpha(\{r \leq r_\infty\})} \right] = C \sup_{\{r \leq r_\infty\}} w_0|u_\infty| < \infty.$$

So $u_\infty \in C_{\gamma,\delta}^{2,\alpha}(\{r \leq r_\infty\})$ is a nonzero element of the kernel of $\Delta - X$. If $r_\infty = \infty$, this contradicts Theorem 5.2. If $r_\infty < \infty$, noting that u_∞ also satisfies the Dirichlet boundary condition, this contradicts Lemma 5.1.

Case 2: $\sup_i \epsilon_i^{-1}|z_i| < \infty$.

Let $\zeta_i := p_{\epsilon_i}(z_i) \in J_G$; we can suppose that $\zeta_i \rightarrow \zeta_\infty \in J_G$.

Let $v_i := \epsilon_i^\gamma(p_{\epsilon_i})_* u_i$. By definition the weight $w_{\epsilon_i,\gamma,\delta}$ is equal to the Joyce weight $\epsilon_i^\gamma p_{\epsilon_i}^* w_\gamma^-$ on the larger and larger domains $\{R \leq \epsilon^{-1}r_\epsilon\}$ of J_G , so

$$\sup_{\{R \leq \epsilon^{-1}r_\epsilon\}} w_\gamma^- |v_i| = w_\gamma^-(\zeta_i) |v_i(\zeta_i)| = 1. \quad (11.2)$$

On the other hand, since $\epsilon^{-2}\omega_\epsilon = \omega^-$ for $R \leq \epsilon^{-1}r_\epsilon$, the control (11.1) translates into

$$\sup_{\{R \leq \epsilon^{-1}r_\epsilon\}} w_{\gamma+2}^- |(\Delta_{\omega^-} - \epsilon^2 X)v_i| = \sup_{\{r \leq r_\epsilon\}} w_{\epsilon,\gamma+2,\delta} |(\Delta_{\omega_\epsilon} - X)u_i| \longrightarrow 0. \quad (11.3)$$

The operators $\Delta_{\omega^-} - \epsilon^2 X$ are uniformly locally controlled, so by local elliptic regularity [GT01, Theorems 8.32-3], as in Case 1, we can extract a subsequence $C^{1,\alpha}$ -converging on each compact domain of J_G . Let v_∞ be its limit. By (11.2) and (11.3),

$$\Delta_{\omega^-} v_\infty = 0, \quad \sup w_\gamma^- |v_\infty| = 1.$$

We conclude as before, using Proposition 6.6 (the Schauder estimate on the Joyce manifold) and Proposition 6.5 (the proof of isomorphism).

Case 3: We may pass to a subsequence with $z_i \rightarrow 0$ and $\epsilon_i^{-1}|z_i| \rightarrow \infty$.

In this case we perform a similar extraction on the cone \mathbb{C}^n/G itself, by scaling by $\lambda_i = |z_i|$. As in Case 2, we consider a sequence of homotheties which restrict the points to a compact region. In this case we take the homotheties p_{λ_i} , and introduce rescaled functions $v_i := \lambda_i^\gamma (p_{\lambda_i})_* u_i$. We can suppose $\zeta_i = \frac{z_i}{\lambda_i} \rightarrow \zeta_\infty \in \mathbb{C}^n/G$. Finally, now $\lambda_i^{-2}(p_{\lambda_i})_* \omega_{\epsilon_i}$ converges to the flat cone metric on \mathbb{C}^n/G , and the weights $\lambda_i^\gamma p_{\lambda_i}^* w_{\epsilon_i,\gamma,\delta}$ converge to the cone weight $w_\gamma(R) = R^\gamma$, in both instances because $1 \gg \lambda_i \gg \epsilon_i$.

The next part of the proof is unchanged from Case 2: the operators $\Delta_{\lambda_i^{-2}(p_{\lambda_i})_*\omega_{\epsilon_i}} - \lambda_i^2 X$ are uniformly locally controlled, and we get a nonzero solution v_∞ of $\Delta v_\infty = 0$ on the whole flat cone \mathbb{C}^n/G , with $\sup_{\mathbb{C}^n/G} R^\gamma |v_\infty| < \infty$.

Finally, since $\gamma \in (0, 2n - 2)$ is not a critical weight of the Laplacian, $\Delta : C_{\gamma}^{2,\alpha}(\mathbb{C}^n/G) \rightarrow C_{\gamma+2}^{\alpha}(\mathbb{C}^n/G)$ is an isomorphism and such a v_∞ cannot exist. \square

Proposition 11.3. *For each $0 < \delta < 1$, each $0 < \gamma < 2n - 2$, and each $0 < \alpha < 1$, there exists a constant c , such that for all ϵ sufficiently small, the operator $\Delta_\epsilon - X : C_{\epsilon,\gamma,\delta}^{2,\alpha}(M_G) \rightarrow C_{\epsilon,\gamma+2,\delta}^{\alpha}(M_G)$ is an isomorphism, and for all $u \in C_{\epsilon,\gamma,\delta}^{2,\alpha}(M_G)$,*

$$\|u\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} \leq c \|(\Delta_\epsilon - X)u\|_{C_{\epsilon,\gamma+2,\delta}^{0,\alpha}}. \quad (11.4)$$

Proof. First observe that the estimate (11.4), and the injectivity of the operator, follow from the C^0 estimate in Proposition 11.2, combined with the uniform Schauder estimate Proposition 9.1.

We now show the surjectivity of $\Delta - X$. This is the same argument as in Theorem 4.3. We want to solve $(\Delta - X)u = v$. On each of a sequence of bounded domains $\{t \leq t_i\}$, one can solve $(\Delta - X)u_i = v$ with $u_i|_{r=r_i} = 0$, since the operator $\Delta - X$, being a compact perturbation of Δ , has index zero for the Dirichlet problem, and is injective (by the maximum principle).

By Proposition 11.2, together with the Schauder estimate Proposition 4.1, we have that for all t_0 and all i such that $t_i \geq t_0 + 1$,

$$\|u_i\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}(\{t \leq t_0\})} \leq C \|v\|_{C_{\epsilon,\gamma+2,\delta}^{\alpha}(M_G)},$$

where the constant C is independent both of t_0 and of i . Therefore when $i \rightarrow \infty$ we can extract a C_{loc}^2 -limit u still satisfying $\|u\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} \leq c \|v\|_{C_{\epsilon,\gamma+2,\delta}^{\alpha}}$, and solving $(\Delta - X)u = v$. \square

Corollary 11.4. *Proposition 11.3 is also valid for the family of operators $\Delta_\epsilon - e^{f_\epsilon} X$.*

Proof. It suffices to check that the norm of the operator

$$(e^{f_\epsilon} - 1)X : C_{\epsilon,\gamma,\delta}^{2,\alpha}(M_G) \longrightarrow C_{\epsilon,\gamma+2,\delta}^{\alpha}(M_G)$$

goes to zero, so that adding it to $\Delta_\epsilon - X$ does not perturb the invertibility and the estimate on the inverse. This follows easily from Corollary 10.2. \square

12 Second-order term

Let us write

$$T_\epsilon(\Psi) = T_\epsilon(0) + L_\epsilon(\Psi) + Q_\epsilon(\Psi),$$

where $L_\epsilon = \frac{1}{4}(\Delta_\epsilon - e^{f_\epsilon} X)$ is the linearization of T_ϵ at the origin.

Proposition 12.1. *There exists constants $c, c' > 0$ such that if $\|\Psi\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}}, \|\Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} \leq c\epsilon^{2+\gamma}$, then*

$$\|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C_{\epsilon,\gamma+2,\delta}^{\alpha}} \leq c'\epsilon^{-2-\gamma} \|\psi - \psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} (\|\psi\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} + \|\psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}}).$$

Proof. Such an estimate is obvious on a ball, without weights, if the geometry of the metric is uniformly controlled: it comes down to saying that the second derivative D^2T remains bounded on a ball.

We can apply this remark on balls on the Cao part of the manifold ($r \geq 1$). The weighted norm on balls is just the usual Hölder norm, multiplied by the factor w_ϵ :

$$\|u\|_{C_{\epsilon,\gamma,\delta}^{k,\alpha}(B(x_0,1))} \sim w_\epsilon(x_0) \|u\|_{C^{k,\alpha}(B(x_0,1))}.$$

Then, for $\|\Psi\|_{C^{2,\alpha}(B(x_0,1))}, \|\Psi'\|_{C^{2,\alpha}(B(x_0,1))} \leq c$ we have the standard estimate

$$\|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C^\alpha(B(x_0,1))} \leq c' \|\Psi - \Psi'\|_{C^{2,\alpha}(B(x_0,1))} (\|\Psi\|_{C^{2,\alpha}(B(x_0,1))} + \|\Psi'\|_{C^{2,\alpha}(B(x_0,1))}).$$

Therefore,

$$\begin{aligned} \|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C_{\epsilon,\gamma+2,\delta}^\alpha(B(x_0,1))} \\ \leq c' w_\epsilon(x_0)^{-1} \|\Psi - \Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}(B(x_0,1))} (\|\Psi\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}(B(x_0,1))} + \|\Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}(B(x_0,1))}) \end{aligned}$$

which gives the required estimate since $w_\epsilon(x_0) \geq 1$.

The situation is quite different on the part where we glued the Joyce manifold, because the injectivity radius and the weight become small. First observe that

$$T_\epsilon(\Psi) = \frac{(\frac{\omega_\epsilon}{\epsilon^2} + \frac{i}{2} \partial \bar{\partial} \frac{\Psi}{\epsilon^2})^n}{(\frac{\omega_\epsilon}{\epsilon^2})^n} - e^{f_\epsilon + \frac{\epsilon^2}{4} X \frac{\Psi}{\epsilon^2}},$$

and recall that on the Joyce region $r \leq r_\epsilon$, $\frac{\omega_\epsilon}{\epsilon^2} = \omega_-$, the metric on J_G , and $f_\epsilon = -\epsilon^2 p_\epsilon^* \mu^-$. Therefore on that region

$$T_\epsilon(\Psi) = \frac{(\omega_- + \frac{i}{2} \partial \bar{\partial} \frac{\Psi}{\epsilon^2})^n}{\omega_-^n} - e^{\epsilon^2[-\mu^- + \frac{1}{4} X \frac{\Psi}{\epsilon^2}]}.$$

On a large compact part $K \subset J_G$, we have as above, for norms taken with respect to ω_- ,

$$\|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C^\alpha} \leq c' \left\| \frac{\Psi}{\epsilon^2} - \frac{\Psi'}{\epsilon^2} \right\|_{C^{2,\alpha}} \left(\left\| \frac{\Psi}{\epsilon^2} \right\|_{C^{2,\alpha}} + \left\| \frac{\Psi'}{\epsilon^2} \right\|_{C^{2,\alpha}} \right)$$

for $\|\frac{\Psi}{\epsilon^2}\|_{C^{2,\alpha}}, \|\frac{\Psi'}{\epsilon^2}\|_{C^{2,\alpha}} \leq c$. But on K we have

$$\|u\|_{C_{\epsilon,\gamma,\delta}^{k,\alpha}} \sim \epsilon^\gamma \|u\|_{C^{k,\alpha}(\omega_-)}.$$

So we deduce that for $\|\Psi\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}}, \|\Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} \leq c\epsilon^{2+\gamma}$, we have on K the estimate

$$\|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C_{\epsilon,\gamma+2,\delta}^\alpha} \leq c' \epsilon^{-2-\gamma} \|\Psi - \Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} (\|\Psi\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}} + \|\Psi'\|_{C_{\epsilon,\gamma,\delta}^{2,\alpha}}).$$

This is again the required estimate.

The case of the conical region between K and $\{r \geq 1\}$ is similar and we will not write the details. The worst constants are obtained on the K part of the manifold, because this is the part of the manifold with smallest injectivity radius and weight. This explains the constants in the statement of the Proposition. \square

13 Inverse function theorem

The proof of Theorem 1.1 will now follow directly from the following quantitative version of the inverse function theorem, previously used in [BM11]:

Lemma 13.1. *Let $T : E \rightarrow F$ be a smooth map between Banach spaces and define $Q := T - T(0) - DT|_0$. Suppose that there are positive constants q , r_0 and c , such that*

1. $\|Q(x) - Q(y)\| \leq q\|x - y\|(\|x\| + \|y\|)$ for every x and y in $B_E(0, r_0)$;
2. $DT|_0$ is an isomorphism with inverse bounded by c ;
3. $\|T(0)\| < \frac{1}{2c} \min(r_0, \frac{1}{2qc})$.

Then the equation $T(x) = 0$ admits a unique solution x in $B_E(0, \min(r_0, \frac{1}{2qc}))$.

Proof of Theorem 1.1. For fixed $0 < \delta < 1$ and $0 < \gamma < 2n - 2$, consider the family of operators

$$T_\epsilon : C_{\epsilon, \gamma, \delta}^{2, \alpha}(M_G) \rightarrow C_{\epsilon, \gamma+2, \delta}^{0, \alpha}(M_G).$$

We will show that for ϵ sufficiently small, the operator T_ϵ satisfies the hypotheses of Lemma 13.1.

By Corollary 10.2, for ϵ sufficiently small

$$\|T_\epsilon(0)\|_{C_{\epsilon, \gamma+2, \delta}^{0, \alpha}(M_G)} = \|1 - e^{f_\epsilon}\|_{C_{\epsilon, \gamma+2, \delta}^{0, \alpha}(M_G)} \leq C\epsilon^{(4+\gamma)\frac{n}{n+1}}.$$

By Corollary 11.4, for ϵ sufficiently small

$$L_\epsilon = \frac{1}{4}[\Delta_a - e^{f_a}X] : C_{\epsilon, \gamma, \delta}^{2, \alpha}(M_G) \rightarrow C_{\epsilon, \gamma+2, \delta}^{0, \alpha}(M_G)$$

is an isomorphism, and its inverse is bounded by a constant independent of ϵ .

By Proposition 12.1, for ϵ sufficiently small, for all Ψ, Ψ' with $\|\Psi\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}}, \|\Psi'\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}} \leq C\epsilon^{\gamma+2}$ the map $Q_\epsilon := T_\epsilon - T_\epsilon(0) - L_\epsilon$ satisfies,

$$\|Q_\epsilon(\Psi) - Q_\epsilon(\Psi')\|_{C_{\epsilon, \gamma+2, \delta}^{0, \alpha}} \leq C\epsilon^{-(\gamma+2)}\|\Psi - \Psi'\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}} (\|\Psi\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}} + \|\Psi'\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}}).$$

Thus if we take $q := C\epsilon^{-(\gamma+2)}$ and $r_0 := C\epsilon^{\gamma+2}$, then (1) and (2) are satisfied, and moreover

$$\begin{aligned} \frac{1}{2c} \min(r_0, \frac{1}{2qc}) &\geq C^{-1}\epsilon^{\gamma+2} \\ &\geq C^{-1}\epsilon^{-\frac{2n-2-\gamma}{n+1}}\|T_a(0)\|. \end{aligned}$$

Since $\gamma < 2n - 2$, for ϵ sufficiently small we then have that $C^{-1}\epsilon^{-\frac{2n-2-\gamma}{n+1}} \geq 1$, and so (3) is satisfied for ϵ sufficiently small.

Therefore by Lemma 13.1 there exists a solution Ψ_ϵ to the equation $T_\epsilon(\Psi_\epsilon) = 0$, i.e.,

$$(\omega_\epsilon + \frac{i}{2}\partial\bar{\partial}\Psi_\epsilon)^n = e^{(X\Psi_\epsilon)/4+f_\epsilon}\Omega \wedge \bar{\Omega}.$$

The solution Ψ_ϵ satisfies, $\|\Psi_\epsilon\|_{C_{\epsilon, \gamma, \delta}^{2, \alpha}(M_G)} \leq C\epsilon^{2+\gamma}$.

Since $\Psi \in C_{\epsilon, \gamma, \delta}^{2, \alpha}(M_G)$ is a solution to the equation $T_\epsilon(\Psi) = 0$, its differential $d\Psi$ is a weak solution to a quasilinear elliptic equation

$$0 = dT_\epsilon(\Psi) = \Delta_{\omega_\epsilon + \frac{i}{2}\partial\bar{\partial}\Psi} d\Psi + F(\nabla^2\Psi, d\Psi).$$

Bootstrapping using the Schauder estimate Proposition 4.1, we conclude that $\Psi \in C_{\epsilon, \gamma, \delta}^{k, \alpha}(M_G)$ for all k . In particular, by Lemma 7.1, $\omega_\epsilon + \frac{i}{2}\partial\bar{\partial}\Psi$ is a steady Kähler-Ricci soliton. \square

References

- [AIK15] Sigurd B. Angenent, James Isenberg, and Dan Knopf. Degenerate neckpinches in Ricci flow. *J. Reine Angew. Math.*, 709:81–117, 2015.
- [AP06] Claudio Arezzo and Frank Pacard. Blowing up and desingularizing constant scalar curvature Kähler manifolds. *Acta Math.*, 196(2):179–228, 2006.
- [APS11] Claudio Arezzo, Frank Pacard, and Michael Singer. Extremal metrics on blowups. *Duke Math. J.*, 157(1):1–51, 2011.
- [BKR01] Tom Bridgeland, Alastair King, and Miles Reid. The McKay correspondence as an equivalence of derived categories. *J. Amer. Math. Soc.*, 14(3):535–554, 2001.
- [BM11] Olivier Biquard and Vincent Minerbe. A Kummer construction for gravitational instantons. *Comm. Math. Phys.*, 308(3):773–794, 2011.
- [BR15] Olivier Biquard and Yann Rollin. Smoothing singular constant scalar curvature Kähler surfaces and minimal Lagrangians. *Adv. Math.*, 285:980–1024, 2015.
- [Cal79] E. Calabi. Métriques kählériennes et fibrés holomorphes. *Ann. Sci. École Norm. Sup. (4)*, 12(2):269–294, 1979.
- [Cao96] Huai-Dong Cao. Existence of gradient Kähler-Ricci solitons. In *Elliptic and parabolic methods in geometry (Minneapolis, MN, 1994)*, pages 1–16. A K Peters, Wellesley, MA, 1996.
- [CD20] Ronan Conlon and Alix Deruelle. Steady gradient Kähler-Ricci solitons on crepant resolutions of Calabi-Yau cones. arXiv:2006.03100, June 2020.
- [CDS19] Ronan Conlon, Alix Deruelle, and Song Sun. Classification results for expanding and shrinking gradient Kähler-Ricci solitons. arXiv:1904.00147, 2019.
- [CR04] Li Chiang and Shi-Shyr Roan. On hypersurface quotient singularities of dimension 4. *Int. J. Math. Math. Sci.*, (45-48):2547–2581, 2004.
- [DdP07] Panagiota Daskalopoulos and Manuel del Pino. Type II collapsing of maximal solutions to the Ricci flow in \mathbb{R}^2 . *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 24(6):851–874, 2007.
- [DHZ01] Dimitrios I. Dais, Christian Haase, and Günter M. Ziegler. All toric local complete intersection singularities admit projective crepant resolutions. *Tohoku Math. J. (2)*, 53(1):95–107, 2001.
- [DS10] Panagiota Daskalopoulos and Natasa Sesum. Type II extinction profile of maximal solutions to the Ricci flow in \mathbb{R}^2 . *J. Geom. Anal.*, 20(3):565–591, 2010.
- [DW11] Andrew S. Dancer and McKenzie Y. Wang. On Ricci solitons of cohomogeneity one. *Ann. Global Anal. Geom.*, 39(3):259–292, 2011.
- [FIK03] Mikhail Feldman, Tom Ilmanen, and Dan Knopf. Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons. *J. Differential Geom.*, 65(2):169–209, 2003.

- [GT01] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. Reprint of the 1998 edition.
- [GZ08] Hui-Ling Gu and Xi-Ping Zhu. The existence of type II singularities for the Ricci flow on S^{n+1} . *Comm. Anal. Geom.*, 16(3):467–494, 2008.
- [Ham88] Richard S. Hamilton. The Ricci flow on surfaces. In *Mathematics and general relativity (Santa Cruz, CA, 1986)*, volume 71 of *Contemp. Math.*, pages 237–262. Amer. Math. Soc., Providence, RI, 1988.
- [Has11] Robert Haslhofer. A renormalized Perelman-functional and a lower bound for the ADM-mass. *J. Geom. Phys.*, 61(11):2162–2167, 2011.
- [Heb96] Emmanuel Hebey. *Sobolev spaces on Riemannian manifolds*, volume 1635 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1996.
- [Ive93] Thomas Ivey. Ricci solitons on compact three-manifolds. *Differential Geom. Appl.*, 3(4):301–307, 1993.
- [Joy00] Dominic D. Joyce. *Compact manifolds with special holonomy*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000.
- [Kap90] Nicolaos Kapouleas. Complete constant mean curvature surfaces in Euclidean three-space. *Ann. of Math. (2)*, 131(2):239–330, 1990.
- [Kap91] Nicolaos Kapouleas. Compact constant mean curvature surfaces in Euclidean three-space. *J. Differential Geom.*, 33(3):683–715, 1991.
- [Nak01] Iku Nakamura. Hilbert schemes of abelian group orbits. *J. Algebraic Geom.*, 10(4):757–779, 2001.
- [Ngu09] Xuan Hien Nguyen. Translating tridents. *Comm. Partial Differential Equations*, 34(1-3):257–280, 2009.
- [Ngu13] Xuan Hien Nguyen. Complete embedded self-translating surfaces under mean curvature flow. *J. Geom. Anal.*, 23(3):1379–1426, 2013.
- [RS05] Yann Rollin and Michael Singer. Non-minimal scalar-flat Kähler surfaces and parabolic stability. *Invent. Math.*, 162(2):235–270, 2005.
- [Szé12] Gábor Székelyhidi. On blowing up extremal Kähler manifolds. *Duke Math. J.*, 161(8):1411–1453, 2012.
- [Tau92] Clifford Henry Taubes. The existence of anti-self-dual conformal structures. *J. Differential Geom.*, 36(1):163–253, 1992.
- [TY91] Gang Tian and Shing-Tung Yau. Complete Kähler manifolds with zero Ricci curvature. II. *Invent. Math.*, 106(1):27–60, 1991.
- [Yan12] Bo Yang. A characterization of noncompact Koiso-type solitons. *Internat. J. Math.*, 23(5):1250054, 13, 2012.

SORBONNE UNIVERSITÉ AND UNIVERSITÉ PARIS CITÉ, CNRS, IMJ-PRG, F-75005 PARIS
 FORDHAM UNIVERSITY