



# **Bracoviruses, ichnoviruses, and virus-like particles from parasitoid wasps retain many features of their virus ancestors**

Jean-Michel Drezen, Annie Bézier, Gaelen R Burke, Michael R Strand

## **► To cite this version:**

Jean-Michel Drezen, Annie Bézier, Gaelen R Burke, Michael R Strand. Bracoviruses, ichnoviruses, and virus-like particles from parasitoid wasps retain many features of their virus ancestors. *Current Opinion in Insect Science*, 2022, 49, pp.93-100. <10.1016/j.cois.2021.12.003>. <hal-03844933>

**HAL Id: hal-03844933**

**<https://hal.science/hal-03844933v1>**

Submitted on 9 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# **Bracoviruses, ichnoviruses, and virus-like particles from parasitoid wasps retain many features of their virus ancestors**

**Jean-Michel Drezen<sup>1</sup>, Annie Bézier<sup>1</sup>, Gaelen R Burke<sup>2</sup> and Michael R Strand<sup>2</sup>**

## **Addresses**

<sup>1</sup>Institut de Recherche sur la Biologie de l'Insecte, UMR7261 CNRS - Université de Tours,  
Tours, France

<sup>2</sup>Department of Entomology, University of Georgia, Athens, GA 30602, USA

Corresponding authors: Drezen, Jean-Michel ([drezen@univ-tours.fr](mailto:drezen@univ-tours.fr)), Strand, Michael R  
([mrstrand@uga.edu](mailto:mrstrand@uga.edu))

## **Highlights**

- Many viruses can integrate into the genomes of hosts.
- BVs, IVs, and VLPs evolved from viruses that integrated into parasitoid wasps.
- BVs and IVs have evolved similar functions through evolutionary convergence.
- BVs and IVs are domesticated endogenous viruses (DEVs).
- VLPs could also be considered as DEVs.

## **Abstract**

Animal genomes commonly contain genes or sequences that have been acquired from different types of viruses. The vast majority of these endogenous virus elements (EVEs) are inactive or consist of only a small number of components that show no evidence of cooption for new functions or interaction. Unlike most EVEs, bracoviruses (BVs), ichnoviruses (IVs) and virus-like particles (VLPs) in parasitoid wasps have evolved through retention and interaction of many genes from virus ancestors. Here, we discuss current understanding of BV, IV and VLP evolution along with associated implications for what constitutes a virus. We suggest that BVs and IVs are domesticated endogenous viruses (DEVs) that differ in several important ways from other known EVEs.

## **Introduction**

Viruses are often viewed as obligate, intracellular parasites because they depend on the biosynthetic machinery of a host cell to produce genome-containing particles (virions). In turn, the function of virions is usually to horizontally transmit the entire viral genome by infecting other host cells that support the production of progeny virus. Studies dating to the 1970s discovered that parasitoid wasps in two families, the Braconidae and Ichneumonidae, produce massive quantities of DNA-containing virions in calyx cells that are located in the ovaries of females [1]. The virions produced in braconids were thereafter named bracoviruses (BVs), while the virions produced in ichneumonids were named ichnoviruses (IVs) [2]. Together, BVs and IVs were also referred to as polydnviruses [2]. Subsequent work showed that BVs and IVs evolved from different viruses that integrated into the germline of wasps [3<sup>\*\*</sup>,4<sup>\*\*</sup>]. However, BVs and IVs have been similarly repurposed through convergence to produce virions that horizontally transfer virulence genes to the hosts that female wasps parasitize, whereas the viral genome components that produce virions in calyx cells remain integrated. The virulence genes virions deliver further have functions in hosts that enable wasp offspring to successfully develop [3<sup>\*\*</sup>,4<sup>\*\*</sup>]. BVs and IVs thus differ from most viruses in that virions do not transmit all viral genome components to host cells they infect, but are nonetheless essential

for vertical transmission of all the viral genome components that are required to produce virions in wasps. Integration of viruses related to the BV ancestor into two other parasitoids have further resulted in ovary calyx cells producing virus-like particles (VLPs) that also have known or likely functions in parasitism [3<sup>\*\*</sup>,4<sup>\*\*</sup>]. In this review, we summarize findings indicating that many types of viruses can be vertically transmitted. We then discuss key features of BVs, IVs and VLPs, the challenges associated with naming these entities, and future study directions.

### **Viruses commonly integrate into the genome of hosts**

Many viruses are only transmitted horizontally while others have evolved latent phases that enable long-term persistence and vertical transmission. Well-known examples of viruses that are both horizontally and vertically transmitted include many large DNA viruses (phages) that infect bacteria, retroviruses that infect vertebrates, and Herpes virus-6 (HHV-6) that infects humans [5-7]. DNA fragments from many types of viruses have also been identified in the genomes of different animals [8]. These fragments, referred to as endogenous virus elements (EVEs) [8], may derive from all or portions of a viral genome integrating into the germline of a host. Most EVEs are rapidly inactivated and decay under neutral evolution. However, a few genes from retroviruses have been coopted by mammals and insects for new functions [9,10] while some EVEs have known or hypothesized roles in protecting hosts against infection from related viruses [11].

In the case of insects, large DNA viruses with circular genomes in the family *Baculoviridae* produce covert infections through unknown mechanisms, while the circular genome of one member of the family *Nudiviridae*, which is closely related to baculoviruses, has been shown to both integrate and remain circular as an episome in the nuclei of persistently-infected host cells [12]. EVEs from a diversity of RNA and DNA viruses have also been identified in insect genomes. Among these are nudivirus-derived EVEs identified in a planthopper (*Nilaparvata lugens*) [13], a chalcidoid wasp (*Eurytoma brunniventris*) [14], and two aphids (*Aphis glycines*, *Melanaphis sacchari*) [15]. A survey of 240 insect genomes further identified 359 genes of

nudivirus origin in 43 other species [16]. The parasitoid *Leptopilina boulardi* is infected by an unclassified virus currently named L. boulardi Filamentous Virus (LbFV) [17]. Recognizable homologs of LbFV genes have been identified in the genomes of several parasitoid wasp species [18\*,19\*]. Similar to retroviruses, nudiviruses and LbFV-like viruses may be predisposed to integrating into the germline of insects due to life cycle features. The majority of EVEs identified in insects are also inactive or consist of one or a few genes that show no evidence of cooption or interaction [20]. Known exceptions are: 1) BVs that evolved from a nudivirus [21\*\*,22\*\*], 2) IVs that evolved from a still unknown virus ancestor [19\*,23\*\*,24\*], and 3) VLPs comprised of virion components identified in the parasitoids *Venturia canescens* and *Fopius arisanus* that also evolved from nudiviruses [3\*\*,4\*\*]. BVs, IVs and VLPs are much more complex than other known EVEs because many genes from their virus ancestors have been retained by wasps and still function to produce virions or virion components. We thus suggest domesticated endogenous virus (DEV) more accurately describes these entities than the term EVE.

### **Nudiviruses and DEVs in parasitoids exhibit a continuum of interacting genes**

The circular, double-stranded DNA genomes (80-230 kb) of viruses in the *Nudiviridae* and its sister family, the *Baculoviridae*, are comprised of many intronless genes (Table 1). Viruses in both families specifically infect arthropods, are usually pathogenic, and replicate in the nuclei of host cells, which release mature virions comprised of nucleocapsids surrounded by a single envelope (Figure 2) [25]. Baculoviruses and nudiviruses also share a partially overlapping set of core genes with functions in replicating the viral genome and producing virions (Table 1) [26]. These include a DNA polymerase that studies of baculoviruses indicate is primarily responsible for replicating the viral genome. It also includes four genes that produce a DNA dependent RNA polymerase that through promoter recognition specifically transcribes the structural genes that produce the nucleocapsid and envelope proteins that form virions [27] (Table 1).

## Bracoviruses

BVs assemble in the nuclei of ovary calyx cells that lyse to release mature virions into the lumen of the lateral oviducts. Transcriptome data from ovaries together with the proteomic analysis of virions from two braconids, *Cotesia congregata* and *Chelonus inanitus*, provided the first evidence that proteins in BV virions are homologs of nudivirus genes, which strongly suggested BVs evolved from a nudivirus ancestor [21<sup>\*\*</sup>,28] (Figure 1). Comparing BVs to nudiviruses indicates they also share several morphological features (Figure 2). Phylogenetic studies support that the nudivirus ancestor of BVs integrated into the genome of a braconid ~100 million years ago (MYA) [29<sup>\*</sup>,30], which was followed by speciation events that have resulted in an estimated ~50,000 BV-producing wasp species existing today [31]. Collectively, these BV-producing wasps form a monophyletic assemblage called the Microgastroid complex that is divided into six subfamilies [32] (Sharanowski *et al.*, this issue).

Insights into BV genome evolution and function primarily derive from studies of two species, *Microplitis demolitor* and *C. congregata*, in the subfamily Microgastrinae [21<sup>\*\*</sup>,22<sup>\*\*</sup>,33<sup>\*\*</sup>,34,35<sup>\*</sup>,36<sup>\*\*</sup>,37<sup>\*</sup>]. Unlike nudiviruses, genome sequencing of initially *M. demolitor* and then *C. congregata* showed that BV genome components are globally dispersed in the genomes of wasps [33<sup>\*\*</sup>,34<sup>\*</sup>,36<sup>\*\*</sup>]. However, these components function as two units, which underlies how virions have been repurposed to package only virulence genes. The first of these units consists of nudivirus genes that despite being dispersed are coordinately expressed and interact to produce virions [22<sup>\*\*</sup>,36<sup>\*\*</sup>] (Table 1). Included in this gene set are the four nudivirus core genes that produce the RNA polymerase, which are expressed early in calyx cells. This is followed by the expression of several other nudivirus genes that encode nucleocapsid and envelope proteins which the nudivirus RNA polymerase transcribes [22<sup>\*\*</sup>,36<sup>\*\*</sup>,38] (Table 1). The fact that almost none of these genes has gained introns after 100 million years suggests strong evolutionary constraints continue to operate in regulating transcription of these viral genes probably in virogenic stroma where virion assembly occurs [39] (Figure 2). Almost half of these genes reside in a very similar ~100 kb domain in the genomes of *M. demolitor* and

*C. congregata*, which may be a remnant of the nudivirus that integrated into the wasp ancestor, but the remainder are widely and disparately dispersed [33<sup>\*\*</sup>,36<sup>\*\*</sup>]. Notably absent, however, is a nudivirus DNA polymerase that as a core gene would have been present in the ancestor of BVs [33<sup>\*\*</sup>,36<sup>\*\*</sup>]. Loss of this gene early in evolution has been suggested as potentially critical in preventing the ancestor from being able to replicate its genome and cause disease in wasps [4<sup>\*\*</sup>]. The second functional unit is a collection of DNA domains containing virulence genes called proviral segments. Most proviral segments are tandemly arrayed in macroloci with orthology relationships suggesting organization was already established before *M. demolitor* and *C. congregata* diverged 53 MYA [36<sup>\*\*</sup>]. Proviral segments are strongly amplified and then processed into circular, double stranded DNAs that range from ~6->40 kb that are individually packaged into virions but have aggregate sizes of >200 kb [40,41<sup>\*</sup>,42<sup>\*</sup>]. Amplification of proviral segments requires a wasp DNA polymerase that remains unidentified [33<sup>\*\*</sup>,36<sup>\*\*</sup>,43,44]. However, recombinases and conserved sequences flanking proviral segments were both likely inherited from the nudivirus ancestor, and are known to also be required for processing functions that produce the circular, dsDNAs that are packaged into nucleocapsids [22<sup>\*\*</sup>,40,41<sup>\*\*</sup>]. Some of the virulence genes on proviral segments originate from wasps [37<sup>\*</sup>,45,46] while others derive from transposable elements [36<sup>\*\*</sup>,37<sup>\*</sup>,47,48] or potentially the nudivirus ancestor but have diverged to the point that their phylogenetic relationship is no longer detectable. However, most BV virulence genes are of unknown origin [37<sup>\*</sup>]. Females inject eggs and BVs into hosts, which is followed by the former hatching and the latter rapidly infecting different cell types followed by expression of virulence genes. Functional studies further implicate several BV virulence genes in altering host immune defenses and growth in ways that enable wasp larvae to develop [49-51].

## **Ichnoviruses**

Several thousand species in two subfamilies of the Ichneumonidae (Campopleginae, Banchinae) produce IVs in calyx cells that morphologically are very similar to one another but are distinct from BVs [23<sup>\*\*</sup>,52]. Insights into IV evolution and function derive from largely similar



approaches to those used to characterize BVs. Studies that initially focused on the campoplegine *Hyposoter didymator* identified several genes whose products are detected in IV virions, which suggests these genes encode virion structural proteins [23\*\*]. None of these genes share significant homology with genes in any known virus but they reside in domains of the wasp genome named IV structural protein encoding regions (IVSPERs), which exhibit features that strongly suggest a viral origin [23\*\*,53\*\*]. Whole genome sequencing of *H. didymator* and another campoplegine, *Campoletis sonorensis*, indicate that gene content and order in IVSPERs are conserved while proviral segments containing virulence genes exhibit considerable dispersion and are more variable [54\*\*]. Wasps in the subfamily Banchinae are less studied, but genes in the IVSPERs of campoplegine wasps have been identified in the banchine *Glypta fumiferanae* that also exhibit similar order [24\*]. However, phylogenetic data indicate that the Campopleginae and Banchinae are separated by several other subfamilies of ichneumonids that do not produce IVs. These findings thus overall favor the interpretation that IVs arose twice from the same or a closely related virus that is either extinct or undiscovered [19\*,32].

### **Virus-like particles**

Endogenized nudiviruses in *V. canescens* and *F. arisanus* have resulted in a different strategy of parasitoid wasp manipulation of host physiology: producing VLPs that contain proteins but no nucleic acid [3\*\*,19\*]. Compared to BVs, the nudiviruses associated with *V. canescens* and *F. arisanus* have also been more recently acquired and thus provide additional insights into nudiviruses domestication. Sequencing of *F. arisanus* shows that the nudivirus genes which produce the RNA polymerase and several structural genes that encode envelope and nucleocapsid components are expressed in calyx cells while other nudivirus core genes are pseudogenized or absent, suggesting decay (Table 1). In turn, the VLPs produced in calyx cells consist of a partially formed nucleocapsid that is surrounded by an envelope [4\*\*]. Results from *V. canescens* likewise indicate the four nudivirus genes that encode the RNA polymerase subunits and several nudivirus structural genes that encode predicted envelope components

are expressed [3\*\*]. Among the latter are several '*per os* infectivity factor (*pif*)' genes that are also encoded by nudiviruses and baculoviruses. Experimental studies in baculoviruses further indicate several *pif* gene products interact and have functions in infecting midgut cells [55]. In contrast, that nudivirus genes encoding nucleocapsid components are pseudogenized (Table 1) [43] is consistent with formation of VLPs comprised of an envelope that surrounds at least three proteins of wasp origin [3\*\*]. Assays further indicate the VLPs produced by *V. canescens* bind to the surface of wasp eggs while the proteins inside envelopes have functions in protecting eggs from the host's immune system [56]. The nudivirus genes in the *V. canescens* and *F. arisanus* genomes are much less dispersed than in the wasps that produce BVs; overall suggesting the early stages of domestication after integration of a complete nudivirus genome include pseudogenization and decay of genes that are not selected for, while genes that remain functional progressively disperse via chromosomal rearrangements. Like BV-producing wasps, *F. arisanus* and *V. canescens* also both lack a nudivirus DNA polymerase suggesting loss of this gene is an important early step in nudivirus domestication while retention of the genes that produce the RNA polymerase is key to producing virions or VLPs that have been repurposed for functions in parasitizing hosts.

### **The viral nature of BVs, IVs and VLPs**

Altogether, nudiviruses have commonly given rise to EVEs that in most insects show no evidence of being functional. In contrast, nudiviruses coopted by a number of parasitoids continue to produce virions (BVs), VLPs comprised of a nucleocapsid and viral envelope or VLPs that consist of a viral envelope alone. IVs derived from an unknown virus further exhibit many functional features of BVs. Coming to a consensus about naming BVs, IVs and VLPs is a challenge. At present, the International Committee on Taxonomy of Viruses (ICTV) defines a virus *sensu stricto* as a type of mobile genetic element that encodes at least one protein that is a major component of the virion encasing the nucleic acid of the respective mobile genetic element (<https://talk.ictvonline.org/information/w/ictv-information/383/ictv-code>). In a broader definition provided in the same text by the ICTV, mobile genetic elements that are of

evolutionary descent from a virus can also be considered viruses. BVs and IVs could thus be formally considered viruses given the strong evidence supporting their evolution from viral ancestors. In contrast, we suggest VLP would be a more appropriate term for the particles produced by *V. canescens* and *F. arisanus* which also are of viral origin but are not mobile genetic elements. BVs and IVs are clearly no longer autonomous in an evolutionary sense given that there is no replicating viral genome that is distinct from the wasp genome. However, forgoing the name “virus” when referring to BVs or IVs detracts from the fact that both produce massive numbers of infectious virions in wasp calyx cells that package many genes through the coordinated interaction of many other genes that were inherited from endogenized virus ancestors. Most BV gene products also continue to interact as they do in baculoviruses and nudiviruses, which further suggests these entities are better described as DEVs than EVEs which are predominantly non-functional sequences of viral origin in the genomes of animals. DEV would be appropriate for VLPs as well if a slightly more flexible definition of viruses was accepted, given that BVs and VLPs both have clear phylogenetic links to nudiviruses (Figure 1). BVs and IVs have also been called “symbiotic viruses” by analogy to vertically transmitted bacterial symbionts that have given rise to organelles like mitochondria and chloroplasts [57]. While bacterial symbionts still contain remnants of their original genome, BV, IV, and VLP genome components are all integrated in the genomes of wasps. However, the comparison of DEVs to other symbionts of microbial origin does provide another perspective when considering what to call BVs and IVs.

Other challenges in the study of DEVs also exist. For example, while many nudivirus-derived EVEs have been identified in insects, to date all examples of endogenized nudiviruses that continue to produce virions are in parasitoids. Does this reflect more intensive study of parasitoids owing to the discovery of BVs and IVs in the 1970s, or does the life cycle of these insects, which obligately involves the development of offspring in another arthropod and horizontal transfer of many factors that promote offspring development from wasps to hosts, lend itself to coopting viruses for similar purposes? A second area of need is to deorphanize the

function of many BV and IV genes in order to understand how transcription is controlled to assure that virions are only produced in calyx cells. In summary, additional comparative data on BVs and IVs are likely to further enhance insights about genome organization and function. Other comparative studies will also likely identify further examples of endogenized virus domestication.

### **Conflict of interest statement**

Nothing declared.

### **Acknowledgements**

We would like to thank Matthieu Leobold for discussions on virogenic stroma and Ange Lorenzi for taking the electron microscopy image showing nudivirus virions. We also would like to thank the French National Research Agency ANR (ABC Papogen project ANR-12-ADAP-0001 and CoteBio ANR17-CE32-0015-02), US National Science Foundation (DEB-1916788), US Department of Agriculture National Institute of Food and Agriculture Hatch Program (GEO0772) and Pulliam Endowment for support.

### **References and recommended reading**

Papers of particular interest published within the period of review and related to the main topic, have been highlighted as:

• of special interest

•• of outstanding interest

1. Vinson SB, Scott JR: **Particles containing DNA associated with the oocyte of an insect parasitoid.** *J Inverteb. Pathol* 1975, **25**:375-378.
2. Stoltz DB, Krell PJ, Summers MD, Vinson SB: **Polydnviridae - A proposed family of insect viruses with segmented, double-stranded, circular DNA genomes.** *Intervirology* 1984, **21**:1-4.
3. Pichon A, Bézier A, Urbach S, Aury JM, Jouan V, Ravallec M, Guy J, Cousserans F, Thézé J, Gauthier J, et al.: **Recurrent DNA virus domestication leading to different parasite virulence strategies.** *Sci Adv* 2015, **1**:e1501150.

- Presents the first evidence that an endogenized nudivirus produces *Venturia canescens* VLPs, which were initially described in 1966.
4. Burke GR, Simmonds TJ, Sharanowski BJ, Geib SM: **Rapid Viral Symbiogenesis via Changes in Parasitoid Wasp Genome Architecture.** *Mol Biol Evol* 2018, **35**:2463-2474.
  - This paper presents results that an endogenized nudivirus produces *Fopius arisanus* VLPs, which structurally differ from the VLPs produced by *V. canescens* by consisting of both nucleocapsid and envelope proteins.
  5. De Paepe M, Tournier L, Moncaut E, Son O, Langella P, Petit MA: **Carriage of lambda Latent Virus Is Costly for Its Bacterial Host due to Frequent Reactivation in Monoxenic Mouse Intestine.** *PLoS Genet* 2016, **12**:e1005861.
  6. Eiden MV: **Endogenous retroviruses - Aiding and abetting genomic plasticity.** *Cell Mol Life Sci* 2008, **65**:3325-3328.
  7. Arbuckle JH, Medveczky MM, Luka J, Hadley SH, Luegmayer A, Ablashi D, Lund TC, Tolar J, De Meirleir K, Montoya JG, et al.: **The latent human herpesvirus-6A genome specifically integrates in telomeres of human chromosomes *in vivo* and *in vitro*.** *Proc Natl Acad Sci U S A* 2010, **107**:5563-5568.
  8. Katzourakis A, Gifford RJ: **Endogenous viral elements in animal genomes.** *PLoS Genet* 2010, **6**:e1001191.
  9. Lavalie C, Cornelis G, Dupressoir A, Esnault C, Heidmann O, Vernochet C, Heidmann T: **Paleovirology of 'syncytins', retroviral env genes exapted for a role in placentation.** *Philos Trans R Soc Lond B Biol Sci* 2013, **368**:20120507.
  10. Ashley J, Cordy B, Lucia D, Fradkin LG, Budnik V, Thomson T: **Retrovirus-like Gag protein Arc1 binds RNA and traffics across synaptic boutons.** *Cell* 2018, **172**:262-274 e211.
  11. Suzuki Y, Baidaliuk A, Miesen P, Frangeul L, Crist AB, Merklings SH, Fontaine A, Lequime S, Moltini-Conclois I, Blanc H, et al.: **Non-retroviral Endogenous Viral Element Limits Cognate Virus Replication in *Aedes aegypti* Ovaries.** *Curr Biol* 2020, **30**:3495-3506 e3496.

12. Lin CL, Lee JC, Chen SS, Wood HA, Li ML, Li CF, Chao YC: **Persistent Hz-1 virus infection in insect cells: evidence for insertion of viral DNA into host chromosomes and viral infection in a latent status.** *J Virol* 1999, **73**:128-139.
  13. Cheng RL, Xi Y, Lou YH, Wang Z, Xu JY, Xu HJ, Zhang CX: **Brown planthopper nudivirus DNA integrated in its host genome.** *J Virol* 2014, **88**:5310-5318.
  14. Zhang Y, Wang J, Han GZ: **Chalcid wasp paleoviruses bridge the evolutionary gap between bracoviruses and nudiviruses.** *Virology* 2020, **542**:34-39.
  15. Liu S, Coates BS, Bonning BC: **Endogenous viral elements integrated into the genome of the soybean aphid, *Aphis glycines*.** *Insect Biochem Mol Biol* 2020, **123**:103405.
  16. Cheng RL, Li XF, Zhang CX: **Nudivirus Remnants in the Genomes of Arthropods.** *Genome Biol Evol* 2020, **12**:578-588.
  17. Lepetit D, Gillet B, Hughes S, Kraaijeveld K, Varaldi J: **Genome Sequencing of the Behavior Manipulating Virus LbFV Reveals a Possible New Virus Family.** *Genome Biol Evol* 2016, **8**:3718-3739.
  18. Di Giovanni D, Lepetit D, Guinet B, Bennetot B, Boulesteix M, Coute Y, Bouchez O, Ravallec M, Varaldi J: **A Behavior-Manipulating Virus Relative as a Source of Adaptive Genes for *Drosophila* Parasitoids.** *Mol Biol Evol* 2020, **37**:2791-2807.
- Identification of an endogenous filamentous virus that provides genes potentially used in the production of *Leptopilina boulardi* VLPs described since 1990.
19. Burke GR, Hines HM, Sharanowski BJ: **The Presence of Ancient Core Genes Reveals Endogenization from Diverse Viral Ancestors in Parasitoid Wasps.** *Genome Biol Evol* 2021, **13**:evab105.
- This study presents comparative genomics data that better resolve the placement in the family Ichneumonidae of the parasitoid lineages that produce IVs.
20. Drezen JM, Gauthier J, Josse T, Bézier A, Herniou E, Huguet E: **Foreign DNA acquisition by invertebrate genomes.** *J Invertebr Pathol* 2017, **147**:157-168.

21. Bézier A, Annaheim M, Herbinière J, Wetterwald C, Gyapay G, Bernard-Samain S, Wincker P, Roditi I, Heller M, Belghazi M, et al.: **Polydnaviruses of braconid wasps derive from an ancestral nudivirus**. *Science* 2009, **323**:926-930.
- This study used a multiomics approach that provided the first evidence that BVs evolved from an endogenized nudivirus.
22. Burke GR, Thomas SA, Eum JH, Strand MR: **Mutualistic polydnaviruses share essential replication gene functions with pathogenic ancestors**. *PLoS Pathog* 2013, **9**:e1003348.
- Homology to baculovirus core genes provided inferential evidence on the possible function of some BV genes but prior studies provided no experimental data. This paper in contrast presents the first experimental results on the function of several nudivirus genes in BV formation.
23. Volkoff AN, Jouan V, Urbach S, Samain S, Bergoin M, Wincker P, Demetere E, Cousserans F, Provost B, Coulibaly F, et al.: **Analysis of virion structural components reveals vestiges of the ancestral ichnovirus genome**. *PLoS Pathog* 2010, **6**:e1000923.
- Genomic data generated and analyzed in this study discovered that several gene products present in the campoplegine wasp *Hyposoter didymator* reside in IVSPERs. This result enabled comparative studies showing that gene products that produce IVs in other species also reside in IVSPERs.
24. Béliveau C, Cohen A, Stewart D, Periquet G, Djoumad A, Kuhn L, Stoltz D, Boyle B, Volkoff AN, Herniou EA, et al.: **Genomic and proteomic analyses indicate that banchine and campoplegine polydnaviruses have similar, if not identical, viral ancestors**. *J Virol* 2015, **89**:8909-8921.
- Identification of viral genes involved in IV production in a wasp species in the subfamily Banchinae.
25. Harrison RL, Herniou EA, Bézier A, Jehle JA, Burand JP, Theilmann DA, Krell PJ, van Oers MM, Nakai M, Ictv Report C: **ICTV Virus Taxonomy Profile: Nudiviridae**. *J Gen Virol* 2020, **101**:3-4.

26. Bézier A, Thézé J, Gavory F, Gaillard J, Poulain J, Drezen JM, Herniou EA: **The genome of the nucleopolyhedrosis-causing virus from *Tipula oleracea* sheds new light on the Nudiviridae family.** *J Virol* 2015, **89**:3008-3025.
  27. Guarino LA, Xu B, Jin J, Dong W: **A virus-encoded RNA polymerase purified from baculovirus-infected cells.** *J Virol* 1998, **72**:7985-7991.
  28. Wetterwald C, Roth T, Kaeslin M, Annaheim M, Wespi G, Heller M, Maser P, Roditi I, Pfister-Wilhelm R, Bézier A, et al.: **Identification of bracovirus particle proteins and analysis of their transcript levels at the stage of virion formation.** *J Gen Virol* 2010, **91**:2610-2619.
  29. Murphy N, Banks JC, Whitfield JB, Austin AD: **Phylogeny of the parasitic microgastroid subfamilies (Hymenoptera: Braconidae) based on sequence data from seven genes, with an improved time estimate of the origin of the lineage.** *Mol Phylogenet Evol* 2008, **47**:378-395.
- Phylogenetic analysis dating the origin of the monophyletic group of wasps associated with bracoviruses.
30. Thézé J, Bézier A, Periquet G, Drezen JM, Herniou EA: **Paleozoic origin of insect large dsDNA viruses.** *Proc Natl Acad Sci U S A* 2011, **108**:15931-15935.
  31. Rodriguez JJ, Fernandez-Triana JL, Smith MA, Janzen DH, Hallwachs W, Erwin TL, Whitfield JB: **Extrapolations from field studies and known faunas converge on dramatically increased estimates of global microgastrine parasitoid wasp species richness (Hymenoptera: Braconidae).** *Insect Conservation and Diversity* 2013, **6**:530-536.
  32. Sharanowski BJ, Ridenbaugh RD, Piekarski PK, Broad GR, Burke GR, Deans AR, Lemmon AR, Moriarty Lemmon EC, Diehl GJ, Whitfield JB, et al.: **Phylogenomics of Ichneumonoidea (Hymenoptera) and implications for evolution of mode of parasitism and viral endogenization.** *Mol Phylogenet Evol* 2021, **156**:107023.
  33. Burke GR, Walden KK, Whitfield JB, Robertson HM, Strand MR: **Widespread genome reorganization of an obligate virus mutualist.** *PLoS Genet* 2014, **10**:e1004660.



\*\* First genome of a BV-producing wasp. Results further showed that the ancestral nudvirus genome is highly dispersed in the genome of the wasp.

34. Burke GR, Walden KKO, Whitfield JB, Robertson HM, Strand MR: **Whole Genome Sequence of the Parasitoid Wasp *Microplitis demolitor* That Harbors an Endogenous Virus Mutualist.** *G3 (Bethesda)* 2018, **8**:2875-2880.

35. Bézier A, Louis F, Jancek S, Periquet G, Thézé J, Gyapay G, Musset K, Lesobre J, Lenoble P, Dupuy C, et al.: **Functional endogenous viral elements in the genome of the parasitoid wasp *Cotesia congregata*: insights into the evolutionary dynamics of bracoviruses.** *Philos Trans R Soc Lond B Biol Sci* 2013, **368**:20130047.

\* Comparative genomics of BV proviral segments in four wasp species in the subfamily Microgastrinae that diverged 20 MYA millions years ago.

36. Gauthier J, Boulain H, van Vugt J, Baudry L, Persyn E, Aury JM, Noel B, Bretaudeau A, Legeai F, Warris S, et al.: **Chromosomal scale assembly of parasitic wasp genome reveals symbiotic virus colonization.** *Commun Biol* 2021, **4**:104.

\*\* First genome from a BV-producing wasp that shows how viral genome components are organized on a chromosomal scale.

37. Desjardins CA, Gundersen-Rindal DE, Hostetler JB, Tallon LJ, Fadrosch DW, Fuester RW, Pedroni MJ, Haas BJ, Schatz MC, Jones KM, et al.: **Comparative genomics of mutualistic viruses of *Glyptapanteles* parasitic wasps.** *Genome Biol* 2008, **9**:R183.

\* First study to sequence and characterize BV proviral segments encoding virulence genes. Results showed that most proviral segments are tandemly arrayed in loci, which comparative studies later showed to also be the case in other BV-producing wasps.

38. Arvin MJ, Lorenzi A, Burke GR, Strand MR: **MdBVe46 is an envelope protein that is required for virion formation by *Microplitis demolitor* bracovirus.** *J Gen Virol* 2021, **102**:001565.

39. Vanarsdall AL, Mikhailov VS, Rohrmann GF: **Characterization of a baculovirus lacking the DBP (DNA-binding protein) gene.** *Virology* 2007, **364**:475-485.

40. Desjardins CA, Gundersen-Rindal DE, Hostetler JB, Tallon LJ, Fuester RW, Schatz MC, Pedroni MJ, Fadrosch DW, Haas BJ, Toms BS, et al.: **Structure and evolution of a proviral locus of *Glyptapanteles indiensis* bracovirus**. *BMC Microbiol* 2007, **7**:61.
41. Louis F, Bézier A, Periquet G, Ferras C, Drezen JM, Dupuy C: **The bracovirus genome of the parasitoid wasp *Cotesia congregata* is amplified within 13 replication units, including sequences not packaged in the particles**. *J Virol* 2013, **87**:9649-9660.
- Mapping of bracovirus replication units and identification of an unexpected patterns of amplification resembling that reported for poxviruses.
42. Burke GR, Simmonds TJ, Thomas SA, Strand MR: ***Microplitis demolitor* bracovirus proviral loci and clustered replication genes exhibit distinct DNA amplification patterns during replication**. *J Virol* 2015, **89**:9511-9523.
- Long read sequencing and analysis of BV replication intermediates that revealed further complexity in how BV proviral segments are amplified during replication.
43. Leobold M, Bézier A, Pichon A, Herniou EA, Volkoff AN, Drezen JM: **The Domestication of a Large DNA Virus by the Wasp *Venturia canescens* Involves Targeted Genome Reduction through Pseudogenization**. *Genome Biol Evol* 2018, **10**:1745-1764.
- Identifies nudivirus core genes that are present in the genome of *V. canescens* but are no longer involved in VLP production.
44. Burke GR, Strand MR: **Deep sequencing identifies viral and wasp genes with potential roles in replication of *Microplitis demolitor* Bracovirus**. *J Virol* 2012, **86**:3293-3306.
45. Burke GR, Strand MR: **Systematic analysis of a wasp parasitism arsenal**. *Mol Ecol* 2014, **23**:890-901.
46. Gasmi L, Boulain H, Gauthier J, Hua-Van A, Musset K, Jakubowska AK, Aury JM, Volkoff AN, Huguet E, Herrero S, et al.: **Recurrent Domestication by Lepidoptera of Genes from Their Parasites Mediated by Bracoviruses**. *PLoS Genet* 2015, **11**:e1005470.
47. Drezen JM, Bézier A, Lesobre J, Huguet E, Cattolico L, Periquet G, Dupuy C: **The few virus-like genes of *Cotesia congregata* bracovirus**. *Arch Insect Biochem Physiol* 2006, **61**:110-122.

48. Zhang HH, Zhou QZ, Wang PL, Xiong XM, Luchetti A, Raoult D, Levasseur A, Santini S, Abergel C, Legendre M, et al.: **Unexpected invasion of miniature inverted-repeat transposable elements in viral genomes.** *Mob DNA* 2018, **9**:19.
49. Thoetkiattikul H, Beck MH, Strand MR: **Inhibitor kappaB-like proteins from a polydnavirus inhibit NF-kappaB activation and suppress the insect immune response.** *Proc Natl Acad Sci U S A* 2005, **102**:11426-11431.
50. Beck MH, Strand MR: **A novel polydnavirus protein inhibits the insect prophenoloxidase activation pathway.** *Proc Natl Acad Sci U S A* 2007, **104**:19267-19272.
51. Pruijssers AJ, Strand MR: **PTP-H2 and PTP-H3 from Microplitis demolitor Bracovirus localize to focal adhesions and are antiphagocytic in insect immune cells.** *J Virol* 2007, **81**:1209-1219.
52. Drezen JM, Leobold M, Bézier A, Huguet E, Volkoff AN, Herniou EA: **Endogenous viruses of parasitic wasps: variations on a common theme.** *Curr Opin Virol* 2017, **25**:41-48.
53. Lorenzi A, Ravallec M, Eychenne M, Jouan V, Robin S, Darboux I, Legeai F, Gosselin-Grenet AS, Sicard M, Stoltz D, et al.: **RNA interference identifies domesticated viral genes involved in assembly and trafficking of virus-derived particles in ichneumonid wasps.** *PLoS Pathog* 2019, **15**:e1008210.

•• Presents the first experimental results on the function of genes in IVSPERs that are required for IV formation.

54. Legeai F, Santos BF, Robin S, Bretaudeau A, Dikow RB, Lemaitre C, Jouan V, Ravallec M, Drezen JM, Tagu D, et al.: **Genomic architecture of endogenous ichnoviruses reveals distinct evolutionary pathways leading to virus domestication in parasitic wasps.** *BMC Biol* 2020, **18**:89.

•• Results show that IVSPERs encoding structural genes are more stable in the wasp genome than proviral segments.

55. Peng K, van Oers MM, Hu Z, van Lent JW, Vlak JM: **Baculovirus *per os* infectivity factors form a complex on the surface of occlusion-derived virus.** *J Virol* 2010, **84**:9497-9504.
56. Reineke A, Asgari S, Schmidt O: **Evolutionary origin of *Venturia canescens* virus-like particles.** *Arch Insect Biochem Physiol* 2006, **61**:123-133.
57. Federici BA, Bigot Y: **Origin and evolution of polydnaviruses by symbiogenesis of insect DNA viruses in endoparasitic wasps.** *J Insect Physiol* 2003, **49**:419-432.

## Figure legends

### Figure 1

Phylogenetic tree for baculoviruses, nudiviruses and endogenous nudiviruses. The colors and abbreviations are those defined in Table 1. The tree was constructed using previously described sequences [3], which include 17 nudivirus core genes that are indicated in bold in Table 1. Amino acid alignments were individually performed with default parameters using the MAFFT alignment plugin v7.0450 from Geneious Prime 2019.2.3 software. A consensus tree was then constructed using Geneious Tree builder on concatenated and refined multiple alignments with the following parameters: genetic distance model = Jukes-Cantor, tree build method = Neighbor- Joining, resampling method = bootstrap, number of replicates = 1000. A nimavirus (white spot syndrome virus (WSSV, AF369029)) was used as outgroup based on gene content. Numbers on tree nodes indicate bootstrap supports (1000 replicates).

### Figure 2

Electron micrographs showing that nudivirus and BV virions exhibit several similarities. (a) Morphogenesis of the betanucleivirus *Heliothis zea* nudivirus -1 (HzNV-1) in the nucleus of an infected host cell. Spheroidal envelopes (E) *de novo* assemble in proximity to virogenic stroma (VS), which is followed by the *de novo* formation of a single electron dense, cylindrical nucleocapsid within an envelope to form a virion (V). Scale bar = 500  $\mu$ m. (b). Morphogenesis of *Microplitis demolitor* bracovirus (MdBV) in the nucleus of a calyx cell. Individual, crescent-shaped envelopes (E) and electron dense nucleocapsids (N) *de novo* assemble in proximity to virogenic stroma (VS). This is followed by envelopes individually surrounding cylindrical nucleocapsids (E+N) to produce virions (V) that form parallel arrays before calyx cell lysis releases virions into the lateral oviducts. HzNV-1 nucleocapsids are larger than MdBV nucleocapsids, which likely reflects that the HzNV-1 genome (228 kb) is much larger than any of the viral segments that are individually packaged into MdBV virions (6-40 kb). In contrast, the number of MdBV virions that form in the nuclei of calyx cells is much larger than the number of HzNV-1 virions that form in the nuclei of permissive host cells.

Table 1

# Retention of the 32 nudivirus core genes among baculoviruses, nudiviruses and endogenized nudiviruses from parasitoid wasps

GENE FUNCTION gene name	BACULOVIRUS				NUDIVIRUS					ENDOGENOUS NUDIVIRUS				
	α	β	δ	γ	α	β	δ	γ	ε*	Bracovirus			VLP	
	Ac MNPV	Cp GV	Nese NPV	Cuni NPV	Or NV	Hz NV-1	To NV	Pm NV	Dh NV	Cc BV	Md BV	Ci BV	Vc ENV	Fa ENV
<b>DNA replication/processing</b>														
1 DNA polymerase DNApol	●	●	●	●	●	●	●	●	●	○	○	/	⊗	/
2 helicase	●	●	●	●	●	●	●	●	●	●	●	/	●	●
3 helicase 2	○	●	○	○	●	●	●	●	●	○	○	/	○	○
4 integrase (int)	○	○	○	○	●	●	●	●	●	●	●	●	⊗	/
5 flap endonuclease fen-1	○	○	○	○	●	●	●	●	●	●	●	/	⊗	/
<b>Transcription/RNA polymerase</b>														
6 RNA polymerase p47	●	●	●	●	●	●	●	●	●	●	●	/	⊗	●
7 RNA polymerase lef-4 <sup>(1)</sup>	●	●	●	●	●	●	●	●	●	●	●	/	⊗	●
8 RNA polymerase lef-5	●	●	●	●	●	●	●	●	●	●	●	/	●	●
9 RNA polymerase lef-8	●	●	●	●	●	●	●	●	●	●	●	/	●	●
10 RNA polymerase lef-9	●	●	●	●	●	●	●	●	●	●	●	/	●	●
<b>Nucleotide metabolism</b>														
11 thymidine kinase tk1	○	○	○	○	●	●	●	●	●	/	/	/	/	/
12 thymidine kinase tk2	○	○	○	○	●	●	●	●	●	/	/	/	/	/
13 thymidine kinase tk3	○	○	○	○	●	●	●	●	●	/	/	/	/	/
<b>Structure/Capsid components</b>														
14 very late factor vlf-1 <sup>(2)</sup>	●	●	●	●	●	●	●	●	●	●	●	●	⊗	○
15 capsid protein vp39	●	●	●	●	●	●	●	●	●	●	●	●	⊗	●
16 p6.9 <sup>(3)</sup>	●	●	●	●	●	●	●	●	●	●	●	/	⊗	/
17 phosphatase 38K	●	●	●	●	●	●	●	●	●	●	●	●	⊗	●
<b>Per os infectivity factors &amp; envelope components</b>														
18 pif-0 (p74)	●	●	●	●	●	●	●	●	●	●	●	●	⊗	●
19 pif-1	●	●	●	●	●	●	●	●	●	●	●	●	⊗	●
20 pif-2	●	●	●	●	●	●	●	●	●	●	●	●	●	●
21 pif-3	●	●	●	●	●	●	●	●	●	●	●	/	●	●
22 pif-4 (19 kDa)	●	●	●	●	●	●	●	●	●	●	●	●	●	●
23 pif-5 (odv-e56)	●	●	●	●	●	●	●	●	●	●	●	●	●	○
24 pif-6 (ac68)	●	●	●	●	●	●	●	●	●	●	●	/	●	●
25 pif-8 (vp91)	●	●	●	●	●	●	●	●	●	●	●	●	●	●
26 sulphydryloxidase p33 (Ac92) <sup>(4)</sup>	●	●	●	●	●	●	●	●	●	●	●	/	●	●
27 ac81	●	●	●	●	●	●	●	●	●	○	○	/	●	●
<b>Other particle components</b>														
28 11K	○	○	○	○	●	●	●	●	●	●	●	●	●	○
29 alpha/beta hydrolase GbNVorf19-like <sup>(5)</sup>	○	○	○	○	●	●	●	●	●	●	●	/	●	●
30 GbNVorf51-like	○	○	○	○	●	●	●	●	●	○	○	/	●	●
31 GbNVorf58-like <sup>(6)</sup>	○	○	○	○	●	●	●	●	●	○	○	/	●	●
32 GbNVorf67-like <sup>(7)</sup>	○	○	○	○	●	●	●	●	●	●	●	/	●	●

Gene functions were assigned on the basis of homology with baculovirus genes and/or proteomic analysis of virions. Four baculoviruses in green (Autographa californica MNPV, Cydia pomonella GV, Neodiprion sertifer NPV and Culex nigripalpus NPV) and five nudiviruses in dark blue have been used as representatives of the genera currently recognized by ICTV (Oryctes rhinoceros NV, Heliothis zea NV-1, Tipula oleacera NV and Penaeus monodon NV) or that have been proposed as new genera (\*) in recently published studies (Dikerogammarus haemobaphes NV). Two complete (Cotesia congregata BV and Microplitis demolitor BV) bracoviruses and one partial (Chelonius inanitus BV) bracovirus are shown in red. Two other endogenized nudiviruses that produce virus-like particles (Venturia canescens ENV and Fopius arisanus ENV) are shown in purple and light blue, respectively. Filled circles (●) indicate presence of the corresponding gene whereas expanded filled circles (●) indicate the gene has expanded into a multimer family. Open circles (○) indicate the absence of the gene and cross filled circles (⊗) indicate detection of a remnant of the ancestral integrated gene. Finally, slashes (/) indicate either that no information is available from the wasp genome (*Chelonius inanitus*), the gene remnant was not deeply searched for (*Fopius arisanus*), or genes are too closely related to wasp genes to unambiguously determine their origin (e.g. *thymidine kinases*). <sup>(1)</sup> Newly identified *lef-4* pseudogene within VcENV cluster 6 (accession number KP972600), from position 3,891 to 5,798. <sup>(2)</sup> For the *vlf-1* family, *vlf-1* and *HzNV-1 orf140-like* were considered to belong to the same recombinase gene family. <sup>(3)</sup> A thorough search for the *p6.9* core gene revealed a putative candidate from position 61,992 to 62,192 in the DhNV genome and two putative candidates from positions 1,679,579 to 1,679,358 and 1,735,989 to 1,736,246 within *M. demolitor* scaffold\_0102 (accession number [NW\\_014463878.1](#)). <sup>(4)</sup> The presence of two copies of *p33/ac92* within the DhNV genome was not confirmed. # P33/Ac92 is detected in both baculovirus envelope and nucleocapsid fractions but is present in higher quantities within the former. <sup>(5)</sup> A *GbNVorf19-like* homolog was identified from position 1,730,226 to 1,731,254 within the same *M. demolitor* scaffold\_0102. <sup>(6)</sup> A *GbNVorf58-like* homolog was identified from position 62,198 to 62,365 within the DhNV genome. <sup>(7)</sup> A *GbNVorf67-like* homolog was identified from position 1,449 to 2,756 within *M. demolitor* scaffold\_0267\_contig3 (accession number AZMT02018993.1). The core genes conserved for the virus phylogeny are indicated in bold (see Figure 1).

Figure 1




Figure 2

