

Protein profiling by MS shows evidence of royal protein pheromones in the cuticle extracts of the termite Reticulitermes flavipes

Guillaume Gabant, Matej Nemcic, Fanny Ruhland, Timothée Toussaint, Christophe Lucas, Martine Cadene

▶ To cite this version:

Guillaume Gabant, Matej Nemcic, Fanny Ruhland, Timothée Toussaint, Christophe Lucas, et al.. Protein profiling by MS shows evidence of royal protein pheromones in the cuticle extracts of the termite Reticulitermes flavipes. Analytics 2022, Sep 2022, Nantes, France. 2022. hal-03845797

HAL Id: hal-03845797 https://hal.science/hal-03845797

Submitted on 20 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Protein profiling by MS shows evidence of protein pheromones in the cuticle extracts of the termite **Reticulitermes flavipes**



Viridiplantae / Mucuna pruni Fungi / Aspergillus nidulans Fungi / Aspergillus nidulans Fungi / Peniophora Metazoa / Insecta / Termitoidae / Coptotermes

Guillaume Gabant, Matej Nemcic, Fanny Ruhland, Timothée Toussaint, Christophe Lucas, Martine Cadene Centre de Biophysique Moléculaire, CNRS UPR4301, 3 Av. de la Recherche Scientifique, CS80054, 45071 Orléans cedex 2

Abstract

Introduction

Proteins are known to work as social interaction signals in many species in the animal kingdom [1]. While they are common mediators of behavior in mammals and aquatic species, proteins have seldom been identified as such in insects. In social insects in particular, individuals have numerous physical contacts, providing many opportunities for the exchange of contact molecules as interaction mediators. However, their role in social interactions is largely unexplored: are they rare or simply underestimated? We set out to find out using molecular profiling of cuticular extracts from different castes and sexes of the termite Reticulitermes flavipes, which can be correlated with behavioral effects in these termites.

Methods

Cuticle extracts of workers and reproductive individuals were obtained by washing live termites in water. Influence of cuticular extracts on workers behavior was assessed by video recordings. Molecular profiling of cuticular extracts was performed by MALDI-TOF MS using the ultra-thin layer spotting method [2]. Identification of proteins was carried out by top-down MS sequencing using MALDI-TOF/TOF LIFT and ESI-QTOF CID fragmentation.

Results

Using MS/MS methods, we show that aqueous cuticle extracts from both reproductive and worker contain proteins and/or peptides. Molecular profiling of protein from cuticular extracts using MALDI-TOF mass spectrometry reveals higher complexity in reproductives than in workers and a sex-specific composition exclusive to reproductives. Reproductives extracts impact workers body-shaking, a vibratory behavior involved in reproductives recognition, while extracts from workers do not. Thus, the difference in protein composition of cuticle extracts may underlie the specific effect of reproductives' extracts on workers behavior. Further MS characterization is necessary to identify the specific protein(s) responsible for this effect.

Conclusions

These results indicate that social signaling may not be limited to cuticular hydrocarbons or other volatile chemicals and that proteins could play an important role as pheromones in insect societies. It also opens new perspectives to understand the role of proteins in reproductives' replacement, fertility signaling, or interspecific insect communication.

MALDI-TOF MS profiling

Reproductives extracts are more complex than workers extracts

PLS-DA analyses based on the relative areas of CPCs discriminate extracts according to their caste and sex



ranging from *m/z* 1 000 to 25 000

Fig. 1: Averaged mass spectra of cuticular aqueous extracts 675 from MALD-TOF analyses. Spectra are shown in positive/negative display for reproductives versus workers for both (A) females and (B) males. The spectra are based on a sum up of 15 spectra for each caste and sex. Annotated peaks are the first-quartile of the 87 major CPCs identified as discriminant by the PLS-DA analyses with VIP values greater than 1. Note that the 3 major CPCs 4/26, 10056 and 10196 are also represented in the figure to support the discussion even if they are not in the first-nuaritie.



Fig. 3: Relative areas of the first-quartile of the 87 major CPCs identifies as discriminant by PLS-DA analyses for reproductives (A) and workers (B) of both seves: finalle reproductive (RF), male reproductives (RM), female workers (WF) and male workers (MW). Relative areas are represented as man (1 SEM) and are calculated on all the spectra - indicates significant differences (Wilcoxon signed-rank test, p=0.05; Multiple compensions were corrected with FDR as the p-value adjustement method).



Fig. 2: PLS-DA plot based on the means of the areas of the 197 CPCs identified from culcular aqueous extracts (n=15 per group; peaks were selected using the sum algorithm with s N > 4 and a peak width of D_{DA} (A) Score plot of female reproductives (FR light turquoise), male reproductives (FR, dark green), and mel workers (WM, dark orange). (B) Score plot of reproductives (dark green) and workers (light orange). Score plot display centroids and dispersion ellipses (cellipser 1-5).



Fig. 4: Compounds repartition across castes and sexes of the first-quartile of the 87 major CPCs. CPCs were attributed to a group if more than one individual present the compound

Aqueous cuticle extracts contain proteins

Top-down MS sequencing CPC-10056 Fig. 5: Fragmentation mass spectrum of the CPC-10056 using UHR-ESI-QTOF-MS/MS CID. The precursor ion *m*/z 914.4 (11+) is nted with a blue lozenge



A0A6L2PEY0 Uncharacterized protein G12

Entry Protein Name A0A371IC31 Protein GIGANTEA A0A1U80M93 ADSL_C domain-containing pro C658619 ADSL_C domain-containing pro A0A553WGF4 RNA-directed RNA polymerase

A0A6L2P7Q1

Uncharacterized protein

Database search

Table 1: Eukaryota hits identified with the sequence tag I/L/LDDV/I/AVI/L. The Peptide Search tool from Uniprot was used to interrogate the SwissProt and TrEMBL databases. Isoleucine and Leucine are treated as equivalent by Peptide Search.



Compoundsy Average New Tool Procursor MSMS Protein database Nucleotide database hit CPC-4428 4427.7 4425.1 LIFT match No significant No significant CPC-10056 10055.8 1171 CID match MSSignificant MSSignificant							
CPC-4428 4427.7 263 1 LIFT No significant match No significant No significant CPC-10056 10055.8 171 CID No significant No significant No significant	Compounds	Average m/z [M+H]+	Precu m/z	rsor z	MS/MS method	Protein database hit	Nucleotide database hit
CPC-10056 10055.8 914.4 11 CID No significant match match JK366975.1 from Coptotermess formosanus metatranscriptomic cDNA library, score = 81	CPC-4428	4427.7	4425. 263	1	LIFT	No significant match	No significant match
	CPC-10056	10055.8	914.4 171	11	CID	No significant match	JK369675.1 from Coptotermes formosanus metatranscriptomic cDNA library, score = 81

Body-shaking vary over time

Reproductives extracts impact workers behavior

Reproductives extracts impact workers body shaking, a vibratory behavior involved in reproductives recognition, while extracts from . workers do not



Fig. 7: Number of body-shaking of workers after 1h (A) and 24h (B) for the different treatments (n=10). In absence of reproductives (R-), in the presence of reproductives (R-), with extracts of workers (We) or reproductives (R-). Box plots show the median, upper and lower quartiles, and range; dots, outliers; crosses, means; different letters indicated significant differences (LMAs,

=0.05)



Fig. 8: Number of body-shaking of workers after 1 h and 24h for the diffe treatments (n=10). In absence of reproductives (R+), in the presence of reproductives (R+), with extracts of workers (Wo) or reproductives (R+) Box plots show the median, upper and lower quartiles, and range; dots, outliers; crosses, means; * indicates significant differences (Wilcoxon signed-rank test, p<0.05).

Discussion - Conclusion

Analyses of termite aqueous extracts of termite revealed a complex mixture of CPCs, which was caste and sex dependent Analyses of termite aqueous extracts of termite revealed a complex mixture of CPCs, which was caste and sex dependent. Mass spectrometry fragmentation confirmed that the profiled compounds are indeed proteins, with at least one example of an ortholog of G12, a digestive protein in termites. CPCs composition was more complex in reproductives than workers, with sex-specific CPCs for both male and female reproductives. This diversity of CPCs, which sex and caste specific compositions, could encode complex communication signals like the cuticular hydrocarbons (CHCs) which are well known to be involved in several social interactions.

Cuticular aqueous extracts from reproductives were enough to trigger body-shaking, while extracts from workers did not. Culcular adjuedus extracts inom reproductives were enough to ingger body-shaking winte extracts inom workers did not. Time dependent comparisons showed an increase of body-shaking events at 24 ho compared to 1h in all treatments, except for reproductives extracts where it was only marginally significant. This general increase, for both the mean values and the data variances, may reflect a behavioral instability due to social isolation. Indeed, tested micro-nests are fragments of larger mother colonies displaying rich interactions. Therefore, workers might be more behaviorally disturbed after 24h of isolation than after 1h. At 24h, the presence of reproductives still induce body-shaking but the reproductives extracts do not anymore. This short effect of the protein extracts might be due to a lack of stability overtime.

The combination of protein profiling with behavioral studies represents a powerful tool to explore unknown aspects of insect communications, which could help to answer unresolved questions about social communication

Future studies will be needed to sequence and identify reproductive-specific CPCs and to test their behavioral functions, either individually or in synergy, to determine their role in the regulation of social interactions in insects, as has been already observed in vertebrates or aquatic organisms,

Acknowledgments





References

[1] Wyatt TD. (2014) Anim. Behav. 97, 273 [2] Cadene M. and Chait B.T. (2000) Anal nem. 72. 5655-5658

Douy-snaking vary over time Time dependent comparisons showed an increase of body-shaking events at 24h compared to 1h in all treatments, except for reproductives extracts where it was only marginally significant (z = 8.5, p = 0.059)