

$\label{eq:choronomy} \begin{array}{l} \text{Two Different Compounds in One Crystal:} \\ \text{di-}\mu\text{-chlorobis}[2-(5\text{-methyl-pyridin-}2\text{-yl-}\kappa\mathbf{N})\text{phenyl-}\\ \kappa\mathbf{C}(1)]\text{diplatinum(II) and} \\ \text{chloro}(\mathbf{N}, \mathbf{N}\text{-dimethylformamide-}\kappa\mathbf{O})[2-(5\text{-methyl-pyridin-}2\text{-yl-}\kappa\mathbf{N})\text{phenyl-}\kappa\mathbf{C}(1)]\text{platinum(II)} \end{array}$

Isabelle Sasaki, Sonia Mallet-Ladeira

▶ To cite this version:

Isabelle Sasaki, Sonia Mallet-Ladeira. Two Different Compounds in One Crystal: di- μ -chlorobis[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]diplatinum(II) and chloro(N,N-dimethylformamide- κ O)[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]platinum(II). Journal of Chemical Crystallography, 2023, 53, pp.293-298. 10.1007/s10870-022-00970-0. hal-03846487

HAL Id: hal-03846487 https://hal.science/hal-03846487

Submitted on 10 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Two different compounds in one crystal: di-μ-chlorobis[2-(5-methyl-pyridin-2-yl-κN)phenyl-κC(1)]diplatinum(II) and chloro(N,N-dimethylformamide-κO)[2-(5-methylpyridin-2-yl-κN)phenyl-κC(1)]platinum(II)

Isabelle Sasaki¹, Sonia Mallet-Ladeira^{2,3}

Corresponding author: isabelle.sasaki@u-bordeaux.fr

Abstract: The crystal structures of di- μ -chlorobis[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)] diplatinum(II) (1) and chloro(N,N-dimethylformamide- κ O)[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]platinum(II) (2) have been determined by means of the X-ray diffraction. Both complexes cocrystallize in the monoclinic space group C2/c with a = 39.5457(18) Å, b = 7.2482(4) Å, c = 21.2269(10) Å and $\beta = 121.460(2)^{\circ}$. The asymmetric unit consists of half a centrosymmetric compound (complex 1) and one full independent compound (complex 2).

Gaphical Abstract

The X-ray structures of co-crystallized di- μ -chlorobis[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]diplatinum(II) (1) as starting material and of the derivative chloro(N,N-dimethylformamide- κ O)[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]platinum(II) (2) are reported



Keywords: Platinum, cyclometalated, crystal structure, di-chloro bridged

Introduction

For decades, cyclometalated platinum complexes have been the subject of intensive research because of their outstanding luminescence properties [1,2]. Consequently, chemistry has been developed for diverse applications

[3] to include them as potential candidates in the elaboration of OLED [4], as memory devices [5], with NLO properties [6], as catalyst [7], as luminescent probes in biology [8] or as anticancer prodrugs [9,10]. To obtain orthoplatinated compounds, a number of syntheses start from K₂PtCl₄ and the bidentate C^N ligands leading to the dimer precursor $[Pt(\mu-Cl)(C^N)]_2$. The, neutral monomeric platinum(II) complexes Pt(II)(C^N)LCl (L and Cl as ligands) are produced by splitting the bridge upon dissolution to give complexes containing a solvent molecule as ancillary ligand or by using unidentate donor ligands [11,12,13]. Depending on the targeted complex, the ancillary ligand may be displaced [14] by a monodentate ligand of interest. In other cases, bidentate ligands L^L can convert the dimer to Pt(II)(C^N) (L^L) [15,16].

In a regular way, when the C^N is a phenylpyridine-type ligand, the neutral monomeric species are formed by dissolution of the dimer in DMSO [11, 12, 17]. RX analyses have showed that the DMSO is coordinated to the platinum atom through the S atom and is *trans* to the N atom of the cyclometalated complex (Scheme 1) [17].

To date, despite attempts by Sakai et al. [18] and although it has been routinely mentioned, the structure of the dimer has never been described. Herein we report the syntheses and structures of chloro(N,N-dimethylformamide- κ O)[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]platinum(II) obtained from the binuclear diµ-chlorobis[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]diplatinum(II).

Experimental

General

The ligand and complex 1 were synthesized as described in the literature [17]. K_2PtCl_4 (Strem), was purchased and used as received. DMF was of reagent grade. NMR spectra were recorded on a Bruker Avance 500 spectrometer.

Synthesis

Complex **2** was directly obtained by heating complex **1** in DMF at 50°C for 48h. Isolation of the complex was carried out after evaporation of the DMF, precipitation with diethyloxide, and filtration to give a pale green powder. ¹ H NMR (300, DMFd₇): δ 8.81 (dd, 1H, J = 0.8 Hz, J _{H-Pt} \approx 42 Hz, Py6); 8.08-8.14 (m, 2H); 8.02 (dd, 1H, J = 7 and 1.8 Hz); 7.74 (dd, 1H, J = 7.2 and 2.1Hz), 7.14-7.23 (m, 2H); 3.14 and 2.98 (s, Me of DMF); 2.46 (s, Me). ¹³C{¹H} NMR (75.47MHz, DMFd₇): δ 206.17; 162.48; 146.57; 145.08; 139.64; 134.89; 133.15; 132.03; 128.39; 123.21; 123.19; 118.80; 17.61. Due to the importance of the residual peaks of the non-deuterated DMF, it was not possible to identify the ¹³C signals from the methyl groups of the DMF ligand.

In order to get crystals, the synthesis was followed as described above, before isolation. After leaving the DMF solution at room temperature for few days, yellow crystals of suitable size and quality for single X-ray diffraction analysis were obtained.

X-ray Structure Determination

Diffraction data were collected at low temperature on a Bruker Kappa APEX II diffractometer using a graphitemonochromated Mo K α radiation ($\lambda = 0.71073$ Å) and equipped with an Oxford Cryosystems Cryostream cooler device. Phi- and Omega- scans were used. The structure was solved by direct methods with SHELXS 97 [19] and refined by full-matrix least-squares on F² [20]. All non-hydrogen atoms were refined with anisotropic displacement parameters. The hydrogen atoms were refined isotropically at calculated positions using a riding model with their isotropic displacement parameters constrained to be equal to 1.5 times the equivalent isotropic displacement parameters of their pivot atoms for terminal carbon sp^3 atoms and 1.2 times for all other carbon atoms. Crystal and experimental data are listed in Table 1.

Results and Discussion

Synthesis

The syntheses of complexes 2 (with DMF) and 3 (with DMSO) are depicted on Scheme 1. For discussion purposes, the synthesis of complex 3 is added.



Scheme 1 Preparation of complexes 2 and 3.

Depending on the nature of the solvent used, the bridge-splitting does not produce similar complexes. As usually described when using DMSO, the solvent molecule is coordinated to the platinum atom through the sulfur atom and is *trans* to the Pt-N bond [11,17] as depicted on Scheme 1.

Cyclometalated complexes with DMF as ancillary ligand are scarce [18, 21]. The synthesis of complex 2 was carried out by gentle heating of complex 1 in DMF for two days. It was isolated after elimination of the excess of solvent and precipitation with ether. The X-ray analysis revealed a κ O-DMF is coordination to the platinum atom (*vide infra*).

X-ray Structure

The starting complex di- μ -chlorobis[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]diplatinum(II) **1** and the final complex chloro(N,N-dimethylformamide- κ O)[2-(5-methyl-pyridin-2-yl- κ N)phenyl- κ C(1)]platinum(II) **2** cocrystallize in DMF.

Views of complexes 1 and 2 are shown on Fig. 1 and 2 respectively and selected bond lengths and angles are given in Table 2. In both complexes the platinum atom is located in a square-planar environment. The ligand coordinated to the platinum shows a certain degree of planarity: the dihedral angles between the two aromatics rings of the ligand in complexes 1 and 2 are 5.95° and 3.33° , respectively.

The Pt-C (1.981(9) - 1.954(7)Å), Pt-N (1.986(7) - 2.002(6) Å) and Pt-Cl (2.447(2)/2.317(2) - 2.298(2) Å) distances are in accordance with bond lengths reported in other analogous platinum complexes. Analysis of the molecular packing shows the presence of rows of **1** which are interleaved by rows of **2** (Fig. 3). Viewed along the crystallographic *c*-axis, the structure is composed of stacks of **1** and **2** complexes (Fig. 4).

In complex **1** we can notice that both Pt-Cl bonds are not similar: indeed, the bond length of Pt2-Cl2i is slightly shorter (2.317(2) Å) than the Pt2-Cl2 bond (2.447(2) Å), revealing the *trans* influence of the σ -carbon. This observation was previously reported on chloro-bridge dimers[22-25].

Thanks to the methyl substituent on the pyridine ring, the positions of the C and the N atoms are determined unambiguously. Examination of the structure reveals a κ O-DMF coordination *trans* to the Pt-C Only one structure of DMF coordinated to a platinum atom with this type of ligand has been published [18]. The authors described the cyclometalated platinum complex with phenylpyridine (ppy) as ligand. The locations of N(ppy) and C(ppy) donors were determined by examining their thermal parameters.

Starting from the same binuclear species, depending of which solvent is used for the following step, a striking difference between the complexes obtained is observed, as depicted on scheme 1. When using DMSO, the solvent is coordinated through the S atom, *trans* to the Pt-N bond, as with DMF the coordination of the solvent occurs by the oxygen atom on the Pt atom *trans* to the Pt-C bond.

Chemical formula	C ₂₄ H ₂₀ Cl ₂ N ₂ Pt ₂ ,2(C ₁₅ H ₁₇ ClN ₂ OPt)		
Formula weight	1741.15		
Crystal system	Monoclinic		
Space group	C 2/c		
T (K)	180(2)		
<i>a</i> (Å)	39.5457(18)		
<i>b</i> (Å)	7.2482(4)		
<i>c</i> (Å)	21.2269(10)		
β (°)	121.460(2)		
$V(Å^3)$	5190.0(5)		
Z	4		
pcalc (gcm ⁻³)	2.228		
F (000)	3264		
μ (mm ⁻¹)	11.002		
Crystal size (mm)	0,12 x 0,1 x 0,06		
θ range (°)	1.92-25.35		
hkl range	$-47 \le h \le 47$		
	$-8 \leq k \leq 8$		
	$-25 \le 1 \le 25$		
Reflections collected/unique	42024 / 4757		
Rint	0.0233		
Number of parameters	320		
R1 $[I \ge 2\sigma(I)]^a$	0.0338		
$wR^2 [I \ge 2\sigma(I)]^b$	0.0813		
R1 [all data]	0.0382		
wR^2 [all data]	0.0841		
Goodness of fit	1.191		
Δρmin/Δρmax (eÅ ⁻³)	2.72 / -2.01		
${}^{a}R1 = \Sigma Fo - Fc / \Sigma Fo $. ${}^{b}wR2 = [\Sigma [w(Fo^{2} - Fc^{2})^{2}] / \Sigma [w(Fo^{2})^{2}]]^{1/2}$.			

Table 1Crystal data, data collection and structure refinement of 1 and2

In Table **3** are gathered selected bonds of complexes **2**, **3** and of the complex described by Sakai [18]. At a first glance, structures of complex **2** and of Sakai's one are similar, with slightly longer bonds in the latter. The Pt-O bond length in complex **2** is 2.141(5) A°, longer than those reported for the Pt–O distances trans to a pyridyl donor (e.g. 2.051(6) A°)[26] as previously observed [18]. When comparing the bond lengths of Pt-Cl in complex **2** (2.298(2) Å) and in complex **3** (2.4054(10) Å) we can notice that it is longer (0.107 Å) when located *trans* to the carbon atom. All these observations highlight the strong *trans* effect of the σ -carbon, as previously mentioned. However, the bond lengths of the Pt-N in both complexes are similar regardless the nature of the atom *trans* to this bond: Pt-Cl for complex **2** and Pt-S for complex **3**.

Deacon and Edwards have studied the mechanism of addition of unidentate ligands to cycloplatinated complexes [11] by a systematic investigation of the bridge cleavage. In fact in the dimeric complex, due to the *trans* effect of the σ -carbon, the Pt-Cl bond *trans* to the aryl substituent should be cleaved first leading to the isomer with the halogen *cis* to the carbon ligand, as observed with DMF in complex **2**. In complex **3** the coordination of the DMSO *trans* to the Pt-N bond, results from a rearrangement to the more stable configuration (through an associative mechanism via a five-coordinate transition state or intermediate).



Fig. 1 Molecular structure of complex 1 with atom labeling scheme. Thermal ellipsoids are drawn at the 30% probability level. All hydrogen atoms are omitted. [Symmetry code : (i) -x,-y,1-z]



Fig. 2 Molecular structure of complex **2** with atom labeling scheme. Thermal ellipsoids are drawn at the 30% probability level. All hydrogen atoms are omitted.

Therefore it should be pointed out that further substitution of the DMF ancillary ligand by molecules of biological interest may lead to cyclometalated complexes not yet explored, as the usual starting complex is complex **3** to date [27, 28].

angles (°) for complexes 1 and 2		
1		
Pt2-C16	1.981(9)	
Pt2-N3	1.986(7)	
Pt2-Cl2	2.447(2)	
Pt2-Cl2 ⁱ	2.317(2)	
C16-Pt2-N3	80.9(3)	
C16-Pt2-Cl2 ⁱ	97.3(3)	
C16-Pt2-Cl2	177.4(2)	
N3-Pt2-Cl2 ⁱ	178.2(2)	
2		
Pt1-C1	1.954(7)	
Pt1-N1	2.002(6)	
Pt1-Cl1	2.298(2)	
Pt1-O1	2.141(5)	
C1-Pt1-N1	81.9(3)	
N1-Pt1-O1	89.8(2)	
C1-Pt1-O1	171.7(2)	
N1-Pt1-Cl1	178.46(17)	

Table 2 Selected bond lengths (Å) and

Symmetry code : (i) -x,-y,1-z



Fig. 3 View of the crystal packing along the b-axis. All hydrogen atoms are omitted. For clarity, the platinum atom of 1 is represented as black spheres and the platinum atom of 2 as grey spheres.



Fig. 4 View of the crystal packing along the *c*-axis. All hydrogen atoms are omitted. For clarity, the platinum atom of 1 is represented by black spheres and the platinum atom of 2 by grey spheres.

	2	[18]	3
Pt-C	1.954(7)	1.958(7); 1.974(8)	2.013(4)
Pt-N	2.002(6)	2.005(6); 2.010(6)	2.068(3)
Pt-Cl	2.298(2)	2.305(2); 2.313(2)	2.4054(10)
Pt-O	2.141(5)	2.158(5); 2.149(5)	-
Pt-S	-	-	2.2219(10)

Table 3 Selected bond lengths (Å) for complexes 2, 3 and from ref [18].

Conclusions

We were able to characterize the routinely mentioned dichloro bridge dinuclear cyclometalated platinum complex with a phenylpyridine-type ligand by X-ray analysis and the resulting mononuclear species after bridge-splitting with DMF. Since the number of cyclometalated platinum complexes with DMF as ancillary ligand are limited, the structural insight achieved from this study provides impulse for further exploration of this complex as starting material to produce new compounds in medicinal chemistry.

Supplementary Material

CCDC-2206545 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Center via <u>www.ccdc.cam.ac.uk/data_request/cif</u>.

Acknowledgements We thank CNRS for financial support.

Compliance with Ethical Standards

Conflict of Interest There is no conflict of interest/competing interests to declare.

References

[1] Brooks J, Babayan Y, Lamansky S, Djurovich PI, Tsyba I, Bau R, Thompson ME (2002) Synthesis and Characterization of Phosphorescent Cyclometalated Platinum Complexes. Inorg. Chem. https://doi.org/10.1021/ic0255508

[2] Kalinowski J, Fattori V, Cocchi M, Williams JAG (2011) Light-emitting devices based on organometallic platinum complexes as emitters. Coord. Chem. Rev. <u>https://doi:10.1016/j.ccr.2011.01.049</u>

[3] Haque A, El Moll H, Alenezi KM, Khan MS, Wong WY (2021) Functional Materials Based on Cyclometalated Platinum(II) β -Diketonate Complexes: A Review of Structure–Property Relationships and Applications. Materials <u>https://doi.org/10.3390/ma14154236</u>

[4] Cebrián C, Mauro M (2018) Recent advances in phosphorescent platinum complexes for organic lightemitting diodes. Beilstein J. Org. Chem. <u>https://doi.org/10.3762/bjoc.14.124</u>

[5] Poh WC, Au-Yeung HL, Chan AKW, Hong EYH, Cheng YH, Leung MY, Lai SL, Low KH, Yam VW (2021) Cyclometalated Platinum(II) Complexes with Donor-Acceptor-Containing Bidentate Ligands and Their Application Studies as Organic Resistive Memories. Chem. Asian J. <u>https://doi.org/10.1002/asia.202100897</u>

[6] Colombo A, Dragonetti C, Guerchais V, Roberto D (2021) An excursion in the second-order nonlinear optical properties of platinum complexes. Coord Chem Rev <u>https://doi.org/10.1016/j.ccr.2021.214113</u>

[7] Dobrynin MV, Sokolova EV, Kinzhalov MA, Smirnov AS, Starova GL, Kukushkin VY, Islamova (2021) RM Cyclometalated Platinum(II) Complexes Simultaneously Catalyze the Cross-Linking of Polysiloxanes and Function as Luminophores. ACS Appl. Polym. Mater. <u>https://doi.org/10.1021/acsapm.0c01190</u>

[8] Koo CK, Wong KL, Man CWY, Lam YW, So LKY, Tam HL, Tsao SW, Cheah KW, Lau KC, Yang YY. (2009) A Bioaccumulative Cyclometalated Platinum(II) Complex with Two-Photon-Induced Emission for Live Cell Imaging. Inorg. Chem. <u>https://doi.org/10.1021/ic801261h</u>

[9] Cutillas N, Yellol GS, de Haro C, Vicente C, Rodríguez V, Ruiz J (2013) Anticancer cyclometalated complexes of platinum group metals and gold. Coord. Chem. Rev. <u>https://doi.org/10.1016/j.ccr.2013.03.024</u>

[10] Babak MV, Pfaffeneder-Kmen, Meier-Menches SM, Legina MS, Theiner S, Licona C, Orvain C, Hejl M, Hanif M, Jakupec MA, Keppler BK, Gaiddon C, Hartinger CG (2018) Rollover Cyclometalated Bipyridine Platinum Complexes as Potent Anticancer Agents: Impact of the Ancillary Ligands on the Mode of Action. Inorg. Chem. <u>https://doi.org/10.1021/acs.inorgchem.7b03210</u>

[11] Black DStC, Deacon GB, Edwards GL (1994) Observations on the Mechanism of Halogen-Bridge Cleavage by Unidentate Ligands in Square Planar Palladium and Platinum Complexes Aust. J. Chem. https://doi.org/10.1071/CH9940217

[12] Godbert N, Pugliese T, Aiello I, Bellusci A, Crispini A, Ghedini M (2007) Efficient, Ultrafast, Microwave-Assisted Syntheses of Cycloplatinated Complexes. Eur. J. Inorg. Chem. <u>https://doi:10.1002/ejic.200700639</u>

[13] Edwards GL, Black DStC, Deacon GB, Wakelin LPG (2005) In vitro and in vivo studies of neutral cyclometallated complexes against murine leukæmias Can. J. Chem. <u>https://doi:10.1139/V05-109</u>

[14] Fernandez-Pampín N, Vaquero M, Gil T, Espino G, Fernandez D, García B, Busto N (2022) Distinct mechanism of action for antitumoral neutral cyclometalated Pt(II)-complexes bearing antifungal imidazolyl-based drugs. J. Inorg. Biochem. <u>https://doi.org/10.1016/j.jinorgbio.2021.111663</u>

[15] Spencer M, Santoro A, Freeman GR, Díez A, Murray PR, Torroba J, Whitwood AC, Yellowlees LJ, Williams JAG, Bruce DW. (2012) Phosphorescent, liquid-crystalline complexes of platinum(II): influence of the β -diketonate co-ligand on mesomorphism and emission properties. Dalton Trans. https://doi:10.1039/c2dt31525d [16] Kabir E, Wu CH, Wu JIC, Teets TS (2016) Heteroleptic Complexes of Cyclometalated Platinum with Triarylformazanate Ligands. Inorg. Chem. <u>https://doi:10.1021/acs.inorgchem.5b02595</u>

[17] Sasaki I, Bijani C, Ladeira S, Bourdon V, Faller P, Hureau C (2012) Interference of a new cyclometallated Pt compound with Cu binding to amyloid-β peptide. Dalton Trans. <u>https://doi/10.1039/c2dt12177h</u>

[18] Kobayashi M., Masaoka S., Sakai K. (2009) Synthesis, crystal structure, solution and spectroscopic properties, and hydrogen-evolving activity of [K(18-crown-6)][Pt(II)(2-phenylpyridinato)Cl₂ Photochem. Photobiol. Sci <u>https://doi.org/10.1039/B814939A</u>

[19] Sheldrick GM (1990) Phase annealing in SHELX-90: direct methods for larger structures. Acta Cryst. https://doi.org/10.1107/S0108767390000277

[20] Sheldrick GM (2015) Crystal structure refinement with SHELXL. Acta Cryst. https://doi.org/10.1107/S2053229614026540

[21] Yamaguchi S, Shinokubo H, Osuka A (2009) Double Cleavage of sp² C-H and sp³ C-H Bonds on One Metal Center: DMF-Appended Cyclometalated Platinum(II) and -(IV) Porphyrins. Inorg. Chem. https://doi:10.1021/ic8019149

[22] Cave GWV, Fanizzi FP, Deeth RJ, Errington W, Rourke JP (2000) C-H Activation Induced by Water. Monocyclometalated to Dicyclometalated: CANAC Tridentate Platinum Complexes. Organometallics. https://doi:10.1021/om9910423

[23] Crosby SH, Clarkson GJ, Deeth RJ, Rourke JP (2010) Platinum(IV) DMSO Complexes: Synthesis, Isomerization, and Agostic Intermediates. Organometallics. <u>https://doi:10.1021/om901087m</u>

[24] Shaw PA, Clarkson GJ, Rourke JP (2017) Reversible C–C bond formation at a triply cyclometallated platinum(IV) centre. Chem. Sci. <u>https://doi:10.1039/c7sc01361b</u>

[25] Ranieri AM, Burt LK, Stagni S, Zacchini S, Skelton BW, Ogden MI, Bissember AC, Massi M (2018) Anionic Cyclometalated Platinum(II) Tetrazolato Complexes as Viable Photoredox Catalysts. Organometallics. https://doi.org/10.1021/acs.organomet.8b00913

[26] Rochon FD, Kong PC, Melanso R (1980) The molecular and crystal structure of transdichloro(dimethylformamide)(2,6-lutidine)platinum(II). Can. J. Chem. <u>https://doi.org/10.1139/v80-016</u>

[27] Fereidoonnezhad M, Niazi M, Ahmadipour Z, Mirzaee T, Faghih Z, Faghih Z, Shahsavari HR (2017) Cyclometalated Platinum(II) Complexes Comprising 2-(Diphenylphosphino)pyridine and Various Thiolate Ligands: Synthesis, Spectroscopic Characterization, and Biological Activity Eur. J. Inorg. Chem. https://DOI:10.1002/ejic.201601

[28] Samouei H, Rashidi M, Heinemann FW (2014) Cyclometalated platinum (II) complexes containing monodentate phosphines: antiproliferative study J. Iran. Chem. Soc, <u>https://doi.org/10.1007/s13738-013-0389-z</u>

Affiliations

Isabelle Sasaki¹. Sonia Mallet-Ladeira^{2,3}

- 1 Université de Bordeaux, Institut des Sciences Moléculaires (CNRS UMR 5255) 33405 Talence, France
- 2 Laboratoire de Chimie de Coordination du CNRS, 205, route de Narbonne, BP 44099, F-3107 Toulouse Cedex 4, France
- 3 Institut de Chimie de Toulouse, Université de Toulouse, UPS, INPT, F-31077 Toulouse Cedex 4, France