



HAL
open science

A novel mechanism for left-right asymmetry establishment involving tissue remodeling and MyoID

Bénédicte Lefèvre, Marine Delvigne, Josué Vidal, Virginie Courtier-Orgogozo, Michael Lang

► **To cite this version:**

Bénédicte Lefèvre, Marine Delvigne, Josué Vidal, Virginie Courtier-Orgogozo, Michael Lang. A novel mechanism for left-right asymmetry establishment involving tissue remodeling and MyoID. 2023. hal-03848954

HAL Id: hal-03848954

<https://hal.science/hal-03848954>

Preprint submitted on 2 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title: A novel mechanism for left-right asymmetry establishment involving tissue remodeling and MyoID

Authors: Bénédicte M. Lefèvre^{1,2}, Marine Delvigne¹, Josué Vidal, Virginie Courtier-Orgogozo^{1*}, and Michael Lang^{1*}

Affiliations:

¹Université de Paris, CNRS, Institut Jacques Monod, F-75013 Paris, France.

²current address: Team “Stem Cells and Tissue Homeostasis”, Institut Curie, CNRS, UMR3215, INSERM U934, PSL Research University, 26 rue d’Ulm, 75248 Paris Cedex 05

*Correspondence to: michael.lang@ijm.fr, virginie.courtier@normalesup.org

#co-last authors

Abstract: How novel left-right asymmetries evolve in animals is unknown. *Drosophila pachea* males display a unique left-right asymmetry in the genital organs and mate in a right-sided position onto the female. In flies, male genitalia undergo a clockwise 360° rotation during development. To test if this tissue remodeling process is linked to asymmetry development in the male genitalia of *Drosophila pachea*, we developed CRISPR and transgenesis in this singular species. As in *D. melanogaster*, mutants of the MyoID gene homolog undergo reverse rotation of the developing genitalia. In addition, they have mirror-image asymmetric genitalia. Although their genital asymmetry is reversed mutants still adopt a right-sided copulation posture. Our results reveal a novel mechanism for establishing left-right asymmetry in animals, involving complex tissue remodeling and MyoID.

One Sentence Summary: *D. pachea* genital left-right size asymmetry is established through MyoID-mediated tissue remodelling and does not determine the direction of mating posture.

Main Text:

While the body of most bilaterian animals is superficially left-right symmetric, some internal organs display left-right asymmetries in position, shape or size (1). The generation of asymmetric organs during development starts with an initial symmetry-breaking step, which can involve several mechanisms: motile cilia and directional fluid flow, orientation of the cell division plane and distribution of molecules within cells (1). How over the course of evolution initially symmetrical organs become asymmetric remains elusive. Males of the fruitfly *Drosophila pachea* have an asymmetric phallus with a right-sided opening for sperm release (gonopore) and a pair of asymmetric external genital lobes, with the left lobe being 1.5 times longer than the right lobe (Fig. 1 A,B) (2–4). In addition, *D. pachea* males mate in a right-sided posture with the male being shifted about 6–8° towards the right side of the female (3–5). The lobes are considered an evolutionary novelty since they are not present in the three closely related sister species (6). The asymmetric genitalia of *D. pachea* likely evolved recently within the past 3–6

40 Ma from ancestral symmetry (6) and this species is therefore a suitable organism to search for the evolutionary origins of genital left-right asymmetry.

45 During development, the male genitalia of a wide range of shizophoran flies undergoes a 360° clockwise rotation (7, 8). In *D. melanogaster*, the direction of this rotation is driven by the myosin MyoID, encoded by the gene *myo31DF* (9), for simplicity here referred to as *myoID*. Loss of function of *myoID* in *D. melanogaster* leads to a counter-clockwise 360° genitalia rotation direction, due to the function of another counter-acting myosin, MyoIC (7, 9, 10). MyoID also controls the chirality of other organs, such as the gut and testes (9, 11–17). Furthermore, MyoID appears to be a conserved key factor among different mechanisms to establish left-right asymmetry in vertebrates and invertebrates (18, 19), through interaction with adherens junctions (10, 11), the actin cytoskeleton (10, 20, 21), planar cell polarity pathway members (13, 18) and Jun kinase pathway regulated apoptosis (12, 15, 22, 23). Ectopically expressed, MyoID and MyoIC are sufficient to induce *de novo* directional twisting at the cell, organ, or body scale in *D. melanogaster* (20). However, the evolutionary transitions to novel left-right size asymmetries remain to be uncovered at the molecular, cellular and tissue scale.

55 We wondered if MyoID is involved in the establishment of left-right asymmetry in *D. pachea* male genitalia. To test this, we generated a mutation in the coding sequence of the *D. pachea myoID* using CRISPR/Cas9 mediated gene editing (24, 25) (*myoID^{mut}* allele, Fig. S1). The induced mutation caused a frameshift in the seventh coding exon, leading to a premature stop codon. In homozygous *myoID^{mut/mut}* males, the length ratio of left and right genital lobes was reversed with a longer right lobe and a shorter left lobe in most (81/82) dissected individuals (Data S1, Figure 1 C). Furthermore, the phallus asymmetry was reversed in *myoID^{mut/mut}* males with the gonopore located at the left tip of the phallus (29/29 dissected genitalia), compared to heterozygous *myoID^{wt/mut}* and *myoID^{wt/wt}* males, where the gonopore was located consistently on the right side (8/8 and 13/13 dissected genitalia, respectively; Fig. S2). The locus *myoID* is autosomal (Fig. S1) and the mutation appears to be recessive since *myoID^{wt/mut}* males did not have visible alterations of male genitalia morphology (Fig. 1 B). Apart from phallus and lobe morphology, homozygous mutant males revealed a genitalia rotation phenotype, with varying orientations of the male genitalia in adults, with respect to the antero-posterior midline (Fig. 1 A,D). In wild-type *D. pachea*, male claspers and genital lobes are pointing towards the ventral side (Fig. 1 A), while *myoID^{mut/mut}* males revealed variable orientations of genitalia, which varied among individuals between 0 and 270° (Fig. 1 D). This latter orientation phenotype was similar to previous observations in *D. melanogaster myoID* mutants (9). Our data shows that MyoID is essential to determine left-right identity in *D. pachea* male genitalia, including external genital lobes and the male phallus. This function might be mediated through its role in determining the direction of male genitalia rotation. In one *myoID^{mut/mut}* males, genital lobe asymmetry was not reversed although genitalia were mis-oriented and in one individual lobes were strongly bent. However, it was unclear if their genitalia rotated clockwise or counter-clockwise during development. These mutants may have had remnant activity and expression of the *myoID^{mut}* allele, leading to a partial rotation in clockwise direction. Alternatively, other unidentified environmental or genetic factors may also contribute to the determination of genital asymmetry in *D. pachea* and were variable in our experiment.

80 To monitor genitalia rotation we inserted a DE-Cadherin::YFP fusion construct (DE-Cad::YFP) (Fig. S3) into the *myoID^{mut/mut}* stock using piggybac transposon mediated germline transformation (26) (Data S2, Fig. S4). We examined male genitalia rotation in pupae that were heterozygous for DE-Cad::YFP to reduce potentially deleterious effects of the piggybac insert

85 (Fig. S5). We observed that *myoID*^{wt/wt} and *myoID*^{wt/mut} males underwent clockwise genitalia
rotation direction (n = 9/9, 25/25, respectively), as observed in most schizophoran flies (7,
8) whereas in *myoID*^{mut/mut} males, genitalia rotated counter-clockwise (n = 9/9) (Fig. 1E, Data S1).
90 In 9 monitored males that had developed at least to the pharate adult stage, we were able to
dissect male genitalia and to determine the orientation and the lobe length ratio of male genitalia
(Fig. S2). All these *myoID*^{mut/mut} males with counter-clockwise genitalia rotation direction had a
reversed lobe length ratio with a longer right lobe and a shorter left lobe, although in one
observation lobes were only rudimentarily developed and the measurements were not informative
about the lobe length ratio (Fig. 1E). In contrast, *myoID*^{wt/wt} and *myoID*^{wt/mut} males had a wild-type
95 lobe ratio (Fig. 1E). We conclude that in *D. pachea* males MyoID controls genitalia rotation
direction, such as previously observed in *D. melanogaster* and that in addition, MyoID
determines left-right identity of genitalia.

Apart from its establishment during development, the role of genital asymmetry with
respect to reproduction is not well understood (27). Genital asymmetry has been hypothesized to
100 co-evolve or to evolve in response to changes in mating position (27). Alternatively, asymmetric
genitalia may cause lateralized mating behavior since the asymmetric parts mediate the coupling
of the female and male abdomen and therefore potentially direct the relative orientation of the
female and male bodies. We investigated if males with reversed genital asymmetry would adopt
a reverted copulation posture by monitoring copulation behavior (Fig. S6, Data S3) of *myoID*^{wt/wt},
myoID^{wt/mut}, or *myoID*^{mut/mut} males with wildtype *myoID*^{wt/wt} females. We annotated mating
105 behavior (Data S4) and copulation postures (Data S5, Fig. S7). A copulation was considered to
occur when the male mounted the female and achieved to settle into an invariant copulation
position for at least 30 seconds. Alternatively, males were sometimes observed to mount the
female, but leg and abdomen movements did not settle and eventually the male separated from
the female within a few minutes, which we considered to be a failed mounting attempt (106/140
110 total attempts, Fig. S6). Only 26% (10/38) of the couples with *myoID*^{mut/mut} males that mounted
the female also adopted a stable copulation position, compared to 11/11 and 13/13 for *myoID*^{wt/wt}
and *myoID*^{wt/mut} males, respectively (Fig. S6). Genitalia orientation angles of *myoID*^{mut/mut} males
that managed to adopt a stable copulation posture were found to deviate at most by 16.7° from
wild-type orientations, while it was higher in the majority of *myoID*^{mut/mut} males (18/29) that
115 failed to copulate (Fig. 1D, Fig. S6). Such defects in genitalia orientation were previously
reported to be detrimental for copulation and efficient genitalia coupling for *D. melanogaster*
myoID mutants (28). Regardless of the genotype and direction of genital asymmetry, *myoID*^{wt/wt},
myoID^{wt/mut} and *myoID*^{mut/mut} males consistently adopted a right-sided average copulation posture
(Fig. 2). We conclude that the right-sided mating posture in *D. pachea* does not depend on the
120 direction of male genital asymmetry.

Overall, our study reveals functional conservation of the unconventional myosin MyoID
to control clockwise genitalia rotation in developing *D. pachea* male genitalia. In addition, the
same protein got co-opted during evolution of *D. pachea* to contribute to determination of the
125 direction of a novel male genital asymmetry. The latter function may be mediated through the
genitalia rotation process itself since rotation direction and genital asymmetry direction are
perfectly associated in our live-imaging trials (Fig. 1E). Mechanical forces between rotating
genitalia and non-rotating outer tissue may act in opposite direction at the left and the right side
and in function of the sense of rotation. Male genital asymmetry in *D. pachea* does not determine
the posture itself since wildtype males and also mutant males with reversed genital asymmetry
130 consistently mate in a right-sided copulation posture. Right-sided mating is therefore female-

controlled or may be a hard-wired behavior that could have possibly favored the evolution of morphological asymmetry as an instructive cue to optimize genital contacts (27). This study provides the first experimental insights into *de novo* Evolution of left-right asymmetry. It involves recruitment of a conserved left-right regulator and is associated with a pre-existing tissue remodeling process.

References and Notes:

1. H. Hamada, P. Tam, *F1000Research*, in press, doi:10.12688/f1000research.21670.1.
2. S. Pitnick, W. Heed, New Species of Cactus-Breeding *Drosophila* (Diptera: Drosophilidae) in the Nannoptera Species Group. *Ann. Entomol. Soc. Am.* **87**, 307–310 (1994).
3. M. Lang, V. Orgogozo, Distinct copulation positions in *Drosophila pachea* males with symmetric or asymmetric: External genitalia. *Contrib. to Zool.* **81**, 87–94 (2012).
4. A. Acurio, F. T. Rhebergen, S. E. Paulus, V. Courtier-Orgogozo, M. Lang, Repeated evolution of asymmetric genitalia and right-sided mating behavior in the *Drosophila nannoptera* species group. *BMC Evol. Biol.* **19**, 109 (2019).
5. F. T. Rhebergen, V. Courtier-Orgogozo, J. Dumont, M. Schilthuisen, M. Lang, *Drosophila pachea* asymmetric lobes are part of a grasping device and stabilize one-sided mating. *BMC Evol. Biol.* **16**, 176 (2016).
6. M. Lang, M. Polihronakis Richmond, A. E. Acurio, T. A. Markow, V. Orgogozo, Radiation of the *Drosophila nannoptera* species group in Mexico. *J. Evol. Biol.* **27**, 575–584 (2014).
7. M. Suzanne, A. G. Petzoldt, P. Spéder, J.-B. Coutelis, H. Steller, S. Noselli, Coupling of Apoptosis and L/R Patterning Controls Stepwise Organ Looping. *Curr. Biol.* **20**, 1773–1778 (2010).
8. H. J. Feuerborn, Das Hypopygium ‘inversum’ und ‘circumversum’ der Dipteren. *Zool. Anz.* **55**, 89–213 (1922).
9. P. Spéder, G. Ádám, S. Noselli, Type ID unconventional myosin controls left–right asymmetry in *Drosophila*. *Nature.* **440**, 803–807 (2006).
10. A. G. Petzoldt, J.-B. Coutelis, C. Géminard, P. Spéder, M. Suzanne, D. Cerezo, S. Noselli, DE-Cadherin regulates unconventional Myosin ID and Myosin IC in *Drosophila* left-right asymmetry establishment. *Development.* **139**, 1874–1884 (2012).
11. K. Taniguchi, R. Maeda, T. Ando, T. Okumura, N. Nakazawa, R. Hatori, M. Nakamura, S. Hozumi, H. Fujiwara, K. Matsuno, Chirality in Planar Cell Shape Contributes to Left-Right Asymmetric Epithelial Morphogenesis. *Science (80- .)* **333**, 339–341 (2011).
12. S. Benitez, C. Sosa, N. Tomasini, A. Macias, Both JNK and apoptosis pathways regulate growth and terminalia rotation during *Drosophila* genital disc development. *Int. J. Dev. Biol.* **54**, 643–653 (2010).

13. N. González-Morales, C. Géminard, G. Lebreton, D. Cerezo, J.-B. Coutelis, S. Noselli, The Atypical Cadherin Dachshous Controls Left-Right Asymmetry in *Drosophila*. *Dev. Cell.* **33**, 675–689 (2015).
- 170 14. S. Hozumi, R. Maeda, K. Taniguchi, M. Kanai, S. Shirakabe, T. Sasamura, P. Spéder, S. Noselli, T. Aigaki, R. Murakami, K. Matsuno, An unconventional myosin in *Drosophila* reverses the default handedness in visceral organs. *Nature.* **440**, 798–802 (2006).
15. K. Taniguchi, S. Hozumi, R. Maeda, M. Ooike, T. Sasamura, T. Aigaki, K. Matsuno, D-JNK signaling in visceral muscle cells controls the laterality of the *Drosophila* gut. *Dev. Biol.* **311**, 251–263 (2007).
- 175 16. R. Hatori, T. Ando, T. Sasamura, N. Nakazawa, M. Nakamura, K. Taniguchi, S. Hozumi, J. Kikuta, M. Ishii, K. Matsuno, Left–right asymmetry is formed in individual cells by intrinsic cell chirality. *Mech. Dev.* **133**, 146–162 (2014).
17. T. Okumura, T. Sasamura, M. Inatomi, S. Hozumi, M. Nakamura, R. Hatori, K. Taniguchi, N. Nakazawa, E. Suzuki, R. Maeda, T. Yamakawa, K. Matsuno, Class I Myosins Have Overlapping and Specialized Functions in Left-Right Asymmetric Development in *Drosophila*. *Genetics.* **199**, 1183–1199 (2015).
- 180 18. T. Juan, C. Géminard, J.-B. Coutelis, D. Cerezo, S. Polès, S. Noselli, M. Fürthauer, Myosin1D is an evolutionarily conserved regulator of animal left–right asymmetry. *Nat. Commun.* **9**, 1942 (2018).
- 185 19. M. Tingler, S. Kurz, M. Maerker, T. Ott, F. Fuhl, A. Schweickert, J. M. LeBlanc-Straceski, S. Noselli, M. Blum, A Conserved Role of the Unconventional Myosin 1d in Laterality Determination. *Curr. Biol.* **28**, 810-816.e3 (2018).
20. G. Lebreton, C. Géminard, F. Lapraz, S. Pyrpassopoulos, D. Cerezo, P. Spéder, M. Ostap E, S. Noselli, Molecular to organismal chirality is induced by the conserved myosin 1D. *Science (80-.).* **362**, 949–952 (2018).
- 190 21. A. Chougule, F. Lapraz, I. Földi, D. Cerezo, J. Mihály, S. Noselli, The *Drosophila* actin nucleator DAAM is essential for left-right asymmetry. *PLOS Genet.* **16**, e1008758 (2020).
22. A. Macias, N. M. Romero, F. Martin, L. Suarez, A. L. Rosa, G. Morata, PVF1/PVR signaling and apoptosis promotes the rotation and dorsal closure of the *Drosophila* male terminalia. *Int. J. Dev. Biol.* **48**, 1087–1094 (2004).
- 195 23. R. Rousset, S. Bono-Lauriol, M. Gettings, M. Suzanne, P. Spéder, S. Noselli, The *Drosophila* serine protease homologue Scarface regulates JNK signalling in a negative-feedback loop during epithelial morphogenesis. *Development.* **137**, 2177–2186 (2010).
- 200 24. M. Jinek, K. Chylinski, I. Fonfara, M. Hauer, J. A. Doudna, E. Charpentier, A Programmable Dual-RNA–Guided DNA Endonuclease in Adaptive Bacterial Immunity. *Science (80-.).* **337**, 816–821 (2012).
25. A. R. Bassett, C. Tibbit, C. P. Ponting, J.-L. Liu, Highly Efficient Targeted Mutagenesis of *Drosophila* with the CRISPR/Cas9 System. *Cell Rep.* **4**, 220–228 (2013).

- 205 26. C. Horn, N. Offen, S. Nystedt, U. Häcker, E. A. Wimmer, *Genetics*, in press (available at <http://www.genetics.org/content/163/2/647.abstract>).
27. B. A. Huber, Mating positions and the evolution of asymmetric insect genitalia. *Genetica*. **138**, 19–25 (2010).
- 210 28. M. Inatomi, D. Shin, Y.-T. Lai, K. Matsuno, Proper direction of male genitalia is prerequisite for copulation in *Drosophila*, implying cooperative evolution between genitalia rotation and mating behavior. *Sci. Rep.* **9**, 210 (2019).
29. W. B. Heed, H. W. Kircher, Unique Sterol in the Ecology and Nutrition of *Drosophila* pachea. *Science* (80-.). **149**, 758–761 (1965).
- 215 30. A. Suvorov, B. Y. Kim, J. Wang, E. E. Armstrong, D. Peede, E. R. R. D'Agostino, D. K. Price, P. Wadell, M. Lang, V. Courtier-Orgogozo, J. R. David, D. Petrov, D. R. Matute, D. R. Schrider, A. A. Comeault, Widespread introgression across a phylogeny of 155 *Drosophila* genomes. *bioRxiv* (2021),
31. B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).
- 220 32. H. Oda, T. Uemura, Y. Harada, Y. Iwai, M. Takeichi, A *Drosophila* Homolog of Cadherin Associated with Armadillo and Essential for Embryonic Cell-Cell Adhesion. *Dev. Biol.* **165**, 716–726 (1994).
33. A. Pacquelet, L. Lin, P. Rørth, Binding site for p120/ δ -catenin is not required for *Drosophila* E-cadherin function in vivo . *J. Cell Biol.* **160**, 313–319 (2003).
- 225 34. D. G. Gibson, L. Young, R.-Y. Chuang, J. C. Venter, C. A. Hutchison, H. O. Smith, Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods.* **6**, 343–345 (2009).
35. O. Nagy, I. Nuez, R. Savaisaar, A. E. Peluffo, A. Yassin, M. Lang, D. L. Stern, D. R. Matute, J. R. David, V. Courtier-Orgogozo, Correlated Evolution of Two Copulatory Organs via a Single cis-Regulatory Nucleotide Change. *Curr. Biol.* **28**, 3450-3457.e13 (2018).
- 230 36. C. S. Thummel, A. M. Boulet, H. D. Lipshitz, Vectors for *Drosophila* P-element-mediated transformation and tissue culture transfection. *Gene.* **74**, 445–456 (1988).
37. M. V Matz, A. F. Fradkov, Y. A. Labas, A. P. Savitsky, A. G. Zaraisky, M. L. Markelov, S. A. Lukyanov, Fluorescent proteins from nonbioluminescent Anthozoa species. *Nat. Biotechnol.* **17**, 969–973 (1999).
- 235 38. H. Ochman, A. S. Gerber, D. L. Hartl, Genetic Applications of an Inverse Polymerase Chain Reaction. *Genetics.* **120**, 621–623 (1988).
39. S. P. Bainbridge, M. Bownes, Staging the metamorphosis of *Drosophila melanogaster*. *Development.* **66**, 57–80 (1981).
- 240 40. J. F. McAlpine, in *Manual of Nearctic Diptera* (McAlpine JF, Peterson BV, Shewell GE, Teskey HJ, Vockeroth JR, Wood DM, *Agricultur.*, 1981), vol. 1, pp. 9–63.

41. S. Pitnick, Operational sex ratios and sperm limitation in populations of *Drosophila* *pachea*. *Behav. Ecol. Sociobiol.* **33**, 383–391 (1993).
42. B. M. Lefèvre, D. Catté, V. Courtier-Orgogozo, M. Lang, Male genital lobe morphology affects the chance to copulate in *Drosophila* *pachea*. *BMC Ecol. Evol.* **21**, 23 (2021).

245

Acknowledgments: We thank Stephane Prigent for preparation of *Drosophila* genitalia. We thank the CNRS ImagoSeine Imaging platform at the Institut Jacques Monod for assistance in microscopy. We are grateful to Nicolas Gompel for initial help with *Drosophila* germline transfection. **Funding:** BL was supported by a pre-doctoral fellowship Sorbonne Paris Cité of the Université de Paris and by a fellowship from the Labex "Who am I?" ["Initiatives d'excellence", Idex ANR-18-IDEX-0001, ANR-11-LABX-0071]. This work was further supported by the CNRS, by a grant of the European Research Council under the European Community's Seventh Framework Program [FP7/2007-2013 Grant Agreement no. 337579] given to VCO and by a grant of the Agence Nationale pour la recherche [ANR-20-CE13-0006] given to ML. **Author contributions:** ML and VCO designed the study. ML and MD generated and characterized *Drosophila* CRISPR mutants and transgenic stocks. BL prepared the membrane marker construct. ML and BL performed time-lapse microscopy experiments. ML, JV and BL carried out mating assays. ML and JV genotyped flies, ML, BL and JV analyzed the data, ML and VCO wrote the manuscript. **Competing interests:** Authors declare no competing interests. **Data and materials availability:** Data supporting this article is available at the DRYAD database (doi:10.5061/dryad.bzkh189bd), except for DNA sequence data of the *D. pachea* *myoID* coding sequence, which was deposited at NCBI (accession number OM240650). Plasmid pBAC-ECFP-15xQUAS_TATA-mcd8-GFP-SV40 (Addgene 104878) was provided by addgene (MTA, addgene order 482018), and plasmid 3XP3::EYFP was a kind gift of D. Stern.

250

255

260

265

Supplementary Materials:

Materials and Methods

270

Figures S1-S7

Table S1

Movies S1-S2

External Databases S1-S5





Fig. 1. MyoID mutants display reversed genital asymmetry and genitalia rotation defects.

(A) Wild-type *D. pachea* male in ventral view. The orientation of genitalia is measured as an angle between the midpoint of male genital claspers relative to the male midline, as depicted by the dashed polar diagram. The scale is 100 μm . (B) External adult male genitalia of (B) a wild-type male *myoID*^{wt/wt} with asymmetric genital lobes at the ventral side (arrows). (C) External adult male genitalia of a *myoID*^{mut/mut} mutant male with reversed genital asymmetry. (B-C) Lobes are indicated by arrows and lobe lengths measurement landmark are shown as red dots. The scale is 100 μm . (D-E) Male genitalia orientation and lobe length asymmetry ratio in wild-type *myoID*^{wt/wt} males (blue), heterozygous *myoID*^{wt/mut} mutant males (grey) and homozygous *myoID*^{mut/mut} mutant males (orange). (D) The polar diagram illustrates genitalia orientation angles. (E) The direction of genital lobe asymmetry is associated with the direction of male genitalia rotation. Left and right lobe lengths are plotted for individual males monitored by time-lapse microscopy. Genitalia rotation direction is indicated as +: clockwise, and as -: counter-clockwise. The diagonal red dashed line indicates the 1:1 lobe length ratio. The star indicates an observation with rudimentarily developed lobes in form of buds on both sides.



290

Fig. 2. The right-sided *D. pachea* mating posture does not depend on the direction of male genital asymmetry. Mating angles (α) (Supplementary Materials) of *D. pachea* copulating with a male being either (A) *myoID*^{wt/wt} with the wild-type lobe length ratio, (B) *myoID*^{wt/mut} heterozygous males with wildtype lobe length ratio, (C) *myoID*^{mut/mut} males with reverted lobe asymmetry ratio. Females had a *myoID*^{wt/wt} genotype in all conditions. The hypothesis angle = 0 was rejected for each male genotype (GLM fit angle ~ genotype, all $p < 1 \times 10^{-10}$). Data acquisition and processing is illustrated in Figs. S6, S7. Positive angle values correspond to right-sided orientations of the male head relative to the female midline. Data points that correspond to the same trial are connected by grey lines. The number of trials are indicated by n, a drawing of the external male genitalia of each genotype is shown on the right.

295