



# Non trivial zeros of the Zeta function using the differential equations

W Oukil

## ► To cite this version:

W Oukil. Non trivial zeros of the Zeta function using the differential equations. 2023. hal-03849860v10

HAL Id: hal-03849860

<https://hal.science/hal-03849860v10>

Preprint submitted on 8 Dec 2023 (v10), last revised 20 May 2024 (v18)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Nontrivial zeros of the Zeta function using the differential equations

W. Oukil

Faculty of Mathematics.

University of Science and Technology Houari Boumediene.

BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

December 8, 2023

## Abstract

We investigate a relation between the differential equations and the Zeta function. We find a more flexible form of the Zeta function in the critical strip. We prove that all the nontrivial zeros lie on the critical line.

**Keywords:** Zeta function, Non trivial zeros, Riemann functional equation, Differential equations.

**AMS subject classifications:** 00A05

## 1 Main result

Consider the representation of the Riemann Zeta function  $\zeta$  defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -\frac{s}{1-s} - s \int_1^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}, \quad (1)$$

where  $\{t\}$  is the fractional part of the real  $t$ . In order to simplify the notation, denote  $B \subset \mathbb{C}$  the critical strip, defined as

$$B := \{s \in \mathbb{C} : \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}\},$$

and define the critical line as  $s \in B$  such that  $1 - 2\Re(s) = 0$ . We prove the following theorems,

**Theorem 1.** Consider the function  $\zeta$  given by the Equation (1). For every  $s \in B$  we have

$$\begin{aligned} \frac{\zeta(s)}{s(1+s)}t^s &= -\frac{1}{2}\left(\frac{t}{1-s} + \frac{1}{s}\right) \\ &+ 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (j2\pi)^{2k} t^{2k-1}}{2k-1-s}, \quad \forall t \in (0, 1]. \end{aligned} \quad (2)$$

**Theorem 2.** All the zeros of the Zeta function on  $B$ , lie on the critical line.

## 2 Main Proposition

For every  $s \in B$ , the Equation (1) is equivalent to,

$$\frac{\zeta(s)}{s} = -\frac{1}{1-s} - \int_1^{+\infty} u^{-1-s} \{u\} du.$$

The aim is to studies the differential equation of solutions the functions

$$t \mapsto \psi_s(z, t) := t^s \left[ z + \int_1^t u^{-1-s} \{u\} du \right], \quad z \in \mathbb{C}, \quad t \geq 1.$$

We focus only on the bounded solutions (there is a unique bounded solution. All other solutions diverge to infinity in norm). More precisely, the strategy to prove the Theorem 1, is to find the initial condition  $z_s \in \mathbb{C}$  such that  $\sup_{t \geq 1} |\psi_s(z_s, t)| < +\infty$ . Since  $\sup_{t \geq 1} |\psi_s(z, t)| < +\infty$  if and only if  $z = -\int_1^{+\infty} u^{-1-s} \{u\} du$ , we obtain  $-\int_1^{+\infty} u^{-1-s} \{u\} du = z_s$ .

For every  $s \in B$  we consider the following differential equation

$$\begin{aligned} \frac{d}{dt}x &= st^{-1}x + t^{-1}\{t\}, \\ t \in [1, +\infty)/\mathbb{N}, \quad x(1) &= z, \quad z \in \mathbb{C}, \quad x : [1, +\infty) \rightarrow \mathbb{C}. \end{aligned} \quad (3)$$

**Lemma 3.** Let be  $s \in B$ . For every  $z \in \mathbb{C}$  there exists a unique continuous solution  $\psi_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  of the differential equation (3). Further,

$$\psi_s(z, t) = t^s \left[ z + \int_1^t u^{-1-s} \{u\} du \right], \quad \forall t \geq 1.$$

*Proof.* Let be  $s \in B$  fixed. The function

$$t \mapsto t^s \int_1^t u^{-1-s} \{u\} du, \quad t \geq 1,$$

is  $C^\infty$  on  $[1, +\infty)/\mathbb{N}$  and continuous on  $[1, +\infty)$ . The Equation (3) is a non-homogeneous linear differential equation. For every  $z \in \mathbb{C}$  the unique continuous solution  $\psi_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  such that  $\psi_s(z, 1) = z$  is given by

$$\psi_s(z, t) = t^s \left[ z + \int_1^t u^{-1-s} \{u\} du \right], \quad \forall t \geq 1.$$

□

**Proposition 4.** Let be  $s \in B$ . Let be  $z \in \mathbb{C}$  and consider the continuous solution  $\psi_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  of the differential equation (3). Then  $\sup_{t \geq 1} |\psi_s(z, t)| < +\infty$ , if and only if

$$z = -\frac{1}{2} \frac{1}{s} + 2(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}.$$

In other words,

$$\begin{aligned} \int_1^{+\infty} u^{-1-s} \{u\} du &= \frac{1}{2} \frac{1}{s} \\ &\quad - 2(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}. \end{aligned}$$

**Lemma 5.** For every  $s \in B$ , there exists a unique  $z_s \in \mathbb{C}$  such that the solution  $\psi_s(z_s, t) : [1, +\infty) \rightarrow \mathbb{C}$  of the differential equation (3) satisfies  $\sup_{t \geq 1} |\psi_s(z_s, t)| < +\infty$ . Further,

$$z_s = - \int_1^{+\infty} u^{-1-s} \{u\} du, \quad \text{and } \psi_s(z_s, t) = -t^s \int_t^{+\infty} u^{-1-s} \{u\} du, \quad \forall t \geq 1.$$

*Proof.* Let be  $s \in B$  fixed. There is a unique bounded solution. All other solutions diverge to infinity in norm. Let be  $z \in \mathbb{C}$  and suppose that

$$\sup_{t \geq 1} |\psi_s(z, t)| < +\infty.$$

Since  $\Re(s) \in (0, 1)$ , by the Lemma 3, we have

$$\sup_{t \geq 1} |\psi_s(z, t)| < +\infty \implies \left| z + \int_1^{+\infty} u^{-1-s} \{u\} du \right| = 0,$$

Then

$$z_s = - \int_1^{+\infty} u^{-1-s} \{u\} du.$$

Prove that  $\sup_{t \geq 1} |\psi_s(z_s, t)| < +\infty$ . Since

$$\left| z_s + \int_1^{+\infty} u^{-1-s} \{u\} du \right| = 0,$$

then

$$z_s + \int_1^t u^{-1-s} \{u\} du = - \int_t^{+\infty} u^{-1-s} \{u\} du, \quad \forall t > 1.$$

By the Lemma 3, we get

$$\psi_s(z_s, t) = -t^s \int_t^{+\infty} u^{-1-s} \{u\} du, \quad \forall t \geq 1.$$

Then  $\sup_{t \geq 1} |\psi_s(z_s, t)| \leq \frac{1}{\Re(s)}$ .  $\square$

**Lemma 6.** Let be  $s \in B$ . For every  $z \in \mathbb{C}$  there exists a unique  $\tilde{z} \in \mathbb{C}$  such that the solution  $\psi_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  of the differential equation (3) satisfies

$$\psi_s(z, t) = \psi_s(z_s, t) + t^s \tilde{z}, \quad \forall t \geq 1.$$

where  $t \mapsto \psi_s(z_s, t)$  is the bounded solution given by the Lemma 5. In addition,

$$\tilde{z} = z + \int_1^{+\infty} u^{-1-s} \{u\} du.$$

*Proof.* Consider

$$\tilde{z} = z + \int_1^{+\infty} u^{-1-s} \{u\} du,$$

By the Lemma 3, we have

$$\psi_s(z, t) = -t^s \int_t^{+\infty} u^{-1-s} \{u\} du + t^s \tilde{z}, \quad \forall t \geq 1.$$

By the Lemma 5, we get

$$\psi_s(z, t) = \psi_s(z_s, t) + t^s \tilde{z}, \quad \forall t \geq 1.$$

$\square$

**Notation 7.** Denote

$$p(t) := \int_1^t \left( \{u\} - \frac{1}{2} \right) du, \quad \forall t \geq 1,$$

where we recall that  $\{u\}$  is the fractional part of the real  $u$ .

**Lemma 8.** *The function  $p$  is a continuous 1-periodic function and satisfies*

$$p(t) = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} (\cos(j2\pi t) - 1), \quad \forall t \geq 1.$$

*Proof.* The function  $u \mapsto \{u\}$  is 1-periodic, then there exists a continuous 1-periodic function  $p : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\int_1^t \{u\} du = (t-1) \int_1^2 \{u\} du + p(t), \quad \forall t \geq 1.$$

Since

$$\int_1^2 \{u\} du = \int_0^1 \{u\} du = \int_0^1 u du = \frac{1}{2},$$

we get

$$\int_1^t \{u\} du = (t-1)\frac{1}{2} + p(t), \quad \forall t \geq 1.$$

The function  $p$  is a piecewise  $C^\infty$ , continuous on  $\mathbb{R}$  and 1-periodic. By Dirichlet Theorem, the Fourier series

$$n \mapsto \sum_{k=-n}^n a_k \exp(ik2\pi t),$$

converge uniformly on  $\mathbb{R}_+$  to the function  $t \mapsto p(t)$ , where  $(a_k)_k \subset \mathbb{C}$  are the Fourier coefficients of the function  $p$ .

$$\int_0^t \left( \{v\} - \frac{1}{2} \right) dv = \sum_{j \in \mathbb{Z}} a_j \exp(ij2\pi t), \quad \forall t \geq 0.$$

By definition of the Fourier coefficients we have

$$\begin{aligned} a_j &= \int_0^1 \exp(-ij2\pi u) \left( \int_0^u \left( \{v\} - \frac{1}{2} \right) dv \right) du \\ &= \frac{1}{2} \int_0^1 \exp(-ij2\pi u) u(u-1) du = \frac{1}{(2j\pi)^2}, \quad \forall j \in \mathbb{Z}^*, \end{aligned}$$

and

$$a_0 = \frac{1}{2} \int_0^1 u(u-1) du = -\frac{1}{12} = -\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2}.$$

The function  $p$  satisfies

$$p(t) = \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2} (\exp(ij2\pi t) - 1), \quad \forall t \geq 0.$$

□

*Proof of the Proposition 4.* Let be  $s \in B$ . Let be  $z \in \mathbb{C}$  and consider the continuous solution  $\psi_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  of the differential equation (3). Define the function  $\tilde{\psi}_s(z, t) : [1, +\infty) \rightarrow \mathbb{C}$  as

$$\tilde{\psi}_s(z, t) := \int_1^t \psi_s(z, u) du, \quad t \geq 1.$$

The Equation (3) implies

$$\frac{d}{dt} \tilde{\psi}_s(z, t) = t^{-1}(1+s)\tilde{\psi}_s(z, t) + t^{-1} \int_1^t \{u\} du + t^{-1}z, \quad \forall t \geq 1.$$

By the Lemma 8 we have

$$\begin{aligned} \int_1^t \left( \{u\} - \frac{1}{2} \right) du &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \left( \cos(j2\pi t) - 1 \right), \\ &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} t^{2k}, \quad \forall t \geq 1. \end{aligned}$$

Then

$$\begin{aligned} \frac{d}{dt} \tilde{\psi}_s(z, t) &= t^{-1}(1+s)\tilde{\psi}_s(z, t) + \frac{1}{2}(1 - (1-2z)t^{-1}) \\ &\quad + 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} t^{2k-1}, \quad \forall t \geq 1. \end{aligned} \quad (4)$$

Use the fact  $|\tilde{\psi}_s(z, 1)| = 0$ , integrate, to obtain

$$\begin{aligned} \tilde{\psi}_s(z, t) &= \frac{1}{2} \left( -\frac{1}{s} t + \frac{1-2z}{1+s} \right) \\ &\quad + 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s} t^{2k} \\ &\quad - t^{1+s} \left[ \frac{1}{2} \left( -\frac{1}{s} + \frac{1-2z}{1+s} \right) + 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s} \right]. \end{aligned} \quad (5)$$

By Lemma 6 and the definition of  $\tilde{\psi}_s(z, t)$  we have

$$\begin{aligned} \tilde{\psi}_s(z, t) &= \int_1^t \psi_s(z, u) du \\ &= \int_1^t \psi_s(z_s, u) du - \frac{\tilde{z}}{1+s} + t^{1+s} \frac{\tilde{z}}{1+s}, \quad \forall t \geq 1. \end{aligned} \quad (6)$$

where  $t \mapsto \psi_s(z_s, t)$  is the bounded solution given by the Lemma 5 and where

$$\tilde{z} = z + \int_1^{+\infty} u^{-1-s} \{\psi_s(u)\} du. \quad (7)$$

Equations (5) and (6) implies

$$\begin{aligned} & 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s} t^{2k} \\ &= t^{1+s} \omega_s(z, \tilde{z}) - \frac{1}{2} \left( -\frac{1}{s} t + \frac{1-2z}{1+s} \right) + \int_1^t \psi_s(z_s, u) du - \frac{\tilde{z}}{1+s}. \end{aligned} \quad (8)$$

where

$$\begin{aligned} \omega_s(z, \tilde{z}) &:= \frac{\tilde{z}}{1+s} + \frac{1}{2} \left( -\frac{1}{s} + \frac{1-2z}{1+s} \right) \\ &+ 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}. \end{aligned}$$

The aim is to prove that  $|\omega_s(z, \tilde{z})| = 0$ . The convergence of the above series can be deduced by the approximation in the differential Equation (4), as

$$\left| \sum_{j=n}^{+\infty} \frac{1}{(j2\pi)^2} (\cos(j2\pi t) - 1) \right| \leq \frac{1}{4\pi^2} \sum_{j=n}^{+\infty} j^{-2}, \quad \forall n \geq 1.$$

In order to simplify the notation, denote

$$\theta(t) = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s} t^{2k}.$$

We have

$$\begin{aligned} & t \frac{d}{dt} \Re(\theta(t)) - (1 + \Re(s)) \Re(\theta(t)) + \Im(s) \Im(\theta(t)) \\ &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} t^{2k} \\ &= \int_1^t \left( \{\psi_s(u)\} - \frac{1}{2} \right) du, \end{aligned} \quad (9)$$

By Equation (8), we have

$$\theta(t) = t^{1+s} \omega_s(z, \tilde{z}) - \frac{1}{2} \left( -\frac{1}{s} t + \frac{1-2z}{1+s} \right) + \int_1^t \psi_s(z_s, u) du - \frac{\tilde{z}}{1+s}. \quad (10)$$

Since  $\sup_{t \geq 1} |\psi_s(z_s, t)| < +\infty$  and  $\Re(s) \in (0, 1)$ , the Equations (9) and(10) implies  $|\omega_s(z, \tilde{z})| = 0$ . In other words,

$$\begin{aligned} \frac{\tilde{z}}{1+s} &= -\frac{1}{2}\left(-\frac{1}{s} + \frac{1-2z}{1+s}\right) \\ &\quad - 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}, \end{aligned}$$

Using the Equation (7), we obtain

$$\begin{aligned} \int_1^{+\infty} u^{-1-s} \{u\} du &= \frac{1}{2} \frac{1}{s} \\ &\quad - 2(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}. \end{aligned}$$

By the Lemma 5, we have  $\sup_{t \geq 1} |\psi_s(z, t)| < +\infty$ , if and only if

$$\begin{aligned} z = z_s &= - \int_1^{+\infty} u^{-1-s} \{u\} du = -\frac{1}{2} \frac{1}{s} \\ &\quad + 2(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}. \end{aligned}$$

□

### 3 Proof of the Theorem 1

*Proof of the Theorem 1.* Let be  $s \in B$ . Recall the Equation (1),

$$\zeta(s) = -\frac{s}{1-s} - s \int_1^{+\infty} u^{-1-s} \{u\} du.$$

Thanks to Main Proposition 4,

$$\begin{aligned} \zeta(s) &= -\frac{s}{1-s} - \frac{1}{2}s \\ &\quad + 2s(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k}}{(2k)!} \frac{1}{2k-1-s}. \end{aligned} \tag{11}$$

In one hand, for every  $t \in (0, 1]$  we have

$$\begin{aligned} \int_t^1 u^{-2-s} \left[ \int_0^u (v - \frac{1}{2}) dv \right] du &= \frac{1}{2} \int_t^1 u^{-2-s} (u^2 - u) du \\ &= \frac{1}{2s(1-s)} - \frac{1}{2} t^{-s} \left( \frac{t}{1-s} + \frac{1}{s} \right). \end{aligned} \tag{12}$$

We recall that the function  $p : \mathbb{R} \rightarrow \mathbb{R}$ , defined as

$$p(t) := \int_0^t \left( \{u\} - \frac{1}{2} \right) du, \quad \forall t \in \mathbb{R}$$

is a continuous 1-periodic function and satisfies

$$\begin{aligned} p(t) &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \left( \cos(j2\pi t) - 1 \right) \\ &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k} t^{2k}}{(2k)!}, \quad \forall t \in \mathbb{R}. \end{aligned}$$

On the other hand, for every  $t \in (0, 1]$  we have

$$\begin{aligned} &\int_t^1 u^{-2-s} \left[ \int_0^u \left( v - \frac{1}{2} \right) dv \right] du \\ &= 2 \int_t^1 u^{-2-s} \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} (-1)^k \frac{(j2\pi)^{2k} u^{2k}}{(2k)!} du \\ &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (j2\pi)^{2k}}{2k-1-s} \\ &\quad - 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (j2\pi)^{2k} t^{2k-1-s}}{2k-1-s}. \end{aligned} \tag{13}$$

Using the Equation (11), the Equations (12) and (13) implies that

$$\begin{aligned} \frac{\zeta(s)}{s(1+s)} &= 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (j2\pi)^{2k} t^{2k-1-s}}{2k-1-s} \\ &\quad - \frac{1}{2} \left( \frac{t^{1-s}}{1-s} + \frac{t^{-s}}{s} \right), \quad \forall t \in (0, 1]. \end{aligned}$$

□

## 4 Proof of the Theorem 2

*Proof of the Theorem 2.* Let  $s \in B$ . In order to simplify the notation, denote

$$\forall t \in (0, 1] : \quad \varphi_s(n, t) := \sum_{j=1}^n f_s(j, t), \quad \forall n \in \mathbb{N}^*.$$

where

$$f_s(z, t) := z^{-1+s} \int_0^z u^{-2-s} (\cos(2\pi tu) - 1) du, \quad \forall z \in \mathbb{R}_+^*, \quad \forall t \in (0, 1]. \quad (14)$$

Since

$$f_s(z, t) = \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k z^{2k-2} (2\pi t)^{2k}}{2k-1-s}, \quad \forall z \in \mathbb{R}_+^*, \quad \forall t \in (0, 1],$$

by the Theorem 1, we get

$$t^{1+s} 2\pi^2 \frac{\zeta(s)}{s(1+s)} = \lim_{n \rightarrow +\infty} \varphi_s(n, t) - \pi^2 \left( \frac{t^2}{1-s} + \frac{t}{s} \right), \quad \forall t \in (0, 1]. \quad (15)$$

For every fixed  $t \in (0, 1]$ , use the Euler-Maclaurin summation formula, given by

$$\begin{aligned} \varphi_s(n, t) &= \frac{1}{2} [f_s(1, t) - f_s(n, t)] + \int_1^n f_s(z, t) dz \\ &\quad + \int_1^n (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz, \quad \forall n \in \mathbb{N}^*. \end{aligned}$$

By the integration part formula, we have

$$\begin{aligned} \int_1^n f_s(z, t) dz &= \frac{1}{s} n f_s(n, t) - \frac{1}{s} \int_0^1 z^{-2-s} (\cos(2\pi tz) - 1) dz \\ &\quad - \frac{1}{s} \int_1^n u^{-2} (\cos(2\pi tu) - 1) du, \quad \forall n \in \mathbb{N}^*. \end{aligned}$$

Then

$$\begin{aligned} \varphi_s(n, t) &= \frac{1}{2} [f_s(1, t) - f_s(n, t)] \\ &\quad + \frac{1}{s} n f_s(n, t) - \frac{1}{s} \int_0^1 z^{-2-s} (\cos(2\pi tz) - 1) dz \\ &\quad - \frac{1}{s} \int_1^n u^{-2} (\cos(2\pi tu) - 1) du \\ &\quad + \int_1^n (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz, \quad \forall n \in \mathbb{N}^*. \end{aligned} \quad (16)$$

Prove that  $\lim_{z \rightarrow +\infty} z f_s(z, t) = 0$ . In order to simplify the notation, denote

$$\begin{aligned} \phi_s(t, z) &:= -\frac{1}{2\pi t} \left[ (1+s) z f_s(z, t) + z^{-1} (\cos(2\pi tz) - 1) \right] \\ &= z^s \int_0^z u^{-1-s} \sin(2\pi tu) du, \quad \forall z \in [1, +\infty). \end{aligned} \quad (17)$$

We have

$$\phi_s(t, z) := \sum_{k \in \mathbb{N}^*} \frac{1}{(2k-1)!} \frac{(-1)^k z^{2k-1} (2\pi t)^{2k-1}}{2k-1-s}.$$

The function  $z \mapsto \phi_s(t, z)$  is the unique solution of the following deferential equation

$$\begin{aligned} \frac{d}{dz} \phi_s(t, z) &= sz^{-1} \phi_s(t, z) + z^{-1} \sin(2\pi t z), \\ \phi_s(t, 1) &= \sum_{k \in \mathbb{N}^*} \frac{1}{(2k-1)!} \frac{(-1)^k (2\pi t)^{2k-1}}{2k-1-s}, \quad \forall z \in [1, +\infty). \end{aligned}$$

Using the same method as in the proof of the Proposition 4, we prove that

$$\int_1^{+\infty} u^{-1-s} \sin(2\pi t u) du = - \sum_{k \in \mathbb{N}^*} \frac{1}{(2k-1)!} \frac{(-1)^k (2\pi t)^{2k-1}}{2k-1-s} = -\phi_s(t, 1),$$

and the solution,  $z \mapsto \phi_s(t, z)$  can be written as

$$\begin{aligned} \phi_s(t, z) &= -z^s \int_z^{+\infty} u^{-1-s} \sin(2\pi t u) du \\ &= \frac{(1+s)}{2\pi t} z^s \int_z^{+\infty} u^{-2-s} (\cos(2\pi t u) - \cos(2\pi t z)) du, \quad \forall z \geq 1. \end{aligned}$$

Then we have  $\lim_{z \rightarrow +\infty} |\phi_s(t, z)| = 0$ . By the Equation (17) we get  $\lim_{z \rightarrow +\infty} |zf_s(z, t)| = 0$ . The Equation (16), implies

$$\begin{aligned} \lim_{n \rightarrow +\infty} \varphi_s(n, t) &= \frac{s-2}{2s} \int_0^1 z^{-2-s} (\cos(2\pi t z) - 1) dz \\ &\quad - \frac{1}{s} \int_1^{+\infty} u^{-2} (\cos(2\pi t u) - 1) du \\ &\quad + \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz \end{aligned}$$

The Equation (15), implies

$$\begin{aligned} t^{1+s} 2\pi^2 \frac{\zeta(s)}{s(1+s)} &= \frac{s-2}{2s} \int_0^1 z^{-2-s} (\cos(2\pi t z) - 1) dz \\ &\quad - \frac{1}{s} \int_1^{+\infty} u^{-2} (\cos(2\pi t u) - 1) du \\ &\quad + \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz \\ &\quad - \pi^2 \left( \frac{t^2}{1-s} + \frac{t}{s} \right), \quad \forall t \in (0, 1]. \end{aligned}$$

Suppose that  $s \in B$  is a zero of the function  $\zeta$ . The above Equation implies that

$$\pi^2 t + \int_1^{+\infty} u^{-2} (\cos(2\pi t u) - 1) du = \delta(s, t), \quad \forall t \in (0, 1]. \quad (18)$$

where

$$\begin{aligned} \delta(s, t) := & -\pi^2 \frac{t^2 s}{1-s} + \frac{s-2}{2} \int_0^1 u^{-2-s} (\cos(2\pi t u) - 1) du \\ & + s \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz, \quad \forall t \in (0, 1]. \end{aligned} \quad (19)$$

Now we develop the last term of the right side of the previous equality. The aim is to find the following limit

$$\lim_{t \rightarrow 0} t^{-2} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz.$$

By the Equation (14), we have

$$\begin{aligned} \frac{d}{dz} f_s(z, t) &:= -(1-s) z^{-1} f_s(z, t) + z^{-3} (\cos(2\pi t z) - 1), \\ \frac{d}{dz} [z^{-1} f_s(z, t)] &= -(2-s) z^{-2} f_s(z, t) + z^{-4} (\cos(2\pi t z) - 1). \end{aligned}$$

Then

$$\begin{aligned} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz &= -(1-s) \int_1^{+\infty} (\{z\} - \frac{1}{2}) z^{-1} f_s(z, t) dz \\ &+ \int_1^{+\infty} (\{z\} - \frac{1}{2}) z^{-3} (\cos(2\pi t z) - 1) dz. \end{aligned}$$

Use the integration part for the first term of the right member.

$$\begin{aligned} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz &= -(1-s)(2-s) \int_1^{+\infty} p(z) z^{-2} f_s(z, t) z \\ &+ (1-s) \int_1^{+\infty} p(z) z^{-4} (\cos(2\pi t z) - 1) dz \\ &+ \int_1^{+\infty} (\{z\} - \frac{1}{2}) z^{-3} (\cos(2\pi t z) - 1) dz. \end{aligned}$$

where

$$p(z) = \int_1^z (\{u\} - \frac{1}{2}) du = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \left( \cos(j2\pi z) - 1 \right).$$

Use the integration part part for the last term of the right member

$$\begin{aligned} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz &= -(1-s)(2-s) \int_1^{+\infty} p(z) z^{-2} f_s(z, t) z \\ &\quad - (2+s) \int_1^{+\infty} p(z) z^{-4} (\cos(2\pi t z) - 1) dz \\ &\quad - 2\pi t \int_1^{+\infty} p(z) z^{-3} \sin(2\pi t z) dz. \end{aligned}$$

By Lebesgue Theorem, we get

$$\begin{aligned} \lim_{t \rightarrow 0} t^{-2} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz \\ = - (1-s)(2-s) \int_1^{+\infty} p(z) z^{-2} \lim_{t \rightarrow 0} t^{-2} f_s(z, t) z \\ - (2+s) \int_1^{+\infty} p(z) z^{-4} \lim_{t \rightarrow 0} t^{-2} (\cos(2\pi t z) - 1) dz \\ - 2\pi \int_1^{+\infty} p(z) z^{-3} \lim_{t \rightarrow 0} t^{-1} \sin(2\pi t z) dz. \end{aligned}$$

We get

$$\begin{aligned} \lim_{t \rightarrow 0} t^{-2} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz \\ = - 2\pi^2 (1-s)(2-s) \int_1^{+\infty} p(z) z^{-3+s} \int_0^z u^{-s} du dz \\ - 2(2+s)\pi^2 \int_1^{+\infty} p(z) z^{-2} dz \\ - 4\pi^2 \int_1^{+\infty} p(z) z^{-2} dz. \end{aligned}$$

Equivalent to

$$\lim_{t \rightarrow 0} t^{-2} \int_1^{+\infty} (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz = 4\pi^2 \int_1^{+\infty} p(z) z^{-2} du dz. \quad (20)$$

Thanks to the Riemann functional equation, the point  $1 - \bar{s}$  is a zero of the function  $\zeta$ . The Equation (18), implies

$$\pi^2 t + \int_1^{+\infty} u^{-2} (\cos(2\pi t u) - 1) du = \delta(1 - \bar{s}, t), \quad \forall t \in (0, 1].$$

Then

$$|\delta(s, t) - \delta(1 - \bar{s}, t)| = 0, \quad \forall t \in (0, 1].$$

Implies

$$\lim_{t \rightarrow 0} t^{-2} |\delta(s, t) - \delta(1 - \bar{s}, t)| = 0, \quad \forall t \in (0, 1].$$

Since

$$\lim_{t \rightarrow 0} t^{-2} \left[ -\pi^2 \frac{t^2 s}{1-s} + \frac{s-2}{2} \int_0^1 u^{-2-s} (\cos(2\pi t u) - 1) du \right] = 2\pi^2,$$

which does not depend on  $s$ . By definition of the function  $\delta$  in the Equation (19), we get

$$\begin{aligned} 0 &= \lim_{t \rightarrow 0} t^{-2} |\delta(s, t) - \delta(1 - \bar{s}, t)| \\ &= \lim_{t \rightarrow 0} t^{-2} \left| \int_0^{+\infty} (\{z\} - \frac{1}{2}) \left[ s \frac{d}{dz} f_s(z, t) - (1 - \bar{s}) \frac{d}{dz} f_{1-\bar{s}}(z, t) \right] dz \right| \end{aligned}$$

By the Equation (20), we obtain

$$\begin{aligned} 0 &= \lim_{t \rightarrow 0} t^{-2} |\delta(s, t) - \delta(1 - \bar{s}, t)| \\ &= 4\pi^2 \int_1^{+\infty} p(z) z^{-2} du z \left| s - (1 - \bar{s}) \right| \\ &= -4\pi^2 |1 - 2\Re(s)| \int_1^{+\infty} p(z) z^{-2} du z. \end{aligned}$$

□

## References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).