



Functional equation and nontrivial zeros of the Zeta function

W Oukil

► To cite this version:

| W Oukil. Functional equation and nontrivial zeros of the Zeta function. 2023. hal-03849860v11

HAL Id: hal-03849860

<https://hal.science/hal-03849860v11>

Preprint submitted on 18 Dec 2023 (v11), last revised 20 May 2024 (v18)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Functional equation and nontrivial zeros of the Zeta function

W. Oukil

Faculty of Mathematics.

University of Science and Technology Houari Boumediene.
BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

December 18, 2023

Abstract

We obtain a more flexible functional equation of the Zeta function. Together with the Riemann functional equation, we prove that any non-trivial zeros s of the function ζ satisfies $2\Re(s) \geq 1$.

Keywords: Zeta function, Non trivial zeros, Riemann functional equation, Differential equations.

AMS subject classifications: 00A05

1 Main results

Consider the representation of the Riemann Zeta function ζ defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -\frac{s}{1-s} - s \int_1^{+\infty} \{u\} u^{-1-s} du, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}, \quad (1)$$

where $\{u\}$ is the fractional part of the real u . In order to simplify the notation, denote $B \subset \mathbb{C}$ the critical strip, defined as

$$B := \{s \in \mathbb{C} : \Re(s) \in (0, 1), \Im(s) \in \mathbb{R}\}.$$

We prove the following theorems,

Theorem 1. Consider the function ζ given by the Equation (1). For every $s \in B$ we have

$$\begin{aligned} \frac{2\pi^2\zeta(s)}{1+s} &= \alpha(\zeta, 1-\bar{s})\beta_-(s) \\ &\quad + \alpha(\zeta, s)\left[\beta_+(s) - \beta_-(s)\right], \quad 2\Re(s) < 1, \\ \frac{2\pi^2\zeta(1-\bar{s})}{2-\bar{s}} &= \alpha(\zeta, 1-\bar{s})\beta_+(1-\bar{s}) \\ &\quad + \alpha(\zeta, s)\left[\beta_-(s) - \beta_-(1-\bar{s})\right], \quad 2\Re(s) > 1. \end{aligned}$$

where

$$\begin{aligned} \alpha(\zeta, s) &:= -\frac{\zeta(\bar{s})}{\bar{s}} - \frac{1}{1-\bar{s}}, \quad \forall s \in B, \\ \beta_-(s) &:= s(1-s) \int_0^1 u^{-2-s}(\cos(2\pi u) - 1)du, \quad \forall s \in B, \\ \beta_+(s) &:= s(1-s) \int_0^{+\infty} u^{-2-s}(\cos(2\pi u) - 1)du, \quad \forall s \in B \end{aligned}$$

Thanks to the Riemann functional equation, we obtain the following theorem,

Theorem 2. Let be $s \in B$ and suppose that s is a zero of the function ζ . Then $2\Re(s) \geq 1$.

2 Main Proposition

We prove in this section the following Proposition

Proposition 3. For every $s \in B$ and $t \in (0, 1)$ we have

$$\frac{2\pi^2}{1+s}\zeta(s)t^{-\frac{1}{2}+s} - \frac{2\pi^2}{2-\bar{s}}\zeta(1-\bar{s})t^{\frac{1}{2}-\bar{s}} = \delta(s, t) - \delta(1-\bar{s}, t),$$

where

$$\begin{aligned} \delta(s, t) &:= -\pi^2 \frac{s}{1-s} t^{\frac{1}{2}} + (1-s)t^{-\frac{3}{2}} \int_1^{+\infty} z^{-2-s}(\cos(2\pi tz) - 1)dz \\ &\quad + st^{-\frac{3}{2}} \int_1^{+\infty} \{z\} \frac{d}{dz} f_s(z, t) dz, \quad \forall s \in B, \forall t \in (0, 1]. \end{aligned}$$

and where

$$s \in B, \forall z \in \mathbb{R}_+^*, \forall t \in (0, 1) : \\ f_s(z, t) = -z^{-1+s} \int_z^{+\infty} u^{-2-s} (\cos(2\pi ut) - 1) du$$

Notation 4. Denote

$$p(t) := \int_1^t \left(\{u\} - \frac{1}{2} \right) du, \quad \forall t \geq 1,$$

where we recall that $\{u\}$ is the fractional part of the real u .

Lemma 5. *The function p is a continuous 1-periodic function and satisfies*

$$p(t) = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \left(\cos(j2\pi t) - 1 \right), \quad \forall t \geq 1.$$

Proof. The function $u \mapsto \{u\}$ is 1-periodic, then there exists a continuous 1-periodic function $p : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\int_1^t \{u\} du = (t-1) \int_1^2 \{u\} du + p(t), \quad \forall t \geq 1.$$

Since

$$\int_1^2 \{u\} du = \int_0^1 \{u\} du = \int_0^1 u du = \frac{1}{2},$$

we get

$$\int_1^t \{u\} du = (t-1) \frac{1}{2} + p(t), \quad \forall t \geq 1.$$

The function p is a piecewise C^∞ , continuous on \mathbb{R} and 1-periodic. By Dirichlet Theorem, the Fourier series

$$n \mapsto \sum_{k=-n}^n a_k \exp(ik2\pi t),$$

converge uniformly on \mathbb{R}_+ to the function $t \mapsto p(t)$, where $(a_k)_k \subset \mathbb{C}$ are the Fourier coefficients of the function p .

$$\int_0^t \left(\{v\} - \frac{1}{2} \right) dv = \sum_{j \in \mathbb{Z}} a_j \exp(ij2\pi t), \quad \forall t \geq 0.$$

By definition of the Fourier coefficients we have

$$\begin{aligned} a_j &= \int_0^1 \exp(-ij2\pi u) \left(\int_0^u \left(\{v\} - \frac{1}{2} \right) dv \right) du \\ &= \frac{1}{2} \int_0^1 \exp(-ij2\pi u) u(u-1) du = \frac{1}{(2j\pi)^2}, \quad \forall j \in \mathbb{Z}^*, \end{aligned}$$

and

$$a_0 = \frac{1}{2} \int_0^1 u(u-1) du = -\frac{1}{12} = -\sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2}.$$

The function p satisfies

$$p(t) = \sum_{j \in \mathbb{Z}^*} \frac{1}{(j2\pi)^2} \left(\exp(ij2\pi t) - 1 \right), \quad \forall t \geq 0.$$

□

Lemma 6. Consider the function ζ given by the Equation (1). For every $s \in B$ we have

$$\begin{aligned} 2\pi^2 \frac{\zeta(s)}{1+s} t^{1+s} &= -\pi^2 \left(\frac{s}{1-s} t^2 + t \right) \\ &\quad - s \sum_{j \in \mathbb{N}^*} j^{-1-s} \int_j^{+\infty} u^{-2-s} (\cos(2\pi ut) - 1) du, \quad \forall t \in (0, 1). \end{aligned}$$

Proof. By the Equation (1). For every $s \in B$ we have

$$\begin{aligned} -\frac{\zeta(s)}{s} &= \frac{1}{1-s} t^{1-s} + \int_t^{+\infty} \{u\} u^{-1-s} du \\ &= \frac{1}{1-s} t^{1-s} + \frac{1}{2s} t^{-s} + \int_t^{+\infty} (\{u\} - \frac{1}{2}) u^{-1-s} du, \quad \forall t \in (0, 1). \end{aligned}$$

Use the integration by part formula

$$-\frac{\zeta(s)}{s} = \frac{1}{1-s} t^{1-s} + \frac{1}{2s} t^{-s} + (1+s) \int_t^{+\infty} p(u) u^{-2-s} du - p(t) t^{-1-s},$$

where the function $p(u)$ is given by the Lemma 5. Use the fact $p(t) = \frac{1}{2}(t^2 - t)$ for every $t \in (0, 1)$ we get

$$\begin{aligned} -\frac{\zeta(s)}{s} &= \frac{1}{1-s} t^{1-s} + \frac{1}{2s} t^{-s} + (1+s) \int_t^{+\infty} p(u) u^{-2-s} du \\ &\quad - \frac{1}{2}(t^2 - t) t^{-1-s}, \quad \forall t \in (0, 1). \end{aligned}$$

Since

$$p(t) = 2 \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} (\cos(j2\pi t) - 1), \quad \forall t \in \mathbb{R}.$$

Use the fact

$$\begin{aligned} & \lim_{n \rightarrow +\infty} \left| \int_t^{+\infty} u^{-2-s} \sum_{j=n}^{+\infty} \frac{1}{(j2\pi)^2} (\cos(j2\pi u) - 1) du \right| \\ & \leq \int_t^{+\infty} u^{-2-s} du \lim_{n \rightarrow +\infty} \sum_{j=n}^{+\infty} \frac{1}{(j2\pi)^2} = 0, \end{aligned}$$

then

$$\begin{aligned} \zeta(s) &= -\frac{s}{2} \frac{1+s}{1-s} t^{1-s} - \frac{1}{2} (1+s) t^{-s} \\ &\quad - 2s(1+s) \sum_{j \in \mathbb{N}^*} \frac{1}{(j2\pi)^2} \int_t^{+\infty} u^{-2-s} (\cos(j2\pi u) - 1) du \quad \forall t \in (0, 1). \end{aligned}$$

Use the change of variable $u \mapsto u = tj^{-1}v$ to obtain

$$\begin{aligned} \zeta(s) &= -\frac{s}{2} \frac{1+s}{1-s} t^{1-s} - \frac{1}{2} (1+s) t^{-s} \\ &\quad - \frac{s}{2\pi^2} (1+s) t^{-1-s} \sum_{j \in \mathbb{N}^*} j^{-1-s} \int_j^{+\infty} u^{-2-s} (\cos(2\pi ut) - 1) du. \end{aligned}$$

□

Proof of the Proposition 3. Let $s \in B$ fixed. In order to simplify the notation, denote

$$\forall t \in (0, 1] : \quad \varphi_s(n, t) := \sum_{j=1}^n f_s(j, t), \quad \forall n \in \mathbb{N}^*.$$

where

$$f_s(z, t) = -z^{-1+s} \int_z^{+\infty} u^{-2-s} (\cos(2\pi ut) - 1) du, \quad \forall z \in \mathbb{R}_+^*, \forall t \in (0, 1), \quad (2)$$

by the Lemma 6, we get

$$\frac{2\pi^2 \zeta(s)}{1+s} t^{1+s} = s \lim_{n \rightarrow +\infty} \varphi_s(n, t) + \pi^2 \left(\frac{s}{1-s} t^2 + t \right), \quad \forall t \in (0, 1). \quad (3)$$

For every fixed $t \in (0, 1]$, use the Euler-Maclaurin summation formula, given by

$$\begin{aligned}\varphi_s(n, t) &= \frac{1}{2}[f_s(1, t) - f_s(n, t)] + \int_1^n f_s(z, t) dz \\ &\quad + \int_1^n (\{z\} - \frac{1}{2}) \frac{d}{dz} f_s(z, t) dz, \quad \forall n \in \mathbb{N}^*,\end{aligned}$$

or,

$$\begin{aligned}\varphi_s(n, t) &= f_s(1, t) - \frac{1}{2}f_s(n, t) + \int_1^n f_s(z, t) dz \\ &\quad + \int_1^n \{z\} \frac{d}{dz} f_s(z, t) dz, \quad \forall n \in \mathbb{N}^*.\end{aligned}$$

Using the Equation (2) and the integration by parts formula, we obtain

$$\begin{aligned}\int_1^n f_s(z, t) dz &= \frac{1}{s} n f_s(n, t) + \frac{1}{s} \int_1^{+\infty} z^{-2-s} (\cos(2\pi t z) - 1) dz \\ &\quad - \frac{1}{s} \int_1^n u^{-2} (\cos(2\pi t u) - 1) du, \quad \forall n \in \mathbb{N}^*.\end{aligned}$$

Then

$$\begin{aligned}\varphi_s(n, t) &= f_s(1, t) - \frac{1}{2}f_s(n, t) \\ &\quad + \frac{1}{s} n f_s(n, t) + \frac{1}{s} \int_1^{+\infty} z^{-2-s} (\cos(2\pi t z) - 1) dz \\ &\quad - \frac{1}{s} \int_1^n u^{-2} (\cos(2\pi t u) - 1) du \\ &\quad + \int_1^n \{z\} \frac{d}{dz} f_s(z, t) dz, \quad \forall n \in \mathbb{N}^*.\end{aligned}$$

The Equation (2) implies $\lim_{z \rightarrow +\infty} z f_s(z, t) = 0$. By consequence,

$$\begin{aligned}\lim_{n \rightarrow +\infty} \varphi_s(n, t) &= \frac{1-s}{s} \int_1^{+\infty} z^{-2-s} (\cos(2\pi t z) - 1) dz \\ &\quad - \frac{1}{s} \int_1^{+\infty} z^{-2} (\cos(2\pi t z) - 1) dz \\ &\quad + \int_1^{+\infty} \{z\} \frac{d}{dz} f_s(z, t) dz\end{aligned}$$

The Equation (3), implies

$$\frac{2\pi^2\zeta(s)}{1+s}t^{1+s} = -\pi^2t - \int_1^{+\infty} z^{-2}(\cos(2\pi tz) - 1)dz + \delta(s, t), \quad \forall t \in (0, 1].$$

where

$$\begin{aligned} \delta(s, t) := & -\pi^2 \frac{t^2 s}{1-s} + (1-s) \int_1^{+\infty} z^{-2-s} (\cos(2\pi tz) - 1) dz \\ & + s \int_1^{+\infty} \{z\} \frac{d}{dz} f_s(z, t) dz, \quad \forall t \in (0, 1]. \end{aligned}$$

□

3 Proof of the Theorem 1

Proof of the Theorem 1. Let be $s \in B$. Thanks to the Proposition 3 we have

$$\frac{2\pi^2}{1+s}\zeta(s)t^{-\frac{1}{2}+s} - \frac{2\pi^2}{2-\bar{s}}\zeta(1-\bar{s})t^{\frac{1}{2}-\bar{s}} = \delta(s, t) - \delta(1-\bar{s}, t), \quad (4)$$

where

$$\begin{aligned} \delta(s, t) := & -\pi^2 \frac{s}{1-s} t^{\frac{1}{2}} + (1-s) t^{-\frac{3}{2}} \int_1^{+\infty} z^{-2-s} (\cos(2\pi tz) - 1) dz \\ & + st^{-\frac{3}{2}} \int_1^{+\infty} \{z\} \frac{d}{dz} f_s(z, t) dz, \quad \forall t \in (0, 1]. \end{aligned} \quad (5)$$

and where

$$\begin{aligned} s \in B, \quad & \forall z \in \mathbb{R}_+^*, \quad \forall t \in (0, 1) : \\ f_s(z, t) = & -z^{-1+s} \int_z^{+\infty} u^{-2-s} (\cos(2\pi ut) - 1) du \end{aligned} \quad (6)$$

Since

$$\lim_{t \rightarrow 0} t^{-\frac{3}{2}} \left[\pi^2 \frac{t^2 s}{1-s} + (1-s) \int_0^1 u^{-2-s} (\cos(2\pi tu) - 1) du \right] = 0$$

then by definition of the function δ in the Equation (5) and the Equation (4), we get

$$\begin{aligned} 2\pi^2 \lim_{t \rightarrow 0} & \left[\frac{\zeta(s)}{1+s} t^{-\frac{1}{2}+s} - \frac{\zeta(1-\bar{s})}{2-\bar{s}} t^{\frac{1}{2}-\bar{s}} \right] \\ & = \lim_{t \rightarrow 0} t^{-\frac{3}{2}} \int_1^{+\infty} \{z\} \left[s \frac{d}{dz} f_s(z, t) - (1-\bar{s}) \frac{d}{dz} f_{1-\bar{s}}(z, t) \right] dz. \end{aligned} \quad (7)$$

From the Equation (6) , we have

$$f_s(z, t) = -t^{1+s} z^{-1+s} \int_{zt}^{+\infty} v^{-2-s} (\cos(2\pi v) - 1) dv, \quad \forall z \in \mathbb{R}_+^*, \quad \forall t \in (0, 1),$$

Then

$$\begin{aligned} \frac{d}{dz} f_s(z, t) &= t^{1+s} (1-s) z^{-2+s} \int_{zt}^{+\infty} v^{-2-s} (\cos(2\pi v) - 1) dv \\ &\quad + z^{-3} (\cos(2\pi zt) - 1), \quad \forall z \in \mathbb{R}_+^*, \quad \forall t \in (0, 1), \end{aligned}$$

In order to simplify the notation, denote

$$\alpha_s(v) := \int_v^{+\infty} \{u\} u^{-1-\bar{s}} du, \quad v \geq 1, \quad s \in B, \quad (8)$$

and denote

$$\begin{aligned} \beta(s) &:= s(1-s) \int_0^{+\infty} v^{-2-s} (\cos(2\pi v) - 1) dv \\ \eta_s(u) &:= -s(1-s) \int_0^u v^{-2-s} (\cos(2\pi v) - 1) dv, \quad u > 0, \quad s \in B, \quad (9) \end{aligned}$$

The Equation (7), implies that

$$\begin{aligned} 2\pi^2 \lim_{t \rightarrow 0} t^{-\frac{1}{2}+s} &\left[\frac{\zeta(s)}{1+s} - \beta(s) \alpha_{1-\bar{s}}(1) - \int_1^{+\infty} \{z\} z^{-2+s} \eta_s(zt) dz \right] \\ &= \lim_{t \rightarrow 0} t^{\frac{1}{2}-\bar{s}} \left[\frac{\zeta(1-\bar{s})}{2-\bar{s}} - \beta(1-\bar{s}) \alpha_s(1) - \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \eta_{1-\bar{s}}(zt) dz \right] \\ &\quad + (2\Re(s) - 1) t^{-\frac{3}{2}} \int_1^{+\infty} \{z\} z^{-3} (\cos(2\pi zt) - 1) dz. \end{aligned}$$

Use the fact

$$\begin{aligned} \lim_{t \rightarrow 0} t^{-\frac{3}{2}} &\left| \int_1^{+\infty} \{z\} z^{-3} (\cos(2\pi zt) - 1) dz \right| \\ &= \lim_{t \rightarrow 0} t^{\frac{1}{2}} \left| \int_t^{+\infty} \left\{ \frac{z}{t} \right\} z^{-3} (\cos(2\pi z) - 1) dz \right| \\ &\leq \lim_{t \rightarrow 0} t^{\frac{1}{2}} \left| \int_1^{+\infty} \left\{ \frac{z}{t} \right\} z^{-3} (\cos(2\pi z) - 1) dz \right| \\ &\quad + \lim_{t \rightarrow 0} t^{\frac{1}{2}} \left| \int_t^1 \left\{ \frac{z}{t} \right\} z^{-3} (\cos(2\pi z) - 1) dz \right| \\ &\leq 2\pi \lim_{t \rightarrow 0} t^{\frac{1}{2}} \left[\int_1^{+\infty} z^{-3} dz + 2\pi \int_t^1 z^{-1} dz \right] \\ &= 0 \end{aligned}$$

then

$$\begin{aligned} & 2\pi^2 \lim_{t \rightarrow 0} t^{-\frac{1}{2}+s} \left[\frac{\zeta(s)}{1+s} - \beta(s)\alpha_{1-\bar{s}}(1) - \int_1^{+\infty} \{z\} z^{-2+s} \eta_s(zt) dz \right] \\ &= \lim_{t \rightarrow 0} t^{\frac{1}{2}-\bar{s}} \left[\frac{\zeta(1-\bar{s})}{2-\bar{s}} - \beta(1-\bar{s})\alpha_s(1) - \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \eta_{1-\bar{s}}(zt) dz \right]. \end{aligned}$$

For every $s \in B$ such that $2\Re(s) < 1$ we obtain

$$\frac{\zeta(s)}{1+s} - \beta(s)\alpha_{1-\bar{s}}(1) = \lim_{t \rightarrow 0} t^{-\frac{1}{2}+s} \int_1^{+\infty} \{z\} z^{-2+s} \eta_s(zt) dz. \quad (10)$$

By definition of the function η_s in the Equation (14), we have

$$\begin{aligned} & t^{-\frac{1}{2}+s} \int_1^{+\infty} \{z\} z^{-2+s} \eta_s(zt) dz \\ &= -s(1-s)t^{-\frac{1}{2}+s} \int_1^{+\infty} \{z\} z^{-2+s} \int_0^{zt} v^{-2-s} (\cos(2\pi v) - 1) dv dz \\ &= s(1-s)t^{-\frac{1}{2}+s} \int_1^{+\infty} \{z\} z^{-2+s} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (2\pi)^2 k (zt)^{2k-1-s}}{2k-1-s} dz \\ &= s(1-s)t^{\frac{1}{2}} \int_1^{+\infty} \{z\} z^{-2+s} \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k (2\pi)^2 k z^{2k-3} t^{2k-2}}{2k-1-s} dz. \end{aligned} \quad (11)$$

Since

$$\begin{aligned} & s(1-s)z^{-2+s} \int_0^{zt} v^{-2-s} (\cos(2\pi v) - 1) dv \\ &= s(1-s) \sum_{k \in \mathbb{N}^*} \frac{1}{(2k)!} \frac{(-1)^k z^{2k-3} (2\pi t)^{2k}}{2k-1-s} \\ &= \sum_{k \in \mathbb{N}^*} \frac{(2k-2)(2k-1)}{(2k)!} \frac{(-1)^k z^{2k-3} (2\pi t)^{2k}}{2k-1-s}. \end{aligned}$$

In order to simplify the notation, denote

$$X(s) := 2\pi^2 \frac{\zeta(s)}{1+s} - \beta(s)\alpha_s(1), \quad \forall s \in B. \quad (12)$$

The Equations (7) and (11), implies that

$$\begin{aligned} & \lim_{t \rightarrow 0} \left[X(s)t^{-\frac{1}{2}+s} - X(1-\bar{s})t^{\frac{1}{2}-\bar{s}} \right] \\ &= (1-2\Re(s)) \lim_{t \rightarrow 0} t^{\frac{1}{2}-\bar{s}} \left[\int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz \right]. \end{aligned} \quad (13)$$

where

$$\xi_x(\mu) := \sum_{k \geq 2} \frac{(2k-1)(2k-2)}{(2k)!} \frac{(-1)^k (2\pi)^{2k} \mu^{2k-2+\bar{x}}}{(2k-2+\bar{x})(2k-1-s)}, \quad \forall x \in B. \quad (14)$$

Using the fact

$$\frac{d^2}{du^2} \left[u^{-1} (\cos(2\pi u) - 1) \right] = \sum_{k \geq 2} \frac{(2k-1)(2k-2)}{(2k)!} (-1)^k (2\pi)^{2k} u^{2k-3},$$

then ξ_s can be written as,

$$\xi_s(v) = \int_0^v z^{-2+2\Re(s)} \int_0^z u^{1-s} \frac{d^2}{du^2} \left[u^{-1} (\cos(2\pi u) - 1) \right] du dz, \quad (15)$$

For a fixed $t \in (0, 1)$ we have

$$\begin{aligned} \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz &= \alpha_s(1) \xi_s(1) \\ &\quad + \int_1^{+\infty} \{z\} z^{-1-\bar{s}} (\xi_s(zt) - \xi_s(1)) dz. \end{aligned}$$

Use the integration part formula for the last term of the right side of the last equality, we get

$$\int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz = \alpha_s(1) \xi_s(1) - t \int_1^{+\infty} \alpha_s(z) \xi'_s(zt) dz. \quad (16)$$

By the Equation (15), we have

$$\xi'_s(v) = v^{-2+2\Re(s)} \int_0^v u^{1-s} \frac{d^2}{du^2} \left[u^{-1} (\cos(2\pi u) - 1) \right] du.$$

In order to simplify the notation, denote

$$\theta(u) := \frac{d^2}{du^2} \left[u^{-1} (\cos(2\pi u) - 1) \right], \quad \forall u > 0, \quad (17)$$

and denote

$$\beta_x(v) := \int_0^v u^{1-x} \theta(u) du, \quad v > 0, \quad x \in B. \quad (18)$$

Then

$$\xi'_s(v) = v^{-2+2\Re(s)} \beta_s(v).$$

In order to simplify the notation, denote

$$\tilde{\alpha}_x(v) := - \int_v^{+\infty} \alpha_x(u) u^{-2+2\Re(x)} du \quad v \geq 1, \quad x \in B.$$

The Equation (16) implies

$$\begin{aligned} \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz &= \alpha_s(1) \xi_s(1) + \tilde{\alpha}_s(1) \beta_s(1) t^{-1+2\Re(s)} \\ &\quad + t^{-1+2\Re(s)} \int_1^{+\infty} \alpha_s(z) z^{-2+2\Re(s)} (\beta_s(zt) - \beta_s(1)) dz. \end{aligned}$$

Use the integration part formula for the last term of the right side of the last equality, we obtain

$$\begin{aligned} \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz &= \alpha_s(1) \xi_s(1) + \tilde{\alpha}_s(1) \beta_s(1) t^{-1+2\Re(s)} \\ &\quad + t^{1+\bar{s}} \int_1^{+\infty} \tilde{\alpha}_s(z) z^{1-s} \theta(zt) dz. \end{aligned} \quad (19)$$

By definition of the function θ in the Equation (17), we have

$$\begin{aligned} \theta(u) &= \frac{d^2}{du^2} [u^{-1}(\cos(2\pi u) - 1)] \\ &= 2u^{-3}(\cos(2\pi u) - 1) - 4\pi^2 u^{-1} \cos(2\pi u), \end{aligned}$$

That implies

$$\begin{aligned} &\lim_{t \rightarrow 0} t^{\frac{3}{2}} \left| \int_1^{+\infty} \tilde{\alpha}_s(z) z^{1-s} \theta(zt) dz \right| \\ &\leq 2 \lim_{t \rightarrow 0} t^{\frac{3}{2}} \left| \int_1^{+\infty} \tilde{\alpha}_s(z) z^{-2-s} (\cos(2\pi u) - 1) dz \right| \\ &\quad + 4\pi^2 \lim_{t \rightarrow 0} t^{\frac{3}{2}} \left| \int_1^{+\infty} \tilde{\alpha}_s(z) z^{-s} \cos(2\pi u) dz \right| \\ &= 2 \lim_{t \rightarrow 0} t^{\frac{3}{2}} \left| \int_1^{+\infty} \tilde{\alpha}_s(z) z^{-2-s} (\cos(2\pi u) - 1) dz \right| \\ &\quad + |s(1+s)| \lim_{t \rightarrow 0} t^{\frac{3}{2}} \left| \int_1^{+\infty} \tilde{\alpha}_s(z) z^{-2-s} (\cos(2\pi u) - 1) dz \right| \\ &\leq (2 + |s(1+s)|) \lim_{t \rightarrow 0} t^{\frac{1}{2}} \int_1^{+\infty} |\tilde{\alpha}_s(z)| z^{-2-r} dz = 0. \end{aligned}$$

The Equation (19), implies

$$\lim_{t \rightarrow 0} \left[t^{\frac{1}{2}-\bar{s}} \int_1^{+\infty} \{z\} z^{-1-\bar{s}} \xi_s(zt) dz - V(s, t) \right] = 0.$$

where

$$V(s, t) := \alpha_s(1) \xi_s(1) t^{\frac{1}{2}-\bar{s}} + \tilde{\alpha}_s(1) \beta_s(1) t^{-\frac{1}{2}+s}.$$

The Equation (13) and by the notation of X in the Equation (12), we get

$$\begin{aligned} 2\pi^2 \frac{\zeta(s)}{1+s} &= \beta(s) \alpha_s(1) + (1 - 2\Re(s)) \tilde{\alpha}_s(1) \beta_s(1), \quad 2\Re(s) < 1, \\ 2\pi^2 \frac{\zeta(1-\bar{s})}{2-\bar{s}} &= \beta(1-\bar{s}) \alpha_{1-\bar{s}}(1) + (1 - 2\Re(s)) \alpha_s(1) \xi_s(1), \quad 2\Re(s) > 1. \end{aligned}$$

By definition of the function ζ in the Equation (1) and the function α_s in the Equation (8), we have

$$\begin{aligned} \alpha_s(1) &= \int_1^{+\infty} \{u\} u^{-1-\bar{s}} du = -\frac{\zeta(\bar{s})}{\bar{s}} - \int_0^1 \{u\} u^{-1-\bar{s}} du \\ &= -\frac{\zeta(\bar{s})}{\bar{s}} - \frac{1}{1-\bar{s}}, \quad \forall s \in B, \end{aligned}$$

same,

$$(1 - 2\Re(s)) \tilde{\alpha}_x(1) = \alpha_{1-\bar{s}}(1) - \alpha_s(1)$$

Recall that, the function β_s is given by the Equation (18), and we have

$$\beta_s(1) := \int_0^1 u^{1-s} \theta(u) du = s(1-s) \int_0^1 u^{-2-s} (\cos(2\pi u) - 1) du, \quad \forall s \in B.$$

Finally recall that ξ_s is given by the Equation (15), using the integration by part formula, we obtain

$$\begin{aligned} (1 - 2\Re(s)) \xi_s(1) &= \int_0^1 u^{1-s} \theta(u) du + \int_0^1 z^{\bar{s}} \theta(u) du \\ &= \beta_s(1) - \beta_{1-\bar{s}}(1), \quad \forall s \in B. \end{aligned}$$

□

4 Proof of the Theorem 2

Proof of the Theorem 2. Let be $x \in B$ such that $2\Re(x) < 1$ and suppose that x is a zero of the function ζ . Thanks to the Riemann functional equation the point $1 - \bar{x}$ is also a zero. By the first item of the Theorem 1, we obtain

$$\frac{1 - 2\Re(x)}{x} \beta_-(x) = -\beta_+(x),$$

where we recall

$$\begin{aligned}\beta_-(s) &:= s(1-s) \int_0^1 u^{-2-s} (\cos(2\pi u) - 1) du, \quad \forall s \in B, \\ \beta_+(s) &:= s(1-s) \int_0^{+\infty} u^{-2-s} (\cos(2\pi u) - 1) du, \quad \forall s \in B\end{aligned}$$

Then

$$\frac{1 - \bar{x}}{x} \int_0^1 u^{-2-x} (\cos(2\pi u) - 1) du = - \int_1^{+\infty} u^{-2-x} (\cos(2\pi u) - 1) du.$$

We apply the second item to the point $1 - \bar{x}$ we get

$$\begin{aligned}- \int_1^{+\infty} u^{-2-x} (\cos(2\pi u) - 1) du &= \frac{\bar{x}(1 - \bar{x})^2}{x^2(1 - x)} \int_0^1 u^{-3 + \bar{x}} (\cos(2\pi u) - 1) du \\ &\quad + \frac{1 - 2\Re(x)}{x} \int_0^1 u^{-2-x} (\cos(2\pi u) - 1) du.\end{aligned}$$

Implies

$$\begin{aligned}x^2(1 - x) \int_0^1 u^{-2-x} (\cos(2\pi u) - 1) du \\ = \bar{x}(1 - \bar{x})^2 \int_0^1 u^{-3 + \bar{x}} (\cos(2\pi u) - 1) du\end{aligned}$$

which can be written as

$$x^2(1 - x) \sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}}{2k - 1 - x} = \bar{x}(1 - \bar{x})^2 \sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}}{2k - 2 + \bar{x}},$$

or again

$$\begin{aligned}\sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}(2k - 2)(2k - 1)^2}{2k - 1 - s} \\ = \sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}(2k - 2)(2k - 1)^2}{2k - 2 + \bar{s}}.\end{aligned}$$

Implies

$$(1 - 2\Re(s)) \sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}(2k-2)(2k-1)^2}{(2k-1-s)(2k-2+\bar{s})} = 0.$$

Since $2\Re(s) < 1$ then

$$\sum_{k \in \mathbb{N}^*} \frac{(-1)^k}{(2k)!} \frac{(2\pi)^{2k}(2k-2)(2k-1)^2}{(2k-1-s)(2k-2+\bar{s})} = 0.$$

For the imaginary part, we obtain

$$\sum_{k \in \mathbb{N}^*} (-1)^k u_{s,k} = 0, \quad u_{s,k} := \frac{(2\pi)^{2k}}{(2k)!} \frac{(4k-3)(2k-2)(2k-1)^2}{|(2k-1-s)(2k-2+\bar{s})|^2}. \quad (20)$$

Since

$$\forall k \geq 12 : \quad u_{s,k} > u_{s,k+1}.$$

Then

$$\sum_{k=12}^{+\infty} (-1)^k u_{s,k} > 0.$$

Since

$$u_{s,2k} - u_{s,2(k+1)} > u_{s,13-2k} - u_{s,12-2k} \quad \forall k \in \llbracket 1, 2 \rrbracket \quad \text{and} \quad u_{s,6} > u_{s,7},$$

then $\sum_{k=2}^{11} (-1)^k u_{s,k} > 0$. We deduce that $\sum_{k \in \mathbb{N}^*} (-1)^k u_{s,k} > 0$. Contradiction with the Equation (20). \square

References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).