



# Non trivial zeros of the Zeta function

W Oukil

## ► To cite this version:

| W Oukil. Non trivial zeros of the Zeta function. 2023. hal-03849860v2

HAL Id: hal-03849860

<https://hal.science/hal-03849860v2>

Preprint submitted on 15 Jan 2023 (v2), last revised 26 Apr 2024 (v18)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Non trivial zeros of the Zeta function

W. Oukil

Faculty of Mathematics.

University of Science and Technology Houari Boumediene.

BP 32 EL ALIA 16111 Bab Ezzouar, Algiers, Algeria.

January 15, 2023

## Abstract

We prove that  $|\zeta(\frac{1}{2} + r + i\tau)| \neq 0$  for every  $\tau \in \mathbb{R}^*$  and  $r \in (-\frac{1}{2}, 0)$ , where  $\zeta$  is the Riemann Zeta function.

**Keywords:** Zeta function, non trivial zeros, Riemann hypothesis.

**AMS subject classifications:** 00A05

## 1 Main result

Consider the representation of the Riemann Zeta function  $\zeta$  defined by the Abel summation formula [[1], page 14 Equation 2.1.5] as

$$\zeta(s) := -s \int_0^{+\infty} \frac{\{t\}}{t^{1+s}} dt, \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*, \quad (1)$$

where  $\{t\}$  is the fractional part of the real  $t$ . We prove the following Theorem.

**Theorem 1.** Consider the function  $\zeta$  defined by the Equation (1). For every  $\tau \in \mathbb{R}^*$  and  $r \in (-\frac{1}{2}, 0)$  we have

$$|\zeta(\frac{1}{2} + r + i\tau)| \neq 0.$$

By the Riemann functional equation we deduce that any *non trivial zero* of the Zeta function has a real part equal to  $\frac{1}{2}$ , where the *non trivial zeros* of the function  $\zeta$  are defined in the following sense

**Definition 2.** Consider the function  $\zeta$  defined by the Equation (1). Let be  $s \in \mathbb{C}$ . We say that  $s$  is a *non trivial zero of the function  $\zeta$*  if

$$|\zeta(s)| = 0 \quad \text{and} \quad \Re(s) \in (0, 1), \quad \Im(s) \in \mathbb{R}^*.$$

## 2 Proof of the Theorem 1

**Notation 3.** In order to simplify the notation, for every  $r \in (-\frac{1}{2}, 0)$  denote

$$f_r(u) := \{u\}u^{-\frac{1}{2}-r}, \quad u > 0.$$

By the Equation (1), remark that

$$\lim_{t \rightarrow +\infty} \left| \int_0^t u^{-i\tau-1} f_r(u) du \right| := \left| \frac{\zeta(\frac{1}{2} + r + i\tau)}{\frac{1}{2} + r + i\tau} \right|.$$

**Notation 4.** Denote

$$p(t) := \int_0^t \left( \{u\} - \frac{1}{2} \right) du, \quad \forall t \geq 0,$$

where we recall that  $\{u\}$  is the fractional part of the real  $u$ . Denote

$$\omega := 2 \max_{t \in [0,1]} |p(t)|.$$

**Lemma 5.** *The function  $p$  is a continuous 1-periodic function.*

*Proof.* The function  $u \mapsto \{u\}$  is 1-periodic, then there exists a continuous 1-periodic function  $p : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$\int_0^t \{u\} du = t \int_0^1 \{u\} du + p(t), \quad \forall t \geq 0.$$

Since

$$\int_0^1 \{u\} du = \int_0^1 u du = \frac{1}{2},$$

we get

$$\int_0^t \{u\} du = t \frac{1}{2} + p(t), \quad \forall t \geq 0.$$

□

**Notation 6.** Let  $g : \mathbb{R}_+ \rightarrow \mathbb{R}$  be a continuous function. For every  $t > s > 0$  we denoted

$$\int_{s,k}^t g(u) du = \underbrace{\int_s^t \frac{1}{u_1} \int_s^{u_1} \frac{1}{u_2} \cdots \int_s^{u_{k-1}} \frac{1}{u_k} g(u_k) du_k du_{k-1} \cdots du_1}_{k \text{ times}}, \quad \forall k \in \mathbb{N}^*.$$

For every  $r \in (-\frac{1}{2}, 0)$  and  $\tau \in \mathbb{R}^*$  denote

$$\begin{aligned}\underline{\Psi}_{\tau,r}(k,t) &= \int_{\frac{1}{t},k}^t u^{i\tau} \int_0^u v^{-i\tau-1} f_r(v) dv du, & \forall t > 1, \forall k \geq 1, \\ \overline{\Psi}_{\tau,r}(k,t) &= \int_{\frac{1}{t},k}^t u^{i\tau} \int_u^{+\infty} v^{-i\tau-1} f_r(v) dv du, & \forall t > 1, \forall k \geq 1.\end{aligned}$$

*Proof of the Theorem 1.* Let be  $\tau \in \mathbb{R}^*$  and  $r \in (-\frac{1}{2}, 0)$ . Prove the Theorem by contradiction. Suppose that

$$\left| \zeta\left(\frac{1}{2} + r + i\tau\right) \right| = 0.$$

By the Equation (1) we have

$$\left| \frac{\zeta(\frac{1}{2} + r + i\tau)}{\frac{1}{2} + r + i\tau} \right| = \left| \int_0^{+\infty} u^{-i\tau-1} f_r(u) du \right|.$$

Then

$$\left| \int_0^{+\infty} u^{-i\tau-1} f_r(u) du \right| = 0,$$

Implies

$$\int_0^t u^{-i\tau-1} f_r(u) du = - \int_t^{+\infty} u^{-i\tau-1} f_r(u) du, \quad \forall t > 0.$$

By the notation of  $\overline{\Psi}_{\tau,r}$  and  $\underline{\Psi}_{\tau,r}$  in the Notation 6 we get

$$|\overline{\Psi}_{\tau,r}(k,t)| = |\underline{\Psi}_{\tau,r}(k,t)|, \quad \forall t > 1, \quad \forall k \geq 1. \quad (2)$$

Since  $0 \leq f_r(u) \leq u^{\frac{1}{2}-r}$  for every  $u > 0$  then for every  $k \geq 1$  we have

$$\begin{aligned}|\underline{\Psi}_{\tau,r}(k,t)| &= \left| \int_{\frac{1}{t},k}^t u^{i\tau} \int_0^u v^{-i\tau-1} f_r(v) dv du \right| \leq \int_{\frac{1}{t},k}^t \int_0^u v^{-1} f_r(v) dv du \\ &\leq \frac{1}{\frac{1}{2}-r} \int_{\frac{1}{t},k}^t u^{\frac{1}{2}-r} du, \quad \forall k \geq 1, \forall t > 1.\end{aligned} \quad (3)$$

Consider the function  $p$  given in the Notation 4. By Lemma 5, the function  $p$  is 1-periodic, then it is bounded. Use the integration part formula

$$\begin{aligned}&\left| \int_u^{+\infty} v^{-i\tau-1} \left( f_r(v) - \frac{1}{2} v^{-\frac{1}{2}-r} \right) dv \right| \\ &= \left| \int_u^{+\infty} v^{-i\tau-1} \left( \{v\} - \frac{1}{2} \right) v^{-\frac{1}{2}-r} dv \right| \\ &= \left| \left( i\tau + \frac{3}{2} + r \right) \int_u^{+\infty} v^{-i\tau-\frac{5}{2}-r} (p(v) - p(u)) dv \right|.\end{aligned}$$

Since

$$|p(v) - p(u)| < \omega, \quad \forall u > 0, \quad \forall v > 0,$$

where  $\omega$  is given by the Notation 4, we get

$$\begin{aligned} \left| \int_u^{+\infty} v^{-i\tau-1} \left( f_r(v) - \frac{1}{2} v^{-\frac{1}{2}-r} \right) dv \right| &\leq \omega(|\tau| + 2) \int_u^{+\infty} v^{-\frac{5}{2}-r} dv \\ &\leq \omega(|\tau| + 2) u^{-\frac{3}{2}-r}, \quad \forall u > 0. \end{aligned}$$

Then

$$\begin{aligned} \left| \int_{\frac{1}{t},k}^t u^{i\tau} \int_u^{+\infty} v^{-i\tau-1} \left( f_r(v) - \frac{1}{2} v^{-\frac{1}{2}-r} \right) dv du \right| \\ \leq \omega(|\tau| + 2) \int_{\frac{1}{t},k}^t u^{-\frac{3}{2}-r} du, \quad \forall t > 1, \quad \forall k \geq 1. \end{aligned}$$

Since

$$u^{-i\tau} \int_u^{+\infty} v^{-i\tau-\frac{3}{2}-r} dv = \frac{1}{\frac{1}{2} + r - i\tau} u^{-\frac{1}{2}-r}, \quad \forall u > 0,$$

Then

$$|\bar{\Psi}_{\tau,r}(k)| \geq \frac{1}{2} \frac{1}{|\frac{1}{2} + r - i\tau|} \int_{\frac{1}{t},k}^t u^{-\frac{1}{2}-r} du - \omega(|\tau| + 2) \int_{\frac{1}{t},k}^t u^{-\frac{3}{2}-r} du. \quad (4)$$

Since  $r \in (-\frac{1}{2}, 0)$ , then there exists  $t > 1$  such that

$$\lim_{k \rightarrow +\infty} \frac{\int_{\frac{1}{t},k}^t u^{-\frac{1}{2}-r} du}{\int_{\frac{1}{t},k}^t u^{\frac{1}{2}-r} du} = +\infty \quad \text{and} \quad \lim_{k \rightarrow +\infty} \frac{\int_{\frac{1}{t},k}^t u^{-\frac{3}{2}-r} du}{\int_{\frac{1}{t},k}^t u^{\frac{1}{2}-r} du} = 0.$$

The Equations (3) and (4) implies that

$$\lim_{k \rightarrow +\infty} \left| \frac{\bar{\Psi}_{\tau,r}(k,t)}{\underline{\Psi}_{\tau,r}(k,t)} \right| \geq \frac{1}{2} \frac{\frac{1}{2} - r}{|\frac{1}{2} + r - i\tau|} \lim_{k \rightarrow +\infty} \frac{\int_{\frac{1}{t},k}^t u^{-\frac{1}{2}-r} du}{\int_{\frac{1}{t},k}^t u^{\frac{1}{2}-r} du} = +\infty.$$

Contradiction with the Equation (2).  $\square$

## References

- [1] E.C. Titchmarsh, The Theory of the Riemann Zeta-Function (revised by D.R. Heath-Brown), Clarendon Press, Oxford. (1986).