



HAL
open science

Glycerol, a possible new player in the biology of trypanosomes

Frédéric Bringaud, Nicolas Plazolles, Erika Pineda, Corinne Asencio, Oriana Villafraz, Yoann Millerioux, Loïc Rivière, Emmanuel Tetaud

► **To cite this version:**

Frédéric Bringaud, Nicolas Plazolles, Erika Pineda, Corinne Asencio, Oriana Villafraz, et al.. Glycerol, a possible new player in the biology of trypanosomes. PLoS Pathogens, 2021, 17, 10.1371/journal.ppat.1010035 . hal-03851321

HAL Id: hal-03851321

<https://hal.science/hal-03851321>

Submitted on 14 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PEARLS

Glycerol, a possible new player in the biology of trypanosomes

Frédéric Bringaud ^{*}, Nicolas Plazolles , Erika Pineda , Corinne Asencio , Oriana Villafraz , Yoann Millerioux , Loïc Rivière , Emmanuel Tetaud 

Microbiologie Fondamentale et Pathogénicité (MFP), UMR 5234, Bordeaux University, CNRS, Bordeaux, France

* frederic.bringaud@u-bordeaux.fr



Introduction

Trypanosoma brucei is a unicellular eukaryote that causes human African trypanosomiasis, also known as sleeping sickness [1]. Parasite transmission between mammals is ensured by a haematophagous insect vector of the genus *Glossina*, also called the tsetse fly. Trypanosomes adapt to their natural hosts, in particular to the available carbon sources required to fuel central metabolism and to produce ATP. For instance, in the digestive tract of the insect vector, procyclic forms (PCFs) of *T. brucei* use proline abundantly present in the insect [2]. It has recently been proposed that other carbon sources may also be required for the parasite in the fly, such as glucose at the onset of infection or intermediates of the tricarboxylic acid cycle [3,4]. By contrast, the bloodstream forms (BSF) rely on glucose present at homeostatic levels in all mammalian fluids [2]. In addition to proline and glucose, glycerol can also fuel central carbon metabolism of the parasite, but its relevance has received little attention until recently. This Pearl article will highlight recent data on glycerol metabolism and their implications for understanding trypanosome biology.

Procyclic trypanosomes (PCF) prefer glycerol over glucose

As mentioned above, PCF evolve in the glucose-depleted midgut of their insect vector, where they rely on proline [5,6], but, nevertheless, use glucose as the primary carbon source and down-regulate proline consumption up to 7 times under *in vitro* conditions [7,8]. The molecular mechanism of this glucose preference over proline is currently unknown. Surprisingly, the parasite also developed an absolute preference for glycerol over glucose, already described in the 1960s by Ryley [9] and recently revisited by our group [10]. Indeed, glucose is not consumed as long as glycerol is present in the medium. As far as we know, *T. brucei* is the only unicellular eukaryote to date reported to prefer a nonglycolytic carbon source to glucose.

We characterised the molecular mechanism of this glycerol preference, which was called “metabolic contest” since it is based on competition between 2 kinases (hexokinase [HK] and glycerol kinase [GK]) for the same substrate (ATP) [10]. When glucose and glycerol enter the cells via glucose transporters and aquaglyceroporins, they are first phosphorylated to glucose 6-phosphate (G6P) and glycerol 3-phosphate (Gly3P) by HK (step 1 in Fig 1) and GK (step 19), respectively. The phospho group prevents G6P, Gly3P, and the derived phosphorylated metabolites from leaving the cell through transport processes, participates in ATP production, and facilitates enzyme binding and activity. In trypanosomes, HK and GK, as well as most other enzymes involved in glycolysis and glycerol metabolism, are located in peroxisome-related organelles, named glycosomes, which show limited or no nucleotide exchange with the cytosol on a metabolic timescale. Therefore, consumption and production of ATP are tightly

OPEN ACCESS

Citation: Bringaud F, Plazolles N, Pineda E, Asencio C, Villafraz O, Millerioux Y, et al. (2021) Glycerol, a possible new player in the biology of trypanosomes. *PLoS Pathog* 17(12): e1010035. <https://doi.org/10.1371/journal.ppat.1010035>

Editor: Laurie Read, University at Buffalo School of Medicine and Biomedical Sciences, UNITED STATES

Published: December 2, 2021

Copyright: © 2021 Bringaud et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: FB’s team is supported by the Centre National de la Recherche Scientifique (CNRS, <https://www.cnrs.fr/>) (including NP, CA and ET salary), the Université de Bordeaux (<https://www.u-bordeaux.fr/>) (including LR salary), the Agence National de Recherche (ANR, <https://anr.fr/>) through the grants GLYCONOV (grant number ANR-15-CE15-0025-01) (including EP salary) and ADIPOTRYP (grant number ANR19-CE15-0004-01) (including EP salary) and the Laboratoire d’Excellence (<https://www.enseignementsup-recherche.gouv.fr/cid51355/laboratoires-d-excellence.html>) through the LabEx ParaFrap (grant number ANR-11-LABX-0024) (including OV and YM salary) and the “Fondation pour le

Recherche Médicale[®] (FRM, <https://www.frm.org/>) ("Equipe FRM", grant n°EQU201903007845) (including EP salary). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist

balanced within the organelle [11], with each ATP molecule required to supply GK and HK being regenerated by phosphoenolpyruvate carboxykinase (PEPCK, step 11) and pyruvate phosphate dikinase (PPDK, step 15) in PCF glycosomes (Fig 1A and 1B). Hence, the limitation of the glycosomal ATP pool available to glycosomal kinases offers a situation where a significant excess of one kinase (here GK) can theoretically abolish the metabolic flux through another one (here HK). In this context, the very large excess of GK activity compared to HK activity (74-fold excess) explains the absolute glycerol preference confirmed by genetic manipulations and metabolic approaches [10]. The competition of given enzymes for a common substrate is a well-known process used to finely tune metabolic fluxes, in particular at metabolic branch points [12]. However, this is the first example of a complete repression of one enzymatic catalysis (HK) by the large excess of another one (GK), as a mechanism to control nutrient utilisation.

Metabolic contest is a new concept to describe metabolic choices that resembles the well-characterised catabolic repression or carbon catabolite repression observed in prokaryotes, yeasts, and fungi, albeit based on a completely different molecular mechanism. The mechanisms by which carbon catabolite repression is imposed are quite variable. They follow a




Fig 1. Mechanism of metabolic contest between glycosomal HK and GK. Pathways leading to excretion of end products (succinate, acetate, and pyruvate, highlighted in white rectangle) from metabolism of glucose and glycerol are indicated in red and blue, respectively. The production and consumption of ATP within the glycosomes are indicated and maintenance of the glycosomal ATP balance is highlighted by black arrows. The thickness of the arrows reflects metabolic fluxes and metabolic steps detected only at relatively very low activity or not at all are represented by dotted lines. The GK/HK activity ratio for procytic and bloodstream forms is indicated below the corresponding schemes (A–B and C–D, respectively). Enzymes are (those underlined are glycosomal kinases producing or consuming ATP) the following: 1, hexokinase (HK); 2, glucose-6-phosphate isomerase; 3, phosphofructokinase; 4, aldolase; 5, triose-phosphate isomerase; 6, glyceraldehyde-3-phosphate dehydrogenase; 7, phosphoglycerate kinase (PGK); 8, phosphoglycerate mutase; 9, enolase; 10, pyruvate kinase; 11, phosphoenolpyruvate carboxykinase (PEPCK); 12, malate dehydrogenase; 13, fumarase; 14, NADH-dependent fumarate reductase; 15, pyruvate phosphate dikinase (PPDK); 16, pyruvate dehydrogenase complex; 17, acetate: succinate CoA-transferase; 18, acetyl-CoA thioesterase; 19, glycerol kinase (GK); 20, glycerol-3-phosphate dehydrogenase; 21, fructose-1,6-bisphosphatase. Abbreviations are: BSF, bloodstream forms; DHAP, dihydroxyacetone phosphate; G3P, glyceraldehyde 3-phosphate; G6P, glucose 6-phosphate; Gly3P, glycerol 3-phosphate; PCF, procytic form; PEP, phosphoenolpyruvate.

<https://doi.org/10.1371/journal.ppat.1010035.g001>

general rule, however, with complex sensory systems relying mostly on protein kinases and phosphatases [13], which include the sugar uptake phosphotransferase system characterised in the 1960s [14]. The advantage of the metabolic contest *versus* carbon catabolite repression mechanism is mainly an immediate switch to the less preferred carbon source when the preferred one is exhausted. Theoretically, a metabolic contest could be found in any organism provided that 3 conditions are met, *i.e.*, (i) the sequestration of both metabolic pathways in the same subcellular compartment; (ii) the competition for the same substrate between 2 enzymes, one in each pathway; and (iii) an unbalanced activity between the competing enzymes.

The physiological role of glycerol metabolism *in vivo* in the fly is still an open question. The preference of glycerol over glucose is based on the huge overexpression of GK, which could be reduced by at least 90% without affecting the glycerol metabolic flux [10]. Incidentally, glycerol also induces a 2.3-fold reduction of GK expression, without affecting metabolic contest, suggesting that the very high GK level in glucose-rich conditions is not compatible with glycerol-rich conditions, may be due to accumulation of Gly3P or other downstream metabolites, and/or a sharp decrease in the glycosomal ATP/ADP ratio [10]. It is also noteworthy that, in *T. brucei*, GK is encoded by 5 duplicated *GK* genes, while the genome of almost all other trypanosomatids contains a single *GK* gene, suggesting that *GK* gene duplication was positively selected in *T. brucei* to allow the preference for glycerol. Although, one cannot exclude that the high GK level has other reasons (see next section).

Glycerol has also been shown *in vitro* to prevent differentiation of early PCF, which expressed the surface protein GPEET, to late PCF, which are GPEET negative [15]. GPEET is expressed during midgut infection, and its down-regulation correlates with trypanosomes crossing the peritrophic matrix and colonising the ectoperitrophic space [16], suggesting that glycerol may be present in the digestive tract of the insect to prevent down-regulation of GPEET expression. Unfortunately, with the exception of amino acids [5], the metabolite content in the midgut and other organs of the tsetse has not been studied so far. Thus, further investigation should be done to determine whether glycerol plays a role in the biology of trypanosomes in the insect vector.

Bloodstream trypanosomes (BSF) also metabolise glycerol efficiently

The topology of the central carbon metabolic network is similar between PCF and BSF, with one main exception, *i.e.*, the cytosolic and glycosomal localisation of phosphoglycerate kinase (PGK, step 7), respectively [17]. Another major difference resides in the 10-fold higher glycolytic flux in BSF (for example, HK activity is up-regulated 28-fold [10]), with pyruvate representing up to 95% of the end products excreted from metabolism of glucose [18] (Fig 1C), while PCF mainly convert glucose to excreted acetate and succinate [19] (Fig 1A). Consequently, PGK is responsible for the regeneration of glycosomal ATP in BSF (Fig 1C), rather than PEPCK/PPDK as mentioned above for PCF (Fig 1A). It is noteworthy that the high glycolytic flux described in *T. brucei* BSF, as well as in plant-infecting *Phytomonas* spp., remains the exception in the trypanosomatid world as most of the trypanosomatid parasitic forms developed a “procyclic-like” form of glycolysis, *i.e.*, relative low glycolytic flux and pyruvate being further metabolised in the glycosomes and the mitochondrion [2,20].

Because of the intronless and polycistronic expression nature of trypanosome genes, differential expression of glycolytic enzymes should be controlled posttranscriptionally [21]. No RNA-binding protein directly involved in this process has been identified so far; however, knockout of a major type I protein arginine methyltransferase (PMRT1) induced changes that resemble the metabolic remodelling that occurs during *T. brucei* life cycle progression to PCF, including down-regulation of HK expression [22].

Besides glucose, glycerol utilisation as substrate for ATP production was first proposed decades ago [23], but the possibility of prolonged growth with this substrate has not been addressed until recently. In fact, glycerol was considered a poison, because of its high toxicity for BSF grown in anaerobiosis or in combination with drugs targeting the alternative oxidase [24], since glycerol prevents ATP production by reversal of the GK that must occur under these conditions to maintain the glycosomal ATP/ADP balance. However, in 2018, 2 groups independently reported that glycerol can support growth of BSF *in vitro*, so far believed to be exclusively dependent on glucose for growth [25,26]. This resonates with the recent discovery that the parasite colonises and propagates in the skin and adipose tissues of its mammalian hosts, where adipocytes produce significant amounts of glycerol [27–29], suggesting that glycerol metabolism may play a role for the *in vivo* development of BSF. It is to be noted that the low GK/HK activity ratio (2.2) provides a rational explanation for the absence of metabolic contest for ATP between GK and HK, resulting in concomitant consumption of glucose and glycerol when both are present (Fig 1D). Indeed, although GK is constitutively expressed in *T. brucei* grown under glucose-rich conditions, HK activity is 28-fold higher in BSF than in PCF, as mentioned above [10].

The glycerol-dependent reduction of GK expression mentioned above for PCF also occurs in BSF [25]. Considering the absence of glycerol preference in this parasitic form, this observation suggests that the reason for the large excess of GK level may be related to glycolysis. Indeed, high GK activity may be required for production of glycerol from glycolysis, for instance, when oxygen is limiting, since GK's specific activity for glycerol production is much lower than that for glycerol phosphorylation [30]. However, it cannot be excluded that extravascular trypanosomes require high GK activity to adapt to glycerol-rich tissue.

The relatively low abundance of glycerol in the bloodstream (50 to 100 μ M) [31,32] compared to glucose (5 mM) and the slight preference of BSF for glucose over glycerol [25], imply that glucose is indeed the main source of ATP for BSF in mammalian fluids. However, in the interstitial fluid of mammalian tissues, the extravascular trypanosomes meet a glucose-rich environment (in the range of 3 mM) containing from 0.2 to 3 mM glycerol, depending on the report and the tissues analysed [31,32], suggesting that trypanosomes may adapt and benefit from this potentially glycerol-rich environment. In this context, the resulting *in vivo* glycerol gradient between the intra- and extravascular compartments could influence the parasite tropism to particular tissues *via* specific sensing pathways, such as the social motility phenomenon described in PCF in the insect midgut [33]. The ongoing analyses of these recently discovered extravascular trypanosomes will certainly reveal fascinating new features, especially on carbon source utilisation. Although the exact role of glycerol metabolism in BSF *in vivo* is not understood yet, these data open novel avenues for developing new diagnostic tools and/or treatments based on unexplored molecular targets.

References

1. Buscher P, Cecchi G, Jamonneau V, Priotto G. Human African trypanosomiasis. *Lancet*. 2017; 390:2397–409. [https://doi.org/10.1016/S0140-6736\(17\)31510-6](https://doi.org/10.1016/S0140-6736(17)31510-6) PMID: 28673422
2. Bringaud F, Riviere L, Coustou V. Energy metabolism of trypanosomatids: adaptation to available carbon sources. *Mol Biochem Parasitol*. 2006; 149:1–9. <https://doi.org/10.1016/j.molbiopara.2006.03.017> PMID: 16682088
3. Villafraz O, Biran M, Pineda E, Plazolles N, Cahoreau E, Ornitz Oliveira Souza R, et al. Procytic trypanosomes recycle glucose catabolites and TCA cycle intermediates to stimulate growth in the presence of physiological amounts of proline. *PLoS Pathog*. 2021; 17:e1009204. <https://doi.org/10.1371/journal.ppat.1009204> PMID: 33647053
4. Naguleswaran A, Fernandes P, Bevk S, Rehmann R, Nicholson P, Roditi I. Developmental changes and metabolic reprogramming during establishment of infection and progression of *Trypanosoma*

- brucei brucei* through its insect host. PLoS Negl Trop Dis. 2021; 15:e0009504. <https://doi.org/10.1371/journal.pntd.0009504> PMID: 34543277
5. Balogun RA. Studies on the amino acids of the tsetse fly, *Glossina morsitans*, maintained on *in vitro* and *in vivo* feeding systems. Comp Biochem Physiol Comp Physiol. 1974; 49:215–22. [https://doi.org/10.1016/0300-9629\(74\)90110-8](https://doi.org/10.1016/0300-9629(74)90110-8) PMID: 4153811
 6. Mantilla BS, Marchese L, Casas-Sanchez A, Dyer NA, Ejeh N, Biran M, et al. Proline metabolism is essential for *Trypanosoma brucei brucei* survival in the tsetse vector. PLoS Pathog. 2017; 13:e1006158. <https://doi.org/10.1371/journal.ppat.1006158> PMID: 28114403
 7. Lamour N, Riviere L, Coustou V, Coombs GH, Barrett MP, Bringaud F. Proline metabolism in procyclic *Trypanosoma brucei* is down-regulated in the presence of glucose. J Biol Chem. 2005; 280:11902–10. <https://doi.org/10.1074/jbc.M414274200> PMID: 15665328
 8. Coustou V, Biran M, Breton M, Guegan F, Riviere L, Plazolles N, et al. Glucose-induced remodeling of intermediary and energy metabolism in procyclic *Trypanosoma brucei*. J Biol Chem. 2008; 283:16342–54. <https://doi.org/10.1074/jbc.M709592200> PMID: 18430732
 9. Ryley JF. Studies on the metabolism of protozoa. 9. Comparative metabolism of bloodstream and culture forms of *Trypanosoma rhodesiense*. Biochem J. 1962; 85:211–23. <https://doi.org/10.1042/bj0850211> PMID: 13983265
 10. Allmann S, Wagnies M, Plazolles N, Cahoreau E, Biran M, Morand P, et al. Glycerol suppresses glucose consumption in trypanosomes through metabolic contest. PLoS Biol. 2021; 19:e3001359. <https://doi.org/10.1371/journal.pbio.3001359> PMID: 34388147
 11. Haanstra JR, Bakker BM, Michels PA. In or out? On the tightness of glycosomal compartmentalization of metabolites and enzymes in *Trypanosoma brucei*. Mol Biochem Parasitol. 2014; 198:18–28. <https://doi.org/10.1016/j.molbiopara.2014.11.004> PMID: 25476771
 12. Siebert G, Pfaender P, Kesselring K. Competition of several enzymes for a common substrate: a possible model of cellular events. Adv Enzyme Regul. 1969; 7:131–48. [https://doi.org/10.1016/0065-2571\(69\)90015-6](https://doi.org/10.1016/0065-2571(69)90015-6) PMID: 5367801
 13. Galinier A, Deutscher J. Sophisticated Regulation of Transcriptional Factors by the Bacterial Phosphoenolpyruvate: Sugar Phosphotransferase System. J Mol Biol. 2017; 429:773–89. <https://doi.org/10.1016/j.jmb.2017.02.006> PMID: 28202392
 14. Kundig W, Ghosh S, Roseman S. Phosphate Bound to Histidine in a Protein as an Intermediate in a Novel Phospho-Transferase System. Proc Natl Acad Sci U S A. 1964; 52:1067–74. <https://doi.org/10.1073/pnas.52.4.1067> PMID: 14224387
 15. Vassella E, Den Abbeele JV, Butikofer P, Renggli CK, Furger A, Brun R, et al. A major surface glycoprotein of *Trypanosoma brucei* is expressed transiently during development and can be regulated post-transcriptionally by glycerol or hypoxia. Genes Dev. 2000; 14:615–26. <https://doi.org/10.1101/gad.14.5.615> PMID: 10716949
 16. Van Den Abbeele J, Claes Y, Van Bockstaele D, Le Ray D, Coosemans M. *Trypanosoma brucei* spp. development in the tsetse fly: characterization of the post-mesocyclic stages in the foregut and proboscis. Parasitology. 1999; 118:469–78. <https://doi.org/10.1017/s0031182099004217> PMID: 10363280
 17. Osinga KA, Swinkels BW, Gibson WC, Borst P, Veeneman GH, Van Boom JH, et al. Topogenesis of microbody enzymes: a sequence comparison of the genes for the glycosomal (microbody) and cytosolic phosphoglycerate kinases of *Trypanosoma brucei*. EMBO J. 1985; 4:3811–7. <https://doi.org/10.1002/j.1460-2075.1985.tb04152.x> PMID: 3004970
 18. Mazet M, Morand P, Biran M, Bouyssou G, Courtois P, Daulouede S, et al. Revisiting the central metabolism of the bloodstream forms of *Trypanosoma brucei*: production of acetate in the mitochondrion is essential for parasite viability. PLoS Negl Trop Dis. 2013; 7:e2587. <https://doi.org/10.1371/journal.pntd.0002587> PMID: 24367711
 19. Bringaud F, Biran M, Millerioux Y, Wagnies M, Allmann S, Mazet M. Combining reverse genetics and NMR-based metabolomics unravels trypanosome-specific metabolic pathways. Mol Microbiol. 2015; 96:917–26. <https://doi.org/10.1111/mmi.12990> PMID: 25753950
 20. Sanchez-Moreno M, Laszity D, Coppens I, Opperdoes FR. Characterization of carbohydrate metabolism and demonstration of glycosomes in a *Phytomonas* sp. isolated from *Euphorbia characias*. Mol Biochem Parasitol. 1992; 54:185–99. [https://doi.org/10.1016/0166-6851\(92\)90111-v](https://doi.org/10.1016/0166-6851(92)90111-v) PMID: 1435859
 21. Clayton CE. Gene expression in Kinetoplastids. Curr Opin Microbiol. 2016; 32:46–51. <https://doi.org/10.1016/j.mib.2016.04.018> PMID: 27177350
 22. Kafková L, Tu C, Pazzo KL, Smith KP, Debler EW, Paul KS, et al. *Trypanosoma brucei* PRMT1 Is a Nucleic Acid Binding Protein with a Role in Energy Metabolism and the Starvation Stress Response. mBio. 2018;9. <https://doi.org/10.1128/mBio.02430-18> PMID: 30563898

23. Ryley JF. Studies on the metabolism of the Protozoa. 7. Comparative carbohydrate metabolism of eleven species of trypanosome. *Biochem J.* 1956; 62:215–22. <https://doi.org/10.1042/bj0620215> PMID: [13293175](https://pubmed.ncbi.nlm.nih.gov/13293175/)
24. Clarkson AB, Brohn FH. Trypanosomiasis: an approach to chemotherapy by the inhibition of carbohydrate catabolism. *Science.* 1976; 194:204–6. <https://doi.org/10.1126/science.986688> PMID: [986688](https://pubmed.ncbi.nlm.nih.gov/986688/)
25. Pineda E, Thonnus M, Mazet M, Mourier A, Cahoreau E, Kulyk H, et al. Glycerol supports growth of the *Trypanosoma brucei* bloodstream forms in the absence of glucose: Analysis of metabolic adaptations on glycerol-rich conditions. *PLoS Pathog.* 2018; 14:e1007412. <https://doi.org/10.1371/journal.ppat.1007412> PMID: [30383867](https://pubmed.ncbi.nlm.nih.gov/30383867/)
26. Kovarova J, Nagar R, Faria J, Ferguson MAJ, Barrett MP, Horn D. Gluconeogenesis using glycerol as a substrate in bloodstream-form *Trypanosoma brucei*. *PLoS Pathog.* 2018; 14:e1007475. <https://doi.org/10.1371/journal.ppat.1007475> PMID: [30589893](https://pubmed.ncbi.nlm.nih.gov/30589893/)
27. Capewell P, Cren-Travaille C, Marchesi F, Johnston P, Clucas C, Benson RA, et al. The skin is a significant but overlooked anatomical reservoir for vector-borne African trypanosomes. *Elife.* 2016;5. <https://doi.org/10.7554/eLife.17716> PMID: [27653219](https://pubmed.ncbi.nlm.nih.gov/27653219/)
28. Caljon G, Van Reet N, De Trez C, Vermeersch M, Perez-Morga D, Van Den Abbeele J. The dermis as a delivery site of *Trypanosoma brucei* for tsetse flies. *PLoS Pathog.* 2016; 12:e1005744. <https://doi.org/10.1371/journal.ppat.1005744> PMID: [27441553](https://pubmed.ncbi.nlm.nih.gov/27441553/)
29. Trindade S, Rijo-Ferreira F, Carvalho T, Pinto-Neves D, Guegan F, Aresta-Branco F, et al. *Trypanosoma brucei* parasites occupy and functionally adapt to the adipose tissue in mice. *Cell Host Microbe.* 2016; 19:837–48. <https://doi.org/10.1016/j.chom.2016.05.002> PMID: [27237364](https://pubmed.ncbi.nlm.nih.gov/27237364/)
30. Krakow JL, Wang CC. Purification and characterization of glycerol kinase from *Trypanosoma brucei*. *Mol Biochem Parasitol.* 1990; 43:17–25. [https://doi.org/10.1016/0166-6851\(90\)90126-7](https://doi.org/10.1016/0166-6851(90)90126-7) PMID: [2290444](https://pubmed.ncbi.nlm.nih.gov/2290444/)
31. Maggs DG, Jacob R, Rife F, Lange R, Leone P, During MJ, et al. Interstitial fluid concentrations of glycerol, glucose, and amino acids in human quadriceps muscle and adipose tissue. Evidence for significant lipolysis in skeletal muscle. *J Clin Invest.* 1995; 96:370–7. <https://doi.org/10.1172/JCI118043> PMID: [7615807](https://pubmed.ncbi.nlm.nih.gov/7615807/)
32. Vestergaard ET, Moller N, Jorgensen JO. Acute peripheral tissue effects of ghrelin on interstitial levels of glucose, glycerol, and lactate: a microdialysis study in healthy human subjects. *Am J Physiol Endocrinol Metab.* 2013; 304:E1273–80. <https://doi.org/10.1152/ajpendo.00662.2012> PMID: [23592479](https://pubmed.ncbi.nlm.nih.gov/23592479/)
33. Saada EA, DeMarco SF, Shimogawa MM, Hill KL. “With a little help from my friends”—social motility in *Trypanosoma brucei*. *PLoS Pathog.* 2015; 11:e1005272. <https://doi.org/10.1371/journal.ppat.1005272> PMID: [26679190](https://pubmed.ncbi.nlm.nih.gov/26679190/)