



**HAL**  
open science

## **Pesticide responses of Arctic and temperate microalgae differ in relation to ecophysiological characteristics**

Juan Du, Disney Izquierdo, Jonathan Naoum, Leanne Ohlund, Lekha Sleno, Beatrix Beisner, Johann Lavaud, Philippe Juneau

### ► **To cite this version:**

Juan Du, Disney Izquierdo, Jonathan Naoum, Leanne Ohlund, Lekha Sleno, et al.. Pesticide responses of Arctic and temperate microalgae differ in relation to ecophysiological characteristics. *Aquatic Toxicology*, 2022, pp.106323. 10.1016/j.aquatox.2022.106323 . hal-03851537

**HAL Id: hal-03851537**

**<https://hal.science/hal-03851537>**

Submitted on 14 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Pesticide responses of Arctic and temperate microalgae differ in relation to  
2 ecophysiological characteristics

3  
4 Juan Du<sup>1</sup>, Disney Izquierdo<sup>2</sup>, Jonathan Naoum<sup>2</sup>, Leanne Ohlund<sup>3</sup>, Lekha Sleno<sup>3</sup>,  
5 Beatrix E. Beisner<sup>4</sup>, Johann Lavaud<sup>5,6</sup> and Philippe Juneau<sup>2\*</sup>

6 <sup>1</sup> Department of Biological Sciences, Université du Québec à Montréal, GRIL-TOXEN,  
7 Succ Centre-Ville, Montreal, Quebec, H3C 3P8, Canada<sup>2</sup> Department of Biological  
8 Sciences, Université du Québec à Montréal, GRIL-EcotoQ-TOXEN, Succ Centre-  
9 Ville, Montreal, Quebec, H3C 3P8, Canada<sup>3</sup> Chemistry Department, Université du  
10 Québec à Montréal, EcotoQ-TOXEN, Succ Centre-Ville, Montreal, Quebec, H3C  
11 3P8, Canada

12 <sup>4</sup> Department of Biological Sciences, Groupe de recherche interuniversitaire en  
13 limnologie (GRIL), Université du Québec à Montréal, Succ Centre-Ville, Montreal,  
14 Quebec, H3C 3P8, Canada

15 <sup>5</sup> TAKUVIK International Research Laboratory IRL3376, Université Laval (Canada) -  
16 CNRS (France), Pavillon Alexandre-Vachon, 1045 av. de la Médecine, local 2064,  
17 G1V 0A6 Québec, Canada

18 <sup>6</sup> LEMAR-Laboratory of Environmental Marine Sciences, UMR6539, CNRS/Univ  
19 Brest/Ifremer/IRD, Institut Universitaire Européen de la Mer, Technopôle Brest-  
20 Iroise, rue Dumont d'Urville, 29280 Plouzané, France.

21 **\*Corresponding author:**

22 E-mail addresses: [juneau.philippe@uqam.ca](mailto:juneau.philippe@uqam.ca) (P. Juneau)

23 Full postal address:

24 Department of Biological Sciences

25 Université du Québec à Montréal

26 CP8888, Succ Centre-Ville

27 Montréal (QC)

28 H3C 3P8

29 Canada

30

31

## 32 **Highlights**

- 33 ● Arctic microalgae have distinct ecophysiological characteristics relative to their  
34 temperate counterparts.
- 35 ● Unexpectedly, one Arctic microalgae was more tolerant to pesticides than its  
36 temperate counterparts, which disagrees that the greater ecological risk of  
37 pollutants in polar ecosystems is always higher than in temperate regions.
- 38 ● Sensitivity variation to the four pesticides appears to be mainly due to differences  
39 in the protective mechanisms between Arctic and temperate microalgae.

## 40 **Abstract**

41 Polar ecosystems play an important role in global primary production. Microalgae have  
42 adaptations that enable them to live under low temperature environments where  
43 irradiance and day length change drastically. Their adaptations, leading to different  
44 ecophysiological characteristics relative to temperate species, could also alter their  
45 sensitivity to pollutants such as pesticides. This study's objective was to understand  
46 how different ecophysiological characteristics influence the response of Arctic  
47 phytoplankton to pesticides in relation to the responses of their temperate counterparts.  
48 Ecophysiological endpoints were related to growth, cell biovolume, pigment content,  
49 photosynthetic activity, photoprotective mechanisms (NPQ, antioxidant enzyme  
50 activities), and reactive oxygen species (ROS) content. The Arctic species *Micromonas*  
51 *polaris* was more resistant to atrazine and simazine than its temperate counterpart  
52 *Micromonas bravo*. However, the other Arctic species *Chaetoceros neogracilis* was  
53 more sensitive to these herbicides than its temperate counterpart *Chaetoceros*  
54 *neogracile*. With respect to two other pesticide toxicity, both temperate microalgae were  
55 more sensitive to trifluralin, while Arctic microalgae were more sensitive to  
56 chlorpyrifos (insecticide). All differences could be ascribed to differences in the eco-  
57 physiological features of the two microalgal groups, which can be explained by cell  
58 size, pigment content, ROS content and protective mechanisms (NPQ and antioxidant  
59 enzymes).

60 **Keywords:** microalgae, *Micromonas*, *Chaetoceros*, ecophysiology, photosynthesis,  
61 pesticides, toxicity mechanism

62

63 Abbreviations:

64 T-CN (Temperate *Chaetoceros neogracile*), T-MB (temperate *Micromonas bravo*) and  
65 A-MP (Arctic *Micromonas polaris*), Arctic *Chaetoceros neogracilis* (A-CN), Car,  
66 carotenoids; Chl *a*, chlorophyll *a*; PSI, photosystem I; PSII, photosystem II; NPQ, non-  
67 photochemical quenching; Q<sub>A</sub>, primary electron acceptor of PSII; Q<sub>B</sub>, secondary  
68 electron acceptor of PSII; RC, reaction center, Φ<sub>M</sub>, PSII maximum quantum yield; Φ'<sub>M</sub>,  
69 PSII operational quantum yield; NPQ<sub>max</sub>, Maximum ability for dissipation of excess  
70 energy; E<sub>k</sub>, Light saturation coefficient; α, Maximum light efficiency use ROS, reactive  
71 oxygen species; SOD, superoxide dismutase; CAT, catalase; logK<sub>ow</sub>, a ratio between  
72 the concentration of a chemical in the octanol phase to its concentration in the water  
73 phase.

## 74 **1. Introduction**

75 Arctic habitats are subjected to harsh environmental conditions. Nevertheless, they  
76 provide a major contribution to global primary production and the Arctic Ocean net  
77 primary production has increased recently (Ardyna and Arrigo 2020). More than 75%  
78 of the Arctic phytoplankton biomass is composed of diatoms and small flagellate  
79 prasinophytes (*Micromonas* sp.) (Balzano et al. 2012, Lovejoy et al. 2007), and they  
80 thus play an essential role in the Arctic food web (Frey et al. 2018). Microalgae in polar  
81 regions have adaptations that enable them to grow well in these regions where  
82 temperatures are permanently low, but irradiance and day length are extremely variable  
83 (Handler 2017). Microalgae are impacted by cold environments in many ways,  
84 including lower enzyme activity (Wiebe et al. 1992), altered membrane fluidity (White  
85 et al. 2000), nutrient availability, balancing the usage and absorption of energy (Parker  
86 and Armbrust 2005), and the capacity to grow (Margesin 2007). There are differences  
87 in taxonomy, genetics, and ecology between Arctic microalgae and their temperate  
88 counterparts, while photo-physiology is not well documented (Lacour et al. 2017).  
89 Although there have been great developments in the taxonomy, genetics, and ecology  
90 of polar phytoplankton, their photo-physiological properties are not yet well  
91 documented, and the understanding of aquatic contaminant effects on their  
92 ecophysiology is lacking (Lyon and Mock 2014).

93 Application of pesticides leads to a substantial loss from agricultural fields leading  
94 to their detection in the aquatic environment, due to leaching, runoff and spray-drift  
95 (Larsbo et al. 2016, Zhang et al. 2018). Some authors have reported that Arctic waters  
96 are contaminated with pesticides applied in southern regions due to the long-distance  
97 aerial and marine transport of chemicals (Cabrerizo et al. 2019, Ma et al. 2018, Muir et  
98 al. 2013, Muir and de Wit 2010). Moreover, owing to the accumulation of pesticides in  
99 Arctic ice cover and snow over the years, pesticide concentrations in Arctic waters  
100 should increase over time, as accumulated ice is melting at an unprecedented rate due  
101 to global warming (Pućko et al. 2017). Chlorpyrifos, diazinon, trifluralin, endosulfan  
102 and lindane are some of the typical pesticides found in surface and groundwater near

103 agricultural lands across Canada and USA (Vorkamp and Riget 2014), as well as in  
104 Arctic waters (Hoferkamp et al. 2010, Vorkamp and Riget 2014, Weber et al. 2010).

105       Among the main classes of pesticides, herbicides are the most widely used (Balmer  
106 et al. 2019). Numerous herbicides have detrimental effects on photosynthesis due to  
107 cellular oxidative damage induced by the accumulation of reactive oxygen species  
108 (ROS). This accumulation promotes lipid peroxidation, which results in the destruction  
109 of membranes, such as the photosynthetic ones (Chalifour et al. 2014, DeLorenzo 2001).  
110 Insecticides, although not designed to affect plants, have been demonstrated to have  
111 toxic effects on the growth, photosynthesis, biovolume, pigment and lipid contents of  
112 phytoplankton (Asselborn et al. 2015, Yadav 2015). It is well known that the sensitivity  
113 to pesticides of temperate algal species varies considerably, and several factors may  
114 contribute to this species-specificity. For example, damage to photosynthetic apparatus  
115 caused by pesticides can be minimized by various photoprotective mechanisms,  
116 including antioxidant system designed to eliminate the excess ROS (Medithi et al. 2021)  
117 and non-photochemical quenching (NPQ) energy dissipation processes related to the  
118 capacity to modulate light absorption and dissipate excess energy as heat (Moustakas  
119 et al. 2022). Furthermore, cells can also adjust pesticide uptake by modifying their  
120 surface to biovolume ratio (Larras et al. 2013, Tang et al. 1998).

121       Most of our understanding about the physiological characteristics and pesticide  
122 effects on microalgae is from temperate phytoplankton species. Indeed, the  
123 physiological features and potential impacts of pesticides on Arctic phytoplankton are  
124 very scarce (Kottuparambil et al. 2017). However, some authors have predicted that  
125 Arctic ecosystems and their organisms are likely more sensitive to contaminants than  
126 those at temperate latitudes (CARC 1990, Kottuparambil et al. 2017). In that case,  
127 ecophysiological properties of Arctic microalgae evolved under the extreme conditions  
128 found in the Arctic Ocean and this result in adaptations that may permit these algae to  
129 cope with pesticide stress. Although there have been many reports on the impacts of  
130 pesticides on the physiology and growth of temperate phytoplankton (Singh et al. 2016,  
131 Vonk and Kraak 2020), very little is known about their potential impacts on microalgae  
132 having extreme low temperatures and rapid environmental change due to global

133 warming such as for Arctic phytoplankton. Moreover, the comparison of the tolerance  
134 mechanisms and ecophysiological characteristics between the phytoplankton of Arctic  
135 with temperate regions is lacking. This knowledge will permit to gain insights for future  
136 development of algal bioassays for Arctic regions since using temperate species could  
137 not be appropriate. We thus compared pesticide responses in Arctic microalgae and their  
138 temperate counterparts to four pesticides (atrazine, simazine, trifluralin and  
139 chlorpyrifos). Arctic microalgae grow under the extreme conditions of the Arctic Ocean,  
140 and should have different physiological characteristics from temperate microalgae.  
141 Therefore, we examined which ecophysiological adaptation might benefit to Arctic  
142 organisms by exploring differences in their sensitivities and response mechanisms.

143

## 144 **2. Materials and methods**

### 145 2.1 Microalgal species and growth conditions

146 We compared the responses in two temperate species: *Chaetoceros neogracile* (T-  
147 CN; CCMP1425), *Micromonas bravo* (T-MB; CCMP1646), and two Arctic species  
148 strains *Micromonas polaris* (A-MP; CCMP2099), *Chaetoceros neogracilis* (A-CN;  
149 RCC2279). The first three species were purchased from National Contract Management  
150 Association (NCMA), while the latter Arctic strain was obtained from the Roscoff  
151 culture collections. All species were cultivated in marine L1 medium (Guillard et al.,  
152 1993) with a total volume of 100 mL medium in species-specific 250 mL Erlenmeyer  
153 flasks. The cultures were grown at  $100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  under a 14:10 h light: dark  
154 illumination cycle with daily gentle shaking. Temperate and Arctic species were grown  
155 at 18 °C and 4 °C respectively. Algal cells were periodically transferred (for at least  
156 eight generations) into fresh medium to maintain their exponential growth phase. The  
157 cell concentrations were measured with a Multisizer 3 Coulter Counter particle analyzer  
158 (Beckman Coulter Inc., USA). The growth rate ( $\mu$ ) was determined as follow:  $\mu =$   
159  $(\ln N_n) - (\ln N_0) / T$ , where  $\mu =$  Average specific growth rate,  $N_0$ ,  $N_n$  indicate cell density  
160 (cells/mL) at the beginning of test and at the end of the treatment (3 days), T expresses

161 the exposure time (3 days).

## 162 2.2 Pesticide preparation and treatment

163 All pesticides (Table 1) used in the present study were obtained from Sigma-  
164 Aldrich (PESTANAL®, Canada). Pesticides (stock solutions in acetone) were added to  
165 the growth media for the exposure experiments at final acetone concentration never  
166 exceeding 0.01%, a concentration of acetone not inducing any measurable effect on the  
167 parameters assayed. From the original eight pesticides, the impacts of four of them were  
168 further investigated (atrazine, simazine, trifluralin and chlorpyrifos), since the other  
169 four pesticides showed no (clopyralid-1000 µg/L, metolachlor-3.5 µM/L and lindane-  
170 1000 µg/L) or very little (< 10%) toxicity (endosulfan-1000 µg/L) on the photosynthetic  
171 activity at very high concentrations for the studied microalgae (data not shown).  
172 Concentrations of atrazine measured in the medium at the beginning of the experiment  
173 were 0 µg/L, 6.44 µg/L, 25.93 µg/L, 50.68 µg/L, 99.00 µg/L and 245.14 µg/L. For  
174 simazine the measured concentrations were 0 µg/L, 5.55 µg/L, 25.23 µg/L, 50.57 µg/L,  
175 101.61 µg/L and 248.62 µg/L. We indicated these atrazine and simazine concentrations  
176 in the figures and tables as 0 µg/L, 5 µg/L, 25 µg/L, 50 µg/L, 100 µg/L and 250 µg/L.  
177 Measured trifluralin concentrations tested were 0 µg/L, 198.47 µg/L and 500.93 µg/L  
178 and for chlorpyrifos 0 µg/L, 189.13 µg/L and 475.56 µg/L (noted in the figures and  
179 tables as 0 µg/L, 200 µg/L and 500 µg/L). No significant differences in the measured  
180 concentrations were observed between the two experimental temperatures for the  
181 different pesticides. Concentrations of pesticides were evaluated using a QTRAP 5500  
182 mass spectrometer (Sciex, Concord, ON, Canada) according to Chalifour et al. (2016)  
183 and Takishita et al. (2021). The four studied pesticides (atrazine, simazine, trifluralin  
184 and chlpyrifos) are known to have relatively long half-lives in water, ranging from  
185 months to years (Bai et al. 2015, Vonk and Kraak 2020), and therefore are stable and  
186 do not degrade during our experiments (Du et al. 2022, in preparation).

187 Exponentially growing microalgae were transferred into 1L Erlenmeyer flasks at  
188 a cell density of  $2.5 \times 10^5$  (*Chaetoceros*) and  $2.5 \times 10^6$  (*Micromonas*) cells mL<sup>-1</sup>  
189 respectively, and then exposed to different concentrations of pesticides for 72 h. All



190 treatments were done in triplicate. Cell densities and cell biovolumes were assessed at  
191 the beginning and the end of the experiment with a particle counter (Multisizer 3  
192 Coulter Counter, Beckman Coulter Inc., USA).

193 Table 1. The chemical families and mode of action for pesticides used in this study  
194 (adapted from [www.irac-online.org](http://www.irac-online.org)).

Class	Substance	Chemical family	Mode of action
Herbicide	Atrazine	Triazine	inhibition of photosynthesis at photosystem II
	Simazine	Triazine	inhibition of photosynthesis at photosystem II
	Trifluralin	Dinitroaniline	inhibition of cell mitosis
	Clopyralid	Pyridinecarboxylic acid	inhibit cell division and growth
	Metolachlor	Chloroacetanilide	inhibition biosynthesis of chlorophyll, proteins, fatty acids and lipids
Insecticide	Chlorpyrifos	Organophosphate	inhibition Nervous System (acetylcholine esterase (AChE))
	Lindane	Organochlorine	inhibition Nervous System (GABA receptor)
	Endosulfan	Organochlorine	inhibition Nervous System (GABA receptor)

195

### 196 2.3 Pigment measurements

197 Algal cultures (25 mL) were harvested, under dim green light, 72 h after the  
198 beginning of the treatments by gentle filtration on 0.8µm filter membrane  
199 (Polytetrafluoroethylene; Xingya Purifying Materials Factory; Shanghai, China), and  
200 placed in 2 mL Eppendorf tubes covered with aluminum foil, then rapidly immersed  
201 into liquid nitrogen and kept at -80 °C until analysis. Extractions of the pigments were  
202 done by adding 2 mL of acetone 90% overnight at -20 °C prior to analysis. Ultrasonic  
203 probe was used to break the cells (3 W/cm<sup>2</sup> for 20 s; Sonic dismembrator Model 100,  
204 Fisher Scientific). The extracts were centrifuged at 4 °C for 10 min (10000×g) and the  
205 supernatant was kept for quantification of chlorophyll (Chl *a*) and carotenoid (Car).  
206 Using Cary 300 UV spectrophotometer (Varian, USA) each extract was scanned  
207 between 400–750nm. Independent triplicates were sampled for each culture. The  
208 contents of Chl *a* and carotenoids were calculated according to Jeffrey and Humphrey  
209 (1975) and Seely et al. (1972) respectively.

### 210 2.4 Fluorescence measurements

211 The photosynthetic light curves were obtained using a PAM fluorometer (Water-  
212 PAM, Walz, Germany) according to Du et al. (2019), with saturation pulses (3000 µmol

213 photons  $\text{m}^{-2} \text{s}^{-1}$ , 800ms) and 8 levels of actinic light intensities (0, 46, 105, 188, 276,  
214 427, 635, 906, and 1207  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ). The samples (3mL) were dark  
215 acclimated for 20 minutes before measurements and all samples were measured at their  
216 incubation temperature (4 °C and 18 °C). The maximum ( $\Phi_M$ ) and operational ( $\Phi'_M$ )  
217 PSII quantum yields, and the non-photochemical quenching (NPQ) were determined  
218 from this light curve when the actinic light was 0  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for  $\Phi_M$  and 105  
219  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  for  $\Phi'_M$  and NPQ. Their evaluation was done using the following  
220 equations:  $\Phi_M = (F_M - F_0) / F_M$  (Kitajima and Butler, 1975);  $\Phi'_M = (F'_M - F_s) / F'_M$  (Genty et  
221 al., 1989);  $\text{NPQ} = (F_M - F'_M) / F'_M$  (Bilger and Björkman, 1990). The maximal electron  
222 transport rate ( $\text{ETR}_{\text{MAX}}$ ), light saturation coefficient ( $E_k$ ) and light efficiency use ( $\alpha$ )  
223 was calculated according to Lacour et al. (2017). The Plant Efficiency Analyzer (PEA,  
224 Hansatech, Instruments Ltd, UK) was used to determine the polyphasic rise in  
225 fluorescence transients. Transients were induced by a 2s red (maximal emission at 650  
226 nm) light pulse with 3600  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  (Strasser et al., 1995). The O-J-I-P  
227 curves of the microalgae were determined and functional parameters evaluating the  
228 PSII energy fluxes under environmental stresses were calculated. All parameter  
229 definitions are in the Supplementary Material (Table S1).

## 230 2.5 Reactive oxygen species (ROS) measurement

231 Intracellular ROS was evaluated by BD Accuri C6 flow cytometer (Biosciences,  
232 San Jose, CA, USA) using the fluorescent dye 2',7'-dichlorodihydrofluorescein  
233 diacetate ( $\text{H}_2\text{DCFDA}$ ) (Molecular probes, Eugene, OR, USA) as described in  
234 (Stachowski-Haberkorn et al. 2013). Cells were analyzed by forward and side scatters  
235 and measured the fluorescence channel FL1 with excitation wavelength of 488 nm and  
236 emission wavelength of 530 nm. To eliminate potential signal alterations brought on by  
237 pesticide effects on FL1 fluorescence, we presented the results as FL1 ratios  
238 ( $\text{H}_2\text{DCFDA}$ -stained samples FL1 value divided by non-contaminated samples FL1  
239 values).

## 240 2.6 Antioxidant enzyme activity measurements

241 After pesticide exposure (72h), microalgal cultures (50 mL) were centrifuged at  
242 15000×g for 25 min at 4 °C, and the pellet was kept into a 2 mL microtube covered  
243 with aluminum foil. After adding 1 mL extraction buffer, samples were immediately  
244 plunged into liquid nitrogen and kept at -80 °C until analysis. For each extracted sample  
245 enzyme activities were determined with Cary 300 UV spectrophotometer (Varian,  
246 USA). Cells were broken with the help of liquid nitrogen, grinding one time, and were  
247 then centrifuged at 15000×g for 25 min at 4 °C prior to analysis. Each sample was  
248 divided into three replicates for analyzing the superoxide dismutase (SOD) and catalase  
249 (CAT) according to Vitoria et al. (2001) and Rao et al. (1996) respectively.

## 250 2.7 Statistical analyses

251 JMP software 10.0 (SAS Institute Inc) was used for statistical evaluations. Data  
252 were verified for normality (Shapiro–Wilk test) and homogeneity (Bartlett test) and  
253 then statistically evaluated using either one or two-way analysis of variance (ANOVA).  
254 Interactions between pesticide concentrations and different species were considered in  
255 2-way ANOVA. When there were significant differences in the response variables  
256 between treatments, contrast analysis (Tukey's HSD test) was used. The EC<sub>50</sub>  
257 (concentration needed to induce 50% of the maximum effect) values for response  
258 variables (growth,  $\Phi_M$ , and  $\Phi'_M$ ) were calculated from the nonlinear least-square fits by  
259 using the inverse of the regression curve (Juneau et al. 2001).


## 260 3. Results

### 261 3.1 Effects of pesticides exposure on cell growth and cell biovolume

#### 262 3.1.1 Effects of atrazine and simazine

263 The presence of atrazine and simazine for 72h significantly inhibited the growth  
264 of all algal species (Tukey's HSD,  $P < 0.05$ ), a growth inhibition that was further  
265 exacerbated at increased atrazine and simazine concentration (Fig. 1). The growth-EC<sub>50</sub>  
266 for temperate *C. neogracile* (T-CN), Arctic *C. neogracilis* (A-CN), temperate *M. bravo*  
267 (T-MB) and Arctic *M. polaris* (A-MP) was 143, 86, 52 and 82 µg/L respectively for

268 atrazine, and 166, 171, 69 and 111  $\mu\text{g/L}$  for simazine (Table 2). Cellular biovolume of  
 269 the T-CN ( $150 \mu\text{m}^3$ ) was intrinsically nearly three times that of A-CN ( $50 \mu\text{m}^3$ ), while  
 270 the biovolume of temperate *M. bravo* was almost the same as Arctic *M. polaris*. The  
 271 treatment of atrazine and simazine tend to increase the cell biovolumes of all species  
 272 by 2-12%, although not significantly for all studied species/treatments (Tukey's HSD,  
 273  $P > 0.05$ , Fig. S3).



274  
 275 Figure 1. The effects of atrazine and simazine on the growth of four species, including  
 276 (red color) temperate *C. neogracile* (T-CN), temperate *M. bravo* (T-MB), Arctic *C.*  
 277 *neogracilis* (A-CN) and (blue color) Arctic *M. polaris* (A-MP) after 72 h exposure. Data  
 278 are expressed as means  $\pm$  SD ( $n = 6$ ).

279 Table 2. The  $\text{EC}_{50}$  of atrazine and simazine based on the growth, the maximal PSII  
 280 quantum yield ( $\Phi_M$ ), the operational PSII quantum yield ( $\Phi'_M$ ). The same column  
 281 superscript letters (a-d) showed significant differences among strains for  $\text{EC}_{50}$ . n.d. =  
 282 not determined. Data are expressed as means  $\pm$  SD ( $n = 6$ )

Species	Pesticides	EC <sub>50</sub> -Φ <sub>M</sub>		EC <sub>50</sub> -Φ' <sub>M</sub>		EC <sub>50</sub> -growth rate	
		Atrazine	Simazine	Atrazine	Simazine	Atrazine	Simazine
T-CN		n.d.	n.d.	66±1.4 <sup>c</sup>	142±6.7 <sup>d</sup>	275±10.6 <sup>d</sup>	306±39.2 <sup>d</sup>
A-CN		n.d.	4073±56.3 <sup>c</sup>	37±2.1 <sup>b</sup>	62±2.1 <sup>c</sup>	188±24.3 <sup>c</sup>	128±20.7 <sup>c</sup>
T-MB		104±5.6 <sup>a</sup>	46±1.5 <sup>a</sup>	31±2.3 <sup>a</sup>	49±1.8 <sup>b</sup>	55±3.3 <sup>a</sup>	68±2.7 <sup>a</sup>
A-MP		156±14.3 <sup>b</sup>	703±67.3 <sup>b</sup>	36±1.7 <sup>b</sup>	46±1.9 <sup>a</sup>	75±2.8 <sup>b</sup>	111±6.8 <sup>b</sup>

283

### 284 3.1.2 Effects of trifluralin and chlorpyrifos

285 The growth of Arctic microalgae (*C. neogracilis*, *M. polaris*) and temperate  
286 microalgae (*C. neogracile*, *M. bravo*) was drastically inhibited in the presence of  
287 chlorpyrifos at 200 µg/L and 500 µg/L, the exception was for *M. bravo* at 200 µg/L  
288 chlorpyrifos (Tukey's HSD,  $P < 0.05$ ). Overall, Arctic microalgae showed greater  
289 decline in their growths than their temperate counterparts in the presence of  
290 chlorpyrifos (Table S2). The chlorpyrifos treatment induced a significant increase  
291 (Tukey's HSD,  $P < 0.05$ ) in the cell biovolume of all species. Overall, the biovolumes  
292 of Arctic microalgae increased by more than 100% compared to the temperate  
293 counterparts except for the A-CN at 200 µg/L chlorpyrifos (Table S1).


294 The growth of all species was significantly decreased by trifluralin (Tukey's HSD,  
295  $P < 0.05$ ), but Arctic microalgae had smaller growth reductions than did their temperate  
296 counterparts in the presence of trifluralin, which is contrary to what was observed in  
297 the presence of chlorpyrifos. Cell biovolume of all species except for *M. bravo*  
298 increased in the presence of trifluralin (Table S1).

### 299 3.2 Effects of pesticides on pigment contents

#### 300 3.2.1 Effects of atrazine and simazine

301 Pesticides had various impacts on pigment contents of the four studied microalgae  
302 after 72 h treatment. The cellular Chl *a* and Car contents of temperate and Arctic  
303 *Chaetoceros* did not show significant changes with increasing atrazine and simazine  
304 concentrations, except for simazine at 250 µg/L (Tukey's HSD,  $P > 0.05$ , Fig. 2),

305 resulting in an unchanged Car/Chl *a* ratio. For *M. bravo*, Chl *a* and Car increased  
 306 slightly at low atrazine concentrations and then significantly decreased at higher  
 307 concentrations (>100 µg/L, Tukey's HSD, P < 0.05, Fig. 2), while atrazine did not affect  
 308 the pigment composition of *M. polaris*. Simazine treatments did not induce a significant  
 309 change in Chl *a* and Car contents after 72 h of exposure for the temperate and Arctic  
 310 *Micromonas* (Tukey's HSD, P > 0.05). Under control conditions, Chl *a* for the temperate  
 311 *Chaetoceros* and *Micromonas* were respectively almost 2.5 and 2.3 times the  
 312 concentrations of the Arctic *Chaetoceros* and *Micromonas*, and carotenoids 1.5 and 1.2  
 313 times higher.



314  
 315 Figure 2. The effects of atrazine (A, C) and simazine (B, D) on the pigment contents of  
 316 four species, including (red color) temperate *C. neogracile* (T-CN) and temperate *M.*  
 317 *bravo* (T-MB), (blue color) Arctic *C. neogracilis* (A-CN) and Arctic *M. polaris* (A-MP)  
 318 after 72 h exposure. Data are expressed as means ± SD (n = 6).

### 319 3.2.2 Effects of trifluralin and chlorpyrifos


320 Chl *a* and Car contents of both temperate microalgae diminished in the presence of  
 321 chlorpyrifos (Table S1). On the other hand, both Arctic microalgae increased their

322 contents in Chl *a* and Car except for *M. polaris* where the Car was unchanged. Car of  
323 both diatoms increased and Chl *a* increased for the lower concentrations of trifluralin,  
324 but decreased for the higher concentrations. In contrast, both prasinophytes decreased  
325 their Car contents, but Chl *a* content was not affected.

### 326 3.3 Effects of pesticides on PSII activity and the energy fluxes pathways

#### 327 3.3.1 Effects of atrazine and simazine on PSII activity and the energy fluxes pathways


328 The maximum PSII quantum yields ( $\Phi_M$ ) of the diatoms (T-CN and A-CN) were  
329 not affected by atrazine and simazine except at high concentrations (100 and 250  $\mu\text{g/L}$ ;  
330 Tukey's HSD,  $P > 0.05$ ) (Fig. 3). On the other hand,  $\Phi_M$  of the two studied prasinophytes  
331 (T-MB and A-MP) were reduced (at variable extents) at any tested concentrations  
332 except 5  $\mu\text{g/L}$  (Fig. 4). The operational PSII quantum yield ( $\Phi'_M$ ) significantly  
333 decreased (Tukey's HSD,  $P < 0.05$ ) with increasing atrazine and simazine  
334 concentrations for all studied species (Fig. 3 and 4). Both  $\Phi_M$  and  $\Phi'_M$  of A-CN  
335 decreased more than T-CN. In contrast,  $\Phi_M$  and  $\Phi'_M$  of A-MP decreased less than T-  
336 MB. The  $\text{EC}_{50}$  of  $\Phi'_M$  and  $\Phi_M$  also confirmed this result,  $\Phi'_M\text{-EC}_{50}$  of T-CN and A-CN  
337 are respectively 66.2, 36.7 and 142.2, 62.3 for atrazine and simazine;  $\Phi'_M\text{-EC}_{50}$  of T-  
338 MB and A-MP are respectively 30.7, 35.8 and 48.7, 45.8 for atrazine and simazine  
339 (Table 2). Therefore, similar trend was observed for  $\Phi_M\text{-EC}_{50}$  and the  $\Phi'_M\text{-EC}_{50}$ .



340

341 Figure 3. The effects of atrazine, simazine, chlorpyrifos and trifluralin on the maximal  
 342 PSII quantum yield ( $\Phi_M$ ) and the operational PSII quantum yield ( $\Phi'_M$ ) of temperate *C.*  
 343 *neogracile* (red color-T-CN) and Arctic *C. neogracililis* (blue color-A-CN) after 72 h  
 344 exposure. Chl and Tri represent chlorpyrifos and trifluralin respectively. Data are  
 345 expressed as means  $\pm$  SD (n = 6).





346

347 Figure 4. The effects of atrazine, simazine, chlorpyrifos and trifluralin on the maximal  
 348 PSII quantum yield ( $\Phi_M$ ), the operational PSII quantum yield ( $\Phi'_M$ ) and non-  
 349 photochemical quenching (NPQ) of temperate *M. bravo* (red color-T-MB) and Arctic  
 350 *M. polaris* (blue color-A-MP) after 72 h exposure. Chl and Tri represent chlorpyrifos  
 351 and trifluralin respectively. Data are expressed as means  $\pm$  SD (n = 6).


352

353 Table 3. The effects of atrazine and simazine on  $\Phi_M$ ,  $\Phi'_M$ ,  $\alpha$ , ETRmax, and  $E_k$  of  
 354 temperate *C. neogracile* (T-CN), *M. bravo* (T-MB) and Arctic *C. neogracilis* (A-CN),  
 355 Arctic *M. polaris* (A-MP) after 72 h exposure. The same column superscript letters (a-  
 356 d) showed significant differences among strains under different pesticide  
 357 concentrations. ND = not determined (Showing that calculations are strongly  
 358 influenced by treatment). Data expressed as means  $\pm$  SD (n = 6).

359


Parameter	Species T-CN							Simazine							
	A-CN	T-MB	A-MP	T-CN	A-CN	T-MB	A-MP	T-CN	A-CN	T-MB	A-MP	T-CN	A-CN	T-MB	A-MP
	Atrazine														
	0	0.65±0.00 <sup>a</sup>	0.58±0.00 <sup>a</sup>	0.65±0.01 <sup>a</sup>	0.62±0.04 <sup>a</sup>	0.66±0.01 <sup>b</sup>	0.58±0.01 <sup>a</sup>	0.66±0.01 <sup>b</sup>	0.58±0.01 <sup>a</sup>	0.64±0.00 <sup>a</sup>	0.63±0.00 <sup>a</sup>	0.64±0.00 <sup>a</sup>	0.58±0.01 <sup>a</sup>	0.64±0.00 <sup>a</sup>	0.63±0.02 <sup>a</sup>
$\Phi_M$	5	0.63±0.00 <sup>a</sup>	0.58±0.01 <sup>a</sup>	0.63±0.00 <sup>a</sup>	0.61±0.00 <sup>b</sup>	0.63±0.01 <sup>a</sup>	0.61±0.00 <sup>b</sup>	0.63±0.01 <sup>a</sup>	0.58±0.00 <sup>a</sup>	0.64±0.01 <sup>a</sup>	0.63±0.00 <sup>a</sup>	0.64±0.00 <sup>a</sup>	0.58±0.00 <sup>a</sup>	0.64±0.00 <sup>a</sup>	0.63±0.00 <sup>a</sup>
	25	0.62±0.01 <sup>b</sup>	0.56±0.01 <sup>a</sup>	0.61±0.01 <sup>a</sup>	0.60±0.00 <sup>b</sup>	0.61±0.01 <sup>ab</sup>	0.60±0.00 <sup>b</sup>	0.60±0.01 <sup>ab</sup>	0.57±0.00 <sup>ab</sup>	0.62±0.00 <sup>b</sup>	0.60±0.01 <sup>a</sup>	0.62±0.00 <sup>b</sup>	0.57±0.00 <sup>ab</sup>	0.62±0.00 <sup>b</sup>	0.60±0.01 <sup>a</sup>
	50	0.63±0.01 <sup>b</sup>	0.53±0.01 <sup>b</sup>	0.50±0.02 <sup>b</sup>	0.50±0.02 <sup>b</sup>	0.54±0.00 <sup>c</sup>	0.50±0.02 <sup>b</sup>	0.54±0.00 <sup>c</sup>	0.65±0.01 <sup>ab</sup>	0.55±0.00 <sup>b</sup>	0.56±0.00 <sup>c</sup>	0.56±0.00 <sup>c</sup>	0.55±0.00 <sup>b</sup>	0.56±0.00 <sup>c</sup>	0.55±0.01 <sup>b</sup>
	100	0.64±0.00 <sup>a</sup>	0.51±0.00 <sup>b</sup>	0.26±0.03 <sup>c</sup>	0.26±0.03 <sup>c</sup>	0.44±0.01 <sup>d</sup>	0.65±0.01 <sup>ab</sup>	0.44±0.01 <sup>ab</sup>	0.65±0.01 <sup>ab</sup>	0.52±0.00 <sup>c</sup>	0.48±0.00 <sup>d</sup>	0.48±0.00 <sup>d</sup>	0.52±0.00 <sup>c</sup>	0.48±0.00 <sup>d</sup>	0.47±0.01 <sup>c</sup>
	250	0.66±0.00 <sup>a</sup>	0.46±0.01 <sup>c</sup>	0.21±0.02 <sup>d</sup>	0.17±0.02 <sup>e</sup>	0.67±0.00 <sup>b</sup>	0.17±0.02 <sup>e</sup>	0.67±0.00 <sup>b</sup>	0.50±0.01 <sup>d</sup>	0.21±0.01 <sup>e</sup>	0.42±0.01 <sup>d</sup>	0.50±0.01 <sup>d</sup>	0.50±0.01 <sup>d</sup>	0.42±0.01 <sup>d</sup>	
	Simazine														
	0	0.65±0.00 <sup>a</sup>	0.55±0.01 <sup>a</sup>	0.56±0.00 <sup>a</sup>	0.49±0.04 <sup>a</sup>	0.66±0.01 <sup>a</sup>	0.49±0.04 <sup>a</sup>	0.66±0.01 <sup>a</sup>	0.56±0.00 <sup>a</sup>	0.56±0.00 <sup>a</sup>	0.49±0.02 <sup>a</sup>	0.56±0.00 <sup>a</sup>	0.56±0.00 <sup>a</sup>	0.49±0.02 <sup>a</sup>	
$\Phi'_M$	5	0.59±0.00 <sup>b</sup>	0.48±0.01 <sup>b</sup>	0.52±0.01 <sup>b</sup>	0.43±0.00 <sup>b</sup>	0.57±0.06 <sup>b</sup>	0.43±0.00 <sup>b</sup>	0.57±0.06 <sup>b</sup>	0.50±0.01 <sup>b</sup>	0.53±0.01 <sup>b</sup>	0.46±0.01 <sup>a</sup>	0.53±0.01 <sup>b</sup>	0.50±0.01 <sup>b</sup>	0.46±0.01 <sup>a</sup>	
	25	0.49±0.01 <sup>c</sup>	0.34±0.01 <sup>c</sup>	0.35±0.02 <sup>c</sup>	0.32±0.01 <sup>c</sup>	0.58±0.01 <sup>b</sup>	0.32±0.01 <sup>c</sup>	0.58±0.01 <sup>b</sup>	0.42±0.01 <sup>c</sup>	0.42±0.02 <sup>b</sup>	0.36±0.01 <sup>b</sup>	0.42±0.02 <sup>b</sup>	0.42±0.01 <sup>c</sup>	0.36±0.01 <sup>b</sup>	
	50	0.36±0.01 <sup>d</sup>	0.25±0.01 <sup>d</sup>	0.14±0.02 <sup>d</sup>	0.18±0.01 <sup>d</sup>	0.52±0.01 <sup>b</sup>	0.14±0.02 <sup>d</sup>	0.18±0.01 <sup>d</sup>	0.52±0.01 <sup>b</sup>	0.33±0.01 <sup>d</sup>	0.24±0.01 <sup>c</sup>	0.27±0.01 <sup>c</sup>	0.33±0.01 <sup>d</sup>	0.24±0.01 <sup>c</sup>	
	100	0.24±0.00 <sup>e</sup>	0.13±0.00 <sup>e</sup>	0.03±0.02 <sup>e</sup>	0.05±0.00 <sup>e</sup>	0.37±0.01 <sup>c</sup>	0.03±0.02 <sup>e</sup>	0.05±0.00 <sup>e</sup>	0.37±0.01 <sup>c</sup>	0.18±0.00 <sup>e</sup>	0.08±0.01 <sup>d</sup>	0.14±0.01 <sup>d</sup>	0.18±0.00 <sup>e</sup>	0.08±0.01 <sup>d</sup>	
	250	0.15±0.00 <sup>f</sup>	0.08±0.01 <sup>f</sup>	0.02±0.01 <sup>e</sup>	0.02±0.01 <sup>e</sup>	0.23±0.00 <sup>e</sup>	0.02±0.01 <sup>e</sup>	0.23±0.00 <sup>e</sup>	0.12±0.01 <sup>f</sup>	0.02±0.01 <sup>e</sup>	0.05±0.01 <sup>c</sup>	0.12±0.01 <sup>f</sup>	0.02±0.01 <sup>e</sup>	0.05±0.01 <sup>c</sup>	
	0	0.68±0.01 <sup>a</sup>	0.55±0.02 <sup>a</sup>	0.60±0.02 <sup>a</sup>	0.52±0.05 <sup>a</sup>	0.69±0.01 <sup>a</sup>	0.52±0.05 <sup>a</sup>	0.69±0.01 <sup>a</sup>	0.53±0.01 <sup>a</sup>	0.59±0.02 <sup>a</sup>	0.52±0.05 <sup>a</sup>	0.59±0.02 <sup>a</sup>	0.53±0.01 <sup>a</sup>	0.52±0.05 <sup>a</sup>	
a	5	0.62±0.01 <sup>b</sup>	0.46±0.01 <sup>b</sup>	0.54±0.00 <sup>b</sup>	0.48±0.00 <sup>b</sup>	0.63±0.06 <sup>a</sup>	0.48±0.00 <sup>b</sup>	0.63±0.06 <sup>a</sup>	0.47±0.03 <sup>b</sup>	0.52±0.01 <sup>b</sup>	0.50±0.01 <sup>a</sup>	0.52±0.01 <sup>b</sup>	0.47±0.03 <sup>b</sup>	0.50±0.01 <sup>a</sup>	
	25	0.51±0.02 <sup>c</sup>	0.33±0.01 <sup>c</sup>	0.35±0.02 <sup>c</sup>	0.34±0.02 <sup>c</sup>	0.60±0.01 <sup>a</sup>	0.34±0.02 <sup>c</sup>	0.60±0.01 <sup>a</sup>	0.41±0.01 <sup>c</sup>	0.43±0.02 <sup>c</sup>	0.37±0.01 <sup>b</sup>	0.43±0.02 <sup>c</sup>	0.41±0.01 <sup>c</sup>	0.37±0.01 <sup>b</sup>	
	50	0.36±0.01 <sup>d</sup>	0.23±0.02 <sup>d</sup>	0.13±0.02 <sup>d</sup>	0.17±0.01 <sup>d</sup>	0.53±0.02 <sup>b</sup>	0.13±0.02 <sup>d</sup>	0.17±0.01 <sup>d</sup>	0.31±0.01 <sup>d</sup>	0.27±0.01 <sup>d</sup>	0.24±0.01 <sup>c</sup>	0.27±0.01 <sup>d</sup>	0.31±0.01 <sup>d</sup>	0.24±0.01 <sup>c</sup>	
	100	0.22±0.01 <sup>e</sup>	0.11±0.00 <sup>e</sup>	ND	0.06±0.00 <sup>e</sup>	0.38±0.02 <sup>c</sup>	ND	0.06±0.00 <sup>e</sup>	0.38±0.02 <sup>c</sup>	0.17±0.01 <sup>e</sup>	0.09±0.01 <sup>d</sup>	0.14±0.01 <sup>e</sup>	0.17±0.01 <sup>e</sup>	0.09±0.01 <sup>d</sup>	
	250	0.11±0.01 <sup>f</sup>	0.10±0.01 <sup>e</sup>	ND	ND	0.20±0.00 <sup>d</sup>	ND	0.20±0.00 <sup>d</sup>	0.11±0.01 <sup>f</sup>	ND	0.05±0.01 <sup>d</sup>	0.11±0.01 <sup>f</sup>	ND	0.05±0.01 <sup>d</sup>	
	0	1337±439 <sup>a</sup>	1037±394 <sup>a</sup>	827±227 <sup>a</sup>	404±96 <sup>a</sup>	1366±540 <sup>a</sup>	404±96 <sup>a</sup>	1366±540 <sup>a</sup>	1147±80 <sup>a</sup>	679±107 <sup>a</sup>	434±163 <sup>a</sup>	1147±80 <sup>a</sup>	679±107 <sup>a</sup>	434±163 <sup>a</sup>	
ETRmax	5	1291±515 <sup>a</sup>	869±312 <sup>a</sup>	522±16 <sup>a</sup>	501±35 <sup>a</sup>	1481±833 <sup>a</sup>	501±35 <sup>a</sup>	1481±833 <sup>a</sup>	801±332 <sup>a</sup>	507±10 <sup>a</sup>	394±40 <sup>a</sup>	801±332 <sup>a</sup>	507±10 <sup>a</sup>	394±40 <sup>a</sup>	
	25	1601±804 <sup>a</sup>	708±35 <sup>a</sup>	490±249 <sup>a</sup>	325±70 <sup>b</sup>	1496±305 <sup>a</sup>	325±70 <sup>b</sup>	1496±305 <sup>a</sup>	741±261 <sup>a</sup>	631±212 <sup>a</sup>	370±103 <sup>a</sup>	741±261 <sup>a</sup>	631±212 <sup>a</sup>	370±103 <sup>a</sup>	
	50	1220±102 <sup>a</sup>	394±46 <sup>b</sup>	142±67 <sup>b</sup>	111±10 <sup>c</sup>	1550±491 <sup>a</sup>	142±67 <sup>b</sup>	111±10 <sup>c</sup>	1550±491 <sup>a</sup>	595±199 <sup>b</sup>	234±24 <sup>a</sup>	595±199 <sup>b</sup>	595±199 <sup>b</sup>	234±24 <sup>a</sup>	
	100	479±73 <sup>b</sup>	114±19 <sup>bc</sup>	2±1 <sup>b</sup>	28±4 <sup>c</sup>	938±256 <sup>a</sup>	2±1 <sup>b</sup>	28±4 <sup>c</sup>	938±256 <sup>a</sup>	219±17 <sup>b</sup>	44±6 <sup>b</sup>	219±17 <sup>b</sup>	219±17 <sup>b</sup>	44±6 <sup>b</sup>	
	250	160±13 <sup>b</sup>	18±1 <sup>c</sup>	0±0 <sup>b</sup>	6±2 <sup>c</sup>	378±2 <sup>b</sup>	6±2 <sup>c</sup>	378±2 <sup>b</sup>	61±16 <sup>c</sup>	0±0 <sup>c</sup>	14±3 <sup>b</sup>	61±16 <sup>c</sup>	0±0 <sup>c</sup>	14±3 <sup>b</sup>	
	0	1972±663 <sup>a</sup>	1872±703 <sup>a</sup>	1389±422 <sup>a</sup>	764±116 <sup>a</sup>	1993±785 <sup>a</sup>	764±116 <sup>a</sup>	1993±785 <sup>a</sup>	2151±128 <sup>ab</sup>	1141±154 <sup>a</sup>	820±229 <sup>a</sup>	2151±128 <sup>ab</sup>	1141±154 <sup>a</sup>	820±229 <sup>a</sup>	
Ek	5	2065±783 <sup>a</sup>	1898±666 <sup>a</sup>	972±29 <sup>a</sup>	1040±70 <sup>b</sup>	2493±1065 <sup>a</sup>	1040±70 <sup>b</sup>	2493±1065 <sup>a</sup>	1695±728 <sup>ab</sup>	978±1 <sup>a</sup>	796±66 <sup>a</sup>	1695±728 <sup>ab</sup>	978±1 <sup>a</sup>	796±66 <sup>a</sup>	
	25	3129±1502 <sup>a</sup>	2138±194 <sup>a</sup>	1416±704 <sup>a</sup>	945±170 <sup>c</sup>	2490±550 <sup>a</sup>	945±170 <sup>c</sup>	2490±550 <sup>a</sup>	1829±679 <sup>ab</sup>	1487±531 <sup>a</sup>	1003±258 <sup>a</sup>	1829±679 <sup>ab</sup>	1487±531 <sup>a</sup>	1003±258 <sup>a</sup>	
	50	3350±183 <sup>a</sup>	1735±219 <sup>a</sup>	1036±356 <sup>a</sup>	654±30 <sup>d</sup>	2899±882 <sup>a</sup>	1036±356 <sup>a</sup>	654±30 <sup>d</sup>	1943±657 <sup>ab</sup>	1347±385 <sup>a</sup>	980±63 <sup>a</sup>	1943±657 <sup>ab</sup>	1347±385 <sup>a</sup>	980±63 <sup>a</sup>	
	100	2185±390 <sup>a</sup>	1068±209 <sup>ab</sup>	ND	479±79 <sup>c</sup>	2510±790 <sup>a</sup>	ND	479±79 <sup>c</sup>	1265±785 <sup>c</sup>	974±129 <sup>a</sup>	512±98 <sup>b</sup>	1265±785 <sup>c</sup>	974±129 <sup>a</sup>	512±98 <sup>b</sup>	
	250	1476±68 <sup>a</sup>	184±29 <sup>b</sup>	ND	ND	1908±35 <sup>a</sup>	ND	1908±35 <sup>a</sup>	561±183 <sup>c</sup>	ND	320±104 <sup>b</sup>	561±183 <sup>c</sup>	ND	320±104 <sup>b</sup>	
	0	0.99±0.22 <sup>a</sup>	2.40±0.35 <sup>a</sup>	1.76±0.17 <sup>a</sup>	6.48±2.36 <sup>a</sup>	1.07±0.25 <sup>a</sup>	6.48±2.36 <sup>a</sup>	1.07±0.25 <sup>a</sup>	2.62±0.68 <sup>a</sup>	1.76±0.17 <sup>a</sup>	0.78±0.07 <sup>a</sup>	2.62±0.68 <sup>a</sup>	1.76±0.17 <sup>a</sup>	0.78±0.07 <sup>a</sup>	
NPQmax	5	0.84±0.14 <sup>ab</sup>	2.04±0.29 <sup>a</sup>	1.69±0.27 <sup>a</sup>	2.80±0.23 <sup>b</sup>	0.77±0.35 <sup>a</sup>	2.80±0.23 <sup>b</sup>	0.77±0.35 <sup>a</sup>	1.64±0.09 <sup>a</sup>	1.69±0.27 <sup>a</sup>	0.72±0.05 <sup>b</sup>	1.64±0.09 <sup>a</sup>	1.69±0.27 <sup>a</sup>	0.72±0.05 <sup>b</sup>	
	25	0.76±0.09 <sup>ab</sup>	1.72±0.14 <sup>a</sup>	2.21±0.07 <sup>a</sup>	2.75±0.37 <sup>bc</sup>	0.90±0.22 <sup>a</sup>	2.75±0.37 <sup>bc</sup>	0.90±0.22 <sup>a</sup>	1.68±0.14 <sup>a</sup>	2.21±0.07 <sup>a</sup>	0.92±0.05 <sup>bc</sup>	1.68±0.14 <sup>a</sup>	2.21±0.07 <sup>a</sup>	0.92±0.05 <sup>bc</sup>	
	50	0.53±0.05 <sup>b</sup>	1.81±0.17 <sup>a</sup>	0.83±0.54 <sup>b</sup>	0.49±0.08 <sup>bcd</sup>	0.90±0.24 <sup>a</sup>	0.83±0.54 <sup>b</sup>	0.49±0.08 <sup>bcd</sup>	1.32±0.18 <sup>b</sup>	1.32±0.18 <sup>b</sup>	0.70±0.08 <sup>bd</sup>	1.32±0.18 <sup>b</sup>	1.32±0.18 <sup>b</sup>	0.70±0.08 <sup>bd</sup>	
	100	0.25±0.12 <sup>bc</sup>	4.88±0.51 <sup>b</sup>	0.14±0.04 <sup>c</sup>	0.07±0.01 <sup>c</sup>	0.65±0.08 <sup>b</sup>	0.14±0.04 <sup>c</sup>	0.07±0.01 <sup>c</sup>	2.76±0.15 <sup>c</sup>	2.76±0.15 <sup>c</sup>	0.36±0.04 <sup>de</sup>	2.76±0.15 <sup>c</sup>	2.76±0.15 <sup>c</sup>	0.36±0.04 <sup>de</sup>	
	250	0.12±0.01 <sup>d</sup>	2.32±0.35 <sup>c</sup>	0.07±0.03 <sup>c</sup>	0.03±0.01 <sup>c</sup>	0.40±0.08 <sup>c</sup>	0.03±0.01 <sup>c</sup>	0.40±0.08 <sup>c</sup>	3.33±0.86 <sup>c</sup>	0.07±0.03 <sup>c</sup>	0.22±0.00 <sup>de</sup>	3.33±0.86 <sup>c</sup>	0.07±0.03 <sup>c</sup>	0.22±0.00 <sup>de</sup>	

361           Calculated parameters acquired from the rapid Chl *a* fluorescence kinetics provide  
362 useful indication on how pesticides may affect energy fluxes within PSII (Force et al.,  
363 2003). The electron transport rate per active reaction center ( $ET_0/RC$ ) of all species  
364 significantly decreased under the treatment of atrazine and simazine by stopping  
365 electron flow between  $Q_A$  and  $Q_B$  (Fig. 5 and 6). The energy conservation parameter of  
366  $PI_{ABS}$  also declined under these treatment conditions, but to an extent that was species-  
367 dependent. We found that  $PI_{ABS}$  of the A-CN was more affected than T-CN (Fig. 5A-  
368 D); in contrast, the  $PI_{ABS}$  of the temperate *M. bravo* was more affected than in the Arctic  
369 *M. polaris* (Fig. 6A-D). The effective dissipation per reaction center ( $DI_0/RC$ ) of the  
370 four species increased because of the high dissipation of the inactive RCs in the  
371 presence of atrazine and simazine. Similarly, the absorption flux per reaction center  
372 ( $ABS/RC$ ), a proxy of the PSII antenna size, was increased by up to 202% in the  
373 presence of atrazine or simazine, except for the temperate *C. neogracile*. The  $DI_0/RC$   
374 was increased in agreement with the change in  $ABS/RC$  in presence of atrazine and  
375 simazine, and the effect of  $DI_0/RC$  on Arctic diatom *C. neogracilis* was stronger (4.5  
376 and 1.6 times) than for the temperate *C. neogracile*; for the prasinophytes we observed  
377 the contrary (the effect on  $DI_0/RC$  of the temperate *M. bravo* was stronger than for *M.*  
378 *polaris*). The maximal rate at which excitons are caught by the active reaction centers  
379 ( $TR_0/RC$ ) was only altered by simazine for *M. bravo*.



380

381 Figure 5. The effects of atrazine and simazine on the chlorophyll fluorescence  
 382 parameters of temperate *C. neogracile* (T-CN) and Arctic *C. neogracilis* (A-CN) after  
 383 72 h exposure.



384

385 Figure 6. The effects of atrazine and simazine on the chlorophyll fluorescence  
 386 parameters of temperate *M. bravo* (T-MB) and Arctic *M. polaris* (A-MP) after 72 h  
 387 exposure.


### 388 3.3.2 Effects of trifluralin and chlorpyrifos on PSII activity and the energy fluxes 389 pathways

390 The  $\Phi_M$  and  $\Phi'_M$  of all species declined in the presence of high concentrations (200  
 391 and 500  $\mu\text{g/L}$ ) of chlorpyrifos and trifluralin (Fig. 3 and 4), but the declines differed  
 392 between temperate (*C. neogracile* and *M. bravo*) and Arctic species (*C. neogracilis* and  
 393 *M. polaris*). Indeed, both  $\Phi_M$  and  $\Phi'_M$  of A-CN decreased more than T-CN when  
 394 exposed to chlorpyrifos, but this trend was reversed in the presence of trifluralin. For  
 395 the prasinophytes, we observed similar effect where both  $\Phi_M$  and  $\Phi'_M$  of Arctic *M.*  
 396 *polaris* decreased more in the presence of chlorpyrifos than ones of the temperate *M.*  
 397 *bravo*. This trend was reversed in the presence of trifluralin for Arctic *M. polaris* which  
 398 declined less than did the temperate *M. bravo*. These results indicate greater sensitivity  
 399 of Arctic species to chlorpyrifos than temperate counterparts, and vice versa with

400 trifluralin. In the presence of chlorpyrifos and trifluralin,  $ET_0/RC$  and  $PI_{ABS}$  of both  
401 *Chaetoceros* species decreased with increasing pesticide concentration concomitantly  
402 with increasing  $DI_0/RC$  (Fig. S1). In response to trifluralin, Arctic and temperate  
403 *Micromonas* demonstrated opposite  $DI_0/RC$  and  $ABS/RC$  trends: these parameters  
404 increased for the temperate *M. bravo* and declined for the Arctic *M. polaris* (Fig. S1).  
405 Similarly,  $ET_0/RC$  and  $DI_0/RC$  responses in the two *Micromonas* species showed  
406 opposite response to chlorpyrifos.


#### 407 3.4 Effects of pesticides on reactive oxygen species, antioxidant enzyme activity and 408 protein content

409 At the lowest concentrations of atrazine and simazine (5 and 25  $\mu\text{g/l}$ ), the ROS  
410 content of all species was unchanged compared to control conditions (Tukey's HSD,  
411  $P>0.05$ ) (Fig. 7). However, the ROS content significantly increased (Tukey's HSD,  
412  $P<0.05$ ) at higher concentrations of pesticides (50, 100 and 250  $\mu\text{g/l}$ ). Despite similar  
413 trends, in the presence of atrazine, simazine and trifluralin, the ROS content of A-CN  
414 increased more than for the temperate *C. neogracile*, while the opposite occurred in the  
415 presence of chlorpyrifos. We observed a less marked increase of ROS content in the  
416 presence of pesticides for the Arctic *M. polaris* compared to the temperate *M. bravo*.  
417 Moreover, under control conditions, the ROS content of Arctic microalgae was lower  
418 than its temperate counterparts (Table S1). The activity of SOD and CAT of all species  
419 strongly increased with atrazine, simazine and trifluralin concentrations (Fig. 8).  
420 Concomitantly, the protein concentration per cell significantly decreased in the  
421 presence of pesticides, indicating that pesticides induced oxidative stress and a  
422 subsequent induction of mechanisms involved in the removal  $O_2^-$  and  $H_2O_2$ .  
423 Surprisingly, for chlorpyrifos, the activity of SOD and CAT, and the total protein  
424 content of temperate *C. neogracile* and *M. bravo* decreased, suggesting that the  
425 antioxidant enzyme system was insufficient to cope with the impact of these pesticides  
426 or at least was not the main protective measure.



427

428 Figure 7. The effects of atrazine (A, C) and simazine (B, D) on the reactive oxygen  
 429 species (ROS) of four species, including (red color) temperate *C. neogracile* (T-CN)  
 430 and temperate *M. bravo* (T-MB), (blue color) Arctic *C. neogracilis* (A-CN) and Arctic  
 431 *M. polaris* (A-MP) after 72 h exposure. Data are expressed as means  $\pm$  SD (n = 6).



432

433 Figure 8. The effects of atrazine (A, C) and simazine (B, D) on the catalase (CAT) and  
 434 superoxide dismutase (SOD) of four species, including (red color) temperate *C.*

435 *neogracile* (T-CN) and temperate *M. bravo* (T-MB), (blue color) Arctic *C. neogracilis*  
436 (A-CN) and Arctic *M. polaris* (A-MP) after 72 h exposure. Data are expressed as means  
437  $\pm$  SD (n = 6).

438

## 439 **4. Discussion**

440 4.1 Different physiological characteristics between Arctic microalgae and their  
441 temperate counterparts

442 It is well accepted that at low temperatures Rubisco activity is the limiting step for  
443 growth (Young et al. 2015). Similarly, the cell biovolume of microalgae can be affected  
444 by growth temperature (Daufresne et al. 2009), and smaller cells have a larger  
445 surface/volume ratio compared to bigger cells, which facilitates better utilization of  
446 resources such as light and nutrients for growth (Wirth et al. 2019). In our study, the  
447 temperate and Arctic prasinophytes have the same cell biovolume (5-8  $\mu\text{m}^3$ ), since  
448 prasinophyte morphological and biochemical characteristics are highly conserved and  
449 may not be strongly modified by the long term growth conditions (McKie-Krisberg and  
450 Sanders 2014). The growth rate of temperate *M. bravo* (T-MB) is almost twice that of  
451 the Arctic *M. polaris* (A-MP), which is consistent with previous studies, indicating that  
452 growth rates of polar green algae are usually lower than for temperate one  
453 (Kottuparambil et al. 2017, Lacour et al. 2017). Thus, not surprisingly, A-MP had lower  
454 ETR<sub>max</sub>, E<sub>k</sub> and  $\alpha$  compared to T-MB, indicating that the utilization efficiency of light  
455 energy and photosynthetic capacity of A-MP were weaker than those of T-MB (Table  
456 3). This was also supported by the lower maximal and operational PSII quantum yields  
457 ( $\Phi_M$  and  $\Phi'_M$ ) in A-MP (Table 3). Interestingly, these differences were not found in the  
458 comparison between congeneric Arctic (*C. neogracilis*-A-CN) and temperate (*C.*  
459 *neogracile*-T-CN) diatoms. Indeed, Arctic and temperate diatoms had similar growth  
460 rate, ETR<sub>max</sub>, E<sub>k</sub> and  $\alpha$  (Table 3), indicating that A-CN, while growing at a low  
461 temperature, has developed some strategies to improve the ability to harvest light  
462 energy and light utilization efficiency to optimize growth. However, these strategies did



463 not prevent lower  $\Phi_M$  and  $\Phi'_M$  in A-CN compared to T-CN (Table 3), which is partly  
464 due to the presence of sustained NPQ (Lacour et al. 2018). Furthermore, A-CN with a  
465 much lower cell biovolume compared to T-CN can make better use of resources to  
466 benefit growth, which may partially explain the same growth rates of A-CN and T-CN.

467 The growth rates of both diatoms were higher than for *Micromonas*, as previously  
468 shown in other species' comparisons (the diatom *Thalassiosira hyalina* with  
469 *Micromonas pusilla*) (Hoppe et al. 2018), probably due to the stronger ability of  
470 diatoms to increase the Rubisco gene expression when its activity is reduced at low  
471 temperature (Young et al. 2015). However, it is uncertain if prasinophytes also exhibit  
472 these acclimation responses (Hoppe et al. 2018). Lacour et al. (2017) observed that for  
473 a given Chl *a* content, polar diatoms grow more slowly than do temperate ones,  
474 suggesting that this difference is related to energy allocation. Interestingly, in our study,  
475 A-CN and T-CN had the same growth rate, but A-CN had a much lower Chl *a* content  
476 compared to its temperate counterpart. This indicates that different polar diatoms may  
477 have developed different adaptation strategies in concordance with their growth  
478 environment. Indeed, diatoms have different inherent NPQ abilities in response to their  
479 respective habitats (Croteau et al. 2021, Croteau et al. 2022). Lower Chl *a* content in  
480 both Arctic microalgae helps reduce excitation pressure on the photosynthetic reaction  
481 center, particularly in situations constraining growth, such as cold environments  
482 (Halsey and Jones 2015). Recent studies have demonstrated that for the majority of  
483 polar microalgal groups, NPQ is an essential element of the species-specific  
484 photoadaptive strategies (Croteau et al. 2021, Galindo et al. 2017). Together with low  
485 Chl *a*, polar microalgae also can induce high NPQ through an efficient de-epoxidation  
486 process to jointly protect the photosynthetic apparatus against environmental stress  
487 such as sudden increase in light intensity and temperature modifications (Lacour et al.  
488 2020, Ni et al. 2017). In comparison, both Arctic microalgae, having lower ROS content  
489 and high CAT and SOD activities, should be more likely to adapt and survive at low  
490 Arctic habitat temperatures (Blanc et 2012). In summary, Arctic species appear to have  
491 evolved different ecophysiological characteristics than their temperate counterparts:

492 Arctic microalgae have (1) lower Chl *a* and carotenoid contents, (2) a much higher  
493 intrinsic NPQ<sub>max</sub>, (3) lower ROS content and (4) higher CAT and SOD activities,  
494 compared to temperate species.

#### 495 4.2 Effects of pesticides on Arctic microalgae and their temperate counterparts

##### 496 4.2.1 Effects of pesticides on diatoms (A-CN and T-CN)

497 PSII inhibiting herbicides (such as atrazine and simazine) that can bind to the Q<sub>B</sub>  
498 site on the D1 protein of PSII, inhibit the PSII-PSI electron transport resulting in high  
499 excitation pressure on PSII and ROS generation (Bai et al. 2015). If the photoprotective  
500 processes, NPQ and ROS scavenging system, are ineffective, reduced energy  
501 production and cellular damages occur, and ultimately algal growth might be reduced.  
502 Our study demonstrated that atrazine and simazine significantly inhibited the growth of  
503 both diatoms, as seen previously for *Navicula pelliculosa* (Chalifour and Juneau 2011).  
504 Our observations were linked to the significant reduction in  $\Phi'_M$  and the electron  
505 transport rate per active RC (ET<sub>0</sub>/RC). Concomitantly, dissipation of excess light  
506 energy (DI<sub>0</sub>/RC) increased but was not sufficient to protect the PSII since  $\Phi_M$ , a proxy  
507 of the PSII RC integrity, was affected, as previously shown when *Phaeodactylum*  
508 *tricornutum* was treated with PSII inhibiting herbicides (Debenest et al. 2010). This  
509 result is also supported by the significant decrease in PI<sub>ABS</sub>, a sensitive indicator of plant  
510 health (Bayat et al. 2018) (Fig. 5A-D). The other unchanged PSII energy fluxes  
511 (ABS/RC and TR<sub>0</sub>/RC) at low pesticide concentrations, together with the minor  
512 variations in the carotenoid contents, indicate that the protection of the photosynthetic  
513 electron transport chain from the ROS produced in the presence of atrazine and  
514 simazine, are likely due to other protective strategies (like antioxidant enzymatic  
515 systems). One could expect that, as the first line of defense against the excess of light  
516 energy under stressful conditions, NPQ would be activated (Kress and Jahns 2017,  
517 Müller et al. 2001). However, NPQ<sub>max</sub> decreased with increasing atrazine and  
518 simazine concentrations (Fig. 3). As shown in previous studies, the decline of NPQ  
519 in the presence of pesticides is attributed to the low buildup of the proton gradient across  
520 the thylakoid membranes, since electron transport is decreased (Chalifour and Juneau

521 2011, Gomes and Juneau 2017). Therefore, we supposed that the inhibition of the  $\Delta\text{pH}$ -  
522 dependant non-photochemical energy dissipation mechanism leads to a reduced ability  
523 to decrease excitation pressure at PSII RC, resulting in higher ROS production.  
524 Although SOD and CAT activities were significantly increased with increasing atrazine  
525 and simazine concentrations, it was not sufficient to cope entirely with the ROS  
526 production induced by increasing atrazine and simazine concentrations (Fig. 7).

527 Overall, as shown by the investigation of the physiological parameters in the  
528 presence of atrazine and simazine and the determined  $\text{EC}_{50}$  (Table 2), A-CN was more  
529 sensitive than the T-CN to atrazine and simazine. We suspect that the more pronounced  
530 PSII RC inactivation of A-CN results from its smaller cell biovolume, thus increasing  
531 overall contact with the pesticide molecules and resulting in an enhanced absorption of  
532 the contaminants (Weiner et al. 2004). However, the antioxidant enzyme system (SOD  
533 and CAT) induced in A-CN was insufficient to cope with the ROS production in the  
534 presence of atrazine and simazine. Furthermore, as previously showed in psychrophilic  
535 diatoms, PSII repair rates are slower than the ones found in temperate diatoms (Petrou  
536 et al. 2010), since lower temperatures decreased enzyme activity and metabolism  
537 (Morgan-Kiss et al. 2006). On the contrary, compared to the Arctic diatom, the  
538 temperate diatom potentially showed lower absorption of atrazine and simazine, the  
539 capacity for an efficient antioxidant enzyme system and probably higher rate of PSII  
540 repair cycle, which ultimately lead to its lower sensitivity to these herbicides.

541 For trifluralin, interestingly, we observed that A-CN was more tolerant than T-CN  
542 according to the growth, and photosynthetic activity ( $\Phi_M$  and  $\Phi'_M$ ; Fig. 3). According  
543 to our data (Table S1, Fig. S1), the greater decrease of  $\text{PI}_{\text{ABS}}$  and  $\text{ABS/RC}$  for T-CN  
544 compared to A-CN, indicated that PSII RC was more damaged in T-CN. However, the  
545 Arctic diatom has more effective antioxidant capacity (SOD and CAT activities) than  
546 its temperate counterpart under the same concentration of trifluralin, indicating that its  
547 tolerance to trifluralin seems to mainly depend on the high efficiency of the antioxidant  
548 system. Although insecticides are not intended to affect plants and algae, chlorpyrifos  
549 has been shown to induce some deleterious impacts at the cellular and population levels

550 (Asselborn et al. 2015), leading to the impairment of cell morphology and growth  
551 (Asselborn et al. 2006, Garrido et al. 2019), and the decrease in diversity of diatoms  
552 (Stratton, 1987). Similar effects were seen in our study, where chlorpyrifos not only  
553 affected the growth, and photosynthesis of both diatoms (Fig. 3; A-CN was more  
554 affected than T-CN), but also caused oxidative stress. The observed difference in the  
555 sensitivity to chlorpyrifos of diatoms was mainly reflected at the electron transport level,  
556 where chlorpyrifos induced the accumulation of  $Q_A^-$  and prevented electron transfer  
557 downstream of  $Q_A$  (revealed by the more pronounced increase in O-J and J-I phases for  
558 the Arctic diatom, Fig. S2), also evidence in the significant decrease of  $\Phi'_M$ , ETo/RC  
559 and  $PI_{ABS}$  for A-CN (Fig. S1). In comparison, this impact on electron transport was  
560 accompanied by a weaker ability to dissipate excess energy (Dio/RC) in the Arctic  
561 diatom, resulting in its greater sensitivity to chlorpyrifos. In addition, some authors have  
562 suggested that insecticides disturb the cell cycle of *Selenastrum capricornutum* since  
563 they observed the inhibition of cell-separation, resulting in the intracellular  
564 accumulation of macromolecules, which are responsible for increasing the biovolume  
565 in A-CN (Fernandez et al. 2021, Rioboo et al. 2002).

#### 566 4.2.2 Effects of pesticides on prasinophytes (A-MP and T-MB)

567 Prasinophytes responded to atrazine and simazine similarly to diatoms when  
568 growth, pigment composition and photosynthetic efficiency were evaluated. In contrast  
569 to diatoms, A-MP was more tolerant than T-MB to atrazine and simazine (Table 2 and  
570 Fig. 6). The more pronounced damage to the PSII RC ( $\Phi_M$ ), drastic inhibition of  
571 photosynthetic electron transport ( $\Phi'_M$  and ETo/RC) and decreased light conversion  
572 efficiency ( $PI_{ABS}$ ) at the PSII RC level in T-MB eventually induced higher production  
573 of ROS, which further damaged these photosynthetic components. We propose several  
574 reasons why A-MP was more tolerant to atrazine and simazine. First, A-MP has lower  
575 Chl *a* content than T-MB in the absence of pesticides, indicating that A-MP may have  
576 lower PSII content and therefore fewer available molecular targets for atrazine and  
577 simazine (DeLorenzox et al. 2004). Previous studies have demonstrated that Arctic  
578 *Micromonas* have lower active PSII levels than their temperate counterparts (Ni et al.

2017). Concomitantly, the lower Chl *a* content in A-MP helps to reduce excitation pressure on the PSII RC caused by atrazine and simazine. Second, the temperate species produced more ROS in the presence of atrazine and simazine compared to the Arctic one, but the induction of CAT and SOD were insufficient to prevent oxidative damage. In comparison, the Arctic species produced lower levels of ROS and have relatively lower CAT and SOD activities compared to the temperate species, which implies that these enzymes can successfully scavenge ROS. ROS has to reach a threshold level to increase the activity of the antioxidant enzyme system (Anu et al. 2016), which may indicate why the temperate species were more affected by the ROS production than the Arctic species in the presence of atrazine and simazine. Finally, A-MP had higher intrinsic NPQ to which was only slightly decreased by atrazine and simazine compared to T-MB (Fig. 4), which would protect PSII (Bai et al. 2015). In summary, T-MB showed higher sensitivity to atrazine and simazine than A-MP, due to the non-effectiveness of its NPQ and antioxidant enzymes to cope with excess light energy and oxidative stress, resulting in photosynthetic damage. In comparison, A-MP seems to mainly rely on NPQ rather than antioxidant enzymes under atrazine and simazine stress.

We observed that growth, photosynthetic efficiency, and electron transfer of the T-MB were more sensitive to the herbicide trifluralin than for the A-MP. Furthermore, significantly increased  $Df_0/RC$ ,  $ABS/RC$  and decreased  $ET_0/RC$  of T-MB in the presence of trifluralin showed that the excess excitation energy caused by a certain number of inactivated RCs, was mostly dissipated. We propose that the different sensitivities of the prasinophytes to trifluralin are mainly determined by the antioxidant enzyme system, since the Arctic species induced five times higher SOD and CAT activities, even if it produced less ROS than its temperate counterpart (Table S1). The insecticide chlorpyrifos, in a way similar to what was observed for diatoms, inhibited the growth of *Micromonas* (A-MP was more affected than T-MB) and doubled the cell biovolume of A-MP (Table S1), for the same reasons that we proposed for diatoms. For T-MB, chlorpyrifos did not affect the photosynthetic efficiency. Furthermore, the increased  $PI_{ABS}$  and  $ET_0/RC$ , and unchanged kinetics of  $Q_A-Q_B$  reduction compared to

608 the control (Fig. S1), indicate that PSII RCs and the whole photosynthetic electron  
609 transfer chain were protected under exposure to chlorpyrifos. By comparison,  
610 photosynthesis of A-MP was strongly affected by this insecticide, as indicated by the  
611 strong reduction of the active PSII RC population and suppression of electron transfer  
612 between  $Q_A$  and  $Q_B$  (Fig. S2).

#### 613 4.2.3 Comparative effect of pesticides on Arctic microalgae and their temperate 614 counterparts

615 In our study, based on the  $EC_{50}$  values for  $\Phi'_M$  and growth, the species sensitivity  
616 sequence was T-MB>A-MP>A-CN>T-CN in the presence of atrazine and simazine. For  
617 trifluralin, based on the impact on  $\Phi_M$  and  $\Phi'_M$  (since no  $EC_{50}$  value cannot be obtained  
618 even at very high concentrations) the species sensitivity sequence was T-MB>A-MP>T-  
619 CN>A-CN and A-CN>T-CN>A-MP>T-MB for chlorpyrifos (Fig. 3 and 4). We found  
620 that diatoms *Chaetoceros* were more tolerant to atrazine, simazine and trifluralin than  
621 the prasinophytes *Micromonas* for both temperate and Arctic strains. Diatoms are  
622 known to be dominant in most aquatic environments (Serôdio and Lavaud 2020), likely  
623 owing to their specific ecophysiological characteristics, like their high PSII/PSI ratio  
624 (Strzepek and Harrison 2004), their high non-photochemical energy dissipation  
625 potential (Lavaud and Lepetit 2013), and the presence of fucoxanthin that can prevent  
626 photooxidation (Tuchman et al. 2006). Furthermore, diatoms can more efficiently  
627 control the ATP/NADPH ratio during photosynthesis compared to other photosynthetic  
628 organisms, permitting them to optimize their carbon fixation and growth (Bailleul et al.  
629 2015). Indeed, we found that  $\Phi'_M$ , pigment, ETR<sub>max</sub>,  $E_k$  and  $\alpha$  in both diatoms were  
630 higher than in prasinophytes, suggesting that diatoms have higher intrinsic  
631 photosynthetic capacity and light utilization efficiency. Therefore, if we consider only  
632 the impact of pesticides, we propose that *Chaetoceros* would have a greater chance of  
633 survival and would become the dominant species in temperate and Arctic ecosystems  
634 contaminated with atrazine, simazine and trifluralin in relation to *Micromonas*. On the  
635 other hand, according to the  $EC_{50}$  of  $\Phi'_M$  and growth (Table 2), atrazine induced  
636 significantly more damage than simazine to all tested algae even though both molecules

637 have the same mode of action on the Q<sub>B</sub> site of D1 protein of PSII. Previous studies  
638 have shown that irgarol with high octanol/water partition coefficient (logK<sub>ow</sub>) is more  
639 toxic than diuron for marine microalgae as PSII inhibitor, and the higher logK<sub>ow</sub> of  
640 irgarol promotes its affinity for the Q<sub>B</sub> binding site, leading to its relative higher toxicity  
641 (Coquille et al. 2018, Dupraz et al. 2016, Kottuparambil et al. 2017). Therefore, we can  
642 assume that the higher toxicity of atrazine compared to simazine (logK<sub>ow</sub>=2.3) could be  
643 due to its higher logK<sub>ow</sub> (2.7) (Ronka 2016) and therefore to its higher affinity for the  
644 Q<sub>B</sub> binding site. Similarly, in the presence of atrazine, CAT and SOD activities were  
645 higher and the total protein content decreased more than in the presence of simazine,  
646 resulting in a decrease in the available protein for photoprotection processes, such as  
647 NPQ and the PSII repair cycle (Bai et al. 2015).

648

#### 649 **ACKNOWLEDGEMENTS**

650 This work was supported by a DFO grant (MECTS-#3789712) obtained by Philippe  
651 Juneau, Johann Lavaud and Beatrix Beisner and the Natural Science and Engineering  
652 Research Council of Canada (NSERC) (RGPIN-2017-06210) awarded to Philippe  
653 Juneau. The authors thanks also the GRIL (Groupe de recherche interuniversitaire en  
654 limnologie) for funding to PJ and BEB that supported Juan Du's PhD.

655 **References**

- 656 Anu, P.R., Bijoy Nandan, S., Jayachandran, P.R. and Don Xavier, N.D. (2016) Toxicity  
657 effects of copper on the marine diatom, *Chaetoceros calcitrans*. *Regional Studies in*  
658 *Marine Science* 8, 498-504.
- 659 Ardyna, M. and Arrigo, K., R. (2020) Phytoplankton dynamics in a changing Arctic  
660 Ocean. *Nature Climate Change* 10(10), 892-903.
- 661 Asselborn, V., Fernandez, C., Zalocar, Y. and Parodi, E.R. (2015) Effects of chlorpyrifos  
662 on the growth and ultrastructure of green algae, *Ankistrodesmus gracilis*. *Ecotoxicology*  
663 *and Environmental Safety* 120, 334-341.
- 664 Asselborn, V.M., Zalocar, Y. and Parody, E. (2006) Efectos del insecticida  
665 organofosforado clorpirifos sobre el crecimiento y morfología de *Selenastrum*  
666 *capricornutum* Printz (Chlorophyta).
- 667 Bai, X., Sun, C., Xie, J., Song, H., Zhu, Q., Su, Y., Qian, H. and Fu, Z. (2015) Effects  
668 of atrazine on photosynthesis and defense response and the underlying mechanisms in  
669 *Phaeodactylum tricornutum*. *Environmental Science and Pollution Research* 22(17),  
670 499-507.
- 671 Bailleul, B., Berne, N., Murik, O., Petroutsos, D., Prihoda, J., Tanaka, A., Villanova, V.,  
672 Bligny, R., Flori, S., Falconet, D., Krieger-Liszkay, A., Santabarbara, S., Rappaport, F.,  
673 Joliot, P., Tirichine, L., Falkowski, P.G., Cardol, P., Bowler, C. and Finazzi, G. (2015)  
674 Energetic coupling between plastids and mitochondria drives CO<sub>2</sub> assimilation in  
675 diatoms. *Nature* 524(7565), 366-369.
- 676 Balmer, J.E., Morris, A.D., Hung, H., Jantunen, L., Vorkamp, K., Rigét, F., Evans, M.,  
677 Houde, M. and Muir, D.C.G. (2019) Levels and trends of current-use pesticides (CUPs)  
678 in the arctic: An updated review, 2010–2018. *Emerging Contaminants* 5, 70-88.
- 679 Balzano, S., Marie, D., Gourvil, P. and Vaultot, D. (2012) Composition of the summer  
680 photosynthetic pico and nanoplankton communities in the Beaufort Sea assessed by T-  
681 RFLP and sequences of the 18S rRNA gene from flow cytometry sorted samples. *ISME*  
682 *Journal* 6(8), 1480-1498.
- 683 Bayat, L., Arab, M., Aliniaiefard, S., Seif, M., Lastochkina, O. and Li, T. (2018) Effects  
684 of growth under different light spectra on the subsequent high light tolerance in rose  
685 plants. *AoB Plants* 10(5), ply052.
- 686 Cabrerizo, A., Muir, D.C.G., Teixeira, C., Lamoureux, S.F. and Lafreniere, M.J. (2019)  
687 Snow Deposition and Melting as Drivers of Polychlorinated Biphenyls and  
688 Organochlorine Pesticides in Arctic Rivers, Lakes, and Ocean. *Environmental Science*  
689 *Technology* 53(24), 14377-14386.
- 690 CARC (1990) Canadian Arctic Resources Committee, Northern perspectives 18.



691 Chalifour, A., Arts, M.T., Kainz, M.J. and Juneau, P. (2014) Combined effect of  
692 temperature and bleaching herbicides on photosynthesis, pigment and fatty acid  
693 composition of *Chlamydomonas reinhardtii*. European Journal of Phycology 49(4),  
694 508-515.

695 Chalifour, A. and Juneau, P. (2011) Temperature-dependent sensitivity of growth and  
696 photosynthesis of *Scenedesmus obliquus*, *Navicula pelliculosa* and two strains of  
697 *Microcystis aeruginosa* to the herbicide atrazine. Aquatic Toxicology 103(1-2), 9-17.

698 Chalifour, A., LeBlanc, A., Sleno, L. and Juneau, P. (2016) Sensitivity of *Scenedesmus*  
699 *obliquus* and *Microcystis aeruginosa* to atrazine: effects of acclimation and mixed  
700 cultures, and their removal ability. Ecotoxicology 25(10), 1822-1831.

701 Coquille, N., Menard, D., Rouxel, J., Dupraz, V., Eon, M., Pardon, P., Budzinski, H.,  
702 Morin, S., Parlanti, E. and Stachowski-Haberkorn, S. (2018) The influence of natural  
703 dissolved organic matter on herbicide toxicity to marine microalgae is species-  
704 dependent. Aquatic Toxicology 198, 103-117.

705 Croteau, D., Guérin, S., Bruyant, F., Ferland, J., Campbell, D.A., Babin, M. and Lavaud,  
706 J. (2021) Contrasting nonphotochemical quenching patterns under high light and  
707 darkness aligns with light niche occupancy in Arctic diatoms. Limnology and  
708 Oceanography 66(S1).

709 Croteau, D., Lacour, T., Schiffrine, N., Morin, P.I., Forget, M.H., Bruyant, F., Ferland,  
710 J., Lafond, A., Campbell, D.A., Tremblay, J.É., Babin, M. and Lavaud, J. (2022) Shifts  
711 in growth light optima among diatom species support their succession during the spring  
712 bloom in the Arctic. Journal of Ecology.

713 Daufresne, M., Lengfellner, K. and Sommer, U. (2009) Global warming benefits the  
714 small in aquatic ecosystems. Proceedings of the National Academy of Sciences 106(31),  
715 12788-12793.

716 Debenest, T., Silvestre, J., Coste, M. and Pinelli, E. (2010) Effects of pesticides on  
717 freshwater diatoms. Reviews of Environmental Contamination and Toxicology 203, 87-  
718 103.

719 DeLorenzo, M.E. (2001) toxicity of pesticides to aquatic microorganisms: a review.  
720 Environmental Toxicology and Chemistry 20 (1), 84-98.

721 DeLorenzox, M.E., Leatherbury, M., Weiner, J.A., Lewitus, A.J. and Fulton, M.H.  
722 (2004) Physiological factors contributing to the species-specific sensitivity of four  
723 estuarine microalgal species exposed to the herbicide atrazine. Aquatic Ecosystem  
724 Health Management 7(1), 137-146.

725 Dupraz, V., Coquille, N., Menard, D., Sussarellu, R., Haugarreau, L. and Stachowski-  
726 Haberkorn, S. (2016) Microalgal sensitivity varies between a diuron-resistant strain and  
727 two wild strains when exposed to diuron and irgarol, alone and in mixtures.

728 Chemosphere 151, 241-252.

729 Fernandez, C., Asselborn, V. and Parodi, E.R. (2021) Toxic effects of chlorpyrifos,  
730 cypermethrin and glyphosate on the non-target organism *Selenastrum capricornutum*  
731 (Chlorophyta). Anais da Academia Brasileira de Ciencias 93(4).

732 Frey, K., Fansh, L. and Ghhihu, H. (2018) Arctic Ocean Primary Productivity: The  
733 Response of Marine Algae to Climate Warming and Sea Ice Decline. Arctic Report Card.

734 Galindo, V., Gosselin, M., Lavaud, J., Mundy, C.J., Else, B., Ehn, J. and Rysgaard, S.  
735 (2017) Pigment composition and photoprotection of Arctic sea ice algae during spring.  
736 Marine Ecology Progress Series 585, 49-69.

737 Garrido, S., Linares, M., Campillo, J.A. and Albentosa, M. (2019) Effect of  
738 microplastics on the toxicity of chlorpyrifos to the microalgae *Isochrysis galbana*, clone  
739 t-ISO. Ecotoxicology and Environmental Safety 173, 103-109.

740 Gomes, M.P. and Juneau, P. (2017) Temperature and Light Modulation of Herbicide  
741 Toxicity on Algal and Cyanobacterial Physiology. Frontiers in Environmental Science  
742 5.

743 Halsey, K.H. and Jones, B.M. (2015) Phytoplankton strategies for photosynthetic  
744 energy allocation. Annual Review of Marine Science 7, 265-297.

745 Handler, E. (2017) Responses to Light Intensity and Regimes by an Arctic strain of the  
746 picophytoplankton *Micromonas* CCMP2099.

747 Hoferkamp, L., Hermanson, M.H. and Muir, D.C. (2010) Current use pesticides in  
748 Arctic media; 2000-2007. Science of the Total Environment 408(15), 2985-2994.

749 Hoppe, C.J.M., Flintrop, C.M. and Rost, B. (2018) The Arctic picoeukaryote  
750 *Micromonas pusilla* benefits synergistically from warming and ocean acidification.  
751 Biogeosciences 15(14), 4353-4365.

752 Juneau, P., David, D. and Saburo, M. (2001) Evaluation of different algal species  
753 sensitivity to mercury and metolachlor by PAM-fluorometry. Chemosphere 45(4-5),  
754 589-598.

755 Kottuparambil, S., Brown, M.T., Park, J., Choi, S., Lee, H., Choi, H.G., Depuydt, S.  
756 and Han, T. (2017) Comparative assessment of single and joint effects of diuron and  
757 Irgarol 1051 on Arctic and temperate microalgae using chlorophyll a fluorescence  
758 imaging. Ecological Indicators 76, 304-316.

759 Kress, E. and Jahns, P. (2017) The Dynamics of Energy Dissipation and Xanthophyll  
760 Conversion in Arabidopsis Indicate an Indirect Photoprotective Role of Zeaxanthin in  
761 Slowly Inducible and Relaxing Components of Non-photochemical Quenching of  
762 Excitation Energy. Frontiers in Plant Science 8, 2094.

- 763 Lacour, T., Babin, M. and Lavaud, J. (2020) Diversity in xanthophyll cycle pigments  
764 content and related nonphotochemical quenching (NPQ) among microalgae:  
765 implications for growth strategy and ecology. *Journal of Phycology* 56(2), 245-263.
- 766 Lacour, T., Larivière, J. and Babin, M. (2017) Growth, Chl *a* content, photosynthesis,  
767 and elemental composition in polar and temperate microalgae. *Limnology and*  
768 *Oceanography* 62(1), 43-58.
- 769 Lacour, T., Larivière, J., Ferland, J., Bruyant, F., Lavaud, J. and Babin, M. (2018) The  
770 Role of Sustained Photoprotective Non-photochemical Quenching in Low Temperature  
771 and High Light Acclimation in the Bloom-Forming Arctic Diatom *Thalassiosira*  
772 *gravida*. *Frontiers in Marine Science* 5.
- 773 Larras, F., Lambert, A.S., Pesce, S., Rimet, F., Bouchez, A. and Montuelle, B. (2013)  
774 The effect of temperature and a herbicide mixture on freshwater periphytic algae.  
775 *Ecotoxicology and Environmental Safety* 98, 162-170.
- 776 Larsbo, M., Sandin, M., Jarvis, N., Etana, A. and Kreuger, J. (2016) Surface Runoff of  
777 Pesticides from a Clay Loam Field in Sweden. *J Environ Qual* 45(4), 1367-1374.
- 778 Lavaud, J. and Lepetit, B. (2013) An explanation for the inter-species variability of the  
779 photoprotective non-photochemical chlorophyll fluorescence quenching in diatoms.  
780 *Biochimica Biophysica Acta* 1827(3), 294-302.
- 781 Lovejoy, C., Vincent, W.F., Bonilla, S., Roy, S., Martineau, M.J., Terrado, R., Potvin,  
782 M., Massana, R. and Pedrós-Alió, C. (2007) Distribution, Phylogeny, and Growth of  
783 Cold-Adapted Picoprasinophytes in Arctic Seas. *Journal of Phycology* 43(1), 78-89.
- 784 Lyon, B.R. and Mock, T. (2014) Polar Microalgae: New Approaches towards  
785 Understanding Adaptations to an Extreme and Changing Environment. *Biology (Basel)*  
786 3(1), 56-80.
- 787 Ma, Y., Adelman, D.A., Bauerfeind, E., Cabrerizo, A., McDonough, C.A., Muir, D.,  
788 Soltwedel, T., Sun, C., Wagner, C.C., Sunderland, E.M. and Lohmann, R. (2018)  
789 Concentrations and Water Mass Transport of Legacy POPs in the Arctic Ocean.  
790 *Geophysical Research Letters* 45(23).
- 791 Margesin, R. (2007) Alpine microorganisms: useful tools for low-temperature  
792 bioremediation *Journal of Microbiology* 45(4), 281-285.
- 793 McKie-Krisberg, Z.M. and Sanders, R.W. (2014) Phagotrophy by the picoeukaryotic  
794 green alga *Micromonas*: implications for Arctic Oceans. *ISME Journal* 8(10), 1953-  
795 1961.
- 796 Medithi, S., Jonnalagadda, P.R. and Jee, B. (2021) Predominant role of antioxidants in  
797 ameliorating the oxidative stress induced by pesticides. *Archives of Environmental &*  
798 *Occupational Health* 76(2), 61-74.

799 Morgan-Kiss, R.M., Priscu, J.C., Pocock, T., Gudynaite-Savitch, L. and Huner, N.P.  
800 (2006) Adaptation and acclimation of photosynthetic microorganisms to permanently  
801 cold environments. *Microbiology and Molecular Biology Reviews* 70(1), 222-252.

802 Moustakas, M., Moustaka, J. and Sperdouli, I. (2022) Hormesis in photosystem II: a  
803 mechanistic understanding. *Current Opinion in Toxicology*.

804 Muir, D., Kurt-Karakus, P. and Stow, J. (2013) Canadian Arctic Contaminants  
805 Assessment Report On Persistent Organic Pollutants. NCP (Northern Contaminants  
806 Program).

807 Muir, D.C. and de Wit, C.A. (2010) Trends of legacy and new persistent organic  
808 pollutants in the circumpolar arctic: overview, conclusions, and recommendations.  
809 *Science of the Total Environment* 408(15), 3044-3051.

810 Müller, P., Li, X.P. and Niyogi, K.K. (2001) Non-photochemical quenching. A response  
811 to excess light energy *Plant physiology* 125(4), 1558-1566.

812 Ni, G., Zimbalatti, G., Murphy, C.D., Barnett, A.B., Arsenault, C.M., Li, G., Cockshutt,  
813 A.M. and Campbell, D.A. (2017) Arctic *Micromonas* uses protein pools and non-  
814 photochemical quenching to cope with temperature restrictions on Photosystem II  
815 protein turnover. *Photosynthesis Research* 131(2), 203-220.

816 Parker, M.S. and Armbrust, E.V. (2005) Synergistic Effects of Light, Temperature, and  
817 Nitrogen Source on Transcription of Genes for Carbon and Nitrogen Metabolism in the  
818 Centric Diatom *Thalassiosira Pseudonana* (Bacillariophyceae)1. *Journal of Phycology*  
819 41(6), 1142-1153.

820 Petrou, K., Hill, R., Brown, C.M., Campbell, D.A., Doblin, M.A. and Ralph, P.J. (2010)  
821 Rapid photoprotection in sea-ice diatoms from the East Antarctic pack ice. *Limnology  
822 and Oceanography* 55(3), 1400-1407.

823 Pućko, M., Stern, G.A., Burt, A.E., Jantunen, L.M., Bidleman, T.F., Macdonald, R.W.,  
824 Barber, D.G., Geilfus, N.X. and Rysgaard, S. (2017) Current use pesticide and legacy  
825 organochlorine pesticide dynamics at the ocean-sea ice-atmosphere interface in resolute  
826 passage, Canadian Arctic, during winter-summer transition. *Science of the Total  
827 Environment* 580, 1460-1469.

828 Rioboo, C., González, O., Herrero, C. and Cid, A. (2002) Physiological response of  
829 freshwater microalga (*Chlorella vulgaris*) to triazine and phenylurea herbicides.  
830 *Aquatic Toxicology* 59(3-4), 225-235.

831 Ronka, S. (2016) Removal of triazine-based herbicides on specific polymeric sorbent:  
832 batch studies. *Pure and Applied Chemistry* 88(12), 1167-1177.


833 Serôdio, J. and Lavaud, J. (2020) Diatoms and their ecological importance. *Life Below  
834 Water*, 1-9.

- 835 Singh, Z., Jasminder, K. and Ravneet, K. (2016) Toxic Effects of Organochlorine  
836 Pesticides: A Review. *American Journal of BioScience* 4(3).
- 837 Stachowski-Haberkorn, S., Jerome, M., Rouxel, J., Khelifi, C., Rince, M. and Burgeot,  
838 T. (2013) Multigenerational exposure of the microalga *Tetraselmis suecica* to diuron  
839 leads to spontaneous long-term strain adaptation. *Aquatic Toxicology* 140-141, 380-  
840 388.
- 841 Strzeppek, R.F. and Harrison, P.J. (2004) Photosynthetic architecture differs in coastal  
842 and oceanic diatoms. *Nature* 431(7009), 684-689.
- 843 Takishita, Y., Souleimanov, A., Bourguet, C., Ohlund, L.B., Arnold, A., A., Sleno, L.  
844 and Smith, D.L. (2021) *Pseudomonas entomophila* 23S Produces a Novel Antagonistic  
845 Compound against *Clavibacter michiganensis* subsp. *michiganensis*, a Pathogen of  
846 Tomato Bacterial Canker. *Applied Microbiology* 1(1), 60-73.
- 847 Tang, J., Hoagland, K.D. and Siegfried, B.D. (1998) Uptake and bioconcentration of  
848 atrazine by selected freshwater algae. *Environmental Toxicology and Chemistry* 17(6),  
849 1085-1090.
- 850 Tuchman, N.C., Schollett, M.A., Rier, S.T. and Geddes, P. (2006) Differential  
851 Heterotrophic Utilization of Organic Compounds by Diatoms and Bacteria under Light  
852 and Dark Conditions. *Hydrobiologia* 561(1), 167-177.
- 853 Vonk, J.A. and Kraak, M.H.S. (2020) Herbicide Exposure and Toxicity to Aquatic  
854 Primary Producers. *Reviews of Environmental Contamination and Toxicology* 250,  
855 119-171.
- 856 Vorkamp, K. and Riget, F.F. (2014) A review of new and current-use contaminants in  
857 the Arctic environment: evidence of long-range transport and indications of  
858 bioaccumulation. *Chemosphere* 111, 379-395.
- 859 Weber, J., Halsall, C.J., Muir, D., Teixeira, C., Small, J., Solomon, K., Hermanson, M.,  
860 Hung, H. and Bidleman, T. (2010) Endosulfan, a global pesticide: a review of its fate  
861 in the environment and occurrence in the Arctic. *Science of the Total Environment*  
862 408(15), 2966-2984.
- 863 Weiner, J.A., DeLorenzo, M.E. and Fulton, M.H. (2004) Relationship between uptake  
864 capacity and differential toxicity of the herbicide atrazine in selected microalgal species.  
865 *Aquatic Toxicology* 68(2), 121-128.
- 866 White, P.L., Wynn-Williams, D.D. and Russell, N.J. (2000) Diversity of thermal  
867 responses of lipid composition in the membranes of the dominant culturable members  
868 of an Antarctic fellfield soil bacterial community *Antarctic Science* 12(3), 386-393.
- 869 Wiebe, W.J., Sheldon, J.W.M. and Pomeroy, L.R. (1992) Bacterial growth in the cold:  
870 evidence for an enhanced substrate requirement *Applied and Environmental*

- 871 Microbiology 58(1), 359-364.
- 872 Wirth, C., Limberger, R. and Weisse, T. (2019) Temperature x light interaction and  
873 tolerance of high water temperature in the planktonic freshwater flagellates  
874 *Cryptomonas* (Cryptophyceae) and *Dinobryon* (Chrysophyceae). *Journal of Phycology*  
875 55(2), 404-414.
- 876 Yadav, N.R. (2015) Toxic effect of chlorpyrifos and dimethoate on protein and  
877 chlorophyll-a content of *spirulina platensis*. *International journal of engineering science*  
878 & advanced research 1, 24-26.
- 879 Young, J.N., Goldman, J.A., Kranz, S.A., Tortell, P.D. and Morel, F.M. (2015) Slow  
880 carboxylation of Rubisco constrains the rate of carbon fixation during Antarctic  
881 phytoplankton blooms. *New Phytologist* 205(1), 172-181.
- 882 Zhang, X., Luo, Y. and Goh, K.S. (2018) Modeling spray drift and runoff-related inputs  
883 of pesticides to receiving water. *Environmental Pollution* 234, 48-58.
- 884


885 **Supplementary Material**

886 Figure S1. The effects of chlorpyrifos and trifluralin on energy fluxes within PSII of  
 887 four species, including temperate *C. neogracile* (T-CN), Arctic *C. neogracile* (A-CN)  
 888 for A and B, temperate *M. bravo* (T-MB) and Arctic *M. polaris* (A-MP) for C and D  
 889 after 72 h exposure. Data expressed as means  $\pm$  SD (n = 6).




890

891 Figure S2. Fast fluorescence kinetics of T-CN (A), A-CN (B), T-MB (C) and A-MP (D),  
 892 exposed to chlorpyrifos and trifluralin for 72h.



893

894 Figure S3. The effects of atrazine and simazine after 72 h exposure on the cell  
 895 biovolume. Data are means  $\pm$  SD. for 72h.



896



Table S1. Definitions for the fluorescence parameters used in this study.

<b>Parameters</b>	<b>Definition</b>
$\Phi_M$	Maximal PSII quantum yield
$\Phi'_M$	Operational PSII quantum yield
NPQ	Non-photochemical quenching
NPQ <sub>max</sub>	Maximum ability for dissipation of excess energy
rETR <sub>max</sub>	Maximum relative photosynthetic electron transport rate
$\alpha$	Maximum light efficiency use
$E_K$	Light saturation coefficient
<b>Specific energy fluxes (per <math>Q_A</math> reducing PSII RC)</b>	
ABS/RC	Absorption flux (of antenna Chls) per RC (also a measure of PSII apparent antenna size)
TRo/RC	Trapped energy flux (leading to $Q_A$ reduction) per RC
ETo/RC	Electron transport flux (further than $Q_A^-$ ) per RC
DIo/RC	Dissipated energy flux per RC
<b>Performance index</b>	
$PI_{ABS}=(RC/ABS)$	Performance index (potential) for energy conservation from photons absorbed by PSII to the reduction of intersystem electron acceptors

899 Table S2. The effects of trifluralin and chlorpyrifos after 72 h exposure on the growth  
 900 rate, cell volume,  $\Phi_M$ ,  $\Phi'_M$ , ROS content, and Chl *a* pigment. The percentages of the  
 901 control value are shown by the numbers in parentheses. Data are means  $\pm$  SD.

Parameter	Pesticide concentration ( $\mu\text{g/l}$ )	T-CN	A-CN	T-MB	A-MP	
Growth rate	Trifluralin	0	0.622 $\pm$ 0.01 (100)	0.655 $\pm$ 0.01 (100)	0.375 $\pm$ 0.01 (100)	0.245 $\pm$ 0.02 (100)
		200	0.201 $\pm$ 0.04 (28.5)	0.533 $\pm$ 0.01 (69.3)	0.049 $\pm$ 0.04 (37.6)	0.027 $\pm$ 0.02 (52.0)
	Chlopyrifos	500	0.081 $\pm$ 0.08 (22.2)	0.346 $\pm$ 0.15 (60.4)	-0.093 $\pm$ 0.08 (28.2)	-0.195 $\pm$ 0.28 (46.4)
		200	0.608 $\pm$ 0.00 (96.0)	0.523 $\pm$ 0.01 (67.4)	0.382 $\pm$ 0.02 (102)	-0.083 $\pm$ 0.02 (37.4)
		500	0.391 $\pm$ 0.01 (50.1)	0.102 $\pm$ 0.03 (19.1)	0.326 $\pm$ 0.05 (86.9)	-0.098 $\pm$ 0.01 (35.8)
	Volume ( $\mu\text{m}^3$ )	Trifluralin	0	144.4 $\pm$ 0.95 (100)	45.8 $\pm$ 2.89 (100)	9.48 $\pm$ 0.20 (100)
200			218.9 $\pm$ 2.51 (151)	53.1 $\pm$ 1.28 (116)	8.43 $\pm$ 0.77 (89.4)	7.62 $\pm$ 0.08 (119)
Chlopyrifos		500	221.8 $\pm$ 20.9 (154)	58.8 $\pm$ 1.66 (129)	8.84 $\pm$ 1.02 (93.3)	8.69 $\pm$ 0.47 (136)
		200	140.7 $\pm$ 1.00 (97.2)	51.2 $\pm$ 0.61 (115)	10.67 $\pm$ 0.04 (112)	12.66 $\pm$ 2.18 (199)
		500	143.8 $\pm$ 5.32 (99.3)	91.3 $\pm$ 9.95 (199)	10.49 $\pm$ 0.21 (111)	12.85 $\pm$ 0.08 (200)
Fv/Fm		Trifluralin	0	0.654 $\pm$ 0.008 (100)	0.624 $\pm$ 0.025 (100)	0.660 $\pm$ 0.002 (100)
	200		0.574 $\pm$ 0.004 (87.8)	0.591 $\pm$ 0.011 (94.8)	0.492 $\pm$ 0.057 (74.5)	0.538 $\pm$ 0.012 (89.1)
	Chlopyrifos	500	0.524 $\pm$ 0.006 (80.2)	0.536 $\pm$ 0.003 (86.0)	0.462 $\pm$ 0.052 (70.0)	0.489 $\pm$ 0.010 (80.9)
		200	0.638 $\pm$ 0.004 (97.6)	0.548 $\pm$ 0.006 (87.8)	0.653 $\pm$ 0.001 (99.0)	0.538 $\pm$ 0.005 (89.0)
		500	0.558 $\pm$ 0.007 (85.4)	0.344 $\pm$ 0.045 (50.8)	0.645 $\pm$ 0.006 (97.9)	0.512 $\pm$ 0.013 (84.7)
	$\Phi'_m$	Trifluralin	0	0.657 $\pm$ 0.003 (100)	0.610 $\pm$ 0.025 (100)	0.555 $\pm$ 0.011 (100)
200			0.552 $\pm$ 0.011 (84.0)	0.579 $\pm$ 0.011 (95.1)	0.356 $\pm$ 0.027 (64.1)	0.406 $\pm$ 0.021 (82.8)
Chlopyrifos		500	0.483 $\pm$ 0.009 (73.5)	0.507 $\pm$ 0.003 (83.3)	0.332 $\pm$ 0.024 (60.0)	0.357 $\pm$ 0.016 (72.9)
		200	0.636 $\pm$ 0.002 (96.7)	0.477 $\pm$ 0.006 (78.4)	0.551 $\pm$ 0.007 (99.3)	0.379 $\pm$ 0.013 (77.2)
		500	0.365 $\pm$ 0.016 (55.6)	0.243 $\pm$ 0.045 (38.9)	0.533 $\pm$ 0.011 (96.0)	0.341 $\pm$ 0.007 (69.4)
ROS content		Trifluralin	0	0.917 $\pm$ 0.236 (100)	0.709 $\pm$ 0.074 (100)	0.128 $\pm$ 0.040 (100)
	200		1.07 $\pm$ 0.052 (117)	0.870 $\pm$ 0.064 (123)	0.508 $\pm$ 0.030 (397)	0.947 $\pm$ 0.197 (393)
	Chlopyrifos	500	1.20 $\pm$ 0.203 (131)	3.46 $\pm$ 0.985 (488)	0.689 $\pm$ 0.125 (539)	0.915 $\pm$ 0.153 (379)
		200	2.16 $\pm$ 0.205 (236)	0.975 $\pm$ 0.120 (138)	1.72 $\pm$ 0.215 (1344)	0.662 $\pm$ 0.151 (274)
		500	2.80 $\pm$ 0.112 (306)	1.02 $\pm$ 0.114 (143)	2.81 $\pm$ 0.581 (2195)	0.492 $\pm$ 0.153 (204)
	Chla ( $10^6$ cells $\mu\text{g/ml}$ )	Trifluralin	0	3.818 $\pm$ 0.168 (100)	1.08 $\pm$ 0.060 (100)	0.540 $\pm$ 0.005 (100)
200			12.01 $\pm$ 1.824 (330)	1.21 $\pm$ 0.039 (109)	0.315 $\pm$ 0.008 (58.0)	0.341 $\pm$ 0.018 (110)
Chlopyrifos		500	6.752 $\pm$ 0.468 (145)	1.12 $\pm$ 0.059 (104)	0.745 $\pm$ 0.043 (137)	0.306 $\pm$ 0.097 (94.1)
		200	1.16 $\pm$ 0.053 (30.3)	1.04 $\pm$ 0.027 (96.3)	0.497 $\pm$ 0.059 (91.4)	0.390 $\pm$ 0.036 (124)
		500	1.20 $\pm$ 0.267 (31.8)	1.75 $\pm$ 0.038 (157)	0.424 $\pm$ 0.025 (80.0)	0.388 $\pm$ 0.012 (123)
CAT ( $\text{U mg}^{-1}\text{min protein}^{-1}$ )		Trifluralin	0	0.090 $\pm$ 0.001 (100)	0.134 $\pm$ 0.003 (100)	0.145 $\pm$ 0.003 (100)
	200		0.101 $\pm$ 0.004 (105)	0.185 $\pm$ 0.004 (138)	0.148 $\pm$ 0.002 (102)	0.353 $\pm$ 0.014 (161)
	Chlopyrifos	500	6.752 $\pm$ 0.003 (113)	0.290 $\pm$ 0.002 (217)	0.281 $\pm$ 0.003 (193)	0.400 $\pm$ 0.087 (182)
		200	0.087 $\pm$ 0.001 (96.8)	0.131 $\pm$ 0.002 (97.6)	0.112 $\pm$ 0.059 (76.6)	0.311 $\pm$ 0.006 (119)
		500	0.057 $\pm$ 0.002 (63.1)	1.089 $\pm$ 0.038 (812)	0.163 $\pm$ 0.025 (112)	0.264 $\pm$ 0.032 (101)