



HAL
open science

Laboratory experiments for measuring kinetic parameters

Max R McGillen

► **To cite this version:**

Max R McGillen. Laboratory experiments for measuring kinetic parameters. Doctoral. France. 2022. hal-03852541

HAL Id: hal-03852541

<https://hal.science/hal-03852541>

Submitted on 15 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Laboratory experiments for measuring kinetic parameters



Dr. Max McGillen

Institut de Combustion, Aérothermique,
Réactivité et Environnement, CNRS-Orléans,
France



Lecture outline

- Why should we care about chemical kinetics?
- Lifetime in the atmosphere
- The nature of the rate coefficient, k
 - What is it?
 - The gas-kinetic limit
 - The Arrhenius equation
- Methods for measuring k
- Dependence of k on T and P
- The problem with parameter-space


Kinetic parameters of the atmosphere

Carte du jour

- Photolysis:
 - $O_3 + hv \rightarrow O(^1D) + O_2$
- Thermal decomposition:
 - $CH_3C(O)OONO_2 \rightarrow CH_3C(O)OO + NO_2$
- Radical–molecule:
 - $OH + \text{isoprene} \rightarrow \text{products}$
- Radical–radical:
 - $RO_2 + RO_2 \rightarrow RO + RO + O_2$
- Termolecular reactions:
 - $OH + SO_2 + M \rightarrow H_2SO_4 + M$
- Equilibria:
 - $NO_2 + NO_3 \rightleftharpoons N_2O_5$
- Phase transfer:
 - Gas-phase \rightarrow aerosol-phase
- Heterogeneous reactions:
 - $2NO_2 + H_2O + \text{surface} \rightarrow HONO + HNO_3$
- ... + more




Why we study chemical kinetics

- So, the atmosphere has lots of different chemical reactions that can happen
- Many of these reactions are in competition with each other
- Normally, the faster the rate of reaction, the more important it is
- That is, the more competitive it is



Types of competition


- For a simple compound like methane, we could ask:
which oxidant is most important?
- For more complex compounds, we could ask:
which reactive sites are more important?
- For compounds of a difference size, we could ask:
which mechanism is most important?



Atmospheric lifetime

- Atmospheric chemists often use the lifetime with respect to OH as a general measure of lifetime:

$$\tau = \frac{1}{k[\text{OH}]}$$



Why is the lifetime so important?

- Lifetime is a major control on the distance over which pollution can be transported
- The pollution from one country could affect the air quality of another
- Furthermore, for certain chemicals, vertical transport (and persistence) becomes important (e.g. CFCs, HCFCs)

For example:

$$\text{ODP}_i = \frac{n_{\text{Cl}}}{3} \frac{f_i}{f_{\text{CFC-11}}} \frac{\tau_i}{\tau_{\text{CFC-11}}} \frac{M_{\text{CFC-11}}}{M_i}$$

For example:

$$\text{AGWP}_i(H) = A_i \tau \left(1 - \exp\left(-\frac{H}{\tau}\right) \right).$$

Why is the lifetime so important?


stratosphere

ozone layer

Very slow transport to the stratosphere

troposphere

Transport to neighbouring cities / countries



The total atmospheric lifetime

- Reaction with OH is generally important for destroying pollution
- But there are many other processes that can be important
- Therefore, it is more useful to consider the lifetime as follows:

$$\frac{1}{\tau_{\text{total}}} = \frac{1}{\tau_{\text{OH}}} + \frac{1}{\tau_{\text{NO}_3}} + \frac{1}{\tau_{\text{O}_3}} + \frac{1}{\tau_{\text{Cl}}} + \frac{1}{\tau_{\text{photolysis}}} + \frac{1}{\tau_{\text{hydrolysis}}} + \dots$$

- Also, depending on the location in the atmosphere, individual processes may dominate, e.g. photolysis is faster in the stratosphere, but hydrolysis is very slow, so we can express lifetime as follows:

$$\frac{1}{\tau_{\text{total}}} = \frac{1}{\tau_{\text{troposphere}}} + \frac{1}{\tau_{\text{stratosphere}}} + \frac{1}{\tau_{\text{mesosphere}}}$$

Either way, it all depends on k (or J)

$$\tau = \frac{1}{k[\text{OH}]} \quad \tau_{\text{O}_3} = \frac{1}{k[\text{O}_3]} \quad \textit{etcetera ...}$$

$$\tau_{\text{oxidant}} = \frac{1}{k_{\text{oxidant}}[\text{oxidant}]}$$

$$\textit{similarly ...} \quad \tau_{\text{photolysis}} = \frac{1}{J}$$

The rate coefficient, k

- Simple atmospheric chemical reactions are first-order, $A \rightarrow B$ (e.g. unimolecular, photolysis):

$$-\frac{d[A]}{dt} = k[A] \longrightarrow \ln \frac{[A]_0}{[A]_t} = kt$$

Relative concentration change with time (convenient)

- Many atmospheric reactions are bi-molecular and second-order, $A + B \rightarrow C$ (e.g. OH addition):

$$-\frac{d[A]}{dt} = k[A][B] \longrightarrow \frac{1}{[A]_0 - [B]_0} \ln \frac{[A]_t[B]_0}{[A]_0[B]_t} = kt$$

Absolute concentration change with time (inconvenient)

- Kinetics prefer to work under pseudo-first-order conditions, where:

$$[B] \gg [A] \therefore [B]_0 \approx [B]_t$$

$$-\frac{d[A]}{dt} = k_{obs}[A] \longrightarrow k_{obs} = k \times [B]$$

Changes in concentration and time...

The rate coefficient, k

- But what is k ?
- k is a parameter that describes efficiency
- If k is big, the reaction process is efficient
- If k is small, the reaction process is inefficient
- If k is 100% efficient, then every collision between molecule A and molecule B results in a reaction
- This is known as the “gas kinetic limit” or “collision limit”

The gas kinetic limit

- Begin by considering how fast a gas molecule can move
- Let's consider something simple like N₂ at room temperature:

$$\bar{v} = \sqrt{\frac{8k_B T}{\pi m}} = \sqrt{\frac{8 \times 1.38 \times 10^{-23} (m^2 kg s^{-2} K^{-1}) \times 298 (K)}{3.14 \dots \times 4.65 \times 10^{-26} (kg)}} = 474.7 m s^{-1}$$

- Next, consider how far a molecule travels before it hits another molecule (mean free path):

$$\lambda = \frac{RT}{\sqrt{2}\pi d^2 N_A P} = \frac{8.314 (m^3 \cdot Pa \cdot K^{-1} \cdot mol^{-1}) \times 298 (K)}{\sqrt{2}\pi (3.64 \times 10^{-10})^2 (m^2) \times 6.02 \times 10^{23} (1 mol) \times 101325 (Pa)} = 6.898 \times 10^{-8} m$$


- Now we can calculate how many collisions are happening:

$$Z = \frac{\bar{v}}{\lambda} = \frac{474.7 (m s^{-1})}{6.898 \times 10^{-8} (m)} = 6.88 \times 10^9 s^{-1}$$

- And, if you divide by the number density of 1 atmosphere of gas at room temperature:

$$k = \frac{Z}{n} = \frac{6.88 \times 10^9 (s^{-1})}{2.46 \times 10^{19} (molecule cm^{-3})} = 2.79 \times 10^{-10} cm^3 molecule^{-1} s^{-1}$$

What controls the **size** of k ?



- Some reactions are highly efficient, i.e. \approx gas kinetic limit (e.g. reactions with chlorine and alkenes, $k = 10^{-10}$)
- Others are a lot less efficient (e.g. alkene ozonolysis, $k = 10^{-22}$)
- So, some reactions are a trillion times more efficient than others!
- Where does this inefficiency come from?
- There are several possibilities...

The Arrhenius equation

- Many reactions have an activation energy
- This imparts a temperature dependence:


$$k(T) = Z \exp(-E_a/RT)$$

- Other (steric) factors are also important:


$$k(T) = Z\rho \exp(-E_a/RT)$$

- Leading to the familiar Arrhenius equation:

$$k(T) = A \exp(-E_a/RT)$$



Svante Arrhenius
(1889)



How do we measure kinetic parameters?

Let's consider a bi-molecular reaction ($A + B \rightarrow C$), we generally need three different things:

A. An environment where A and B can react together (a reactor)

- Appropriate temperature/ pressure, absence of secondary chemistry


B. An analytical tool to measure (at least one of) A, B or C

- Needs to be sensitive enough, and rapid enough $-\frac{d[A]}{dt} = k_{obs}[A]$

C. A measurement of time



- A way of exposing A to B for a known amount of time $-\frac{d[A]}{dt} = k_{obs}[A]$

Laser Induced Fluorescence




- Temperature range: 220–373 K
- Pressure range: 30–600 Torr
- Bath gases: He, Ar, N₂, O₂
- OH precursors: H₂O₂, HNO₃
- On-line measurement of [SO₂]
- Supporting RR measurements (N₂, Ar, O₂, Air)


Laser Induced Fluorescence



$$\ln \left(\frac{[\text{OH}]_t}{[\text{OH}]_0} \right) = \ln \left(\frac{S_t}{S_0} \right) = -(k[\text{reactant}] + k_d)t = -k't$$



Laser photolysis coupled to CRDS






$$\tau_0 = \frac{L}{c(1-R)}$$

$$\tau_{abs} = \frac{L}{c\{(1-R) + [abs]\sigma_{abs}L\}}$$

$$[abs]\sigma_{abs} = \frac{1}{c} \left(\frac{1}{\tau_{abs}} - \frac{1}{\tau_0} \right)$$

Convenient chemistry




355 nm probe light

T range:
P range:
Ring-down
Reaction


$[\text{Cl}]_0: \sim 2 \times 10^{12} \text{ (molec. cm}^{-3}\text{)}$

nd
gas



mirror with N₂ purge
thermocouple
pressure gauge

heater




Fit function: $\frac{dN}{dt} = -2k_{obs}N^2 - k_pN$

$$N(t) = \frac{k_2 N(t_0)}{k_p e^{k_p t} - 2k_{obs} N(t_0) + 2k_{obs} N(t_0) e^{k_p t}}$$


$$\Delta\kappa(t) = \frac{k_p}{\frac{k_p}{\Delta\kappa(t_0)} e^{k_p t} - (k') \left(\frac{2L}{cd}\right) + (k') \left(\frac{2L}{cd}\right) e^{k_p t}}$$

Relative rate kinetics



- Another approach is to use the relative rate approach
- Here we introduce two compounds that are allowed to react simultaneously:
$$A + OH \rightarrow \text{product}$$
$$B + OH \rightarrow \text{product}$$
- So, if we know the reactivity of B (for example), then we can deduce that of A

Relative rate kinetics



- This is done with the relative rate approach:



$$\ln\left(\frac{[A]_0}{[A]_t}\right) - k_d t = \frac{k_A}{k_B} \ln\left(\frac{[B]_0}{[B]_t}\right) - k_d t$$

- This can be very precise and effective, but it is limited by the accuracy of the reference




T-dependence of k

- Many reactions are temperature dependent
- And over a small temperature range, they may obey the **Arrhenius equation**
- But over larger ranges, we observe curvature
- And this curvature can be described by the **modified Arrhenius equation**
- This is referred to as non-Arrhenius behaviour




Non-Arrhenius behaviour




- Several reasons for this:
- Multiple (*i*) reactions sites, each with their own different T-dependence
- Since the total reaction rate is the sum of each:

$$k_{\text{total}}(T) = \sum_i \kappa_i A_i e^{-E_i/RT}$$



Non-Arrhenius behaviour

- A typical atmospheric emission such as *cis*-3-hexenol possesses **several reaction sites**
- These sites react through **several mechanisms**, which respond differently to temperature changes
- Some sites have a positive T-dep, some sites have a negative T-dep, some sites have positive and negative T-dep
- Wait... negative T-dep?



Negative temperature-dependence of k

- Remember what I said earlier?
- Well, it is true, but it isn't the whole truth...



The Arrhenius equation

- Many reactions have an activation energy
- This emparts a temperature dependence:


$$k(T) = Z \exp(-E_a/RT)$$

- Other (steric) factors are also important:


$$k(T) = Zp \exp(-E_a/RT)$$


- Leading to the familiar Arrhenius equation:

$$k(T) = A \exp(-E_a/RT)$$









Svante Arrhenius
(1889)






OH + oxygenate




Negative temperature-dependence of k

- We see negative temperature dependence for many atmospheric reactions
- Atmospheric chemistry transforms reduced molecules into oxygenated molecules
- This is an opportunity for **van der Waals complexes** to form
- In fact, non-Arrhenius behaviour is not exceptional, it is common




Almost all atmospheric degradation products are oxygenated

Non-Arrhenius behaviour



Zhou *et al.* (2011) identified 8 reaction channels in a reaction that possesses 5 reaction sites

McGillen *et al.* (2013) used 3 reaction channels to describe a reaction that possesses 2 reaction sites




Pressure-dependence of k

- There are several very important reactions in the atmosphere that exhibit pressure dependence
- Pressure dependence in atmospheric reactions (mostly) involves energy transfer with the N_2 and O_2
- These are referred to as M, a third body that doesn't participate direction in a reaction:

For example: $SO_2 + OH \rightleftharpoons [HOSO_2]^* + M \rightarrow HOSO_2 + M \rightarrow \rightarrow H_2SO_4$



- This is a competition: can $[HOSO_2]^*$ collide with M before it falls apart?
- Therefore, this reaction depends on $[M]$ (\propto pressure)

kinetics in the fall-off regime




- Where atmospheric pressure dependence is important, reactions are usually in the fall-off region
- At some point, k cannot get any larger, it has reached its high-pressure limit (k_∞)
- The onset of k_∞ can vary widely <mBar – 100s of Bar
- This depends of the stability of the intermediate

P- and T-dependence in OH + SO₂




Pressure dependence in ozonolysis



- Ozonolysis is a unique chemical system with a large number of local minima on the PES
- There is a cascade of intermediate species, which are losing energy by colliding with M
- It becomes a race to the bottom of the potential energy surface

Estimating k

- I have talked a lot about measuring k so far
- But measuring k is often very difficult and time-consuming
- Yet there are many reactions rates that we would like to know
- This is generally done by parameterizing aspects of molecules that control reactivity
- These are the structure-activity relationships (SARs)




Structure-activity relationships (SARs)

- There are many types of SARs that can be used
- The most popular approach is the group-additivity method
- This method makes use of the idea that the observed k is the summation of all reaction channels:

$$k_{\text{total}}(T) = \sum_i \kappa_i A_i e^{-E_i/RT}$$

- This has the advantage of providing estimates for product branching ratios
- This is ideal for chemical models, where we want to know the lifetime of a chemical, but also the secondary pollutants that are formed



$$\begin{aligned}
 k_{\text{total}} = & \{ 2k_{\text{sec}} F(-\text{CH}_2-) F(>\text{CH-}) F_5 F_6 \\
 & + k_{\text{tert}} [F(-\text{CH}_2-)]^2 F(>\text{CH-}) F_5 F_5 F_6 \\
 & + k_{\text{tert}} [F(-\text{CH}_2-)]^2 F(>\text{C}<) F_5 F_5 F_6 \\
 & + k_{\text{sec}} [F(>\text{CH-})]^2 F_5 F_5 \\
 & + k_{\text{tert}} F(>\text{C}<) F(>\text{CH-}) F(-\text{CH}_3) F_5 F_6 \\
 & + 2k_{\text{prim}} F(>\text{C}<) + k_{\text{prim}} F(>\text{CH-}) \}
 \end{aligned}$$


Example calculation from: Kwok and Atkinson (1995)

Structure-activity relationship principles

- Fundamentally, these are parameterizations, so, regardless of the approach, the idea is to describe the process of interest in a simplified way
- For SARs, this is linked to the molecular structure
- But the molecular property that is chosen for parameterization will change depending on the reaction mechanism
- For example, for the hydrogen abstraction reaction: $\text{RH} + \text{OH} \rightarrow \text{H}_2\text{O} + \text{R}$, a description of the strength of the R–H bond may be best
- On the other hand, for OH addition reactions:
 $\text{R}_1(\text{R}_2)\text{C}=\text{C}(\text{R}_3)\text{R}_4 + \text{OH} \rightarrow \text{R}_1(\text{R}_2)(\text{OH})\text{C}=\text{C}(\text{R}_3)\text{R}_4$, a description of the electron density of the C=C bond may be the best approach

Structure-activity relationship performance

- High quality measurements may determine k within $\sim 5\%$
- SARs tend to be accurate to within a factor of 2
- There is no replacement for high-quality laboratory data, but models need to contain reasonable information



Source: Carter (2020)

Databases to find kinetic data

NIST Chemical Kinetics Database:

<https://kinetics.nist.gov/kinetics/>

IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation:

<https://iupac-aeris.ipsl.fr/>

NASA/ JPL data evaluation:


<https://jpldataeval.jpl.nasa.gov/>

Database for the kinetics of the gas-phase atmospheric reactions of organic compounds:

<https://data.eurochamp.org/data-access/kin/#/home>

Further reading

- This was a really quick introduction to the science of kinetics, if you would like to learn more about it, consider the following book:



Conclusions

- Kinetics has important effects in all aspects of atmospheric chemistry
- Kinetics is a fundamental quantity that can be measured with a variety of analytical and experimental techniques
- This quantity often varies according to the atmospheric conditions (temperature and pressure effects)
- Atmospheric chemistry produces a huge variety of chemical products
- This leads to an enormous parameter-space (too many to measure)
- So, we also need to think how we can estimate those things that we don't have time to measure