



**HAL**  
open science

## New perspectives on the regulation of germinal center reaction via $\alpha\nu\beta 8$ - mediated activation of $TGF\beta$

Sébastien This, H. Paidassi

► **To cite this version:**

Sébastien This, H. Paidassi. New perspectives on the regulation of germinal center reaction via  $\alpha\nu\beta 8$ -mediated activation of  $TGF\beta$ . *Frontiers in Immunology*, 2022, 13, 10.3389/fimmu.2022.942468 . hal-03852949


**HAL Id: hal-03852949**

**<https://hal.science/hal-03852949>**

Submitted on 15 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## OPEN ACCESS

## EDITED BY

Rachel Maurie Gerstein,  
University of Massachusetts Medical  
School, United States

## REVIEWED BY

Eun Jeong Park,  
Mie University, Japan  
Jason G. Cyster,  
University of California, San Francisco,  
United States  
Marita Chakhtoura,  
Independent Researcher, NJ,  
Sicklerville, United States

## \*CORRESPONDENCE

Helena Paidassi  
helena.paidassi@inserm.fr

## SPECIALTY SECTION

This article was submitted to  
B Cell Biology,  
a section of the journal  
Frontiers in Immunology

RECEIVED 12 May 2022

ACCEPTED 25 July 2022

PUBLISHED 22 August 2022

## CITATION

This S and Paidassi H (2022) New perspectives on the regulation of germinal center reaction *via*  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$ . *Front. Immunol.* 13:942468. doi: 10.3389/fimmu.2022.942468

## COPYRIGHT

© 2022 This and Paidassi. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# New perspectives on the regulation of germinal center reaction *via* $\alpha_v\beta_8$ -mediated activation of TGF $\beta$

Sébastien This<sup>1,2,3</sup> and Helena Paidassi<sup>1\*</sup>

<sup>1</sup>Centre International de Recherche en Infectiologie (CIRI), Univ Lyon, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, ENS de Lyon, Lyon, France, <sup>2</sup>Centre de Recherche de l'Hôpital Maisonneuve-Rosemont, Montréal, QC, Canada, <sup>3</sup>Département de microbiologie, immunologie et infectiologie, Université de Montréal, Montréal, QC, Canada

Transforming growth factor- $\beta$  (TGF $\beta$ ) is a long-known modulator of immune responses but has seemingly contradictory effects on B cells. Among cytokines, TGF $\beta$  has the particularity of being produced and secreted in a latent form and must be activated before it can bind to its receptor and induce signaling. While the concept of controlled delivery of TGF $\beta$  signaling *via*  $\alpha_v\beta_8$  integrin-mediated activation has gained some interest in the field of mucosal immunity, the role of this molecular mechanism in regulating T-dependent B cell responses is just emerging. We review here the role of TGF $\beta$  and its activation, in particular by  $\alpha_v\beta_8$  integrin, in the regulation of mucosal IgA responses and its demonstrated and putative involvement in regulating germinal center (GC) B cell responses. We examine both the direct effect of TGF $\beta$  on GC B cells and its ability to modulate the functions of helper cells, namely follicular T cells (Tfh and Tfr) and follicular dendritic cells. Synthetizing recently published works, we reconcile apparently conflicting data and propose an innovative and unified view on the regulation of the GC reaction by TGF $\beta$ , highlighting the role of its activation by  $\alpha_v\beta_8$  integrin.

## KEYWORDS

TGF $\beta$  activation, alpha(v)-beta8 integrin ( $\alpha_v\beta_8$ ), IgA B cell response, germinal center (GC) reaction, follicular T helper cells (Tfh), follicular regulatory helper T cell (Tfr), follicular dendritic cell (FDC)

**Abbreviations:** TGF $\beta$ , Transforming growth factor- $\beta$ ; LTBP, Latent TGF $\beta$  binding protein; GARP, Glycoprotein-A repetitions predominant protein; LAP, Latency-associated peptide; TLR, Toll like receptor; BCR, B cell receptor; RA, Retinoic Acid; Ig, Immunoglobulin; Ag, Antigen; cDC, Conventional dendritic cells; FDC, Follicular dendritic cells; Tfh, Follicular helper T cells; Tfr, Follicular regulatory T cell; Treg, Regulatory T cells; PC, Plasma cells; memB, Memory B cells; MLN Mesenteric lymph nodes; PP, Peyer's patches; SILP, Small Intestinal *lamina propria*; TD, T-cell dependent [B cell responses]; GCR, Germinal center reaction; GC, Germinal Center; LZ, Light Zone; DZ, Dark Zone; CSR, Class-switch recombination; SHM, Somatic hypermutation; KO, Knock-out.

## Introduction

The humoral arm of adaptive immunity has been at the center of discussion in the recent global COVID-19 pandemic. Efforts to detect and correlate SARS-CoV-2 neutralizing antibodies to protection or to generate these antibodies through vaccination are at the center of the current research landscape (1, 2). More generally, the different processes regulating the B cell responses have long been harnessed for targeted immunotherapies, vaccination being the first and best known. Defects in the regulation of humoral responses are also linked to many human pathologies: B cell hyperplasia, antibody (Ab)-mediated autoimmune disorders, graft rejection, allergy... (3, 4). Understanding the mechanisms underlying the regulation of humoral response reaction is an important goal to identify potential targets to improve vaccine efficacy, develop new therapeutic options for Ab-mediated disorders or propose novel immunotherapeutic strategies.

Pathogen clearance and formation of long-lasting humoral protection through antibody production, requires the activation of B lymphocytes and subsequent differentiation into antibody-secreting plasma cells (PC) and memory B cells (memB). In the vast majority of T-cell dependent (TD) responses, after initial protein antigen (Ag) encounter, antigen-specific B cells interact with antigen-specific T cells and form complex micro-anatomical structures called Germinal Centers (GC).

GC are the main sites where affinity maturation of the antibody response and generation of B cell memory most generally take place. While some B cell responses are GC-independent, the GC reaction (GCR) is key in determining the quality, the amplitude and persistence of the humoral response to TD Ag (5). This complex sequence of events is tightly coordinated by several interdependent cellular and molecular mechanisms which have been extensively described and reviewed over the years (6–10). Among these, cytokine-mediated regulation of the GC, either directly by their action on GC B cells or indirectly to control GC helper cells, is growing in importance.

In this review, we focus on the role of the Transforming Growth Factor  $\beta$  (TGF $\beta$ ) in the regulation of humoral immunity in the context of T-dependent response. The complex nature of TGF $\beta$  biology, and in particular the requirement for its activation, has limited the ability to study this cytokine in the context of the GC regulation. While such concept gained some interest in the field of mucosal immunity, in particular for the regulation of intestinal T cell and IgA responses, the role of  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation for regulation of the GCR has been poorly studied. Here, we will highlight the role of TGF $\beta$  activation in the regulation of B cell responses, particularly during the GCR.

## TGF $\beta$ , a complex pleiotropic cytokine

TGF $\beta$ , historically identified as a soluble molecule promoting fibroblast transformation and formation of growth

colonies, has since been shown to be a key cytokine for the regulation of the immune system, both for the maintenance of immune homeostasis and the regulation of inflammatory responses (11).

The three isoforms of TGF $\beta$ , namely TGF $\beta$ 1, TGF $\beta$ 2 and TGF $\beta$ 3, are produced by a wide diversity of cells, signaling through the TGF $\beta$  receptor (TGF $\beta$ R) on a large range of immune and non-immune cells. In the immune system, it is well accepted that the TGF $\beta$  cytokine, TGF $\beta$ 1 in particular, exerts strong immunomodulatory functions, acts as chemoattractant, promotes immune cell death and, is a critical regulator of immune cell differentiation, particularly of T cells, B cells, Natural Killer T cells and dendritic cells (DC) (12) Overall TGF $\beta$  plays a critical role in fine-tuning the immune responses inasmuch as it regulates magnitude and polarization of immune response, contributes to resolution of inflammation, and is critical for immune tolerance (13).

Besides, it is important to note that the studies on the regulation of immune responses by TGF $\beta$  have often been limited to the analysis of its secretion and downstream effects. But, among cytokines, TGF $\beta$  has a very peculiar position as it is produced and secreted in a latent form and needs to be activated to bind to its receptor and signal (14). Its latent nature is however often overlooked. We will give a brief introduction on the mechanisms of TGF $\beta$  activation,

While the role of TGF $\beta$  in T cell responses has been extensively studied (15), its implication in the regulation of humoral response has thus far mainly focused on Ig Class A (IgA) Class Switch Recombination (CSR) at mucosal surfaces. The overall function of TGF $\beta$  in modulating B cell responses, and in particular during the Germinal Center reaction (GCR), still remains elusive.

In this review, we will first review the instances where regulation of the GCR by TGF $\beta$  has been demonstrated, both from the standpoint of its direct effect on B cells and of its ability to modulate the functions of helper cells, namely follicular T cells and follicular dendritic cells (FDC). We will then review what is known of TGF $\beta$  activation and IgA CSR as a means to give insight in the direct or indirect regulation of GC by TGF $\beta$ -activation. Based on all these studies, we propose several hypotheses for the mechanism(s) by which TGF $\beta$  might be activated for the regulation of GC B cell responses.

## Regulation of the GC reaction by TGF $\beta$

### Overall considerations for the TGF $\beta$ -mediated regulation of the GC

The GC reaction is a tightly coordinated cascade of events, initiated and driven by the presence of the cognate Ag. The

general mechanisms underlying proliferation, affinity maturation and differentiation of B cells in the GC have been largely reviewed elsewhere and will not be described here (5). We highlight here two important parameters of the GCR that relate to its regulation by TGF $\beta$  and its activation.

1/The regulation of the GC involves interaction of B cells with multiple cell partners at different steps of the GCR; follicular T cells (namely follicular helper T cells (T<sub>fh</sub>) and follicular regulatory T cells (T<sub>fr</sub>)) and stromal cells, such as Follicular Dendritic Cells (FDC) (6–10). These cell/cell interactions occur *via* delivery of membrane-bound receptor/ligand signals (thus requiring physical contact and to some extent immune synapse formation) as well as *via* soluble factors. It is now well accepted that the type and stability of the synapses formed between B cells and other cell players of the GC are critical in determining the output of the GC (16). In that particular context, and given the mechanisms for TGF $\beta$  activation described below, we believe that controlled delivery of active TGF $\beta$  in the context of different combinations of cytokines and/or co-stimulatory factors delivered at cell/cell contact would account for the apparent pleiotropic effect of TGF $\beta$  on GC B cell responses.

2/The process of CSR, allowing B cells to switch from IgM and IgD expression to downstream isotypes (such as IgG, IgA, IgE), has long been thought to be an integral part of GC. However, Rocco and colleagues recently extended early observation by Toellner *et al.* to establish that CSR occurs infrequently within the GC (17, 18) and is instead initiated prior to entry into the B cell follicle at the T cell/B cell border (18). This review will detail to some extent the importance of TGF $\beta$  in IgA CSR in the mucosal compartment. This subject will be treated separately from the GCR, but we believe that similar mechanisms regulate IgA CSR and the GC reaction *via* activation of TGF $\beta$ .

## Direct TGF $\beta$ stimulation of B cell influences the GC output

TGF $\beta$  is a long-known modulator of antibody responses (19) (Table 1). Early *in vitro* studies in the 1980s and 1990s have highlighted the importance of TGF $\beta$  signaling for the regulation of B cell responses both in human and mouse. TGF $\beta$  was shown to inhibit B cell proliferation *via* growth arrest, to regulate B cell survival *via* induction of apoptosis, to inhibit Ig synthesis and to suppress CSR towards most Ig classes in favor of promoting the IgA class [reviewed in (12, 19, 49)]. In 2000, most of these observations were confirmed *in vivo* by Cazac and Roes (24). Using conditional deletion of TGF $\beta$ RII on B cells (*Cd19<sup>Cre</sup>.Tgfb2<sup>fllox/fllox</sup>*), they showed that deficient TGF $\beta$  signaling in B cells led to reduced lifespan of B2 B cells, expansion of B1-peritoneal B cells, elevated IgG3 responses to a weakly immunogenic antigen, generation of anti-dsDNA

antibodies and overall B cell hyperresponsiveness but did not lead to overt clinical autoimmunity. B cell specific TGF $\beta$ RII deficiency was also associated with a complete loss of IgA. Following these preliminary observations, many studies have further investigated the role of TGF $\beta$  for the regulation of mucosal B cell responses, and TGF $\beta$  is now widely considered as the master regulator of IgA CSR [reviewed in (29, 30)] (Table 1). Conceivably because of this, the study of TGF $\beta$ -mediated regulation of B cell responses has often been restricted to its ability to promote IgA CSR and to inhibit B cell proliferation.

Studies investigating TGF $\beta$  signaling specifically on GC B cells reveal that its action might be more refined than initially thought. In agreement with its known inhibitory functions, TGF $\beta$  stimulation promotes apoptosis of GC B cells through regulation of Bcl2 family members and inhibition of BCR-mediated rescue from apoptosis (25, 26). Accordingly, the loss of TGF $\beta$  signaling in B cell responses induced GC hyperplasia in the PP compartment, chronically stimulated by the gut microflora (24). Interestingly, stimulation of the TGF $\beta$ RI by BMP7 (member of the TGF $\beta$ -superfamily) also promotes apoptosis of human GC B cells through the canonical Smad-signaling pathways (50), showing multiple redundant pathways for controlling GC B cell survival.

More recently, a study analyzing *in situ* pSmad2 signaling in mice by confocal microscopy, revealed that TGF $\beta$  signaling in GC B cells promotes the Light Zone (LZ) to Dark Zone (DZ) transition (31). Accordingly, GC B cell specific-TGF $\beta$ RII deficiency induced accumulation of cells in the LZ, led to decreased apoptosis and in turn decreased antibody affinity. Albright and colleagues thus propose that the specific TGF $\beta$  delivery to GC B cells, inside established GC structures, might be driving GC LZ to DZ trafficking rather than IgA CSR (31).

It is important to note that most of these studies did not investigate the source of TGF $\beta$  production and delivery to B or GC B cells. Albright and colleagues did however suggest, using conditional deletion of stromal cells, that GC B cell pSmad signaling is at least partially mediated by FDC.

## FoxP3-expressing T cells regulate the humoral response through TGF $\beta$ -dependent mechanisms.

FoxP3<sup>+</sup> T cells have previously been involved in the regulation of multiple aspect of humoral responses through, among other, B cell apoptosis, CSR or Ig production [reviewed in (51, 52)]. Furthermore, several studies demonstrated that FoxP3<sup>+</sup> T cell-mediated suppression of humoral responses is at least partially mediated by TGF $\beta$  (53–56). More recently, it has been demonstrated that CD25<sup>+</sup>Lag3<sup>+</sup> Treg secreting TGF $\beta$ 3 suppressed antibody secretion, through mechanisms inhibiting crucial pathways for B cell differentiation and survival (57, 58).

TABLE 1 Pleiotropic functions of TGF $\beta$  in the regulation of humoral responses.

		Immunostimulating functions of TGF $\beta$		Immunosuppressive functions of TGF $\beta$	
		Effects of TGF $\beta$	Reference	Effects of TGF $\beta$	Reference
<b>Direct effect of TGF<math>\beta</math> on B cells</b>				<i>In vitro</i> , TGF $\beta$ limits B cell proliferation by inducing growth arrest	(20–23) (reviewed in (19))
		TGF $\beta$ is required for B2B cell survival	(24)	<i>In vitro</i> , TGF $\beta$ decreases B cell survival by inducing apoptosis	(25–28) (reviewed in (19))
		TGF $\beta$ induces IgA CSR	(24) (reviewed in [(29, 30)])	TGF $\beta$ controls IgG3 responses, limits expansion of peritoneal B1-B cells, limits B cell responsiveness, limits GC in PP	(24)
		TGF $\beta$ increases Ab affinity during GC by promoting LZ/DZ transition	(31)	<i>In vitro</i> , TGF $\beta$ suppresses CSR toward Ig in favor of IgA	(32, 33)
<b>Indirect effects of TGF<math>\beta</math> on GC via</b>	<b>Tfh</b>	TGF $\beta$ is required for Tfh induction in viral influenza infection	(34)	TGF $\beta$ inhibits IgG synthesis	(22)
		TGF $\beta$ is required for human Tfh differentiation	(39, 39, 40)	TGF $\beta$ limits Tfh frequency and suppresses Tfh function ( <i>in vitro</i> and <i>in vivo</i> )	(35–38)
	<b>Tfr</b>			<i>In vitro</i> , TGF $\beta$ inhibits mouse Tfh differentiation	(39, 41, 42)
				TGF $\beta$ induces Foxp3 expression by Tfr <i>in vitro</i>	(43)
	<b>FDC</b>			TGF $\beta$ limits auto-immunity by promoting Tfr development	(32, 36, 44)
				TGF $\beta$ limits conversion of Tfr into Tfh	(45)
		<i>In vitro</i> , TGF $\beta$ prevents TNF $\alpha$ -induced apoptosis of FDC, which may promote a functional FDC network during the GC	(46)		
		TGF $\beta$ promotes GC B cell survival by inducing PG production FDC	(47, 48)		

In the light of the more recent literature, it is however important to consider that these instances of TGF $\beta$ -mediated regulation of B cells by FoxP3<sup>+</sup> T cells could be mediated by the recently characterized Tfr rather than conventional Treg.

Indeed, 10 years ago, a new subset of Foxp3<sup>+</sup> regulatory T cells named follicular regulatory T cell (Tfr) has been discovered, that inhibits the GC reaction by directly suppressing Tfh and/or B cells. Since the initial description of Tfr in 2011 (59–61), a growing number of studies has documented the role of these cells in the regulation of GC B cell responses [reviewed in (8, 62, 63)]. While several studies have shown a role for CTLA4 and PD1 in the direct regulation of GC B cells and Tfh, indirect evidence suggest that TGF $\beta$  may also be part of the Tfr immunomodulatory machinery. Sayin *et al.* demonstrated that Tfr express elevated levels of the TGF $\beta$ -binding protein GARP, at higher levels than conventional Treg (64, 65). McCarron and colleagues also showed that Tfr display significant staining for TGF $\beta$  (35). Furthermore, Miles and colleagues, demonstrated that HIV infected tonsil cells displayed a higher frequency of TGF $\beta$  expressing Tfr (36), inhibiting B cell responses indirectly through inhibition of Tfh proliferation, ICOS expression and cytokine secretion. More recently, Turner *et al.* showed that the specific deletion of TGF $\beta$ 1 in all FoxP3-expressing cells (Tfr and conventional Treg) resulted in a high increase of Tfh frequency,

suggesting that Tfr-derived TGF $\beta$  could regulate Tfh proliferation. This result is in agreement with the study from McCarron and Marie in which they show that TGF $\beta$  receptor-deficiency in all T cells leads to the aberrant accumulation of Tfh cells in mice (35, 38). Altogether, these studies point to a critical role of TGF $\beta$  in Tfr immunomodulatory functions, even though the precise mechanisms involved in active TGF $\beta$  delivery remain to be established.

## TGF $\beta$ indirectly regulates GC B cell through regulation of helper cell differentiation and/or function

### TGF $\beta$ a newly established regulator of Tfr development

It is well established that TGF $\beta$  is a critical cytokine for the induction of Foxp3<sup>+</sup> Treg (iTreg) in the periphery and the maintenance of FoxP3 expression by Foxp3<sup>+</sup> Treg cells (15, 66). A few recent studies also suggest a role for TGF $\beta$  in Tfr development. In *ex vivo* models of HIV or HCV infection, important Tfr expansion is detected. Specifically, during HIV infection, this increase in Tfr frequency is only partly mediated by TGF $\beta$ R signaling, while exosomes secreted by HCV infected

hepatocytes, containing large quantities of TGF $\beta$ , appear to act directly on T cells to induce Tfr from activated human CD4<sup>+</sup> T cells (36, 67). Despite some efforts, the authors however failed to fully demonstrate (i) whether this was due to direct TGF $\beta$  signaling to T cell or other cells supporting Tfr differentiation (DC, B cells, ...) and (ii) whether this was due to *de novo* Tfr differentiation or proliferation of pre-existing Tfr. In parallel, in a mouse model of spontaneous autoimmune development, IL2 and TGF $\beta$  synergize *in vivo* to promote Tfr development in the periphery in naïve mice (44). Interestingly, TGF $\beta$  stimulation of naïve T cells induces the miR-10a-5p micro-RNA. This micro-RNA, is expressed at high levels on Treg and Tfr and constrains their conversion into Tfh (45). Altogether, these studies suggest that TGF $\beta$  signaling on Tfr is important for their induction, their expansion and/or their persistence during the GC reaction.

More recently, Jacobsen and colleagues proposed a mechanism by which Tfh acquire FoxP3 expression and an intermediate CD25<sup>-</sup> Tfr phenotype in the late stages of the GCR (43). They proposed that these late Foxp3-expressing Tfh participate in the termination of the GCR reaction. TGF $\beta$  stimulation of Tfh *in vitro* was capable of inducing FoxP3 expression but the *in vivo* demonstration remains to be made (43).

Overall, it seems that TGF $\beta$ , in synergy with other cytokines and signals, is a novel regulator of Tfr differentiation. Some contradicting evidence, such as the study by McCarron and Marie demonstrating that T-cell specific TGF $\beta$ R deficiency did not lead to a decreased Tfr population (35), however highlights that the mechanism by which TGF $\beta$  controls Tfr differentiation has not been fully resolved. It is important to note that Tfr remain a recently-described T cell population and that later studies have demonstrated that Tfr encompass multiple subtypes, derived from either natural thymic Treg, peripherally induced iTreg or Tfh, and generated under different stimulation or immunization scheme and pathological models [summarized in (63)]. It is likely that the cellular origin, localization, and mechanisms of induction of each Tfr subtype might dictate a different requirement for TGF $\beta$ .

## The controversial role of TGF $\beta$ in the differentiation of Tfh

In the study of human Tfh differentiation it is now well accepted that TGF $\beta$  is a critical factor for the differentiation of naïve T cells into Tfh. TGF $\beta$  stimulation induces robust *Bcl6*, *Cxcr5*, *Pdcd1* (coding for PD1), *Icos* and *Il21* expression by peripheral blood CD4<sup>+</sup> T cells (34, 39, 40).

In comparison, studies in mice show a more controversial role of TGF $\beta$ . On one hand, early *in vitro* studies, demonstrated that TGF $\beta$  stimulation of naïve murine T cells, inhibits the expression of *Il21*, *Bcl6*, *Icos* and *Cxcr5*, and thus the differentiation of Tfh (39, 41, 42). However, blocking TGF $\beta$  signaling *in vivo*, using either anti-TGF $\beta$  blocking antibody or the TGF $\beta$ RII CD4<sup>+</sup> T cell conditional-dominant negative (DN)

mouse model (*dnTGF $\beta$ RII*), did not alter the frequency of Tfh in models of NP-KLH immunization and influenza infection (35, 41, 68). Interestingly, while *dnTGF $\beta$ RII* mice have a similar frequency of Tfh than their WT counterpart upon immunization with NP-KLH, these mice spontaneously develop autoimmune symptoms with an important accumulation of Tfh. The authors propose that, while TGF $\beta$  does not appear to control Tfh induction, TGF $\beta$  signaling may be required for the regulation of their survival (35).

On the other hand, in the context of acute LCMV infection, Marshall and colleagues revealed an important TGF $\beta$  signature on Tfh associated with the strong upregulation of genes generally associated with Treg such as *Nt5e* (CD73), *Folr4* (folate receptor 4), *Foxp3*, and *Ikzf2* (Helios) among other (34). Investigating the chromatin organizer *Satb1*, several studies have shown that TGF $\beta$  stimulation of murine T cells silences *Satb1* expression, which in turn promotes Tfh differentiation (40, 69) Using a TCR transgenic TGF $\beta$ RII CD4<sup>+</sup> T cells conditional-KO reveals that *in vivo* generation of Tfh following influenza infection required TGF $\beta$  signaling on T cells. In this context, TGF $\beta$  inhibits mTOR signaling in T cells and dampens IL2 responsiveness allowing for Tfh differentiation (34). It is important to note that the influenza neuraminidase enzyme can promote the cleavage of latent TGF $\beta$  in the lung mucosae (70, 71). While this might not be sufficient to explain the discrepancy in TGF $\beta$  requirement for Tfh induction in the study by Marshall *et al*, as compared to the other murine approaches, it is important to note that a wide variety of potentially unknown biases can be at play between these studies. Thus, while TGF $\beta$  is clearly required for human Tfh differentiation, it is, to this day, difficult to understand the disparity found in the role of TGF $\beta$  for murine Tfh differentiation.

## Effects of TGF $\beta$ on Tfh function

In addition to its controversial role in the differentiation of Tfh, TGF $\beta$  also has direct suppressive effects on the function of Tfh, which thus indirectly impacts GC B cell responses. In a recent study using co-cultures of GC-Tfh and GC-B cell isolated from human tonsils, O'Connor and colleagues show that addition of TGF $\beta$  is sufficient to inhibit the secretion of IL-21 and sCD40L by Tfh. This is associated with a decreased production of IgG. While a direct effect of TGF $\beta$  on GC B cells cannot be excluded in this co-culture model, this study suggests that TGF $\beta$ -mediated suppression of Tfh function would be sufficient to regulate GC B cell responses (37).

This hypothesis is supported by earlier studies both *in vitro* and *in vivo*. As discussed earlier, human tonsil Tfr have the ability *in vitro* to inhibit Tfh proliferation, ICOS expression and cytokine secretion (36). Interestingly, neutralization of TGF $\beta$  in co-cultures of Tfh and Tfr restored most of IL-21 production by Tfh, a crucial B cell help cytokine (36). In addition, *in vivo*, mice

with a TGF $\beta$  deficiency in FoxP3-expressing cells (including Tfr) (*Foxp3<sup>cre</sup>.Tgfb1<sup>lox/lox</sup>*) or with impaired TGF $\beta$  signaling in T cells (including Tfh) (*Cd4<sup>cre</sup>.Tgfb2<sup>lox/lox</sup>*) develop fatal autoimmunity associated with increased frequency of Tfh and GC B cells (35, 38). McCarron and Marie show that accumulation of Tfh wasn't due to an excessive proliferation of Tfh but rather that Tfh lacking TGF $\beta$  signaling are resistant to apoptosis (35). In parallel, O'Connor and colleagues recently identified a new regulatory Innate Lymphoid Cell (ILC) population in the follicles of human tonsils and LN named follicular regulatory ILC (ILCfr), which inhibits the ability of Tfh to secrete IL-21 and sCD40L *in vitro* (37). Interestingly, ILCfr secretes TGF $\beta$  upon activation and in co-cultures of human tonsil Tfh, GC B and ILCfr cells, TGF $\beta$  neutralization was sufficient to restore IL-21 secretion by Tfh as well as IgG levels by GC B cells (37).

Altogether, these studies demonstrate that TGF $\beta$  affects Tfh function (Tfh survival, Tfh expression and secretion of critical B cell help factors) and thus indirectly regulates GC B cell response and autoimmunity. The relative contribution of Tfr and ILCfr in the secretion of TGF $\beta$  for the regulation of the ability of GC Tfh to provide B cell help remains to be established.

### TGF $\beta$ -dependent regulation of FDC survival and function

Follicular Dendritic Cells (FDC) form a dense network of stromal cells involved in different aspects of GC B cell proliferation and selection. These cells are critical in the formation of the GC and provide multiple soluble factors instrumental for the correct development of the GCR (9).

Apical LZ FDC have been shown to express the TGF $\beta$ BR (72, 73). Following these early observations, Park *et al.* first described *in vitro* that TGF $\beta$ -induced Smad2 signaling in FDC-like cell lines induced a decreased expression of the death receptor signaling pathway: Fas and Caspase 8 (46). TGF $\beta$  signaling may thus play a role in preventing Fas-mediated apoptosis of FDC and thus to maintain a functional FDC network throughout the GCR.

Additionally, studies report that TGF $\beta$  stimulation of human FDC-like cell promotes, in synergy with IL1 $\beta$ , the production of the Prostaglandin-endoperoxide synthase 2 (Cox-2) enzyme, and in consequence promotes Prostaglandin production (47, 48). Prostaglandin presentation by FDC-like cells *in vitro* actively participates in promoting GC B cell survival (74). Therefore, TGF $\beta$  may also indirectly participates in the promotion of the GCR *via* regulation of FDC functions.

Overall, TGF $\beta$  appears to have multiple and potentially opposing effects on the GCR. When acting directly on B cells, for example TGF $\beta$  has been shown to induce GC B cell apoptosis, and to promote LZ/DZ trafficking and thus increased Ab affinity. Regarding the indirect effects of TGF $\beta$ , on one hand TGF $\beta$  promotes FDC survival and therefore supports the GC reaction, while on the other hand it promotes

Tfr expansion and persistence, which inhibit GC B cell. Finally, its role in regulating Tfh development remains controversial. Altogether these observations highlight that whether acting directly on B cells or indirectly *via* GC helper cells, the role of TGF $\beta$  is to this day not entirely resolved and requires further investigation. We believe that these apparent contradictions might be due to the specific nature of TGF $\beta$  that needs to be activated to signal.

## $\alpha_v\beta 8$ -mediated TGF $\beta$ activation is key for bioavailability of TGF $\beta$ in the immune system

### Latent-TGF $\beta$ requires activation prior to signaling

TGF $\beta$  has a very peculiar position among cytokines, as it is produced in tissues in a latent form and must be activated in order to bind to its receptor and enable subsequent signaling and functions. The biology of TGF $\beta$  production and activation has been extensively reviewed in several fields of biology (75–78). Figure 1 recapitulates the mechanism for TGF $\beta$  production, sequestration in the tissues, activation and signaling. It is important to note that TGF $\beta$ , is overall largely available in tissues and in the serum in a latent form (Figure 1A) (81). TGF $\beta$  can be found both in the extracellular matrix and bound to cell surface through the recently identified Glycoprotein-A Repetitions Predominant protein (GARP) (Figure 1B) (75, 77, 79). Hence, despite some regulation of TGF $\beta$  signaling being made at the level of TGF $\beta$  secretion by immune cells, most of the regulation of TGF $\beta$  bioavailability and downstream signaling is done at the level of its activation.

Many pathways for TGF $\beta$  activation have been described (75), however  $\alpha_v$  integrin-mediated activation of latent TGF $\beta$  appears to be one of the most prominent pathway for TGF $\beta$  activation *in vivo*. Integrins are heterodimeric transmembrane adhesion molecules, composed of one alpha ( $\alpha$ ) and one beta ( $\beta$ ) subunit, that mediate cell-cell and cell-extracellular matrix interactions (82).  $\alpha_v$  is the most promiscuous of  $\alpha$  subunits, pairing with five different  $\beta$  integrins ( $\beta 1$ ,  $\beta 3$ ,  $\beta 5$ ,  $\beta 6$  and  $\beta 8$ ). The  $\alpha_v$  integrins have been implicated in many different cell functions, but their ability to bind and activate latent TGF $\beta$  is of particular importance to immune regulation, licensing  $\alpha_v$ -expressing cells to the control of several immune processes (75–77).

Briefly, binding of TGF $\beta$  by  $\alpha_v$  integrins is mediated through an Arginine-Glycine-Aspartate (RGD) tripeptide present at the surface of the LAP of latent TGF $\beta$  (Figure 1C). Activation of latent TGF $\beta$  follows its binding to  $\alpha_v$  integrins. While many ways of TGF $\beta$  activation have been described *in vitro* (proteolytic degradation, deglycosylation, or physicochemical




FIGURE 1

$\alpha_v\beta_8$  integrins regulate TGF $\beta$  bioavailability in the immune system. TGF $\beta$  is produced by cells as an inactive complex and must be activated in order to bind to its receptor and signal. (A) Transcription of *TGF $\beta$*  produces a homodimeric propeptide containing the active TGF $\beta$  molecule (blue) and the Latent Associated Peptide (LAP; in red). In the endoplasmic reticulum, association with the LTBP or GARP 'chaperone' proteins (green) enhances proper folding of the latent complex. In the Golgi apparatus, LAP-TGF $\beta$  is cleaved by Furin-like enzymes, but active TGF $\beta$  stays non-covalently bound to the LAP and forms the Small Latency Complex or latent TGF $\beta$  (79) (B) Upon secretion, latent TGF $\beta$  is sequestered to the ECM through binding of the LTBP or anchored at the plasma membrane by GARP. (C) Binding of  $\alpha_v$  integrins to the RGD tripeptide motif in the LAP induces the dissociation of TGF $\beta$  from the LAP via the recruitment of metalloproteases such as MMP14 (78). Alternatively, conformational changes can allow TGF $\beta$  binding to its receptor without the release of active TGF $\beta$  (80). (D) TGF $\beta$  binding to the TGF $\beta$  receptor induces signaling via the canonical phospho-Smad (pSmad) pathway, or through the alternative MAPK, Small GTPases and PIP3K pathways.

factors (ROS, low pH condition or UV radiation) (75),  $\alpha_v$  integrin-mediated activation of latent TGF $\beta$  appears to be the most important pathway for TGF $\beta$  activation *in vivo*. Mutation in the RGD sequence in the TGF $\beta$  LAP, which disrupts the  $\alpha_v$  integrin-binding site (*Tgfb1*<sup>RGE/RGE</sup> mice), indeed recapitulates many of the phenotypes of the TGF $\beta$  knockout (KO) mouse (83). Thus, through their involvement in the conversion of latent TGF $\beta$  to a form that binds and signals on its receptor,  $\alpha_v$  integrins have the intriguing ability to regulate TGF $\beta$  activation and bioavailability.

While  $\alpha_v\beta_1$ ,  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$  have been shown to bind and activate latent TGF $\beta$  in the context of fibrosis and fibroblast differentiation and function (84–89) its contribution to the regulation of TGF $\beta$ -dependent immune responses *in vivo* has

not been demonstrated to this day. On the contrary, deletion of the high affinity binding integrins,  $\alpha_v\beta_6$  or  $\alpha_v\beta_8$ , causes failure of effective TGF $\beta$  signaling *in vivo* as both  $\beta_6$ - and  $\beta_8$ -deficient mice develop inflammation (90, 91). Thus both  $\alpha_v\beta_6$  and  $\alpha_v\beta_8$  integrins have been shown to activate TGF $\beta$  for the regulation of immune responses (78) (76). However, while  $\beta_6$  KO mice only have a mild phenotype,  $\beta_8$ -deficient mice phenocopies mice with a selective loss of  $\alpha_v$  integrin-mediated TGF $\beta$ 1 activation (83). This difference in phenotype might be explained by two observations. First,  $\alpha_v\beta_8$  integrin has a divergent cytoplasmic domain compared to other  $\alpha_v$  integrins and its constitutionally open conformation likely confers an advantage for the binding of TGF $\beta$  without prior inside-out signaling (92–94). Furthermore,  $\alpha_v\beta_6$  expression seems to be restricted to



epithelial cells, contrary to  $\alpha_v\beta_8$  that has been shown to be expressed by inflammatory fibroblasts and specific immune cell populations (cDC1, Treg). Therefore, because  $\alpha_v\beta_8$ -activation of TGF $\beta$  is emerging as critical regulator of TGF $\beta$  immune responses, and because  $\alpha_v\beta_6$  has, to this day, never been associated with the regulation of B cell, only the role of  $\alpha_v\beta_8$ -mediated activation will thus be reviewed here.

Currently two main mechanisms for  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation are now being accepted, both involving a third molecular partner [Reviewed in (95)] (Figure 1C). On one hand, it has been demonstrated *in vitro* that the Matrix metalloproteinase-14 (MMP14, also called MT1-MMP) is required for the  $\alpha_v\beta_8$ -dependent activation of TGF $\beta$ , suggesting a cleavage-dependent local release of active TGF $\beta$  (Figure 1C) (96). Since then, the importance of MMP14 in the *in vitro* and *in vivo* activation of TGF $\beta$  has been demonstrated in multiple context (endothelial function, bone development and pathology, senescence and cancer progression...) (97–102). On the other hand, a few studies have demonstrated the importance of the GARP molecule in “chaperoning” the latent TGF $\beta$  molecule for  $\alpha_v\beta_8$ -mediated activation (103–105). The crystal structure of the GARP-TGF $\beta$  complex was recently elucidated and showed that LAP binding by GARP allows the further binding of TGF $\beta$  by  $\alpha_v\beta_8$  in a conformation allowing an important flexibility (80, 106). The authors suggest that presentation of latent TGF $\beta$  by GARP to  $\alpha_v\beta_8$  integrin might allow for active TGF $\beta$  to be presented to the TGF $\beta$ R without the need of active soluble TGF $\beta$  release at a cell/cell synapse, TGF $\beta$  thus remaining membrane bound.

## Regulation of $\alpha_v\beta_8$ expression, a checkpoint for the modulation of TGF $\beta$ -dependent immune responses

As mentioned earlier,  $\alpha_v\beta_8$  is a decisive factor for understanding TGF $\beta$ -regulated immune responses, especially in the mucosal interfaces. We and others have previously established the key role of  $\alpha_v\beta_8$  on DC for regulation of intestinal T cell responses *via* the presentation of activated TGF $\beta$  to naïve T cells (107–109). Additionally,  $\alpha_v\beta_8$  expression by Treg themselves is also critical for regulating overt effector T cell-mediated inflammation in the gut (110). Since then,  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  has been involved in the regulation of other TGF $\beta$ -dependent immune processes such as IEL generation, regulation of intestinal inflammation, etc. (111–117). This is partly reviewed in (78).

These studies, however revealed two key properties of  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation that, we believe, will be important to understand the TGF $\beta$ -dependent regulation of the GCR. First  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation requires cognate interaction between DC and T cells (116). Hence, reinforcing our assumption that understanding cell/cell contact in the GC is

critical. Second,  $\alpha_v\beta_8$  is expressed in a stable extended-closed conformation, which is not affected by ligand binding or ‘inside-out’ signals, hence excluding the possibility of a contractility-dependent mechanism for  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation. The key mechanism for regulating TGF $\beta$  activation and thus TGF $\beta$  signaling is therefore through the expression of  $\beta_8$  gene (*Itgb8*) (108, 118–121).

In summary, the large availability of TGF $\beta$  in tissues and serum, mostly in a latent form (81), and the various pathways for its activation have been powerful arguments to establish that the regulation of TGF $\beta$  bioavailability is not only dependent on its secretion but also and most importantly on its activation. In the context of the immune system,  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation is emerging as a dominant, if not major pathway for the *in vivo* regulation of activated TGF $\beta$  availability and regulation of TGF $\beta$ -dependent T cell responses. However, very little is known about the role of  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation in regulating B cell responses.

## $\alpha_v\beta_8$ -mediated activation of TGF $\beta$ is required for optimal mucosal IgA responses

TGF $\beta$  is the master regulator of IgA class switch recombination but the importance of its activation for this purpose was, until recently, unknown (29, 30). We will focus on the demonstrated and putative mechanisms of  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  by the various B cell partners during IgA response, namely, conventional DC (cDC), T cells (FoxP3-expressing T cells in particular), and FDC (Figure 2). It is important to note that despite the high abundance of TGF $\beta$  in the IgA PC residency niche, the regulation of PC biology by TGF $\beta$  remains to be established. Here, we will thus focus on the well-established role of TGF $\beta$  in the induction of IgA responses.

Of note, the origin of latent TGF $\beta$  is not precisely investigated but can likely be attributed to mucosal cDC, Tfh and/or FDC, or alternatively non-immune cells such as epithelial cells and fibroblasts, an important sources of latent TGF $\beta$  in gut and lung tissues (29, 128–131). Additionally, B cell can also produce TGF $\beta$  (131), and this ability has been linked to their cell-intrinsic ability to promote IgA response, placing B cell as a likely important source of TGF $\beta$  for IgA responses (132). Furthermore, mature B cells can express the TGF $\beta$ -docking GARP protein following B cell receptor and Toll Like Receptor (TLR) stimulation (126, 127), with GARP expression by B cells being important for optimal fecal IgA responses (126, 127). Campbell and colleagues additionally propose that  $\alpha_v\beta_8$  activation of GARP-bound TGF $\beta$  can promote cis-presentation of active-TGF $\beta$  on the TGF $\beta$ -producing cell (See Figure 1D) (80). Altogether, these observations suggest that GARP physically takes TGF $\beta$  produced by B cells to the




FIGURE 2

Demonstrated and putative roles of  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  in the regulation of mucosal IgA responses. (A) At steady state, Reboldi and colleagues suggest that  $\alpha_v\beta_8$  expression by Peyer's Patches CD103<sup>+</sup> cDC is required for the induction of IgA response to the commensal flora (122). Additionally, TGF $\beta$  activation by FDC through mechanisms yet to be determined might also be a mechanism for the promotion of IgA response in the PP (123) (B) In the context of intestinal rotavirus infection, MLN cDC1, migrating from the LP, are required for optimal RV-specific IgA response, in part promoted via  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  (124). (C) In lung-draining LN, the ability of individual cDC subpopulation to induce IgA response to microbiota via TGF $\beta$  production is correlated with their expression of the *Itgb8* transcript (125). (D) Despite lack of formal demonstration, we propose that, given their molecular arsenal and TGF $\beta$ -dependent function in the intestinal mucosae,  $\alpha_v\beta_8$ -expressing Foxp3<sup>+</sup> T cells (Treg) could activate GARP-bound TGF $\beta$  on B cells for induction of IgA responses (104, 110). More generally, the expression of GARP-TGF $\beta$  complexes by B cells suggest that activated B cells themselves could be a physiologically relevant source of latent TGF $\beta$  (126, 127).

immune synapse and that GARP-bound TGF $\beta$  is an important source of TGF $\beta$  in the regulation of IgA responses (Figure 2).

## $\alpha_v\beta_8$ -mediated activation of TGF $\beta$ by mucosal DC regulates IgA responses

We and others have shown that mucosal cDC, and in particular mesenteric lymph nodes (MLN) migratory cDC, i.e. migrating from the intestinal *lamina propria* (LP) to the MLN, specifically express  $\alpha_v\beta_8$  integrin, which licenses them to activate TGF $\beta$  and regulate TGF $\beta$ -dependent immune T cell responses (108, 109, 119). More recently, using a newly established reporter mouse model for  $\beta_8$  integrin (*Itgb8*) gene expression, we have shown that *Itgb8* is preferentially expressed by a large proportion of MLN migratory type 1 cDC (cDC1) and to a smaller extent by ~10% of MLN migratory type 2 cDC (cDC2). Additionally, a small fraction (~10%) of Peyer's Patches (PP) cDC1 also express the  $\beta_8$  integrin subunit (124). Ruane and

colleagues also demonstrated that  $\alpha_v\beta_8$  is expressed, at least at the RNA level, by cDC1 and cDC2 in the lung draining lymph nodes (125). It is interesting to note that  $\alpha_v\beta_8$  expression by cDC in naïve mice is restricted to mucosal associated lymph nodes as it not expressed in the spleen (119, 125). Consistent with this observation, we have shown that factors, commonly associated with the mucosal compartment such as Retinoic Acid (RA), microbial components, TLR ligands – CpG in particular - and TGF $\beta$  itself, promotes *Itgb8* expression on non-mucosal cDC (119, 122). These observations suggest a functional specialization of mucosal cDC for the regulation of mucosal-associated TGF $\beta$ -dependent immune responses.

Three studies recently started addressing the role of  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  in the regulation of IgA B cell responses. First, Ruane and colleagues showed in the lung mucosae, that both lung cDC1 and cDC2 promote the generation of protective IgA responses in a TGF $\beta$ -dependent manner. This ability correlates with higher expression of the  $\beta_8$  subunit as compared to other antigen presenting subsets,

suggesting a role for  $\alpha_v\beta_8$  expression in promoting lung anti-microbiota IgA responses (125). Second, Reboldi and colleagues have shown *in vitro* using a cDC:B cell coculture model that murine PP cDC from naïve mice are able to promote IgA response in a  $\beta_8$ -dependent manner. *In vivo*, they have further demonstrated that conditional KO of *Itgb8* on all CD11c-expressing cells, including cDC (*Cd11c<sup>Cre</sup> x Itgb8<sup>fllox</sup>* mice) or injection of an anti- $\beta_8$  blocking antibody impairs the formation of steady state IgA<sup>+</sup> GC B cells (122). Importantly the relative contribution of CD11c-expressing cells (cDC1, cDC2 or other CD11c<sup>+</sup> cells, including subsets of T cells and ILCs) responsible for  $\alpha_v\beta_8$ -mediated regulation of PP IgA response requires further investigating. Finally, our group has shown that MLN cDC1 but not MLN cDC2 can promote IgA responses *in vitro* via  $\alpha_v\beta_8$ -mediated binding and activation of latent TGF $\beta$ . Interestingly, targeted deletion of *Itgb8* in cDC1 (*Xcr1<sup>Cre</sup> x Itgb8<sup>fllox</sup>*) did not significantly alter IgA responses in naïve mice. However, in the context of viral enteric infection, such as the intestinal rotavirus infection,  $\alpha_v\beta_8$  expression by MLN cDC1 is required for the optimal generation of anti-viral IgA responses (124).

Furthermore, as alluded to earlier, it is thought that the timing of TGF $\beta$  delivery and the type of synapse established is an important factor to consider when thinking about TGF $\beta$  activation. Interestingly, Reboldi and colleagues propose that the  $\alpha_v\beta_8$ -mediated presentation of activated TGF $\beta$  occurs in the sub-epithelial dome of the PP, where prolonged cell/cell conjugates between CD11c<sup>+</sup> cells and B cells can be observed, prior to the entry of activated B cells into the GC (122). This is consistent with the description that CSR occurs infrequently inside the GC but rather prior to the entry into the B cell follicle (17, 18). While Roco and colleagues show that CSR is initiated at the T cell/B cell border and suggest that the signal for CSR are given by pre-follicular helper T cells, the study by Reboldi and colleagues further suggests that cDC could also provide important and complementary factors for IgA CSR following initial antigen encounter in the mucosal compartment. While we have not investigated the timing of active TGF $\beta$  delivery by MLN cDC1 to B cells during rotavirus infection (124), we propose that cDC, migrating from the lamina propria to the MLN, could present activated-TGF $\beta$  to B cells concomitantly with the presentation of native antigen coated on their surface, as previously demonstrated (133, 134).

Here it is critical to note that these studies have not formally demonstrated that the  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation is required for the induction of IgA responses from naïve B cells *per se* – through the quantification of  $\alpha$ -germline transcript or  $\alpha$ -circle transcript (CT $\alpha$ ) – but rather shows that  $\alpha_v\beta_8$  is required for the promotion of optimal IgA responses as a whole. Additionally,  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation is not the only mechanism by which cDC can support IgA responses; other mechanisms of TGF $\beta$  activation might be in place as well as other TGF $\beta$ -independent redundant and complementary

mechanisms which have already been described (BAFF, APRIL, IL6 and RA secretion) reviewed in (135).

Nevertheless, these studies point to a critical role of  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation in the molecular toolkit used by mucosal DC, both cDC1 and cDC2, for the promotion of IgA responses. Besides, it appears that the nature of the initial trigger (i.e viral or bacterial) and the tissue in which the immune response is initiated (i.e draining lymph node, PP or mucosa) will determine the type of cDC subset (i.e cDC1 vs cDC2) that mediates the  $\alpha_v\beta_8$ -mediated control of IgA responses.

## Putative role for T cells and FDC in $\alpha_v\beta_8$ -mediated control of IgA responses

Despite their conventional immunomodulatory function, intestinal FoxP3-expressing Treg are an important regulator of IgA responses. In 2009, Cong and colleagues first formally demonstrated the importance of CD4<sup>+</sup>CD25<sup>+</sup> Treg in the generation of IgA responses to flagellin in a TGF $\beta$ -dependent manner (136). Further studies confirmed that FoxP3<sup>+</sup> T cells and thymus derived Treg promote the generation of robust and diverse IgA responses in the gut in a TGF $\beta$ -dependent manner (137, 138). Kawamoto and colleagues, further propose that FoxP3<sup>+</sup> T cells differentiate into Tfr to control IgA production in the intestine (139, 140). Intestinal Treg, as well as Tfr, have been shown to express the  $\alpha_v\beta_8$  integrin (51, 110), the expression of which licenses Treg to activate GARP-bound latent TGF $\beta$  (104, 110). While  $\alpha_v\beta_8$  expressed by FoxP3-expressing T cells is dispensable for maintenance of intestinal immune homeostasis in naïve mice, it was shown to be required for suppression of T-cell-mediated intestinal inflammation. While no data is currently available, the expression of  $\alpha_v\beta_8$  by Treg and/or Tfr and their ability to control TGF $\beta$ -dependent responses, could be one of the mechanisms licensing them to modulate IgA responses and needs further investigation.

In addition to T cell, B cells also interact with stromal cells, and especially FDC which are known to secrete TGF $\beta$  (72, 73, 123, 141). In PP, Suzuki and colleagues have shown that FDC-M1<sup>+</sup> cells (which includes FDC and contaminating MFG-E8<sup>+</sup> macrophages) express at high level molecules associated with TGF $\beta$  activation ( $\alpha_v$  integrin subunit, Matrix Metalloproteases, CD36) (123). The authors further show that these TGF $\beta$ -activating molecules can be robustly induced in PLN FDC after stimulation with mucosal associated factors such as Retinoic Acid (RA) and TLR ligands. Like PP FDC-M1<sup>+</sup> cells, these “mucosa-imprinted” FDC then display an increased ability to promote IgA responses *in vitro*. In addition, stimulation with RA and TLR ligands reduces the level of LAP-TGF $\beta$ 1 present at the surface of PLN FDC, suggesting that active TGF $\beta$  is cleaved and shed from the FDC surface. Finally, PP FDC isolated from *Myd88*<sup>-/-</sup> mice or from mice fed with a vitamin A-deficient diet display increased LAP-TGF $\beta$ 1 present at the cell surface and

these mice display markedly reduced intestinal IgA<sup>+</sup> populations. Altogether these results suggest that FDC are licensed to promote IgA responses through TGFβ activation, most likely in an α<sub>v</sub>β8-dependent manner.

As CSR infrequently occurs in the GC (17, 18), it seems likely that FDC may play a role for induction of IgA CSR in primary follicles during the initial activation of B cells. It is important to note that the studies discussed here don't directly show whether TGFβ secretion, activation and/or presentation to B cells by FDC is required for the proper induction of IgA CSR. It is also possible that sustained TGFβ signaling by FDC from the initiation (pre-GC) to the termination of the GCR is required for induction of optimal IgA responses. The precise role of FDC-mediated TGFβ activation in the promotion of IgA responses therefore requires further investigation. The role of FDC-mediated activation of TGFβ for regulation of the GCR, outside of induction of IgA CSR, will be discussed in the next chapter.

To summarize, while the role of α<sub>v</sub>β8-mediated activation of TGFβ by DC for IgA responses is now well established, several studies suggest that follicular T cells and FDC could also activate and present TGFβ to B cells for the promotion of IgA CSR (Figure 2).

## Fine regulation of the GCR by TGFβ activation

As described earlier (Table 1), TGFβ has seemingly contradictory effects on B cell responses during the GC reaction. This reminds of the multiple effects of TGFβ on T cell responses in the gut, where induction of regulatory or inflammatory Th17 cell is dependent on the context of the synapse where TGFβ is presented to naïve T cells. Thus, the apparent multiple roles of TGFβ in the regulation of the GC are likely dependent on the synapse in which TGFβ is presented and delivered to the target cells.

Similar to induction of mucosal IgA responses, regulation of the GC involves a plethora of interaction between many cell types (B cells, Tfh, Tfr and FDC) which influence different steps of the GC reaction. Accordingly, regulation of their differentiation, activation, function and/or survival, is critical for an optimal GCR. Here we question the involvement of TGFβ activation, in particular *via* α<sub>v</sub>β8 integrin, in the fine-tuning of GC B cell responses *via* regulation of both the differentiation and the function of these follicular cell populations (Tfh, Tfr and FDC) (Figure 3).

## TGFβ-mediated regulation of the GCR by FDC *via* TGFβ activation

The relationship between TGFβ and FDC is complex, as TGFβ was proposed to be a critical regulator of FDC survival

and function *in vitro* as well as TGFβ being part of the molecular toolkit of FDC in their promotion of the GC and IgA CSR (31, 46, 47, 123). The question remains as to the mechanism of TGFβ activation for the regulation of GC B cells and FDC biology.

As stated earlier, mucosal associated factors (RA and TLR ligands) have been shown to induce expression of TGFβ-activating molecules in FDC-M1<sup>+</sup> cells (123). Importantly, FDC are of mesenchymal origin; other mesenchymal cell, such as fibroblasts, have also been shown to activate TGFβ through α<sub>v</sub>- and more specifically α<sub>v</sub>β8-specific mechanisms in the context of lung and liver fibrosis (144, 145). So even though the formal demonstration that FDC can indeed activate TGFβ and present it to B cells remains to be made, these results suggest that FDC may be able to provide active TGFβ for regulation of the GC reaction (123).

While the role of TGFβ in promoting FDC survival and function needs to be confirmed *in vivo*, it is also important to note that the source of latent TGFβ and its mode of activation is not known to this day. Although further investigations are warranted, we speculate that FDC might be able to activate TGFβ in an autocrine manner to promote their survival, at least in the context of mucosal GC. Alternatively, intestinal stromal cell have also been shown to express *Itgb8* and could participate in the paracrine regulation of the FDC network (146). More generally, outside of mucosal associated immune response, the question of α<sub>v</sub>β8-mediated regulation of FDC network remains entirely open.

Overall, the relationship between TGFβ and its activation and FDC populations is two-fold. First, FDC are a likely source of TGFβ production and activation for presentation to B cells during the GCR and second, TGFβ is an important factor in the regulation of FDC survival and cytokine production, thus promoting indirectly the GCR, potentially through the regulation of FDC efficacy but also FDC expansion and contraction throughout the GCR.

## Putative function of α<sub>v</sub>β8 expression by follicular regulatory T cells for TGFβ-mediated regulation of GC B cell responses

Foxp3<sup>+</sup> Treg and Tfr regulate the GC, at least in part *via* TGFβ (36, 53–56, 64, 65). Furthermore, several clues point towards a role for α<sub>v</sub>β8-mediated TGFβ activation in Tfr function. First, analysis of the differentially expressed genes between Tfr and naïve or effector T cells (microarray) reveals a preferential expression of *Itgb8* on Tfr as compared to naïve and effector T cells (61). Next, in 2017, investigating the transcriptional activity of different regulatory T cells population in a peptide antigens model immunization, it was shown that *Itgb8* is expressed on Tfr and that its expression is controlled through the mTORC1 pathway (64, 142). It is

interesting to note that, Jacobsen *et al.* recently proposed that the late-stage contraction of the GC is partially mediated by FoxP3-expressing GC T cells. These cells, in part differentiated from Tfh, display a phenotype closely resembling CD25<sup>-</sup> Tfr (43).

Altogether, these studies suggest that Tfr have the potential to produce, bind and activate cell-bound TGF $\beta$  complexes to control GC B cell responses, potentially in relation with the GC contraction. However, formal demonstration is still pending that  $\alpha_v\beta_8$  expression by Tfr is required for their immunoregulatory function is required. Similarly, determining which cells (GC B cells and/or Tfh) are directly targeted by Tfr-activated TGF $\beta$  during the GCR remains an open question.

In addition, TGF $\beta$  is, at least partially, required for the development of some Tfr (Table 1). Unfortunately, to this day, there has been no investigation into the mechanisms by which TGF $\beta$  is activated and presented to T cell in the context of Tfr differentiation. Given the importance of these cells in regulating the GCR, especially in the context of autoimmune reactions, a better characterization of the TGF $\beta$ -dependent mechanisms for Tfr differentiation is highly warranted.

## TGF $\beta$ activation in the context of Tfh differentiation

Given the critical role of myeloid cell populations for the activation and presentation of TGF $\beta$  to T cells in the context of induced Treg generation in the mucosal environment (108, 109, 119), a couple of studies have investigated the potential role for cDC and macrophages in presenting TGF $\beta$  to naïve T cells at the T:B border.

Looking at human pediatric tonsils, Schmitt and colleagues, showed that CD11c<sup>+</sup> DC present in the T cell zones near GC were positive for TGF $\beta$  immunofluorescent staining and that T cells in the vicinity showed robust pSmad staining, indicating active TGF $\beta$ R signaling (39). In 2019, it was further demonstrated that human tonsil cDC2 and CD14<sup>+</sup> macrophages are potent inducers of Tfh cells *in vitro*. They further demonstrate that these cells preferentially express OX40L and secrete TGF $\beta$  which is required for CXCL13 production by Tfh. Furthermore, both these subsets express the  $\beta_8$  integrin subunit *ex vivo* and after TLR stimulation




FIGURE 3

Model for the regulation of the GC reaction via the controlled delivery of active TGF $\beta$  by  $\alpha_v\beta_8$  integrin. (A) TGF $\beta$  signaling in FDC may promote their survival and cytokine production and as such indirectly supports the GC reaction. In addition, given their expression of TGF $\beta$  activating molecule, mucosal FDC themselves are a likely source of active TGF $\beta$  (123). Furthermore, Albright and colleagues provided evidence that TGF $\beta$  produced by FDC may be important for LZ to DZ trafficking of GC B cells. The mechanisms involved may include  $\alpha_v\beta_8$  although formal demonstration is lacking (31). (B) Evidence suggests that TGF $\beta$  is required, at least in certain situations, for the induction of Foxp3<sup>+</sup> follicular T cells (Tfr) (36, 44, 67). In addition, Tfr may be able to regulate GC B cell responses via  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  (61, 64, 142). (C)  $\alpha_v\beta_8$ -mediated activation of TGF $\beta$  by follicular myeloid populations has been involved in the induction of Tfh (143). Additionally, a model in which TGF $\beta$  promotes the survival of Tfh in the context of immunization have been proposed (35). In these contexts, the source of active TGF $\beta$  is still currently unknown.

(143). These data suggest that TGF $\beta$  activation by cDC2 and macrophages could potentially regulate Tfh differentiation and/or activation in the human GC environment. Given that cDC2 are found in the T cell zone while CD14<sup>+</sup> macrophages are rather found in the B cell follicle, the latter colocalizing with Tfh, the authors suggest that these two subsets have a complementary and sequential role in Tfh induction. cDC2 could be responsible for priming pre-Tfh in the T cell zone while CD14<sup>+</sup> macrophages might instruct the maturation or survival of Tfh in the B cell follicle.

The controversial requirement of TGF $\beta$  for Tfh induction and the demonstration that myeloid cells might promote Tfh induction through TGF $\beta$  activation warrants further investigation. It is enticing to consider that TGF $\beta$  requirement for mouse (and potentially human) Tfh development might be different depending on the type of immunogen and associated danger signals being presented to T cells and that the context (tissue localization, time, co-stimulation, environmental context, etc.) in which TGF $\beta$  is activated and presented to naïve T cells, pre-Tfh and mature Tfh might be key to resolve these seemingly contradictory roles of TGF $\beta$  on follicular T cell biology.

## Conclusion

From the current literature, it is clear that there is more to TGF $\beta$ -mediated control of B cells than just simply IgA CSR and inhibition of B cell proliferation, as TGF $\beta$  also regulates Ig CSR decision, Ig production, Ag responsiveness and, in GC B cells, the LZ to DZ transition. The differentiation and/or survival of follicular cell populations (Tfh, Tfr and FDC), which are integral to the GCR, also appear to be regulated by TGF $\beta$ , although some controversy remains. Additionally, accumulating evidence points toward a role for  $\alpha_v\beta_8$  in licensing these populations, along with cDC, for TGF $\beta$ -mediated regulation of the GCR. However, TGF $\beta$  activation in the context of the regulation of humoral responses, is severely under-investigated and the complete sequence of events allowing the secretion, activation, and presentation of TGF $\beta$  for each of these cell types remains elusive. This search is made particularly challenging by the constant discovery of additional layers of regulation of the GCR by TGF $\beta$ . For instance, the very recent demonstration that intratumoral CD8<sup>+</sup> T cells could recruit Tfh to the tumor microenvironment *via* CXCL13 secretion following TGF $\beta$  stimulation (147) or the discovery of novel immune populations, such as follicular regulatory Innate Lymphoid Cells (ILCfr), which appear to inhibit the GCR in a TGF $\beta$ -dependent manner (37).

It is critical to consider that the regulation of the humoral response involves the interaction of B cells with many different cell types, each of which finely regulates the different steps of the GC reaction (SHM, CSR, proliferation, differentiation, ...). Similar to what has been described in the gut, where the

induction of Treg and/or inflammatory Th17 cells is dependent on the context encoded at the T cell/DC synapse, the regulation of humoral responses by TGF $\beta$  might be dependent on the context in which TGF $\beta$  is activated and presented to B cells or to cells regulating B cell responses (FDC, Tfh, Tfr) (78, 148). Because of the production of TGF $\beta$  in a latent form, several important parameters must be taken into account to fully understand the importance of TGF $\beta$  in regulating the GC: 1/the conditions of production and sequestration of latent TGF $\beta$  in the extracellular matrix or at the cell surface; 2/the cells involved and the mechanism by which latent TGF $\beta$  is activated; 3/the necessity of TGF $\beta$  activation and presentation through cell/cell contact or paracrine secretion, which implies to consider the context of TGF $\beta$  delivery, i.e. in conjunction with the presentation of other membrane-bound or soluble factor; 4/the time of delivery, both in the context of individual cell activation and differentiation (e.g. on naïve CD4 T cells, pre-Tfh or fully differentiated Tfh) and in the context of the whole GC (i.e. during initiation, the effector phase or the contraction of the GC)) and 5/which tissue (lymphoid organs vs tissues, mucosal vs peripheral compartment,...) and inflammatory context (steady state vs infection, bacterial vs viral vs fungal vs parasitic infection,...) in which TGF $\beta$  is delivered. We believe that only when taking all these considerations of location, timing of cellular interaction and mechanisms of TGF $\beta$  activation into account that we will be able to reconcile and resolve the ambiguities of the multiple roles of TGF $\beta$  in the regulation of humoral responses in particular in the context of the GCR.

This field of research is rapidly expanding; further studies will be critical in establishing the precise mechanisms of TGF $\beta$ -mediated regulation of the GCR, which will surely inform new avenues of treatment for autoimmunity, graft rejection, allergy ... Interestingly, animal studies have previously demonstrated that  $\alpha_v\beta_8$  targeting is possible to treat inflammatory disorders such as encephalitis or lung inflammation and to overcome tumor immune evasion (92, 116, 149, 150). Altogether, these studies suggest that targeting  $\alpha_v\beta_8$ -mediated TGF $\beta$  activation could represent a valid strategy to develop immunotherapies for human inflammatory pathologies and cancer (151). Whether this could be extended to humoral responses and could open new therapeutic interventions to re-establish tolerance in antibody-mediated disorders (autoimmunity, allergy, graft rejection) or on the contrary to boost and/or optimize humoral immune responses (in the context of immunotherapy or vaccination) is currently unknown and remains to be investigated.

It is important to note that both B and T cell physiological developments are dependent on TGF $\beta$  (19, 152). While it is not discussed in this review, we cannot exclude that some of the experimental setups and evidence described here involve, at least partly, mechanisms linked to development of B or T cell progenitors.

## Author contributions

ST et HP wrote the manuscript. ST designed the figures. HP supervised the study. All authors contributed to the article and approved the submitted version.

## Funding

HP is supported by the Agence Nationale de la Recherche (ANR-20-CE15-0015). ST was supported by a PhD fellowship from the French Ministry of Higher Education and the Cole Foundation Postdoctoral Fellowship.

## Acknowledgments

We would like to thank Dr Thierry Defrance and Prof Olivier Thaumat for many fruitful discussions and critical reading of the manuscript. Servier Medical Art illustrations

(under Creative Commons Attribution 3.0 Unported License) were used and modified to make some of the figures in this article (<https://smart.servier.com/>).

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Post N, Eddy D, Huntley C, Van Schalkwyk MCI, Shrotri M, Leeman D, et al. Antibody response to SARS-CoV-2 infection in humans: A systematic review. *PLoS One* (2020) 15:e0244126. doi: 10.1371/journal.pone.0244126
- Chvatal-Medina M, Mendez-Cortina Y, Patiño PJ, Velilla PA, Rugeles MT. Antibody responses in COVID-19: A review(2021) (Accessed April 18, 2022).
- Etzioni A, Ochs HD. The hyper IgM syndrome—an evolving story. *Pediatr Res* (2004) 56:519–25. doi: 10.1203/01.PDR.0000139318.65842.4A
- Yong PFK, Salzer U, Grimbacher B. The role of costimulation in antibody deficiencies: ICOS and common variable immunodeficiency. *Immunol Rev* (2009) 229:101–13. doi: 10.1111/j.1600-065X.2009.00764.x
- Mesin L, Ersching J, Victora GD. Germinal center b cell dynamics. *Immunity* (2016) 45:471–82. doi: 10.1016/j.immuni.2016.09.001
- Crotty S. T Follicular helper cell biology: A decade of discovery and diseases. *Immunity* (2019) 50:1132–48. doi: 10.1016/j.immuni.2019.04.011
- Sage PT, Sharpe AH. T Follicular regulatory cells in the regulation of b cell responses. *Trends Immunol* (2015) 36:410–8. doi: 10.1016/j.it.2015.05.005
- Sage PT, Sharpe AH. T Follicular regulatory cells. *Immunol Rev* (2016) 271:246–59. doi: 10.1111/imr.12411
- Aguzzi A, Kranich J, Krautler NJ. Follicular dendritic cells: origin, phenotype, and function in health and disease. *Trends Immunol* (2014) 35:105–13. doi: 10.1016/j.it.2013.11.001
- Zhang Y, Garcia-Ibanez L, Toellner K-M. Regulation of germinal center b-cell differentiation. *Immunol Rev* (2016) 270:8–19. doi: 10.1111/imr.12396
- Moses HL, Roberts AB, Derynck R. The discovery and early days of TGF- $\beta$ : A historical perspective. *Advanced Online Article Cold Spring Harb Perspect Biol* (2016) 8. doi: 10.1101/cshperspect.a021865
- Kelly A, Houston SA, Sherwood E, Casulli J, Travis MA. Chapter four - regulation of innate and adaptive immunity by TGF $\beta$ . In: FW Alt, editor. *Advances in immunology*. Elsevier, Amsterdam, Netherlands: Academic Press (2017). p. 137–233. doi: 10.1016/bs.ai.2017.01.001
- Sanjabi S, Oh SA, Li MO. Regulation of the immune response by TGF- $\beta$ : From conception to autoimmunity and infection. *Cold Spring Harb Perspect Biol* (2017) 9:a022236. doi: 10.1101/cshperspect.a022236
- Hinck AP, Mueller TD, Springer TA. Structural biology and evolution of the TGF- $\beta$  family. *Cold Spring Harb Perspect Biol* (2016) 8:a022103. doi: 10.1101/cshperspect.a022103
- Li MO, Flavell RA. TGF- $\beta$ : A master of all T cell trades. *Cell* (2008) 134:392–404. doi: 10.1016/j.cell.2008.07.025
- Stebegg M, Kumar SD, Silva-Cayetano A, Fonseca VR, Linterman MA, Graca L. Regulation of the germinal center response. *Front Immunol* (2018) 9:2469. doi: 10.3389/fimmu.2018.02469
- Toellner KM, Gulbranson-Judge A, Taylor DR, Sze DM, MacLennan IC. Immunoglobulin switch transcript production *in vivo* related to the site and time of antigen-specific b cell activation. *J Exp Med* (1996) 183:2303–12. doi: 10.1084/jem.183.5.2303
- Roco JA, Mesin L, Binder SC, Nefzger C, Gonzalez-Figueroa P, Canete PF, et al. Class-switch recombination occurs infrequently in germinal centers. *Immunity* (2019) 51:337–50.e7. doi: 10.1016/j.immuni.2019.07.001
- Tamayo E, Alvarez P, Merino R. TGF $\beta$  superfamily members as regulators of b cell development and function—implications for autoimmunity. *Int J Mol Sci* (2018) 19:3928. doi: 10.3390/ijms19123928
- Kehrl JH, Roberts AB, Wakefield LM, Jakowlew S, Sporn MB, Fauci AS. Transforming growth factor beta is an important immunomodulatory protein for human b lymphocytes. *J Immunol* (1986) 137:3855–60.
- Petit-Koskas E, Génot E, Lawrence D, Kolb J-P. Inhibition of the proliferative response of human b lymphocytes to b cell growth factor by transforming growth factor-beta. *Eur J Immunol* (1988) 18:111–6. doi: 10.1002/eji.1830180117
- Kehrl JH. Transforming growth factor- $\beta$ : An important mediator of immunoregulation. *Int J Cell Cloning* (1991) 9:438–50. doi: 10.1002/stem.1991.5530090502
- Warner GL, Ludlow JW, Nelson DA, Gaur A, Scott DW. Anti-immunoglobulin treatment of murine b-cell lymphomas induces active transforming growth factor beta but pRB hypophosphorylation is transforming growth factor beta independent. *Cell Growth Differ* (1992) 3:175–81.
- Czac BB, Roes J. TGF- $\beta$  receptor controls b cell responsiveness and induction of IgA. *In Vivo. Immun* (2000) 13:443–51. doi: 10.1016/S1074-7613(00)00044-3
- Holder MJ, Knox K, Gordon J. Factors modifying survival pathways of germinal center b cells. glucocorticoids and transforming growth factor- $\beta$ , but not cyclosporin a or anti-CD19, block surface immunoglobulin-mediated rescue from apoptosis. *Eur J Immunol* (1992) 22:2725–8. doi: 10.1002/eji.1830221037

26. Spender LC, O'Brien DI, Simpson D, Dutt D, Gregory CD, Allday MJ, et al. TGF- $\beta$  induces apoptosis in human b cells by transcriptional regulation of BIK and BCL-X l. *Cell Death Differentiation* (2009) 16:593–602. doi: 10.1038/cdd.2008.183
27. Lømo J, Blomhoff HK, Beiske K, Stokke T, Smeland EB. TGF-beta 1 and cyclic AMP promote apoptosis in resting human b lymphocytes. *J Immunol* (1995) 154:1634–43.
28. Amoroso SR, Huang N, Roberts AB, Potter M, Letterio JJ. Consistent loss of functional transforming growth factor  $\beta$  receptor expression in murine plasmacytomas. *Proc Natl Acad Sci* (1998) 95:189–94. doi: 10.1073/pnas.95.1.189
29. Cerutti A. The regulation of IgA class switching. *Nat Rev Immunol* (2008) 8:421–34. doi: 10.1038/nri2322
30. Stavnezer J, Kang J. The surprising discovery that TGF $\beta$  specifically induces the IgA class switch. *J Immunol* (2009) 182:5–7. doi: 10.4049/jimmunol.182.1.5
31. Albright AR, Kabat J, Li M, Raso F, Reboldi A, Muppidi JR. TGF $\beta$  signaling in germinal center b cells promotes the transition from light zone to dark zone. *J Exp Med* (2019) 216:2531–45. doi: 10.1084/jem.20181868
32. Coffman R, Leberman D, Shrader B. Transforming growth factor beta specifically enhances IgA production by lipopolysaccharide-stimulated murine b lymphocytes. *J Exp Med* (1989) 170:1039–44.
33. Sonoda E, Matsumoto R, Hitoshi Y, Ishii T, Sugimoto M, Araki S, et al. Transforming growth factor  $\beta$  induces IgA production and acts additively with interleukin 5 for IgA production. *J Immunol* (1989) 182:14–9.
34. Marshall HD, Ray JP, Laidlaw BJ, Zhang N, Gawande D, Staron MM, et al. The transforming growth factor beta signaling pathway is critical for the formation of CD4 T follicular helper cells and isotype-switched antibody responses in the lung mucosa. *eLife* (2015) 4:e04851. doi: 10.7554/eLife.04851
35. McCarron MJ, Marie JC. TGF- $\beta$  prevents T follicular helper cell accumulation and b cell autoreactivity. *J Clin Invest* (2014) 124(10):4375–86. doi: 10.1172/JCI76179
36. Miles B, Miller SM, Folkvord JM, Kimball A, Chamanian M, Meditz AL, et al. Follicular regulatory T cells impair follicular T helper cells in HIV and SIV infection. *Nat Commun* (2015) 6:8608. doi: 10.1038/ncomms9608
37. O'Connor MH, Muir R, Chakhtoura M, Fang M, Moysi E, Moir S, et al. A follicular regulatory innate lymphoid cell population impairs interactions between germinal center tfh and b cells. *Commun Biol* (2021) 4:1–13. doi: 10.1038/s42003-021-02079-0
38. Turner JA, Stephen-Victor E, Wang S, Rivas MN, Abdel-Gadir A, Harb H, et al. Regulatory T cell-derived TGF- $\beta$ 1 controls multiple checkpoints governing allergy and autoimmunity. *Immunity* (2020) 53:1202–14.e6. doi: 10.1016/j.immuni.2020.10.002
39. Schmitt N, Liu Y, Bentebibel S-E, Munagala I, Bourdery L, Venuprasad K, et al. TGF- $\beta$  co-opts STAT3-STAT4 signaling to promote human T follicular helper cell differentiation. *Nat Immunol* (2014) 15:856–65. doi: 10.1038/ni.2947
40. Chaurio RA, Anadon CM, Lee Costich T, Payne KK, Biswas S, Harro CM, et al. TGF- $\beta$ -mediated silencing of genomic organizer SATB1 promotes tfh cell differentiation and formation of intra-tumoral tertiary lymphoid structures. *Immunity* (2022) 55:115–28.e9. doi: 10.1016/j.immuni.2021.12.007
41. Nurieva RI, Chung Y, Hwang D, Yang XO, Kang HS, Ma L, et al. Generation of T follicular helper cells is mediated by interleukin-21 but independent of T helper 1, 2, or 17 cell lineages. *Immunity* (2008) 29:138–49. doi: 10.1016/j.immuni.2008.05.009
42. Suto A, Kashiwakuma D, Kagami S, Hirose K, Watanabe N, Yokote K, et al. Development and characterization of IL-21-producing CD4+ T cells. *J Exp Med* (2008) 205:1369–79. doi: 10.1084/jem.20072057
43. Jacobsen JT, Hu W, R. Castro TB, Solem S, Galante A, Lin Z, et al. Expression of Foxp3 by T follicular helper cells in end-stage germinal centers. *Science* (2021) 373:eabe5146. doi: 10.1126/science.abe5146
44. Li L, Yang S-H, Yao Y, Xie Y-Q, Yang Y-Q, Wang Y-H, et al. Block of both TGF- $\beta$  and IL-2 signaling impedes neuropilin-1+ regulatory T cell and follicular regulatory T cell development. *Cell Death Dis* (2016) 7:e2439. doi: 10.1038/cddis.2016.348
45. Takahashi H, Kanno T, Nakayama S, Hirahara K, Sciumè G, Muljo SA, et al. TGF- $\beta$  and retinoic acid induce the microRNA miR-10a, which targets bcl-6 and constrains the plasticity of helper T cells. *Nat Immunol* (2012) 13:587–95. doi: 10.1038/ni.2286
46. Park S-M, Kim S, Choi J-S, Hur D-Y, Lee W-J, Lee M-S, et al. TGF- $\beta$  inhibits fas-mediated apoptosis of a follicular dendritic cell line by down-regulating the expression of fas and caspase-8: Counteracting role of TGF- $\beta$  on TNF sensitization of fas-mediated apoptosis. *J Immunol* (2005) 174:6169–75. doi: 10.4049/jimmunol.174.10.6169
47. Cho W, Kim Y, Kim J, Park S, Park D, Kim B-C, et al. Suppressor of cytokine signaling 1 is a positive regulator of TGF- $\beta$ -induced prostaglandin production in human follicular dendritic cell-like cells. *J Immunol* (2015) 194:4287–97. doi: 10.4049/jimmunol.1401615
48. Choe J, Park J, Lee S, Kim Y-M, Jeoung D. Opposing roles of TGF- $\beta$  in prostaglandin production by human follicular dendritic cell-like cells. *Mol Immunol* (2016) 76:41–8. doi: 10.1016/j.molimm.2016.06.008
49. Letterio JJ, Roberts AB. REGULATION OF IMMUNE RESPONSES BY TGF- $\beta$ . *Annu Rev Immunol* (1998) 16:137–61. doi: 10.1146/annurev.immunol.16.1.137
50. Bollum LK, Huse K, Oksvold MP, Bai B, Hilden VI, Forfang L, et al. BMP-7 induces apoptosis in human germinal center b cells and is influenced by TGF- $\beta$  receptor type I ALK5. *PLoS One* (2017) 12:e0177188. doi: 10.1371/journal.pone.0177188
51. Wing JB, Sakaguchi S. Foxp3+ treg cells in humoral immunity. *Int Immunol* (2014) 26:61–9. doi: 10.1093/intimm/dxt060
52. Weingartner E, Golding A. Direct control of b cells by tregs: An opportunity for long-term modulation of the humoral response. *Cell Immunol* (2017) 318:8–16. doi: 10.1016/j.cellimm.2017.05.007
53. Nakamura K, Kitani A, Strober W. Cell contact-dependent immunosuppression by CD4(+)CD25(+) regulatory T cells is mediated by cell surface-bound transforming growth factor beta. *J Exp Med* (2001) 194:629–44. doi: 10.1084/jem.194.5.629
54. Lim HW, Hillsamer P, Banham AH, Kim CH. Cutting edge: Direct suppression of b cells by CD4+CD25+ regulatory T cells. *J Immunol* (2005) 175:4180–3. doi: 10.4049/jimmunol.175.7.4180
55. Eddahri F, Oldenhove G, Denanglaire S, Urbain J, Leo O, Andris F. CD4+CD25+ regulatory T cells control the magnitude of T-dependent humoral immune responses to exogenous antigens. *Eur J Immunol* (2006) 36:855–63. doi: 10.1002/eji.200535500
56. Xu A, Liu Y, Chen W, Wang J, Xue Y, Huang F, et al. TGF- $\beta$ -Induced regulatory T cells directly suppress b cell responses through a noncytotoxic mechanism. *J Immunol* (2016) 196:3631–41. doi: 10.4049/jimmunol.1501740
57. Okamura T, Sumitomo S, Morita K, Iwasaki Y, Inoue M, Nakachi S, et al. TGF- $\beta$ -expressing CD4 + CD25 - LAG3 + regulatory T cells control humoral immune responses. *Nat Commun* (2015) 6:1–14. doi: 10.1038/ncomms7329
58. Tsuchida Y, Sumitomo S, Ishigaki K, Suzuki A, Kochi Y, Tsuchiya H, et al. TGF- $\beta$ 3 inhibits antibody production by human b cells. *PLoS One* (2017) 12:e0169646. doi: 10.1371/journal.pone.0169646
59. Wollenberg I, Agua-Doce A, Hernández A, Almeida C, Oliveira VG, Faro J, et al. Regulation of the germinal center reaction by Foxp3+ follicular regulatory T cells. *J Immunol* (2011) 187:4553–60. doi: 10.4049/jimmunol.1101328
60. Chung Y, Tanaka S, Chu F, Nurieva R, Martinez GJ, Rawal S, et al. Follicular regulatory T (Tfr) cells with dual Foxp3 and Bcl6 expression suppress germinal center reactions. *Nat Med* (2011) 17:983–8. doi: 10.1038/nm.2426
61. Linterman MA, Pierson W, Lee SK, Kallies A, Kawamoto S, Rayner TF, et al. Foxp3+ follicular regulatory T cells control T follicular helper cells and the germinal center response. *Nat Med* (2011) 17:975–82. doi: 10.1038/nm.2425
62. Miles B, Connick E. Control of the germinal center by follicular regulatory T cells during infection. *Front Immunol* (2018) 9:2704. doi: 10.3389/fimmu.2018.02704
63. Fazilleau N, Aloulou M. Several follicular regulatory T cell subsets with distinct phenotype and function emerge during germinal center reactions. *Front Immunol* (2018) 9:1792. doi: 10.3389/fimmu.2018.01792
64. Wing JB, Kitagawa Y, Locci M, Hume H, Tay C, Morita T, et al. A distinct subpopulation of CD25- T-follicular regulatory cells localizes in the germinal centers. *Proc Natl Acad Sci U.S.A.* (2017) 114:E6400–9. doi: 10.1073/pnas.1705551114
65. Sayin I, Radtke AJ, Vella LA, Jin W, Wherry EJ, Buggert M, et al. Spatial distribution and function of T follicular regulatory cells in human lymph nodes. *J Exp Med* (2018) 215:1531–42. doi: 10.1084/jem.20171940
66. Marie JC, Letterio JJ, Gavin M, Rudensky AY. TGF- $\beta$ 1 maintains suppressor function and Foxp3 expression in CD4+CD25+ regulatory T cells. *J Exp Med* (2005) 201:1061–7. doi: 10.1084/jem.20042276
67. Cobb DA, Kim O-K, Golden-Mason L, Rosen HR, Hahn YS. Hepatocyte-derived exosomes promote T follicular regulatory cell expansion during hepatitis c virus infection. *Hepatology* (2018) 67:71–85. doi: 10.1002/hep.29409
68. León B, Bradley JE, Lund FE, Randall TD, Ballesteros-Tato A. FoxP3+ regulatory T cells promote influenza-specific tfh responses by controlling IL-2 availability. *Nat Commun* (2014) 5:3495. doi: 10.1038/ncomms4495
69. Stephen TL, Payne KK, Chaurio RA, Allegranza MJ, Zhu H, Perez-Sanz J, et al. SATB1 expression governs epigenetic repression of PD-1 in tumor-reactive T cells. *Immunity* (2017) 46:51–64. doi: 10.1016/j.immuni.2016.12.015
70. Carlson CM, Turpin EA, Moser LA, O'Brien KB, Cline TD, Jones JC, et al. Transforming growth factor- $\beta$ : Activation by neuraminidase and role in highly pathogenic H5N1 influenza pathogenesis. *PLoS Pathog* (2010) 6:e1001136. doi: 10.1371/journal.ppat.1001136



71. Roberson EC, Tully JE, Guala AS, Reiss JN, Godburn KE, Pociask DA, et al. Influenza induces endoplasmic reticulum stress, Caspase-12-dependent apoptosis, and c-jun n-terminal kinase-mediated transforming growth factor- $\beta$  release in lung epithelial cells. *Am J Respir Cell Mol Biol* (2012) 46:573–81. doi: 10.1165/rncmb.2010.0460OC
72. Imai Y, Yamakawa M. Morphology, function and pathology of follicular dendritic cells. *Pathol Int* (1996) 46:807–33. doi: 10.1111/j.1440-1827.1996.tb03555.x
73. Yamada K, Yamakawa M, Imai Y, Tsukamoto M. Expression of cytokine receptors on follicular dendritic cells. *Blood* (1997) 90:4832–41. doi: 10.1182/blood.V90.12.4832
74. Kim J, Lee S, Kim Y-M, Jeoung D-I, Choe J. Human follicular dendritic cells promote germinal center b cell survival by providing prostaglandins. *Mol Immunol* (2013) 55:418–23. doi: 10.1016/j.molimm.2013.03.019
75. Annes JP, Munger JS, Rifkin DB. Making sense of latent TGF $\beta$  activation. *J Cell Sci* (2003) 116:217–24. doi: 10.1242/jcs.00229
76. Worthington JJ, Klementowicz JE, Travis MA. TGF $\beta$ : a sleeping giant awoken by integrins. *Trends Biochem Sci* (2011) 36:47–54. doi: 10.1016/j.tibs.2010.08.002
77. Robertson IB, Rifkin DB. Regulation of the bioavailability of TGF- $\beta$  and TGF- $\beta$ -Related proteins. *Cold Spring Harb Perspect Biol* (2016) 8:a021907. doi: 10.1101/cshperspect.a021907
78. McEntee CP, Gunaltay S, Travis MA. Regulation of barrier immunity and homeostasis by integrin-mediated transforming growth factor  $\beta$  activation. *Immunology* (2019) 160:139–48. doi: 10.1111/imm.13162
79. Metelli A, Salem M, Wallace CH, Wu BX, Li A, Li X, et al. Immunoregulatory functions and the therapeutic implications of GARP-TGF- $\beta$  in inflammation and cancer. *J Hematol Oncol* (2018) 11:24. doi: 10.1186/s13045-018-0570-z
80. Campbell MG, Cormier A, Ito S, Seed RI, Bondesson AJ, Lou J, et al. Cryo-EM reveals integrin-mediated TGF- $\beta$  activation without release from latent TGF- $\beta$ . *Cell* (2020) 180:1–12. doi: 10.1016/j.cell.2019.12.030
81. Khan SA, Joyce J, Tsuda T. Quantification of active and total transforming growth factor- $\beta$  levels in serum and solid organ tissues by bioassay. *BMC Res Notes* (2012) 5:636. doi: 10.1186/1756-0500-5-636
82. Barczyk M, Carracedo S, Gullberg D. Integrins. *Cell Tissue Res* (2010) 339:269–80. doi: 10.1007/s00441-009-0834-6
83. Yang Z, Mu Z, Dabovic B, Jurukovski V, Yu D, Sung J, et al. Absence of integrin-mediated TGF $\beta$ 1 activation *in vivo* recapitulates the phenotype of TGF $\beta$ 1-null mice. *J Cell Biol* (2007) 176:787–93. doi: 10.1083/jcb.200611044
84. Munger JS, Harpel JG, Giaccotti FG, Rifkin DB. Interactions between growth factors and integrins: latent forms of transforming growth factor-beta are ligands for the integrin alphavbeta1. *Mol Biol Cell* (1998) 9:2627–38. doi: 10.1091/mbc.9.9.2627
85. Ludbrook SB, Barry ST, Delves CJ, Horgan CMT. The integrin alphavbeta3 is a receptor for the latency-associated peptides of transforming growth factors beta1 and beta3. *Biochem J* (2003) 369:311–8. doi: 10.1042/BJ20020809
86. Asano Y, Ihn H, Yamane K, Jinnin M, Tamaki K. Increased expression of integrin  $\alpha$ v $\beta$ 5 induces the myofibroblastic differentiation of dermal fibroblasts. *Am J Pathol* (2006) 168:499–510. doi: 10.2353/ajpath.2006.041306
87. Tatler AL, John AE, Jolly L, Habgood A, Porte J, Brightling C, et al. Integrin  $\alpha$ v $\beta$ 5 mediated TGF- $\beta$  activation by airway smooth muscle cells in asthma. *J Immunol* (2011) 187:6094–107. doi: 10.4049/jimmunol.1003507
88. Sarraza V, Koehler A, Chow ML, Zimina E, Li CX, Kato H, et al. Integrins  $\alpha$ v $\beta$ 5 and  $\alpha$ v $\beta$ 3 promote latent TGF- $\beta$ 1 activation by human cardiac fibroblast contraction. *Cardiovasc Res* (2014) 102:407–17. doi: 10.1093/cvr/cvu053
89. Reed NI, Jo H, Chen C, Tsujino K, Arnold TD, DeGrado WF, et al. The  $\alpha$ v $\beta$ 1 integrin plays a critical *in vivo* role in tissue fibrosis. *Sci Transl Med* (2015) 7:288ra79. doi: 10.1126/scitranslmed.aaa5094
90. Munger JS, Huang X, Kawakatsu H, Griffiths MJD, Dalton SL, Wu J, et al. A mechanism for regulating pulmonary inflammation and fibrosis: The integrin  $\alpha$ v $\beta$ 6 binds and activates latent TGF  $\beta$ 1. *Cell* (1999) 96:319–28. doi: 10.1016/S0092-8674(00)80545-0
91. Aluwihare P, Mu Z, Zhao Z, Yu D, Weinreb PH, Horan GS, et al. Mice that lack activity of  $\alpha$ v $\beta$ 6- and  $\alpha$ v $\beta$ 8-integrins reproduce the abnormalities of Tgfb1- and Tgfb3-null mice. *J Cell Sci* (2009) 122:227–32. doi: 10.1242/jcs.035246
92. Minagawa S, Lou J, Seed RI, Cormier A, Wu S, Cheng Y, et al. Selective targeting of TGF- $\beta$  activation to treat fibroinflammatory airway disease. *Sci Transl Med* (2014) 6:241ra79. doi: 10.1126/scitranslmed.3008074
93. Wang J, Dong X, Zhao B, Li J, Lu C, Springer TA. Atypical interactions of integrin  $\alpha$ v $\beta$ 8 with pro-TGF- $\beta$ 1. *PNAS* (2017), 114(21):201705129. doi: 10.1073/pnas.1705129114
94. Cormier A, Campbell MG, Ito S, Wu S, Lou J, Marks J, et al. Cryo-EM structure of the  $\alpha$ v $\beta$ 8 integrin reveals a mechanism for stabilizing integrin extension. *Nat Struct Mol Biol* (2018) 25:698–704. doi: 10.1038/s41594-018-0093-x
95. Song G, Luo B-H. Atypical structure and function of integrin  $\alpha$ v $\beta$ 8. *J Cell Physiol* (2020) 236(7):4874–87. doi: 10.1002/jcp.30242
96. Mu D, Cambier S, Fjellbirkeland L, Baron JL, Munger JS, Kawakatsu H, et al. The integrin  $\alpha$ v $\beta$ 8 mediates epithelial homeostasis through MT1-MMP-dependent activation of TGF- $\beta$ 1. *J Cell Biol* (2002) 157:493–507. doi: 10.1083/jcb.200109100
97. Dallas SL, Rosser JL, Mundy GR, Bonewald LF. Proteolysis of latent transforming growth factor- $\beta$  (TGF- $\beta$ )-binding protein-1 by osteoclasts: A CELLULAR MECHANISM FOR RELEASE OF TGF- $\beta$  FROM BONE MATRIX\*. *J Biol Chem* (2002) 277:21352–60. doi: 10.1074/jbc.M111663200
98. Liang HPH, Xu J, Xue M, Jackson CJ. Matrix metalloproteinases in bone development and pathology: current knowledge and potential clinical utility. *MNM* (2016) 3:93–102. doi: 10.2147/MNM.S92187
99. Tatti O, Vehviläinen P, Lehti K, Keski-Oja J. MT1-MMP releases latent TGF-beta1 from endothelial cell extracellular matrix *via* proteolytic processing of LTBP-1. *Exp Cell Res* (2008) 314:2501–14. doi: 10.1016/j.yexcr.2008.05.018
100. Sounni NE, Dehne K, van Kempen L, Egeblad M, Affara NI, Cuevas I, et al. Stromal regulation of vessel stability by MMP14 and TGFbeta. *Dis Model Mech* (2010) 3:317–32. doi: 10.1242/dmm.003863
101. Gutiérrez-Fernández A, Soria-Valles C, Osorio FG, Gutiérrez-Abril J, Garabaya C, Aguirre A, et al. Loss of MT1-MMP causes cell senescence and nuclear defects which can be reversed by retinoic acid. *EMBO J* (2015) 34:1875–88. doi: 10.15252/embj.201490594
102. Knapinska AM, Fields GB. The expanding role of MT1-MMP in cancer progression. *Pharmaceuticals* (2019) 12:77. doi: 10.3390/ph12020077
103. Wang R, Zhu J, Dong X, Shi M, Lu C, Springer TA. GARP regulates the bioavailability and activation of TGF $\beta$ . *MBOC* (2012) 23:1129–39. doi: 10.1091/mbc.e11-12-1018
104. Edwards JP, Thornton AM, Shevach EM. Release of active TGF- $\beta$ 1 from the latent TGF- $\beta$ 1/GARP complex on T regulatory cells is mediated by integrin  $\beta$ 8. *J Immunol* (2014) 193:2843–9. doi: 10.4049/jimmunol.1401102
105. Sun L, Jin H, Li H. GARP: a surface molecule of regulatory T cells that is involved in the regulatory function and TGF- $\beta$  releasing. *Oncotarget* (2016) 7:42826–36. doi: 10.18632/oncotarget.8753
106. Liénart S, Merceron R, Vanderaa C, Lambert F, Colau D, Stockis J, et al. Structural basis of latent TGF- $\beta$ 1 presentation and activation by GARP on human regulatory T cells. *Science* (2018) 362:952–6. doi: 10.1126/science.aau2909
107. Travis MA, Reizis B, Melton AC, Masteller E, Tang Q, Proctor JM, et al. Loss of integrin  $\alpha$ v $\beta$ 8 on dendritic cells causes autoimmunity and colitis in mice. *Nature* (2007) 449:361–5. doi: 10.1038/nature06110
108. Paidassi H, Acharya M, Zhang A, Mukhopadhyay S, Kwon M, Chow C, et al. Preferential expression of integrin  $\alpha$ v $\beta$ 8 promotes generation of regulatory T cells by mouse CD103+ dendritic cells. *Gastroenterology* (2011) 141:1813–20. doi: 10.1053/j.gastro.2011.06.076
109. Worthington JJ, Czajkowska BI, Melton AC, Travis MA. Intestinal dendritic cells specialize to activate transforming growth factor- $\beta$  and induce Foxp3+ regulatory T cells *via* integrin  $\alpha$ v $\beta$ 8. *Gastroenterology* (2011) 141:1802–12. doi: 10.1053/j.gastro.2011.06.057
110. Worthington JJ, Kelly A, Smedley C, Bauché D, Campbell S, Marie JC, et al. Integrin  $\alpha$ v $\beta$ 8-mediated TGF- $\beta$  activation by effector regulatory T cells is essential for suppression of T-Cell-Mediated inflammation. *Immunity* (2015) 42:903–15. doi: 10.1016/j.immuni.2015.04.012
111. Luda KM, Joeris T, Persson EK, Rivollier A, Demiri M, Sitnik KM, et al. IRF8 transcription-Factor-Dependent classical dendritic cells are essential for intestinal T cell homeostasis. *Immunity* (2016) 44:860–74. doi: 10.1016/j.immuni.2016.02.008
112. Grueter B, Petter M, Egawa T, Laule-Kilian K, Aldrian CJ, Wuerch A, et al. Runx3 regulates integrin  $\alpha$ E/CD103 and CD4 expression during development of CD4+CD8+ T cells. *J Immunol* (2005) 175:1694–705. doi: 10.4049/jimmunol.175.3.1694
113. Worthington JJ, Klementowicz JE, Rahman S, Czajkowska BI, Smedley C, Waldmann H, et al. Loss of the TGF $\beta$ -activating integrin  $\alpha$ v $\beta$ 8 on dendritic cells protects mice from chronic intestinal parasitic infection *via* control of type 2 immunity. *PLoS Pathog* (2013) 9:e1003675. doi: 10.1371/journal.ppat.1003675
114. Steel N, Faniyi AA, Rahman S, Swietlik S, Czajkowska BI, Chan BT, et al. TGF $\beta$ -activation by dendritic cells drives Th17 induction and intestinal contractility and augments the expulsion of the parasite trichinella spiralis in mice. *PLoS Pathog* (2019) 15:e1007657. doi: 10.1371/journal.ppat.1007657

115. Melton AC, Bailey-Bucktrout SL, Travis MA, Fife BT, Bluestone JA, Sheppard D. Expression of  $\alpha\beta$  integrin on dendritic cells regulates Th17 cell development and experimental autoimmune encephalomyelitis in mice. *J Clin Invest* (2010) 120:4436–44. doi: 10.1172/JCI43786
116. Acharya M, Mukhopadhyay S, Paidassi H, Jamil T, Chow C, Kissler S, et al.  $\alpha\beta$  integrin expression by DCs is required for Th17 cell differentiation and development of experimental autoimmune encephalomyelitis in mice. *J Clin Invest* (2010) 120:4445–52. doi: 10.1172/JCI43796
117. Kudo M, Melton AC, Chen C, Engler MB, Huang KE, Ren X, et al. IL-17A produced by  $\alpha\beta$  T cells drives airway hyper-responsiveness in mice and enhances mouse and human airway smooth muscle contraction. *Nat Med* (2012) 18:547–54. doi: 10.1038/nm.2684
118. Paidassi H, Acharya M, Lacy-Hulbert A. Alpha ( $\alpha$ ) integrins license regulatory T cells to apoptotic cells and self-associated antigens. *Ann New York Acad Sci* (2010) 1209:68–76. doi: 10.1111/j.1749-6632.2010.05783.x
119. Boucard-Jourdin M, Kugler D, Ahanda M-LE, This S, Calisto JD, Zhang A, et al.  $\beta$ 8 integrin expression and activation of TGF- $\beta$  by intestinal dendritic cells are determined by both tissue microenvironment and cell lineage. *J Immunol* (2016) 197(5):1968–78. doi: 10.4049/jimmunol.1600244
120. Chanab D, Boucard-Jourdin M, Paidassi H. Gut inflammation in mice leads to reduction in  $\alpha\beta$  integrin expression on CD103+CD11b- dendritic cells. *J Crohns Colitis* (2017) 11:258–9. doi: 10.1093/ecco-jcc/jjw140
121. Zhang A, Paidassi H, Lacy-Hulbert A, Savill J. Apoptotic cells induce CD103 expression and immunoregulatory function in myeloid dendritic cell precursors through integrin  $\alpha\beta$  and TGF- $\beta$  activation. *PLoS One* (2020) 15: e0232307. doi: 10.1371/journal.pone.0232307
122. Reboldi A, Arnon TI, Rodda LB, Atakilit A, Sheppard D, Cyster JG. IgA production requires b cell interaction with subepithelial dendritic cells in peyer's patches. *Science* (2016) 352:aaf4822. doi: 10.1126/science.aaf4822
123. Suzuki K, Maruya M, Kawamoto S, Sitnik K, Kitamura H, Agace WW, et al. The sensing of environmental stimuli by follicular dendritic cells promotes immunoglobulin a generation in the gut. *Immunity* (2010) 33:71–83. doi: 10.1016/j.immuni.2010.07.003
124. Nakawesi J, This S, Hütter J, Boucard-Jourdin M, Barateau V, Muleta KG, et al.  $\alpha\beta$ 8 integrin-expression by BATF3-dependent dendritic cells facilitates early IgA responses to rotavirus. *Mucosal Immunol* (2020), 14, 1–15. doi: 10.1038/s41385-020-0276-8
125. Ruane D, Chorny A, Lee H, Faith J, Pandey G, Shan M, et al. Microbiota regulate the ability of lung dendritic cells to induce IgA class-switch recombination and generate protective gastrointestinal immune responses. *J Exp Med* (2016) 213:53–73. doi: 10.1084/jem.20150567
126. Wallace CH, Wu BX, Salem M, Ansa-Addo EA, Metelli A, Sun S, et al. B lymphocytes confer immune tolerance via cell surface GARP-TGF- $\beta$  complex. *JCI Insight* (2018) 3:e99863. doi: 10.1172/jci.insight.99863
127. Dedobbeleer O, Stockis J, van der Woning B, Coulie PG, Lucas S. Cutting edge: Active TGF- $\beta$ 1 released from GARP/TGF- $\beta$ 1 complexes on the surface of stimulated human b lymphocytes increases class-switch recombination and production of IgA. *J Immunol Author Choice* (2017) 199:391. doi: 10.4049/jimmunol.1601882
128. Bauché D, Marie JC. Transforming growth factor  $\beta$ : a master regulator of the gut microbiota and immune cell interactions. *Clin Transl Immunol* (2017) 6: e136. doi: 10.1038/cti.2017.9
129. Ihara S, Hirata Y, Koike K. TGF- $\beta$  in inflammatory bowel disease: a key regulator of immune cells, epithelium, and the intestinal microbiota. *J Gastroenterol* (2017) 52:777–87. doi: 10.1007/s00535-017-1350-1
130. Saito A, Horie M, Nagase T. TGF- $\beta$  signaling in lung health and disease. *Int J Mol Sci* (2018) 19:2460. doi: 10.3390/ijms19082460
131. Huai G, Markmann JF, Deng S, Rickert CG. TGF- $\beta$ -secreting regulatory b cells: unsung players in immune regulation. *Clin Transl Immunol* (2021) 10:e1270. doi: 10.1002/cti2.1270
132. Gros MJ, Naquet P, Guinamard RR. Cell intrinsic TGF- $\beta$ 1 regulation of b cells. *J Immunol* (2008) 180:8153–8. doi: 10.4049/jimmunol.180.12.8153
133. Heath WR, Kato Y, Steiner TM, Caminschi I. Antigen presentation by dendritic cells for b cell activation. *Curr Opin Immunol* (2019) 58:44–52. doi: 10.1016/j.coi.2019.04.003
134. Kato Y, Steiner TM, Park H-Y, Hitchcock RO, Zaid A, Hor JL, et al. Display of native antigen on cDC1 that have spatial access to both T and b cells underlies efficient humoral vaccination. *J Immunol* (2020) 205:1842–56. doi: 10.4049/jimmunol.2000549
135. Tezuka H, Ohteki T. Regulation of IgA production by intestinal dendritic cells and related cells. *Front Immunol* (2019) 10:1891. doi: 10.3389/fimmu.2019.01891
136. Cong Y, Feng T, Fujihashi K, Schoeb TR, Elson CO. A dominant, coordinated T regulatory cell-IgA response to the intestinal microbiota. *Proc Natl Acad Sci U.S.A.* (2009) 106:19256–61. doi: 10.1073/pnas.0812681106
137. Wang L, Ray A, Jiang X, Wang J-Y, Basu S, Liu X, et al. T Regulatory cells and b cells cooperate to form a regulatory loop that maintains gut homeostasis and suppresses dextran sulfate sodium-induced colitis. *Mucosal Immunol* (2015) 8:1297–312. doi: 10.1038/mi.2015.20
138. Gribonika I, Eliasson DG, Chandode RK, Schön K, Strömberg A, Bemark M, et al. Class-switch recombination to IgA in the peyer's patches requires natural thymus-derived tregs and appears to be antigen independent. *Mucosal Immunol* (2019) 12:1268–79. doi: 10.1038/s41385-019-0202-0
139. Kawamoto S, Maruya M, Kato LM, Suda W, Atarashi K, Doi Y, et al. Foxp3 + T cells regulate immunoglobulin a selection and facilitate diversification of bacterial species responsible for immune homeostasis. *Immunity* (2014) 41:152–65. doi: 10.1016/j.immuni.2014.05.016
140. Rescigno M. Tfr cells and IgA join forces to diversify the microbiota. *Immunity* (2014) 41:9–11. doi: 10.1016/j.immuni.2014.06.012
141. Munoz O, Fend F, de Beaumont R, Husson H, Astier A, Freedman AS. TGFbeta-mediated activation of Smad1 in b-cell non-hodgkin's lymphoma and effect on cell proliferation. *Leukemia* (2004) 18:2015–25. doi: 10.1038/sj.leu.2403485
142. Xu L, Huang Q, Wang H, Hao Y, Bai Q, Hu J, et al. The kinase mTORC1 promotes the generation and suppressive function of follicular regulatory T cells. *Immunity* (2017) 47:538–51.e5. doi: 10.1016/j.immuni.2017.08.011
143. Durand M, Walter T, Pirnay T, Naessens T, Gueguen P, Goudot C, et al. Human lymphoid organ cDC2 and macrophages play complementary roles in T follicular helper responses. *J Exp Med* (2019) 216:1561–81. doi: 10.1084/jem.20181994
144. Kitamura H, Cambier S, Somanath S, Barker T, Minagawa S, Markovics J, et al. Mouse and human lung fibroblasts regulate dendritic cell trafficking, airway inflammation, and fibrosis through integrin  $\alpha\beta$ 8-mediated activation of TGF- $\beta$ . *J Clin Invest* (2011) 121:2863–75. doi: 10.1172/JCI45589
145. Henderson NC, Arnold TD, Katamura Y, Giacomini MM, Rodriguez JD, McCarty JH, et al. Targeting of  $\alpha\beta$  integrin identifies a core molecular pathway that regulates fibrosis in several organs. *Nat Med* (2013) 19:1617–24. doi: 10.1038/nm.3282
146. Cording S, Wahl B, Kulkarni D, Chopra H, Pezoldt J, Buettner M, et al. The intestinal micro-environment imprints stromal cells to promote efficient treg induction in gut-draining lymph nodes. *Mucosal Immunol* (2014) 7:359–68. doi: 10.1038/mi.2013.54
147. Niogret J, Berger H, Rebe C, Mary R, Ballot E, Truntzer C, et al. Follicular helper-T cells restore CD8+-dependent antitumor immunity and anti-PD-L1/PD-1 efficacy. *J Immunother Cancer* (2021) 9:e002157. doi: 10.1136/jitc-2020-002157
148. Nolte M, Margadant C. Controlling immunity and inflammation through integrin-dependent regulation of TGF- $\beta$ . *Trends Cell Biol* (2020) 30:49–59. doi: 10.1016/j.tcb.2019.10.002
149. Stockis J, Liénart S, Colau D, Collignon A, Nishimura SL, Sheppard D, et al. Blocking immunosuppression by human tregs *in vivo* with antibodies targeting integrin  $\alpha\beta$ 8. *PNAS* (2017) 114:E10161–8. doi: 10.1073/pnas.1710680114
150. Takasaka N, Seed RI, Cormier A, Bondesson AJ, Lou J, Elattma A, et al. Integrin  $\alpha\beta$ 8-expressing tumor cells evade host immunity by regulating TGF- $\beta$  activation in immune cells. *JCI Insight* (2018) 3:e122591. doi: 10.1172/jci.insight.122591
151. Batlle E, Massagué J. Transforming growth factor- $\beta$  signaling in immunity and cancer. *Immunity* (2019) 50:924–40. doi: 10.1016/j.immuni.2019.03.024
152. Oh SA, Li MO. TGF- $\beta$ : Guardian of T cell function. *J Immunol* (2013) 191:3973–9. doi: 10.4049/jimmunol.1301843