



HAL
open science

The Alzheimer's disease risk factor APOE4 drives pro-inflammation in human astrocytes via HDAC-dependent repression of TAGLN3

Philippe Benech

► **To cite this version:**

Philippe Benech. The Alzheimer's disease risk factor APOE4 drives pro-inflammation in human astrocytes via HDAC-dependent repression of TAGLN3. 2022. hal-03854275

HAL Id: hal-03854275

<https://hal.science/hal-03854275v1>

Preprint submitted on 15 Nov 2022


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Extended Data Figure 4.

NF- κ B activity underlies the APOE4-specific proinflammatory phenotype in astrocytes. Cell lysates from [*APOE3/E3*] (ND-1 [3/3]) and their respective [*APOE4/E4*]-Ki isogenic astrocytes (ND-1 Ki [4/4/]), either in unstimulated or stimulated (Stim) conditions, were prepared for cell fractioning. In **a**) representative Western blots to validate the fractioning (Cytosolic and nuclear fractions). Note that the nuclear Histone 3 was almost exclusively detected in the nuclear fraction, as expected. **b**) From the two fractions, different members of the NF- κ B cascade, namely p50 (top), Phosphorylated p65 (P-p65, middle) and p52 (bottom) were analysed by Western blots. Representative immunoblots (left panels) and quantitative analyses (bar charts, right) are shown. Astrocytes from the original ND-1 [*APOE3/E3*] line and two independent isogenic clones [*APOE4/E4*]-Ki were analysed, with 2 independent batches of differentiation for each line. Data are presented as mean \pm SEM using unpaired two-tailed Student's t-Test. In **b**, A.U. stands for arbitrary units (see methods).

Extended Data Fig.5




Extended Data Figure 5.

TAGLN3 stands as a new neuroinflammatory modulator in human astrocytes.

a) Quantification of *TAGLN3* mRNA levels, using qPCR analyses, from [*APOE3/E3*], [*APOE4/E4*]-Ki and [*APOE*-KO] astrocytes. **b-d)** Stimulated [*APOE4/E4*]-Ki astrocytes were treated with 5 or 20 $\mu\text{g/mL}$ of TAGLN3, as recombinant protein (TAGLN3-rec), or with its control vehicle, 24 hours prior to being analysed by Western blot analyses for TAGLN3, IL-6 and IL-8 (**b**). Quantification of Western blot analyses are shown in **c** and **d**. In all different assays, astrocytes from the original ND-1 [*APOE3/E3*] line and three independent isogenic clones [*APOE4/E4*]-Ki and APOE-KO (when applicable) were analysed, with 3 to 7 independent batches of differentiation for each line. Data are presented as mean \pm SEM using one-way ANOVA followed by post hoc Dunnett's test (a, c and d). In **c** and **d**, A.U. stands for arbitrary units (see methods).

Extended Data Fig.6




Extended Data Figure 6.

TAGLN3 downregulation in patient-specific astrocytes carrying the *APOE4* allele

a,b) *TAGLN3* mRNA expression levels from astrocytes derived from hiPSCs of individuals carrying different APOE polymorphisms [*APOE3/E3*], [*APOE3/E4*] or [*APOE4/E4*]. For each genotype, the *TAGLN3* expression of different hiPSC lines and different batches of differentiation were pooled (**a**) or represented in a patient/donor-specific manner (**b**). **c,d)** Western blotting analyses for TAGLN3 from hiPSC-astrocytes from individuals carrying different APOE polymorphisms [*APOE3/E3*], [*APOE3/E4*] or [*APOE4/E4*]. Non-stimulated (Stim-) and stimulated (Stim+) astrocytes were analysed.

Representative Western Blot (**c**) and quantitative analyses (**d**) are shown. For each hiPSC line, 2 to 3 independent batches of differentiation were analysed. Data are presented as mean \pm SEM using one-way ANOVA followed by post hoc Dunnett's test. In **d**, A.U. stands for arbitrary units (see methods).

Extended Data Fig.7



Extended Data Figure 7. TAGLN3 intracellular distribution and a comprehensive gene

interactions map underlying the novel APOE4-TAGLN3 axis. a) Representative western Blot

revealing the presence of TAGLN3 in in the cytosolic (C) and nuclear (N) fractions of [APOE3/E3]

astrocytes. TAGLN3 was barely detected in the membrane (M) fraction and totally absent from the

cytoskeleton (Ck) fraction. ACTIN, VIMEMTIN and HISTONE 3 were used as positive control of the

C, Ck and N fractions respectively. **b)** Gene interactions map revealing TAGLN3 downregulation as a

novel candidate controlling inflammation in an APOE-dependent manner. [APOE3/E3] astrocytes and

their respective isogenic [APOE4/E4]-Ki astrocytes, either nonstimulated (No Stim) or stimulated

(Stim) with a pro-inflammatory cocktail, were prepared for DNA microarray analysis. Raw data were

analysed to identify upregulated and downregulated genes in each experimental condition. Fold

changes (FC) cut-off used for analyses were ≥ 1.5 and ≤ 0.6 for upregulated and downregulated genes,

respectively. Leveraging on those analyses and using PredictSearch® as a powerful text mining

software, we established a gene network linking different genes found dysregulated in an APOE-

isoform dependent manner to identify possible mechanisms at play in the pro-inflammatory phenotype

observed in astrocytes carrying the APOE4 allele. The numbers indicate references (see main and

supplementary references).

Supplementary references (related to Extended data Figure 7)

64. Fan, L., Jaquet, V., Dodd, P. R., Chen, W. & Wilce, P. A. Molecular cloning and characterization of hNP22: a gene up-regulated in human alcoholic brain. *J Neurochem* **76**, 1275–1281 (2001).
65. Feil, S., Hofmann, F. & Feil, R. SM22 α Modulates Vascular Smooth Muscle Cell Phenotype During Atherogenesis. *Circulation Research* **94**, 863–865 (2004).
66. Wamhoff, B. R. *et al.* A G/C Element Mediates Repression of the SM22 α Promoter Within Phenotypically Modulated Smooth Muscle Cells in Experimental Atherosclerosis. *Circulation Research* **95**, 981–988 (2004).
67. Dresselhaus, E. *et al.* Class I HDAC inhibition is a novel pathway for regulating astrocytic apoE secretion. *PLoS ONE* **13**, e0194661 (2018).
68. Prasad, H. & Rao, R. The Na⁺/H⁺ exchanger NHE6 modulates endosomal pH to control processing of amyloid precursor protein in a cell culture model of Alzheimer disease. *J Biol Chem* **290**, 5311–5327 (2015).