



**HAL**  
open science

# Recent Advances in C(sp<sup>3</sup>)–C(sp<sup>3</sup>) and C(sp<sup>3</sup>)–C(sp<sup>2</sup>) Bond Formation through Cathodic Reactions: Reductive and Convergent Paired Electrolyses

Aurélie Claraz, Géraldine Masson

► **To cite this version:**

Aurélie Claraz, Géraldine Masson. Recent Advances in C(sp<sup>3</sup>)–C(sp<sup>3</sup>) and C(sp<sup>3</sup>)–C(sp<sup>2</sup>) Bond Formation through Cathodic Reactions: Reductive and Convergent Paired Electrolyses. ACS Organic & Inorganic Au, 2021, 2 (2), pp.126 - 147. 10.1021/acscorginorgau.1c00037 . hal-03854744

**HAL Id: hal-03854744**

**<https://hal.science/hal-03854744v1>**

Submitted on 16 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Recent Advances in $C(sp^3)-C(sp^3)$ and $C(sp^3)-C(sp^2)$ Bond Formation through Cathodic Reactions: Reductive and Convergent Paired Electrolyses

Aur lie Claraz\* and G raldine Masson\*


Cite This: *ACS Org. Inorg. Au* 2022, 2, 126–147

Read Online

ACCESS |

Metrics &amp; More

Article Recommendations



**ABSTRACT:** The formation of  $C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bonds is one of the major research goals of synthetic chemists. Electrochemistry is commonly considered to be an appealing means to drive redox reactions in a safe and sustainable fashion and has been utilized for C–C bond-forming reactions. Compared to anodic oxidative methods, which have been extensively explored, cathodic processes are much less investigated, whereas it can pave the way to alternative retrosynthetic disconnections of target molecules and to the discovery of new transformations. This review provides an overview on the recent achievements in the construction of  $C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bonds via cathodic reactions since 2017. It includes electrochemical reductions and convergent paired electrolyses.

**KEYWORDS:** *electrosynthesis, electrochemical synthesis, C–C bond formation, electroreduction, convergent paired electrolysis, cathodic reaction, carbon-centered radical, electrocatalysis*

## 1. INTRODUCTION

$C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bonds are inherently omnipresent in most natural and synthetic organic molecules, rendering their formation a central task in organic synthesis. In recent decades, intensive efforts have been devoted to develop efficient C–C bond formation, among which radical-based transformations offer a powerful alternative approach to the traditional polar reactions. In this context, electrochemistry has recently gained renewed interest as a safe and environmentally friendly technique to generate carbon-centered radicals. By carefully controlling the amount of charge and the cell potential, substrates or catalysts can be selectively activated to prohibit unwanted side reactions such as overoxidations or over-reductions that can easily occur with more classical chemical reaction conditions.<sup>1–6</sup> In electrosynthesis, three different approaches can be distinguished: (i) net oxidative anodic electrolysis whereby hydrogen evolution is the most frequent counter-reaction; (ii) net reductive cathodic electrolysis whereby a sacrificial reductant (typically a simple amine, the anode itself, or in few cases the solvent) is oxidized to counterbalance

the process; (iii) redox neutral paired electrolysis during which two desirable half-reactions are taking place at both electrodes. As such, in each case, the use of hazardous or difficult to handle redox reagents is avoided. Compared to electro-oxidative methods, electrochemical synthesis employing a cathode material as the working electrode remains relatively less studied.<sup>7</sup> However, several successful approaches have recently been accomplished for the building of  $C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bonds through cathodic reductions, allowing significant improvement of the previously established procedures as well as the discovery of new transformations. Although electrochemical organic synthesis was comprehensively surveyed by Baran et al. in 2017,<sup>1</sup> this review describes recent

Received: October 25, 2021

Revised: December 2, 2021

Accepted: December 3, 2021

Published: December 22, 2021



advances in  $C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bond-forming reactions through cathodic transformations that have been reported since July 2017. It covers cathodic reductive processes as well as convergent paired electrolyses, whereby both cathode and anode materials are working electrodes. To aid the reader, electrochemical parameters—including cell type (divided or undivided), mode of electrolysis (constant current CCE or constant potential CPE electrolysis), and electrode compositions—are graphically depicted (Figure 1a). For instance, Figure

1a, and each part is divided according to the initial electrolytic event that has enabled the generation of either a  $C(sp^3)$  radical, carbanion, or radical anion.


## 2. $C(sp^3)-C(sp^3)$ BOND FORMATION

### 2.1. Reductive Decarboxylative Process

Carboxylic acids are ubiquitous in natural products and constitute therefore an ideal source of versatile  $C(sp^3)$ -centered radicals with structural diversity upon decarboxylative processes. Oxidative decarboxylation of aliphatic acids on platinum anodes is a well-known electrochemical transformation (Kolbe electrolysis) wherein the resulting alkyl radicals dimerize (homocoupling) to form a new  $C-C$  bond.<sup>8</sup> Even though heterodimerizations between two different carboxylic acids (mixed Kolbe electrolysis) or radical cyclizations have been reported,<sup>9,10</sup> the synthetic potential of the alkyl radical for the construction of  $C(sp^3)-C(sp^3)$  bonds was not fully leveraged in such oxidative conditions. Recently, the electroreductive decarboxylation of redox-active *N*-hydroxyphthalimide (NHP)<sup>11</sup> esters became a complementary method to produce alkyl radicals. While primary alkyl iodides and bromides exhibit two-electron reduction at  $-2.3$  and  $-2.6$  V vs SCE in DMF, respectively,<sup>12,13</sup> NHP esters derived from primary carboxylic acids feature single-electron reduction potential at  $-1.6$  V vs SCE in DMF.<sup>7</sup> As such, the desired alkyl radical can be generated under relatively mild reaction conditions upon a rapid  $N-O$  fragmentation/decarboxylation sequence. In 2020, Wang et al. reported a decarboxylative Giese-type reaction between NHP esters **1** and various Michael esters **2** under constant potential electrolysis in an undivided cell using two inexpensive graphite electrodes.<sup>14,15</sup> A stoichiometric amount of the reductant Hantzsch ester (**3**) was necessary to reduce the cell potential, thus minimizing side reactions. The reaction conditions tolerated a wide range of NHP esters derived from primary, secondary, and tertiary carboxylic acids and various Michael acceptors such as vinyl sulfones and  $\alpha,\beta$ -unsaturated ketones, esters, and amides. As depicted in Scheme 1, mechanistic investigations suggested that the transformation was initiated by cathodic reduction of NHP ester **1** to form the radical anion **5**. After a fragmentation/decarboxylation sequence, the resulting alkyl radical **6** underwent Giese-type addition to the Michael acceptor **2** and generated radical **7**. Meanwhile, anodic oxidation of the HE **3** furnished radical cation **8**. Subsequent deprotonation would trigger SET between **9** and **7** followed by protonation of anion **11** to yield the desired product **4**.

Interestingly, this transformation has been extended to the deoxygenative fragmentation of *N*-phthalimidoyl oxalates **12** derived from secondary and tertiary alcohols (Scheme 2). Herein, a hydrogen atom transfer (HAT) between in situ generated radical **7** and HE **3** was suggested at the end of the proposed mechanism.<sup>15</sup>


Recently, Baran, Blackmond, and Reisman et al. disclosed the radical coupling between redox active NHP esters **13** derived from primary carboxylic acids and aldehydes **14**.<sup>16</sup> Notably, only a catalytic loading of  $CrCl_3$  was employed contrary to a previous chemical transformation which required 4 equiv of toxic chrome reagents.<sup>17</sup> This electrochemical reaction took place in an undivided cell using an inexpensive Al-based sacrificial anode and a Ni-foam cathode. Scheme 3 displays the proposed mechanism supported by a combination of detailed kinetic investigations, cyclic voltammetry, and UV-visible spectroelec-




**Figure 1.** Cell notation and interpretation of the color coding used in the proposed mechanisms.

1b (left) displays a constant current electrolysis performed in an undivided cell with a carbone cathode and a magnesium anode, whereas Figure 1b (right) corresponds to a constant potential electrolysis in a divided cell with two platinum electrodes. In agreement with the proposed mechanism, a color code is employed to distinguish each fragments of the products. As such, turquoise, pink, and purple are utilized for the reactants that undergo a single-electron transfer (SET) reduction, a SET oxidation, and the attack of either a radical or a carbanion, respectively. Orange color is used for other reactants in multicomponent reactions. For example, in Figure 1c (left), radical  $R^1\bullet$  **II** is produced at the cathode by SET reduction of reagent **I** and adds to the alkene **III** to form the purposed  $C-C$  bond. A second SET reduction of the resulting radical **IV** furnished anion **V**, which reacts with the electrophile  $E^+$ . As such,  $R^1$  is drawn in turquoise, the alkene **III** in purple, and the electrophile  $E^+$  in orange. Conversely, in Figure 1c (right), radicals  $R^1\bullet$  **II** is formed at the cathode by SET reduction of reagent **I**, whereas radical  $R^2\bullet$  **VIII** is generated at the anode by SET oxidation of reagent **VII**. Radicals **II** and **VIII** combine to form the  $C-C$  bond and yield product **IX**. As such,  $R^1$  and  $R^2$  are drawn in turquoise and pink, respectively.  $C(sp^3)-C(sp^3)$  and  $C(sp^3)-C(sp^2)$  bond-forming processes are discussed sequen-


## Scheme 1. Electroreductive Decarboxylative Giese-Type Reaction




## proposed mechanism




## selected examples




## Scheme 2. Electroreductive Deoxygenative Giese-Type Reaction



## proposed final step




## selected examples




trochemistry. Cr(III) was initially reduced at the cathode to produce Cr(II) which in turn reduced NHP ester **13** to furnish alkyl radical **16**. Coordination with Cr(II) followed by addition to the aldehyde **14** gave rise to Cr alkoxide species **18**. Subsequent transmetalation with oxophilic triethylsilyl chloride


## Scheme 3. Electroreductive Radical Coupling between NHP Esters and Aldehydes



## proposed mechanism



## selected examples



(TESCI) regenerated Cr(III) catalyst and delivered protected aldols **19** in moderate to good yields. Aromatic as well as aliphatic aldehydes were well-tolerated.


## 2.2. SET Reduction of Alkyl Halides

Alkyl halides stand out as another important source of alkyl radicals via SET reduction. However, the direct electrolysis of alkyl halides requires highly reducing potentials that could lead to undesired side reactions. As such, indirect processes have been recently developed.

Wilden et al. have described a Giese-type addition of alkyl iodides **20** to activated alkenes **21** under constant potential electrolysis in a divided cell at mildly reducing potential ( $-1.0$  V vs Ag quasi-reference electrode) in a buffered acidic semi-aqueous solvent system.<sup>18</sup> Water and—intriguingly—a catalytic amount of molecular oxygen were essential for the success of this reductive transformation. Primary, secondary, and tertiary alkyl iodides were well-tolerated. However, poor yield was achieved with alkyl bromides. Mechanistic studies identified in situ generated hydroxyl radical and ozone as mediators to generate the  $\text{C}(\text{sp}^3)$  radicals **24** from alkyl iodides **20**. The reaction was initiated by the oxidation of water in the anodic chamber to the hydroxyl radical, which in turn reacted with molecular oxygen to generate ozone. The latter slowly diffused in the cathodic chamber to activate the alkyl iodide **20** via unstable trioxide intermediate **23** that subsequently fragments into the corresponding alkyl radical **24**,  $\text{IO}^\bullet$  radical, and molecular oxygen. Under the slightly acidic and aerobic conditions, a second

pathway produced even more alkyl radicals **24** via two interlocking cycles and the generation of unstable I(II) species **25** from hydroxyl radical  $\text{OH}^\bullet$ . Addition of nucleophilic radical **24** to activated alkenes **21** generated radical **26** which underwent cathodic SET reduction and protonation to deliver the desired product **22** (Scheme 4).<sup>19</sup>


#### Scheme 4. Reactive Oxygen Species-Mediated Electroreductive Giese-Type Addition of Alkyl Iodides to Electron-Deficient Alkenes




##### proposed mechanism

##### cathodic chamber


##### pathway A:





##### pathway B:




##### then




##### anodic chamber



#### Scheme 5. Ni-Electrocatalyzed Reductive Cyclization




##### proposed mechanism



Ni complexes can be employed as electrocatalysts.<sup>20</sup> For instance, recently, Toste and Chang et al. have shown that the new Ni complex **30** featuring a redox-active pentapyridine ligand could activate alkyl iodide **28** toward the synthesis of cyclopentane **31** in the presence of the halogen atom donor **29**.<sup>21</sup> Based on synthetic, electroanalytical, and computational studies, it was proposed that the electroactive reduced Ni complex has its excess spin delocalized throughout the  $\pi$ -system of the redox-active ligand, allowing the activation of the alkyl iodide **29** via outer-sphere electron transfer. The resulting alkyl radical **32** rapidly recombined with the Ni complex to form the  $[\text{LNi}(\text{R})]$  species. The latter reduced another equivalent of **28** to regenerate the  $\text{Ni}^+$  complex and form radical **32**. Subsequent cyclization and trapping by **29** produced iodo-containing cyclopentane **31** (Scheme 5).

Condon et al. published an electrocatalytic reductive Giese-type addition of simple alkyl bromides **34** to activated alkenes **35** with low to moderate yields using an in situ generated cobalt-Salen complex as the catalyst.<sup>22</sup> The reaction was performed under simple reaction conditions using constant current electrolysis in an undivided cell with a cheap nickel grid cathode and an iron-based sacrificial anode. The test of other sacrificial anodes resulted in lower yields, indicating a putative role of in situ generated iron salts. Better yields were obtained with the secondary and tertiary alkyl bromides compared to the primary ones (Scheme 6).


#### Scheme 6. Co-Electrocatalyzed Reductive Giese-Type Addition of Alkyl Bromides to Activated Alkenes




The intermolecular electrochemical difunctionalization of alkenes is an elegant method to access molecular complexity in a rapid and sustainable fashion. Most of the previous methods rely on electro-oxidative processes.<sup>23-25</sup> The development of electroreductive transformations appears as an attractive means to allow different functionalizations. In 2020, Lin et al. established an elegant electroreductive regioselective carbofor-

mylation and carbocarboxylation of styrene derivatives using secondary and tertiary alkyl bromides **37** as alkyl radical precursors and cheap dimethylformamide and carbon dioxide as electrophiles.<sup>26</sup> A simple undivided cell was employed with carbon graphite cathode and Mg-based sacrificial anode under constant current electrolysis. Cyclic voltammetry analysis and control experiments suggested a radical/polar crossover mechanism wherein SET reduction of alkyl bromide **37** delivered an alkyl radical **41** which then added regioselectively to the alkene **38**. Subsequent reduction of the resulting radical **42** to anion **43** followed by addition to the electrophile delivered the dicarboxylated adducts **39/40** in good yields. It is also worth noting that this strategy has been extended to the anti-Markovnikov hydroalkylation of alkenes using acetonitrile as proton source (Scheme 7).

### Scheme 7. Electroreductive Carboformylation and Carbocarboxylation of Alkenes



#### proposed mechanism




### 2.3. Generation of Fluoroalkyl Radicals


The incorporation of a trifluoromethyl group onto target molecules can dramatically impact their physical, chemical, and biological properties. Electrochemistry represents an appealing approach to introduce such a group.<sup>27</sup> While most of the electrochemical methods utilized an oxidative process with  $\text{CF}_3\text{SO}_2\text{Na}$  (Langlois reagent) as trifluoromethyl reagent,<sup>28</sup> the trifluoromethylation involving cathodic processes for  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  bond formation has been only recently described. Wang et al. reported on the use of IMDN- $\text{SO}_2\text{CF}_3$  **45** for the cathodic generation of a trifluoromethyl radical.<sup>29</sup> This approach enabled the synthesis of trifluoromethyl-containing isoquinoline-1,3-diones and oxindoles **47** via a redox-neutral trifluoromethylation/cyclization sequence of activated alkenes **46**. The proposed mechanism started with the SET reduction of **45** followed by the regioselective addition of the resulting trifluoromethyl radical **48** to the alkene **46**. Further radical cyclization on the aromatic ring followed by anodic oxidation and deprotonation provided the desired functionalized heterocycles **47** through a linear paired electrolysis (Scheme 8).

Very recently, Hu et al. developed an electrochemical hydrodifluoroalkylation of conjugated alkenes **52** by cathodic reduction of difluoroalkyl sulfones **51**.<sup>30</sup> The reaction took place


### Scheme 8. Electrochemical Construction of Trifluoromethyl-Containing Heterocycles with IMDN- $\text{SO}_2\text{CF}_3$



#### proposed mechanism



#### selected examples




in an undivided cell under simple constant current electrolysis using cheap carbon graphite electrodes in the presence of triethylamine. Difluoromethyl radical as well as various difluoroalkyl radicals could be efficiently introduced to provide the difluoroalkyl-containing compounds in moderate to high yields. Control experiments supported a radical/polar crossover mechanism. As such, difluoroalkyl radical **54** was initially generated at the cathode and regioselectively added to the alkene **52** to form the new radical **55**. A subsequent SET reduction and protonation yielded the target product **53**. The oxidation of triethylamine at the cathode counterbalanced the overall process (Scheme 9).

### 2.4. SET Reduction of Ketones


Electrochemical reduction of ketones is a powerful tool to generate ketyl radicals which might undergo further reduction to the corresponding carbanions and nucleophilic addition, direct radical addition, or dimerization depending on the reaction conditions. It constitutes therefore a highly attractive synthetic method for  $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$  bond formation.

In 2019, Kise et al. published a C2-functionalization of chromones **57** via an electroreductive coupling of various diaryl ketones **58** in the presence of an excess of trimethylsilyl chloride in a divided cell under constant current electrolysis.<sup>31</sup> Moderate to high yields were achieved. However, no reaction took place with electron-rich 4,4'-dimethoxybenzophenone as the partner due to its higher reduction potential (absolute value) in comparison to that of the chromone. Based on cyclic voltammetry analysis, the authors proposed a mechanism, as depicted in Scheme 10. The reaction was initiated by two consecutive SET reductions of diaryl ketone **58** in the presence of trimethylsilyl chloride to generate carbanion **60**. Subsequent


## Scheme 9. Electroreductive Hydrodifluoroalkylation of Conjugated Alkenes




proposed mechanism




## Scheme 10. C2-Functionalization of Chromones

cathodic chamber: Pt(-); 57 and 58, Et<sub>3</sub>N, TMSCl, nBu<sub>4</sub>ClO<sub>4</sub>, THFanodic chamber: Pt(+); Et<sub>3</sub>NOTs, DMF

proposed mechanism



selected examples




nucleophilic conjugate addition to the chromone 57 delivered 2-substituted chromone 59. Interestingly, the functionalization of 3-methylchromone and 3-arylchromone produced selectively the *cis*-adducts (Scheme 10). Soon after, the authors extended this procedure to the C2-functionalizations of 4-quinolones.<sup>32</sup>


Baran and Minter et al. reported an electroreductive coupling between aliphatic ketones 61 and unactivated alkenes 62 for the synthesis of tertiary alcohols 63 in good yields.<sup>33</sup> The electrolysis was performed in an undivided cell using a zinc-based sacrificial anode and tin cathode. According to kinetic studies and voltammetry analysis, the transformation was initiated by SET reduction of the ketone 61 to form the ketyl radical anion 64, which underwent nucleophilic addition of the alkene 62. This step would be facilitated by the binding of the reactants to the tin cathode. A second SET reduction to dianion 66 followed by protonation formed the tertiary alcohol 63. The authors

highlighted the importance of the sacrificial Zn anode which would facilitate the SET events by providing Zn<sup>2+</sup> (Scheme 11).


## Scheme 11. Electroreductive Coupling between Ketones and Unactivated Alkenes



proposed mechanism




selected examples



One year later, Noël et al. described a divergent paired electrochemical transformation of furfural 67 allowing simultaneously the access to 2(*5H*)-furanone 68 in the anodic chamber and furfuryl alcohol 69 and mostly hydrofuroin 70 in the cathodic part. Interestingly, the electrolysis was performed in water in a divided-cell flow microreactor (Scheme 12).<sup>34</sup>


## Scheme 12. Divergent Paired Electrolysis of Furfural




Peters et al. designed an electrocatalyst 72 comprising a redox cobaltocene and a tethered Brønsted base to enable the reductive pinacol coupling of acetophenone 71. The electrolysis was performed at a constant current potential using boron-doped diamond working electrode. In the presence of a Brønsted acid, the electrocatalyst favored a proton-coupled electron transfer (PCET) event leading to the neutral radical 75. In contrast, a direct reduction on the electrode in the absence of this electrocatalyst required a higher bias, resulting in adsorption and passivation issues and furnished the desired pinacol product 73 in lower yield. Also of note, a simple cobaltocene electrocatalyst was totally inefficient to deliver the pinacol product due to the competing hydrogen evolution reaction.

Such observations underpinned the crucial role of both the cobaltocene electrocatalyst and the tethered Brønsted base (Scheme 13).<sup>35</sup>

### Scheme 13. Electrocatalytic Reductive Pinacol Coupling of Acetophenone




#### proposed mechanism



Recently, Zhang et al. investigated the electroreductive pinacol coupling of aromatic carbonyl compounds and found that carbon fiber paper was optimal as the working electrode.<sup>36</sup> Xu et al. studied the electroreductive cross-coupling of benzaldehyde **76** and furfural **77** on Cu and Pb surfaces. They demonstrated that Cu featured a higher selectivity for cross-coupling product **79** (heterodimerization), whereas Pb favored furfural coupling **78b** (homodimerization). This difference of selectivity was explained by a much stronger adsorption of benzaldehyde on Pb compared to furfural (Scheme 14).<sup>37</sup>

### Scheme 14. Electroreductive Cross-Coupling of Furfural and Benzaldehyde



### 2.5. SET Reduction of Alkenes


Alkenes feature very negative reduction potential, making it difficult to generate the corresponding radical anion. The electrohydrodimerization of acrylonitrile ( $E_p \approx -2$  V in H<sub>2</sub>O vs SCE)<sup>38</sup> to produce adiponitrile, a precursor of Nylon-6,6, is one of the most well-known industrial synthetic electrochemical transformation. Optimizing such a process by identifying the key factors influencing the selectivity toward adiponitrile is worth studying. Ramachandran et al. developed a mathematical model to elucidate the optimal concentration, voltage, and current as long as the effect of mass transfer coefficients between electrolyte and electrodes.<sup>39</sup> Modestino et al. used a systematic

approach to study such parameters.<sup>40</sup> Their investigations have shown that the selectivity of the process toward the formation of adiponitrile was governed by mass transport. Indeed, the formation of propionitrile as the byproduct is favored at high current densities when the reactant concentration in the electrical double layer (EDL) is low. On the other hand, oligomers are formed at low current densities when the reactant concentration in the EDL is high. As such, maintaining intermediate concentration of reactants in the EDL at high current densities is key to enhance the performance of this process. To do so, Modestino et al. combined pulsed electrolysis with artificial intelligence.<sup>41</sup> With pulsed organic electrosynthesis, the composition of the EDL is periodically renewed, enabling an effective process at high current densities, and the diffusion and migration rates of reactants, products, electrolytes, and intermediates are better controlled. Such techniques improved the selectivity by 325% and adiponitrile production rate by 30%.

### 2.6. Regioselective Radical Silylation of an Alkene

The addition of a radical to an alkene generates a C(sp<sup>3</sup>) radical primed for further transformations. Radical silylation of alkenes is a reliable strategy to produce valuable organosilicon molecules with various potential applications in materials science and medicinal chemistry. In this context, Lin et al. developed an elegant new synthetic way to generate silyl radicals via the electroreduction of chlorosilanes **80**.<sup>42</sup> They originally applied this strategy to the vicinal disilylation of alkenes and further extended it to the cyclization of styrene **81** bearing a tethered leaving group, allowing the formation of a C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond by a tandem process. The electrolysis was performed in an undivided cell using a graphite working electrode and a Mg-based sacrificial anode under constant current electrolysis in THF. Mechanistic studies suggested that the reaction was initiated by SET reduction of chlorosilane **80**, which would be facilitated by magnesium salts arising from the oxidation of the sacrificial anode. Subsequent regioselective addition to the alkene **81** enabled the construction of a Si–C bond and the generation of C(sp<sup>3</sup>) radical **85**. A second SET reduction followed by intramolecular nucleophilic substitution delivered the silylated pentacycle **82** in moderate yield (Scheme 15).

### Scheme 15. Tandem Si–C and C–C Bond Formation via Cathodic Generation of Trimethylsilyl Radical






### 3. C(sp<sup>3</sup>)-C(sp<sup>2</sup>) BOND FORMATION

#### 3.1. Reductive Decarboxylative Process

The transition-metal-catalyzed electrochemical process is an efficient method for constructing C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bonds. In 2018, Bio et al. developed a Ni-catalyzed electroreductive coupling between NHP esters **87** and aryl iodides **88**, enabling the formation of a new C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bond.<sup>43</sup> According to the authors, the matching between the reduction potentials of the electrocatalyst and the reactants allowed both the direct decarboxylative cathodic reduction of NHP esters **87** to the corresponding C(sp<sup>3</sup>) radical **91** and the reduction of Ni(I) to Ni(0). Subsequent oxidative addition of the aryl halide **88** and interception of the alkyl radical **91** afforded Ni(III) complex **93**. Further reductive elimination delivered the coupling product **89** and closed the catalytic cycle. Triethylamine was employed as a sacrificial reductant, and a divided cell was required to prevent the anodic oxidation of in situ generated low-valent Ni species. Improved performances were achieved under continuous flow conditions, allowing an easy scale-up (Scheme 16). One year

**Scheme 16. Ni-Catalyzed Electroreductive Cross-Coupling between NHP Esters and Aryl Iodides**




later, Loren et al. designed a similar transformation during which NHP esters are in situ generated from the corresponding carboxylic esters by premixing with tetramethyluronium hexafluorophosphate. Using a Zn-based sacrificial anode, their procedure had the advantage to employ a simple undivided cell.<sup>44</sup>

Minisci-type reaction is an efficient method for the formation of C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bonds, through radical addition of nucleophilic carbon radicals on electron-deficient azacycles.

In 2019, Zeng et al. published a Ni-catalyzed redox-neutral cross-coupling between NHP esters **94** and quinaxolin-2-ones **95**. A Ni(II) complex was used as the electrocatalyst in an undivided cell with a carbon felt anode and a nickel foam cathode. Contrary to the work reported above, the authors proposed here that the Ni(II) species would promote the indirect reduction of NHP esters **94** to generate the alkyl radical **97** (path A). The latter would add to quinaxolinone **95** via complex **98**. Subsequent deprotonation and oxidation would lead to functionalized quinaxolinone **96** and release the electrocatalyst. An alternative mechanism involving the direct reduction of NHP esters **94** was also proposed (path B). Oxidation of triethylamine took place at the anode (Scheme 17).<sup>45</sup> Afterward, Wang et al. reported a similar transformation under electrocatalyst- and triethylamine-free reaction conditions in an undivided cell with two graphite electrodes.<sup>46</sup>


**Scheme 17. Ni-Catalyzed Electroreductive Cross-Coupling between NHP Esters and Quinaxolinones**



In 2019, Lei et al. explored an electrochemical decarboxylative Minisci-type alkylation of azaarenes **101** using NHP esters **100** as alkyl radical precursors, providing the corresponding functionalized heterocycles **102** in moderate to high yields.<sup>47</sup> The electrolysis was performed in an undivided cell employing two simple graphite felt electrodes and *p*-toluene sulfonic acid as an acidic additive. Based on cyclic voltammetry and RPE experiments, the authors suggested a mechanism involving a linear paired electrolysis. As depicted in Scheme 15, the transformation started by the direct cathodic decarboxylative generation of alkyl radicals **103** from NHP esters **100**. Subsequent regioselective radical addition to protonated azacycles **105** furnished radical cation **106**, which underwent

deprotonation and anodic oxidation to deliver the desired products **102** (Scheme 18).


**Scheme 18. Electrochemical Decarboxylative Minisci-Type Alkylation of Azaarenes with NHP Esters**



### 3.2. Reductive Deaminative Process

During their investigations on electrochemical Minisci-type reaction, Lei et al. employed the Katritzky *N*-cyclohexylpyridinium salt **108** as the alkyl radical precursor for the C2-functionalization of 4-methylquinoline **109** through a cathodic deaminative process (Scheme 19).<sup>48</sup>

**Scheme 19. Electrochemical Deaminative Minisci-Type Alkylation of Quinoline **109** with Kratritzky Salt **108****




### 3.3. Generation of Fluoroalkyl Radicals

In 2018, Studer et al. disclosed a radical trifluoromethylation of isonitriles **111** to build CF<sub>3</sub>-containing phenanthridine **113**.<sup>48</sup> The reaction was initiated by the cathodic reduction of Togni reagent **112** to form trifluoromethyl radical **115**. After radical addition to isonitrile **111** and cyclization, the resulting cyclohexadienyl radical **117** is deprotonated by in situ generated *ortho*-iodobenzoate **114**, leading to radical anion **119**. The latter would be a sufficient oxidant species to enable SET, with Togni reagent **112** delivering the desired heterocycle **113** and another

trifluoromethyl radical **115**. As such, this radical chain mechanism allowed the transformation to proceed with a catalytic amount of electric charge (0.075*F*) (Scheme 20).


**Scheme 20. Electroreduction of Togni Reagent-Initiated Trifluoromethylation of Isonitrile**



In 2019, Kappe and Cantillo et al. explored an electro-reductive approach for the radical trifluoromethylation of electron-rich (hetero)cycles using readily available triflyl chloride **120** and triethylamine in an undivided cell with two cheap carbon graphite electrodes. Based on NMR experiments and density functional theory (DFT) calculations, the authors suggested that the trifluoromethyl radical **125** was formed upon SET cathodic reduction of in situ generated triflyltriethylammonium complex **123**. Subsequent addition to aromatic cycle **121** formed radical **126**, which rearomatized via either hydrogen atom abstraction or SET oxidation/deprotonation processes to yield the CF<sub>3</sub>-containing aromatic cycles **122** in modest to good yields. Interestingly, this cathodic event enabled the electrochemical trifluoromethylation of compounds, which would be difficult to achieve under an oxidative procedure with thiacycles or substrates bearing aldehydes (Scheme 21).<sup>49</sup> In 2020, Luo and Nguyen et al. established an alternated current electrolysis for the trifluoromethylation of (hetero)arenes with triflyl chloride in the presence of an inorganic base.<sup>50</sup>

Hisaeda et al. employed a vitamin B<sub>12</sub> derivative as a cobalt-based electrocatalyst for the cathodic perfluoroalkylation of electron-rich (hetero)arenes **128** with cheap perfluoroalkyl iodides **127** under visible light irradiation. The electrolysis was performed in an undivided cell equipped with a reference

Scheme 21. Cathodic Trifluoromethylation of (Hetero)arenes with Triflyl Chloride




electrode to fix the potential at  $-0.8$  V vs Ag/AgCl. Herein, Co(II) was initially reduced to Co(I) species. The latter underwent oxidative addition of perfluoroalkyl iodides 127 to form Co(III)–R<sub>F</sub> complex 130 and subsequently produce perfluoroalkyl radical upon visible light irradiation. Further radical addition to the arene and rearomatization delivered the target functionalized product. A very low catalyst loading was employed, but the slow addition of a large excess of perfluoroalkyl reagent (9 equiv) was required (Scheme 22).<sup>51</sup>


### 3.4. SET Reduction of Alkenes

Alkenes have emerged as an attractive source of alkyl radical anion through an electrochemical reductive process in cross-coupling with CO<sub>2</sub> as the electrophile. Recently, Buckley and Malkov et al. reported an electrochemical highly regioselective hydroxycarboxylation of styrenes 133, giving rise to  $\beta$ -carboxylation products 134 (Scheme 23).<sup>52,53</sup> The electrolysis proceeded in a simple undivided cell under constant current using carbon electrodes and triethanolamine (TEOA) as a proton source. This protocol is compatible with a wide range of substrates including mono-, di-, and trisubstituted alkenes. Based on cyclic voltammetry and control experiments, the proposed mechanism involved an electrochemical chemical reduction of the alkene 133 on the cathode, followed by SET reduction to the corresponding radical anion 135, subsequent carboxylation to regioselectively form the C–C bond by favoring the formation of benzylic radical 136, and further SET reduction and protonation. However, due to the small difference of reduction potential between carbon dioxide and styrene ( $E_{1/2} = -2.21$  V and  $-2.58$  V vs SCE in DMF, respectively), an alternative mechanism involving the direct reduction of carbon dioxide to the corresponding radical anion  $\cdot\text{CO}_2^-$  and subsequent addition to the styrene to form the common intermediate 136 could not be ruled out (Scheme 20). This procedure has been subsequently applied to the selective  $\alpha,\delta$ -hydrocarboxylation

Scheme 22. Co-Electrocatalyzed Perfluoroalkylation of Arenes with Perfluoroalkyl Iodides under Light Irradiation



Scheme 23. Electroreductive Hydrocarboxylation of Styrenes




of conjugated dienes<sup>53</sup> and to the selective  $\beta$ -carboxylation of  $\alpha,\beta$ -unsaturated esters.<sup>54</sup>

Wang and Zhang et al. reported the electrochemical  $\beta$ -carboxylation of  $\alpha,\beta$ -unsaturated ketones using silica nanowires as the working electrode under light irradiation. This specific electrode would absorb light energy to reduce the consumption of electrical energy and enhance the reaction efficiency.<sup>55</sup>

Zhou et al. developed a regioselective defluorinative/carboxylation of  $\alpha$ -trifluoromethyl alkenes **138** under constant current electrolysis with two platinum electrodes.<sup>56</sup> Herein, the anion **140** resulting from the second cathodic SET reduction underwent rapid fluoride elimination via an E1CB-type mechanism instead of protonation leading to valuable *gem*-difluoroalkenes **139** in good yields. Notably, a wide range of  $\alpha$ -CF<sub>3</sub> alkenes **138** could be efficiently employed, including those with  $\alpha$ -aryl/alkyl/alkynyl substituents and trisubstituted alkenes (Scheme 24).


**Scheme 24. Electroreductive Carboxylation of  $\alpha$ -CF<sub>3</sub> Alkenes for the Synthesis of *gem*-Difluoroalkenes**



Cyanopyridines are often used as partners in a radical–radical coupling reaction to access pyridines, which are found in a broad variety of biologically active molecules. Findlater et al. described an efficient regioselective hydroypyridylation of electron-deficient alkenes **142** with 4-cyanopyridine **143**.<sup>57</sup> This electroreductive decyanative alkylation reaction was facilitated by a catalytic amount of Ni(acac)<sub>2</sub>. The authors suggested two putative mechanisms involving the initial reduction of the Michael acceptor to the corresponding radical anion followed by either radical addition to Ni-activated 4-cyanopyridine or radical cross-coupling with in situ generated 4-cyanopyridyl radical anion (Scheme 25).


Very recently, Jiang and Xu et al. employed  $\alpha,\beta$ -unsaturated thioesters **145** as electron-deficient alkenes for a similar hydroypyridylation reaction under Ni-free electrolysis. Interestingly,  $\beta$ -aryl- $\beta$ -alkyl-substituted thioesters were compatible substrates, allowing the construction of quaternary carbons  $\alpha$

**Scheme 25. Electroreductive Hydroypyridylation of Electron-Deficient Alkenes**



to pyridines. Contrary to the work described above, cyclic voltammetry experiments strongly supported the cathodic reduction of both reaction partners **145** and **143** and further radical combination. The oxidation of dimethylsulfoxide (DMSO) was suggested to be the counter-reaction occurring at the anode (Scheme 26).<sup>58</sup>

**Scheme 26. Electroreductive Hydroypyridylation of  $\alpha,\beta$ -Unsaturated Alkenes**




### 3.5. SET Reduction of C=X Bonds (X = O, NR)


Rovis and Lehnher et al. demonstrated an elegant electro-synthetic access to quaternary carbon centers bearing a challenging primary or secondary amino group.<sup>59</sup> Their approach relied on the reductive coupling between iminium salts **147** and cyanoazaarenes **148**. A simple undivided cell was employed under constant current electrolysis. A zinc-based cathode featuring a high overpotential for hydrogen evolution was employed to preclude the loss of H<sup>+</sup>. Cyclic voltammetry analyses and DFT calculations supported a mechanism involving cathodic generation of both transient  $\alpha$ -amino radical **150** from iminium **147** via a SET reduction and persistent neutral radical **151** from cyanoazaarene **148** via a PCET event. Further radical cross-coupling afforded the *ipso*-substituted products **149** with moderate to excellent yields. The PCET process allowed a better matching between the reduction potential of both reactive partners but imposed unavoidably the use of N-heterocycles (Scheme 27).

Similarly, an electroreductive coupling between carbonyl compounds **153** and cyano(aza)arenes **154** has been disclosed by Xia and Yang et al.<sup>60</sup> Aromatic and aliphatic aldehydes as well as aromatic ketones were suitable substrates for this *ipso*-substitution transformation, delivering the corresponding secondary and tertiary alcohols **155** in high yields. DABCO was employed as a sacrificial reductant. Herein, the higher reduction potential (absolute value) of the carbonyl compounds compared to that of the iminium salts matched more easily with those of the cyano(hetero)arenes. As such, unlike Rovis's work, a PCET process was not required and a radical–radical cross-coupling between cathodically generated ketyl radical **156** and persistent radical anion **157** was proposed. Accordingly, azaarenes were not mandatory, making electron-poor hydrocarbon-based cyanoarenes, such as 1,4-dicyanobenzene, additional successful coupling partners (Scheme 28). A similar coupling between ketones or aldehydes and 1,4-dicyanoarenes was demonstrated by Findlater, Zhang, and Xu et al. The latter

Scheme 27. Electroreductive Coupling between Iminium Salts and Cyanoazaarenes



Scheme 28. Electroreductive Coupling between Carbonyl Compounds and Cyano(aza)arenes




proposed an alternative mechanism involving reduction of the 1,4-dicyanoarene to the corresponding radical anion 157 with further nucleophilic addition to the aldehyde, second SET reduction, and decyanation.<sup>61</sup>

### 3.6. Generation of a Vinyl Organometallic Species and Further Addition to Aldehydes

Recently, Baran, Blackmond, and Reisman et al. developed a new electrochemical Nozaki–Hiyama–Kishi (NHK) reaction between vinyl halides 158 and aldehydes 159.<sup>16</sup> The cathodic reduction of Cr(III) avoids the use of superstoichiometric amount of reducing agents, which are traditionally required to render this transformation catalytic in Cr. Although early electrochemical NHK couplings exhibited limited substrate scope (only one example of aliphatic aldehyde) and issues due to difficult setup and the use of expensive electrodes,<sup>62–64</sup> this new protocol features operationally simple reaction conditions: undivided cell, cheap Al-based sacrificial anode, and Ni-foam cathode and potentiostatic electrolysis without a reference electrode. Choosing Cp<sub>2</sub>ZrCl<sub>2</sub> as the oxophilic additive to trap the alkoxide at this end of the catalytic cycle was key for the success of this transformation. A broad substrate scope was

demonstrated with aromatic as well as aliphatic aldehydes using a catalytic amount of NiCl<sub>2</sub> and CrCl<sub>2</sub>. Also noteworthy is an asymmetric version of this transformation using a chiral ligand (Scheme 29).

Scheme 29. Electrocatalytic NHK Reaction




Durandetti et al. have developed a nickel-catalyzed electroreductive intramolecular carbometalation of alkynes 161 and subsequent reaction with aldehydes 162.<sup>65</sup> This domino process formed consecutively a C(sp<sup>2</sup>)–C(sp<sup>2</sup>) bond and a C(sp<sup>2</sup>)–C(sp<sup>3</sup>) bond to furnish various functionalized dihydrobenzofurans, (iso)chromans, indoles, and indanes 163 in good yields. The electrolysis was performed in an undivided cell with cheap nickel foam cathode and a sacrificial aluminum anode under constant current. Cyclic voltammetry studies supported the initial two-electron reduction of Ni(II) catalyst to Ni(0). Subsequent oxidative addition of the aryl iodide 161 and stereoselective 5-*exo*-dig cyclization produced the vinylnickel species 165. Final addition to the aldehyde 162 and transmetalation with anodically generated Al(III) furnished the aluminum alkoxide species 167 and closed the nickel catalytic cycle (Scheme 30).


### 3.7. Two Consecutive SET Reductions for the Generation of a Carbanion or a Nucleophilic Organometallic Species and Further Carboxylation

As mentioned above (see section 3.4), the use of carbon dioxide as a C1 source for the direct access to carboxylic acid is a very appealing approach. Various procedures have recently been established or improved for the carboxylation of cathodically generated carbanion intermediates 170 via two consecutive SET reductions of the substrate 168 and C–X bond cleavage.<sup>66</sup> In some cases, the competitive SET reduction of carbon dioxide could take place at the cathode, resulting in the need of excess electricity or an alternative mechanism. In 2018, He et al. developed a core–shell structured silver nanowire/nitrogen-doped carbon electrocatalyst for the electrocarboxylation of organic bromides.<sup>67</sup> In 2018, Senboku et al. reported the synthesis of mandelic acid derivatives through the electrocarboxylation of the corresponding benzal diacetates.<sup>68</sup> Recently, the same authors described the synthesis of *N*-Boc- $\alpha$ -amino acids in an electroreductive C–S bond cleavage of the *N*-Boc- $\alpha$ -aminosulfones/carboxylation sequence in DMF with a platinum plate cathode and a magnesium rod anode.<sup>69</sup> In 2020,


Scheme 30. Intramolecular Carbonickelation of Alkynes and Subsequent Reaction with Aldehydes



proposed mechanism



selected examples





Guirado et al. reported the synthesis of Ibuprofen and Naproxen via the electrocarboxylation of the corresponding benzyl chlorides. They used an ionic liquid as the solvent/electrolyte system and a silver cathode to decrease the reduction potential value of the organic halide.<sup>70,71</sup> Very recently, Manthiram et al. employed anhydrous magnesium bromide as a source of Mg<sup>2+</sup> cations to inhibit the nucleophilicity of the obtained carboxylate salts, allowing the electrocarboxylation of various benzylic and aliphatic halides to take place with a platinum anode instead of a sacrificial anode (Scheme 31a).<sup>72</sup> Also of note, Senboku et al. utilized a continuous flow system to succeed in the electroreductive carboxylation of in situ generated *N*-aryl aldimines 174, giving rise to *N*-aryl- $\alpha$ -amino acids 175 in moderate to high yields (Scheme 31b).<sup>73</sup>

In 2018, Mei et al. reported an efficient Pd-catalyzed electroreductive carboxylation of allylic esters 176 under constant current electrolysis, giving rise to  $\alpha$ -aryl carboxylic acids 177 with good yields and regioselectivity. A simple undivided cell was used with platinum cathode and magnesium-based sacrificial anode. Cyclic voltammetry measurements supported an initial oxidative addition of allylic ester 176 to in situ generated Pd(0) to form cationic  $\pi$ -allyl Pd(II) complex


Scheme 31. Two Consecutive SET Electroreductions and Subsequent Carboxylations

a) reductive C-X cleavage


b) reduction of *in-situ* generated aldimine

178. The latter equilibrated to the favored branched  $\eta^1$ -allyl Pd(II) species 179, which underwent two cathodic SET reductive events. The resulting nucleophilic Pd(0) species 180 reacted with carbon dioxide to deliver magnesium carboxylate 181 to regenerate Pd(0). The addition of 1 equiv of ethanol was essential for the success of this transformation, but its role was not elucidated. Remarkably, an asymmetric variant was proposed with up to 67% ee using a chiral phosphine (Scheme 32).<sup>74</sup> In 2020, Ackermann et al. reported a similar elegant

Scheme 32. Pd-Catalyzed Electroreductive Carboxylation of Allylic Esters

L<sub>3</sub> = DPPPH = 1,2-bis(diphenylphosphino)benzene

proposed mechanism




carboxylation of allylic chloride using earth-abundant Co(II) electrocatalyst with good yields and low to moderate regioselectivity. A Co(III)/Co(I) catalytic cycle was proposed.<sup>75</sup>

### 3.8. Ni-Electrocatalyzed Alkylation of C(sp<sup>2</sup>) Centers with Alkyl Halides

Ni-electrocatalyzed cross-electrophile coupling represents an attractive strategy for the formation of C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bonds.<sup>76,77</sup> In 2019, Hansen et al. reported a Ni-catalyzed reductive cross-coupling of aryl bromides 183 with alkyl bromides 182.<sup>78</sup> The electrolysis was conducted in acetonitrile in a divided cell with reticulated vitreous carbon (RVC)

electrodes. In contrast with their previous work, simple diisopropylamine was employed as an organic sacrificial reductant instead of a zinc-based sacrificial anode.<sup>79</sup> With careful adjustments of the charge (3.3*F*) and the catalyst loading (7 mol %), yields up to 80% could be achieved at high current (50 mA) with short reaction times on large scales. Importantly, a combination of two ligands was crucial for the success, but the optimal ratio was substrate-dependent (Scheme 33).


**Scheme 33. Ni-Electrocatalyzed Reductive Cross-Coupling of Aryl Bromides with Alkyl Bromides in Acetonitrile**



In 2020, Sevov et al. investigated the same reaction and discovered an electrocatalytic system which was operationally simple (electrolysis at room temperature in an undivided cell), demonstrably scalable (>17 g), and exhibited a broad substrate scope including primary and challenging secondary alkyl bromides with high to excellent yields.<sup>80</sup> The salient feature of this new procedure is the use of Ni complex Ni·MeBPI·OAc, featuring a tridentate ligand MeBPI in combination with the redox shuttle Ni(MeBPI)<sub>2</sub>, which served as an overcharge protector of the active Ni catalytic species. Indeed, intensive cyclic voltammetry analyses and evaluation of the reactivity of isolated organometallic complexes revealed that over-reduction of the on-cycle active Ni catalyst species occurred at a potential ( $E = -2.0$  V vs Fc/Fc<sup>+</sup>) which was only 100 mV higher (absolute value) than the potential required to initiate the cross-coupling ( $E = -1.9$  V vs Fc<sup>+</sup>/Fc). In a constant current electrolysis, this reductive degradation of the catalyst was overcome thanks to the redox shuttle Ni(MeBPI)<sub>2</sub> that could short circuit the electrochemical cell by carrying the electrons from the cathode to the anode when the rate of reduction (current density *i*) was becoming superior to the rate of cross-coupling (*k*<sub>cat</sub>) (Scheme 34).

Mei et al. and Rueping et al. independently developed a Ni-catalyzed electroreductive coupling between aryl halides **186** and alkyl bromides **185**, whereby a chain-walking process took place. After the formation of the Ni(II)(alkyl)(aryl) complex, an iterative β-hydride elimination, migratory insertion, and reductive elimination afforded products in good yields and excellent regioselectivity.<sup>81,82</sup> The electrolyses were performed under mild reaction conditions in an undivided cell. The two

**Scheme 34. Ni-Electrocatalyzed Reductive Cross-Coupling of Aryl Bromides with Alkyl Bromide in the Presence of an Overcharge Protector**




methyl groups at the *ortho* position on the ligands were critical to achieve the desired regioselectivity. The two research groups suggested a Ni(0)/Ni(II) catalytic cycle in which Ni(0) species underwent oxidative addition of aryl halide **186**. According to Rueping et al., the resulting Ni(II) intermediate **189** was reduced at the cathode. SET between the Ni(I)–Ar complex **190** and alkyl bromide **185** generated the corresponding primary alkyl radical **191** and re-formed Ni(II) intermediate **189**. Upon a second SET cathodic event, the trapping of radical **191** by the Ni(I)–Ar complex **190** afforded Ni(II)(alkyl)(aryl) complex **192**. Iterative β-H elimination/migratory insertion sequences took place until the thermodynamically more stable branched benzylic Ni(II) complex **193** was attained. Final reductive elimination eventually led to the desired 1,1-diaryllkane **187** and regenerated the active Ni(0) species (Scheme 35).

Subsequently, Mei et al. harnessed such a chain-walking process for the electroreductive cross-coupling of alkyl bromides **194** and in situ generated alkyl anhydrides to access to dialkyl ketones **196** in good yields and high regioselectivity. A control experiment with a stoichiometric amount of Ni(cod)<sub>2</sub> indicated a putative Ni(0) species as the active catalyst. Based on cyclic voltammetry measurements, the proposed mechanism involved the oxidative addition of in situ generated anhydride to Ni(0) to form acyl Ni(II) complex **199**. Upon cathodic SET reduction, trapping of alkyl radical **198** led to Ni(II) complex **200**. Subsequent β-H elimination, migratory insertion, and reductive elimination yielded dialkyl ketones **196** and the active Ni(0) complex. The latter reduced the alkyl bromide **194** to generate the alkyl radical **198**. A second SET cathodic reduction regenerated the active Ni(0) catalyst species. It is worth mentioning that addition of magnesium salts was critical to the success of this transformation, probably facilitating the oxidative addition of alkyl anhydride (Scheme 36).<sup>83</sup>






Scheme 38. Ni-Electrocatalyzed Redox-Neutral *ortho*-C–H Alkylation of 8-Aminoquinoline

with each other to yield the product. Although very appealing, this redox-neutral approach implies several challenges that need to be leveraged to achieve the desired transformation over side reactions. Most importantly, it requires (i) properly matching the reaction rate of anodic oxidation and cathodic reduction and (ii) ensuring an effective mass transfer of reactive intermediates from the surface of the electrodes to the bulk solution. To succeed in the formation of a  $C(sp^3)$ – $C(sp^2)$  bond via convergent paired electrolysis, two general strategies have recently emerged involving the anodic generation of a  $C(sp^3)$  radical and subsequent combination with (i) a cathodically produced persistent aryl radical anion or (ii) a Ni(II)ArX complex.

**3.9.1. Combination with a Cathodically Produced Persistent Radical Anion.** In 2019, Ye et al. accomplished an electrochemical redox neutral  $\alpha$ -arylation of tertiary amines **210** with benzonitriles **209**.<sup>92</sup> Using TEMPO as electrocatalyst in the presence of 2,6-lutidine, this  $C(sp^3)$ –H bond functionalization proceeded under constant current electrolysis with two RVC electrodes. The best yields were obtained with tertiary arylamines bearing electron-donating groups on the aromatic ring, while electron-withdrawing substituents were not tolerated. Interestingly, the arylation reactions of cyclic tertiary aliphatic amines took place regioselectively at the cyclic position. Cyanoheteroarenes, such as 1,4-dicyanopyridines, were not compatible. From a mechanistic point of view, SET anodic oxidation of TEMPO afforded  $TEMPO^+$  which in turn oxidized tertiary amines **203** to form  $\alpha$ -amino radical **212** in the presence of 2,6-lutidine. Meanwhile, persistent radical anion **214** was generated at the cathode via SET reduction of benzonitrile **209**. Subsequent radical cross-coupling delivered the *ipso*-substitution product **211** upon elimination of cyanide anion and aromatization.  $\alpha$ -Cyanation byproduct **217** was also obtained in a small amount with over anodic oxidation of radical **213** to iminium **216**. It is worth mentioning that an electrolysis on a 10 mmol scale was performed with a three-electrode system, RVC(+)|RVC(-)|RVC(+), to increase the concentration of an  $\alpha$ -amino radical at the anode and the collision frequency of anodic and cathodic radicals (Scheme 39). A similar transformation was realized by Mo and Lu et al. in the absence of TEMPO by employing an engineered microfluidic cell ( $\mu$ RN-eChem) equipped with two glassy carbon electrodes.<sup>93</sup>


Scheme 39. Electrochemical Redox-Neutral  $\alpha$ -Amino C( $sp^3$ ) Arylation

In their continuous effort to develop electrochemical access to diaryl methanol derivatives (see section 3.5), Findlater, Zhang, and Xu et al. disclosed an electrochemical  $\alpha$ -arylation of benzylic alcohols **219** with 1,4-dicyanobenzene **218** using a graphite anode and a nickel cathode.<sup>61</sup> Primary and secondary alcohols could be employed, albeit with lower yields in the latter case. Cyclic voltammety analyses provided evidence of both anodic oxidation of benzylic alcohol **219** and cathodic reduction of 1,4-dicyanobenzene **218**, leading to ketyl radicals **223** and radical anion **221**, respectively. Further radical cross-coupling gave rise to intermediate **224** and eventually to the *ipso*-substitution product **220** upon spontaneous decyanation. This  $C(sp^3)$ –H bond functionalization was performed in the presence of valeraldehyde to trap the toxic cyanide anion (Scheme 40).


An elegant thiol-catalyzed allylic  $C(sp^3)$ –H arylation of alkenes **225** with 1,4-dicyanobenzene **218** was demonstrated by Mo et al. by taking advantage of their  $\mu$ RN-eChem flow cell.<sup>93</sup> Anodic oxidation of triisopropylsilanethiol catalyst **226** generated electrophilic thiyl radical **228** which regioselectively abstracted a hydrogen atom at the allylic position of alkene **226**. The resulting allylic radical **229** coupled with cathodically generated persistent radical anion **221** to build the  $C(sp^2)$ – $C(sp^3)$  bond upon *ipso*-substitution (Scheme 41).

This homemade  $\mu$ RN-eChem flow cell was also applied to the electrochemical decarboxylative and decyanative cross-coupling

### Scheme 40. Electrochemical Redox-Neutral $\alpha$ -Arylation of Benzylic Alcohols




### Scheme 41. Thiol-Electrocatalyzed Redox-Neutral Allylic C-H Arylation



between carboxylic acids **232** and electron-poor cyanobenzenes **231**.<sup>94</sup> Six equivalents of carboxylic acids were necessary to obtain high yields. Tetrabutyl ammonium hydroxide was


employed as a base to generate the carboxylate salts. As such, the reaction medium was sufficiently conductive, making an additional electrolyte unnecessary. The transformation involved the concomitant anodic generation of an alkyl radical via oxidative decarboxylation of carboxylate salts and cathodic generation of a persistent aromatic radical anion via SET reduction of **231** and further radical combination. Such a convergent paired electrolysis precluded the preactivation of carboxylic acids under the form of redox-active esters. Primary and secondary alkyl radicals reacted smoothly, but no example was reported with the tertiary ones (Scheme 42).

### Scheme 42. Electrochemical Redox-Neutral Decarboxylative Arylation of Carboxylic Acids



A last application of this electrochemical flow cell involved the deboronative arylation of organic trifluoroborate salts **234** with 1,4-dicyanobenzene **218**.<sup>94</sup> In a similar process as the decarboxylative arylation reaction mentioned above, an alkyl radical was generated at the anode upon SET oxidation of **234** and further deboronation (Scheme 43).


### Scheme 43. Electrochemical Redox-Neutral Deboronative Arylation of Organic Trifluoroborate Salts with 1,4-Dicyanobenzene



**3.9.2. Combination with a Cathodically Produced Ni(II)ArX Complex.** Liu et al. reported a Ni-catalyzed electrochemical cross-coupling of benzyltrifluoroborate salts **238** with (hetero)aryl bromides or chloride **236** and  $\beta$ -bromostyrene **237**. Two RVC electrodes were utilized under galvanostatic electrolysis.<sup>95</sup> Remarkably, electron-rich and electron-poor substituents were well-tolerated on the aromatic rings of both **236** and **238**. Mechanistically, cyclic voltammetry

analysis ruled out a Ni(I)/Ni(0) redox couple at the cathode. The catalytic active Ni(I) species **241** would more likely undergo oxidative addition of the organic halide **236/237** to provide Ni(III) complex **242**. Concomitant cathodic SET reduction of the latter and anodic SET deboronative oxidation of **238** afforded Ni(II) complex **243** and benzylic radical **244**, respectively. Subsequent radical combination and reductive elimination of the resulting high valent Ni(III) species **245** delivered the cross-coupling products **239/240** and recovered the active catalytic Ni(I) species (Scheme 44).


**Scheme 44. Ni-Electrocatalyzed Redox-Neutral Cross-Coupling of Organic Trifluoroborate Salts with Organic Halides**



In 2020, Hu et al. established a Ni-catalyzed electrochemical redox-neutral cross-coupling between toluene derivatives **247** and (hetero)aryl bromides **246**.<sup>95,96</sup> This attractive benzylic C(sp<sup>3</sup>)-H arylation proceeded under mild constant current electrolysis but required a specific fluorine-doped tin oxide-coated glass anode. Electron-rich as well as electron-poor aryl bromides **246** were suitable coupling partners for this transformation. However, a large excess of toluene derivatives **247** was necessary to achieve high yields (typically 3 equiv of 4-methylanisole and 10 equiv of other less electron-rich partners). Cyclic voltammetry measurements and control experiments supported the anodic generation of a benzylic radical. The latter


was introduced in the nickel catalytic cycle similarly to Liu's work (see Scheme 44) (Scheme 45).

**Scheme 45. Ni-Electrocatalyzed Redox Neutral Cross-Coupling of Toluene Derivatives with Aryl Bromides**



In the same vein, Zhang and Wang et al. have very recently developed a nickel-catalyzed electrochemical difluoromethylation of aryl iodides **249** by employing sodium difluoromethanesulfinate **250** as source of fluoroalkyl radicals with simple graphite felt electrodes in DMSO. Phenanthroline derivative **L9** was the best ligand in the presence of dimethylamino pyridine (DMAP). Moderate to good yields were obtained but no reaction occurred with aryl bromides. It is worth mentioning that this procedure could be extended to the monofluoromethylation of aryl iodides (Scheme 46).<sup>97</sup>


**Scheme 46. Ni-Electrocatalyzed Redox-Neutral Difluoromethylation of Aryl Iodides**



Very recently, Li et al. have developed an elegant electrochemical dehydroxylative coupling of primary and secondary alcohols **253** with aryl bromides **252** under Ni(II) catalysis.<sup>98</sup> The electrolysis was performed under constant current with cheap graphite anode and nickel foam cathode in NMP/LiBr as the solvent electrolyte system in the presence of an excess of triphenylphosphine. Control experiments and analysis of reaction byproducts provided evidence for the in situ generation of alkyl bromides **257**. Accordingly, two consecutive SET oxidations of bromide anion would occur first at the anode to generate Br<sub>2</sub> which would be trapped by the triphenylphosphine to form bromotriphenylphosphonium salt **255**. Further reaction with alcohol **253** would produce the alkyl bromide **257** via the intermediate **256** (Appel-type reaction). Subsequently, at the cathode, the coupling with aryl bromide **252** under Ni(II) electrocatalysis formed the C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond, in agreement

with Hansen's and Sevov's works (see section 3.8).<sup>78–80</sup> More specifically, the authors proposed the electrochemical generation of low-valent Ni(0) species via two SET reductions of NiBr<sub>2</sub>. Oxidative addition of aryl bromide **252** and combination with alkyl radical **259** gave rise to Ni(III) complex **260**. Subsequent reductive elimination afforded the coupling product **254** and Ni(I) complex **261**. The latter further reduced alkyl bromide **257** to produce the alkyl radical **259** and to regenerate the Ni(II) catalyst. A stoichiometric amount of DIPEA was required to neutralize the in situ generated HBr (Scheme 47).

### Scheme 47. Ni-Electrocatalyzed Redox-Neutral Dehydroxylative Coupling of Alcohol with Aryl Bromide



## 4. CONCLUSION

Important achievements have been recently accomplished in electrochemical cathodic processes for the formation of C(sp<sup>3</sup>)-C(sp<sup>3</sup>) and C(sp<sup>3</sup>)-C(sp<sup>2</sup>). Electroreductive transformations as well as convergent paired electrolysis have been performed under mild reaction conditions, enabling a great step toward green and sustainable chemistry. Such successes have been possible thanks to intensive mechanistic studies, the development of electrocatalytic systems, and the amelioration of electrolytic setups such as electrode materials.<sup>82</sup> The use of flow cell systems has also greatly contributed to these recent accomplishments. Indeed, by shortening the distance between the anode and cathode, such engineering minimizes the Ohmic drop and increases mass transfer, allowing therefore the application of higher currents, the reduction of the concentration of supporting electrolytes, and an easier scale-up.<sup>99,100</sup>

Additionally, the recent standardization of apparatus either in batch<sup>101,102</sup> or flow cells<sup>103–105</sup> should accelerate the recourse to the electrochemistry in both academic and industrial research. However, despite significant advances, unconquered challenges still need to be met to access more complex molecules. The design of cascade reactions and the combination with photocatalytic processes<sup>106,107</sup> are examples of avenues worth exploring. Moreover, the “holy grail” in the formation of C–C bonds is the control of the enantioselectivity. Rare examples of catalytic asymmetric electroreductive processes have been reported to date and much more development in the highly challenging enantioselective electrochemical cathodic reactions has to be accomplished. We hope that this review will stimulate further progress in this exciting area of electrochemical C–C bond-forming reactions through cathodic transformations.

## AUTHOR INFORMATION

### Corresponding Authors

**Aurélien Claraz** – Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Saclay, Gif-sur-Yvette 91198 Cedex, France; Email: aurelien.claraz@cnrs.fr

**Géraldine Masson** – Institut de Chimie des Substances Naturelles, CNRS UPR 2301, Université Paris-Saclay, Gif-sur-Yvette 91198 Cedex, France; [orcid.org/0000-0003-2333-7047](https://orcid.org/0000-0003-2333-7047); Email: geraldine.masson@cnrs.fr

Complete contact information is available at: <https://pubs.acs.org/10.1021/acsorginorgau.1c00037>

### Notes

The authors declare no competing financial interest.

## REFERENCES

- Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic organic electrochemical methods since 2000: On the verge of a renaissance. *Chem. Rev.* **2017**, *117*, 13230–13319.
- Jiang, Y.; Xu, K.; Zeng, C. Use of Electrochemistry in the Synthesis of Heterocyclic Structures. *Chem. Rev.* **2018**, *118*, 4485–4540.
- Mohle, S.; Zirbes, M.; Rodrigo, E.; Gieshoff, T.; Wiebe, A.; Waldvogel, S. R. Modern electrochemical aspects for the synthesis of value-added organic products. *Angew. Chem., Int. Ed.* **2018**, *57*, 6018–6041.
- Schotten, C.; Nicholls, T. P.; Bourne, R. A.; Kapur, N. N.; Nguyen, B.; Willans, C. E. Making electrochemistry easily accessible to the synthetic chemist. *Green Chem.* **2020**, *22*, 3358–3375.
- Wang, H.; Gao, X.; Lv, Z.; Abdelilah, T.; Lei, A. Recent Advances in oxidative R1-H/R2-H cross-coupling with hydrogen evolution via photo-/electrochemistry. *Chem. Rev.* **2019**, *119*, 6769–6787.
- Novaes, L. F. T.; Liu, J.; Shen, Y.; Lu, L.; Meinhardt, J. M.; Lin, S. Electrocatalysis as an Enabling Technology for Organic Synthesis. *Chem. Soc. Rev.* **2021**, *50*, 7941–8002.
- Park, S. H.; Ju, M.; Ressler, A. J.; Shim, J.; Kim, H.; Lin, S. Reductive electrocatalysis: a new dawn. *Aldrichimica Acta* **2021**, *54*, 17–27.
- Schäfer, H.-J. Recent contributions of Kolbe electrolysis to organic synthesis. *Top. Curr. Chem.* **1990**, *152*, 91–151.
- Moeller, K. D. Synthetic applications of anodic electrochemistry. *Tetrahedron* **2000**, *56*, 9527–9554.
- Lebreux, F. D. R.; Buzzo, F.; Marko, I. N. Synthesis of five- and six-membered-ring compounds by environmentally friendly radical cyclizations using Kolbe electrolysis. *Synlett* **2008**, *2008*, 2815–2820.
- Parida, S. K.; Mandal, T.; Das, S.; Hota, S. K.; De Sarkar, S.; Murarka, S. Single electron transfer-induced redox processes involving N-(acyloxy)phthalimides. *ACS Catal.* **2021**, *11*, 1640–1683.
- Cleary, J. A.; Mubarak, M. S.; Vieira, K. L.; Anderson, M. R.; Peters, D. G. Electrochemical reduction of alkyl halides at vitreous

carbon cathodes in dimethylformamide. *J. Electroanal. Chem. Interfacial Electrochem.* **1986**, *198*, 107–124.

(13) Andrieux, C. P.; Gallardo, I.; Savéant, J.-M. Outer-Sphere Electron-Transfer Reduction of Alkyl Halides. A Source of Alkyl Radicals or of Carbanions? Reduction of Alkyl Radicals. *J. Am. Chem. Soc.* **1989**, *111*, 1620–1626.

(14) Barton, L. M.; Chen, L.; Blackmond, D. G.; Baran, P. S. Electrochemical borylation of carboxylic acids. *Proc. Natl. Acad. Sci. U. S. A.* **2021**, *118*, e2109408118.

(15) Chen, X.; Luo, X.; Peng, X.; Guo, J.; Zai, J.; Wang, P. Catalyst-free decarboxylation of carboxylic acids and deoxygenation of alcohols by electro-induced radical formation. *Chem. - Eur. J.* **2020**, *26*, 3226–3230.

(16) Gao, Y.; Hill, D. E.; Hao, W.; McNicholas, B. J.; Vantourout, J. C.; Hadt, R. G.; Reisman, S. E.; Blackmond, D. G.; Baran, P. S. Electrochemical Nozaki–Hiyama–Kishi coupling: scope, applications, and mechanism. *J. Am. Chem. Soc.* **2021**, *143*, 9478–9488.

(17) Ni, S.; Padiyal, N. M.; Kingston, C.; Vantourout, J. C.; Schmitt, D. C.; Edwards, J. T.; Kruszyk, M. M.; Merchant, R. R.; Mykhailiuk, P. K.; Sanchez, B. B.; Yang, S.; Perry, M. A.; Gallego, G. M.; Mousseau, J. J.; Collins, M. R.; Cherney, R. J.; Lebed, P. S.; Chen, J. S.; Qin, T.; Baran, P. S. A radical approach to anionic chemistry: synthesis of ketones, alcohols, and amines. *J. Am. Chem. Soc.* **2019**, *141*, 6726–6739.

(18) Li, D.; Ma, T.-K.; Scott, R. J.; Wilden, J. D. Electrochemical radical reactions of alkyl iodides: a highly efficient, clean, green alternative to tin reagents. *Chem. Sci.* **2020**, *11*, 5333–5338.

(19) Ma, T.-K.; Li, D.; Wilden, J. Mechanistic studies of reactive oxygen species mediated electrochemical radical reactions of alkyl iodides. *Chem. Commun.* **2021**, *57*, 8356–8359.

(20) Dunach, E.; Medeiros, M. J.; Olivero, S. Electrochemical cyclizations of organic halides catalyzed by electrogenerated nickel(I) complexes: towards environmentally friendly methodologies. *Electrochim. Acta* **2017**, *242*, 373–381.

(21) Wuttig, A.; Derrick, J. S.; Loipersberger, M.; Snider, A.; Head-Gordon, M.; Chang, C. J.; Toste, F. D. Controlled single-electron transfer via metal–ligand cooperativity drives divergent nickel-electrocatalyzed radical pathways. *J. Am. Chem. Soc.* **2021**, *143*, 6990–7001.

(22) Condon, S.; Cannes, C.; Bedioui, F. Cobalt-salen catalyzed electroreductive alkylation of activated olefins. *J. Chem.* **2019**, *2019*, 1–6.

(23) For a recent review, see: Martins, G. M.; Shirinfar, B.; Hardwick, T.; Ahmed, N. A green approach: vicinal oxidative electrochemical alkene difunctionalization. *ChemElectroChem* **2019**, *6*, 1300–1315.

(24) For a recent review, see: Mei, H.; Yin, Z.; Liu, J.; Sun, H.; Han, J. Recent advances on the electrochemical difunctionalization of alkenes/alkynes. *Chin. J. Chem.* **2019**, *37*, 292–301.

(25) For a recent review, see: Sauer, G. S.; Lin, S. An electrocatalytic approach to the radical difunctionalization of alkenes. *ACS Catal.* **2018**, *8*, 5175–5187.

(26) Zhang, W.; Lin, S. Electroreductive carbonyl functionalization of alkenes with alkyl bromides via a radical-polar crossover mechanism. *J. Am. Chem. Soc.* **2020**, *142*, 20661–20670.

(27) Kisukuri, C. M.; Fernandes, V. A.; Delgado, J. A. C.; Häring, A. P.; Paixão, M. W.; Waldvogel, S. R. Electrochemical installation of CFH<sub>2</sub>–, CF<sub>2</sub>H–, CF<sub>3</sub>–, and perfluoroalkyl groups into small organic molecules. *Chem. Rec.* **2021**, *1*, 2502–2525.

(28) Pioneering work: Tommasino, J.-B.; Brondex, A.; Médebielle, M.; Thomalla, M.; Langlois, B. R.; Billard, T. Trifluoromethylation reactions with potassium trifluoromethanesulfinate under electrochemical oxidation. *Synlett* **2002**, 1697–1699.

(29) Guo, Y.; Wang, R.; Song, H.; Liu, Y.; Wang, Q. Electrochemical trifluoromethylation/cyclization for the synthesis of isoquinoline-1,3-diones and oxindoles. *Chem. Commun.* **2021**, *57*, 8284–8287.

(30) Zhou, X.; Ni, C.; Deng, L.; Hu, J. Electrochemical reduction of fluoroalkyl sulfones for radical fluoroalkylation of alkenes. *Chem. Commun.* **2021**, *57*, 8750–8753.

(31) Kise, N.; Nagamine, H.; Sakurai, T. Electroreductive intermolecular coupling of chromones with benzophenones: synthesis

of 2-diarylmethylchromones and tetrasubstituted furans. *Eur. J. Org. Chem.* **2019**, *2019*, 3662–3676.

(32) Kise, N.; Yoshimura, Y.; Manto, T.; Sakurai, T. Electroreductive intermolecular coupling of 4-quinolones with benzophenones: synthesis of 2-substituted 4-quinolones. *ACS Omega* **2019**, *4*, 20080–20093.

(33) Hu, P.; Peters, B. K.; Malapit, C. A.; Vantourout, J. C.; Wang, P.; Li, J.; Mele, L.; Echeverria, P.-G.; Minter, S. D.; Baran, P. S. Electroreductive olefin–ketone coupling. *J. Am. Chem. Soc.* **2020**, *142*, 20979–20986.

(34) Cao, Y.; Knijff, J.; Delparish, A.; d'Angelo, M. F. N.; Noël, T. A divergent paired electrochemical process for the conversion of furfural using a divided-cell flow microreactor. *ChemSusChem* **2021**, *14*, 590–594.

(35) Chalkley, M. J.; Garrido-Barros, P.; Peters, J. C. A molecular mediator for reductive concerted proton-electron transfer via electrocatalysis. *Science* **2020**, *369*, 850–854.

(36) Liu, C.; Li, R.; Zhou, W.; Liang, Y.; Shi, Y.; Li, R.-L.; Ling, Y.; Yu, Y.; Li, J.; Zhang, B. Selectivity origin of organic electrosynthesis controlled by electrodes material: a case study of pinacols. *ACS Catal.* **2021**, *11*, 8958–8967.

(37) Anibal, J.; Xu, B. Electroreductive C–C coupling of furfural and benzaldehyde on Cu and Pb Surfaces. *ACS Catal.* **2020**, *10*, 11643–11653.

(38) Thangavelu, S.; Subbiah, P.; Udupa, K. S. Electrochemical reduction of acrylonitrile on glassy carbon electrode in aqueous solutions. *B. Electrochem.* **1985**, *1*, 137–140.

(39) Suwanvaipattana, P.; Limtrakul, S.; Vatanatham, T.; Ramachandran, P. A. Modeling of electro-organic synthesis to facilitate cleaner chemical manufacturing: Adiponitrile production. *J. Cleaner Prod.* **2017**, *142*, 1296–1308.

(40) Blanco, D. E.; Dookhith, A. Z.; Modestino, M. A. Enhancing selectivity and efficiency in the electrochemical synthesis of adiponitrile. *React. Chem. Eng.* **2019**, *4*, 8–16.

(41) Blanco, D. E.; Lee, B.; Modestino, M. A. Optimizing organic electrosynthesis through controlled voltage dosing and artificial intelligence. *Proc. Natl. Acad. Sci. U. S. A.* **2019**, *116*, 17683–17689.

(42) Lu, L.; Siu, J. C.; Lai, Y.; Lin, S. An electroreductive approach to radical silylation via the activation of strong Si–Cl bonds. *J. Am. Chem. Soc.* **2020**, *142*, 21272–21278.

(43) Li, H.; Breen, C. P.; Seo, H.; Jamison, T. F.; Fang, Y.-Q.; Bio, M. M. Ni-Catalyzed electrochemical decarboxylative C–C couplings in batch and continuous flow. *Org. Lett.* **2018**, *20*, 1338–1341.

(44) Koyanagi, T.; Herath, A.; Chong, A.; Ratnikov, M.; Valiere, A.; Chang, J.; Molteni, V.; Loren, J. One-pot electrochemical nickel-catalyzed decarboxylative sp<sup>2</sup>–sp<sup>3</sup> cross-coupling. *Org. Lett.* **2019**, *21*, 816–820.

(45) Lian, F.; Xu, K.; Meng, W.; Zhang, H.; Tan, Z.; Zeng, C. Nickel-catalyzed electrochemical reductive decarboxylative coupling of N-hydroxyphthalimide esters with quinoxalinones. *Chem. Commun.* **2019**, *55*, 14685–14688.

(46) Niu, K.; Song, L.; Hao, Y.; Liu, Y.; Wang, Q. Electrochemical decarboxylative C3 alkylation of quinoxalin-2(1H)-ones with N-hydroxyphthalimide esters. *Chem. Commun.* **2020**, *56*, 11673–11676.

(47) Liu, Y.; Xue, L.; Shi, B.; Bu, F.; Wang, D.; Lu, L.; Shi, R.; Lei, A. Catalyst-free electrochemical decarboxylative cross-coupling of N-hydroxyphthalimide esters and N-heteroarenes towards C(sp<sup>3</sup>)–C(sp<sup>2</sup>) bond formation. *Chem. Commun.* **2019**, *55*, 14922–14925.

(48) Lübbsmeyer, M.; Leifert, D.; Schäfer, H.; Studer, A. Electrochemical initiation of electron-catalyzed phenanthridine synthesis by trifluoromethylation of isonitriles. *Chem. Commun.* **2018**, *54*, 2240–2243.

(49) Jud, W.; Maljuric, S.; Kappe, C. O.; Cantillo, D. Cathodic C–H trifluoromethylation of arenes and heteroarenes enabled by an in situ-generated triflyltriethylammonium complex. *Org. Lett.* **2019**, *21*, 7970–7975.

(50) Rodrigo, S.; Um, C.; Mixdorf, J. C.; Gunasekera, D.; Nguyen, H. M.; Luo, L. Alternating current electrolysis for organic electrosynthesis:

- trifluoromethylation of (hetero)arenes. *Org. Lett.* **2020**, *22*, 6719–6723.
- (51) Hossain, M. J.; Ono, T.; Wakiya, K.; Hisaeda, Y. A vitamin B12 derivative catalyzed electrochemical trifluoromethylation and perfluoroalkylation of arenes and heteroarenes in organic media. *Chem. Commun.* **2017**, *53*, 10878–10881.
- (52) Alkayal, A.; Tabas, V.; Montanaro, S.; Wright, I. A.; Malkov, A. V.; Buckley, B. R. Harnessing applied potential: selective  $\beta$ -hydrocarboxylation of substituted olefins. *J. Am. Chem. Soc.* **2020**, *142*, 1780–1785.
- (53) Sheta, A. M.; Mashaly, M. A.; Said, S. B.; Elmorsy, S. S.; Malkov, A. V.; Buckley, B. R. Selective  $\alpha,\delta$ -hydrocarboxylation of conjugated dienes utilizing CO<sub>2</sub> and electrosynthesis. *Chem. Sci.* **2020**, *11*, 9109–9114.
- (54) Sheta, A.; Alkayal, A.; Mashaly, M.; Said, S.; Elmorsy, S.; Malkov, A.; Buckley, B. Selective electrosynthetic hydrocarboxylation of  $\alpha,\beta$ -unsaturated esters with carbon dioxide. *Angew. Chem., Int. Ed.* **2021**, *60*, 21832–21837.
- (55) Chen, R.; Tian, K.; He, D.; Gao, T.; Yang, G.; Xu, J.; Chen, H.; Wang, D.; Zhang, Y. Carboxylation of  $\alpha,\beta$ -unsaturated ketones by CO<sub>2</sub> fixation through photoelectro-chemistry. *ACS Appl. Energy Mater.* **2020**, *3*, 5813–5818.
- (56) Gao, X.-T.; Zhang, Z.; Wang, X.; Tian, J.-S.; Xie, S.-L.; Zhou, F.; Zhou, J. Direct electrochemical defluorinative carboxylation of  $\alpha$ -CF<sub>3</sub> alkenes with carbon dioxide. *Chem. Sci.* **2020**, *11*, 10414–10420.
- (57) Zhang, S.; Li, L.; Li, X.; Zhang, J.; Xu, K.; Li, G.; Findlater, M. Electroreductive 4-pyridylation of electron-deficient alkenes with assistance of Ni(acac)<sub>2</sub>. *Org. Lett.* **2020**, *22*, 3570–3575.
- (58) Xu, H.; Liu, J.; Nie, F.; Zhao, X.; Jiang, Z. Metal-free hydroxyarylation of thioester-activated alkenes via electroreductive radical coupling. *J. Org. Chem.* **2021**, *86*, 16204.
- (59) Lehnher, D.; Lam, Y.-H.; Nicastrì, M. C.; Liu, J.; Newman, J. A.; Regalado, E. L.; DiRocco, D. A.; Rovis, T. Electrochemical synthesis of hindered primary and secondary amines via proton-coupled electron transfer. *J. Am. Chem. Soc.* **2020**, *142*, 468–478.
- (60) Zhang, X.; Yang, C.; Gao, H.; Wang, L.; Guo, L.; Xia, W. Reductive arylation of aliphatic and aromatic aldehydes with cyanoarenes by electrolysis for the synthesis of alcohols. *Org. Lett.* **2021**, *23*, 3472–3476.
- (61) Zhang, S.; Li, L.; Li, J.; Shi, J.; Xu, K.; Gao, W.; Zong, L.; Li, G.; Findlater, M. Electrochemical arylation of aldehydes, ketones, and alcohols: from cathodic reduction to convergent paired electrolysis. *Angew. Chem., Int. Ed.* **2021**, *60*, 7275–7282.
- (62) Kuroboshi, M.; Tanaka, M.; Kishimoto, S.; Tanaka, H.; Torii, S. Electrochemical regeneration of chromium(II). Alkenylation of carbonyl compounds. *Synlett* **1999**, *1999*, 69–70.
- (63) Durandetti, M.; Périchon, J.; Nédélec, J.-Y. Nickel- and chromium-catalyzed electrochemical coupling of aryl halides with arenecarboxaldehydes. *Tetrahedron Lett.* **1999**, *40*, 9009–9013.
- (64) Durandetti, M.; Nédélec, J.-Y.; Périchon, J. An electrochemical coupling of organic halide with aldehydes, catalytic in chromium and nickel salts. The Nozaki–Hiyama–Kishi Reaction. *Org. Lett.* **2001**, *3*, 2073–2076.
- (65) Déjardin, C.; Renou, A.; Maddaluno, J.; Durandetti, M. Nickel-Catalyzed Electrochemical Cyclization of Alkynyl Aryl Iodide and the Domino Reaction with Aldehydes. *J. Org. Chem.* **2021**, *86*, 8882–8890.
- (66) For a review, see: Senboku, H. Electrochemical fixation of carbon dioxide: synthesis of carboxylic acids. *Chem. Rec.* **2021**, *21*, 2354.
- (67) Yang, H.-P.; Zhang, H.-W.; Wu, Y.; Fan, L.-D.; Chai, X.-Y.; Zhang, Q.-L.; Liu, J.-H.; He, C.-H. A Core–Shell-Structured Silver Nanowire/Nitrogen-Doped Carbon Catalyst for Enhanced and Multifunctional Electrofixation of CO<sub>2</sub>. *ChemSusChem* **2018**, *11*, 3905–3910.
- (68) Senboku, H.; Sakai, K.; Fukui, A.; Sato, Y.; Yamauchi, Y. A Core–shell-structured silver nanowire/nitrogen-doped carbon catalyst for enhanced and multifunctional electrofixation of CO<sub>2</sub>. *ChemElectroChem* **2019**, *6*, 4158–4164.
- (69) Senboku, H.; Minemura, Y.; Suzuki, Y.; Matsuno, H.; Takakuwa, M. Synthesis of N-Boc- $\alpha$ -amino acids from carbon dioxide by electrochemical carboxylation of N-Boc- $\alpha$ -aminosulfones. *J. Org. Chem.* **2021**, *86*, 16077.
- (70) Mena, S.; Santiago, S.; Gallardo, I.; Guirado, G. Sustainable and efficient electrosynthesis of naproxen using carbon dioxide and ionic liquids. *Chemosphere* **2020**, *245*, 125557–125566.
- (71) Mena, S.; Sanchez, J.; Guirado, G. Electrocarboxylation of 1-chloro-(4-isobutylphenyl)ethane with a silver cathode in ionic liquids: an environmentally benign and efficient way to synthesize Ibuprofen. *RSC Adv.* **2019**, *9*, 15115–15123.
- (72) Corbin, N.; Yang, D.-T.; Lazowski, N.; Steinberg, K.; Manthiram, K. Suppressing carboxylate nucleophilicity with inorganic salts enables selective electrocarboxylation without sacrificial anodes. *Chem. Sci.* **2021**, *12*, 12365–12376.
- (73) Naito, Y.; Nakamura, Y.; Shida, N.; Senboku, H.; Tanaka, K.; Atobe, M. Integrated flow synthesis of  $\alpha$ -amino acids by in situ generation of aldimines and subsequent electrochemical carboxylation. *J. Org. Chem.* **2021**, *86*, 15953.
- (74) Jiao, K.-J.; Li, Z.-M.; Xu, X.-T.; Zhang, L.-P.; Li, Y.-Q.; Zhang, K.; Mei, T.-S. Palladium-catalyzed reductive electrocarboxylation of allyl esters with carbon dioxide. *Org. Chem. Front.* **2018**, *5*, 2244–2248.
- (75) Ang, N. W. J.; Oliveira, J. C. A.; Ackermann, L. Electroreductive cobalt-catalyzed carboxylation: cross-electrophile coupling with atmospheric CO<sub>2</sub>. *Angew. Chem., Int. Ed.* **2020**, *59*, 12842–12847.
- (76) Durandetti, M.; Nédélec, J. Y.; Périchon, J. Nickel-catalyzed direct electrochemical cross-coupling between aryl halides and activated alkyl halides. *J. Org. Chem.* **1996**, *61*, 1748–1755.
- (77) Durandetti, M.; Périchon, J. Nickel-catalyzed electrochemical coupling of aryl, heteroaryl or vinyl halides with activated alkyl chlorides: synthetic and stereochemical aspects. *Synthesis* **2004**, *18*, 3079–3083.
- (78) Perkins, R. J.; Hughes, A. J.; Weix, D. J.; Hansen, E. C. Metal-reductant-free electrochemical nickel-catalyzed couplings of aryl and alkyl bromides in acetonitrile. *Org. Process Res. Dev.* **2019**, *23*, 1746–1751.
- (79) Perkins, R. J.; Pedro, D. J.; Hansen, E. C. Electrochemical nickel catalysis for sp<sup>2</sup>–sp<sup>3</sup> cross-electrophile coupling reactions of unactivated alkyl halides. *Org. Lett.* **2017**, *19*, 3755–3758.
- (80) Truesdell, B. L.; Hamby, T. B.; Sevov, C. S. General C(sp<sup>2</sup>)–C(sp<sup>3</sup>) cross-electrophile coupling reactions enabled by overcharge protection of homogeneous electrocatalysts. *J. Am. Chem. Soc.* **2020**, *142*, 5884–5893.
- (81) Jiao, K.-J.; Liu, D.; Ma, H. X.; Qiu, H.; Fang, P.; Mei, T.-S. Nickel-catalyzed electrochemical reductive relay cross-coupling of alkyl halides to aryl halides. *Angew. Chem., Int. Ed.* **2020**, *59*, 6520–6524.
- (82) Kumar, G. S.; Peshkov, A.; Brzozowska, A.; Nikolaienko, P.; Zhu, C.; Rueping, M. Nickel-catalyzed chain-walking cross-electrophile coupling of alkyl and aryl halides and olefin hydroarylation enabled by electrochemical reduction. *Angew. Chem., Int. Ed.* **2020**, *59*, 6513–6519.
- (83) Jiao, K.-J.; Ma, C.; Liu, D.; Qiu, H.; Cheng, B.; Mei, T.-S. Nickel-catalyzed electrochemical reductive relay cross-coupling of alkyl halides with alkyl carboxylic acids. *Org. Chem. Front.* **2021**, *8*, 6603.
- (84) (a) DeLano, T. J.; Reisman, S. E. Enantioselective electroreductive coupling of alkenyl and benzyl halides via nickel catalysis. *ACS Catal.* **2019**, *9*, 6751–6754.
- (85) Cherney, A. H.; Reisman, S. E. Nickel-catalyzed asymmetric reductive cross-coupling between vinyl and benzyl electrophiles. *J. Am. Chem. Soc.* **2014**, *136*, 14365–14368.
- (86) Recent review: Ghosh, M.; Shinde, V. M.; Rueping, M. A review of asymmetric synthetic organic electrochemistry and electrocatalysis: concepts, applications, recent developments and future directions. *Beilstein J. Org. Chem.* **2019**, *15*, 2710–2746.
- (87) Recent review: Chang, X.; Zhang, Q.; Guo, C. Asymmetric electrochemical transformations. *Angew. Chem.* **2020**, *132*, 12712–12722.
- (88) Recent review: Yamamoto, K.; Kuriyama, M.; Onomura, O. Asymmetric electrosynthesis: Recent advances in catalytic transformations. *Curr. Opin. Electrochem.* **2021**, *28*, 100714–100720.

- (89) Recent review: Margarita, C.; Lundberg, H. Recent advances in asymmetric catalytic electrosynthesis. *Catalysts* **2020**, *10*, 982–1005.
- (90) Samanta, R. C.; Struwe, J.; Ackermann, L. Nickel-electrocatalyzed mild C–H alkylations at room temperature. *Angew. Chem., Int. Ed.* **2020**, *59*, 14154–14159.
- (91) Dicciani, J. B.; Diao, T. Mechanisms of nickel-catalyzed cross-coupling reactions. *Trends Chem.* **2019**, *1*, 830–844.
- (92) Ma, Y.; Yao, X.; Zhang, L.; Ni, P.; Cheng, R.; Ye, J. Direct arylation of  $\alpha$ -amino C(sp<sup>3</sup>)-H bonds by convergent paired electrolysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 16548–16552.
- (93) Mo, Y.; Lu, Z.; Rughoobur, G.; Patil, P.; Gershenfeld, N.; Akinwande, A. I.; Buchwald, S. L.; Jensen, K. F. Microfluidic electrochemistry for single-electron transfer redox-neutral reactions. *Science* **2020**, *368*, 1352–1357.
- (94) Luo, J.; Hu, B.; Wu, W.; Hu, M.; Liu, T. L. Nickel catalyzed electrochemical C(sp<sup>3</sup>)-C(sp<sup>2</sup>) cross-coupling reactions of benzyl trifluoroborate and organic halides. *Angew. Chem., Int. Ed.* **2021**, *60*, 6107–6116.
- (95) Zhang, L.; Hu, X. Nickel catalysis enables convergent paired electrolysis for direct arylation of benzylic C–H bonds. *Chem. Sci.* **2020**, *11*, 10786–10791.
- (96) Zou, Z.; Li, H.; Huang, M.; Zhang, W.; Zhi, S.; Wang, Y.; Pan, Y. Electrochemical-promoted nickel-catalyzed oxidative fluoroalkylation of aryl iodides. *Org. Lett.* **2021**, *23*, 8252.
- (97) Li, Z.; Sun, W.; Wang, X.; Li, L.; Zhang, Y.; Li, C. Electrochemically enabled, nickel-catalyzed dehydroxylative cross-coupling of alcohols with aryl halides. *J. Am. Chem. Soc.* **2021**, *143*, 3536–3543.
- (98) Heard, D. M.; Lennox, A. J. J. Electrode materials in modern organic electrochemistry. *Angew. Chem., Int. Ed.* **2020**, *59*, 18866–18884.
- (99) Noël, T.; Cao, Y.; Laudadio, G. The fundamentals behind the use of flow reactors in electrochemistry. *Acc. Chem. Res.* **2019**, *52*, 2858–2869.
- (100) Maljuric, S.; Jud, W.; Kappe, C. O.; Cantillo, D. Translating batch electrochemistry to single-pass continuous flow conditions: an organic chemist's guide. *J. Flow Chem.* **2020**, *10*, 181–190.
- (101) For selected examples, see: (a) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic organic electrochemistry: calling all engineers. *Angew. Chem., Int. Ed.* **2018**, *57*, 4149–4155.
- (102) Gütz, C.; Klöckner, B.; Waldvogel, S. R. *Org. Process Res. Dev.* **2016**, *20*, 26–32.
- (103) For a selected example, see: Rein, J.; Annand, J. R.; Wismer, M. K.; Fu, J.; Siu, J. C.; Klapars, A.; Strotman, N. A.; Kalyani, D.; Lehnher, D.; Lin, S. Unlocking the potential of high-throughput experimentation for electrochemistry with a standardized microscale reactor. *ACS Cent. Sci.* **2021**, *7*, 1347–1355.
- (104) For a selected example, see: Folgueiras-Amador, A. A.; Philipps, K.; Guillaud, S.; Poelakker, J.; Wirth, T. An easy-to-machine electrochemical flow microreactor: efficient synthesis of isoindolinone and flow functionalization. *Angew. Chem., Int. Ed.* **2017**, *56*, 15446–15450.
- (105) For a selected example, see: Laudadio, G.; de Smet, W.; Struik, L.; Cao, Y.; Noël, T. Design and application of a modular and scalable electrochemical flow microreactor. *J. Flow Chem.* **2018**, *8*, 157–165.
- (106) Barham, J. P.; König, B. Synthetic photoelectrochemistry. *Angew. Chem., Int. Ed.* **2020**, *59*, 11732–11747.
- (107) Buglioni, L.; Raymenants, F.; Slattery, A.; Zondag, S. D. A.; Noël, T. Technological innovations in photochemistry for organic synthesis: Flow chemistry, high-throughput experimentation, scale-up, and photoelectrochemistry. *Chem. Rev.* **2021**, DOI: 10.1021/acs.chemrev.1c00332.

## Recommended by ACS

### Benzylic C(sp<sup>3</sup>)-H Bonds Play the Dual Role of Starting Material and Oxidation Inhibitor for Hydrazides in the Electrochemical Synthesis of Hydrazones

Issa Yavari and Sina Shaabanzadeh

NOVEMBER 08, 2022  
THE JOURNAL OF ORGANIC CHEMISTRY

READ 

### High Electric Field on Water Microdroplets Catalyzes Spontaneous and Ultrafast Oxidative C–H/N–H Cross-Coupling

Dongmei Zhang, Xinxing Zhang, *et al.*

AUGUST 12, 2022  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

### Electroreduction of Carbonyl Compounds Catalyzed by a Manganese Complex

Igor Fokin, Inke Siewert, *et al.*

JULY 06, 2022  
ACS CATALYSIS

READ 

### Electrocatalytic Radical-Polar Crossover Hydroetherification of Alkenes with Phenols

Steve H Park, Hyunwoo Kim, *et al.*

AUGUST 12, 2022  
ACS CATALYSIS

READ 

Get More Suggestions >