


# A Broad View of Structure-Activity Relationship Performance for Alkanes and Haloalkanes




The 26<sup>th</sup> International Symposium on Gas Kinetics and Related Phenomena, Rennes, 2022

**Max McGillen** (CNRS-Orléans, France)



✉ [max.mcgillen@cnrs-orleans.fr](mailto:max.mcgillen@cnrs-orleans.fr)

# The combinatorial explosion


## – A justification for SARs –



- The atmosphere is a complicated place
- Our best view of complexity comes from automated mechanism generation software (AMG)
- AMG produces a large number of molecules
- **About which we don't know very much...**



- A high degree of multifunctionality
- Especially for functional groups that aren't well studied
- Prediction of these rate coefficients is surely a gross extrapolation
- **Clearly this is too much for now...**



- Valorso, R. *et al.* Explicit modelling of SOA formation from  $\alpha$ -pinene photooxidation: sensitivity to vapour pressure estimation. *Atmos. Chem. Phys.* **11**, 6895–6910 (2011).
- McGillen, M. R. *et al.* Database for the kinetics of the gas-phase atmospheric reactions of organic compounds. *Earth System Science Data* **12**, 1203–1216 (2020).




# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons



- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation




# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons



- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons




McGillen, M. R. *et al.* Database for the kinetics of the gas-phase atmospheric reactions of organic compounds. *Earth System Science Data* **12**, 1203–1216 (2020).

- Only consider alkanes and haloalkanes
- Atmospherically important (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons





- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons

No OH addition

No H-bonded complexes




- Michelat et al. *in preparation*.
- El Othmani, H. et al. Gas-Phase Rate Coefficient of OH + 1,2-Epoxybutane Determined between 220 and 950 K. *ACS Earth Space Chem.* **5**, 960–968 (2021).

- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons



$\Delta_f H$

$D_{R-H}$

IP


Quantum  
chem. data

Swain-Lupton  
constants,  $F, R$

- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons



- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- **Highly functionalized**
- Different methods for estimation




# The Sandbox: a unique testing environment containing hydrocarbons and halocarbons



- Only consider alkanes and haloalkanes
- Atmospherically important
- (relatively) high data coverage
- Variation in reactivity
- Mechanistically simpler
- Complementary information
- Highly functionalized
- Different methods for estimation



# Group-additivity methods



- This method (and its derivatives) is the go-to group-additivity method for atmospheric chemistry
- **Despite this, it's not very effective for haloalkanes**

$$k(\text{CH}_3\text{-X}) = k_{\text{prim}} F(\text{X})$$


$$k(\text{X-CH}_2\text{-Y}) = k_{\text{sec}} F(\text{X}) F(\text{Y})$$

$$k\left(\text{X-CH}\begin{matrix} \nearrow \text{Y} \\ \searrow \text{X} \end{matrix}\right) = k_{\text{tert}} F(\text{X}) F(\text{Y}) F(\text{Z})$$

- Kwok, E. & Atkinson, R. Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: An update. *Atmos. Environ.* **29**, 1685–1695 (1995).



# Group-additivity methods




- This approach is less commonly used than the Atkinson approach
- **However, it is clearly superior for haloalkanes, which may relate to its “third-group multiplier”**

$$\log k = \log k(\text{CH}_4) + G_1 \dots G_3$$

- DeMore, W. B. Experimental and Estimated Rate Constants for the Reactions of Hydroxyl Radicals with Several Halocarbons. *J. Phys. Chem.* **100**, 5813–5820 (1996).
- **Update of fitting parameters: this work**

# Group-additivity methods



- We updated an update of the Atkinson method (Calvert) to impose lower limits on individual reaction sites
- **This mimics the “third-group multiplier” and appears to improve the algorithm**

$$k(\text{CH}_3\text{-X}) = k_{\text{prim}} F(\text{X}) \quad \text{if } k(\text{CH}_3\text{-X}) < a = a$$


$$k(\text{X-CH}_2\text{-Y}) = k_{\text{sec}} F(\text{X}) F(\text{Y}) \quad \text{if } k(\text{X-CH}_2\text{-Y}) < b = b$$

$$k\left(\text{X-CH} \begin{array}{l} \swarrow \text{Y} \\ \searrow \text{X} \end{array}\right) = k_{\text{tert}} F(\text{X}) F(\text{Y}) F(\text{Z}) \quad \text{if } k(\text{X-CH}_2 < \frac{\text{Y}}{\text{Z}}) < c = c$$

- Calvert, J. G., Derwent, R. G., Orlando, J. J., Tyndall, G. S. & Wallington, T. J. *Mechanisms of the atmospheric oxidations of the alkanes*. (Oxford University Press, 2008).
- **Algorithm update: this work**



# Linear free-energy relationship methods




- Ionization potential (IP) has been used previously to correlate with  $k_{\text{OH}}$
- **IP calculated using PM6, and corrected against experimental values (where available)**

$$\frac{\log k}{n_{\text{H}}} = m \cdot \text{IP} + c$$

- Percival, C. J., Marston, G. & Wayne, R. P. Correlations between rate parameters and calculated molecular properties in the reactions of the hydroxyl radical with hydrofluorocarbons. *Atmospheric Environment* **29**, 305–311 (1995).
- Stewart, J. J. P. MOPAC2016.
- **Algorithm update, IP calculations: this work**

# Linear free-energy relationship methods




- The OH-Cl relationship has been observed to work well before
- It implies a similar mechanism, probably C-H bond strength
- **Although it is a reasonably good predictor, there are some problems with higher  $k$  values (which can be corrected)**

- T.J. Wallington and co-workers, various works, applied mainly to HFEs
- **This work, focussing on a more complete alkanes and haloalkane dataset**




# Linear free-energy relationship methods



- The OH–O(<sup>3</sup>P) relationship has also been observed to work well before
- As with Cl, it implies a similar mechanism, probably C–H bond strength
- **These types of measurements seem to have gone out of favour, so there is very little new data to correlate**

- Gaffney, J. S. & Levine, S. Z. Predicting gas phase organic molecule reaction rates using linear free-energy correlations. I. O(<sup>3</sup>P) and OH addition and abstraction reactions. *Int. J. Chem. Kinet.* **11**, 1197–1209 (1979).
- **This work, focussing on a more complete alkanes and haloalkane dataset**

# Linear free-energy relationship methods




- Correlations between aqueous-phase and gas-phase rate coefficients have been attempted previously
- **For this set of compounds, it appears (at least with the quality of the data available) that this relationship is too scattered to be useful**

- Klöpffer, W., Kaufmann G. & Frank, R. Phototransformation of Air Pollutants: Rapid Test for the Determination of  $k_{\text{OH}}$ . *Zeitschrift für Naturforschung A* **40**, 686 (1985).
- Wallington, T. J., Dagaut, P. & Kurylo, M. J. Correlation between gas-phase and solution-phase reactivities of hydroxyl radicals towards saturated organic compounds. *J. Phys. Chem.* **92**, 5024–5028 (1988).
- **This work, focussing on a more complete alkanes and haloalkane dataset**




# Linear free-energy relationship methods



- But, it is more compelling if we look at the broader picture
- Similarities between gas- and solution-phase have been observed for  $\geq 100$  years!
- **Alkanes and halocarbons are among the biggest outliers, which may result from difficulties getting them into aqueous solution**

- Daniels, F. & Johnston, E. H. The thermal decomposition of gaseous nitrogen pentoxide. A monomolecular reaction. *J. Am. Chem. Soc.* **43**, 53–71 (1921).
- Lueck, R. H. The thermal decomposition of nitrogen pentoxide in solution. *J. Am. Chem. Soc.* **44**, 757–769 (1922).

# Linear free-energy relationship methods




- It is expected that BDE is one of the primary determinants in this hydrogen abstraction mechanism
- Accurate BDEs are few and far between, a machine-learning method is available
- **This relationship appears to break down for the more reactive molecules ( $E_a$  vs  $A$ ?)**

- Hsu, K.-J. & DeMore, W. B. Temperature-Dependent Rate Constants and Substituent Effects for the Reactions of Hydroxyl Radicals with Three Partially Fluorinated Ethers. *J. Phys. Chem.* **99**, 11141–11146 (1995).
- St. John, P. C., Guan, Y., Kim, Y., Kim, S. & Paton, R. S. Prediction of organic homolytic bond dissociation enthalpies at near chemical accuracy with sub-second computational cost. *Nat. Commun.* **11**, 2328 (2020).




# Linear free-energy relationship methods



- Is the machine-learning approach accurate for BDEs in our dataset?
- **It seems to perform quite well, of course there are also uncertainties in the experimental database...**

- Luo, Y.-R. *Handbook of Bond Dissociation Energies in Organic Compounds*. (CRC Press, 2002). doi:10.1201/9781420039863.
- St. John, P. C., Guan, Y., Kim, Y., Kim, S. & Paton, R. S. Prediction of organic homolytic bond dissociation enthalpies at near chemical accuracy with sub-second computational cost. *Nat. Commun.* **11**, 2328 (2020).

# Linear free-energy relationship methods



- This method requires a lot of independent variables:  $(\Delta_r H, F, R)$
- This isn't very practical! However, it becomes more practical with ML
- **This extends the applicability of Poutsma's approach, which is otherwise quite reliable**


$$\log k(\text{OH}) = -0.000630\Delta_r H^2 - 0.151\Delta_r H - 1.056 \sum F - 1.053 \sum R - 21.26$$

where  $\Delta_r H = \text{BDE}(\text{R-H})_{ML} - \text{BDE}(\text{HO-H})$

• Poutsma, M. L. Evolution of Structure–Reactivity Correlations for the Hydrogen Abstraction Reaction by Hydroxyl Radical and Comparison with That by Chlorine Atom. *The Journal of Physical Chemistry A* **117**, 6433–6449 (2013).



# Linear free-energy relationship methods




- This method was developed at a time when there was less kinetic information available
- **Considering this, it performs quite well, and it can probably be improved in the future**

$$k(T) = n_H 10^{6.55} M^{-1} T^{1.5} \exp[-190(D_{RH} - 95.3)/T]$$




where  $D_{RH} = D_{RH_{ML}} = \text{BDE}(\text{R-H})_{ML}$

- Cohen, N. & Benson, S. W. Empirical correlations for rate coefficients for reactions of hydroxyl radicals with haloalkanes. *J. Phys. Chem.* **91**, 171–175 (1987).
- **This work, extending application with ML BDEs**

# New approach: the electrotopological state



- A new method has been developed, which makes use of the Kier-Hall index
- This describes the electronic character and the connectivity of a hydrogen environment
- **This approach yields site-specific rate coefficients with high accuracy and a low number of fitting parameters**



Hall, L. H., Mohney, B. & Kier, L. B. The electrotopological state: structure information at the atomic level for molecular graphs. *J. Chem. Inf. Model.* **31**, 76–82 (1991).



# Can we combine these results somehow?

- It seems plausible that in a highly uncertain situation, you should make use of as much information as possible
- The following statement may apply to our individual SAR models:

*“Every attempt to represent nature in a set of equations, resolvable on a digital computer, inevitably introduces inaccuracy.” (Hagedorn et al., 2005)*


- Hagedorn et al. were discussing the macroscopic world of numerical weather prediction – not the microscopic world of chemical reactivity
- However, there is an important parallel: we all must extrapolate
- To test the ensemble, I took a simple “multiple measurement” approach:

$$k_{ensemble} = \frac{\sum_{i=1}^N w_i k_i}{\sum_{i=1}^N w_i}$$

where,  $w_i = 1/\sigma_i^2$

Hagedorn, R., Doblas-Reyes, F. J. & Palmer, T. N. The rationale behind the success of multi-model ensembles in seasonal forecasting - I. Basic concept. *Tellus A* **57**, 219–233 (2005).


# Bof ... ?




- The ensemble is less accurate at reproducing the experimental data than some individual approaches
- This was (initially) disappointing
- But, we shouldn't assume that our database is perfect
- **There is an interesting asymmetry to the errors in estimation space, with a tendency towards underprediction**



# Is the ensemble helpful?



- High UV cross sections
- Low vapour pressure
- Single measurements



- Very low  $k$  values
- High sensitivity to impurities
- High(er?) GWPs

# Conclusions and perspectives

- The Sandbox is a useful place to conduct this thought experiment
- There is a diversity of methods for making predictions, several of which are at least as good as the Atkinson G-A approach
- In addition, I present you with an entirely new site-specific method for rate coefficient estimation based on the electrotopological state
- The ensemble appears to be able to identify possibly erroneous data, **this should be checked with new experiments**
- There are several things that I didn't have time to discuss, but which are also very important:
- Site-specific reaction channels; influence of temperature; **consensus in estimation space** – next time

*Thank you for your attention,  
and thanks to my colleagues in the CRC SAR panel for helpful discussions*