

# **Spectral Analysis of the Incompressible Viscous Rayleigh–Taylor System in R3**

Olivier Lafitte, Tiến-Tài Nguyễn

# **To cite this version:**

Olivier Lafitte, Tiến-Tài Nguyễn. Spectral Analysis of the Incompressible Viscous Rayleigh–Taylor System in R3. Water Waves, 2022, 4 (2), pp.259 - 305. 10.1007/s42286-022-00065-5. hal-03857523

# **HAL Id: hal-03857523 <https://hal.science/hal-03857523>**

Submitted on 25 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# <span id="page-1-7"></span>SPECTRAL ANALYSIS OF THE INCOMPRESSIBLE VISCOUS RAYLEIGH-TAYLOR SYSTEM IN R<sup>3</sup>

### <span id="page-1-3"></span><span id="page-1-2"></span>OLIVIER LAFITTE AND TIẾN-TÀI NGUYỄN

Abstract. The linear instability study of the viscous Rayleigh-Taylor model in the neighborhood of a laminar smooth increasing density profile  $\rho_0(x_3)$ amounts to the study of the following ordinary differential equation of order 4:

<span id="page-1-0"></span> $-\lambda^2[\rho_0 k^2 \phi - (\rho_0 \phi')'] = \lambda \mu (\phi^{(4)} - 2k^2 \phi'' + k^4 \phi) - g k^2 \rho'$  $(0.1)$ 

where  $\lambda$  is the growth rate in time,  $k$  is the wave number transverse to the density profile.

In the case of  $\rho'_0 \geq 0$  compactly supported, we provide a spectral analysis showing that in accordance with the results of  $[8]$ , there is an infinite sequence of non trivial solutions  $(\lambda_n, \phi_n)$  of [\(0.1\)](#page-1-0), with  $\lambda_n \to 0$  when  $n \to +\infty$  and  $\phi_n \in H^4(\mathbf{R})$ . In the more general case where  $\rho'_0 > 0$  everywhere and  $\rho_0$ converges at  $\pm\infty$  to finite limits  $\rho_{\pm} > 0$ , we prove that there exist finitely non trivial solutions  $(\lambda_n, \phi_n)$  of  $(0.1)$ . The line of investigation is to reduce both cases to the study of a self-adjoint operator on a compact set.

### <span id="page-1-1"></span>CONTENTS

<span id="page-1-5"></span><span id="page-1-4"></span>

<span id="page-1-6"></span>*Date*:  $21^{st}$  May, 2022 at 06:04.

<sup>2010</sup> *Mathematics Subject Classification.* 34B05, 47A05, 47A55, 47B07, 76D05.

*Key words and phrases.* Rayleigh–Taylor instability, linear growth rate, self-adjoint operator, spectral theory.



#### <span id="page-2-1"></span>DATA AVAILABILITY STATEMENT

All data generated or analysed during this study are included in this published article (and its supplementary information files).

#### 1. INTRODUCTION

<span id="page-2-0"></span>The Rayleigh–Taylor (RT) instability, studied first by Lord Rayleigh in [\[19\]](#page-36-1) and then Taylor [\[20\]](#page-36-2) is well known as a gravity-driven instability in two semi-infinite inviscid and incompressible fluids when the heavy one is on top of the light one. It has attracted much attention due to both its physical and mathematical importance. Two applications of worth mentioning are implosion of inertial confinement fusion capsules [\[13\]](#page-36-3) and core-collapse of supernovae [\[18\]](#page-36-4). For a detailed physical comprehension of the RT instability, we refer to three survey papers  $[10, 21, 22]$  $[10, 21, 22]$  $[10, 21, 22]$  $[10, 21, 22]$  $[10, 21, 22]$ . Mathematically speaking, for the inviscid and incompressible regime in the neighborhood of a smooth density profile, the classical RT instability was investigated by Lafitte  $[12]$ , Guo and Hwang  $[4]$  and Helffer and Lafitte  $[8]$ .

In this paper, we pay attention to the viscous RT instability. One of the first studies on the viscous RT can be seen in the book of Chandrasekhar [\[1,](#page-35-2) Chapter X] considering two uniform viscous fluid separated by a horizontal boundary and generalizing the classical result of Rayleigh and Taylor. We are concerned with the following Navier–Stokes equation describing the motion of a nonhomogeneous incompressible viscous fluid in the presence of a uniform gravitational field in  $\mathbb{R}^3$ ,

<span id="page-2-2"></span>
$$
\begin{cases}\n\partial_t \rho + \operatorname{div}(\rho \vec{u}) = 0, \\
\partial_t (\rho \vec{u}) + \operatorname{div}(\rho \vec{u} \otimes \vec{u}) + \nabla P = \mu \Delta \vec{u} - \rho \vec{g}, \\
\operatorname{div} \vec{u} = 0,\n\end{cases}
$$
\n(1.1)

where  $t \geq 0, x = (x_1, x_2, x_3) \in \mathbb{R}^3$ . The unknowns  $\rho := \rho(x, t), \, \vec{u} := \vec{u}(x, t)$  and  $P := P(x, t)$  denote respectively the density, the velocity and the pressure of the

<span id="page-2-3"></span>

<span id="page-3-2"></span>fluid, while  $\mu > 0$  is the viscosity coefficient and  $\vec{g} := (0,0,g), g > 0$  being the gravitational constant.

Let  $\rho_0 > 0$  be a  $C^1$ -function depending only on  $x_3$  and let  $P_0$  be another function of  $x_3$  given by  $P'_0 = -g\rho_0$  with  $' = d/dx_3$ . Note that it is not a physical solution of the system, because  $P_0 \to \pm \infty$  when  $x_3 \to \pm \infty$ , but, nevertheless, it is a relevant model to study. Then,  $(\rho, \vec{u}, P) = (\rho_0, \vec{0}, P_0)$  is an equilibrium state of [\(1.1\)](#page-2-2). The linear RT instability of [\(1.1\)](#page-2-2) in the vicinity  $(\rho_0(x_3), \vec{0}, P_0(x_3))$  amounts to the investigation of the parameter  $\lambda \in \mathbb{C}$  (Re $\lambda > 0$ ) such that there exists a bounded solution  $\phi \in H^4(\mathbf{R})$  of the following ordinary differential equation

<span id="page-3-0"></span>
$$
-\lambda^2(\rho_0 k^2 \phi - (\rho_0 \phi')') = \lambda \mu (\phi^{(4)} - 2k^2 \phi'' + k^4 \phi) - g k^2 \rho'_0 \phi, \tag{1.2}
$$

where  $k > 0$  is the transverse wave number *being fixed* and  $\phi$  is the third component of the perturbed velocity. Our desired solution  $\phi$  decays to zero at  $\pm \infty$ , i.e.

<span id="page-3-1"></span>
$$
\lim_{x_3 \to \pm \infty} \phi(x_3) = 0. \tag{1.3}
$$

In this case, such a  $\lambda$  is a growth rate of the instability or is a characteristic value of the linearized problem (see [\[1,](#page-35-2) Chapter X, Sections 92-93]).

The derivation of [\(1.2\)](#page-3-0) will be shown in the next section.

Let us discuss about the mathematical study. For the inviscid problem (i.e.  $\mu = 0$ in [\(1.1\)](#page-2-2) and [\(1.2\)](#page-3-0)), (1.2) can be seen as an eigenvalue problem with eigenvalue  $\lambda^2$ . Multiple eigenvalues of the inviscid problem are first mentioned by Cherfils and Lafitte [\[2\]](#page-35-3) through the precise study of the inviscid ODE for a specific profile. Mention also the paper of Mikaelian [\[14\]](#page-36-9) for the connection between the Rayleigh  $(1.2)$  and the Schrödinger equations. Lafitte  $[12]$  then observed that possible growth rates  $\lambda$ 's for the inviscid Rayleigh-Taylor problem (not only the largest one) are such that  $\lambda^2$  is an eigenvalue of a suitable self-adjoint operator and described this spectrum, for large values of  $k$ , as the eigenvalues of a 1D Schrodinger operator, which is generalized by Helffer and Lafitte [\[8\]](#page-36-0).

On the other side, the natural variational structure of the inviscid problem pointed out in [\[1\]](#page-35-2) and used by Guo and Hwang [\[4\]](#page-35-1) describes the square of the maximum growth rate  $\lambda_1^2$  as the maximum of of the Rayleigh quotient

$$
\frac{g k^2 \int_{\mathbf{R}}\rho_0' |\phi|^2 dx_3}{\int_{\mathbf{R}}\rho_0 (|\phi'|^2+k^2|\phi|^2)dx_3}
$$

and is used to obtain a proof of the nonlinear instability. With this argument, it can be noticed that the multiple eigenvalues are given by  $\{\lambda_2, \ldots, \lambda_j, \ldots\}$ , where  $\lambda_j^2$  is equal to

$$
\max_{\phi\in H_k^1} \frac{gk^2\int_{\mathbf{R}}\rho_0'|\phi|^2 dx_3}{\int_{\mathbf{R}}\rho_0(|\phi'|^2+k^2|\phi|^2)dx_3}
$$

where  $H_k^1 = \{ \phi \in H^1(\mathbf{R}), \int_{\mathbf{R}} \phi \phi_l dx_3 = 0, \forall l \leq j \}, \phi_l$  is a nontrivial eigenfunction associated with  $\lambda_l$ . This Rayleigh quotient cannot be used here due to the presence of the viscosity term.

For the viscous problem, Jiang, Jiang and Ni [\[11\]](#page-36-10) used a modified variational approach, described by Guo and Tice [\[5\]](#page-35-4) , where a bootstrap argument yields the largest growth rate and a corresponding solution being regular under the assumption  $\rho'_0 \in C_0^{\infty}(\mathbf{R})$ ,  $\inf_{\mathbf{R}} \rho_0 > 0$  and  $\rho'_0(x_3) > 0$  for some points  $x_3 \in \mathbf{R}$ .

As far as we could find, no other authors performed studies of the discrete spectrum of the inviscid or the viscous linearized RT instability. We aim at the viscous study in the mathematical spirit of Helffer-Lafitte [\[8\]](#page-36-0).

<span id="page-4-0"></span>We then illustrate our spectral analysis.

Throughout this paper, we consider  $\rho_0$  as an increasing function, i.e.  $\rho'_0 \geq 0$ . For an increasing density profile  $\rho_0$ , we show in [A](#page-31-0)ppendix A that  $\lambda$  is real and bounded by  $\sqrt{\frac{g}{L_0}}$ , with  $L_0^{-1} := \sup_{R} \frac{\rho_0'}{\rho_0}$ . Since our goal is to study the unstable modes, we restrict our study to the case  $\lambda > 0$ .

The ODE [\(1.2\)](#page-3-0) rewrites as a linear system of ODEs on  $(\phi, \phi', \phi'', \phi''')^T$ . As the solution of this system must tend to 0 when  $x_3 \to \pm \infty$ , using the fact that the profile  $\rho_0$  goes to  $\rho_+$  at  $\pm \infty$ , we are able to deduce the stable linear space  $S_{+}$  at  $+\infty$  and the unstable linear space  $S_{-}$  at  $-\infty$ , that are vector spaces of dimension two. Hence, we transform the problem for the normal modes on  **into** an ODE problem stated on a compact interval  $(x_-, x_+)$  with appropriate boundary conditions deduced from the outer solutions. In the case of a compactly supported  $\rho'_0$  (the simplest case of a convergence), they are described by  $(D.14)-(D.15)$  $(D.14)-(D.15)$  $(D.14)-(D.15)$ . In the case of a non compactly supported  $\rho'_0$ , they are described by  $(4.7)-(4.8)$  $(4.7)-(4.8)$  $(4.7)-(4.8)$ . Note that, if  $\rho'_0$  is compactly supported, the natural choice of  $x_{\pm}$  is to choose the ends of  $\text{supp}\rho_0'$ . However, if  $\rho_0'$  non compactly supported,  $x_\pm$  are deduced from the behavior at  $\pm\infty$  of the outer solutions and depend also on *k* and  $\lambda$ .

In order to solve  $(1.2)$  on  $(x_-, x_+)$ , the crucial tool in our study is to construct two bilinear forms, that are continuous and coercive,  $B_{\lambda}$  (denoted respectively by  $B_{a,\lambda}$  [\(3.10\)](#page-10-1) for  $\rho'_0$  being compactly supported and by  $B_{x_-,x_+,\lambda}$  [\(4.15\)](#page-21-0) for  $\rho'_0$  being positive everwhere) such that the finding of bounded solutions of Eq. [\(1.2\)](#page-3-0) on the whole line is equivalent to the existence of solutions to the variational problem

$$
\lambda \mathcal{B}_{\lambda}(\phi,\omega) = g k^2 \int_{x_-}^{x_+} \rho'_0 \phi \omega dx_3 \quad \text{for all } \omega \in H^2((x_-,x_+)).
$$

As  $\mathcal{B}_{\lambda}$  is coercive on  $H^2((x_-, x_+)), \sqrt{\mathcal{B}_{\lambda}(\cdot, \cdot)}$  is a norm on  $H^2((x_-, x_+)),$  and one denotes by  $(H^2((x_-, x_+)))'$  the dual of  $H^2((x_-, x_+))$  for the induced topology. In view of the Riesz representation theorem, we thus define an abstract operator

$$
Y_{\lambda} \in \mathcal{L}(H^2((x_-, x_+)), (H^2((x_-, x_+))')
$$

(see  $Y_{a,\lambda}$  in Proposition [3.3](#page-11-0) for  $\rho'_0 \geq 0$  being compactly supported and  $Y_{x_-,x_+,\lambda}$ in Proposition [4.3](#page-22-0) for  $\rho'_0$  being positive everywhere), associated with the norm  $\sqrt{\mathcal{B}_{\lambda}(\cdot,\cdot)}$  to its dual, such that

$$
\mathcal{B}_{\lambda}(f,\omega) = \langle Y_{\lambda}f,\omega \rangle, \quad \text{for all } \omega \in H^{2}((x_{-},x_{+})) \tag{1.4}
$$

Note that it will be crucial to choose  $x_-, x_+$  such that  $\sqrt{\mathcal{B}_\lambda(\cdot, \cdot)}$  is a norm on  $H^2((x_-, x_+))$ . The existence of bounded solutions of Eq. [\(1.2\)](#page-3-0) on **R** is thus equivalent to finding weak solutions  $\phi \in H^2((x_-, x_+)$  of

$$
\lambda Y_{\lambda}\phi = g k^2 \rho'_0 \phi \text{ in } (H^2((x_-, x_+)))'.
$$

A classical bootstrap argument shows that  $\phi$  belongs to  $H^4((x_-, x_+))$ , from which one deduces that  $\phi$  satisfies [\(1.2\)](#page-3-0) on  $(x_-, x_+)$  (in the sense of elements of  $H^4((x_-, x_+)),$ and the boundary conditions [\(D.14\)](#page-1-4)-[\(D.15\)](#page-1-5) or [\(4.7\)](#page-18-0)-[\(4.8\)](#page-18-1) hold. Note that, as  $\phi$  belongs to  $H^4((x_-, x_+))$ , these boundary conditions (involving derivatives  $\phi'', \phi'''$  of  $\phi$  at  $x_+$ ) are well defined.

We further rewrite the problem, after introducing  $M$  the operator of multiplication by  $\sqrt{\rho'_0}$  in  $L^2((x_-, x_+))$ , as finding *v* (which is  $\mathcal{M}^{-1}Y_\lambda \phi$ ) satisfying

$$
\mathcal{M}Y_{\lambda}^{-1}\mathcal{M}v = \frac{\lambda}{gk^2}v.
$$

<span id="page-5-1"></span>We show that  $\mathcal{M}Y_{\lambda}^{-1}\mathcal{M}$  is self-adjoint and compact, which enables to use the theory of self-adjoint and compact operators on the functional space  $H^2((x_-, x_+))$ . It is a generalization of a Sturm-Liouville problem. The discrete spectrum of the operator  $\mathcal{M}Y_{\lambda}^{-1}\mathcal{M}$  is thus an infinite sequence of eigenvalues, denoted by  $\gamma_n(\lambda)$  and let  $v_n$ be an eigenfunction associated with  $\gamma_n(\lambda)$ .

The problem of finding characteristic values of [\(1.2\)](#page-3-0) amounts to solving all the equations

<span id="page-5-0"></span>
$$
\gamma_n(\lambda) = \frac{\lambda}{gk^2}.\tag{1.5}
$$

When  $\rho'_0 \ge 0$  is compactly supported, for each *n*, we will show the existence and uniqueness of a solution  $\lambda_n$  to [\(1.5\)](#page-5-0) owing first to the differentiability in  $\lambda$  of  $\gamma_n(\lambda)$ (see Lemma [3.2\)](#page-14-0), which is an easy extension of Kato's perturbation theory of [\[9\]](#page-36-11), and secondly on the fact that  $\lambda \to \gamma_n(\lambda)$  is decreasing in  $\lambda$ , through the derivative  $\frac{d}{d\lambda}(\frac{1}{\gamma_n(\lambda)})$  (see Lemma [3.3\)](#page-14-1) which exists also thanks to a similar argument of [\[9\]](#page-36-11). Then  $\phi_n = Y_{\lambda_n}^{-1} \mathcal{M} v_n \in H^4((x_-, x_+))$  is glued with the decaying solutions of [\(1.2\)](#page-3-0) in the outer regions by the boundary conditions at  $x_{\pm}$ , which yields a solution of  $(1.2)$ in  $H^4(\mathbf{R})$  associated with  $\lambda_n$ . Hence, we obtain in our first result, Theorem [2.1,](#page-7-0) the existence of an infinite sequence of characteristic values  $(\lambda_n)_{n\geq 1}$ , that decreases towards 0 as  $n \to \infty$ .

When  $\rho'_0 > 0$  everywhere, unlike the first case, we lack the analytical expression of boundary conditions. We thus do not have the decrease of  $\gamma_n$  or any control of  $\gamma_n(\lambda)$  when  $\lambda$  goes to 0. Consequently, for  $\rho'_0 > 0$  everywhere, our arguments only lead to a multiple existence of positive characteristic values  $\lambda$  such that  $\lambda \geq \epsilon_{\star} > 0$ . This is our second result, Theorem [2.2.](#page-7-1)

We then continue our operator method in other contexts. The first one [\[15\]](#page-36-12) is to investigate the nonlinear Rayleigh-Taylor instability of the gravity-driven incompressible Navier-Stokes equations with Navier-slip boundary conditions in a 2D slab domain  $\Omega = 2\pi L \mathbb{T} \times (-1, 1)$  with  $L > 0$  and T is the usual 1D-torus. Based on infinitely unstable modes of the linearized problem, we consider a wide class of initial data to the nonlinear perturbation problem, extending Grenier's framework [\[6\]](#page-35-5), to prove nonlinear Rayleigh-Taylor instability.

Furthermore, as noted in [\[1,](#page-35-2) Chapter X, Section 94(e)], we study the gravity waves problem that occur in an infinitely deep ocean and the occupied domain is bounded above by a moving interface  $\{(t, x) \in \mathbf{R}_+ \times 2\pi L\mathbb{T} \times \mathbf{R} | x_2 = \eta(t, x_1)\}.$ The governing equation is the gravity-driven incompressible Navier–Stokes equation without any effect of surface tension on the free surface. The precise study will be derived in a forthcoming work [\[16\]](#page-36-13).

We organize this paper as follows. In Section [2,](#page-6-0) we describe the physical model and then give the statement of our main theorems. Section [3](#page-8-0) and Section [4](#page-17-0) are devoted to proving Theorem [2.1](#page-7-0) as  $\rho_0$  satisfies [\(2.6\)](#page-6-4)-[\(2.7\)](#page-7-2) and proving Theorem [2.2](#page-7-1) as  $\rho_0$  satisfies  $(2.8)-(2.9)$  $(2.8)-(2.9)$  $(2.8)-(2.9)$ , respectively.

**Notation.** Throughout this paper,  $C^*$  is a generic constant depending on  $\rho_0$  and other physical parameters, independent of  $\lambda$ .

#### 2. The main results

<span id="page-6-7"></span><span id="page-6-1"></span><span id="page-6-0"></span>2.1. **Derivation of the physical model.** Recall that  $(\rho, \vec{u}, P) = (\rho_0, \vec{0}, P_0)$  is an equilibrium state of  $(1.1)$ . The quantities

$$
\sigma = \rho - \rho_0, \quad \vec{u} = \vec{u} - \vec{0}, \quad p = P - P_0
$$

satisfy the following nonlinear perturbation equations

$$
\begin{cases}\n\partial_t \sigma + \vec{u} \cdot \nabla (\rho_0 + \sigma) = 0, \\
(\rho_0 + \sigma) \partial_t \vec{u} + (\rho_0 + \sigma) \vec{u} \cdot \nabla \vec{u} + \nabla p = \mu \Delta \vec{u} - \sigma \vec{g}, \\
\text{div}\vec{u} = 0.\n\end{cases}
$$
\n(2.1)

That implies the following linearized system

$$
\begin{cases}\n\partial_t \sigma + \rho'_0 u_3 = 0, \\
\rho_0 \partial_t \vec{u} + \nabla p = \mu \Delta \vec{u} - \sigma \vec{g}, \\
\text{div}\vec{u} = 0.\n\end{cases}
$$
\n(2.2)

Since  $\rho_0$  depends only on  $x_3$ , we continue the analysis into normal modes as in [\[1,](#page-35-2) Chapter X, Section 91]. Precisely, we seek the perturbations of the form

$$
(\sigma, \vec{u}, p)(t, x) = e^{\lambda t + ik_1 x_1 + ik_2 x_2} (\zeta, -i\psi, -i\theta, \phi, q)(x_3),
$$
\n(2.3)

where  $k \in \mathbb{R}^2$ ,  $\lambda \in \mathbb{C} \setminus \{0\}$  and  $\text{Re}\lambda \geq 0$ . Let  $k = \sqrt{k_1^2 + k_2^2}$ , we arrive at the following system

<span id="page-6-5"></span>
$$
\begin{cases}\n\lambda \zeta + \rho_0' \phi = 0, \\
\lambda \rho_0 \psi - k_1 q + \mu (k^2 \psi - \psi'') = 0, \\
\lambda \rho_0 \theta - k_2 q + \mu (k^2 \theta - \theta'') = 0, \\
\lambda \rho_0 \phi + q' + \mu (k^2 \phi - \phi'') + g \zeta = 0, \\
k_1 \psi + k_2 \theta + \phi' = 0.\n\end{cases}
$$
\n(2.4)

We directly see  $\zeta = -\frac{\rho'_0 \phi}{\lambda}$ . Hence,  $(2.4)_4$  $(2.4)_4$  becomes

<span id="page-6-6"></span>
$$
\lambda^2 \rho_0 \phi + \lambda q' + \lambda \mu (k^2 \phi - \phi') = g \rho'_0 \phi.
$$
 (2.5)

We multiply  $(2.4)_{2}$  $(2.4)_{2}$  by  $k_1$  and  $(2.4)_{3}$  by  $k_2$ , then use  $(2.4)_{4}$  to obtain the equality

$$
\lambda^2 \rho_0 \phi' + k^2 \lambda q + \lambda \mu (k^2 \phi' - \phi''') = 0.
$$

Deriving this equation, and replacing  $\lambda q'$  thanks to [\(2.5\)](#page-6-6), we get the fourth-order ordinary equation  $(1.2)$ . The investigation of multiple growing normal modes  $(D.13)$ amounts to finding regular solutions  $\phi \in H^4(\mathbf{R})$  of [\(1.2\)](#page-3-0). These solutions decay to zero at  $\pm \infty$ , i.e.  $\phi$  satisfies [\(1.3\)](#page-3-1).

<span id="page-6-2"></span>2.2. Main results. Assuming the density profile  $\rho_0$  is increasing, we first show that all characteristic values  $\lambda$  are real and obtain the following upper bound  $\sqrt{\frac{g}{L_0}}$ of  $\lambda$ . The proof is postponed to Appendix [A.](#page-31-0)

<span id="page-6-3"></span>**Lemma 2.1.** All characteristic values  $\lambda$  for Eq. [\(1.2\)](#page-3-0)-[\(1.3\)](#page-3-1) in  $H^4(\mathbf{R})$  are always *real and satisfy that*  $\lambda \leqslant \sqrt{\frac{g}{L_0}}$ .

In view of Lemma [2.1,](#page-6-3) we seek for functions  $\phi$  being real and we only consider vector space of real functions in what follows.

<span id="page-6-4"></span>In the case  $\rho'_0 \geq 0$  compactly supported, our assumption is

 $\rho'_0$  is a nonnegative function of class  $C_0^0(\mathbf{R})$ ,  $\text{supp}(\rho'_0) = [-a, a]$ , (2.6)

Outside  $(-a, a)$ , we denote

<span id="page-7-2"></span>
$$
\rho_0(x_3) = \begin{cases} \rho_- & \text{as } x_3 \in (-\infty, -a], \\ \rho_+ & \text{as } x_3 \in [a, +\infty), \end{cases}
$$
 (2.7)

with  $\rho$   $\lt$   $\rho$  + are two positive constants. This can be seen, physically speaking, as the situation of the toy model with a layer, of size 2*a*, in which there is a mixture of the two fluids of density  $\rho_{-}$  and  $\rho_{+}$ . We have the following result in Section [3.](#page-8-0)

<span id="page-7-0"></span>**Theorem 2.1.** Let  $\rho_0$  satisfy [\(2.6\)](#page-6-4) and [\(2.7\)](#page-7-2). There exist an infinite sequence  $(\lambda_n, \phi_n)_{n \geq 1}$  *with*  $\lambda_n \in (0, \sqrt{\frac{g}{L_0}})$  *and*  $\phi_n \in H^4(\mathbf{R})$  *satisfying* [\(1.2\)](#page-3-0)*. In addition,*  $\lambda_n$ *decreases towards*  $\theta$  *as*  $\eta$  *goes to*  $\infty$ *.* 

We divide the proof of Theorem [2.1](#page-7-0) into various steps. Notice that the rapid convergence is not necessary in this case because  $\rho_0$  is constant on  $(-\infty, -a)$  and on  $(a, +\infty)$ . Hence, in this case  $\phi$  can be found explicitly since [\(1.2\)](#page-3-0) outside  $(-a, a)$  is an ODE with constant coefficients, that will shown in Proposition [3.1.](#page-8-2) After that, we deduce two boundary conditions [\(D.14\)](#page-1-4) and [\(D.15\)](#page-1-5) at  $\mp a$  respectively in Lemma [3.1.](#page-9-0) Eq. [\(1.2\)](#page-3-0) becomes a boundary-value problem on the finite interval  $(-a, a)$ . Our method to solve that is based on the operator approach, namely the theory of compact and self-adjoint operator for  $Y_{a,\lambda}$ , as illustrated in the introduction part.

In the next part, Section [4,](#page-17-0) we consider  $\rho'_0$  no longer compactly supported. The assumptions on  $\rho_0$  are

<span id="page-7-3"></span>
$$
\rho_0 \in C^1(\mathbf{R}), \quad \lim_{x_3 \to \pm \infty} \rho_0(x_3) = \rho_{\pm} \in (0, +\infty)
$$
\n(2.8)

and

<span id="page-7-4"></span>
$$
0 < \rho_0'(x_3) < \rho_m < +\infty \text{ for all } x_3 \in \mathbf{R}.\tag{2.9}
$$

The second theorem is as follows.

<span id="page-7-1"></span>**Theorem 2.2.** Let  $\rho_0$  satisfy [\(2.8\)](#page-7-3) and [\(2.9\)](#page-7-4). For  $0 < \epsilon_{\star} \ll 1$ , there exists  $N(\epsilon_{\star}) \in$  $\mathbb{N}^*$  such that there are at least  $N(\epsilon_{\star})$  values of  $\lambda \in [\epsilon_{\star}, \sqrt{\frac{g}{L_0}}]$  such that  $\phi \in H^4(\mathbf{R})$ *satisfying* [\(1.2\)](#page-3-0)*.*

The proof of Theorem [2.2](#page-7-1) remains the same to that one of the first case, but more complicated. We point out the main differences as follows.

Questions concerning the existence of solutions of  $(1.2)$  being bounded at  $\infty$  are not trivial as in the first case. In Section [4,](#page-17-0) we transform Eq.  $(1.2)$  into a system of ODEs [\(4.3\)](#page-17-2). The matrix  $L(x_3, \lambda)$  has 4 eigenvalues  $\pm k$  and  $\pm \sqrt{k^2 + \lambda \rho_0(x_3)/\mu}$ , that are different for all  $\lambda > 0$ . We then follow [\[3,](#page-35-6) Theorem 8.1, Chapter 3]. whose statement given in Appendix [C,](#page-33-0) to deduce that  $(1.2)$  admits two linearly independent solutions decaying to 0 at  $+\infty$  (respectively  $-\infty$ ). A suitable interval  $(x_{-}, x_{+})$  is thus determined through a precise calculation of the family of solutions decaying exponentially to zero at  $\pm \infty$ , which yield appropriate boundary conditions  $(4.7)$  at  $x_{-}$  and  $(4.8)$  at  $x_{+}$  in Proposition [4.1.](#page-18-2)

We then solve  $(1.2)$  on a finite interval  $(x_-, x_+)$  with boundary conditions  $(4.7)$ – [\(4.8\)](#page-18-1). To do that, in Section [4.2,](#page-20-0) we construct the bilinear form  $\mathcal{B}_{x_-,x_+\lambda}$  in Proposition [4.2](#page-21-1) and continue the same arguments as in Section [3](#page-8-0) to obtain solution in the inner region  $(x_-, x_+)$ . Note that the coercivity of  $\mathcal{B}_{x_-,x_+\lambda}$  relies on the positivity of the terms  $BV_{x+\lambda}$  and  $BV_{x-\lambda}$  stated in Lemma [4.1.](#page-20-1) Due to the lack of analytical expression of boundary conditions in this case, it turns out that the positivity of  $BV_{x_+,\lambda}$  and  $BV_{x_-, \lambda}$  will be derived, in Proposition [4.2,](#page-21-1) by deducing the behavior at  $\pm \infty$  of coefficients  $n_{ij}^{\pm}$  (*i, j* = 1, 2) depending on ( $x_{\pm}$ ,  $\lambda$ ) and appearing in the

boundary conditions [\(4.7\)](#page-18-0)-[\(4.8\)](#page-18-1). Having the bilinear form  $\mathcal{B}_{x_{-},x_{+},\lambda}$ , we continue our arguments in Propositions [4.3](#page-22-0) and [4.4,](#page-23-1) that follows the same line of Section [3,](#page-8-0) to prove Theorem [2.2.](#page-7-1)

Note that, Lemma [4.1](#page-20-1) does not give any control on  $x_{-}$  and  $x_{+}$ . It is interesting for computational purposes as well as for a study of particular profiles (for example profiles decaying exponentially to their limit at  $\pm \infty$ ), to be able to derive an explicit interval on which this is true. Notice that the restriction  $\lambda \geq \epsilon_{\star} > 0$  of Theorem [2.2](#page-7-1) implies that the matrix  $R(\lambda)$  (see [\(4.2\)](#page-17-3)) in Eq. [\(4.3\)](#page-17-2) becomes regular. Hence, for profile  $\rho'_0 > 0$  everywhere, we devote Section [4.4](#page-24-0) to establish a control of  $x_-\,$  and  $x_+$  independent of  $\lambda$ . In Propositions [4.7](#page-28-0) and [4.8,](#page-29-0) through a careful construction of Volterra series, we can obtain refined estimates of bounded solutions of [\(4.3\)](#page-17-2) at  $\infty$  uniformly in  $\lambda \in [\epsilon_{\star}, \sqrt{\frac{g}{L_0}}]$ . They allow us to have refined estimates on the coefficients  $n_{ij}^{\pm}$   $(i, j = 1, 2)$  uniformly in  $\lambda \in [\epsilon_{\star}, \sqrt{\frac{g}{L_0}}]$ . Hence, we obtain a criterion for  $x_{-}$  and  $x_{+}$  in Proposition [4.5](#page-24-1) to fulfill the conditions of Lemma [4.1](#page-20-1) and extend Proposition [4.2.](#page-21-1)

#### 3. The compactly supported profile

<span id="page-8-0"></span>In this section, we consider  $\rho_0$  satisfying [\(2.6\)](#page-6-4) and [\(2.7\)](#page-7-2). We remark that in this section, we use the notations  $\nu_{\pm} = \frac{\rho_{\pm}}{\mu}$  and  $\tau_{\pm}(\lambda) = (k^2 + \lambda \nu_{\pm})^{1/2}$  and throughout this paper, we will use  $x$  instead of  $x_3$  for notational conveniences.

<span id="page-8-1"></span>3.1. The solution in outer regions and reduction to a problem on a finite **interval.** We derive, in this subsection, the precise expression of  $\phi(x)$  as  $|x| \ge a$ .

<span id="page-8-2"></span>Proposition 3.1. *There are two linearly independent solutions of* [\(1.2\)](#page-3-0) *decaying to 0 at*  $+\infty$  *as*  $x \in [a, +\infty)$ , *i.e.* 

<span id="page-8-3"></span>
$$
\phi_1^+(x) = e^{-kx}
$$
 and  $\phi_2^+(x) = e^{-\tau_+(\lambda)x}$ . (3.1)

*and two linearly independent solutions of*  $(1.2)$  *decaying to 0 at*  $-\infty$  *as*  $x \in (-\infty, -a]$ *, i.e.*

$$
\phi_1^-(x) = e^{kx} \quad \text{and} \quad \phi_2^-(x) = e^{\tau_-(\lambda)x}.\tag{3.2}
$$

*All solutions decaying to 0 at*  $+\infty$  *(respectively at*  $-\infty$ *) are spanned by*  $(\phi_1^+, \phi_2^+)$ *(respectively by*  $(\phi_1^- , \phi_2^-)$ *).* 

*Proof.* For  $x \in [a, +\infty)$ , [\(1.2\)](#page-3-0) reduces to

$$
-\lambda \nu_+(k^2 \phi - \phi'') = \phi^{(4)} - 2k^2 \phi'' + k^4 \phi.
$$
\n(3.3)

We seek  $\phi$  as  $\phi(x) = e^{rx}$ . Hence,

$$
-\lambda \nu_+(k^2 - r^2) = r^4 - 2k^2r^2 + k^4,
$$

which yields  $r = \pm k$  or  $r = \pm (k^2 + \lambda \nu_+)^{1/2}$ . Since  $\phi$  tends to 0 at  $+\infty$ , we get two linearly independent solutions

$$
\phi_1^+(x) = e^{-kx}
$$
 and  $\phi_2^+(x) = e^{-\tau_+(\lambda)x}$ .

Hence, all solutions  $\phi$  decaying to 0 at  $+\infty$  are of the form

<span id="page-8-4"></span>
$$
\phi(x) = A_1^+ e^{-k(x-a)} + A_2^+ e^{-\tau_+(\lambda)(x-a)} \tag{3.4}
$$

for all  $x \in [a, +\infty)$  and for some real constants  $A_1^+$  and  $A_2^+$ .

If  $x \in (-\infty, -a]$ , the same calculation implies [\(3.1\)](#page-8-3). Then, all solutions  $\phi$  decaying to 0 at  $-\infty$  are of the form

<span id="page-9-1"></span>
$$
\phi(x) = A_1^- e^{k(x+a)} + A_2^- e^{\tau_-(\lambda)(x+a)} \tag{3.5}
$$

for all  $x \in (-\infty, -a]$  and for some real constants  $A_1^-$  and  $A_2^ \frac{1}{2}$ .

Once it is proven that  $\phi(x)$  outside  $(-a, a)$  is of the form  $(3.4)$  and  $(3.5)$ , we search for  $\phi$  on  $(-a, a)$ . That solution has to match with [\(3.4\)](#page-8-4) and [\(3.5\)](#page-9-1) well, i.e. there is a condition on  $(\phi, \phi', \phi'', \phi'')$  at  $x = \pm a$ . We will show the conditions in the following lemma.

<span id="page-9-0"></span>**Lemma 3.1.** *The boundary condition of* [\(1.2\)](#page-3-0) *at*  $x = -a$ , for  $\phi \in H^4(\mathbf{R})$ , *is* 

$$
\begin{cases}\nk\tau_{-}\phi(-a) - (k + \tau_{-}(\lambda))\phi'(-a) + \phi''(-a) = 0, \\
k\tau_{-}(\lambda)(k + \tau_{-}(\lambda))\phi(-a) - (k^{2} + k\tau_{-}(\lambda) + \tau_{-}^{2}(\lambda))\phi'(-a) + \phi'''(-a) = 0.\n\end{cases}
$$
\n(3.6)

*and*  $at x = a$  *is* 

$$
\begin{cases}\nk\tau_{+}(\lambda)\phi(a) + (k + \tau_{+}(\lambda))\phi'(a) + \phi''(a) = 0, \\
-k\tau_{+}(\lambda)(k + \tau_{+}(\lambda))\phi(a) - (k^{2} + k\tau_{+}(\lambda) + \tau_{+}^{2}(\lambda))\phi'(a) + \phi'''(a) = 0.\n\end{cases}
$$
\n(3.7)

*Proof.* The boundary condition of the solutions  $\phi$  of [\(1.2\)](#page-3-0) at  $x = \pm a$  is equivalent to the fact that  $\phi$  belongs to the space of decaying solutions at  $\pm \infty$ . On the one hand, it can be seen from  $(3.4)$  and  $(3.5)$  that

$$
\begin{pmatrix}\n\phi(x) \\
\phi'(x) \\
\phi''(x) \\
\phi'''(x)\n\end{pmatrix} = A_1^- e^{k(x+a)} \begin{pmatrix}\n1 \\
k \\
k^2 \\
k^3\n\end{pmatrix} + A_2^- e^{\tau_-(\lambda)(x+a)} \begin{pmatrix}\n1 \\
\tau_-(\lambda) \\
\tau_2^2(\lambda) \\
\tau_2^3(\lambda)\n\end{pmatrix} \quad \text{for } x \leq -a
$$

and that

$$
\begin{pmatrix} \phi(x) \\ \phi'(x) \\ \phi''(x) \\ \phi'''(x) \end{pmatrix} = A_1^+ e^{-k(x-a)} \begin{pmatrix} 1 \\ -k \\ k^2 \\ -k^3 \end{pmatrix} + A_2^+ e^{-\tau_+(\lambda)(x-a)} \begin{pmatrix} 1 \\ -\tau_+(\lambda) \\ \tau_+^2(\lambda) \\ -\tau_+^3(\lambda) \end{pmatrix}, \quad \text{for } x \ge a.
$$

On the other hand, the orthogonal complement of the subspace  $S_{-}$  of  $\mathbb{R}^{4}$  spanned by two vectors  $(1, k, k^2, k^3)^T$  and  $(1, \tau_-(\lambda), \tau_-(\lambda), \tau_-(\lambda))^T$  is spanned by

$$
(k\tau_{-}(\lambda), -(k+\tau_{-}(\lambda)), 1, 0)^{T}
$$
 and  $(k\tau_{-}(\lambda)(k+\tau_{-}(\lambda)), -(k^{2}+k\tau_{-}(\lambda)+\tau_{-}(\lambda)^{2}), 0, 1)^{T}$ .

Similarly, the orthogonal complement of the subspace  $S_+$  of  $\mathbb{R}^4$  spanned by two vectors  $(1, -k, k^2, -k^3)^T$  and  $(1, -\tau_+(\lambda), \tau_+^2(\lambda), -\tau_+^3(\lambda))$  is spanned by two vectors  $(k\tau_+(\lambda), k+\tau_+(\lambda), 1, 0)^T$  and  $(-k\tau_+(\lambda)(k+\tau_+(\lambda)), -(k^2+k\tau_+(\lambda)+\tau_+^2(\lambda)), 0, 1)^T$ .

The above arguments allow us to set  $(D.14)$  and  $(D.15)$  as boundary conditions of Eq.  $(1.2)$  on  $(-a, a)$ .

Remark that *S*<sup> $-$ </sup> is also spanned by  $\{(1, k, k^2, k^3)^T, (0, 1, k + \tau_-(\lambda), k^2 + k\tau_-(\lambda) + \tau_-(\lambda)\}$  $(\tau^2(\lambda))^T$ , and that  $S_+$  is also spanned by  $\{(1, -k, k^2, -k^3)^T, (0, 1, k - \tau_+(\lambda), k^2, -k^3)\}$  $k\tau_{+}(\lambda) + \tau_{+}^{2}(\lambda)$ <sup>T</sup>. This choice of generating families yields two uniformly independent families when  $\lambda \to 0$ . We aim at solving  $(1.2)$  on  $(-a, a)$  with the boundary conditions  $(D.14)$ - $(D.15)$ .

<span id="page-10-0"></span>3.2. A bilinear form and a self-adjoint invertible operator. We introduce the bilinear form  $\mathcal{B}_{a,\lambda}$  in the following proposition.

<span id="page-10-5"></span>Proposition 3.2. *Let us denote by*

$$
BV_{-a,\lambda}(\vartheta,\varrho) := \mu \begin{pmatrix} k\tau_{-}(\lambda)(k+\tau_{-}(\lambda))\vartheta(-a)\varrho(-a) - k\tau_{-}(\lambda)\vartheta'(-a)\varrho(-a) \\ -k\tau_{-}(\lambda)\vartheta(-a)\varrho'(-a) + (k+\tau_{-}(\lambda))\vartheta'(-a)\varrho'(-a) \end{pmatrix}
$$
(3.8)

*and by*

$$
BV_{a,\lambda}(\vartheta,\varrho) := \mu \begin{pmatrix} k\tau_+(\lambda)(k+\tau_+(\lambda))\vartheta(a)\varrho(a) - k\tau_+(\lambda)\vartheta'(a)\varrho(a) \\ -k\tau_+(\lambda)\vartheta(a)\varrho'(a) + (k+\tau_+(\lambda))\vartheta'(a)\varrho'(a) \end{pmatrix}.
$$
 (3.9)

*Then,*

<span id="page-10-1"></span>
$$
\mathcal{B}_{a,\lambda}(\vartheta,\varrho) := BV_{a,\lambda}(\vartheta,\varrho) + BV_{-a,\lambda}(\vartheta,\varrho) + \lambda \int_{-a}^{a} \rho_0(k^2 \vartheta \varrho + \vartheta' \varrho') dx + \mu \int_{-a}^{a} (\vartheta'' \varrho'' + 2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx.
$$
\n(3.10)

*is a continuous and coercive bilinear form on*  $H^2((-a, a))$ .

*Furthermore, let*  $(H^2((-a, a)))'$  *be the dual space of*  $H^2((-a, a))$  *associated with the norm*  $\sqrt{\mathcal{B}_{a,\lambda}(\cdot,\cdot)}$ *, there exists a unique operator* 

$$
Y_{a,\lambda} \in \mathcal{L}(H^2((-a,a)), (H^2((-a,a))))'),
$$

*that is also bijective, such that*

<span id="page-10-4"></span>
$$
\mathcal{B}_{a,\lambda}(\vartheta,\varrho)=\langle Y_{a,\lambda}\vartheta,\varrho\rangle\tag{3.11}
$$

*for all*  $\vartheta, \varrho \in H^2((-a, a)).$ 

*Proof.* Clearly,  $\mathcal{B}_{a,\lambda}$  is a bilinear form on  $H^2((-a, a))$  since the terms  $BV_{+a,\lambda}(\vartheta, \varrho)$ are well defined. We then establish the boundedness of  $\mathcal{B}_{a,\lambda}$ . The integral terms of  $B_{a,\lambda}$  are bounded by

<span id="page-10-2"></span>
$$
\max \left(\rho_+ k^2 \sqrt{\frac{g}{L_0}} + \mu k^4, \rho_+ \sqrt{\frac{g}{L_0}} + 2\mu k^2, \mu\right) \|\vartheta\|_{H^2((-a,a))} \|\varrho\|_{H^2((-a,a))}.\tag{3.12}
$$

About the two first terms  $BV_{\pm a,\lambda}(\vartheta,\varrho)$ , it follows from the general Sobolev inequality that

$$
\|\vartheta(y)\|_{C^{0,j}((-a,a))} \leqslant C(j,a)\|\vartheta(y)\|_{H^1((-a,a))} \quad \text{for all } j \in [0,1/2).
$$

Therefore, we obtain

$$
|\vartheta(a)|^2 + |\vartheta(-a)|^2 \leq C^* \|\vartheta\|_{H^1((-a,a))}^2.
$$

Similarly,

$$
|\vartheta'(a)|^2 + |\vartheta'(-a)|^2 \leq C^* \|\vartheta'\|_{H^1((-a,a))}^2.
$$

<span id="page-10-3"></span>Consequently, we get

$$
|BV_{\pm a,\lambda}(\vartheta,\varrho)| \leq C^{\star}(|\vartheta(\pm a)| + \vartheta'(\pm a)|)(|\varrho(\pm a)| + |\varrho'(\pm a)|)
$$
  

$$
\leq C^{\star} \|\vartheta\|_{H^{2}((-a,a))} \|\varrho\|_{H^{2}((-a,a))}.
$$
 (3.13)

In view of  $(3.12)$  and  $(3.13)$ , we find that

$$
|\mathcal{B}_{a,\lambda}(\vartheta,\varrho)| \leq C^{\star} \|\vartheta\|_{H^2((-a,a))} \|\varrho\|_{H^2((-a,a))},
$$
\n(3.14)

i.e.  $\mathcal{B}_{a,\lambda}$  is bounded.

We move to show the coercivity of  $\mathcal{B}_{a,\lambda}$ . We have that

$$
\mathcal{B}_{a,\lambda}(\vartheta,\vartheta) = BV_{a,\lambda}(\vartheta,\vartheta) + BV_{-a,\lambda}(\vartheta,\vartheta) + \lambda \int_{-a}^{a} \rho_0 (k^2 |\vartheta|^2 + |\vartheta'|^2) dx
$$

$$
+ \mu \int_{-a}^{a} (|\vartheta''|^2 + 2k^2 |\vartheta'|^2 + k^4 |\vartheta|^2) dx.
$$

 $BV_{a,\lambda}(\vartheta,\vartheta) \geq 0$  follows from the following equality

$$
\frac{1}{\mu}BV_{a,\lambda}(\vartheta,\vartheta) = k\tau_{+}(\lambda)(k + \tau_{+}(\lambda))|\vartheta(a)|^{2} + 2k\tau_{+}(\lambda)\vartheta(a)\vartheta'(a) \n+ (k + \tau_{+}(\lambda))|\vartheta'(a)|^{2} \n= k\tau_{+}(\lambda)(k + \tau_{+}(\lambda))\left|\vartheta(a) + \frac{\vartheta'(a)}{k + \tau_{+}(\lambda)}\right|^{2} \n+ \frac{k^{2} + k\tau_{+}(\lambda) + \tau_{+}^{2}(\lambda)}{k + \tau_{+}(\lambda)}|\vartheta'(a)|^{2}.
$$

We also obtain that  $BV_{-a,\lambda}(\vartheta,\vartheta) \geq 0$ . Therefore, we deduce that

$$
\mathcal{B}_{a,\lambda}(\vartheta,\vartheta) \geq \mu \min(k^4, 2k^2, 1) \|\vartheta\|_{H^2((-a,a))}^2.
$$
\n(3.15)

It then tells us that  $\mathcal{B}_{a,\lambda}$  is a continuous and coercive bilinear form on  $H^2((-a, a))$ . It follows from Reisz's representation theorem that there is a unique operator  $Y_{a,\lambda}$  $\mathcal{L}(H^2((-a,a)), (H^2((-a,a)))'$ , that is also bijective, satisfying [\(3.11\)](#page-10-4) for all  $\vartheta, \varrho \in$  $H^2((-a, a))$ . Proof of Proposition [3.2](#page-10-5) is complete.

The next proposition is to devoted to studying the properties of  $Y_{a,\lambda}$ .

<span id="page-11-0"></span>Proposition 3.3. *We have the following results.*

(1) *For all*  $\vartheta \in H^2((-a, a)),$ 

$$
Y_{a,\lambda}\vartheta = \lambda(\rho_0 k^2 \vartheta - (\rho_0 \vartheta')') + \mu(\vartheta^{(4)} - 2k^2 \vartheta'' + k^4 \vartheta)
$$

*in*  $\mathcal{D}'((-a, a))$ .

(2) Let  $f \in L^2((-a, a))$  be given, there exists a unique solution  $\vartheta \in H^2((-a, a))$ *of*

<span id="page-11-3"></span>
$$
Y_{a,\lambda}\vartheta = f \ \ in \ (H^2((-a,a)))',\tag{3.16}
$$

*then*  $\vartheta \in H^4((-a, a))$  *and satisfies the boundary conditions* [\(D.14\)](#page-1-4)–[\(D.15\)](#page-1-5)*.* 

*Proof.* Let  $\varrho \in C_0^{\infty}((-a, a))$ , it follows from Proposition [3.2](#page-10-5) that, for  $\vartheta \in H^2(-a, a)$ 

<span id="page-11-1"></span>
$$
\lambda \int_{-a}^{a} \rho_0(k^2 \vartheta \varrho + \vartheta' \varrho') dx + \mu \int_{-a}^{a} (\vartheta'' \varrho'' + 2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx = \langle Y_{a,\lambda} \vartheta, \varrho \rangle. \tag{3.17}
$$

We respectively define  $(\vartheta'')'$  and  $(\vartheta'')''$  in the distributional sense as the first and second derivative of  $\vartheta''$  which is in  $L^2((-a, a))$ . Hence, [\(3.17\)](#page-11-1) is equivalent to

<span id="page-11-2"></span>
$$
\lambda \int_{-a}^{a} \rho_0(k^2 \vartheta \varrho + \vartheta' \varrho') dx + \mu \int_{-a}^{a} (2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx + \langle (\vartheta'')'', \varrho \rangle = \langle Y_{a,\lambda} \vartheta, \varrho \rangle. \tag{3.18}
$$

for all  $\varrho \in C_0^{\infty}((-a, a))$ . We deduce from  $(3.18)$  that

<span id="page-11-4"></span>
$$
\int_{-a}^{a} (\lambda (k^2 \rho_0 \vartheta - (\rho_0 \vartheta')) + \mu (-2k^2 \vartheta'' + k^4 \vartheta)) \varrho dx + \mu \langle (\vartheta'')'', \varrho \rangle = \langle Y_{a,\lambda} \vartheta, \varrho \rangle
$$
 (3.19)

for all  $\varrho \in C_0^{\infty}((-a, a))$ . The first assertion follows.

Let  $f \in L^2((-a, a))$  and  $\vartheta \in H^2((-a, a))$  be the solution of [\(3.16\)](#page-11-3), we then improve the regularity of the weak solution  $\vartheta$  of [\(3.19\)](#page-11-4). Indeed, we rewrite (3.19) as

<span id="page-12-0"></span>
$$
\mu \langle (\vartheta'')'', \varrho \rangle = \int_{-a}^{a} (f + 2\mu k^2 \vartheta'' - \mu k^4 \vartheta - \lambda k^2 \rho_0 \vartheta + \lambda (\rho_0 \vartheta')') \varrho dx \tag{3.20}
$$

for all  $\varrho \in C_0^{\infty}((-a,a))$ . Since  $(f + 2\mu k^2 \vartheta'' - \mu k^4 \vartheta - \lambda k^2 \rho_0 \vartheta + \lambda(\rho_0 \vartheta')')$  belongs to  $L^2((-a, a))$ , it then follows from [\(3.20\)](#page-12-0) that  $(\vartheta'')'' \in L^2((-a, a))$ . Furthermore, by usual distribution theory, we define  $\Psi \in \mathcal{D}'((-a, a))$  such that

<span id="page-12-1"></span>
$$
\langle \Psi, \varrho \rangle = \langle (\vartheta'')'', \zeta_{\rho} \rangle \tag{3.21}
$$

for all  $\varrho \in C_0^{\infty}$ , where  $\zeta_{\rho}(x) = \int_{-a}^x (\varrho(y) - \int_{-a}^a \varrho(s) ds) dy$ . Hence, it can be seen that

 $\langle \Psi',\varrho\rangle=-\langle \Psi,\varrho'\rangle=-\langle (\vartheta'')'',\zeta_{\rho'}\rangle=-\langle (\vartheta'')'',\varrho\rangle$ 

that implies  $(\vartheta'')' - \Psi \equiv \text{constant}$ . In view of  $(\vartheta'')'' \in L^2((-a, a))$  and [\(3.21\)](#page-12-1), we know that  $(\vartheta'')' \in L^2((-a, a))$ . Since  $\vartheta \in H^2((-a, a))$  and  $(\vartheta'')', (\vartheta'')'' \in L^2((-a, a))$ , it tells us that  $\vartheta$  belongs to  $H^4((-a, a))$ .

By exploiting  $(3.20)$ , we then show that  $\vartheta$  satisfies  $(D.14)$  and  $(D.15)$ . Indeed, consider now  $\rho \in C^{\infty}((-a, a))$ , one has, using the integration by parts

$$
\int_{-a}^{a} (\vartheta'')''(x)\varrho(x)dx = (\vartheta'')'(a)\varrho(a) - (\vartheta'')'(-a)\varrho(-a) - (\vartheta'')(a)\varrho'(a) + (\vartheta'')(-a)\varrho'(-a) + \int_{-a}^{a} \vartheta''(x)\varrho''(x)dx.
$$

We perform on the other terms of  $(3.19)$  the integration by parts, which yields

$$
\lambda \int_{-a}^{a} \rho_0 (k^2 \vartheta \varrho + \vartheta' \varrho') dx + \mu \int_{-a}^{a} (\vartheta'' \varrho'' + 2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx - \lambda \rho_0 \vartheta' \varrho \Big|_{-a}^{a} + \mu (\vartheta''' \varrho \Big|_{-a}^{a} - \vartheta'' \varrho' \Big|_{-a}^{a} - 2k^2 \vartheta' \varrho \Big|_{-a}^{a} \Big) = \int_{-a}^{a} (Y_{a,\lambda} \vartheta) \varrho dx.
$$

It then follows from the definition of the bilinear form  $\mathcal{B}_{a,\lambda}$  that

<span id="page-12-2"></span>
$$
BV_{a,\lambda}(\vartheta,\varrho)+BV_{-a,\lambda}(\vartheta,\varrho)=-\lambda\rho_0\vartheta'\varrho\Big|_{-a}^{a}+\mu\Big(\vartheta'''\varrho\Big|_{-a}^{a}-\vartheta''\varrho'\Big|_{-a}^{a}-2k^2\vartheta'\varrho\Big|_{-a}^{a}\Big) (3.22)
$$

for all  $\rho \in C^{\infty}((-a, a)).$ 

By collecting all terms corresponding to  $\varrho(-a)$  in [\(3.22\)](#page-12-2), we deduce that

$$
\mu(k\tau_{-}(\lambda)(k+\tau_{-}(\lambda))\vartheta(-a) - k\tau_{-}(\lambda)\vartheta'(-a))
$$
  
=  $\lambda \rho_0(-a)\vartheta'(-a) - \mu(\vartheta'''(-a) - 2k^2\vartheta'(-a)).$ 

It yields

$$
\vartheta''(-a) - (k^2 + k\tau)(\lambda) + \tau^2(\lambda)\vartheta'(-a) + k\tau(\lambda)(k + \tau)(\lambda)\vartheta(-a) = 0
$$

owing to the definition of  $\tau_{-}(\lambda)$ . Then, we collect all terms corresponding to  $\varrho(a)$ or to  $\varrho'(\pm a)$  in [\(3.22\)](#page-12-2) to conclude that  $\vartheta$  satisfies [\(D.14\)](#page-1-4) and [\(D.15\)](#page-1-5). This ends the proof of Proposition [3.2.](#page-10-5)  $\Box$ 

We have the following proposition on  $Y_{a,\lambda}^{-1}$ .

<span id="page-12-3"></span>**Proposition 3.4.** *The operator*  $Y_{a,\lambda}^{-1}: L^2((-a,a)) \to L^2((-a,a))$  *is compact and self-adjoint.*

<span id="page-12-4"></span>

<span id="page-13-3"></span>*Proof.* It follows from Proposition [3.3](#page-11-0) that  $Y_{a,\lambda}$  admits an inverse operator  $Y_{a,\lambda}^{-1}$ from  $L^2((-a, a))$  to a subspace of  $H^4((-a, a))$  requiring all elements satisfy [\(D.14\)](#page-1-4)– [\(D.15\)](#page-1-5), which is symmetric due to Proposition [3.2.](#page-10-5) We compose  $Y_{a,\lambda}^{-1}$  with the continuous injection from  $H^4((-a, a))$  to  $L^2((-a, a))$ . Notice that the embedding  $H^p((-a, a)) \hookrightarrow H^q((-a, a))$  for  $p > q \geq 0$  is compact. Therefore,  $Y_{a, \lambda}^{-1}$  is compact and self-adjoint from  $L^2((-a, a))$  to  $L^2((-a, a))$ . Proposition [3.4](#page-12-3) is shown.  $\square$ 

Remark 3.1. *In this paper, we choose to define the operator*

$$
\phi \mapsto \lambda(\rho_0 k^2 \phi - (\rho_0 \phi')') + \mu(\phi^{(4)} - 2k^2 \phi'' + k^4 \phi) = Y_{a,\lambda} \phi
$$

*with boundary conditions* [\(D.14\)](#page-1-4)*-*[\(D.15\)](#page-1-5) *through the Riesz representation theorem. We can also define that by the following way.*

*Ya, is well defined on*

 $D(Y_{a,\lambda}) = \{ \phi \in C^4((-a,a)), \phi \text{ verifies (D.14)} - (D.15) \}$  $D(Y_{a,\lambda}) = \{ \phi \in C^4((-a,a)), \phi \text{ verifies (D.14)} - (D.15) \}$  $D(Y_{a,\lambda}) = \{ \phi \in C^4((-a,a)), \phi \text{ verifies (D.14)} - (D.15) \}$  $D(Y_{a,\lambda}) = \{ \phi \in C^4((-a,a)), \phi \text{ verifies (D.14)} - (D.15) \}$  $D(Y_{a,\lambda}) = \{ \phi \in C^4((-a,a)), \phi \text{ verifies (D.14)} - (D.15) \}$ 

*and that we can extend*  $Y_{a,\lambda}$  *over the closure of*  $D(Y_{a,\lambda})$ *. Furthermore,*  $Y_{a,\lambda}$  *with the domain*  $H^4((-a, a))$  *containing functions that satisfy* [\(D.14\)](#page-1-4)*-*[\(D.15\)](#page-1-5) *is symmetric and positive. It follows from Friedrichs extension (see* [\[7,](#page-36-14) Theorem 4.3.1] *e.g.) that Ya, admits a self-adjoint extension.*

<span id="page-13-0"></span>3.3. **A sequence of characteristic values.** We continue considering  $\lambda \in (0, \sqrt{\frac{g}{L_0}}]$ and study the operator  $S_{a,\lambda} := \mathcal{M}Y_{a,\lambda}^{-1}\mathcal{M}$ , where  $\mathcal M$  is the operator of multiplication by  $\sqrt{\rho_0'}$ . Note that this choice prevents to consider a case where  $\rho_0'$  could be negative.

**Proposition 3.5.** *The operator*  $S_{a,\lambda}: L^2((-a,a)) \to L^2((-a,a))$  *is compact and self-adjoint, under the hypothesis* [\(2.6\)](#page-6-4)*.*

*Proof.* Due to the boundedness of  $\rho'_0$ , the operator  $S_{a,\lambda}$  is well-defined from  $L^2((-a,a))$ to itself.  $Y_{a,\lambda}^{-1}$  is compact, so is  $S_{a,\lambda}$ . Moreover, because both the inverse  $Y_{a,\lambda}^{-1}$  and *M* are self-adjoint, the self-adjointness of  $S_{a,\lambda}$  follows.

As a result of the spectral theory of compact and self-adjoint operators, the point spectrum of  $S_{a,\lambda}$  is discrete, i.e. is a decreasing sequence  $\{\gamma_n(k,\lambda)\}_{n\geq 1}$  of positive eigenvalues of  $S_{a,\lambda}$  that tends to 0 as  $n \to \infty$ , associated with normalized orthogonal eigenvectors  $\{\varpi_n\}_{n\geq 1}$  in  $L^2((-a, a))$ . That means

$$
\gamma_n(\lambda)\varpi_n=\mathcal{M}Y_{a,\lambda}^{-1}\mathcal{M}\varpi_n.
$$

so that with  $\phi_n = Y_{a,\lambda}^{-1} \mathcal{M} \varpi_n \in H^4((-a,a))$ , one has

<span id="page-13-1"></span>
$$
\gamma_n(\lambda)Y_{a,\lambda}\phi_n = \rho'_0\phi_n \tag{3.23}
$$

and  $\phi_n$  satisfies [\(D.14\)](#page-1-4)-[\(D.15\)](#page-1-5). [\(3.23\)](#page-13-1) also tells us that  $\gamma_n(\lambda) > 0$  for all *n*. Indeed, we obtain

$$
\gamma_n(\lambda) \int_{-a}^a (Y_{a,\lambda} \phi_n) \phi_n dx = \int_{-a}^a \rho'_0 |\phi_n|^2 dx.
$$

That implies

<span id="page-13-2"></span>
$$
\gamma_n(\lambda)\mathcal{B}_{a,\lambda}(\phi_n,\phi_n) = \int_{-a}^a \rho'_0 |\phi_n|^2 dx.
$$
\n(3.24)

Since  $\mathcal{B}_{a,\lambda}(\phi_n, \phi_n) > 0$  and  $\rho'_0 > 0$  on  $(-a, a)$ , we know that  $\gamma_n(\lambda)$  is positive.

For each *n*, in order to verify that  $\phi_n$  is a solution of [\(1.2\)](#page-3-0), we are left to look for real values of  $\lambda_n$  such that [\(1.5\)](#page-5-0). To solve (1.5), we have to prove that  $\gamma_n(\lambda)$  is differentiable and decreasing in terms of  $\lambda$ , respectively in two next lemmas.

<span id="page-14-0"></span>**Lemma 3.2.** For each *n*, the functions  $\gamma_n(\lambda)$  and  $\phi_n$  are differentiable in terms of  $\lambda \in (0, \sqrt{\frac{g}{L_0}}].$ 

*Proof.* The family  $(Y_{a,\lambda})_{\lambda \in (0,\sqrt{\frac{g}{L_0}}]}$  is a family of bounded operators owing to Propo-sition [3.4.](#page-12-3) It can be seen that the boundary conditions  $(D.14)-(D.15)$  $(D.14)-(D.15)$  $(D.14)-(D.15)$  differentiable in the parameter  $\lambda$  will tell us that  $(Y_{a,\lambda})_{\lambda \in (0,\sqrt{\frac{g}{L_0}}]}$  is also a family of differentiable operators on  $\lambda$  by following a generalized treatment of [\[9,](#page-36-11) Example 1.15, Chapter VII, \$1.6]. Since  $Y_{a,\lambda}^{-1}$  exists for all  $\lambda \in (0, \sqrt{\frac{g}{L_0}}]$ , it follows from [\[9,](#page-36-11) Theorem 2.23, Chapter IV, \$2.6] that  $Y_{a,\lambda}^{-1}$  is differentiable for all  $\lambda \in (0, \sqrt{\frac{g}{L_0}}]$ , so is  $S_{a,\lambda}$ . We then apply the differentiable property of eigenvalues of self-adjoint and compact opera-tors, demonstrated in Appendix [B](#page-32-0) to deduce that  $\gamma_n(\lambda)$  and  $\phi_n$  are differentiable functions. functions.  $\Box$ 

<span id="page-14-1"></span>**Lemma 3.3.** For each *n*, the function  $\gamma_n(\lambda)$  is decreasing in  $\lambda \in (0, \sqrt{\frac{g}{L_0}}]$ .

*Proof.* Let  $z_n = \frac{d\phi_n}{d\lambda}$ . Note that  $\frac{d}{d\lambda}Y_{a,\lambda}u = \rho_0k^2u - (\rho_0u')'$ . It follows from [\(3.23\)](#page-13-1) that

$$
k^2 \rho_0 \phi_n - (\rho_0 \phi'_n)' + Y_{a,\lambda} z_n = \frac{1}{\gamma_n(\lambda)} \rho'_0 z_n + \frac{d}{d\lambda} \Big(\frac{1}{\gamma_n(\lambda)}\Big) \rho'_0 \phi_n \tag{3.25}
$$

<span id="page-14-2"></span>on  $(-a, a)$ . In addition, we have that at  $x = -a$ ,

<span id="page-14-6"></span>
$$
\begin{cases}\nz_n''(-a) - (k + \tau_-(\lambda))z_n'(-a) + k\tau_-(\lambda)z_n(-a) \\
= \frac{\nu_-}{2\tau_-(\lambda)}\phi_n'(-a) - \frac{k\nu_-}{2\tau_-(\lambda)}\phi_n(-a), \\
z_n'''(-a) - (k^2 + k\tau_-(\lambda) + \tau_-^2(\lambda))z_n'(-a) + k\tau_-(\lambda)(k + \tau_-(\lambda))z_n(-a) \\
= \left(\frac{k\nu_-}{2\tau_-(\lambda)} + \nu_-\right)\phi_n'(-a) - \left(\frac{k^2\nu_-}{2\tau_-(\lambda)} + k\nu_-\right)\phi_n(-a)\n\end{cases} (3.26)
$$

and that at  $x = a$ ,

<span id="page-14-7"></span>
$$
\begin{cases}\nz_n''(a) + (k + \tau_+(\lambda))z_n'(a) + k\tau_+(\lambda)z_n(a) \\
= -\frac{\nu_+}{2\tau_+(\lambda)}\phi_n'(a) - \frac{k\nu_+}{2\tau_+(\lambda)}\phi_n(a), \\
z_n'''(a) - (k^2 + k\tau_+(\lambda) + \tau_+^2(\lambda))z_n'(a) - k\tau_+(\lambda)(k + \tau_+(\lambda))z_n(a) \\
= \left(\frac{k\nu_+}{2\tau_+(\lambda)} + \nu_+\right)\phi_n'(a) + \left(\frac{k^2\nu_+}{2\tau_+(\lambda)} + k\nu_+\right)\phi_n(a).\n\end{cases} \tag{3.27}
$$

Multiplying by  $\psi_n$  on both sides of [\(3.25\)](#page-14-2) and then integrating by parts to obtain that

<span id="page-14-5"></span>
$$
\int_{-a}^{a} (k^2 \rho_0 \phi_n - (\rho_0 \phi'_n)') \phi_n dx + \int_{-a}^{a} (Y_{a,\lambda} z_n) \phi_n dx
$$
  
= 
$$
\frac{1}{\gamma_n(\lambda)} \int_{-a}^{a} \rho'_0 z_n \phi_n dx + \frac{d}{d\lambda} \left(\frac{1}{\gamma_n(\lambda)}\right) \int_{-a}^{a} \rho'_0 |\phi_n|^2 dx.
$$
 (3.28)

Thanks to the integration by parts, we have

<span id="page-14-3"></span>
$$
\int_{-a}^{a} (k^2 \rho_0 \phi_n - (\rho_0 \phi'_n)') \phi_n dx = \int_{-a}^{a} \rho_0 (k^2 |\phi_n|^2 + |\phi'_n|^2) dx - (\rho_0 \phi'_n \phi_n) \Big|_{-a}^{a} \quad (3.29)
$$

and

<span id="page-14-4"></span>
$$
\int_{-a}^{a} (Y_{a,\lambda}z_n)\phi_n dx = \int_{-a}^{a} (Y_{a,\lambda}\phi_n)z_n dx + \left(\mu(z_n'''\phi_n - z_n''\phi_n' - 2k^2z_n'\phi_n) - \lambda\rho_0 z_n'\phi_n\right)\Big|_{-a}^{a} - \left(\mu(\phi_n'''z_n - \phi_n''z_n' - 2k^2\phi_n'z_n) - \lambda\rho_0\phi_n'z_n\right)\Big|_{-a}^{a}.
$$
\n(3.30)

Owing to [\(3.29\)](#page-14-3), [\(3.30\)](#page-14-4) and [\(3.23\)](#page-13-1), [\(3.28\)](#page-14-5) becomes

<span id="page-15-0"></span>
$$
\int_{-a}^{a} \rho_0(k^2 |\phi_n|^2 + |\phi_n'|^2) dx + \left( \mu(z_n'''\phi_n - z_n''\phi_n' - 2k^2 z_n' \phi_n) - \lambda \rho_0 z_n' \phi_n \right) \Big|_{-a}^{a}
$$

$$
- \left( \mu(\phi_n''' z_n - \phi_n'' z_n' - 2k^2 \phi_n' z_n) - \lambda \rho_0 \phi_n' z_n \right) \Big|_{-a}^{a} - (\rho_0 \phi_n' \phi_n) \Big|_{-a}^{a} \quad (3.31)
$$

$$
= \frac{d}{d\lambda} \left( \frac{1}{\gamma_n(\lambda)} \right) \int_{-a}^{a} \rho_0' |\phi_n|^2 dx.
$$

Using  $(3.26)$ , we obtain

$$
-\left(\mu(z_n'''\phi_n - z_n''\phi_n' - 2k^2z_n'\phi_n) - \lambda\rho_0 z_n'\phi_n\right)(-a) + \left(\mu(\phi_n''z_n - \phi_n''z_n' - 2k^2\phi_n'z_n) - \lambda\rho_0\phi_n'z_n\right)(-a) + \rho_-\phi_n'(-a)\phi_n(-a) = \mu\left(\frac{k^2\nu_-}{2\tau_-(\lambda)} + k\nu_-\right)|\phi_n(-a)|^2 - \mu\left(\frac{k\nu_-}{2\tau_-(\lambda)} + \nu_-\right)\phi_n'(-a)\phi_n(-a) - \mu\frac{k\nu_-}{2\tau_-(\lambda)}\phi_n(-a)\phi_n'(-a) + \mu\frac{\nu_-}{2\tau_-(\lambda)}|\phi_n'(-a)|^2 + \rho_-\phi_n'(-a)\phi_n(-a).
$$

Keep in mind that  $\mu\nu = \rho_+$ , one has

<span id="page-15-1"></span>
$$
-\left(\mu(z_n'''\phi_n - z_n''\phi_n' - 2k^2 z_n' \phi_n) - \lambda \rho_0 z_n' \phi_n\right)(-a) + \left(\mu(\phi_n'' z_n - \phi_n'' z_n' - 2k^2 \phi_n' z_n) - \lambda \rho_0 \phi_n' z_n\right)(-a) + \rho_-\phi_n'(-a)\phi_n(-a)
$$
 (3.32)  
=  $k\rho_-|\phi_n(-a)|^2 + \frac{\rho_-}{2\tau_-(\lambda)}|\phi_n'(-a) - k\phi_n(-a)|^2.$ 

Similarly, using [\(3.27\)](#page-14-7), we obtain

<span id="page-15-2"></span>
$$
\left(\mu(z_n'''\phi_n - z_n''\phi_n' - 2k^2 z_n'\phi_n) - \lambda \rho_0 z_n'\phi_n\right)(a) \n- \left(\mu(\phi_n''' z_n - \phi_n'' z_n' - 2k^2 \phi_n' z_n) - \lambda \rho_0 \phi_n' z_n\right)(a) - \rho_-\phi_n'(a)\phi_n(a) \n= k\rho_+ |\phi_n(a)|^2 + \frac{\rho_+}{2\tau_+(\lambda)} |\phi_n'(a) - k\phi_n(a)|^2.
$$
\n(3.33)

Combining  $(3.31)$ ,  $(3.32)$  and  $(3.33)$ , we deduce that

$$
\frac{d}{d\lambda} \left(\frac{1}{\gamma_n(\lambda)}\right) \int_{-a}^{a} \rho'_0 |\phi_n|^2 dx
$$
\n
$$
= \int_{-a}^{a} \rho_0(k^2 |\phi_n|^2 + |\phi'_n|^2) dx + k\rho_- |\phi_n(-a)|^2 + \frac{\rho_-}{2\tau_-(\lambda)} |\phi'_n(-a) - k\phi_n(-a)|^2
$$
\n
$$
+ k\rho_+ |\phi_n(a)|^2 + \frac{\rho_+}{2\tau_+(\lambda)} |\phi'_n(a) - k\phi_n(a)|^2. \tag{3.34}
$$

It yields that  $\frac{1}{\gamma_n(\lambda)}$  is increasing in  $\lambda$ , i.e.  $\gamma_n(\lambda)$  is decreasing in  $\lambda$ . This ends the proof of Lemma  $3.3$ .

Now we are in proposition to solve [\(1.5\)](#page-5-0).

<span id="page-15-3"></span>**Proposition 3.6.** For each *n*, there exists only one positive  $\lambda_n$  satisfying [\(1.5\)](#page-5-0). In *addition,*  $\lambda_n$  *decreases towards* 0 *as n goes to*  $\infty$ *.* 

*Proof.* Using [\(3.24\)](#page-13-2), we know that

$$
\frac{1}{\gamma_n(\lambda)}\int_{-a}^a \rho'_0 |\phi_n|^2 dx = \int_{-a}^a (Y_{a,\lambda}\phi_n)\phi_n dx = \mathcal{B}_{a,\lambda}(\phi_n,\phi_n),
$$

that implies

$$
\frac{1}{L_0 \gamma_n(\lambda)} \ge \lambda k^2 + \frac{\mu k^4}{\rho_+}.
$$

Consequently, for all  $n \geq 1$ ,

<span id="page-16-1"></span>
$$
\lim_{\lambda \to \sqrt{\frac{g}{L_0}}} \frac{\lambda}{\gamma_n(\lambda)} > gk^2.
$$
\n(3.35)

As  $0 < \frac{1}{\gamma_n(\lambda)}$  and it is a increasing function,  $\frac{1}{\gamma_n(\lambda)} \leq \frac{1}{\gamma_n(\frac{1}{2}\sqrt{\frac{g}{L_0}})}$  for all  $\lambda \leq \frac{1}{2}\sqrt{\frac{g}{L_0}}$ . That implies

<span id="page-16-2"></span>
$$
\lim_{\lambda \to 0} \frac{\lambda}{\gamma_n(\lambda)} \le \lim_{\lambda \to 0} \frac{\lambda}{\gamma_n(\frac{1}{2}\sqrt{\frac{g}{L_0}})} = 0 \quad \text{for all } n \ge 1.
$$
 (3.36)

Combining [\(3.35\)](#page-16-1), [\(3.36\)](#page-16-2) and using Lemma [3.3,](#page-14-1) we deduce that there is only one  $\lambda_n \in (0, \sqrt{\frac{g}{L_0}})$  satisfying [\(1.5\)](#page-5-0) for each  $n \geq 1$ .

We then prove that the sequence  $(\lambda_n)_{n\geq 1}$  is decreasing. Indeed, if  $\lambda_m < \lambda_{m+1}$ for some  $m \geq 1$ , we have

$$
\gamma_m(\lambda_m) > \gamma_m(\lambda_{m+1}).
$$

Meanwhile, we also have

$$
\gamma_m(\lambda_{m+1}) > \gamma_{m+1}(\lambda_{m+1}).
$$

That implies

$$
\frac{\lambda_m}{gk^2} = \gamma_m(\lambda_m) > \gamma_{m+1}(\lambda_{m+1}) = \frac{\lambda_{m+1}}{gk^2}.
$$

That contradiction tells us that  $(\lambda_n)_{n\geq 1}$  is a decreasing sequence. Suppose that

$$
\lim_{n \to \infty} \lambda_n = c_0 > 0.
$$

Note that

$$
\gamma_n(c_0) \geq \gamma_n(\lambda_n) = \frac{\lambda_n}{gk^2}.
$$

Let  $n \to \infty$ , we get a contradiction that  $0 \ge c_0$ . Hence,  $\lambda_n$  decreases towards 0 as  $n \to \infty$ . We conclude Proposition 3.6.  $n \to \infty$ . We conclude Proposition [3.6.](#page-15-3)

<span id="page-16-0"></span>3.4. **Proof of Theorem [2.1.](#page-7-0)** Let  $\lambda_n$  be found from Proposition [3.6](#page-15-3) and  $\phi_n(x)$  $Y_{a,\lambda_n}^{-1}M\varpi_n(x)$  on  $(-a, a)$ . Keep in mind our computations in Subsection [3.1,](#page-8-1) we extend  $\phi_n$  to the whole line by requiring  $\phi_n$  satisfies [\(3.4\)](#page-8-4) and [\(3.5\)](#page-9-1) for some constants  $A_{n,1}^{\pm}$  and  $A_{n,2}^{\pm}$  as  $\lambda = \lambda_n$ . Those constants  $A_{n,1}^{\pm}$  and  $A_{n,2}^{\pm}$  are defined by

<span id="page-16-3"></span>
$$
\begin{cases}\n\phi_n(a) = A_{n,1}^+ + A_{n,2}^+, \n\phi'_n(a) = k A_{n,1}^+ + \sqrt{k^2 + \lambda_n \nu_+} A_{n,2}^+.\n\end{cases}
$$
\n(3.37)

and by

<span id="page-16-4"></span>
$$
\begin{cases}\n\phi_n(-a) = A_{n,1}^- + A_{n,2}^-,\n\phi'_n(-a) = kA_{n,1}^- + \sqrt{k^2 + \lambda_n \nu_-} A_{n,2}^-.\n\end{cases}
$$
\n(3.38)

Since  $\phi_n$  is not trivial, there exists  $i \geq 0$  such that  $\phi_n^{(i)}(a) \neq 0$ , which ensures that, at least, one of  $A_{n,1}^{\pm}$  is nonzero. Similarly, at least, one of  $A_{n,2}^{\pm}$  is nonzero.

Solving  $(3.37)$  and  $(3.38)$ , we get that

$$
A_{n,1}^{+} = \frac{\sqrt{k^2 + \lambda_n \nu_+} \phi_n(a) - \phi_n'(a)}{\sqrt{k^2 + \lambda_n \nu_+} - k}, \quad A_{n,2}^{+} = \frac{\phi_n'(a) - k\phi_n(a)}{\sqrt{k^2 + \lambda_n \nu_+} - k}.
$$
 (3.39)

<span id="page-16-5"></span>

and that

$$
A_{n,1}^- = \frac{\sqrt{k^2 + \lambda_n \nu_-} \phi_n(-a) - \phi_n'(-a)}{\sqrt{k^2 + \lambda_n \nu_-} - k}, \quad A_{n,2}^- = \frac{\phi_n'(-a) - k\phi_n(-a)}{\sqrt{k^2 + \lambda_n \nu_-} - k}.\tag{3.40}
$$

Therefore, the function  $\phi_n$  is a regular solution of [\(1.2\)](#page-3-0) as  $\lambda = \lambda_n$  for each  $n \ge 1$ . Proof of Theorem [2.1](#page-7-0) is complete.

#### 4. The strictly increasing profile case

<span id="page-17-0"></span>In this section, let  $(e_1, e_2, e_3, e_4)$  be the canonical basis of  $\mathbb{R}^4$  and we pay attention to an increasing profile  $\rho_0$  satisfying  $(2.8)$  and  $(2.9)$ . We will use the Frobenius matrix norm  $\|\cdot\|_F$  and the Euclidean vector norm  $\|\cdot\|_2$ .

# <span id="page-17-1"></span>4.1. Solutions decaying to 0 at infinity and reduction to a problem on a finite interval. Let

<span id="page-17-7"></span>
$$
L(x,\lambda) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -\frac{\lambda k^2 \rho_0(x)}{\mu} - k^4 & 0 & \frac{\lambda \rho_0(x)}{\mu} + 2k^2 & 0 \end{pmatrix}
$$
(4.1)

and

<span id="page-17-3"></span>
$$
R(\lambda) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{g k^2}{\lambda \mu} & \frac{\lambda}{\mu} & 0 & 0 \end{pmatrix}
$$
(4.2)

We set  $U = (\phi, \phi', \phi'', \phi''')^T$  and then rewrite [\(1.2\)](#page-3-0) as

<span id="page-17-2"></span>
$$
U'(x) = (L(x, \lambda) + \rho'_0(x)R(\lambda))U(x),
$$
\n(4.3)

The eigenvalues of  $L(x, \lambda)$ ,  $\pm k$  and  $\pm \sigma_0(x, \lambda)$ , with  $\sigma_0(x, \lambda) = \sqrt{k^2 + \frac{\lambda}{\mu} \rho_0(x)}$  are different for all  $\lambda > 0$  and for all  $x \in \mathbb{R}$ . Furthermore,

<span id="page-17-4"></span>
$$
\int_{-\infty}^{+\infty} |L'(x,\lambda)| dx < +\infty, \quad \int_{-\infty}^{+\infty} |\rho_0'(x)R(\lambda)| dx < +\infty.
$$
 (4.4)

 $(4.4)$  allow us to use  $[3,$  Theorem 8.1, [C](#page-33-0)hapter 3] (see Appendix C for the statement) to find bounded solutions of [\(4.3\)](#page-17-2) near  $\pm \infty$ . We denote  $\sigma_{\pm}(\lambda) = \lim_{x \to \pm \infty} \sigma_0(x, \lambda)$ . Then, there exist  $\bar{x}$ <sup> $\bar{x}$ </sup>  $\leq \bar{x}$ <sup> $\bar{x}$ </sup> such that we have the existence of bounded solutions  $U_{1,2}^+$  on  $(\bar{x}_+, +\infty)$  and  $U_{3,4}^-$  on  $(-\infty, \bar{x}_-)$  such that in a neighborhood of  $+\infty$ ,

<span id="page-17-5"></span>
$$
e^{kx}U_1^+(x,\lambda) \to (-k^{-3}, k^{-2}, -k^{-1}, 1)^T,
$$
  
\n
$$
\exp\left(\int_{\bar{x}_+}^x \sigma_0(y,\lambda)dy\right)U_2^+(x,\lambda) \to (-\sigma_+^{-3}(\lambda), \sigma_+^{-2}(\lambda), -\sigma_+^{-1}(\lambda), 1)^T,
$$
\n(4.5)

and in a neighborhood of  $-\infty$ ,

<span id="page-17-6"></span>
$$
e^{-kx}U_3^-(x,\lambda) \to (k^{-3}, k^{-2}, k^{-1}, 1)^T,
$$
  
\n
$$
\exp\left(-\int_{\bar{x}_-}^x \sigma_0(y,\lambda)dy\right)U_4^-(x,\lambda) \to (\sigma_-^{-3}(\lambda), \sigma_-^{-2}(\lambda), \sigma_-^{-1}(\lambda), 1)^T.
$$
\n(4.6)

Let us prove that  $U_1^+(x,\lambda)$  and  $U_2^+(x,\lambda)$  are linearly independent. Through Cauchy-Lipschitz theorem, if they are linearly dependent at a particular  $x_0$  then they are linearly dependent for all x, that is there exists  $T(\lambda)$  such that

$$
U_1^+(x,\lambda) = T(\lambda)U_2^+(x,\lambda).
$$

In particular, as the limit of  $kx - \int_{\bar{x}_+}^x \sigma_0(y, \lambda) dy$  when  $x \to +\infty$  is equal to  $-\infty$ , one observes that

$$
e^{kx}U_1^+(x,\lambda)=T(\lambda)\text{exp}\Big(kx-\int_{\bar{x}_+}^x\sigma_0(y,\lambda)dy\Big)\Big(U_2^+(x,\lambda)\text{exp}\Big(\int_{\bar{x}_+}^x\sigma_0(y,\lambda)dy\Big)\Big).
$$

The right hand side of this identity converges to 0 when  $x \to +\infty$ , while the left hand side converges to  $(-k^{-3}, k^{-2}, -k^{-1}, 1)^T$  when  $x \to +\infty$ , contradiction. Similarly,  $U_3^-(x,\lambda)$  and  $U_4^-(x,\lambda)$  are linearly independent.

The aim of the next proposition is to reduce the study of Eq.  $(1.2)$  on the real line to its study on a finite interval as in the previous section. This is really the key of our result for a smooth general profile because it entitles us to use the compact injection of  $H^j((a, b))$  into  $H^{j-1}((a, b)), j \geq 1$ .

<span id="page-18-2"></span>**Proposition 4.1.** *There exist*  $x^0 \leq \bar{x}$  *and*  $x^0 + \bar{x}$  *x such that, for all*  $x^- \leq x^0$  $\lim_{x \to \infty} \frac{1}{x}$  is  $\lim_{x \to \infty} \frac{1}{x}$  and  $\lim_{x \to \infty} \frac{1}{x}$  is  $\lim_{x \to \infty} \frac{1}{x}$  and  $\lim_{x \to \infty} \frac{1}{x}$  is  $\lim_{x \to \infty} \frac{1}{x}$  and  $\lim_{x \to \infty} \frac{1}{x}$  is  $\lim_{x \to \infty} \frac{1}{x}$  and  $\lim_{x \to \infty} \frac{1}{x}$  is  $\lim_{x \to \infty$ *that equation*

$$
\lambda^{2}(k^{2}\rho_{0}\phi - (\rho_{0}\phi')') + \lambda\mu(\phi^{(4)} - 2k^{2}\phi'' + k^{4}\phi) = g k^{2} \rho'_{0}\phi,
$$

*where*  $\phi \in H^4((x_-, x_+))$ , supplemented with the boundary conditions at  $x_-$  are

<span id="page-18-0"></span>
$$
\begin{cases}\nn_{11}^{-}\phi(x_{-}) + n_{12}^{-}\phi'(x_{-}) + \phi''(x_{-}) = 0, \\
n_{21}^{-}\phi(x_{-}) + n_{22}^{-}\phi'(x_{-}) + \phi'''(x_{-}) = 0\n\end{cases}
$$
\n(4.7)

and at  $x_+$ , are

<span id="page-18-1"></span>
$$
\begin{cases}\nn_{11}^{+}\phi(x_{+}) + n_{12}^{+}\phi'(x_{+}) + \phi''(x_{+}) = 0, \\
n_{21}^{+}\phi(x_{+}) + n_{22}^{+}\phi'(x_{+}) + \phi'''(x_{+}) = 0,\n\end{cases} \tag{4.8}
$$

*is equivalent to equation*

$$
\lambda^{2}(k^{2}\rho_{0}\Phi - (\rho_{0}\Phi')') + \lambda\mu(\Phi^{(4)} - 2k^{2}\Phi'' + k^{4}\Phi) = g k^{2} \rho'_{0} \Phi,
$$

*where*  $\Phi \in H^4(\mathbf{R})$ *.* 

*In this case*  $\Phi = \phi$  *on*  $(x_-, x_+)$  *and on*  $(x_+, +\infty)$  (respectively  $(-\infty, x_-)$ ),  $\Phi$  *is the first component of a linear combination of*  $U_{1,2}^+(x,\lambda)$  (see [\(4.5\)](#page-17-5)) (respectively  $U_{3,4}^{-}(x,\lambda),$  see  $(4.6)$ .

*Proof.* Notice that if  $\Phi$  is a bounded solution of

$$
\lambda^{2}(k^{2} \rho_{0} \Phi - (\rho_{0} \Phi')') + \lambda \mu (\Phi^{(4)} - 2k^{2} \Phi'' + k^{4} \Phi) = g k^{2} \rho'_{0} \Phi,
$$

 $\Theta = (\Phi, \Phi', \Phi'', \Phi''')^T$  is a bounded solution of [\(4.3\)](#page-17-2). Any decaying solution  $\Psi$  of  $(4.3)$  on  $(\bar{x}_+ , +\infty)$  belongs to the space spanned by  $U_1^+(x, \lambda)$  and  $U_2^+(x, \lambda)$ , which is of dimension 2 because they are linearly independent. It is equivalent to say that, for any  $x_+ \geq \bar{x}_+$ ,

<span id="page-18-3"></span>
$$
U_1^+(x_+,\lambda) \wedge U_2^+(x_+,\lambda) \wedge \Theta(x_+,\lambda) = 0 \tag{4.9}
$$

and

 $\Theta(x_+,\lambda)$  belongs to the space spanned by  $U_1^+(x_+,\lambda)$  and  $U_2^+(x_+,\lambda)$  .

Let us write  $U_i^+ = (U_{i1}^+, U_{i2}^+, U_{i3}^+, U_{i4}^+)^T$  for  $i = 1, 2$  and  $U_i^- = (U_{i1}^-, U_{i2}^-, U_{i3}^-, U_{i4}^-)^T$ for  $i = 3, 4$ . System  $(4.9)$  is a system of four equations on the components of

 $\Theta(x_+,\lambda)$ , hence there exists a couple of equations which are linearly independent (the system being of rank 2). Let us notice that two of these four equations contain, in  $\Phi''$  and  $\Phi'''$ , respectively the term

$$
(U_{11}^+ U_{22}^+ - U_{12}^+ U_{21}^+)(x_+, \lambda) \Phi''(x_+, \lambda)
$$

and

$$
(U_{11}^+ U_{22}^+ - U_{12}^+ U_{21}^+)(x_+, \lambda) \Phi'''(x_+, \lambda).
$$

As the limit, when  $x_+ \to +\infty$ , of

$$
\exp\Bigl(kx_{+}+\int_{\bar{x}_{+}}^{x_{+}}\sigma_{0}(y,\lambda)dy\Bigr)(U_{11}^{+}U_{22}^{+}-U_{12}^{+}U_{21}^{+})(x_{+},\lambda)
$$

is

$$
-\frac{1}{k^3 \sigma_+^2(\lambda)} + \frac{1}{k^2 \sigma_+^3(\lambda)} = -\frac{\lambda \rho_+}{\mu k^3 \sigma_+^3(\lambda)(k + \sigma_+)} < 0,
$$

by continuity there exists a  $x_+^0 \geq \bar{x}_+$  such that, for all  $x_+ \geq x_+^0$ ,

$$
(U_{11}^+ U_{22}^+ - U_{12}^+ U_{21}^+)(x_+, \lambda) < 0
$$

hence the equations for [\(4.9\)](#page-18-3) on the components  $e_1 \wedge e_2 \wedge e_3$  and  $e_1 \wedge e_2 \wedge e_4$  write as  $N^+ \Theta(x_+, \lambda) = 0$  with  $N^+$  is a  $4 \times 2$  matrix of the form

$$
N^{+} = \begin{pmatrix} n_{11}^{+} & n_{12}^{+} & 1 & 0 \\ n_{21}^{+} & n_{22}^{+} & 0 & 1 \end{pmatrix}.
$$

We are now able to write the couple  $N^+ \Theta(x_+, \lambda) = 0$  as

$$
\begin{cases} n_{11}^+\Phi(x_+, \lambda)+n_{12}^+\Phi'(x_+, \lambda)+\Phi''(x_+, \lambda)=0, \\ n_{21}^+\Phi(x_+, \lambda)+n_{22}^+\Phi'(x_+, \lambda)+\Phi'''(x_+, \lambda)=0, \end{cases}
$$

In a similar way, there exist  $n_{ij}^ (i, j = 1, 2)$  depending on  $x_{-}$  and  $\lambda$  such that

$$
\begin{cases} n_{11}^- \Phi(x_-, \lambda) + n_{12}^- \Phi'(x_-, \lambda) + \Phi''(x_-, \lambda) = 0, \\ n_{21}^- \Phi(x_-, \lambda) + n_{22}^- \Phi'(x_-, \lambda) + \Phi'''(x_-, \lambda) = 0, \end{cases}
$$

Hence  $\Phi$  is solution of the ODE on  $(x_-, x_+)$ :

$$
\lambda^{2}(k^{2}\rho_{0}\Phi - (\rho_{0}\Phi')') + \lambda\mu(\Phi^{(4)} - 2k^{2}\Phi'' + k^{4}\Phi) = g k^{2} \rho'_{0} \Phi,
$$

with boundary conditions  $(4.7)$ ,  $(4.8)$ .

Conversely, assume that  $\phi \in H^4((x_-, x_+))$  is a solution of equation

$$
\lambda^{2}(k^{2}\rho_{0}\phi - (\rho_{0}\phi')') + \lambda\mu(\phi^{(4)} - 2k^{2}\phi'' + k^{4}\phi) = g k^{2}\rho'_{0}\phi,
$$

with boundary conditions  $(4.7)-(4.8)$  $(4.7)-(4.8)$  $(4.7)-(4.8)$ . From the boundary conditions, we deduce that there exist  $C_j^+(j = 1, 2), D_k^-(k = 3, 4)$  such that

$$
U(x_+,\lambda) = C_1^+ U_1^+(x_+,\lambda) + C_2^+ U_2^+(x_+,\lambda)
$$

and

$$
U(x_{-},\lambda) = D_3^- U_3^-(x_{-},\lambda) + D_4^- U_4^-(x_{-},\lambda).
$$

Then, through Cauchy-Lipschitz theorem,

$$
U(x,\lambda)=C_1^+U_1^+(x,\lambda)+C_2^+U_2^+(x,\lambda)\quad\text{for all }x\geqslant x_+
$$

and

$$
U(x,\lambda) = D_3^- U_3^-(x,\lambda) + D_4^- U_4^-(x,\lambda) \quad \text{for all } x \leq x_-.
$$

As these are decaying solutions at  $\pm \infty$  respectively, and as there is no jump at  $x_+$  (or *x*<sub>-</sub>) for  $\phi$ ,  $\phi'$ ,  $\phi''$ ,  $\phi'''$  (which have a meaning as  $\phi$  is assumed to be in  $H^4((x_-, x_+)),$ the function

$$
\Phi(x) = \begin{cases} C_1^+ U_{11}^+(x, \lambda) + C_2^+ U_{21}^+(x, \lambda), & \text{as } x \ge x_+ \\ \phi(x), & \text{as } x_- < x < x_+ \\ D_3^- U_{31}^-(x, \lambda) + D_4^- U_{41}^-(x, \lambda), & \text{as } x \le x_- \end{cases}
$$

belongs to  $H^4(\mathbf{R})$  and solves equation

$$
\lambda^2 (k^2 \rho_0 \Phi - (\rho_0 \Phi')') + \lambda \mu (\Phi^{(4)} - 2k^2 \Phi'' + k^4 \Phi) = g k^2 \rho'_0 \Phi,
$$
  
on **R**.

Remark that in this proof we have no additional information on the values of  $x^0$ and of  $x_+^0$ . This will be the aim of Section [4.4.](#page-24-0)

<span id="page-20-0"></span>4.2. A bilinear form and a self-adjoint invertible operator. The aim of Section [4.2](#page-20-0) is to study the boundary terms that come from the bilinear form  $(4.15)$ below. Let

<span id="page-20-3"></span>
$$
BV_{x_+, \lambda}(\vartheta, \varrho) := -\lambda (\rho_0 \vartheta' \varrho)(x_+) - \mu (n_{21}^+(x_+, \lambda) \vartheta(x_+) + n_{22}^+(x_+, \lambda) \vartheta'(x_+)) \varrho(x_+) + \mu (n_{11}^+(x_+, \lambda) \vartheta(x_+) + n_{12}^+(x_+, \lambda) \vartheta'(x_+)) \varrho'(x_+) - 2k^2 \mu (\vartheta' \varrho)(x_+) (4.10)
$$

and by

<span id="page-20-4"></span>
$$
BV_{x_-, \lambda}(\vartheta, \varrho) := \lambda(\rho_0 \vartheta' \varrho)(x_-) + \mu(n_{21}^-(x_-, \lambda)\vartheta(x_-) + n_{22}^-(x_-, \lambda)\vartheta'(x_+))\varrho(x_-) - \mu(n_{11}^-(x_-, \lambda)\vartheta(x_-) + n_{12}^-(x_-, \lambda)\vartheta'(x_-))\varrho'(x_-) + 2k^2\mu(\vartheta'\varrho)(x_-).
$$
\n(4.11)

Lemma 4.1. *Necessary and sufficient conditions to get*

<span id="page-20-5"></span>
$$
BV_{x_+, \lambda}(\vartheta, \vartheta) \geq 0 \text{ and } BV_{x_-, \lambda}(\vartheta, \vartheta) \geq 0 \tag{4.12}
$$

*for all*  $\vartheta \in H^2([x_-, x_+])$  *are* 

<span id="page-20-2"></span>
$$
n_{12}^+(x_+,\lambda) \ge 0, \quad n_{21}^+(x_+,\lambda) \le 0,
$$
  
\n
$$
(n_{11}^+(x_+,\lambda) - n_{22}^+(x_+,\lambda) - k^2 - \sigma_0^2(x_+,\lambda))^2 + 4n_{12}^+n_{21}^+(x_+,\lambda) \le 0,
$$
\n
$$
(4.13)
$$

<span id="page-20-1"></span>*and*

$$
n_{12}^-(x_-,\lambda) \ge 0, \quad n_{21}^-(x_-,\lambda) \le 0,
$$
  
\n
$$
(n_{11}^-(x_-,\lambda) - n_{22}^-(x_-,\lambda) - k^2 - \sigma_0^2(x_-,\lambda))^2 + 4n_{12}^-(x_-,\lambda) \le 0.
$$
\n(4.14)

*Proof of Lemma [4.1.](#page-20-1)* We treat only the case  $BV_{x_+,\lambda}(\vartheta,\vartheta) \geq 0$ . Since  $\mu \sigma_0^2(x,\lambda) =$  $\mu k^2 + \lambda \rho_0(x)$ , we rewrite

$$
\frac{1}{\mu}BV_{x_+, \lambda}(\vartheta, \vartheta) = -n_{21}^+(x_+, \lambda)|\vartheta(x_+)|^2 + n_{12}^+|\vartheta'(x_+)|^2 \n+ (n_{11}^+(x_+, \lambda) - n_{22}^+(x_+, \lambda) - k^2 - \sigma_0^2(x_+, \lambda))\vartheta(x_+)\vartheta'(x_+).
$$

We observe that it is a quadratic polynomial in  $\vartheta(x_+), \vartheta'(x_+)$ . The first case is the case where  $n_{12}^+ = n_{21}^+ = 0$ . The inequality [\(4.13\)](#page-20-2) imply that  $BV_{x_+,\lambda}(\vartheta,\vartheta) = 0$  for all  $\vartheta$ . The second case is the case where at least one of these two real numbers is not zero. For example, if  $n_{12}^+ \neq 0$ , the inequality means that the polynomial

$$
n_{12}^+t^2 + (n_{11}^+ - n_{22}^+ - k^2 - \sigma_0^2)t - n_{21}^+
$$

is always of the sign of  $n_{12}^+$  hence positive, hence  $BV_{x_+,\lambda}(\vartheta,\vartheta) \geq 0$  for all  $\vartheta \in$  $H^2((x_-, x_+))$ . Lemma [4.1](#page-20-1) is proven.

<span id="page-21-1"></span>**Proposition 4.2.** *There exists*  $x^1 \leq x^0, x^1 \geq x^0$  *such that, for any*  $x^- \leq x^1$ *and*  $x_+ \geq x_+^1$  *chosen arbitrarily* 

<span id="page-21-0"></span>
$$
\mathcal{B}_{x_-,x_+,\lambda}(\vartheta,\varrho) := BV_{x_-, \lambda}(\vartheta,\varrho) + BV_{x_+, \lambda}(\vartheta,\varrho) + \lambda \int_{x_-}^{x_+} \rho_0(k^2 \vartheta \varrho + \vartheta' \varrho') dx + \mu \int_{x_-}^{x_+} (\vartheta'' \varrho'' + 2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx,
$$
\n(4.15)

*is a bilinear form on*  $H^2((x_-, x_+))$ , that is continuous and coercive.

*Proof.* Recall  $n_{ij}^{\pm}$  (*i*, *j* = 1, 2) are given in Proposition [4.1](#page-18-2) and that  $BV_{x-1}$ ,  $BV_{x+1}$ , are given in  $(4.10)$ ,  $(4.11)$ . One observes that one needs to prove that

<span id="page-21-2"></span>
$$
|\mathcal{B}_{x_-,x_+,\lambda}(\vartheta,\varrho)| \leq C^{\star} \|\vartheta\|_{H^2((x_-,x_+))} \|\varrho\|_{H^2((x_-,x_+))}
$$
(4.16)

and that

<span id="page-21-3"></span>
$$
\mathcal{B}_{x_-,x_+,\lambda}(\vartheta,\vartheta) \geqslant C^{\star} \|\vartheta\|_{H^2((x_-,x_+))}^2.
$$
\n
$$
(4.17)
$$

For positive  $\lambda, \mu$  and k, we have

$$
\lambda \int_{x_-}^{x_+} \rho_0(k^2 \vartheta \varrho + \vartheta' \varrho') dx_3 + \mu \int_{x_-}^{x_+} (\vartheta'' \varrho'' + 2k^2 \vartheta' \varrho' + k^4 \vartheta \varrho) dx
$$
  

$$
\leq C^* \|\vartheta\|_{H^2((x_-,x_+))} \|\varrho\|_{H^2((x_-,x_+))}
$$

and that

$$
\lambda \int_{x_{-}}^{x_{+}} \rho_{0}(k^{2}|\vartheta|^{2} + |\vartheta'|^{2}) dx + \mu \int_{x_{-}}^{x_{+}} (|\vartheta''|^{2} + 2k^{2}|\vartheta'|^{2} + k^{4}|\vartheta|^{2}) dx \geq C^{\star} \|\vartheta\|_{H^{2}((x_{-},x_{+}))}^{2}.
$$

Note that the condition  $\mu > 0$  is necessary to obtain the coercivity on  $H^2((x_-, x_+))$ . Note also that the case  $\mu = 0$  amounts to the inviscid Rayleigh-Taylor instability, for which similar results are known (and the corresponding problem needs only to be defined in  $H^1$ ). In addition

$$
|BV_{x_+, \lambda}(\vartheta, \varrho)| \le C^{\star}(|\vartheta(x_+)| + |\varrho(x_+)|)(|\vartheta'(x_+)| + |\varrho'(x_+)|),
$$
  

$$
|BV_{x_-, \lambda}(\vartheta, \varrho)| \le C^{\star}(|\vartheta(x_-)| + |\varrho(x_-)|)(|\vartheta'(x_-)| + |\varrho'(x_-)|)
$$

and the Sobolev embedding yields [\(4.16\)](#page-21-2). The continuity of  $\mathcal{B}_{x_-,x_+,\lambda}$  on  $H^2((x_-,x_+))$ follows.

To show  $(4.17)$ , it suffices to prove that  $(4.12)$  holds. In view of Lemma [4.1,](#page-20-1) we verify [\(4.13\)](#page-20-2). Since  $N^+U_1^+(x_+,\lambda) = N^+U_2^+(x_+,\lambda) = 0$ , we have that  $n_{ij}^+$  $(i, j = 1, 2)$  depend on  $x_+$  and  $\lambda$  and satisfy

<span id="page-21-4"></span>
$$
\begin{cases}\nn_{11}^+ U_{11}^+(x_+, \lambda) + n_{12}^+ U_{12}^+(x_+, \lambda) + U_{13}^+(x_+, \lambda) = 0, \\
n_{11}^+ U_{21}^+(x_+, \lambda) + n_{12}^+ U_{22}^+(x_+, \lambda) + U_{23}^+(x_+, \lambda) = 0,\n\end{cases} \tag{4.18}
$$

and

<span id="page-21-5"></span>
$$
\begin{cases}\nn_{21}^+ U_{11}^+(x_+, \lambda) + n_{22}^+ U_{12}^+(x_+, \lambda) + U_{14}^+(x_+, \lambda) = 0, \\
n_{21}^+ U_{21}^+(x_+, \lambda) + n_{22}^+ U_{22}^+(x_+, \lambda) + U_{24}^+(x_+, \lambda) = 0.\n\end{cases} \tag{4.19}
$$

Let  $n^+_{ij}(\lambda)$  be the limit of  $n^+_{ij}(x_+,\lambda)$  as  $x \to +\infty$ . When  $x_+ \to +\infty$ , [\(4.18\)](#page-21-4)-[\(4.19\)](#page-21-5) converge to

$$
\begin{cases}\n-n_{11}^{+}(\lambda)k^{-3} + n_{12}^{+}(\lambda)k^{-2} - k^{-1} = 0, \\
-n_{11}^{+}(\lambda)\sigma_{+}^{-3}(\lambda) + n_{12}^{+}(\lambda)\sigma_{+}^{-2}(\lambda) - \sigma_{+}^{-1}(\lambda) = 0,\n\end{cases}
$$
\n(4.20)

and

$$
\begin{cases}\n-\mathsf{n}_{21}^+(\lambda)k^{-3} + \mathsf{n}_{22}^+(\lambda)k^{-2} + 1 = 0, \\
-\mathsf{n}_{21}^+(\lambda)\sigma_+^{-3}(\lambda) + \mathsf{n}_{22}^+(\lambda)\sigma_+^{-2}(\lambda) + 1 = 0,\n\end{cases}
$$
\n(4.21)

hence

$$
\mathsf{n}_{11}^+(\lambda) = k\sigma_+(\lambda), \quad \mathsf{n}_{12}^+(\lambda) = k + \sigma_+(\lambda)
$$

and

$$
\mathsf{n}_{21}^+(\lambda) = -k\sigma_+(\lambda)(k + \sigma_+(\lambda)), \quad \mathsf{n}_{22}^+(\lambda) = -(k^2 + k\sigma_+(\lambda) + \sigma_+^2(\lambda)).
$$

One thus has

$$
\begin{aligned} & \left( \mathsf{n}_{11}^+(\lambda) - \mathsf{n}_{22}^+(\lambda) - k^2 - \sigma_+^2(\lambda) \right)^2 + 4\mathsf{n}_{12}^+\mathsf{n}_{21}^+(\lambda) \\ &= \left( (k + \sigma_+(\lambda))^2 - k^2 - \sigma_+^2(\lambda) \right)^2 - 4k\sigma_+(\lambda)(k + \sigma_+(\lambda))^2 \\ &= -4k\sigma_+(\lambda)(k^2 + k\sigma_+(\lambda) + \sigma_+^2(\lambda)) < 0. \end{aligned}
$$

Hence, by continuity, there exists  $x_+^1 \ge x_+^0$  such that, for all  $x_+ \ge x_+^1$ ,

$$
(n_{11}^+(x_+,\lambda)-n_{22}^+(x_+,\lambda)-k^2-\sigma_0^2(x_+,\lambda))^2+4n_{12}^+n_{21}^+(x_+,\lambda)<0.
$$

The proof of the existence of  $x^1 \leq x^0$  such that, for all  $x^- \leq x^1$  such that

$$
(n_{11}^-(x_-,\lambda)-n_{22}^-(x_-,\lambda)-k^2-\sigma_0^2(x_-,\lambda))^2+4n_{12}^-\overline{n}_{21}^-(x_-,\lambda)<0
$$

follows the same pattern. Hence, by application of Lemma  $4.1$ ,  $(4.17)$  follow, which ends the proof of Proposition [4.2.](#page-21-1)  $\Box$ 

Mimicking the arguments in Propositions [3.2,](#page-10-5) [3.3,](#page-11-0) [3.4](#page-12-3) we obtain the following proposition.

<span id="page-22-0"></span>**Proposition 4.3.** Let  $(H^2((x_-, x_+)))'$  be the dual space of  $H^2((x_-, x_+))$  associated *with the norm*  $\sqrt{\mathcal{B}_{x_-,x_+,\lambda}(\cdot,\cdot)}$ , there exists a unique operator

$$
Y_{x_-,x_+,\lambda} \in \mathcal{L}(H^2((x_-,x_+)), (H^2((x_-,x_+))'),
$$

*that is also bijective, such that*

$$
\mathcal{B}_{x_-,x_+,\lambda}(\vartheta,\varrho)=\langle Y_{x_-,x_+,\lambda}\vartheta,\varrho\rangle\tag{4.22}
$$

*for all*  $\vartheta, \rho \in H^2((x_-, x_+))$ *. Furthermore, we have* 

(1) *For all*  $\vartheta \in H^2((x_-, x_+)),$ 

$$
Y_{x_-,x_+,\lambda}\vartheta = \lambda(k^2\rho_0\vartheta - (\rho_0\vartheta')') + \mu(\vartheta^{(4)} - 2k^2\vartheta'' + k^4\vartheta)
$$

in 
$$
\mathcal{D}'((x_-, x_+))
$$
.

(2) Let  $f \in L^2((x_-, x_+))$  be given, there exists a unique  $\vartheta \in H^4((x_-, x_+))$ *satisfying the boundary conditions* [\(4.7\)](#page-18-0)*–*[\(4.8\)](#page-18-1)*.*

$$
Y_{x_-,x_+,\lambda}\vartheta = f \ \ in \ (H^2((x_-,x_+)))',\tag{4.23}
$$

(3) The operator  $Y_{x-,x_+,\lambda}^{-1}: L^2((x_-,x_+)) \to L^2((x_-,x_+))$  is compact and self*adjoint.*

It is then straightforwad to obtain the following spectral result. The discrete spectrum of  $\mathcal{M}Y_{x_-,x_+,\lambda}^{-1}\mathcal{M}$  is a sequence of eigenvalues  $(\gamma_n(\lambda))_{n\geq 1}$ . The function  $\gamma_n(\lambda)$  is a continuous function of  $\lambda$  for all *n* from the arguments of Lemma [3.2.](#page-14-0) The problem of finding a characteristic value amounts to solving the equality [\(1.5\)](#page-5-0) as before. However, no control is possible on  $\gamma_n(\lambda)$  when  $\lambda$  goes to 0. In addition, no control of  $x_{\pm}^0$  and  $x_{\pm}^1$  (because it may depend on  $\lambda$ ) is available to have a possibility of having estimates of  $\gamma_n(\lambda)$  as well. Having an explicit (even if not optimal) criterion on  $x_{\pm}^1$  such that the inequalities of coercivity [\(4.13\)](#page-20-2)-[\(4.14\)](#page-20-1) are true is the aim of the refined estimates of the solutions  $U_{1,2}^+$  and  $U_{3,4}^-$  (deducing in

<span id="page-23-5"></span>Propositions [4.7,](#page-28-0) [4.8](#page-29-0) below) which follow. That will be postponed to Section [4.4](#page-24-0) below.

<span id="page-23-0"></span>4.3. The finding of characteristic value  $\lambda_n$  and Proof of Theorem [2.2.](#page-7-1) Let  $\epsilon_{\star} > 0$  be given, we look for  $\lambda_n \in (\epsilon_{\star}, \sqrt{\frac{g}{L_0}})$  satisfying [\(1.5\)](#page-5-0). However, unlike the previous case with  $\rho'_0 \geq 0$  being compactly supported, we do not have here the decrease of  $\gamma_n$  on  $\lambda$  to obtain the uniqueness of  $\lambda_n$ .

<span id="page-23-1"></span>**Proposition 4.4.** *For*  $0 < \epsilon_{\star} \ll 1$ *, there exists*  $N(\epsilon_{\star}) \in \mathbb{N}$  *such that there is at least one positive*  $\lambda_n \in (\epsilon_{\star}, \sqrt{\frac{g}{L_0}})$  *satisfying* [\(1.5\)](#page-5-0) *for each*  $1 \leq n \leq N(\epsilon_{\star})$ *.* 

*Proof.* We still have

<span id="page-23-2"></span>
$$
\lim_{\lambda \to \sqrt{\frac{g}{L_0}}} \frac{\lambda}{\gamma_n(\lambda)} > gk^2.
$$
\n(4.24)

and  $b_n(\epsilon_{\star}) := \inf_{\lambda \geq \epsilon_{\star}} \gamma_n(\lambda) > 0$ . Notice that  $\{b_n(\epsilon_{\star})\}_{n \geq 1}$  is a sequence decreasing to 0 as  $n \to \infty$ . Set

$$
N(\epsilon_\star):=\sup\Big\{n\big|b_n(\epsilon_\star)>\frac{\epsilon_\star}{gk^2}\Big\}\in[1,+\infty).
$$

For  $1 \leq n \leq N(\epsilon_{\star}),$ 

<span id="page-23-3"></span>
$$
\lim_{\lambda \to \epsilon_{\star}} \frac{\lambda}{\gamma_n(\lambda)} \le \lim_{\lambda \to \epsilon_{\star}} \frac{\lambda}{b_n(\epsilon_{\star})} = \frac{\epsilon_{\star}}{b_n(\epsilon_{\star})} < g k^2. \tag{4.25}
$$

It then follows from [\(4.24\)](#page-23-2) and [\(4.25\)](#page-23-3) that we have at least one desired  $\lambda_n \in$  $(\epsilon_{\star}, \sqrt{\frac{g}{L_0}})$  for  $1 \leq n \leq N(\epsilon_{\star}).$ 

Now, we are able to prove Theorem [2.2.](#page-7-1)

*Proof.* Let  $\varpi_n$  be an eigenfunction associated with  $\gamma_n$  of  $\mathcal{M}Y_{x_+,x_-, \lambda}^{-1}\mathcal{M}$ . That means

$$
\mathcal{M}Y_{x_-,x_+,\lambda_n}^{-1}\mathcal{M}\varpi_n=\gamma_n(\lambda_n,k)\varpi_n=\frac{\lambda_n}{gk^2}\varpi_n.
$$

Hence,  $\phi_n = Y_{x_{-},x_{+},\lambda_n}^{-1} \mathcal{M} \varpi_n \in H^4((x_{-},x_{+}))$  satisfies

$$
\lambda_n Y_{x_-,x_+,\lambda_n} \phi_n = g k^2 \rho'_0 \phi_n
$$

on  $(x_-, x_+)$ . In order to conclude that  $\phi_n$  is a solution of [\(1.2\)](#page-3-0), we then extend  $\phi_n$ on  $\bf R$  by continuity.

Let us take  $\lambda = \lambda_n$  in the formulas of  $U_{1,2}^+$  from [\(4.5\)](#page-17-5) and in the formulas of  $U_{3,4}^-$ from [\(4.6\)](#page-17-6). Hence,  $\phi_n$  is of the form

 $\phi_n(x) = B_{n,1}^+ U_{11}^+(x, \lambda_n) + B_{n,2}^+ U_{21}^+(x, \lambda_n)$ 

as  $x \geqslant x_+$  and

$$
\phi_n(x) = B_{n,3}^- U_{31}^-(x,\lambda_n) + B_{n,4}^- U_{41}^-(x,\lambda_n)
$$

as  $x \leq x_{-}$  for some real constants  $B_{n,1}^{+}, B_{n,2}^{+}, B_{n,3}^{-}$  and  $B_{n,4}^{-}$ . The constants  $B_{n,1}^+, B_{n,2}^+$  are defined by

<span id="page-23-4"></span>
$$
\begin{cases}\n\phi_n(x_+) = B_{n,1}^+ U_{11}^+(x_+, \lambda_n) + B_{n,2}^+ U_{21}^+(x_+, \lambda_n) \\
\phi'_n(x_+) = B_{n,1}^+ U_{12}^+(x_+, \lambda_n) + B_{n,2}^+ U_{22}^+(x_+, \lambda_n).\n\end{cases} \tag{4.26}
$$

Similarly, we have the system for  $B_{n,3}^-$  and  $B_{n,4}^-$  is

<span id="page-24-2"></span>
$$
\begin{cases}\n\phi_n(x_-) = B_{n,3}^- U_{31}^-(x_-, \lambda_n) + B_{n,4}^- U_{41}^-(x_-, \lambda_n), \\
\phi'_n(x_-) = B_{n,3}^- U_{32}^-(x_-, \lambda_n) + B_{n,4}^- U_{42}^-(x_-, \lambda_n).\n\end{cases} \tag{4.27}
$$

Since  $\phi_n$  is not trivial, there exists  $i \geq 0$  such that  $\phi_n^{(i)}(x_+)$  is nonzero. It implies that, at least, one of  $B_{n,1}^+$  and  $B_{n,2}^+$  is nonzero.

Solving  $(4.26)$  and  $(4.27)$ , we obtain that

$$
B_{n,1}^{+} = \frac{U_{22}^{+}(x_{+}, \lambda_{n})\phi_{n}(x_{+}) - U_{21}^{+}(x_{+}, \lambda_{n})\phi_{n}'(x_{+})}{(U_{11}^{+}U_{22}^{+} - U_{12}^{+}U_{21}^{+})(x_{+}, \lambda_{n})},
$$

that

$$
B_{n,2}^{+} = \frac{-U_{12}^{+}(x_{+}, \lambda_{n})\phi_{n}(x_{+}) + U_{11}^{+}(x_{+}, \lambda_{n})\phi_{n}'(x_{+})}{(U_{11}^{+}U_{22}^{+} - U_{12}^{+}U_{21}^{+})(x_{+}, \lambda_{n})},
$$

that

$$
B_{n,3}^{-} = \frac{U_{42}^{-}(x_{-},\lambda_{n})\phi_{n}(x_{-}) - U_{41}^{-}(x_{-},\lambda_{n})\phi_{n}'(x_{-})}{(U_{31}^{-}U_{42}^{-} - U_{41}^{-}U_{32}^{-})(x_{-},\lambda_{n})},
$$

and that

$$
B_{n,4}^- = \frac{-U_{32}^-(x_-,\lambda_n)\phi_n(x_-) + U_{31}^-(x_-,\lambda_n)\phi_n'(x_-)}{(U_{31}^+U_{42}^- - U_{41}^-U_{32}^-)(x_-,\lambda_n)}.
$$

Therefore, we get that  $\phi_n$  is a regular solution of [\(1.2\)](#page-3-0) as  $\lambda = \lambda_n$  for each  $1 \le n \le N(\epsilon)$ . This ends the proof of Theorem 2.2.  $N(\epsilon_{\star})$ . This ends the proof of Theorem [2.2.](#page-7-1)

Remark 4.1. *In the forthcoming work* [\[17\]](#page-36-15)*, we will use the above operator approach and a more precise hypothesis on*  $\rho_0 - \rho_{\pm}$  *to ensure that the classical Rayleigh-Taylor instability has an infinite number of discrete spectrum. That will complete the study in* [\[12\]](#page-36-8), where the first author only considered the case  $\frac{\rho'_0}{\rho_0}$  has a nondegenerate *minimum.*

- **Remark 4.2.** (1) *For*  $|x_+|$  *large enough, the investigation of regular solutions to Eq.* [\(1.2\)](#page-3-0) *on the real line is equivalent to that one on*  $(x_-, x_+)$  *with boundary conditions* [\(4.7\)](#page-18-0) *and* [\(4.8\)](#page-18-1) *at*  $x_{\pm}$ *. More computations are required in the second case due to the lack of compact assumption of*  $\rho'_0$ .
	- (2) For  $|x_+|$  large enough, the problem [\(1.2\)](#page-3-0) on  $(x_-, x_+)$  with boundary condi*tions* [\(4.7\)](#page-18-0) *and* [\(4.8\)](#page-18-1) *is equivalent to a weaker version of* [\(1.2\)](#page-3-0) *that can be solved by applying Riesz representation theorem on the bilinear continuous form*  $\mathcal{B}_{x-x+1}$  *and then improving the regularity.*
	- (3) *Generally speaking, a problem on the real line with decaying solutions at infinity and transversality hypotheses is equivalent to a problem with the compact setting when we have enough decays on the solutions at*  $\pm \infty$ *.*

<span id="page-24-0"></span>4.4. Explicit construction and refined estimates of the decaying solutions at infinity. In the regime  $\lambda \geq \epsilon_{\star} > 0$ , we notice again that  $R(\lambda)$  is uniformly bounded. So that, as  $\lambda \geq \epsilon_{\star} > 0$ , we further derive a control of the inner region  $(x_{-},x_{+})$  independent of  $\lambda$  in the following proposition, extending the result of Proposition [4.2.](#page-21-1)

**Proposition 4.5.** Let  $\epsilon_{\star} > 0$  given and let  $\lambda \geq \epsilon_{\star}$ . Let  $z_{+,\epsilon_{\star}}$  (respectively,  $z_{-,\epsilon_{\star}}$ ) *be the sum of two upper bounds, that are functions decreasing towards 0 at*  $+\infty$ *(respectively, at*  $-\infty$ *), on the r.h.s. of* [\(4.48\)](#page-28-1) *and* [\(4.49\)](#page-28-2) *(respectively,* [\(4.52\)](#page-29-1) *and*  $(4.53)$ *). There exist positive constants*  $\Gamma_+(\epsilon_{\star})$  *such that, for all*  $x_+, x_-$  *such that* 

<span id="page-24-1"></span>
$$
z_{+,\epsilon_\star}(x_+) \leq \Gamma_+(\epsilon_\star) \quad \text{and} \quad z_{-,\epsilon_\star}(x_-) \leq \Gamma_-(\epsilon_\star), \tag{4.28}
$$

*we have*  $\mathcal{B}_{x_-,x_+,\lambda}$  *is coercive.* 

The proof of Proposition [4.5](#page-24-1) relies on the refined estimates of the bounded solutions of  $(4.3)$  near  $\infty$ , presented in Propositions [4.7,](#page-28-0) [4.8.](#page-29-0) Before going to the proof of Propositions [4.7,](#page-28-0) [4.8,](#page-29-0) thus Proposition [4.5,](#page-24-1) we present some materials. Notice from [\(4.1\)](#page-17-7) that one has  $L(x, \lambda) = P(x, \lambda)D(x, \lambda)P(x, \lambda)^{-1}$ , where

$$
D(x,\lambda) = \text{diag}(-k, -\sigma_0(x,\lambda), k, \sigma_0(x,\lambda)), \qquad (4.29)
$$
  

$$
P(x,\lambda) = \begin{pmatrix} -k^{-3} & -\sigma_0^{-3}(x,\lambda) & k^{-3} & \sigma_0^{-3}(x,\lambda) \\ k^{-2} & \sigma_0^{-2}(x,\lambda) & k^{-2} & \sigma_0^{-2}(x,\lambda) \\ -k^{-1} & -\sigma_0^{-1}(x,\lambda) & k^{-1} & \sigma_0^{-1}(x,\lambda) \\ 1 & 1 & 1 & 1 \end{pmatrix}
$$
(4.30)

and

$$
P(x,\lambda)^{-1} = \frac{\mu}{2\lambda \rho_0(x)} \begin{pmatrix} -k^3 \sigma_0^2(x,\lambda) & k^2 \sigma_0^2(x,\lambda) & k^3 & -k^2\\ k^2 \sigma_0^3(x,\lambda) & -k^2 \sigma_0^2(x,\lambda) & -\sigma_0^3(x,\lambda) & \sigma_0^2(x,\lambda)\\ k^3 \sigma_0^2(x,\lambda) & k^2 \sigma_0^2(x,\lambda) & -k^3 & -k^2\\ -k^2 \sigma_0^3(x,\lambda) & -k^2 \sigma_0^2(x,\lambda) & \sigma_0^3(x,\lambda) & \sigma_0^2(x,\lambda) \end{pmatrix}
$$
(4.31)

The columns of matrix *P* are denoted by  $P_1$ ,  $P_2$ ,  $P_3$ ,  $P_4$  and  $P_2$ ,  $P_4$  depend on  $(x, \lambda)$ . Note that for every positive  $k, \lambda$  and  $\mu$ ,  $P$  and  $P^{-1}$  are bounded uniformly in **R** and  $P^{-1}$  becomes singular when  $\lambda \to 0$  and x is fixed.

Then, we set 
$$
U(x) = P(x, \lambda)V(x)
$$
, (4.3) becomes  
\n
$$
V'(x) = (D(x, \lambda) + \rho'_0(x)M(x, \lambda))V(x),
$$
\n(4.32)

where

<span id="page-25-3"></span>
$$
M(x,\lambda) = P(x,\lambda)^{-1}R(\lambda)P(x,\lambda) - \frac{d\sigma_0(x)}{d\rho_0(x)}P(x,\lambda)^{-1}\frac{dP(x,\lambda)}{d\sigma_0(x)}.
$$

<span id="page-25-2"></span>Lemma 4.2. *Let*

$$
\delta(\epsilon_{\star}) := \sqrt{k^2 + \frac{\epsilon_{\star}\rho_-}{\mu}} \quad \text{and} \quad \delta_s := \sqrt{k^2 + \sqrt{\frac{g}{L_0}}\frac{\rho_+}{\mu}}.
$$

*For any*  $\lambda \in [\epsilon_{\star}, \sqrt{\frac{g}{L_0}}]$ , there hold

<span id="page-25-0"></span>
$$
\sup_{x \in \mathbf{R}} \|P(x,\lambda)\|_{F} \le \Gamma_{p} := \max\left(1, \frac{1}{k}, \frac{1}{k^{2}}, \frac{1}{k^{3}}\right) \tag{4.33}
$$

*and*

<span id="page-25-1"></span>
$$
\sup_{x \in \mathbf{R}} \|M(x,\lambda)\|_{F} \leq \Gamma_{m}(\epsilon_{\star}) := \frac{1}{\rho_{-}\epsilon_{\star}^{2}} \max\left(g\left(k + \frac{1}{L_{0}}\right), g\left(\frac{k^{2}}{\delta(\epsilon_{\star})} + \frac{1}{L_{0}}\right)\right) + \frac{1}{4\delta(\epsilon_{\star})}\sqrt{\frac{g}{L_{0}}} \max\left(\frac{2k^{2}}{\delta^{2}(\epsilon_{\star})}(k + \delta_{s}), \frac{5k^{2}}{\delta(\epsilon_{\star})} + \delta_{s}, \frac{k^{2}}{\delta(\epsilon_{\star})} + \delta_{s}\right).
$$
\n(4.34)

*Proof.* Due to  $\epsilon_{\star} \leq \lambda \leq \sqrt{\frac{g}{L_0}}$ , we get that  $\delta(\epsilon_{\star}) < \sigma_0(x,\lambda) \leq \delta_s$  for all  $x \in \mathbf{R}$ , it yields [\(4.33\)](#page-25-0). We move to demonstrate [\(4.34\)](#page-25-1). Let

$$
a_{\pm}(\lambda) = g k \pm \lambda^2
$$
,  $b_{\pm}(x,\lambda) = \frac{g k^2}{\sigma_0(x,\lambda)} \pm \lambda^2$ 

and

$$
c_{\pm}(x,\lambda) = k \pm \sigma_0(x,\lambda), \quad d(x,\lambda) = \frac{5k^2}{\sigma_0(x,\lambda)} - \sigma_0(x,\lambda).
$$

Direct computations show that

$$
\frac{d\sigma_0(x,\lambda)}{d\rho_0(x)} = \frac{\lambda}{2\mu} \frac{\rho'_0(x)}{\sigma_0(x,\lambda)},
$$

that

$$
P(x,\lambda)^{-1}R(\lambda)P(x,\lambda) = \frac{1}{2\lambda^2 \rho_0(x)} \begin{pmatrix} a_- & \frac{k^2b_-}{\sigma_0^2} & -a_+ & -\frac{k^2b_+}{\sigma_0^2} \\ -\frac{a_-\sigma_0^2}{k^2} & -b_- & \frac{a_+\sigma_0^2}{k^2} & -b_+ \\ a_- & \frac{k^2b_-}{\sigma_0^2} & -a_+ & -\frac{k^2b_+}{\sigma_0^2} \\ -\frac{a_-\sigma_0^2}{k^2} & -b_- & \frac{a_+\sigma_0^2}{k^2} & -b_+ \end{pmatrix} (x,\lambda)
$$

and that

$$
P(x,\lambda)^{-1} \frac{dP(x,\lambda)}{d\sigma_0(x,\lambda)} = \frac{\mu}{2\lambda \rho_0(x)} \begin{pmatrix} 0 & -\frac{2k^2c_+}{\sigma_0^2} & 0 & \frac{2k^2c_-}{\sigma_0^2} \\ 0 & d & 0 & -\frac{k^2}{\sigma_0} + \sigma_0 \\ 0 & \frac{2k^2c_-}{\sigma_0^2} & 0 & -\frac{2k^2c_+}{\sigma_0^2} \\ 0 & -\frac{k^2}{\sigma_0} + \sigma_0 & 0 & d \end{pmatrix} (x,\lambda).
$$

¨

For all  $x \in \mathbf{R}$ , it is clear that

$$
|a_{\pm}(\lambda)| \le g\Big(k + \frac{1}{L_0}\Big), \quad |b_{\pm}(x,\lambda)| \le g\Big(\frac{k^2}{\delta(\epsilon_\star)} + \frac{1}{L_0}\Big)
$$

and that

$$
|c_{\pm}(x,\lambda)| \leq k + \delta_s, \quad |d(x,\lambda)| \leq \frac{5k^2}{\delta(\epsilon_\star)} + \delta_s.
$$

Therefore,  $(4.34)$  follows. Proof of Lemma [4.2](#page-25-2) is complete.

Let  $\tilde{x}_+$  be chosen such that

$$
\int_{\tilde{x}_+}^{+\infty} \rho_0'(\tau) \|M(\tau,\lambda)\|_F d\tau \le \Gamma_m(\epsilon_\star) (\rho_+ - \rho_0(\tilde{x}_+)) < \frac{1}{2} \tag{4.35}
$$

and  $\tilde{x}$ <sub> $-$ </sub> be chosen such that

$$
\int_{-\infty}^{\tilde{x}_{-}} \rho_{0}'(\tau) \| M(\tau, \lambda) \|_{F} d\tau \leqslant \Gamma_{m}(\epsilon_{\star}) (\rho_{0}(\tilde{x}_{-}) - \rho_{-}) < \frac{1}{2}.
$$
 (4.36)

Let  $\alpha_+(x)$  and  $\beta_+(x,\lambda)$  be defined

$$
\alpha_{\pm}(x) = \pm k(x - \tilde{x}_{\pm}), \quad \beta_{\pm}(x,\lambda) = \pm \int_{\tilde{x}_{\pm}}^{x} \sigma_0(y,\lambda) dy. \tag{4.37}
$$

We then study the solutions of  $(4.32)$  decaying to 0 at  $+\infty$ .

<span id="page-26-1"></span>**Proposition 4.6.** *Eq.* [\(4.32\)](#page-25-3) *on*  $(\tilde{x}_{+}, +\infty)$  *admits a unique solution*  $V_1(x, \lambda)$  *such that*  $e^{-\alpha_+(x)}V_1(x)$  converges to  $e_1$  *as*  $x \to +\infty$  *and a unique solution*  $V_2(x,\lambda)$  *such that*  $e^{-\beta_+(x,\lambda)}V_2(x)$  converges to  $e_2$  *as*  $x \to +\infty$ . Furthermore, we have the following *estimates*

<span id="page-26-0"></span>
$$
\|e^{\alpha_+(x)}V_1(x,\lambda)-e_1\|_2 \le 2\Gamma_m(\epsilon_\star) \left(\begin{matrix} (\rho_+ - \rho_0(x)) + \rho_0(\tilde{x}_+)e^{-(\delta(\epsilon_\star)-k)(x-\tilde{x}_+)} \\ + \left|\rho_0(x) - (\delta(\epsilon_\star)-k)\int_{\tilde{x}_+}^x \rho_0(\tau)e^{-(\delta(\epsilon_\star)-k)(x-\tau)}d\tau \right| \end{matrix}\right) \tag{4.38}
$$

*and*

<span id="page-27-4"></span>
$$
\|e^{\beta_+(x,\lambda)}V_2(x,\lambda) - e_2\|_2 \le 2\Gamma_m(\epsilon_\star)(\rho_+ - \rho_0(x))\tag{4.39}
$$

*for all*  $x \geqslant \tilde{x}_+$ *.* 

*Proof.* We define the matrices

$$
\Psi(x,\lambda) = \text{diag}(e^{-\alpha_+(x)}, e^{-\beta_+(x,\lambda)}, e^{\alpha_+(x)}, e^{\beta_+(x,\lambda)}),
$$

and

$$
\Psi_1(x,\lambda) = \text{diag}(0, e^{-\beta_+(x,\lambda)}, 0, 0)
$$

$$
\Psi_2(x,\lambda) = \text{diag}(e^{-\alpha_+(x)}, 0, e^{\alpha_+(x)}, e^{\beta_+(x,\lambda)}).
$$

Then, we consider the equation

<span id="page-27-0"></span>
$$
V_1(x,\lambda) = e^{-\alpha_+(x)}e_1 + \int_{\tilde{x}_+}^x \Psi_1(x,\lambda)\Psi^{-1}(\tau,\lambda)\rho'_0(\tau)M(\tau,\lambda)V_1(\tau,\lambda)d\tau
$$
  

$$
-\int_x^{+\infty}\Psi_2(x,\lambda)\Psi(\tau,\lambda)^{-1}\rho'_0(\tau)M(\tau,\lambda)V_1(\tau,\lambda)d\tau.
$$
 (4.40)

It can be seen that a solution  $V_1$  of  $(4.40)$  satisfies  $(4.32)$ . We solve  $(4.40)$  by the Picard iteration method. Indeed, let  $V_1^{(0)}(x) = 0$  and

$$
V_1^{(j+1)}(x,\lambda) = e^{-\alpha_+(x)}e_1 + \int_{\tilde{x}_+}^x \Psi_1(x,\lambda)\Psi^{-1}(\tau,\lambda)\rho'_0(\tau)M(\tau,\lambda)V_1^{(j)}(\tau,\lambda)d\tau
$$

$$
-\int_x^{+\infty} \Psi_2(x,\lambda)\Psi^{-1}(\tau,\lambda)\rho'_0(\tau)M(\tau,\lambda)V_1^{(j)}(\tau,\lambda)d\tau.
$$

We have that

$$
\Psi_1(x,\lambda)\Psi^{-1}(\tau,\lambda) = \text{diag}(0, e^{-(\beta_+(x,\lambda)-\beta_+(\tau,\lambda))}, 0, 0)
$$

and that

$$
\Psi_2(x,\lambda)\Psi^{-1}(\tau,\lambda) = \text{diag}(e^{-(\alpha_+(\tau)-\alpha_+(x))}, 0, e^{\alpha_+(\tau)-\alpha_+(x)}, e^{-(\beta_+(x,\lambda)-\beta_+(\tau,\lambda))}).
$$
  
Hence, we can estimate for  $\tilde{x}_+ \le \tau \le x$ ,

$$
\|\Psi_1(x,\lambda)\Psi^{-1}(\tau,\lambda)\|_{F} \le e^{-(\alpha_+(x)-\alpha_+(\tau))+\int_{\tau}^x (k-\sigma_0(s))ds}
$$
  

$$
\le e^{-(\alpha_+(x)-\alpha_+(\tau))-(\delta(\epsilon_\star)-k)(x-\tau)}
$$
(4.41)

<span id="page-27-1"></span>and for  $\tau \geqslant x$ ,

<span id="page-27-2"></span>
$$
\|\Psi_2(x,\lambda)\Psi^{-1}(\tau,\lambda)\|_F \leq e^{-(\alpha_+(x)-\alpha_+(\tau))}.\tag{4.42}
$$

Using  $(4.41)$  and  $(4.42)$ , we get

$$
e^{\alpha_+(x)} \|V_1^{(j+1)}(x,\lambda) - V_1^{(j)}(x,\lambda)\|_2
$$
  
\$\leq \Gamma\_m(\epsilon\_\*) \int\_{\tilde{x}\_+}^{\infty} e^{\alpha\_+(\tau)} \rho'\_0(\tau) \|V\_1^{(j)}(\tau,\lambda) - V\_1^{(j-1)}(\tau,\lambda)\|\_F d\tau.\$

Thanks to the induction, we get for all  $x \geq \tilde{x}_+$  and for all  $j \geq 0$ ,

<span id="page-27-3"></span>
$$
e^{\alpha_+(x)} \|V_1^{(j+1)}(x,\lambda) - V_1^{(j)}(x,\lambda)\|_2 \leqslant \left(\frac{1}{2}\right)^j,\tag{4.43}
$$

yielding the uniform convergence of  $\{V_1^{(j)}(x,\lambda)\}_{j\geqslant0}$  on any interval of  $(\tilde{x}_+,\infty)$ . Let  $V_1(x, \lambda)$  be the limit function.  $V_1^{(j)}(x, \lambda)$  is continuous, so is  $V_1(x, \lambda)$ . Moreover, [\(4.43\)](#page-27-3) implies that

$$
e^{\alpha_+(x)}\|V_1^{(j+1)}(x,\lambda)\|_2\leqslant \sum_{i=0}^j e^{\alpha_+(x)}\|V_1^{(i+1)}(x,\lambda)-V_1^{(i)}(x,\lambda)\|\leqslant \sum_{i=0}^j \Big(\frac{1}{2}\Big)^i.
$$

That tells us for  $x \geq \tilde{x}_+$ ,

<span id="page-28-3"></span>
$$
||V_1(x,\lambda)||_2 \leq 2e^{-\alpha_+(x)}.
$$
\n(4.44)

Once we have [\(4.44\)](#page-28-3), we then prove [\(4.38\)](#page-26-0). Indeed,

$$
e^{\alpha_+(x)}V_1(x,\lambda) - e_1 = e^{\alpha_+(x)} \int_{\tilde{x}_+}^x \Psi_1(x,\lambda)\Psi^{-1}(\tau,\lambda)\rho'_0(\tau)M(\tau,\lambda)V_1(\tau,\lambda)d\tau
$$

$$
-e^{\alpha_+(x)} \int_x^{+\infty} \Psi_2(x,\lambda)\Psi(\tau,\lambda)^{-1}\rho'_0(\tau)M(\tau,\lambda)V_1(\tau,\lambda)d\tau.
$$

We make use of  $(4.42)$  and  $(4.44)$  to have that

<span id="page-28-4"></span>
$$
e^{\alpha_+(x)} \int_x^{+\infty} \|\Psi_2(x,\lambda)\Psi^{-1}(\tau,\lambda)\rho'_0(\tau)M(\tau,\lambda)V_1(\tau,\lambda)\|_2 d\tau \leq 2\Gamma_m(\epsilon_\star)(\rho_+ - \rho_0(x)).
$$
\n(4.45)

From  $(4.41)$  and  $(4.44)$ , we obtain that

<span id="page-28-5"></span>
$$
\begin{split} \|e^{\alpha_{+}(x)} \int_{\tilde{x}_{+}}^{x} \Psi_{1}(x,\lambda) \Psi^{-1}(\tau,\lambda) \rho_{0}'(\tau) M(\tau,\lambda) V_{1}(\tau,\lambda) d\tau \|_{2} \\ &\leq 2 \Gamma_{m}(\epsilon_{\star}) \int_{\tilde{x}_{+}}^{x} \rho_{0}'(\tau) e^{-(\delta(\epsilon_{\star}) - k)(x-\tau)} d\tau. \end{split} \tag{4.46}
$$

After integrating by parts, we get

<span id="page-28-6"></span>
$$
\int_{\tilde{x}_{+}}^{x} \rho_{0}'(\tau) e^{-(\delta(\epsilon_{\star}) - k)(x - \tau)} d\tau = -\rho_{0}(\tilde{x}_{+}) e^{-(\delta(\epsilon_{\star}) - k)(x - \tilde{x}_{+})} + \rho_{0}(x)
$$
\n
$$
- (\delta(\epsilon_{\star}) - k) \int_{\tilde{x}_{+}}^{x} \rho_{0}(\tau) e^{-(\delta(\epsilon_{\star}) - k)(x - \tau)} d\tau.
$$
\n(4.47)

Combining [\(4.45\)](#page-28-4), [\(4.46\)](#page-28-5) and [\(4.47\)](#page-28-6) gives [\(4.38\)](#page-26-0).

By considering the eigenvalue  $-\sigma_0(x, \lambda)$  of  $L(x, \lambda)$ , we continue the idea in Theorem [C.1](#page-33-1) and mimic the above arguments to the solution  $V_2(x, \lambda)$  such that  $e^{\beta + (x,\lambda)}V_2(x)$  converges to  $e_2$  at  $+\infty$  and enjoying [\(4.39\)](#page-27-4). That ends the proof of Lemma [4.6.](#page-26-1)  $\Box$ 

Now, we get back to  $(4.3)$  to find solutions that are bounded near  $+\infty$ .

# <span id="page-28-0"></span>**Proposition 4.7.** *Eq.* [\(4.3\)](#page-17-2) *on*  $(\tilde{x}_+, +\infty)$  *admits*

(1) *a unique solution*  $U_1^+(x,\lambda)$  *satisfying that as*  $x \to +\infty$ ,  $e^{\alpha_+(x)}U_2^+(x,\lambda)$ *converges to*  $(-k^{-3}, k^{-2}, -k^{-1}, 1)^T$  *and that for all*  $x \ge \tilde{x}_+$ *,* 

$$
e^{\alpha_{+}(x)}U_{1}^{+}(x,\lambda) - (-k^{-3}, k^{-2}, -k^{-1}, 1)^{T} \|_{2}
$$
  
\n
$$
\leq 2\Gamma_{p}\Gamma_{m}(\epsilon_{\star}) \left( (\rho_{+} - \rho_{0}(x)) + \rho_{0}(\tilde{x}_{+})e^{-(\delta(\epsilon_{\star}) - k)(x - \tilde{x}_{+})} + |\rho_{0}(x) - (\delta(\epsilon_{\star}) - k)\int_{\tilde{x}_{+}}^{x} \rho_{0}(\tau)e^{-(\delta(\epsilon_{\star}) - k)(x - \tau)}d\tau \right), \quad (4.48)
$$

(2) *a unique solution*  $U_2^+(x,\lambda)$  *satisfying that as*  $x \to +\infty$ ,  $e^{\beta_+(x,\lambda)}U_2^+(x,\lambda)$ *converges to*  $(-\sigma_+^{-3}(\lambda), \sigma_+^{-2}(\lambda), -\sigma_+^{-1}(\lambda), 1)^T$  and that for all  $x \ge \tilde{x}_+$ ,

<span id="page-28-2"></span>
$$
\|e^{\beta_+(x,\lambda)}U_2^+(x,\lambda) - (-\sigma_+^{-3}(\lambda), \sigma_+^{-2}(\lambda), -\sigma_+^{-1}(\lambda), 1)^T\|_2
$$
  
\$\leqslant \left(\sqrt{\frac{g(4\delta^{10}(\epsilon\_\star) + 16\delta^{12}(\epsilon\_\star) + 9\delta\_s^4)}{16L\_0\mu^2\delta^{16}(\epsilon\_\star)} + 2\Gamma\_p\Gamma\_m(\epsilon\_\star)\right)(\rho\_+ - \rho\_0(x)).\tag{4.49}

<span id="page-28-1"></span> $\parallel$ 

*Proof.* We define  $U_j^+(x,\lambda) = P(x,\lambda)V_j(x,\lambda)(j = 1,2)$ , with  $V_1$  and  $V_2$  are two solutions of  $(4.32)$  satisfying  $(4.38)$  and  $(4.39)$  respectively. It can be seen that  $U_1^+(x,\lambda)$  and  $U_2^+(x,\lambda)$  are two solution of [\(4.3\)](#page-17-2).

Note that

$$
e^{\alpha_+(x)}U_1^+(x,\lambda) = P_1 + P(x,\lambda)(e^{\alpha_+(x)}V_1(x,\lambda) - e_1)
$$
  
=  $(-k^{-3}, k^{-2}, -k^{-1}, 1)^T + P(x,\lambda)(e^{\alpha_+(x)}V_1(x,\lambda) - e_1).$ 

[\(4.48\)](#page-28-1) is then clear due to the estimate [\(4.38\)](#page-26-0). According to l'Hôpital's rule, we have

$$
\lim_{x \to +\infty} \frac{\int_{\tilde{x}_+}^x \rho_0(\tau) e^{(\delta(\epsilon_\star) - k)\tau} d\tau}{e^{(\delta(\epsilon_\star) - k)x}} = \lim_{x \to +\infty} \frac{\rho_0(x) e^{(\delta(\epsilon_\star) - k)x}}{(\delta(\epsilon_\star) - k) e^{(\delta(\epsilon_\star) - k)x}} = \frac{\rho_+}{\delta(\epsilon_\star) - k},
$$

that implies

$$
\lim_{x \to +\infty} \left| \rho_0(x) - (\delta(\epsilon_\star) - k) \int_{\tilde{x}_+}^x \rho_0(\tau) e^{-(\delta(\epsilon_\star) - k)(x - \tau)} d\tau \right| = 0.
$$

The behavior of  $U_1^+(x,\lambda)$  at  $+\infty$  follows.

To prove 
$$
(4.49)
$$
, we write

$$
e^{\beta_+(x,\lambda)}U_2^+(x,\lambda) - (-\sigma_+^{-3}(\lambda), \sigma_+^{-2}(\lambda), -\sigma_+^{-1}(\lambda), 1)^T
$$
  
=  $P_2(x,\lambda) - (-\sigma_+^{-3}(\lambda), \sigma_+^{-2}(\lambda), -\sigma_+^{-1}(\lambda), 1)^T + P(x,\lambda)(e^{\beta_+(x,\lambda)}V_2(x,\lambda) - e_2).$ 

Since  $\delta(\epsilon_{\star}) < \sigma_{+}(\lambda) < \delta_{s}$  for all  $\lambda \in [\epsilon_{\star}, \sqrt{\frac{g}{L_{0}}}]$ , we bound that

<span id="page-29-2"></span>
$$
\|P_{2}(x,\lambda) - (-\sigma_{+}^{-3}(\lambda), \sigma_{+}^{-2}(\lambda), -\sigma_{+}^{-1}(\lambda), 1)^{T}\|_{2}^{2}
$$
\n
$$
= \frac{\lambda^{2}}{\mu^{2}}(\rho_{0}(x) - \rho_{+})^{2} \left[ \frac{1}{\sigma_{0}^{2}(x,\lambda)\sigma_{+}^{2}(\lambda)(\sigma_{0}(x,\lambda) + \sigma_{+}(\lambda))^{2}} + \frac{1}{\sigma_{0}^{2}(x,\lambda)\sigma_{+}^{2}(\lambda)} - \left(\frac{\sigma_{0}^{2}(x,\lambda) + \sigma_{0}(x,\lambda)\sigma_{+}(\lambda) + \sigma_{+}^{2}(\lambda)}{\sigma_{0}^{3}(x,\lambda)\sigma_{+}^{3}(\lambda)(\sigma_{0}(x,\lambda) + \sigma_{+}(\lambda))}\right)^{2}
$$
\n
$$
\leq \frac{g(4\delta^{10}(\epsilon_{\star}) + 16\delta^{12}(\epsilon_{\star}) + 9\delta_{s}^{4})}{16L_{0}\mu^{2}\delta^{16}(\epsilon_{\star})} (\rho_{0}(x) - \rho_{+})^{2}.
$$
\n(4.50)

Meanwhile, as a result of [\(4.39\)](#page-27-4),

<span id="page-29-3"></span>
$$
||P(x,\lambda)(e^{\beta_+(x,\lambda)}V_2(x,\lambda)-e_2)||_2 \leq 2\Gamma_p \Gamma_m(\epsilon_\star)(\rho_+-\rho_0(x)). \tag{4.51}
$$

Thanks to [\(4.50\)](#page-29-2) and [\(4.51\)](#page-29-3), we obtain [\(4.49\)](#page-28-2) hence the behavior of  $U_2^+(x, \lambda)$  at  $+\infty$ . Proof of Proposition [4.7](#page-28-0) is complete.  $\Box$ 

We now fix two positive eigenvalues of  $L(x, \lambda)$ , *k* and  $\sigma_0(x, \lambda)$  and thus follow Theorem [C.1](#page-33-1) again. We are able to construct solutions of [\(4.3\)](#page-17-2) that are bounded near  $-\infty$  as in Proposition [4.7.](#page-28-0)

<span id="page-29-0"></span>**Proposition 4.8.** *Eq.* [\(4.3\)](#page-17-2) *on*  $(-\infty, \tilde{x}_-)$  *admits* 

(1) *a unique solution*  $U_3^-(x, \lambda)$  *satisfying that as*  $x \to -\infty$ ,  $e^{\alpha_-(x)}U_3^-(x, \lambda)$ *converges to*  $(k^{-3}, k^{-2}, k^{-1}, 1)^T$  *and that, for all*  $x \leq \tilde{x}$ 

<span id="page-29-1"></span>
$$
\|e^{\alpha_{-}(x)}U_{3}^{-}(x,\lambda)-(k^{-3},k^{-2},k^{-1},1)^{T}\|_{2}
$$
  

$$
\leq 2\Gamma_{p}\Gamma_{m}(\epsilon_{\star})\begin{pmatrix}(\rho_{0}(x)-\rho_{-})+\rho_{0}(\tilde{x}_{-})e^{-(\delta(\epsilon_{\star})-k)(\tilde{x}_{-}-x)}\\+\rho_{0}(x)-(\delta(\epsilon_{\star})-k)\int_{x}^{\tilde{x}_{-}}\rho_{0}(\tau)e^{-(\delta(\epsilon_{\star})-k)(\tau-x)}d\tau\end{pmatrix},\tag{4.52}
$$

(2) *a unique solution*  $U_4^-(x, \lambda)$  *satisfying that as*  $x \to -\infty$ ,  $e^{\beta_-(x, \lambda)}U_4^-(x, \lambda)$ *converges to*  $(\sigma_-^{-3}(\lambda), \sigma_-^{-2}(\lambda), \sigma_-^{-1}(\lambda), 1)^T$  *and that for all*  $x \leq \tilde{x}_-,$ 

<span id="page-30-0"></span>
$$
\|e^{\beta_{-}(x,\lambda)}U_{4}^{-}(x,\lambda) - (\sigma_{-}^{-3}(\lambda), \sigma_{-}^{-2}(\lambda), \sigma_{-}^{-1}(\lambda), 1)^{T}\|_{2}
$$
  
\$\leqslant \left(\sqrt{\frac{g(4\delta^{10}(\epsilon\_{\star}) + 16\delta^{12}(\epsilon\_{\star}) + 9\delta\_{s}^{4})}{16L\_{0}\mu^{2}\delta^{16}(\epsilon\_{\star})} + 2\Gamma\_{p}\Gamma\_{m}(\epsilon\_{\star})\right)(\rho\_{0}(x) - \rho\_{-}). \qquad (4.53)\$

We are now in position to prove Proposition [4.5.](#page-24-1)

*Proof.* We recall  $n_{ij}^+$   $(i, j = 1, 2)$  from  $(4.18)$  and  $(4.19)$  to have that

<span id="page-30-1"></span>
$$
\begin{pmatrix} n_{11}^+ \\ n_{12}^+ \end{pmatrix} (x_+, \lambda) = -\frac{1}{U_{11}^+ U_{22}^+ - U_{21}^+ U_{12}^+} \begin{pmatrix} U_{22}^+ & -U_{12}^+ \\ -U_{21}^+ & U_{11}^+ \end{pmatrix} \begin{pmatrix} U_{13}^+ \\ U_{23}^+ \end{pmatrix} (x_+, \lambda) \tag{4.54}
$$

and

<span id="page-30-2"></span>
$$
\begin{pmatrix} n_{21}^+ \\ n_{22}^+ \end{pmatrix} (x_+, \lambda) = -\frac{1}{U_{11}^+ U_{22}^+ - U_{21}^+ U_{12}^+} \begin{pmatrix} U_{22}^+ & -U_{12}^+ \\ -U_{21}^+ & U_{11}^+ \end{pmatrix} \begin{pmatrix} U_{14}^+ \\ U_{24}^+ \end{pmatrix} (x_+, \lambda). \tag{4.55}
$$

We are ready to prove the estimates  $(4.13)$  needed for Lemma [4.1.](#page-20-1) Now using  $(4.48)$ and  $(4.49)$  into  $(4.54)$  yields that

$$
n_{12}^+(x_+,\lambda) = \frac{U_{21}^+ U_{13}^+ - U_{11}^+ U_{23}^+}{U_{11}^+ U_{22}^+ - U_{21}^+ U_{12}^+}(x_+,\lambda) = \frac{k + \sigma_+(\lambda) + f_1(x_+,\lambda)}{1 + f_2(x_+,\lambda)},
$$

where  $|f_j(x, \lambda)| = O(z_{+,\epsilon_*}(x))$   $(j = 1, 2)$  uniformly in  $\lambda \in [\epsilon_*, \sqrt{\frac{g}{L_0}}]$  as  $x \to \infty$ . Hence, there exists a constant  $\xi_+(\epsilon_{\star}) > 0$  such that

$$
n_{12}^+(x_+, \lambda) \geq k + \sigma_+(\lambda) - \xi_+(\epsilon_\star) z_{+,\epsilon_\star}(x_+) \geq k + \delta(\epsilon_\star) - \xi_+(\epsilon_\star) z_{+,\epsilon_\star}(x_+).
$$

That implies  $n_{12}^+(x_+,\lambda) > 0$  if

<span id="page-30-3"></span>
$$
z_{+,\epsilon_{\star}}(x_{+}) < \frac{k + \delta(\epsilon_{\star})}{\xi_{+}(\epsilon_{\star})}.\tag{4.56}
$$

We then estimate

 $\Delta_+(x_+,\lambda) := (n_{11}^+(x_+,\lambda) - n_{22}^+(x_+,\lambda) - k^2 - \sigma_0^2(x_+,\lambda))^2 + 4n_{12}^+n_{21}^+(x_+,\lambda)$ Using  $(4.48)$  and  $(4.49)$  into  $(4.54)$  and  $(4.55)$  again, we have

$$
n_{11}^+(x_+, \lambda) = k\sigma_+(\lambda) + O(z_{+,\epsilon_\star}(x_+)),
$$
  
\n
$$
n_{22}^+(x_+, \lambda) + k^2 + \sigma_0^2(x_+, \lambda) = -k\sigma_+(\lambda) + O(z_{+,\epsilon_\star}(x_+)),
$$
  
\n
$$
n_{21}^+(x_+, \lambda) = -k\sigma_+(\lambda)(k + \sigma_+(\lambda)) + O(z_{+,\epsilon_\star}(x_+)).
$$

Hence, there exists  $w_+(\epsilon_*) > 0$  such that

$$
\Delta_{+}(x_{+},\lambda) \le -4k\sigma_{+}(\lambda)(k^{2}+k\sigma_{+}(\lambda)+\sigma_{+}^{2}(\lambda))+w_{+}(\epsilon_{\star})z_{+,\epsilon_{\star}}(x_{+})
$$
  

$$
\le -4k\delta(\epsilon_{\star})(k^{2}+k\delta(\epsilon_{\star})+\delta^{2}(\epsilon_{\star}))+w_{+}(\epsilon_{\star})z_{+,\epsilon_{\star}}(x_{+}).
$$

The inequality  $\Delta_+(x_+, \lambda) \leq 0$  is equivalent to

<span id="page-30-4"></span>
$$
z_{+,\epsilon_\star}(x_+) \leqslant \frac{4k\delta(\epsilon_\star)(k^2 + k\delta(\epsilon_\star) + \delta^2(\epsilon_\star))}{w_+(\epsilon_\star)}.\tag{4.57}
$$

Combining  $(4.56)$  and  $(4.57)$ , we take

$$
\Gamma_{+}(\epsilon_{\star}) = \min\left(\frac{k + \delta(\epsilon_{\star})}{\xi_{+}(\epsilon_{\star})}, \frac{4k\delta(\epsilon_{\star})(k^{2} + k\delta(\epsilon_{\star}) + \delta^{2}(\epsilon_{\star}))}{w_{+}(\epsilon_{\star})}\right).
$$

<span id="page-31-4"></span>If  $x_+$  satisfies  $z_{+,\epsilon_*}(x_+) \leq \Gamma_+(\epsilon_*)$ , then one has  $n_{12}^+(x_+,\lambda) > 0 \geq \Delta_+(x_+,\lambda)$ , i.e.  $(4.13)$ . That implies  $BV_{x_+, \lambda}(\vartheta, \vartheta) \geq 0$ .

Similarly, we get that, from [\(4.19\)](#page-21-5), we follow the above arguments to show that there exists  $\Gamma_{-}(\epsilon_{\star}) > 0$  such that for  $z_{-,\epsilon_{\star}}(x_{-}) \leq \Gamma_{-}(\epsilon_{\star})$ , [\(4.14\)](#page-20-1) holds. It yields  $BV_{-} \setminus (\theta, \theta) \geq 0$ . Proposition 4.5 is proven  $BV_{x_{-},\lambda}(\vartheta,\vartheta) \geq 0$ . Proposition [4.5](#page-24-1) is proven.

#### <span id="page-31-1"></span>**ACKNOWLEDGMENTS**

The authors would like to thank Prof. Jean-Marc Delort, Prof. Catherine Sulem and Prof. Jeffrey Rauch for fruitful discussions and Prof. Bernard Helffer and Prof. David Lannes for advice on this study. The second author thanks also to Assoc. Prof. QuËc Anh Ngô for his encouragement. Both authors are deeply grateful to one of anonymous referees for his/her valuable comments to improve the presentation of this paper. Parts of this work were carried out while the second author was visiting Université de Montréal and he would like to thank Université de Montréal for the heartfelt hospitality. This work is supported by a grant from Région Île-de-France.

# Appendix A. Proof of Lemma [2.1](#page-6-3)

<span id="page-31-0"></span>Multiplying by  $\phi$  on both sides of [\(1.2\)](#page-3-0) and then integrating by parts, we obtain that

<span id="page-31-2"></span>
$$
-\lambda^{2} \int_{\mathbf{R}} \left( k^{2} \rho_{0} |\phi|^{2} + \rho_{0} |\phi'|^{2} \right) dx = \lambda \mu \int_{\mathbf{R}} \left( |\phi''|^{2} + 2k^{2} |\phi'|^{2} + k^{4} |\phi|^{2} \right) dx
$$
  

$$
- g k^{2} \int_{\mathbf{R}} \rho'_{0} |\phi|^{2} dx.
$$
 (A.1)

Suppose that  $\lambda = \lambda_1 + i\lambda_2$ , then one deduces from [\(A.1\)](#page-31-2) that

$$
-(\lambda_1^2 - \lambda_2^2) \int_{\mathbf{R}} \left( k^2 \rho_0 |\phi|^2 + \rho_0 |\phi'|^2 \right) dx = \lambda_1 \mu \int_{\mathbf{R}} \left( |\phi''|^2 + 2k^2 |\phi'|^2 + k^4 |\phi|^2 \right) dx
$$

$$
- g k^2 \int_{\mathbf{R}} \rho'_0 |\phi|^2 dx
$$
(A.2)

and that

<span id="page-31-3"></span>
$$
-2\lambda_1\lambda_2\int_{\mathbf{R}} \left(k^2\rho_0|\phi|^2 + \rho_0|\phi'|^2\right)dx = \lambda_2\mu\int_{\mathbf{R}} \left(|\phi''|^2 + 2k^2|\phi'|^2 + k^4|\phi|^2\right)dx. \tag{A.3}
$$

If  $\lambda_2 \neq 0$ , [\(A.3\)](#page-31-3) leads us to

$$
-2\lambda_1 \int_{\mathbf{R}} \left( k^2 \rho_0 |\phi|^2 + \rho_0 |\phi'|^2 \right) dx = \mu \int_{\mathbf{R}} \left( |\phi''|^2 + 2k^2 |\phi'|^2 + k^4 |\phi|^2 \right) dx,
$$

which yields

$$
-(\lambda_1^2 - \lambda_2^2) \int_{\mathbf{R}} \left( k^2 \rho_0 |\phi|^2 + \rho_0 |\phi'|^2 \right) dx = -2\lambda_1^2 \int_{\mathbf{R}} \left( k^2 \rho_0 |\phi|^2 + \rho_0 |\phi'|^2 \right) dx - g k^2 \int_{\mathbf{R}} \rho'_0 |\phi|^2 dx.
$$

Equivalently,

$$
(\lambda_1^2 + \lambda_2^2) \int_{\mathbf{R}} \left( k^2 \rho_0 |\phi|^2 + \rho_0 |\phi'|^2 \right) dx = -g k^2 \int_{\mathbf{R}} \rho'_0 |\phi|^2 dx. \tag{A.4}
$$

That implies

$$
(\lambda_1^2+\lambda_2^2)k^2\inf_{\mathbf{R}}\rho_0\int_{\mathbf{R}^2}|\phi|^2dx\leqslant -gk^2\int_{\mathbf{R}}\rho_0'|\phi|^2dx.
$$

The positivity of  $\rho'_0$  yields a contradiction, then  $\lambda$  is real. Using  $(A.1)$  again, we further get that

$$
\lambda^2 \int_{\mathbf{R}} \rho_0 (k^2 |\phi|^2 + |\phi'|^2) dx \leqslant g k^2 \int_{\mathbf{R}} \rho'_0 |\phi|^2 dx.
$$

It tells us that  $\lambda$  is bounded by  $\sqrt{\frac{g}{L_0}}$ . This finishes the proof of Lemma [2.1.](#page-6-3)

# <span id="page-32-0"></span>Appendix B. Differentiability of eigenvalues of self-adjoint and compact operators

The classical perturbation theory from [\[9,](#page-36-11) Chapter VII, \$3] has shown the continuous property of the eigenvalues for a family of holomorphic self-adjoint operators in an infinite-dimensional Hilbert space. If the operators are only differentiable, we will present a proof of the differentiability of the eigenvalues for compact and self-adjoint operators in an infinite-dimensional Hilbert space deduced from that one for matrix functions in a finite-dimensional space (see [\[9,](#page-36-11) Chapter II, \$5]).

Theorem B.1. *Let I be a closed interval and H be an infinite-dimensional Hilbert* space and  $(A(\lambda))_{\lambda \in I}$  be a family of self-adjoint and compact operators in *H* depend*ing continuously differentiable on*  $\lambda$ . Then, all eigenvalues and all eigenvectors of  $A(\lambda)$  are differentiable functions on  $\lambda$ .

*Proof.* Let  $\lambda_0 \in I$  be fixed. Since  $A(\lambda_0)$  is a self-adjoint and compact operator in *H*, the spectrum of  $A(\lambda_0)$  is discrete. Let  $\gamma_0$  be an arbitrary eigenvalue of  $A(\lambda_0)$  and  $E = \text{Ker}(A(\lambda_0) - \gamma_0 \text{Id}_H)$ , we have the decomposition  $H = E \oplus E^{\perp}$ . Consequently, for all  $\lambda \in I$ ,

$$
A(\lambda) = \begin{pmatrix} \text{Proj}_E(A(\lambda)\text{Proj}_E) & \text{Proj}_E(A(\lambda)\text{Proj}_{E^\perp}) \\ \text{Proj}_{E^\perp}(A(\lambda)\text{Proj}_E) & \text{Proj}_{E^\perp}(A(\lambda)\text{Proj}_{E^\perp}) \end{pmatrix}
$$

that we will denote by  $(A_{ij}(\lambda))_{1\leq i,j\leq 2}$  for brevity. Notice that

$$
A(\lambda_0) = \begin{pmatrix} \gamma_0 \mathrm{Id}_E & 0 \\ 0 & A_{22}(\lambda_0) \end{pmatrix},
$$

and  $A_{22}(\lambda_0) - \gamma_0 \text{Id}_{E^\perp}$  is invertible.

Let  $0 < \varepsilon \ll 1$  and  $\gamma$  be an eigenvalue of  $A(\lambda)$  being close to  $\gamma_0$ , i.e.  $|\gamma - \gamma_0| < \varepsilon$ . We write that

$$
A(\lambda) - \gamma \text{Id} = \begin{pmatrix} A_{11}(\lambda) - \gamma \text{Id}_E & A_{12}(\lambda) \\ A_{21}(\lambda) & A_{22}(\lambda) - \gamma \text{Id}_{E^\perp} \end{pmatrix}.
$$
 (B.1)

If  $x = (y, z)^T$  is a corresponding eigenvector, we obtain

$$
\begin{cases} A_{11}(\lambda)y + A_{12}(\lambda)z = \gamma y, \\ A_{21}(\lambda)y + A_{22}(\lambda)z = \gamma z. \end{cases}
$$

Consequently,

<span id="page-32-1"></span>
$$
\begin{cases}\n(A_{11}(\lambda) - \gamma \mathrm{Id}_E)y + A_{12}(\lambda)z = 0, \\
A_{21}(\lambda)y + (A_{22}(\lambda) - \gamma \mathrm{Id}_{E^\perp})z = 0.\n\end{cases}
$$
\n(B.2)

<span id="page-32-2"></span>

Since  $A_{22}(\lambda_0) - \gamma_0 \text{Id}_{E^\perp}$  is invertible, we have that  $A_{22}(\lambda) - \gamma_0 \text{Id}_{E^\perp}$  is invertible for  $|\lambda - \lambda_0| < \delta \ll 1$ . We further get that  $A_{22}(\lambda) - \gamma \mathrm{Id}_{E^\perp}$  is also invertible for  $|\lambda - \lambda_0| < \delta$  and  $|\gamma - \gamma_0| < \varepsilon$ . Hence, we deduce that [\(B.2\)](#page-32-1) is equivalent to

<span id="page-33-2"></span>
$$
\begin{cases}\nz = -(A_{22}(\lambda) - \gamma \mathrm{Id}_{E^\perp})^{-1} A_{21}(\lambda) y, \\
(A_{11}(\lambda) - A_{12}(\lambda) (A_{22}(\lambda) - \gamma \mathrm{Id}_{E^\perp})^{-1} A_{21}(\lambda)) y = \gamma y.\n\end{cases} (B.3)
$$

If  $y = 0$ ,  $(B.3)<sub>1</sub>$  $(B.3)<sub>1</sub>$  implies  $z = 0$ , which is impossible. We have  $y \neq 0$ , this means that if  $\gamma \in (\gamma_0 - \varepsilon, \gamma_0 + \varepsilon)$  is an eigenvalue of  $A(\lambda)$ ,  $(B.3)_2$  $(B.3)_2$  tells us that  $\gamma$  is also an eigenvalue of  $B(\lambda, \gamma)$  defined by

$$
B(\lambda, \gamma) := A_{11}(\lambda) - A_{12}(\lambda)(A_{22}(\lambda) - \gamma \mathrm{Id}_{E^\perp})^{-1} A_{21}(\lambda) : E \to E
$$

Notice that *E* is finite-dimensional thanks to Riesz's theorem.  $B(\lambda, \gamma)$  turns out to be a matrix, having eigenvalues  $\gamma_i(\lambda, \gamma)$  ( $1 \leq j \leq \dim E$ ). Then, there exists *j* such that  $\gamma_j(\lambda, \gamma) = \gamma$ . It follows from [\[9,](#page-36-11) Chapter II, \$5] that  $\gamma_j(\lambda, \gamma)$  and its associated<br>eigenvector are differentiable at  $\lambda_0$  so is  $\gamma$ eigenvector are differentiable at  $\lambda_0$ , so is  $\gamma$ .

#### <span id="page-33-0"></span>Appendix C. Decaying solutions of the linear system of ODEs

We state here the key ingredient for our analysis in Section [4](#page-17-0) due to E. A. Coddington and N. Levinson [\[3,](#page-35-6) Theorem 8.1, Chapter 3].

<span id="page-33-1"></span>Theorem C.1. *We consider a linear system*

<span id="page-33-3"></span>
$$
W'(y) = (A + L(y) + R(y))W(y).
$$
 (C.1)

Let *A* be a constant matrix with characteristic roots  $\mu_j$ ,  $j = 1, \ldots, n$ , all of which *are distinct. Let the matrix L be differentiable and satisfy*

$$
\int_0^\infty \|L'(y)\| dy < \infty \tag{C.2}
$$

*and let*  $L(y) \to 0$  *as*  $y \to \infty$ *. Let the matrix R be integrable and let* 

$$
\int_0^\infty \|R(y)\| dy < \infty. \tag{C.3}
$$

Let the roots of  $det(A + L(y) - \lambda I_n) = 0$  be denoted by  $\lambda_j(y), j = 1, \ldots, n$ . Clearly, *by reordering the*  $\mu_j$  *if necessary,*  $\lim_{y\to\infty} \lambda_j(y) = \mu_j$ *. For a given h, let* 

$$
d_{hj}(y) = Re(\lambda_h(y) - \lambda_j(y)).
$$

*Suppose all*  $\lambda_i$  ( $1 \leq j \leq n$ ) *fall into one of two classes*  $H_1$  *and*  $H_2$ *, where* 

$$
\lambda_j \in H_1 \quad \text{if } \int_0^y d_{hj}(s)ds \to \infty \text{ as } y \to \infty \text{ and } \int_{y_1}^{y_2} d_{hj}(s)ds \ge -K \quad (y_2 \ge y_1 \ge 0),
$$

*and*

$$
\lambda_j \in H_2 \quad \text{if } \int_{y_1}^{y_2} d_{hj}(s) ds \leq K \quad (y_2 \geq y_1 \geq 0),
$$

*where h is fixed and K is a constant. Let p<sup>h</sup> be the eigenvector corresponding to*  $\mu_h$ *, i.e.*  $Ap_h = \mu_h p_h$ *. Hence, there is a solution*  $\phi_h$  *of* [\(C.1\)](#page-33-3) *and*  $a y_0 \in (0, \infty)$  *such that*

$$
\lim_{y \to \infty} \phi_h(y) \exp \Big[ - \int_{y_0}^y \lambda_h(s) ds \Big] = p_h.
$$

# <span id="page-34-0"></span>Appendix D. A remark on the relation between the formulation on  $[x_-, x_+]$  and the formulation on **R** of the viscous RT problem

<span id="page-34-2"></span><span id="page-34-1"></span>**Proposition D.1.** *For all*  $\phi$  *bounded solution of* [\(1.2\)](#page-3-0) *on* **R** *and for all*  $\theta \in H^2(\mathbf{R})$ *, there holds*

$$
\lambda \int_{-\infty}^{+\infty} \rho_0(k^2 \phi \theta + \phi' \theta') dx + \mu \int_{-\infty}^{+\infty} (\phi'' \theta'' + 2k^2 \phi \theta' + k^4 \phi \theta) dx
$$
  
=  $\mathcal{B}_{x_-,x_+, \lambda}(\phi, \theta) + \int_{\mathbf{R}\backslash [x_-,x_+]}\frac{g k^2 \rho'_0}{\lambda} \phi \theta dx.$  (D.1)

We immediately have two remarks from  $(D.1)$ .

(1) In the case of  $\rho'_0$  being compactly supported (supp $\rho'_0 = [-a, a]$ ), we have

$$
\lambda \int_{-\infty}^{+\infty} \rho_0(k^2 \phi \theta + \phi' \theta') dx + \mu \int_{-\infty}^{+\infty} (\phi'' \theta'' + 2k^2 \phi \theta' + k^4 \phi \theta) dx = \mathcal{B}_{a,\lambda}(\phi, \theta). \quad (D.2)
$$

That means  $\mathcal{B}_{x_-,x_+,\lambda}$  is independent of  $x_{\pm}$  if and only if  $x_{-} \leqslant -a < a \leqslant x_+$ .

(2) In the case  $\rho'_0 > 0$  everywhere, for each  $(x_-, x_+)$ , a penalization of  $\mathcal{B}_{x_-,x_+,\lambda}$ by the term  $\int_{x_-}^{x_+}$  $\frac{g k^2 \rho'_0}{\lambda} \phi \theta dx$  is necessary to obtain the ODE [\(1.2\)](#page-3-0) on the whole space.

<span id="page-34-6"></span>*Proof of Proposition [D.1.](#page-34-2)* To prove [\(D.1\)](#page-34-1), we show two following identities

$$
\int_{x_+}^{+\infty} \lambda \rho_0 (k^2 \phi \theta + \phi' \theta') dx + \mu \int_{x_+}^{+\infty} (k^4 \phi \theta + 2k^2 \phi' \theta' + \phi'' \theta'') dx
$$
  
= 
$$
\int_{x_+}^{+\infty} \frac{g k^2 \rho'_0}{\lambda} \phi \theta dx + B V_{x_+, \lambda}(\phi, \theta).
$$
 (D.3)

<span id="page-34-7"></span>and

$$
\int_{-\infty}^{x_{-}} \lambda \rho_{0} (k^{2} \phi \theta + \phi' \theta') dx + \mu \int_{-\infty}^{x_{-}} (k^{4} \phi \theta + 2k^{2} \phi' \theta' + \phi'' \theta'') dx
$$
  
= 
$$
\int_{-\infty}^{x_{-}} \frac{g k^{2} \rho'_{0}}{\lambda} \phi \theta dx + BV_{x_{+},\lambda}(\phi, \theta).
$$
 (D.4)

We have the following remark that we state on  $(x_+, +\infty)$  (a similar expression holds true for  $(-\infty, x_{-})$  that if  $\phi$  is a bounded solution of  $(1.2)$  on  $(x_{+}, +\infty)$ , we then have from Proposition [4.1](#page-18-2) that  $\phi$  satisfies [\(4.8\)](#page-18-1) at  $x_+$ . We integrate by parts to have that

$$
\int_{x_{+}}^{+\infty} \phi'' \theta'' dx = \phi'' \theta' \Big|_{x_{+}}^{+\infty} - \phi''' \theta \Big|_{x_{+}}^{+\infty} + \int_{x_{+}}^{+\infty} \phi^{(4)} \theta dx
$$
  
= -(\phi'' \theta')(x\_{+}) + (\phi''' \theta)(x\_{+}) + \int\_{x\_{+}}^{+\infty} \phi^{(4)} \theta dx. (D.5)

<span id="page-34-5"></span>Because of  $(4.8)$ , we have

<span id="page-34-3"></span>
$$
\phi''(x_+) = -(n_{11}^+ \phi(x_+) + n_{12}^+ \phi'(x_+))
$$
 (D.6)

and

<span id="page-34-4"></span>
$$
\phi'''(x_+) = -(n_{21}^+ \phi(x_+) + n_{22}^+ \phi'(x_+)).
$$
\n(D.7)

Substituting  $(D.6)$  and  $(D.7)$  into  $(D.5)$ , we obtain

<span id="page-35-7"></span>
$$
\int_{x_+}^{+\infty} \phi'' \theta'' dx = (n_{11}^+ \phi(x_+) + n_{12}^+ \phi'(x_+)) \theta'(x_+) - (n_{21}^+ \phi(x_+) + n_{22}^+ \phi'(x_+)) \theta(x_+) + \int_{x_+}^{+\infty} \phi^{(4)} \theta dx.
$$
\n(D.8)

<span id="page-35-8"></span>Using the integration by parts again, we deduce that

$$
\int_{x_+}^{+\infty} \phi' \theta' dx = \phi' \theta \Big|_{x_+}^{+\infty} - \int_{x_+}^{+\infty} \phi'' \theta dx = -(\phi' \theta)(x_+) - \int_{x_+}^{+\infty} \phi'' \theta dx \tag{D.9}
$$

<span id="page-35-9"></span>and that

$$
\int_{x_+}^{+\infty} \rho_0 \phi' \theta' dx = -(\rho_0 \phi' \theta)(x_+) - \int_{x_+}^{+\infty} (\rho_0 \phi')' \theta dx.
$$
 (D.10)

In view of  $(D.8)$ ,  $(D.9)$  and  $(D.10)$ , we obtain

<span id="page-35-10"></span>
$$
\int_{x_{+}}^{+\infty} \lambda \rho_{0}(k^{2} \phi \theta + \phi' \theta') dx + \mu \int_{x_{+}}^{+\infty} (k^{4} \phi \theta + 2k^{2} \phi' \theta' + \phi'' \theta'') dx
$$
  
\n
$$
= \int_{x_{+}}^{+\infty} (\lambda (k^{2} \rho_{0} \phi - (\rho_{0} \phi')') + \mu (\phi^{(4)} - 2k^{2} \phi'' + k^{4} \phi)) \theta dx
$$
  
\n
$$
+ \mu (n_{11}^{+} \phi(x_{+}) + n_{12}^{+} \phi'(x_{+})) \theta'(x_{+}) - \mu (n_{21}^{+} \phi(x_{+}) + n_{22}^{+} \phi'(x_{+})) \theta(x_{+})
$$
  
\n
$$
- 2k^{2} \mu (\phi' \theta)(x_{+}) - \lambda (\rho_{0} \phi' \theta)(x_{+})
$$
  
\n
$$
= \int_{x_{+}}^{+\infty} \frac{g k^{2} \rho_{0}'}{\lambda} \phi \theta dx + B V_{x_{+},\lambda}(\phi, \theta).
$$
 (D.11)

Hence,  $(D.3)$  follows from  $(D.11)$ . Similarly, we obtain  $(D.4)$ . It follows from  $(D.3)$ and [\(D.4\)](#page-34-7) that

$$
\lambda \int_{-\infty}^{+\infty} \rho_0(k^2 \phi \theta + \phi' \theta') dx + \mu \int_{-\infty}^{+\infty} (\phi'' \theta'' + 2k^2 \phi \theta' + k^4 \phi \theta) dx
$$
  
\n
$$
= \int_{-\infty}^{x_-} \frac{g k^2 \rho'_0}{\lambda} \phi \theta dx + \int_{x_+}^{+\infty} \frac{g k^2 \rho'_0}{\lambda} \phi \theta dx + BV_{x_-,\lambda}(\phi, \theta) + BV_{x_+,\lambda}(\phi, \theta)
$$
  
\n
$$
+ \int_{x_-}^{x_+} \lambda \rho_0(k^2 \phi \theta + \phi' \theta') dx + \mu \int_{x_-}^{x_+} (k^4 \phi \theta + 2k^2 \phi' \theta' + \phi'' \theta'') dx
$$
  
\n
$$
= \int_{-\infty}^{x_-} \frac{g k^2 \rho'_0}{\lambda} \phi \theta dx + \int_{x_+}^{+\infty} \frac{g k^2 \rho'_0}{\lambda} \phi \theta dx + \mathcal{B}_{x_-, x_+,\lambda}(\phi, \theta).
$$
  
\nIt yields (D.1).

#### <span id="page-35-0"></span>**REFERENCES**

- <span id="page-35-2"></span>[1] S. CHANDRASEKHAR, Hydrodynamics and Hydromagnetic Stability, Oxford University Press, London, 1961. [1,](#page-1-7) [2,](#page-2-3) [4,](#page-4-0) [5](#page-5-1)
- <span id="page-35-3"></span>[2] C. CHERFILS AND O. LAFITTE, Analytic solutions of the Rayleigh equation for linear density profiles Phys. Rev. E 62, 2967 (2000) [2](#page-2-3)
- <span id="page-35-6"></span>[3] E. A. CODDINGTON AND N. LEVINSON, Theory of Ordinary Differential Equations, McGraw-Hill, New York, 1955. [6,](#page-6-7) [16,](#page-16-5) [32](#page-32-2)
- <span id="page-35-1"></span>[4] Y. Guo, H. J. Hwang, On the dynamical Rayleigh–Taylor instability, *Arch. Rational Mech. Anal.* 167 (2003), pp. 235–253. [1,](#page-1-7) [2](#page-2-3)
- <span id="page-35-4"></span>[5] Y. Guo, I. Tice, Linear Rayleigh-Taylor instability for viscous, compressible fluids, *SIAM J. Math. Anal.* 42 (2011), pp. 1688–1720. [3](#page-3-2)
- <span id="page-35-5"></span>[6] E. Grenier, On the nonlinear instability of Euler and Prandtl equations, *Commun. Pure Appl. Math.* 53, (2000), pp. 1067–1091. [4](#page-4-0)
- <span id="page-36-14"></span>[7] B. Helffer, Spectral Theory and its Applications (2013) (Cambridge Studies in Advanced Mathematics). Cambridge: Cambridge University Press. doi:10.1017/CBO9781139505727. [12](#page-12-4)
- <span id="page-36-0"></span>[8] B. HELFFER, O. LAFITTE, Asymptotic methods for the eigenvalues of the Rayleigh equation for the linearized Rayleigh-Taylor instability, *Asymptotic Analysis* 33 (2003), pp. 189–235. [1,](#page-1-7) [2,](#page-2-3) [3](#page-3-2)
- <span id="page-36-11"></span>[9] T. Kato, Perturbation theory for linear operators, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition. [4,](#page-4-0) [13,](#page-13-3) [31,](#page-31-4) [32](#page-32-2)
- <span id="page-36-5"></span>[10] H. Kull, Theory of the Rayleigh-Taylor instability, *Phys. Rep.* 206 (1991), pp. 197–325. [1](#page-1-7)
- <span id="page-36-10"></span>[11] F. Jiang, S. Jiang, G. Ni, Nonlinear instability for nonhomogeneous incompressible viscous fluids, *Sci. China Math.* 56 (2013), pp 665–686. [3](#page-3-2)
- <span id="page-36-8"></span>[12] O. Lafitte, Sur la phase linéaire de l'instabilité de Rayleigh-Taylor. Séminaire Equations aux Dérivées Partielles du Centre de Mathématiques de l'Ecole Polytechnique, Année 2000–2001. [1,](#page-1-7) [2,](#page-2-3) [23](#page-23-5)
- <span id="page-36-3"></span>[[1](#page-1-7)3] J. D. LINDL, 1998. Inertial Confinement Fusion. Springer. 1
- <span id="page-36-9"></span>[14] K. O. MIKAELIAN, Connection between the Rayleigh and the Schrödinger equations, *Phys. Rev. E* 53, 3551. [2](#page-2-3)
- <span id="page-36-12"></span>[15] T.-T. NGUYỄN, Linear and nonlinear analysis of the Rayleigh-Taylor system with Navierslip boundary conditions, preprint, https://arxiv.org/abs/2204.09857. [4](#page-4-0)
- <span id="page-36-13"></span>[16] T.-T. Nguyễn, Nonlinear Rayleigh-Taylor instability of viscous wave in an infinitely deep ocean, work in progress. [4](#page-4-0)
- <span id="page-36-15"></span>[17] T.-T. NGUYỄN, Etude des instabilités non linéaires autour de solutions laminaires de systèmes d'EDP de la mécanique des fluides ou des modèles de propagation en biologie, PhD thesis, in preparation. [23](#page-23-5)
- <span id="page-36-4"></span>[18] B. A. Remington, R. P. Drake, H. Takabe, D. Arnett, A review of astrophysics experiments on intense lasers, *Phys. Plasmas* 7 (2000), pp. 1641–1652. [1](#page-1-7)
- <span id="page-36-1"></span>[19] J.W. STRUTT (LORD RAYLEIGH), Investigation of the character of the equilibrium of an incompressible heavy fluid of variable density, *Proc. London Math. Soc.* 14 (1883), pp. 170–177. [1](#page-1-7)
- <span id="page-36-2"></span>[20] G. Taylor, The instability of liquid surfaces when accelerated in a direction perpendicular to their planes, *Proc. R. Soc. Lond. Ser. A* 201 (1950), pp. 192–196. [1](#page-1-7)
- <span id="page-36-6"></span>[21] Y. Zhou, Rayleigh–Taylor and Richtmyer–Meshkov instability induced flow, turbulence, and mixing. I., *Phys. Rep.* 720–722 (2017), pp. 1–136. [1](#page-1-7)
- <span id="page-36-7"></span>[22] Y. Zhou, Rayleigh–Taylor and Richtmyer–Meshkov instability induced flow, turbulence, and mixing. II., *Phys. Rep.* 723–725 (2017), pp. 1–160. [1](#page-1-7)