



HAL
open science

A uniqueness criterion and a counterexample to regularity in an incompressible variational problem

Marcel Dengler, Jonathan J. Bevan

► **To cite this version:**

Marcel Dengler, Jonathan J. Bevan. A uniqueness criterion and a counterexample to regularity in an incompressible variational problem. 2022. hal-03860861

HAL Id: hal-03860861

<https://hal.science/hal-03860861v1>

Preprint submitted on 18 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

A UNIQUENESS CRITERION AND A COUNTEREXAMPLE TO REGULARITY IN AN INCOMPRESSIBLE VARIATIONAL PROBLEM

M. DENGLER AND J.J. BEVAN

ABSTRACT. In this paper we consider the problem of minimizing functionals of the form $E(u) = \int_B f(x, \nabla u) dx$ in a suitably prepared class of incompressible, planar maps $u : B \rightarrow \mathbb{R}^2$. Here, B is the unit disk and $f(x, \xi)$ is quadratic and convex in ξ . It is shown that if u is a stationary point of E in a sense that is made clear in the paper, then u is a unique global minimizer of $E(u)$ provided the gradient of the corresponding pressure satisfies a suitable smallness condition. We apply this result to construct a non-autonomous, uniformly convex functional $f(x, \xi)$, depending smoothly on ξ but discontinuously on x , whose unique global minimizer is the so-called N -covering map, which is Lipschitz but not C^1 .

1. INTRODUCTION

Let $B \subset \mathbb{R}^2$ be the unit ball. For any $u \in W^{1,2}(B, \mathbb{R}^2)$, define the energy $E(u)$ by

$$(1) \quad E(u) = \int_B f(x, \nabla u) dx,$$

where the integrand is quadratic in the gradient argument

$$(2) \quad f(x, \xi) = M(x)\xi \cdot \xi \text{ for any } x \in B \text{ and } \xi \in \mathbb{R}^{2 \times 2},$$

and where $M \in L^\infty(B, \mathbb{R}^{16})$ is symmetric, i.e. $M_{ijkl} = M_{klij}$ for all $i, j, k, l \in \{1, 2\}$.¹ Furthermore, we require that there exists a constant $\nu > 0$ s.t.

$$(3) \quad M(x)\xi \cdot \xi \geq \nu |\xi|^2 \text{ for a.e. } x \in B \text{ and all } \xi \in \mathbb{R}^{2 \times 2},$$

so that $f(x, \xi)$ is uniformly convex in ξ .

Assume that g is the trace of a map $u_0 \in W^{1,2}(B, \mathbb{R}^2)$ that satisfies $\det \nabla u_0 = 1$ a.e. in B , so that the class

$$(4) \quad \mathcal{A}^g := \{u \in W^{1,2}(B, \mathbb{R}^2) : \det \nabla u = 1 \text{ a.e. in } B, u|_{\partial B} = g\}$$

is, in particular, nonempty. The constrained minimization problem that we study in this paper is then to find

$$(5) \quad \min_{u \in \mathcal{A}^g} E(u)$$

in \mathcal{A}^g . Concrete instances of g for which \mathcal{A}^g is nonempty include:

- (a) $g(x) := Ax$, $x \in \partial B$, where A is any constant matrix in $SL^+(2, \mathbb{R})$, and
- (b) $g(\theta) := \frac{1}{\sqrt{N}} e_R(N\theta)$, where $N \in \mathbb{N} \setminus \{0\}$ and $0 \leq \theta \leq 2\pi$.

1991 *Mathematics Subject Classification.* 49K10, 49K20, 73C15, 73C50.

Key words and phrases. Calculus of Variations, Elasticity, Uniqueness, Counterexample to Regularity.

¹ $M \in L^\infty$ is sufficient to guarantee $E(u) < \infty$ for any $u \in W^{1,2}$.

Note that the latter is the trace of the so-called N -covering map

$$(6) \quad u_N(R, \theta) = \frac{R}{\sqrt{N}} e_R(N\theta),$$

expressed in plane polar coordinates, and where we employ the notation $e_R(\theta) = (\cos \theta, \sin \theta)$.

Definition 1.1. (*Stationary point*) *We say that u is a stationary point of $E(\cdot)$ if there exists a function λ , which we shall henceforth refer to as a pressure, belonging to $W^{1,1}(B)$ and such that*

$$(7) \quad \operatorname{div}(\nabla_\xi f(x, \nabla u) + 2\lambda(x) \operatorname{cof} \nabla u) = 0 \quad \text{in } \mathcal{D}'(B).$$

The first main result we obtain shows that if u is a stationary point of the energy E whose corresponding pressure λ satisfies, in addition, the assumption that

$$(8) \quad \|\nabla \lambda(x) R\|_{L^\infty(B, \mathbb{R}^2)} \leq \frac{\sqrt{3}\nu}{2\sqrt{2}},$$

then u is a global minimizer of E . We think of the estimate (8) as characterizing ‘smallness’ of the pressure; concrete examples (such as can be found in [3, Proposition 3.4] or [5], for instance) show that such an estimate need not hold in general. In the following we assume that g has been chosen and fixed so that \mathcal{A}^g is non-empty.

Theorem 1.2 (Uniqueness under small pressure). *Let the energy functional $E(u)$ be given by (1), and let $f(x, \xi)$ be given by (2), where $M \in L^\infty(B, \mathbb{R}^{16})$ is symmetric and satisfies (3) for some $\nu > 0$. Let u be a stationary point of E in the sense of (7) and assume that the corresponding pressure λ satisfies*

$$(9) \quad \|\nabla \lambda(x) R\|_{L^\infty(B, \mathbb{R}^2)} \leq \frac{\sqrt{3}\nu}{2\sqrt{2}}.$$

Then u is a global minimizer of E in \mathcal{A}^g .

Moreover, if the inequality is strict, i.e. $|\nabla \lambda(x) R| < \frac{\sqrt{3}\nu}{2\sqrt{2}}$ on a measurable set $U \subset B$ with $\mathcal{L}^2(U) > 0$ then u is the unique global minimizer of E in \mathcal{A}^g .²

In our second main result we provide an explicit integrand $f(x, \xi)$ of the form (2) whose corresponding energy functional E is minimized in \mathcal{A}^g by the N -covering map u_N . Here, g is the trace of u_N as defined in (6). For its construction we make use of Theorem 1.2. A novelty of our approach is that, in order to apply Theorem 1.2, we develop a method to compute the corresponding pressure explicitly.

Theorem 1.3 (Counterexample to regularity). *Let g be the trace of the N -covering map u_N , let $N \in \mathbb{N} \setminus \{1\}$ and let $a \in (N^2 - N, N^2 + N)$. Define for $x \in B \setminus \{0\}$ and $\xi \in \mathbb{R}^{2 \times 2}$ the function*

$$f(x, \xi) = \nu [a(e_R^T \xi e_R)^2 + (e_R^T \xi e_\theta)^2 + a(e_\theta^T \xi e_R)^2 + (e_\theta^T \xi e_\theta)^2],$$

where $\nu > 0$. Then the following statements are true:

²Here the norm is defined via $\|f\|_{L^\infty(B, \mathbb{R}^2, \mu)} := \max\{\|f_1\|_{L^\infty(B, \mu)}, \|f_2\|_{L^\infty(B, \mu)}\}$ and $\|f_i\|_{L^\infty(B, \mu)} := \lim_{p \rightarrow \infty} \left(\int_B |f_i(x)|^p d\mu \right)^{\frac{1}{p}}$.

(i) There exists $M \in L^\infty(B, \mathbb{R}^{16})$ such that

$$f(x, \xi) = \nu M(x) \xi \cdot \xi$$

for any $x \in B \setminus \{0\}$, $\xi \in \mathbb{R}^{2 \times 2}$ and where M is of the explicit form³

$$\begin{aligned} M(x) = & a((e_R \otimes e_R)(e_R \otimes e_R)) + ((e_R \otimes e_\theta)(e_R \otimes e_\theta)) \\ & + a((e_\theta \otimes e_R)(e_\theta \otimes e_R)) + ((e_\theta \otimes e_\theta)(e_\theta \otimes e_\theta)) \end{aligned}$$

and satisfies $M(x) \geq \nu Id$ for any $x \in B \setminus \{0\}$ and any $N \in \mathbb{N} \setminus \{1\}$.

- (ii) The maps $x \mapsto M(x)$ and $x \mapsto f(x, \xi)$, for any $\xi \in \mathbb{R}^{2 \times 2} \setminus \{0\}$, are discontinuous at 0.
 (iii) The maps $x \mapsto M(x)$ and $x \mapsto f(x, \xi)$, for any $\xi \in \mathbb{R}^{2 \times 2} \setminus \{0\}$, belong to

$$W^{1,q} \setminus W^{1,2} \text{ for any } 1 \leq q < 2$$

with the spaces (B, \mathbb{R}^{16}) and (B) respectively.

(iv) The map

$$(10) \quad u = u_N \in C^{0,1}(B, \mathbb{R}^2) \setminus C^1(B, \mathbb{R}^2)$$

is a stationary point of E , as defined in (7), and the corresponding pressure λ exists and satisfies $\lambda \in W^{1,q}(B)$ for any $1 \leq q < 2$.

- (v) Moreover, the map u_N is the unique global minimizer of E in the class \mathcal{A}^g .
 (vi) The minimum energy is given by

$$\min_{v \in \mathcal{A}^g} E(v) = \frac{\nu\pi}{2}(1+a) \left(\frac{1}{N} + N \right).$$

The problem of studying a functional of the form $E(u)$ is of interest not least because the regularity and/or uniqueness of minimizers of such incompressible problems cannot necessarily be determined *a priori*. Concerning uniqueness in the compressible setting, works including but not limited to John [7], Knops and Stuart [10], Sivaloganathan [16], Zhang [22], and Sivaloganathan and Spector [17] provide conditions under which the uniqueness of a global minimizer can be expected. By contrast, a striking example given by Spadaro [18] clearly demonstrates that global minimizers need not be unique, even under full displacement boundary conditions. When the domain of integration, or reference configuration, is an annulus, a number of papers, including those of John [7], Post and Sivaloganathan [12], Taheri [20], and Morris and Taheri [11, 21], address uniqueness. With the topology of the annulus at their disposal, a multiplicity of solutions/equilibria can be generated by working with certain homotopy classes. For example, Morris and Taheri [11, 21] consider functionals of the form $W(x, s, \xi) = F(|x|^2, |s|^2)|\xi|^2/2$, with $F \in C^2$, on the annulus A and admissible maps $\mathcal{A} = W_{\text{id}}^{1,2}(A, \mathbb{R}^2)$, and show that there are countably many solutions, with exactly one for each homotopy class.

In the homogeneous, incompressible elasticity setting, Knops and Stuart [10, Section 6] (see also [15]) show that the equilibrium solutions they consider are, when subject to affine displacement boundary conditions, global minimizers of the associated energy. Recent results [4] show that there are polyconvex energies with inhomogeneous integrands that, under pure displacement boundary conditions, possess countably many pairs of equal-energy stationary

³Here the multiplication is understood through its action on $\xi \in \mathbb{R}^{2 \times 2}$ which is given by

$$((a \otimes b)(c \otimes d))\xi \cdot \xi = (a \otimes b)_{ij}\xi_{ij}(c \otimes d)_{kl}\xi_{kl} \text{ for } i, j, k, l \in \{1, 2\}.$$

points. It is an open question whether the lowest-energy pair of such stationary points represent global minimizers.

The regularity of equilibrium solutions or minimizers in incompressible variational problems is, like its compressible counterpart, a delicate matter. Ball [1, §2.6] points out that any minimizer of the Dirichlet energy in the class $W^{1,2}(B; \mathbb{R}^2)$, subject to the pointwise (incompressibility) constraint $\det \nabla u = 1$ a.e and boundary condition $u(1, \theta) = \frac{1}{\sqrt{2}}(\cos(2\theta), \sin(2\theta))$, must fail to be C^1 . There is evidence to suggest that the double-covering map u_2 (see (6)) itself may be the global minimizer in that particular problem [3, 5]. A partial regularity result for Lipschitz minimizers that are subject to a type of monotonicity condition was established in [6], and Karakhanyan [8, 9] proves that, in the case of the Dirichlet energy, bounds on the so-called dual pressure lead, by a sophisticated argument, to the conclusion that suitably defined equilibrium solutions must belong to the Hölder class $C_{\text{loc}}^{\frac{1}{2}}$. It is perhaps significant that Karakhanyan's results, like ours, also depend on 'pressure bounds', but more significant still that the maps he deals with are measure-preserving. The double-covering map u_2 mentioned above, and indeed the N -covering maps which form the basis of the counterexample to regularity in Section 3 of this paper, do not preserve \mathcal{L}^2 -measure in the sense of [19, Eq.(24)], for example, and so are less relevant to physically realistic models of elasticity.

It seems that pressure regimes can be used to divide the sorts of incompressible problems we consider into two classes. The double-covering problem introduced by Ball appears to lie in the 'high pressure' regime⁴, whereas the problem we focus on falls, by design, into the 'low pressure' regime, where we can say a bit more.

Let $v, u \in \mathcal{A}^g$ and suppose that u is a stationary point of E in the sense of (7). To compare $E(v)$ and $E(u)$ we set $\eta = v - u$ and expand $E(v) = E(u + \eta)$ as

$$E(v) = E(u) + E(\eta) + \int_B 2M(x) \nabla u(x) \cdot \nabla \eta(x) dx$$

Our problem, as expressed in (5), is made more tractable by the observation made in [3] that the stationarity condition (7) allows us, at the expense of incorporating a pressure term, to rewrite the final, affine-in- $\nabla \eta$ term in the expansion above as a term that is quadratic in $\nabla \eta$, namely

$$\int_B M(x) \nabla u(x) \cdot \nabla \eta(x) dx = \int_B \lambda(x) \det \nabla \eta(x) dx.$$

In particular,

$$E(v) = E(u) + \int_B |\nabla \eta|^2 + 2\lambda \det \nabla \eta dx.$$

For the details, see (12) and the foregoing discussion.

Plan of the paper: The main purpose of Section 2 will be to prove the uniqueness result, Theorem 1.2. We begin by giving two technical lemmas, the first of which enables us to decompose certain expressions in terms of Fourier modes. Section 2 concludes with the proof of Theorem 1.2, together with an argument which shows that the prefactor $3^{\frac{1}{2}}2^{-\frac{3}{2}} \simeq 0.6123$

⁴By which we mean that the pressure λ_2 , say, appearing in the equilibrium equations associated with u_2 obeys $\|R \nabla \lambda_2\|_{\infty} = 3\nu$, when adapted to the notation we use in this paper. The prefactor of ν in the latter exceeds the prefactor $3^{\frac{1}{2}}2^{-\frac{3}{2}}$ appearing in the condition (9) of Theorem 1.2, which is why we refer to this as the 'high pressure' regime.

appearing in (9) can be replaced by 1 when λ depends on just one of the variables R, θ . See Corollary 2.3. The focus of Section 3 is Theorem 1.3. In order to obtain this result we first develop a method to compute the pressure explicitly: this is done for a quite general situation in Lemma 3.1, and then more concretely in Lemmata 3.2-3.3.

1.1. Notation. For a 2×2 -matrix A the cofactor is given by

$$(11) \quad \text{cof } A = \begin{pmatrix} a_{22} & -a_{21} \\ -a_{12} & a_{11} \end{pmatrix},$$

and we define the matrix J via

$$J := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

For two vectors $a \in \mathbb{R}^n, b \in \mathbb{R}^m$ we define the tensor product $a \otimes b \in \mathbb{R}^{n \times m}$ by $(a \otimes b)_{ij} := (ab^T)_{ij} = a_i b_j$ for all $1 \leq i \leq n, 1 \leq j \leq m$. When φ is suitably differentiable, we recall that $\det \nabla \varphi = J \varphi_R \cdot \varphi_\tau$, where φ_R and $\varphi_\tau = \frac{1}{R} \varphi_\theta$ are, respectively, the radial and angular derivatives of φ . We use \mathcal{L}^2 to denote two-dimensional Lebesgue measure. For any $k \in \mathbb{N} \setminus \{0\}$ and $f : B \rightarrow \mathbb{R}$ measurable we define the norm $\|f\|_{L^2(dx/R^k)} := \left(\int_B |f(x)|^2 \frac{dx}{R^k} \right)^{\frac{1}{2}}$. For a measurable vector-valued $f = (f_1, f_2) : B \rightarrow \mathbb{R}^2$ we define $\|f\|_{L^2(dx/R^k)} := \left(\int_B |f_1(x)|^2 + |f_2(x)|^2 \frac{dx}{R^k} \right)^{\frac{1}{2}}$. All other notation is either standard or is defined when it is used.

2. UNIQUENESS IN THE SMALL PRESSURE REGIME

To prove Theorem 1.2 we need two technical lemmas. The first contains basic properties of functions in the class $W^{1,1}(B)$ that satisfy $\|R \nabla \lambda\|_{L^\infty(B, \mathbb{R}^2)} < \infty$, and it relies on a standard Fourier decomposition which, when applied to $\eta \in C^\infty(B, \mathbb{R}^2)$, is given by:

$$\eta(x) = \sum_{j \geq 0} \eta^{(j)}(x), \quad \text{where } \eta^{(0)}(x) = \frac{1}{2} A_0(R), \quad A_0(R) = \frac{1}{2\pi} \int_0^{2\pi} \eta(R, \theta) d\theta$$

and, for any $j \geq 1$,

$$\eta^{(j)}(x) = A_j(R) \cos(j\theta) + B_j(R) \sin(j\theta),$$

where

$$A_j(R) = \frac{1}{2\pi} \int_0^{2\pi} \eta(R, \theta) \cos(j\theta) d\theta \quad \text{and} \quad B_j(R) = \frac{1}{2\pi} \int_0^{2\pi} \eta(R, \theta) \sin(j\theta) d\theta.$$

For later use, we set $\tilde{\eta} := \eta - \eta^{(0)}$.

Lemma 2.1. *Let $\lambda \in W^{1,1}(B)$ and assume that $\|R \nabla \lambda\|_{L^\infty(B, \mathbb{R}^2)} < \infty$. Then the following statements are true:*

- i) $\lambda \in BMO(B)$.
- ii) If $\varphi_n \rightarrow \varphi \in W^{1,2}(B, \mathbb{R}^2)$ then $\int_B \lambda(x) \det \nabla \varphi_n dx \rightarrow \int_B \lambda(x) \det \nabla \varphi dx$.
- iii) It holds $\int_B |\nabla \varphi|^2 dx = \sum_{j \geq 0} \int_B |\nabla \varphi^{(j)}|^2 dx$ for any $\varphi \in W^{1,2}(B, \mathbb{R}^2)$.

iv) $\det \nabla \varphi^{(0)} = 0$ for any $\varphi \in W^{1,2}(B, \mathbb{R}^2)$.

v) $\int_B \lambda(x) \det \nabla \varphi \, dx = -\frac{1}{2} \int_B ((\operatorname{cof} \nabla \varphi) \nabla \lambda(x)) \cdot \varphi \, dx$ for any $\varphi \in W_0^{1,2}(B, \mathbb{R}^2)$.

vi) $\int_B \lambda(x) \det \nabla \varphi \, dx = -\frac{1}{2} \int_B ((\operatorname{cof} \nabla \varphi^{(0)}) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx - \frac{1}{2} \int_B ((\operatorname{cof} \nabla \varphi) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx$ for any $\varphi \in W_0^{1,2}(B, \mathbb{R}^2)$.

Proof. i)-iv) For a proof of these points, see [3, Lem 3.2 and Prop 3.2]. The argument given there still applies if λ depends on x instead of R .

v) Assuming $\varphi \in C_c^\infty(B)$, a computation shows:

$$\begin{aligned}
\int_B \lambda(x) \det \nabla \varphi \, dx &= \int_B \lambda(x) J\varphi_{,R} \cdot \varphi_{,\theta} \frac{dx}{R} \\
&= - \int_B (\lambda(x) J\varphi_{,R})_{,\theta} \cdot \varphi \frac{dx}{R} \\
&= - \int_B \lambda(x)_{,\theta} J\varphi_{,R} \cdot \varphi \frac{dx}{R} - \int_B \lambda(x) J\varphi_{,R\theta} \cdot \varphi \frac{dx}{R} \\
&= - \int_B \lambda(x)_{,\theta} J\varphi_{,R} \cdot \varphi \frac{dx}{R} + \int_B \lambda(x)_{,R} J\varphi_{,\theta} \cdot \varphi \frac{dx}{R} \\
&\quad + \int_B (\lambda(x) J\varphi_{,\theta}) \cdot \varphi_{,R} \frac{dx}{R} \\
&= - \int_B ((\operatorname{cof} \nabla \varphi) \nabla \lambda(x)) \cdot \varphi \, dx - \int_B (\lambda(x) J\varphi_{,R}) \cdot \varphi_{,\theta} \frac{dx}{R}
\end{aligned}$$

The result follows by bringing the rightmost term to the left-hand side and dividing by two. Note, as a last step, that one needs to upgrade the above equation to hold not just for $\varphi \in C_c^\infty(B)$ but instead for all $\varphi \in W_0^{1,2}(B)$. This is slightly delicate because of the weak spaces involved: for a proof, see [3, Lem 3.2.(iv)].

vi) This is a version of (v) in which we emphasise the way that the above expression depends on $\varphi^{(0)}$. Again, we assume $\varphi \in C_c^\infty(B)$, and we start by noting $\varphi_{,\theta} = \tilde{\varphi}_{,\theta}$, hence,

$$\begin{aligned}
\int_B \lambda(x) \det \nabla \varphi \, dx &= \int_B \lambda(x) J\varphi_{,R} \cdot \tilde{\varphi}_{,\theta} \frac{dx}{R} \\
&= - \int_B ((\operatorname{cof} \nabla \varphi) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx + \int_B (\lambda(x) J\tilde{\varphi}_{,\theta}) \cdot \tilde{\varphi}_{,R} \frac{dx}{R}
\end{aligned}$$

then the rightmost term is just

$$\int_B (\lambda(x) J\tilde{\varphi}_{,\theta}) \cdot \tilde{\varphi}_{,R} \frac{dx}{R} = - \int_B (\lambda(x) J\tilde{\varphi}_{,R}) \cdot \tilde{\varphi}_{,\theta} \frac{dx}{R} = \frac{1}{2} \int_B ((\operatorname{cof} \nabla \tilde{\varphi}) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx,$$

together with the above we get

$$\int_B \lambda(x) \det \nabla \varphi \, dx = -\frac{1}{2} \int_B ((\operatorname{cof} \nabla \varphi^{(0)}) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx - \frac{1}{2} \int_B ((\operatorname{cof} \nabla \varphi) \nabla \lambda(x)) \cdot \tilde{\varphi} \, dx.$$

□

The uniqueness condition will be of the form $\|\nabla \lambda(x) R\|_{L^\infty(B, \mathbb{R}^2)} \leq C$, for some constant $C > 0$ and where λ will be the corresponding pressure to some stationary point. A priori, the condition only guarantees the existence of $\nabla \lambda(x) R$ in a suitable space. In the next lemma we show that λ and $\nabla \lambda$ themselves exist in a suitable space, which, in particular, allows one to make use of the technical lemma above.

Lemma 2.2. *Let $\mu : B \rightarrow \mathbb{R}$ be a function satisfying*

$$\|\nabla \mu(x) R\|_{L^\infty(B, \mathbb{R}^2)} < \infty.$$

Then $\mu \in W^{1,p}(B, \mathbb{R}^2)$ for any $1 \leq p < 2$.

Proof. The proof is straightforward. Indeed, it holds that

$$\int_B |\nabla \mu|^p \, dx \leq \|\nabla \mu R\|_{L^\infty(B, \mathbb{R}^2)}^p \int_B R^{1-p} \frac{dx}{R},$$

where the latter integrand is integrable for all $1 \leq p < 2$. □

We are now in a position to prove the main statement of this section.

Proof of Theorem 1.2:

Let $u \in \mathcal{A}^g$ be a stationary point with pressure λ , let $v \in \mathcal{A}^g$ be arbitrary and set $\eta := v - u \in W_0^{1,2}(B, \mathbb{R}^2)$.

We start expanding the energy via

$$E(v) = E(u) + E(\eta) + H(u, \eta),$$

where

$$H(u, \eta) := 2 \int_B M(x) \nabla u \cdot \nabla \eta \, dx$$

denotes the mixed terms.

Expanding the Jacobian of η and exploiting the fact that both u and v satisfy $\det \nabla u = \det \nabla v = 1$ a.e. yields

$$\det \nabla \eta = -\operatorname{cof} \nabla u \cdot \nabla \eta \quad \text{a.e.}$$

By the latter identity and the fact that (u, λ) satisfies (7), H can be written as

$$(12) \quad H(u, \eta) = 2 \int_B \lambda(x) \det \nabla \eta \, dx.$$

By Lemma 2.1.(vi) we have

$$H(u, \eta) = - \int_B (\operatorname{cof} \nabla \eta^{(0)} \nabla \lambda(x)) \cdot \tilde{\eta} \, dx - \int_B (\operatorname{cof} \nabla \eta \nabla \lambda(x)) \cdot \tilde{\eta} \, dx =: (I) + (II).$$

Now by noting that the 0–mode is only a function of R , we get

$$(\operatorname{cof} \nabla \eta^{(0)} \nabla \lambda(x)) \cdot \tilde{\eta} = \frac{\lambda, \theta}{R} (\eta_{1,R}^{(0)} \tilde{\eta}_2 - \eta_{2,R}^{(0)} \tilde{\eta}_1).$$

Instead of just λ, θ on the right hand side of the latter equation we would like to have the full gradient of λ . This can be achieved by using the basic relations $e_\theta \cdot e_\theta = 1$ and $e_R \cdot e_\theta = 0$ to obtain

$$(\operatorname{cof} \nabla \eta^{(0)} \nabla \lambda(x)) \cdot \tilde{\eta} = (\lambda, R R e_R + \lambda, \theta e_\theta) \cdot (\eta_{1,R}^{(0)} \tilde{\eta}_2 - \eta_{2,R}^{(0)} \tilde{\eta}_1) \frac{e_\theta}{R}.$$

Arguing similarly for (II), and a short computation shows

$$(13) \quad \begin{aligned} H(u, \eta) = & - \int_B (\lambda, R R e_R + \lambda, \theta e_\theta) \cdot \left[(\tilde{\eta}_1 \tilde{\eta}_{2, \theta} - \tilde{\eta}_2 \tilde{\eta}_{1, \theta}) \frac{e_R}{R} \right. \\ & \left. + (\tilde{\eta}_2 (\eta_{1,R}^{(0)} + \eta_{1,R}) - \tilde{\eta}_1 (\eta_{2,R}^{(0)} + \eta_{2,R})) e_\theta \right] \frac{dx}{R}. \end{aligned}$$

By Hölder's inequality we get⁵

$$\begin{aligned} H(u, \eta) \geq & - \|\nabla \lambda(x) R\|_{L^\infty(B, \mathbb{R}^2, \frac{dx}{R})} \int_B \left[|\tilde{\eta}_1 \tilde{\eta}_{2, \theta} - \tilde{\eta}_2 \tilde{\eta}_{1, \theta}| \frac{1}{R} \right. \\ & \left. + \left| \tilde{\eta}_2 (\eta_{1,R}^{(0)} + \eta_{1,R}) - \tilde{\eta}_1 (\eta_{2,R}^{(0)} + \eta_{2,R}) \right| \right] \frac{dx}{R}. \end{aligned}$$

By $\|\nabla \lambda(x) R\|_{L^\infty(\frac{dx}{R})} \leq \frac{\sqrt{3}\nu}{2\sqrt{2}}$ and a weighted Cauchy-Schwarz Inequality, we see

$$\begin{aligned} H(u, \eta) \geq & - \frac{\nu\sqrt{3}}{4\sqrt{2}} \left[2a \|\tilde{\eta}_1\|_{L^2(dx/R^2)}^2 + 2a \|\tilde{\eta}_2\|_{L^2(dx/R^2)}^2 \right. \\ & \left. + \frac{1}{a} \int_B \left[\frac{\tilde{\eta}_{2, \theta}^2}{R^2} + (\eta_{2,R}^{(0)} + \eta_{2,R})^2 + (\eta_{1,R}^{(0)} + \eta_{1,R})^2 + \frac{\tilde{\eta}_{1, \theta}^2}{R^2} \right] dx \right]. \end{aligned}$$

Next we recall an elementary Fourier estimate (see, for instance, [3, Proof of Proposition 3.3]), which states that for any $\phi \in C^\infty(B)$ it holds

$$(14) \quad \int_B R^{-2} |\tilde{\phi}, \theta|^2 dx \geq \int_B R^{-2} |\tilde{\phi}|^2 dx.$$

Applying the Cauchy-Schwarz inequality and (14), and then combining some of the norms yields

$$\begin{aligned} H(u, \eta) \geq & - \frac{\nu\sqrt{3}}{4\sqrt{2}} \left[\left(2a + \frac{1}{a}\right) \|\tilde{\eta}_{1, \theta}\|_{L^2(dx/R^2)}^2 + \left(2a + \frac{1}{a}\right) \|\tilde{\eta}_{2, \theta}\|_{L^2(dx/R^2)}^2 + \right. \\ & \left. + \frac{2}{a} \|\eta_{1,R}^{(0)}\|_{L^2(dx)}^2 + \frac{2}{a} \|\eta_{1,R}\|_{L^2(dx)}^2 + \frac{2}{a} \|\eta_{2,R}^{(0)}\|_{L^2(dx)}^2 + \frac{2}{a} \|\eta_{2,R}\|_{L^2(dx)}^2 \right] \end{aligned}$$

⁵Since we are not only interested in a qualitative but rather a quantitative estimate, we need to specify which norm we pick on \mathbb{R}^2 . For this the above Hölder estimate is given more carefully by

$$\int_B f \cdot g d\mu = \int_B f_1 g_1 + f_2 g_2 d\mu \leq \int_B |f_1| |g_1| + |f_2| |g_2| d\mu \leq \max\{\|f_1\|_{L^\infty(B, \mu)}, \|f_2\|_{L^\infty(B, \mu)}\} \int_B (|g_1| + |g_2|) d\mu$$

This is the reason why we defined the norm of f via $\|f\|_{L^\infty(B, \mathbb{R}^2, \mu)} := \max\{\|f_1\|_{L^\infty(B, \mu)}, \|f_2\|_{L^\infty(B, \mu)}\}$.

$$\geq -\frac{\nu\sqrt{3}}{4\sqrt{2}} \left[(2a + \frac{1}{a}) \|\tilde{\eta}_{,\theta}\|_{L^2(dx/R^2)}^2 + \frac{2}{a} \|\eta_{,R}^{(0)}\|_{L^2(dx)}^2 + \frac{2}{a} \|\eta_{,R}\|_{L^2(dx)}^2 \right].$$

Making use of $\tilde{\eta}_{,\theta} = \eta_{,\theta}$, which is true since the zero-mode does not depend on θ , and $\|\eta_{,R}^{(0)}\|_{L^2(dx)}^2 \leq \|\eta_{,R}\|_{L^2(dx)}^2$ we obtain

$$H(u, \eta) \geq -\frac{\nu\sqrt{3}}{4\sqrt{2}} \left[(2a + \frac{1}{a}) \|\eta_{,\theta}\|_{L^2(dx/R^2)}^2 + \frac{4}{a} \|\eta_{,R}\|_{L^2(dx)}^2 \right].$$

Choosing $a = \frac{\sqrt{3}}{\sqrt{2}}$ and again combining norms gives

$$\begin{aligned} H(u, \eta) &\geq -\frac{\nu\sqrt{3}}{4\sqrt{2}} \left[\frac{4\sqrt{2}}{\sqrt{3}} (\|\eta_{,\theta}\|_{L^2(dx/R^2)}^2 + \|\eta_{,R}\|_{L^2(dx)}^2) \right] \\ &= -\nu D(\eta), \end{aligned}$$

where $D(\eta) := \|\nabla\eta\|_{L^2(dx)}^2$ denotes the Dirichlet energy. This yields

$$E(\eta) + H(u, \eta) \geq E(\eta) - \nu D(\eta) \geq 0,$$

which, since $M\xi \cdot \xi \geq \nu|\xi|^2$ for all $\xi \in \mathbb{R}^2$, completes the proof.

The prefactor $\frac{\sqrt{3}}{2\sqrt{2}}$ in (9) is the best we have for general λ at the moment. If, however, circumstances are such that λ depends on only one of R, θ throughout B , then condition (9) can be replaced by the weaker assumption

$$(15) \quad \|\nabla\lambda(x)R\|_{L^\infty(B, \mathbb{R}^2)} \leq \nu.$$

Corollary 2.3. *Let the conditions of Theorem 1.2 be in force, but with (15) replacing (8), and assume that either $\lambda(x) = \lambda(R)$ or $\lambda(x) = \lambda(\theta)$ for all $x \in B$. Then the conclusions of Theorem 1.2 hold.*

Proof. (i) ($\lambda(x) = \lambda(R)$.) This case is significantly simpler and one can argue more along the lines of the proof of [3, Prop.3.3]. The reason is that in this case it still holds that

$$H(v, \eta) = 2 \int_B \lambda(R) \det \nabla\eta \, dx = 2 \int_B \lambda(R) \det \nabla\tilde{\eta} \, dx,$$

where $\tilde{\eta} = \eta - \eta^{(0)}$ eliminating the 0-mode.

Then applying of [3, Lemma 3.2.(iv)] yields

$$H(v, \eta) = \int_B \lambda'(R) R \tilde{\eta} \cdot J \tilde{\eta}_{,\theta} \frac{dx}{R^2}.$$

Using Hölder's inequality, $\|\lambda'(R)R\|_{L^\infty(0,1)} \leq \nu$, and Fourier estimate (14) we get

$$\begin{aligned} H(v, \eta) &\geq -\|\lambda'(R)R\|_{L^\infty(0,1)} \int_B |\tilde{\eta}| |\tilde{\eta}_{,\theta}| \frac{dx}{R^2} \\ &\geq -\nu \left(\int_B |\tilde{\eta}|^2 \frac{dx}{R^2} \right)^{1/2} \left(\int_B |\tilde{\eta}_{,\theta}|^2 \frac{dx}{R^2} \right)^{1/2} \end{aligned}$$

$$\begin{aligned} &\geq -\nu \int_B |\tilde{\eta}_{,\theta}|^2 \frac{dx}{R^2} \\ &\geq -\nu \int_B |\nabla \eta|^2 dx. \end{aligned}$$

Note, as before, that the \sim could be dropped because $\int_B |\nabla \tilde{\eta}|^2 dx \leq \int_B |\nabla \eta|^2 dx$.

(ii) ($\lambda(x) = \lambda(\theta)$.) Here we start with (13) which simplifies to

$$H(u, \eta) = - \int_B \lambda_{,\theta}(\theta) [(\eta_{2,R}^{(0)} + \eta_{2,R}) \tilde{\eta}_1 + (\eta_{1,R}^{(0)} + \eta_{1,R}) \tilde{\eta}_2] \frac{dx}{R}$$

By Hölder's inequality, Inequality (14) and $\|\lambda_{,\theta}\|_{L^\infty(0,2\pi)} \leq \nu$ we get

$$\begin{aligned} H(u, \eta) &\geq - \|\lambda_{,\theta}\|_{L^\infty(0,2\pi)} \int_B |(\eta_{2,R}^{(0)} + \eta_{2,R}) \tilde{\eta}_1 + (\eta_{1,R}^{(0)} + \eta_{1,R}) \tilde{\eta}_2| \frac{dx}{R} \\ &\geq - \frac{\nu}{2} [2\|\tilde{\eta}_{1,\theta}\|_{L^2(dx/R^2)}^2 + 2\|\tilde{\eta}_{2,\theta}\|_{L^2(dx/R^2)}^2 + \|\eta_{,R}^{(0)}\|_{L^2(dx)}^2 + \|\eta_{,R}\|_{L^2(dx)}^2] \end{aligned}$$

Using $\tilde{\eta}_{,\theta} = \eta_{,\theta}$ and $\|\eta_{,R}^{(0)}\|_{L^2(dx)}^2 \leq \|\eta_{,R}\|_{L^2(dx)}^2$ we get

$$\begin{aligned} H(u, \eta) &\geq - \frac{\nu}{2} [2\|\eta_{,\theta}\|_{L^2(dx/R^2)}^2 + 2\|\eta_{,R}\|_{L^2(dx)}^2] \\ &= -\nu D(\eta). \end{aligned}$$

□

Remark 2.4 (Relaxation of the assumptions). *The result of Theorem 1.2 continues to hold if we assume that $f(x; \xi) = M(x)\xi \cdot \xi \geq \nu(|x|)|\xi|^2$ for some $\nu \in L^\infty(\mathbb{R}_+)$, $\nu(R) \geq 0$, and all $\xi \in \mathbb{R}^{2 \times 2}$. Here, $\nu(R) = 0$ is allowed⁶ to be 0. The assumption $\nu = \nu(R)$ is needed because we do not know if the Fourier estimate (14) is still true if ν depends on both R and θ . In this case, the small pressure condition can be relaxed to a pointwise estimate:*

$$|\nabla \lambda(x)R| \leq \frac{\sqrt{3}\nu(R)}{2\sqrt{2}} \quad \text{for a.e. } x \in B,$$

with uniqueness if the inequality is strict on some non-null set.

3. A METHOD FOR COMPUTING THE PRESSURE AND A COUNTEREXAMPLE TO REGULARITY

In this section we construct an explicit functional $E(u)$ of the form (1), where the integrand obeys (2) and (3), such that $u = u_N$ is the global minimizer of E in $\mathcal{A}^{\text{tr}}(u_N)$. The strategy is as follows:

- (i) select a candidate trace function $g \in W^{k+1,p}(\mathbb{S}^1, \mathbb{R}^2)$ for $k \geq 1$ and $1 \leq p \leq \infty$ which obeys⁷ $1 = Jg(\theta) \cdot g'(\theta)$ for a.e. θ in $[0, 2\pi]$ ⁸

⁶Note that in these circumstances, we do not need to verify that a minimizer of the associated functional $E(u)$ exists in order to apply Theorem 1.2. Rather, it is enough to establish that u is a stationary point in the sense of (7).

⁷The condition $Jg \cdot g' = 1$ ensures that the one-homogeneous extension u obeys $\det \nabla u = 1$ a.e. in B

⁸Here, by a slight abuse of notation, we put $\tilde{g}(\theta) = g(\cos \theta, \sin \theta)$ and then promptly drop the $\tilde{}$

- (ii) extend g to a one-homogeneous function $u(R, \theta) := Rg(\theta)$, and compute, in Lemma 3.1, a PDE which must be satisfied by both u and an associated λ in order that u is a stationary point of E in the sense of (7);
- (iii) fix $g = \text{tr}(u_N)$ and construct, in Lemma 3.2, a suitable $f(x, \xi)$ such that the PDE in step (ii) can be solved for λ , and
- (iv) verify that the small pressure condition stipulated in Corollary 2.3 is satisfied by λ , and hence that u_N is the unique global minimizer of the associated energy E .

We first examine conditions on M necessary for a to be a stationary point of E in the sense of (7).

Notation: Recall the notation for 2d polar coordinates

$$\{e_R, e_\theta\} := \{(\cos \theta, \sin \theta), (-\sin \theta, \cos \theta)\}.$$

Additionally, we will use

$$\{e_{NR}, e_{N\theta}\} := \{(\cos(N\theta), \sin(N\theta)), (-\sin(N\theta), \cos(N\theta))\}$$

for any $N \in \mathbb{N}$. Moreover, we will use the notation $M_{ijgk} = (M(e_i \otimes e_j)) \cdot (g \otimes e_k)$ for any combination of $i, j, k \in \{R, \theta\}$ and any map $g \in \mathbb{R}^2$. Especially, if $g = e_{Nl}$ for some $l \in \{R, \theta\}$ we will use $M_{ij(Nl)k}$ for short.

Lemma 3.1 (Representation of the pressure). *Let $1 \leq p \leq \infty$, $k \in \mathbb{N} \setminus \{0\}$ and assume $M \in (L^\infty \cap W^{k,p})(B, \mathbb{R}^{16})$, $g \in W^{k+1,p}(\mathbb{S}^1, \mathbb{R}^2)$ where g obeys $Jg \cdot g' = 1$ a.e. in $[0, 2\pi)$, and let $u = Rg(\theta) \in \mathcal{A}^g$ be a stationary point of the energy E as defined in (7).*

Then there exists a corresponding pressure $\lambda \in W^{k,p}(B, \mathbb{R})$ and it satisfies the following system of equations a.e. in B :

$$\begin{aligned}
\lambda(x)_{,\theta} (Jg \cdot e_R) - \lambda(x)_{,R} R(Jg' \cdot e_R) &= - [M_{R\theta(g+g'')\theta} \\
&\quad + ((M_{,\theta})_{R\theta gR} + (M_{,\theta})_{R\theta g'\theta}) \\
&\quad + R((M_{,R})_{RRgR} + (M_{,R})_{RRg'\theta})] \\
&=: h_1(M, g) \\
\lambda(x)_{,\theta} (Jg \cdot e_\theta) - \lambda(x)_{,R} R(Jg' \cdot e_\theta) &= - [M_{\theta\theta(g+g'')\theta} \\
&\quad + ((M_{,\theta})_{\theta\theta gR} + (M_{,\theta})_{\theta\theta g'\theta}) \\
&\quad + R((M_{,R})_{\theta RgR} + (M_{,R})_{\theta Rg'\theta})] \\
(16) \qquad \qquad \qquad &=: h_2(M, g)
\end{aligned}$$

Proof. Let $u = Rg(\theta) \in \mathcal{A}^g$ be a stationary point. If there exists a corresponding pressure $\lambda \in W^{1,p}$ then u is a solution of

$$(17) \quad \int_B M(x) \nabla u \cdot \nabla \eta \, dx = - \int_B \lambda(x) \text{cof} \nabla u \cdot \nabla \eta \, dx \quad \text{for any } \eta \in C_c^\infty(B, \mathbb{R}^2).$$

For now, let us assume that $\lambda \in W^{1,p}(B, \mathbb{R})$. In order to derive the system of equations above, we enter the explicit form of u and the representation $\eta = (\eta \cdot e_R)e_R + (\eta \cdot e_\theta)e_\theta$ into the stationarity condition. By some further calculations, which are mainly integrations by parts, we obtain (16). In the last step of the proof we discuss the existence of $\lambda \in W^{1,p}$.

Step 1: Computation of left-hand side of (17):

The derivative and the cofactor of the map $u = Rg(\theta) + b$ are given by

$$\begin{aligned}\nabla u &= g \otimes e_R + g' \otimes e_\theta, \\ \text{cof } \nabla u &= Jg \otimes e_\theta - Jg' \otimes e_R.\end{aligned}$$

Plugging the above into the left-hand side of (17) and integrating by parts yields

$$\begin{aligned}(\text{LHS}) &= \int_B M(x)(g \otimes e_R + g' \otimes e_\theta) \cdot \left(\eta_{,R} \otimes e_R + \frac{1}{R} \eta_{,\theta} \otimes e_\theta \right) dx \\ &= - \int_B R(M(x),_R)(g \otimes e_R + g' \otimes e_\theta) \cdot (\eta \otimes e_R) \frac{dx}{R} \\ &\quad - \int_B [M(x)((g + g'') \otimes e_\theta) + M(x),_\theta (g \otimes e_R + g' \otimes e_\theta)] \cdot (\eta \otimes e_\theta) \frac{dx}{R}\end{aligned}$$

Now by expanding $\eta = \alpha e_R + \beta e_\theta$ with $\alpha = (\eta \cdot e_R)$ and $\beta = (\eta \cdot e_\theta)$ and the shorthand introduced above we get

$$\begin{aligned}(\text{LHS}) &= - \int_B R[(M,_R)_{RRgR} + (M,_R)_{RRg'\theta}] \alpha + R[(M,_R)_{\theta RgR} + (M,_R)_{\theta Rg'\theta}] \beta \frac{dx}{R} \\ &\quad - \int_B M_{R\theta(g+g'')\theta} \alpha + [(M,_\theta)_{R\theta gR} + (M,_\theta)_{R\theta g'\theta}] \alpha \\ &\quad + M_{\theta\theta(g+g'')\theta} \beta + [(M,_\theta)_{\theta\theta gR} + (M,_\theta)_{\theta\theta g'\theta}] \beta \frac{dx}{R} \\ &= \int_B h_1 \alpha + h_2 \beta \frac{dx}{R}.\end{aligned}$$

Step 2: Computation of right-hand side of (17):

Now by again using the explicit form of $\text{cof } \nabla u$, and integration by parts we get

$$\begin{aligned}(\text{RHS}) &= - \int_B (\lambda(x)(Jg \cdot \eta_{,\theta}) - \lambda(x)R(Jg' \cdot \eta_{,R})) \frac{dx}{R} \\ &= \int_B \lambda_{,\theta}(x)(Jg \cdot \eta) - \lambda_{,R}(x)R(Jg' \cdot \eta) \frac{dx}{R}.\end{aligned}$$

Further, we use the expression $\eta = \alpha e_R + \beta e_\theta$ with the notation $\alpha = (\eta \cdot e_R)$ and $\beta = (\eta \cdot e_\theta)$ to derive

$$\begin{aligned}(\text{RHS}) &= \int_B (\lambda_{,\theta}(x)(Jg \cdot e_R) - \lambda_{,R}(x)R(Jg' \cdot e_R)) \alpha \\ &\quad + (\lambda_{,\theta}(x)(Jg \cdot e_\theta) - \lambda_{,R}(x)R(Jg' \cdot e_\theta)) \beta \frac{dx}{R}.\end{aligned}$$

Together with Step 1 and the realization that in the above $\alpha, \beta \in C_c^\infty(B)$ are arbitrary, the claimed equations need to be true a.e. in B .

Step 3: Existence of the pressure $\lambda \in W^{1,p}(B, \mathbb{R})$:

The equations above can be rewritten as

$$\int_B \operatorname{div} (\lambda \operatorname{cof} \nabla u) \cdot \eta \, dx = \int_B h(M, g) \cdot \eta \frac{dx}{R},$$

where $h = (h_1, h_2)$. We know that $\operatorname{div} (\lambda \operatorname{cof} \nabla u) \in L^p(dx)$ iff $h(M, g) \in L^p(\frac{dx}{R})$, with obvious notation. Now consider $h_1(M, g)$ (the argument being similar for h_2) and define

$$\begin{aligned} h_{11} &:= -M_{R\theta(g+g'')\theta} \\ h_{12} &:= [((M, \theta)_{R\theta g R} + (M, \theta)_{R\theta g' \theta}) \\ &\quad + R((M, R)_{RRgR} + (M, R)_{RRg' \theta})]. \end{aligned}$$

Then for $h_{11} \in L^p(\frac{dx}{R})$ we need $M \in L^\infty(dx)$ and $g, g'' \in L^p$, which is true by assumption. Now, by Sobolev imbedding, we have $W^{2,p} \hookrightarrow W^{1,\infty}([0, 2\pi], \mathbb{R}^2)$, and hence, in order for $h_{12} \in L^p(\frac{dx}{R})$, and bearing in mind that $g, g' \in L^\infty$, it is enough to require that $\nabla M \in L^p(dx)$. This is exactly how we chose the classes for M and g . This guarantees the existence of $\operatorname{div} (\lambda \operatorname{cof} \nabla u) = (\operatorname{cof} \nabla u) \nabla \lambda \in L^p(dx)$. By further noting that $g, g' \in L^\infty$, it is immediate that $\nabla u \in L^\infty(dx)$, and since $\det \nabla u = 1$ a.e. in B , we may write

$$\nabla \lambda = (\operatorname{cof} \nabla u)^T \frac{h(M, g)}{R} \in L^p(dx).$$

In particular, when M and g are specified, $\nabla \lambda$ is specified and it belongs to the class $L^p(B)$, reverting to the traditional notation. One can argue similarly for the higher integrability. \square

We now specify $g = \operatorname{tr}(u_N)$ and compute the pressure under the assumptions that M depends only on θ and is diagonal with respect to the basis of polar coordinates.

Lemma 3.2 (Representation of the pressure, N-cover, $M(\theta)=\operatorname{diag}$). *For $N \in \mathbb{N} \setminus \{1\}$ let $g = \frac{1}{\sqrt{N}} e_{NR}$ and assume $M \in (L^\infty \cap W^{k,p})(B, \mathbb{R}^{16})$ for some $1 \leq p \leq \infty$ and $k \in \mathbb{N}$, where M is of the specific form*

$$M(x) = \operatorname{diag}(M_{RRRR}, M_{R\theta R\theta}, M_{\theta R\theta R}, M_{\theta\theta\theta\theta}) = \operatorname{diag}(\alpha(\theta), \beta(\theta), \gamma(\theta), \delta(\theta))$$

with $\nu > 0$ and $\alpha, \beta, \gamma, \delta \geq \nu$ for any $\theta \in [0, 2\pi)$. Furthermore, suppose $u = Rg(\theta) \in \mathcal{A}^g$ is a stationary point of the energy E , as defined in (1).

Then there exists a corresponding pressure $\lambda \in W^{k,p}(B)$ and it satisfies the following system of equations a.e. in B :

$$\begin{aligned} (18) \quad -\lambda(x)_{,\theta} \frac{1}{\sqrt{N}} \sin(\theta_{N-1}) + \lambda(x)_{,R} R\sqrt{N} \cos(\theta_{N-1}) &= \sqrt{N} \beta' \sin(\theta_{N-1}) \\ &\quad + \left[\sqrt{N}(N-1)\beta + \sqrt{N}\delta - \frac{\alpha}{\sqrt{N}} \right] \cos(\theta_{N-1}) \\ &=: h_1 \end{aligned}$$

$$\begin{aligned} (19) \quad \lambda(x)_{,\theta} \frac{1}{\sqrt{N}} \cos(\theta_{N-1}) + \lambda(x)_{,R} R\sqrt{N} \sin(\theta_{N-1}) &= -\sqrt{N} \delta' \cos(\theta_{N-1}) \\ &\quad + \left[\sqrt{N}\beta + \sqrt{N}(N-1)\delta - \frac{\gamma}{\sqrt{N}} \right] \sin(\theta_{N-1}) \\ &=: h_2 \end{aligned}$$

where we used the shorthand $\theta_k := k\theta$ for any $k \in \mathbb{R}$.

Proof. By Lemma 3.1, we know that the pressure λ exists and system (16) is satisfied. Now we just have to verify that (16) agrees with the claimed system given by (18) and (19). We start by verifying the first of the equations in the system (16).

Step 1:

That the left-hand side of (18) follows from the left-hand side of the first equation in system (16) is a straightforward calculation. Hence, we focus on the corresponding right-hand side, which we named h_1 . First note that, because M depends only on θ , we are left with

$$(20) \quad h_1 = - [M_{R\theta(g+g'')\theta} + (M, \theta)_{R\theta gR} + (M, \theta)_{R\theta g'\theta}].$$

We have $g = \frac{1}{\sqrt{N}}e_{NR}$, $g' = \sqrt{N}e_{N\theta}$, $g'' = -N\sqrt{N}e_{NR}$, and hence

$$M_{R\theta(g+g'')\theta} = \left(\frac{1}{\sqrt{N}} - \sqrt{N}N \right) M_{R\theta(NR)\theta} = \left(\frac{1}{\sqrt{N}} - \sqrt{N}N \right) (M_{R\theta R\theta}(e_{NR} \cdot e_R) + M_{R\theta\theta\theta}(e_{NR} \cdot e_\theta)).$$

Using that $M_{R\theta\theta\theta} = 0$ and $M_{R\theta R\theta} = \beta$ yields

$$M_{R\theta(g+g'')\theta} = \left(\frac{1}{\sqrt{N}} - \sqrt{N}N \right) \beta \cos(\theta_{N-1}).$$

For the second term of (20), consider

$$M_{R\theta gR, \theta} = (M, \theta)_{R\theta gR} + M_{\theta\theta gR} - M_{RRgR} + M_{R\theta g'R} + M_{R\theta g\theta},$$

which, after a short calculation, gives

$$\begin{aligned} (M, \theta)_{R\theta gR} &= M_{R\theta gR, \theta} - M_{\theta\theta gR} + M_{RRgR} - M_{R\theta g'R} - M_{R\theta g\theta} \\ &= \frac{1}{\sqrt{N}} [\alpha - \beta] \cos(\theta_{N-1}). \end{aligned}$$

Similarly, for the rightmost term of (20) we get

$$\begin{aligned} (M, \theta)_{R\theta g'\theta} &= M_{R\theta g'\theta, \theta} - M_{\theta\theta g'\theta} + M_{RRg'\theta} - M_{R\theta g''\theta} + M_{R\theta g'R} \\ &= -\sqrt{N}\beta' \sin(\theta_{N-1}) + \sqrt{N}[\beta - \delta] \cos(\theta_{N-1}). \end{aligned}$$

Together,

$$h_1 = \sqrt{N}\beta' \sin(\theta_{N-1}) + \left[\sqrt{N}\delta - \frac{\alpha}{\sqrt{N}} + \sqrt{N}(N-1)\beta \right] \cos(\theta_{N-1}).$$

Step 2:

By arguing similarly, we find that

$$h_2 = - [M_{\theta\theta(g+g'')\theta} + (M, \theta)_{\theta\theta gR} + (M, \theta)_{\theta\theta g'\theta}].$$

Then

$$\begin{aligned} M_{\theta\theta(g+g'')\theta} &= \left(\frac{1}{\sqrt{N}} - \sqrt{N}N \right) \delta \sin(\theta_{N-1}) \\ (M, \theta)_{\theta\theta gR} &= M_{\theta\theta gR, \theta} + M_{R\theta gR} + M_{\theta RgR} - M_{\theta\theta g'R} - M_{\theta\theta g\theta} \\ &= \frac{1}{\sqrt{N}} [\gamma - \delta] \sin(\theta_{N-1}) \\ (M, \theta)_{\theta\theta g'\theta} &= M_{\theta\theta g'\theta, \theta} + M_{R\theta g'\theta} + M_{\theta Rg'\theta} - M_{\theta\theta g''\theta} + M_{\theta\theta g'R} \end{aligned}$$

$$=\sqrt{N}\delta' \cos(\theta_{N-1}) + \sqrt{N}[\delta - \beta] \sin(\theta_{N-1})$$

and finally

$$h_2 = -\sqrt{N}\delta' \cos(\theta_{N-1}) + \left[\sqrt{N}\beta + \sqrt{N}(N-1)\delta - \frac{\gamma}{\sqrt{N}} \right] \sin(\theta_{N-1}),$$

completing the proof. \square

Next, we compute the small pressure criteria in the same situation. Moreover, we will provide an explicit form of the pressure.

Lemma 3.3 (Small pressure condition, N-cover, $M(\theta)=\text{diag}$). *Let the assumptions be as above. For any $N \in \mathbb{N} \setminus \{1\}$, let $M = (a, 1, a, 1)\nu$, where we pick a to be constant and in the range*

$$1 \leq N^2 - N < a < N^2 + N.$$

Then for this M the corresponding pressure λ is given by

$$\lambda(x) = c + \left[N - \frac{a}{N} \right] \ln(R) \text{ for any } x \in B$$

for any real constant $c \in \mathbb{R}$, which is independent of R and θ . Moreover, $\lambda \in W^{1,q}(B)$ for any $1 \leq q < 2$ and λ satisfies condition (9) strictly.

Proof. Define first

$$H_1 = \left[\sqrt{N}(N-1)\beta + \sqrt{N}\delta - \frac{\alpha}{\sqrt{N}} \right] \text{ and } H_2 = \left[\sqrt{N}\beta + \sqrt{N}(N-1)\delta - \frac{\gamma}{\sqrt{N}} \right].$$

By solving the system (18) and (19) we obtain

$$\begin{aligned} \lambda_{,R} R &= (\beta' - \delta') \frac{\sin(2\theta_{N-1})}{2} + \frac{1}{\sqrt{N}} (H_1 \cos^2(\theta_{N-1}) + H_2 \sin^2(\theta_{N-1})) \\ \lambda_{,\theta} &= \sqrt{N} (H_2 - H_1) \frac{\sin(2\theta_{N-1})}{2} - N (\beta' \sin^2(\theta_{N-1}) + \delta' \cos^2(\theta_{N-1})). \end{aligned}$$

For the specific case of $M = (a, 1, a, 1)\nu$ they become

$$\lambda_{,R} R = \left[N - \frac{a}{N} \right] \text{ and } \lambda_{,\theta} = 0$$

showing, in particular, that λ depends only on R i.e. $\lambda(x) = \lambda(R)$. Indeed, the pressure is then given by

$$\lambda(x) = c + \left[N - \frac{a}{N} \right] \ln(R) \text{ for any } x \in B$$

and for any real constant $c \in \mathbb{R}$, which is independent of R and θ . The small pressure condition of Corollary 2.3 can now be applied, giving

$$\left| N - \frac{a}{N} \right| < 1.$$

Solving this inequality by case distinction yields the claimed bounds on a . The integrability is then easily deduced, completing the proof. \square

To make it more accessible for the reader we collect what we have shown so far in the following.

Proof of Theorem 1.3

(i), (ii), and (vi) trivial.

- (iii) It is enough to show this point for M . Note, that M only depends on θ , i.e. $M(x) = M(\theta)$. Hence, the gradient is given by

$$(21) \quad \nabla M = \frac{1}{R} \partial_\theta M(\theta) \otimes e_\theta \text{ for any } x \in B \setminus \{0\}.$$

First realise that the derivative with respect to θ only replaces e_R with e_θ (up to sign) and vice versa, and therefore one can still bound the modulus of $\|\partial_\theta M(\theta)\|_{L^\infty(B, \mathbb{R}^{16})} \leq C(a)$ via some real constant $C(a) > 0$. Then, by integrating $|\nabla M|^q$ with respect to dx using (21) and by the latter discussion, the claim follows.

- (iv) As a consequence of $g \in C^\infty$ and point (iii), Lemma 3.2 guarantees that u is a stationary point and the existence of λ in the right spaces.
- (v) By Lemma 3.3 we know that λ satisfies the small pressure criteria strictly. Together, with Theorem 1.2 this implies that $u = u_N$ is indeed the unique global minimizer to the energy E .

Remark 3.4. (i) To summarize the foregoing analysis, we have shown that for the full ball $B \subset \mathbb{R}^2$ and smooth boundary conditions, albeit with a topological change, there is a uniformly convex functional, which depends discontinuously on x , but smoothly on ∇u , such that the corresponding energy is uniquely globally minimised by a map that is everywhere Lipschitz but not $C^1(B)$.

(ii) One might also be interested in this counterexample on the scale of Sobolev spaces. With this in mind, note that for any $N \in \mathbb{N} \setminus \{1\}$ and $b \in \mathbb{R}^2$ we have⁹

$$u = \frac{R}{\sqrt{N}} e_{NR} + b \in W^{2,q}(B, \mathbb{R}^2) \setminus W^{2,2}(B, \mathbb{R}^2) \text{ for any } 1 \leq q < 2.$$

Moreover, one might like to compare our result with the high-order regularity result, given in [2]. They showed that for the special case of the Dirichlet functional and $u \in W^{2,q}(B, \mathbb{R}^2)$ with $q > 2$ being a stationary point satisfying $\det \nabla u = 1$ a.e., then $u \in C^\infty(B, \mathbb{R}^2)$. It is possible that a similar result could be established, for a fairly general non-autonomous p -growth functional with the necessary changes in q . Let's assume for a second that such a result is indeed possible. Intriguingly, this seems to leave a 'gap' at $q = 2$.

(iii) The singular set Σ in our example is, of course, just the origin $\Sigma = \{0\}$. It remains an open question whether there can be other incompressible variational problems, including in incompressible elasticity, where the stationary points/minimizers possess a richer Σ .

REFERENCES

- [1] John M. Ball. *Some Open Problems in Elasticity*, chapter I.1, pages 3–59. Springer-Verlag, NY, 2002.
- [2] P. Bauman, N. C. Owen, and D. Phillips. Maximum Principles and a priori estimates for an incompressible material in nonlinear elasticity. *Communications in Partial Differential Equations*, 17(7):1185–1212, 1992.
- [3] Jonathan J. Bevan. On double-covering stationary points of a constrained Dirichlet energy. *Annales de l'Institut Henri Poincaré (C) Non Linear Analysis*, 31(2):391–411, 2014.
- [4] Jonathan J. Bevan and Jonathan H. B. Deane. A continuously perturbed Dirichlet energy with area-preserving stationary points that 'buckle' and occur in equal-energy pairs. *Non. Dif. Eq. App.*, 28(1), dec 2020.

⁹One might compare this to recent results given in [13,14]. They obtain higher integrability, in particular, in the same spaces as we do, for counterexamples in compressible elasticity. Clearly their result is extraordinarily difficult, since the counterexamples are produced by means of convex integration.

- [5] Jonathan J. Bevan and Jonathan H. B. Deane. Energy minimizing N -covering maps in two dimensions. *Calc. Var. Par. Dif. Eq.*, 60(4), 2021.
- [6] L.C. Evans and R.F. Gariepy. On the partial regularity of energy-minimizing, area-preserving maps. *Calculus of Variations and Partial Differential Equations*, 9(4):357–372, dec 1999.
- [7] F. John. Uniqueness of non-linear elastic equilibrium for prescribed boundary displacements and sufficiently small strains. *Commun. Pure Appl. Math.*, 25:617–634., 1972.
- [8] Aram L. Karakhanyan. Sufficient conditions for regularity of area-preserving deformations. *Manuscripta Mathematica*, 138(3-4):463–476, 2011.
- [9] Aram L. Karakhanyan. Regularity for energy-minimizing area-preserving deformations. *Journal of Elasticity*, 114(2):213–223, 2013.
- [10] R. J. Knops and C. A. Stuart. Quasiconvexity and uniqueness of equilibrium solutions in nonlinear elasticity. *Arch. Rat. Mech. Anal.*, 86(3):233–249, 1984.
- [11] C. Morris and A. Taheri. Twist maps as energy minimisers in homotopy classes: Symmetrisation and the coarea formula. *Nonlinear Analysis: Theory, Methods & Applications*, 152:250–275, 2017.
- [12] K. D. E. Post and J. Sivaloganathan. On homotopy conditions and the existence of multiple equilibria in finite elasticity. *Proc. R. Soc. Ed. Sect. A*, 127(3):595–614, 1997.
- [13] A. Rüländ, C. Zillinger, and B. Zwirnagl. Higher Sobolev Regularity of Convex Integration Solutions in Elasticity: The Dirichlet Problem with Affine Data in $\text{int}(K^{lc})$. *SIAM J. Math. Anal.*, 50(4):3791–3841, 2018.
- [14] A. Rüländ, C. Zillinger, and B. Zwirnagl. Higher Sobolev Regularity of Convex Integration Solutions in Elasticity: The Planar Geometrically Linearized Hexagonal-to-Rhombic Phase Transformation. *Journal of Elasticity*, 138:1–76, 2020.
- [15] S. Shahrokhi-Dehkordi and A. Taheri. Quasiconvexity and Uniqueness of Stationary Points on a Space of Measure Preserving Maps. *Journal of Convex Analysis*, 17:69–79, 2010.
- [16] Jeyabal Sivaloganathan. Uniqueness of Regular and Singular Equilibria for Spherically Symmetric Problems of Nonlinear Elasticity. *Arch. Rat. Mech. Anal.*, 96:97–136., 1986.
- [17] Jeyabal Sivaloganathan and Scott J. Spector. On the Uniqueness of Energy Minimizers in Finite Elasticity. *J. Elas.*, 133(1):73–103, feb 2018.
- [18] E. N. Spadaro. Non-Uniqueness of Minimizers for Strictly Polyconvex Functionals. *Arch. Rat. Mech. Anal.*, 193(3):659–678, 2008.
- [19] Vladimír Šverák. Regularity properties of deformations with finite energy. *Archive for Rational Mechanics and Analysis*, 100(2):105–127, jun 1988.
- [20] Ali Taheri. Minimizing the Dirichlet Energy over a space of measure preserving maps. *Topological Methods in Nonlinear Analysis Journal of the Juliusz Schauder Center*, 33:170–204., 2009.
- [21] Ali Taheri and Charles Morris. On the Uniqueness and Monotonicity of Energy Minimisers in the Homotopy Classes of Incompressible Mappings and Related Problems. *Journal of Mathematical Analysis and Applications*, 473(18):1–26, 2019.
- [22] Kewei Zhang. Energy minimizers in nonlinear elastostatics and the implicit function theorem. *Archive for Rational Mechanics and Analysis*, 114(2):95–117, 1991.

ACKNOWLEDGEMENTS: The first author is appreciative to the Department of Mathematics at the University of Surrey and was funded by the Engineering & Physical Sciences Research Council (EPSRC).

(M. Dengler) FLIEDERWEG 1, 72189 VÖHRINGEN, GERMANY, (CORRESPONDING AUTHOR).
Email address: marci.dengler@web.de

(J. Bevan) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SURREY, GUILDFORD, GU2 7XH, UNITED KINGDOM.
Email address: j.bevan@surrey.ac.uk