

## Modifications of intrinsic foot joint kinetics after total ankle replacement

Alexandre Naaim, Raphaël Dumas, Jean-Luc Besse, Laurence Cheze, Bernhard Devos Bevernage, Thibaut Leemrijse, Paul-André Deleu

### ▶ To cite this version:

Alexandre Naaim, Raphaël Dumas, Jean-Luc Besse, Laurence Cheze, Bernhard Devos Bevernage, et al.. Modifications of intrinsic foot joint kinetics after total ankle replacement. SB 2022, 47eme Congrès de la Société de Biomécanique, Oct 2022, Monastir, Tunisia. pp S223-S225. hal-03862369

## HAL Id: hal-03862369 https://hal.science/hal-03862369

Submitted on 21 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Modifications of intrinsic foot joint kinetics after total ankle replacement

A. Naaïm<sup>a</sup>, R. Dumas<sup>a</sup>, J. L. Besse<sup>a,b</sup>, L. Chèze<sup>a</sup>, B. Devos Bevernage<sup>c</sup>, T. Leemrijse<sup>c</sup> and P. A. Deleu<sup>c</sup>

<sup>a</sup>Univ Lyon, Univ Gustave Eiffel, Univ Claude Bernard Lyon 1, LBMC UMR\_T 9406, Lyon, France; <sup>b</sup>Hospices Civils de Lyon, Centre Hospitalier Lyon-Sud, Service de Chirurgie Orthopédique et Traumatologique, Pierre-Bénite Cédex, France; <sup>c</sup>Foot & Ankle Institute, Brussels, Belgium

#### 1. Introduction

Osteoarthritis (OA) is a disabling pathology that can affect numerous joints but has been less studied for the ankle than in any other joint (Arnolds et al. 2022). Ankle OA has been proved to impact biomechanically both the lower limb and the entire foot (Queen 2017). While conservative treatment can manage the early stage of OA, end-stage may require surgical management. The gold standard has been for a long time the ankle arthrodesis; however, the Total Ankle Replacement (TAR) has been considered lately a better solution. Indeed, TAR might maintain the pre-operative range of motion (ROM) and allows to protect the adjacent joints. Changes in foot joint kinematics after TAR have been assessed in previous studies (Brodsky et al. 2013; Deleu et al. 2022). However, changes in foot joint kinetics have not yet been investigated. Therefore, this prospective study aims at comparing the kinematics and kinetics of the intrinsic foot joints between pre- and post-operative conditions in TAR patients, as well as between patients and a control group. The main hypothesis is that the kinetics of the considered foot joints during gait will have a more typical profile after TAR.

#### 2. Methods

#### 2.1. Participants

15 Participants (Age:  $59 \pm 10$  years, BMI:  $26 \pm 4$  kg/m<sup>2</sup>) suffering from end-stage ankle OA scheduled for primary TAR were recruited in this prospective study. Inclusion criteria were: (1) diagnosis of post traumatic end stage OA (2) age >18 years (3) no history of neurological or systemic diseases (4) being able to walk 100m with no aid. Patients participated to two experimental sessions, the first within 15 days before TAR surgery and one year after surgery. These participants were peer-matched to 15 asymptomatic participants (CTRL group) (Age:  $57 \pm 9$  years, BMI:  $25 \pm 4$  kg/m<sup>2</sup>) according to demographics and post-operative walking speed. All participants signed an informed consent, and this study was approved by the local ethical committee.

#### 2.2. Data collection

To obtain the foot kinematics and kinetics, participants were asked to walk barefoot over a 10 m walkway at a self-selected speed. The walkway was equipped with 10 optoelectronic cameras, a force platform and a pressure platform, all synchronised at 200 Hz. The trajectories of 16 reflective markers were recorded. These markers were positioned on the foot and the shank by a senior clinician according to the multi-segment Rizzoli Foot model (Leardini et al. 2007). Five walking trials were collected for each participant.

#### 2.3. Data analysis

Data analysis was identical with that described by Deleu et al. (2021). Joint kinematics and kinetics were calculated according to the adaptation of the Rizzoli multi-segment foot model proposed by Deschamps et al. (2017) following the ISB recommendations. Joint moments were calculated in the joint coordinate system using a bottom-up Newton-Euler inverse dynamic method. All data necessary for the kinematics and kinetics were filtered using a low-pass zero-lag, 4th order, Butterworth filter, with a cut-off frequency of 10 Hz. Inertia and weight parameters of each foot segment were not considered as the inertia effects were supposed to be negligible during stance in comparison to the external forces. Joint centres location were the one proposed by Deschamps et al (2017) in their adaptation of the model. All calculations were done using an ad-hoc Matlab program. To distribute the force plate data (centre of pressure, forces, and moments) over each foot segment, a validated proportionality scheme was used (Sarasawat et al. 2014). The main characteristic for ankle OA is the alteration of the sagittal plane ankle mechanics. Therefore, our study outcomes were the joint angles and moments about the flexion-extension axis for the Shank-Calcaneus (Sha-Cal), Chopart, Lisfranc and the 1st Metatarso-Phalangeal (MTP1) joint. However, the full power was computed and not only its 'component' about the flexion-extension axes which might be found in other studies but may seem incorrect mechanically. To compare the results to Minimal Detectable Change (MDC) obtained by Eerdekens et al. (2019), the mean difference (MD) between pre- and post-surgery for the sagittal ROM, the peak plantarflexion moment and the peak power generation for the Sha-Cal joint was calculated.

#### 2.4. Statistical analysis

Inter-segmental foot kinematics and kinetics were compared using one-dimensional Statistical Parametric



**Figure 1.** Joint angles, moments and power waveforms representation with  $\pm 1$  standard deviation for the Sha-Cal, Chopart, Lisfranc, and MTP1.

Mapping (Pataky et al. 2016). For comparing the preand post-operative conditions, a paired t-test was used, and to compare the CTRL group to the pre- and postoperative conditions a two-sample t-tests was used. An adjusted *P*-value (0.05/3 = 0.017) was used to control the type 1 error rate.

#### 3. Results and discussion

Walking speed increased significantly after surgery (1.05 m/s pre-operatively; 1.21 m/s post-operatively) being comparable with that of CTRL group (1.22 m/s).

The MD between the pre-operative and postoperative was  $2.5 \pm 4.2^{\circ}$  (MDC:  $2.2^{\circ}$ ) for the Sha-Cal joint sagittal ROM, 0.23 ± 0.3 Nm/kg (MDC: 0.06 Nm/ kg) for the plantar flexion moment and  $0.82 \pm 0.57$ Watt/kg (MDC: 0.39 Watt/kg) for peak power. Consequently, all the MDs found are above the MDC measured by Eerdekens et al. (2019). As seen on Figure 1, and in accordance with the literature (Brodsky et al. 2011), the pre-surgery ROM was maintained, and the walking speed of the patient was increased post-surgery which are both signs of a successful TAR. No significant difference was found between the control and the post-surgery group except for the Sha-Cal joint angle during pre-swing phase. Indeed, even if the moment in the different joints has increased after surgery, they seem normalized relative to the CTRL group. One of the explanations could be that patients were adopting an antalgic walking strategy pre-operatively and that no long-term compensation has been developed. This could be confirmed by the fact that the main difference between the pre-surgery participants and the CTRL group can be found on sagittal moment during the terminal stance, i.e. at push off (50 to 90% of the stance phase). As a result, it seems that the patients might have only reduced their walking speed as a pain coping mechanism to reduce moment in the foot joints.

#### 4. Conclusions

This study is the first to quantify changes in foot joint kinematics and kinetics in patients undergoing a TAR and compare the results to a matched CTRL group using a multi-foot segment model. Our results show that the adaptation of the intrinsic foot joints in terms of kinetics might be mainly due to the reduced walking speed. This reduction in walking speed might be mainly due to the pain induced by OA. As a result, when the pain is removed after surgery, the speed can be normalized. As usual, for marker-based technology, soft tissue artefacts are a limitation on the results obtained here but using fluoroscopy on multiple patients seems costly, cumbersome, and ethically arguable. In addition, more participants should be included, and various levels of ankle OA should be evaluated. However, it might be difficult to find speed matched subjects for this kind of pathology as the gait of CTRL subjects at this speed might lead to an unnatural pattern. At last, to complete this study, other joint axes should also be investigated.

#### **Disclosure statement**

No potential conflict of interest was reported by the authors.

#### References

- Arnolds JB, Bowen CJ, Chapman LS, Gates LS, Golightly YM, Halstead J, Hannan MT, Menz HB, Munteanu SE, Paterson KL, Roddy E, Siddle HJ, Thomas MJ. 2022. International Foot and Ankle Osteoarthritis Consortium review and research agenda for diagnosis, epidemiology, burden, outcome assessment and treatment. Osteorarthritis and Cartilage. In press available online 15 February 2022.
- Brodsky JW, Polo FE, Coleman SC, Bruck N. 2011. Changes in gait following the Scandinavian Total Ankle Replacement. J. Bone Jt Surg. 93:1890–1896
- Brodsky JW, Coleman SC, Smith S, Polo FE, Tenebaum S. 2013. Hindfoot motion following STAR total ankle arthroplasty: a multisegment foot model gait study. Foot Ankle Int. 34:1479–1485

- Deleu PA, Leemrijse T, Cheze L, Naaim A, Dumas R, Devos Bevernage B, Birch I, Besse JL. 2021. Post-sprain versus post-fracture post-traumatic ankle osteoarthritis: Impact on foot and ankle kinematics and kinetics. Gait Posture. 86:278–286
- Deleu PA, Naaim A, Cheze L, Dumas R, Devos Bevernage B, Birch I, Besse JL, Leemrijse T. 2022. Changes in ankle and foot kinematic after fixed-bearing total ankle replacement. J. Biomech. 136:111060
- Deschamps K, Eerdekens M, Desmet D, Matriacali GA, Wuite S, Staes F. 2017. Estimation of foot joint kinetics in three and four segment foot models using an existing proportionality scheme: Application in paediatric barefoot walking. J. Biomech. 61:168–175
- Eerdekens M, Staes F, Matriacali G, Wuite S, Peerlinck K, Deschamps K. 2019. Quantifying clinical misinterpretations associated to one-segment kinetic foot modelling in both a healthy and patient population. Clin Biomech. 67: 160–165

- Leardini A, Benedetti MG, Berti D, Bettinelli D, Nativo R, Gianni S. 2007. Rear-foot, mid-foot and fore-foot motion during the stance phase of gait. Gait Posture. 25:453-462
- Pataky TC, Vanrenterghem J, Robinson MA. 2016. The probability of false positives in zero-dimensional analyses of one-dimensional kinematic, force and EMG trajectories. J. Biomech 49:1468–1476
- Queen R. 2007. Directing clinical care using lower extremity biomechanics in patients with ankle osteoarthritis and ankle arthroplasty. J. Orthop. Res 35:2345–2355
- Sarasawat P, MacWilliams BA, Davis RB, D'Astous JL. 2014. Kinematics and kinetics of normal and planovalgus feet during walking. Gait Posture. 39:339–345

**KEYWORDS** Multi-segment Foot model; total ankle replacement; osteoarthritis

🖾 alexandre.naaim@univ-lyon1.fr