



**HAL**  
open science

# THE CONTRIBUTION OF DEEPLY SUBDUCTED CONTINENTAL CRUST TO MANTLE METASOMATISM AND DEEP CARBON CYCLE

Alessia Borghini, Silvio Ferrero, Gautier Nicoli, Patrick O'Brien, Oscar Laurent,  
Laurent Remusat, Giulio Borghini, Sula Milani

## ► To cite this version:

Alessia Borghini, Silvio Ferrero, Gautier Nicoli, Patrick O'Brien, Oscar Laurent, et al.. THE CONTRIBUTION OF DEEPLY SUBDUCTED CONTINENTAL CRUST TO MANTLE METASOMATISM AND DEEP CARBON CYCLE. GSA Connects 2021 in Portland, Oregon, Oct 2021, Portland, United States. <10.1130/abs/2021AM-367559>. <hal-03865182>

**HAL Id: hal-03865182**

**<https://hal.science/hal-03865182v1>**

Submitted on 22 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# GSA Connects 2021 in Portland, Oregon

Paper No. 233-4

Presentation Time: 2:25 PM

## THE CONTRIBUTION OF DEEPLY SUBDUCTED CONTINENTAL CRUST TO MANTLE METASOMATISM AND DEEP CARBON CYCLE

**BORGHINI, Alessia**<sup>1</sup>, **FERRERO, Silvio**<sup>2</sup>, **NICOLI, Gautier**<sup>1</sup>, **O'BRIEN, Patrick J.**<sup>1</sup>, **LAURENT, Oscar**<sup>3</sup>, **REMUSAT, Laurent**<sup>4</sup>, **BORGHINI, Giulio**<sup>5</sup> and **MILANI, Sula**<sup>5</sup>

(1)Institut für Geowissenschaften, Universität Potsdam, Potsdam, 14476, Germany, (2)Institut für Geowissenschaften, Universität Potsdam, Potsdam, 14476, Germany; Museum für Naturkunde (MfN), Berlin, 10115, Germany, (3)ETH, Zürich, Zürich, 8092, Switzerland; eCNRS, Observatoire Midi-Pyrénées, Toulouse, 31400, France, (4)Museum National d'Histoire Naturelle, Institut de minéralogie, de physique des matériaux et de cosmochimie, Paris, 75005, France, (5)Università degli Studi di Milano Statale, Dipartimento di Scienze della Terra „Ardito Desio“, Milano, 20133, Italy

Primary inclusions of metasomatic melt from melting of continental crust are trapped in garnets of ultra-high pressure (UHP) eclogites of the Erzgebirge, Bohemian Massif (Germany). Here we estimate the contribution of the subducted continental crust to mantle metasomatism and deep carbon fluxes via the characterization of these melt inclusions. They are micrometric and occur in cluster in the inner part of the garnet as both nanogranitoids (i.e. polycrystalline inclusions) and glassy, often with a shrinkage bubble. Nanogranitoids contain kumdykolite, quartz, kokchetavite, biotite, white mica, calcite and rare graphite. The inclusions are in the same microstructural position of polycrystalline quartz, interpreted as pseudomorph after coesite suggesting entrapment at UHP conditions. The melt is granitic, hydrous and slightly peraluminous; its H<sub>2</sub>O and CO<sub>2</sub> content were measured with NanoSIMS. CO<sub>2</sub> concentration, after vapor reintegration, is  $19552 \pm 772$  ppm, the highest content of CO<sub>2</sub> measured so far in crustal melt inclusions. The melt shows trace elements enrichment consistent with a continental origin, high amounts of incompatible elements (Cs in particular), Pb, Th, U, Li and B. This trace elements pattern is similar to other metasomatic melts in mantle eclogites, metasomatized mantle rocks and post-collisional ultrapotassic plutonic rocks in the Bohemian Massif, suggesting widespread mantle metasomatism from melts originated in the continental crust. We argue that a large amount of volatiles, carbon in particular, are stored at the base of the orogen during the deep subduction of the continental crust. We modelled the carbon flux during subduction of the continental crust during the formation of the Bohemian Massif (i.e. Variscan orogeny) and obtained an endogenic flux of  $65 \pm 28$  Mt C yr<sup>-1</sup>, similar, within error, to the one calculated for the oceanic crust. Thus, in collisional settings, the deep continental crust acts as a long-term carbon storage unit, preventing the completion of the carbon cycle.