



**HAL**  
open science

## **Parasitized, non-parasitized, why? A study of factors influencing tick burden in roe deer neonates**

Léa Bariod, Sonia Saïd, Clément Calenge, Stéphane Chabot, Vincent Badeau, G. Bourgoïn

### ► **To cite this version:**

Léa Bariod, Sonia Saïd, Clément Calenge, Stéphane Chabot, Vincent Badeau, et al.. Parasitized, non-parasitized, why? A study of factors influencing tick burden in roe deer neonates. *EvoLyon*, Nov 2021, Lyon, France. . hal-03865224

**HAL Id: hal-03865224**

**<https://hal.science/hal-03865224>**

Submitted on 22 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Parasitized, non-parasitized, why? A study of factors influencing tick burden in roe deer neonates.

Léa Bariod<sup>a,b,c</sup>, Sonia Saïd<sup>b</sup>, Clément Calenge<sup>b</sup>, Stéphane Chabot<sup>b</sup>, Vincent Badeau<sup>d</sup>, Gilles Bourgoïn<sup>a,c</sup>

<sup>a</sup>École nationale vétérinaire VetAgro Sup – Marcy l'Étoile, France; <sup>b</sup>Office Français de la Biodiversité - Birieux, France; <sup>c</sup>Laboratoire de Biométrie et Biologie Évolutive – Lyon, France; <sup>d</sup>Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement – Champenoux, France



## Context of study

- *Ixodes ricinus*, the most common tick in Central Europe, is an important vector of pathogens **threatening human and animal health**<sup>1</sup>.
- Tick abundance and repartition increased during last decades with global environmental changes in the continent, which include **climate and habitat changes**, and **increase of deer densities**<sup>2</sup>.
- Indeed, **cervid species** are identified as **key hosts to ticks**, especially **roe deer** (*Capreolus capreolus*) whose populations have increased in recent years in France and Europe<sup>3</sup>.
- Mammal **newborns** have a naïve and immature immune system and are therefore more vulnerable to tick infestations.
- Thus it is important to **improve knowledge of factors influencing the tick burden of neonates**. Here, we study roe deer fawns monitored during 27 years in a French population with a novel Bayesian model.



## Problematics & Hypotheses

### Biotic and abiotic factors: which ones favor ticks?


Tick burden should increase with:

- H1: age of the fawn**, as they become more mobile and more exposed to questing ticks in the vegetation. We also assume that there will be a renewal of ticks with a number of ticks that will become constant on the fawn<sup>4</sup>.
- H2: Spring weather conditions**, with higher tick burden during **warm and wet conditions**, which favor tick activity<sup>5</sup>.
- H3: habitat structure**, with higher tick burden in habitats with important shrub cover and ground-based vegetation which retains humidity<sup>6</sup>.
- H4: roe deer density**, with higher tick burden when number of adult roe deer is the highest, which there are more hosts to participate in the tick cycle and therefore in its reproduction and dispersal<sup>7</sup>.



## Material & methods

### Bayesian model describing the infestation process of the roe deer fawns by ticks according to several factors



### Study site:

« Territoire d'Étude et d'Expérimentation » of Trois-Fontaines (Grand-Est, France)



An enclosed forest of 1360 ha (48°38'N, 4°54'E)

### Dataset available:

27 years  
1992 → 2018



965 newborn fawns

Captured during breeding period of the female roe deer (late April to mid-June)

## Results


### Bayesian model with the most probable combination of variables to explain the infestation rate of fawns

Instantaneous rate of infestation  $\sim$  Age + Humidity + Year (random factor)

### Hypothese 1 ✓

The instantaneous risk of infestation increases the first 5 days of fawn life and after becomes equal to zero.

Est. of >5 days old: -17;7 [-45.75; -2.6]<sub>CI95%</sub>



### Hypothese 2 (✓)

**Humidity** 6 days prior the fawn capture had significant effects on the tick burden of fawn.

Est. of Humidity: 0.013 [-0.003; 0.03]<sub>CI95%</sub>

**Temperature** 6 days prior the fawn capture had no significant effects on the tick burden of fawn.

Est. of Temperature: -0.005 [-0.047; 0.037]<sub>CI95%</sub>

### Hypotheses 3 and 4 ✗

**Vegetation structure**, where the fawn was found, had no significant effect on the tick burden of fawn.


Est. of habitat1: -0;07 [-0.32; 0.15]<sub>CI95%</sub> Est. of habitat2: 0.06 [-0.16; 0.29]<sub>CI95%</sub> Est. of habitat3: -0.10 [-0.33; 0.11]<sub>CI95%</sub> Est. of habitat4: 0.11 [-0.12; 0.37]<sub>CI95%</sub> Est. of habitat5: -0.01 [-0.24; 0.22]<sub>CI95%</sub>

**Roe deer density** had no significant effect on the tick burden of fawn.

Est. of Roe deer density: 0.002 [-0.001; 0.005]<sub>CI95%</sub>

### Random effect of the year

Vertical red lines delimit three periods according to the Lothar hurricane (December 1999):



Standard deviation of the year random effect for each period:

1 Est.: 0.427 [0.024; 1.112]<sub>CI95%</sub> 2 Est.: 0.807 [0.410; 1.516]<sub>CI95%</sub> 3 Est.: 0.160 [0.01; 0.442]<sub>CI95%</sub>

## Conclusion

- The level of tick burden became constant when fawns were 5 days old and more. This result could be explained by the **depletion of questing ticks in the habitat of fawns** and the **turnover of tick feeding on fawns**. Indeed, ticks take their blood meal on hosts for a few days before falling to the ground, where they molt or lay eggs during few weeks.
- Humidity was significant but no effects of temperature, vegetation structure or roe deer density were found on the tick burden of fawn. However, we noticed a **strong heterogeneity of tick burden between years** suggesting two explanations:
  1. Others variables could be involved, such as **local density of various host species** or **habitat characteristics** of fawn, which may be important for tick activity.
  2. The environmental variables used in this study affected the infection process at **other spatial or temporal scales**.
- This would require **additional data on ticks in the environment**, at different times of the year, which would allow a more precise study in Trois-Fontaines to improve our understanding of tick infection in ungulate newborns.

## Corresponding author

Léa Bariod

PhD student at « École nationale vétérinaire VetAgro Sup », « Laboratoire de Biométrie et de Biologie Évolutive » and « Office Français de la Biodiversité, Direction recherche et Appui scientifique »

E-mail: bariod.lea@orange.fr  
linkedin.com/in/lea-bariod-39047312b/  
researchgate.net/profile/Lea\_Bariod



## Bibliography

1. de la Fuente, J., Estrada-Peña, A., Venzal, J. M., Kocan, K. M., & Sonenshine, D. E. (2008). Overview: ticks as vectors of pathogens that cause disease in humans and animals. *Front Biosci*, 13(13), 6938-6946.
2. Vor, T., Kiffner, C., Hagedorn, P., Niedrig, M., & Rühle, F. (2010). Tick burden on European roe deer (*Capreolus capreolus*). *Experimental and Applied Acarology*, 51(4), 405-417.
3. Kiffner, C., Lüdige, C., Alings, M., Vor, T., & Rühle, F. (2010). Abundance estimation of Ixodes ticks (Acari: Ixodidae) on roe deer (*Capreolus capreolus*). *Experimental and Applied Acarology*, 52(1), 73-84.
4. Taylor, M. A., Coop, R. L., & Walli, R. L. (2015). *Veterinary Parasitology*. John Wiley & Sons.
5. Medlock, J. M., Hansford, K. M., Bormane, A., Derdakova, M., Estrada-Peña, A., George, J. C., ... & Van Bortel, W. (2013). Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasites & vectors*, 6(1), 1-11.
6. Tack, W., Madder, M., Baeten, L., Vanhellemont, M., Gruwez, R., & Verheyen, K. (2012). Local habitat and landscape affect *Ixodes ricinus* tick abundances in forests on poor, sandy soils. *Forest ecology and management*, 265, 30-36.
7. Ruiz-Fons, F., & Gilbert, L. (2010). The role of deer as vehicles to move ticks, *Ixodes ricinus*, between contrasting habitats. *International journal for parasitology*, 40(9), 1013-1020.

## Acknowledgments

This study was conducted in the context of a PhD co-funded by the national veterinary school VetAgro Sup and the French Office for Biodiversity (OFB) under partnership agreement CNV20-0947. We are also grateful to the volunteers and technicians for participating in the fawn captures and data collection at Trois-Fontaines (France).