

Long-term observation of permafrost evolution in European mountains

Jeannette Noetzli, Hanne Hvidtfeldt Christiansen, Bernd Etzelmüller, Mauro Guglielmin, Martin Hoelzle, Ketil Isaksen, Florence Magnin, Cécile Pellet, Marcia Phillips, Paolo Pogliotti

▶ To cite this version:

Jeannette Noetzli, Hanne Hvidtfeldt Christiansen, Bernd Etzelmüller, Mauro Guglielmin, Martin Hoelzle, et al.. Long-term observation of permafrost evolution in European mountains. International Mountain Conference - IMC2022, Sep 2022, Innsbruck, Austria. hal-03865454

HAL Id: hal-03865454 https://hal.science/hal-03865454v1

Submitted on 22 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Long-term observation of permafrost evolution in European mountains

Jeannette Noetzli, WSL Institute for Snow and Avalanche Research SLF, Davos Dorf, Switzerland Hanne H. Christiansen, Geology Department, University Centre in Svalbard, Longyearbyen, Norway Bernd Etzelmueller, Department of Geosciences, University of Oslo, Norway

Mauro Gugliemin, Department of Theoretical and Applied Science, Insubria University, Italy Martin Hoelzle, Department of Geosciences, University of Fribourg, Fribourg, Switzerland Ketil Isaksen, Norwegian Meteorological Institute, Oslo, Norway

Florence Magnin, Laboratoire EDYTEM, CNRS/Université Savoie Mont-Blanc, Le Bourget-du-Lac, France

Cécile Pellet, Department of Geosciences, University of Fribourg, Fribourg, Switzerland Marcia Phillips, WSL Institute for Snow and Avalanche Research SLF, Davos Dorf, Switzerland Paolo Pogliotti, Environmental Protection Agency of Valle d'Aosta, Saint Christophe, Italy

Permafrost, ground material remaining at or below 0 °C for at least two consecutive years, is a key component of the cryosphere in high-latitude and high-altitude regions. As such it is globally observed as an Essential Climate Variable (ECV) of the Global Climate Observation System (GCOS). Mountain permafrost is permafrost in areas with steep topography and accounts for ca. 30% of the global area underlain by permafrost and can be found at low and high latitudes and in the Northern and Southern Hemispheres. We present an overview of long-term permafrost data from European mountain regions (the Alps and the Nordic countries, including Svalbard) – where most mountain permafrost data are available – and describe observed changes and patterns of permafrost evolution.

Permafrost temperatures are typically measured in boreholes. Ranges of permafrost temperature and warming rates in mountains are similar to those observed in lowland polar areas but with high spatial variability in atmospheric, surface and subsurface conditions due to the complex topography with large environmental gradients. The strongest warming was observed in the past decade for bedrock sites at high latitudes and high elevations, where permafrost temperatures are several degrees below 0 °C (e.g., in mountain sites on Svalbard or above 3500 m asl. in the Alps). Ground temperatures in ice-rich permafrost close to the lower permafrost boundary – for example in many rock glaciers in the Alps – increase at a lower rate due to latent heat uptake during ice melt. Significant increase in active layer thickness (ALT) – the thickness of the layer above the permafrost that freezes and thaws annually – was observed. In the European Alps, ALT increased by meters since the start of observations 1–2 decades ago, even doubling at some sites.

In mountain regions, permafrost degradation can affect stability of infrastructure and steep mountain slopes. Although observed parameters and techniques applied to lowland permafrost in general also apply to mountain areas, the long-term collection of robust and comparable permafrost data is challenging due to steep topography, site access, harsh weather conditions, risk of natural hazards and the strong spatial heterogeneity of determining factors. The continuation of time series over decades highly depends on field laboratories permanently operated for long-term climate monitoring. Finally, their coordination in national or regional monitoring networks (such as, for example, PERMOS, SIOS, or PermaFRANCE) is key for standardization and comparability and to describe a consistent picture of the mountain permafrost state and changes.