



# INFERRING THE STRUCTURAL PROPERTIES OF EDDIES IN THE LOG LAYER FROM SPECTRAL STATISTICS

Lionel Agostini, Michael Leschziner

## ► To cite this version:

Lionel Agostini, Michael Leschziner. INFERRING THE STRUCTURAL PROPERTIES OF EDDIES IN THE LOG LAYER FROM SPECTRAL STATISTICS. Tenth International Symposium on Turbulence and Shear Flow Phenomena, Jul 2017, Chicago, France. pp.653-658, 10.1615/tsfp10.1110 . hal-03866112

**HAL Id: hal-03866112**

**<https://hal.science/hal-03866112>**

Submitted on 22 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INFERRING THE STRUCTURAL PROPERTIES OF EDDIES IN THE LOG LAYER FROM SPECTRAL STATISTICS

Lionel Agostini and Michael Leschziner

Aeronautics Department

Imperial College London

Prince Consort Rd. London SW7 2AZ

l.agostini@imperial.ac.uk and mike.leschziner@imperial.ac.uk

## Background

The structure of near-wall layers has been the subject of much research over many years, with conceptual descriptions of the Attached Eddy Hypothesis (AEH, henceforth) by Townsend (1980) and Perry & Chong (1982) being key historical fix points. The fact that major efforts have continued unabated over the past two to three decades reflects the exceptional structural complexity of near-wall layers awaiting insight, as well as new opportunities to investigate open questions as a consequence of outstandingly high-quality experimental and DNS data emerging over the past decade (Marusic *et al.* (2013); Smits *et al.* (2011); Hultmark *et al.* (2013); Rosenberg *et al.* (2013); Jiménez & Hoyas (2008); Lee & Moser (2015)). In particular, the availability of massive amounts of spatially and temporally fully-resolved raw DNS data for fairly high Reynolds numbers has opened new routes to investigating many statistical and structural properties of near-wall layers, with the objective of unravelling a variety of scale-interaction processes.

One subject of particular interest and attention has been the origin and significance of energetic large-scale structures present in the outer parts of the log layer, which are the cause of a plateau of, and – at sufficiently high Reynolds number – an outer (second) maximum in, the streamwise energy in a boundary layer. This behaviour is illustrated by the profiles in Fig. 1, derived from channel-flow DNS data created by Lozano-Durán & Jiménez (2014) and Lee & Moser (2015), respectively.

Among many processes associated with these structures, “footprinting” and “modulation” of small(er)-scale near-wall structures, and implications arising from both to the universality of near-wall turbulence, have been major focal point of recent studies, exploiting experimental data (Mathis *et al.* (2011); Ganapathisubramani *et al.* (2012)) and DNS fields (Agostini & Leschziner (2014), Zhang & Chernyshenko (2016)). A question that is posed by the Reynolds-number-dependent elevation and distortion of the streamwise energy in the log-layer, depicted in Fig. 1, is whether the validity of the AEH in this layer can be defended, in view of the fact that the AEH is associated with a logarithmic decline of the streamwise energy away from the wall. In contrast to the variation in the log-layer, such a decay is observed, experimentally, in the remote outer region of boundary layers at very high Reynolds numbers (Hultmark *et al.* (2013); Rosenberg *et al.* (2013); Vassiliocos *et al.* (2015); Vallikivi *et al.* (2015)), but this region is well beyond the log-layer and populated by “very large-




Figure 1. Wall-normal distribution of the streamwise stress at :  $Re_\tau = 4200$  (red line) and  $Re_\tau = 5200$  (Lee & Moser (2015)). The dashed lines represent the variation  $\overline{u'u'}^+ = -1.26 \log y^+ + B$ , with  $B = 12.2$  and  $12.7$  for  $Re_\tau = 4200$  and  $Re_\tau = 5200$ , respectively.

scale structures”. Also, in this outer region, the spectrum of the streamwise energy shows the variation  $\phi_{uu}(k_x) \propto k_x^{-1}$ , which is compatible with the logarithmic decay of the streamwise energy. On the other hand, recent studies by Davidson *et al.* (2006a); Hwang (2015) and observations by Jiménez & Hoyas (2008) suggest that the AEH is also valid in the intermediate (or meso-) layer covering the log-layer region.

It is against the above background and controversy that the present study set out to examine closely the structural and spectral properties of a channel-flow boundary layer at  $Re_\tau = 4200$ , for which extensive DNS data were generated by Lozano-Durán & Jiménez (2014). The primary focus of the study is on an examination of the statistical properties of sub-ranges of scales within the pre-multiplied wall-normal spectra, and on related analysis of the pre-multiplied derivatives of the second-order structure function, the latter leading to a proposed alternative interpretation of the conventional AEH in the intermediate layer  $100 < y^+ < 2000$ .

## Results

Fig. 2 shows pre-multiplied spectra  $\phi_{uu}(\lambda_x)$  and  $\phi_{uu}(\lambda_z)$ , where  $\lambda_x$  and  $\lambda_z$  are the wave numbers in the streamwise and spanwise directions, respectively. The hor-




Figure 2. Pre-multiplied power spectrum of the streamwise fluctuations, in both streamwise (a) and spanwise direction (b); at  $Re_\tau = 4200$ . The dotted red lines show either the relation  $\lambda_x^+ = 3.5y^+$  or  $\lambda_z^+ = 7y^+ = 2\lambda_x^+$ .




Figure 3. Sub-ranges in spectral map having distinct turbulence characteristics (1c: dominance of streamwise component; 2c: dominance of cross-flow components).

izontal dashed lines identify the meso-layer, on which attention focuses. Although there is an indication that the  $\lambda_x$  and  $\lambda_z$  locations at which the energy begins to rise steeply vary linearly with  $y^+$ , in consonance with the AEH, neither spectral map features a well-defined constant-value plateau within the meso-layer. However, the existence of such a plateau, implying the variation  $\phi_{uu}(k_x) \propto k_x^{-1}$  or  $\phi_{uu}(k_z) \propto k_z^{-1}$  within a triangular region that is bounded by  $\lambda_{x,min}^+ = Cy^+$  and  $\lambda_{x,max}^+ = cst$ , can easily be shown to be

necessary in order to satisfy the AEH-compatible logarithmic decay of the streamwise energy, such as indicated in the outer region of the meso-layer in Fig.1.

In a preliminary part of the present study, the spectra  $\phi_{uu}$ ,  $\phi_{vv}$  and  $\phi_{ww}$  were combined, in a manner identified below, to determine maps of scalar parameters in the  $(\lambda_x^+ - y^+)$ -plane that allowed distinct sub-ranges of eddy scales in the spectra  $\phi_{uu}(\lambda_x)$  and  $\phi_{uu}(\lambda_z)$  to be delineated, with each sub-range having distinct statistical characteristics. The newly proposed scalar parameters express, respectively, the degree of isotropy, via  $\gamma^{3c} \equiv \frac{3|\Phi_{uu}||\Phi_{vv}||\Phi_{ww}|}{|\Phi_{uu}|^3 + |\Phi_{vv}|^3 + |\Phi_{ww}|^3}$ , and the degree anisotropy, via  $\gamma_u^{1c} \equiv \frac{|\Phi_{uu}||\Phi_{vv}|}{|\Phi_{uu}|^2 + |\Phi_{vv}|^2 + |\Phi_{ww}|^2}$ , of the energy density. The eddy-scale sub-regions so identified are shown in Fig.3. Of primary relevance and interest is sub-region C, characterized by a dominance of the streamwise energy component, relative to the two others, in the region in which the length scales of the structures is relatively large. Eddies having smaller length scales, located within region B, are close to isotropic, in terms of energy, do not carry shear stress and are therefore deemed “detached”. In contrast, the state in region C is characterized by anisotropy, with streamwise energy dominating, and the eddies are affected by shear, carry shear stress and comply with the notion of “attached” eddies.

The main part of the study focused on an investigation of maps of the pre-multiplied derivatives of the second-order structure function  $\delta \times \frac{dS_{2,u}(\delta)}{d\delta}$ , where  $\delta$  is the separation, either in  $x$  or  $z$  in  $S_{2,u}(y, \delta) = \langle [u(y, x) - u(y, x + \delta)]^2 \rangle_{z,t}$ , the subscripts  $z, t$  identifying homogeneous averaging directions. Such maps are shown in Fig. 4. The rationale of focusing on these maps arises from the observation by Davidson *et al.* (2006b,a) that there is a close relationship between  $S_{2,u}$  and the pre-multiplied spectra – a subject that will be discussed and clarified in the full paper. However, the advantage of the pre-multiplied derivatives of the structure function is that plateau regions in these map are more pronounced and thus more readily identifiable as being associated with the AEH than in the corresponding spectral maps. A key argument advanced in this paper is that the trapezoidal regions, rather than the conventional triangular areas shown in Fig 4, allow an extended or modified interpretation of the AEH in the meso-layer, as is schematically indicated in the parts (c) and (d) of Fig. 4. This argument will be set out in detail in the full paper. A supporting observation is that profiles of the structure functions themselves,  $S_{2,u}(\delta_x)$  and  $S_{2,u}(\delta_z)$ , at different  $y^+$  levels within the meso-layer collapse onto an almost universal logarithmic variation when  $\delta_x$  and  $\delta_z$  are scaled with  $y$ . This connection will also be pursued in detail in the full paper.

## REFERENCES

- Agostini, L & Leschziner, MA 2014 On the influence of outer large-scale structures on near-wall turbulence in channel flow. *Physics of Fluids* **26** (7), 075107.
- Davidson, PA, Krogstad, P-A, Nickels, TB *et al.* 2006a A refined interpretation of the logarithmic structure function law in wall layer turbulence. *Physics of Fluids* **18** (6), 5112.
- Davidson, PA, Nickels, TB & Krogstad, P- 2006b The logarithmic structure function law in wall-layer turbulence. *Journal of Fluid Mechanics* **550**, 5160.
- Ganapathisubramani, B, Hutchins, N, Monty, JP, Chung, D & Marusic, I 2012 Amplitude and frequency modula-




Figure 4. Conceptual representation of AE self-similarity through the meso-layer : (a) Maps of PMDS2 with  $\delta$  taken in streamwise and spanwise direction, respectively. Conceptual velocity profile associated to (b) the original AEH and (c) revisited AEH.

tion in wall turbulence. *Journal of Fluid Mechanics* **712**, 6191.

Hultmark, Marcus, Vallikivi, M, Bailey, SCC & Smits, AJ 2013 Logarithmic scaling of turbulence in smooth-and rough-wall pipe flow. *Journal of Fluid Mechanics* **728**, 376395.

Hwang, Yongyun 2015 Statistical structure of self-sustaining attached eddies in turbulent channel flow. *Journal of Fluid Mechanics* **767**, 254289.

Jiménez, J. & Hoyas, S. 2008 Turbulent fluctuations above the buffer layer of wall-bounded flows. *Journal of Fluid Mechanics* **611**, 215236.

Lee, M. & Moser, R. D. 2015 Direct numerical simulation of turbulent channel flow up to  $Re_\tau \approx 5200$ . *Journal of Fluid Mechanics* **774**, 395415.

Lozano-Durán, A. & Jiménez, J. 2014 Effect of the computational domain on direct simulations of turbulent channels up to  $Re_\tau = 4200$ . *Physics of Fluids* **26** (1), 011702.

Marusic, Ivan, Monty, Jason P, Hultmark, Marcus & Smits, Alexander J 2013 On the logarithmic region in wall turbulence. *Journal of Fluid Mechanics* **716**, R3.

Mathis, R., Hutchins, N. & Marusic, I. 2011 A predictive innerouter model for streamwise turbulence statistics in wall-bounded flows. *Journal of Fluid Mechanics* **681**, 537566.

Perry, AE & Chong, MS 1982 On the mechanism of wall turbulence. *Journal of Fluid Mechanics* **119**, 173217.

Rosenberg, BJ, Hultmark, Marcus, Vallikivi, M, Bailey, SCC & Smits, AJ 2013 Turbulence spectra in smooth-and rough-wall pipe flow at extreme Reynolds numbers. *Journal of Fluid Mechanics* **731**, 4663.

Smits, Alexander J, McKeon, Beverley J & Marusic, Ivan 2011 High-Reynolds number wall turbulence. *Annual Review of Fluid Mechanics* **43**, 353375.

Townsend, Albert A 1980 *The structure of turbulent shear flow*. Cambridge university press.

Vallikivi, M., Ganapathisubramani, B. & Smits, AJ 2015 Spectral scaling in boundary layers and pipes at very high Reynolds numbers. *Journal of Fluid Mechanics* **771**, 303.

Vassilicos, JC, Laval, J-P, Foucaut, J-M & Stanislas, Michel 2015 The streamwise turbulence intensity in the intermediate layer of turbulent pipe flow. *Journal of Fluid Mechanics* **774**, 324341.

Zhang, Chi & Chernyshenko, Sergei I 2016 Quasisteady quasihomogeneous description of the scale interactions in near-wall turbulence. *Physical Review Fluids* **1** (1), 014401.