



HAL
open science

Spatiotemporal dynamics of surface sediment characteristics and benthic macrofauna compositions in a temperate high-energy River-dominated Ocean Margin

Bastien Lamarque, Bruno Deflandre, Sabine Schmidt, Guillaume Bernard, Nicolas Dubosq, Melanie Diaz, Nicolas Lavesque, Frederic Garabetian, Florent Grasso, Aldo Sottolichio, et al.

► To cite this version:

Bastien Lamarque, Bruno Deflandre, Sabine Schmidt, Guillaume Bernard, Nicolas Dubosq, et al.. Spatiotemporal dynamics of surface sediment characteristics and benthic macrofauna compositions in a temperate high-energy River-dominated Ocean Margin. *Continental Shelf Research*, 2022, 247, pp.104833. 10.1016/j.csr.2022.104833 . hal-03866642

HAL Id: hal-03866642

<https://hal.science/hal-03866642>

Submitted on 23 May 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Spatiotemporal dynamics of surface sediment characteristics and**
2 **benthic macrofauna compositions in a temperate high-energy**
3 **River-dominated Ocean Margin**

4 Bastien Lamarque ^{1, *}, Bruno Deflandre ², Sabine Schmidt ², Guillaume Bernard ¹, Nicolas
5 Dubosq ², Mélanie Diaz ^{2, **}, Nicolas Lavesque ¹, Frédéric Garabetian ¹, Florent Grasso ³, Aldo
6 Sottolichio ², Sylvain Rigaud ⁴, Alicia Romero-Ramirez ¹, Marie-Ange Cordier ², Dominique
7 Poirier ², Martin Danilo ² and Antoine Grémare ¹

8 ¹ UMR EPOC, Université de Bordeaux, CNRS, UMR 5805, Station Marine d'Arcachon, 2 rue du Professeur
9 Jolyet, 33120 Arcachon, France

10 ² UMR EPOC, Université de Bordeaux, CNRS, UMR 5805, Bâtiments B18/B18N, Allée Geoffroy Saint-
11 Hilaire, CEDEX, 33615 Pessac, France

12 ³ IFREMER, DYNECO/DHYSED, Centre de Bretagne, CS 10070, 29280 Plouzané, France

13 ⁴ Université de Nîmes, EA 7352 CHROME, rue du Dr Georges Salan, 30021 Nîmes, France

14 * Correspondence: bastien.lamarque@u-bordeaux.fr

15 ** Present address: Royal Netherlands Institute for Sea Research, Department of Coastal Systems, and Utrecht
16 University, PO Box 59, 1790 AB Den Burg Texel, The Netherlands

17
18 **Abstract:**

19 The benthic compartment of River-dominated Ocean Margins (RiOMar) is largely
20 affected by sedimentary processes, as well as by natural and anthropogenic disturbances.
21 Recent studies have confirmed the major importance of riverine inputs and local
22 hydrodynamics in the spatial structuration of low- and high-energy temperate RiOMar,
23 respectively. Differences in the nature of these structuring factors could also affect the temporal
24 dynamics of these two types of systems. The present study is aiming at: (1) quantifying
25 spatiotemporal changes in surface sediment and benthic macrofauna within the West Gironde
26 Mud Patch (WGMP; high-energy system) over both short (2016-2018) and longer (2010/2016-
27 2018) time scales, (2) identifying the main environmental factors explaining those changes,
28 and (3) achieving a comparison with the Rhône River Prodelta (RRP; low-energy system) in
29 view of further characterizing the functioning of the benthic components of these two temperate
30 RiOMar. Surface sediment characteristics (grain size, quantitative and qualitative descriptors
31 of particulate organic matter) and benthic macrofauna compositions were assessed based on 4
32 seasonal sampling of 5 stations located along a depth gradient within the WGMP. Results
33 highlighted the existence of spatial patterns for both surface sediment and benthic macrofauna,
34 which are both better explained by local hydrodynamics. Most variables presented seasonal
35 changes. Benthic macrofauna compositions also showed pluri-annual changes, which were
36 attributed to a cicatrization process following a major disturbance caused by the 2013-2014
37 series of severe winter storms, and suggests the major role of local hydrodynamics in

1 explaining 2010/2016-2018 temporal changes in WGMP benthic macrofauna compositions.
2 The comparison with the RRP further highlighted major discrepancies between the two systems
3 in the main processes (i.e., hydrodynamics *versus* river hydrological regime) explaining surface
4 sediment characteristics and benthic macrofauna compositions, which supports current
5 RiOMar typologies.

6
7 **Keywords:** RiOMar; West Gironde Mud Patch; North-East Atlantic shelf; Particulate organic
8 matter; Benthic macrofauna; Spatiotemporal changes; Hydrodynamics; Storms

9
10 **Abbreviations:**

11 AFDW : Ash-Free Dry Weight

12 AJD : Annual Julian Days

13 BSS : Bottom Shear Stress

14 BSS₁₀₀ : Bottom Shear Stress integrated over 100-day periods

15 BSS₃₆₅ : Bottom Shear Stress integrated over 365-day periods

16 Chl-*a* : Chlorophyll-*a*

17 CJD : Cumulated Julian Days

18 D_{0.5} : Median diameter of sediment particles

19 dbRDA : distance-based Redundancy Analysis

20 DISTLM : DISTance-based Linear Model

21 DW : Dry Weight

22 EHAA : Enzymatically Hydrolysable Amino Acids

23 Flow₁₀₀ : River flows integrated over 100-day periods

24 Flow₃₆₅ : River flows integrated over 365-day periods

25 nMDS : non-Metric Multidimensional Scaling

26 PCA : Principal Components Analysis

27 Phaeo-*a* : Phaeophytin-*a*

28 POC : Particulate Organic Carbon

29 POM : Particulate Organic Matter

30 RiOMar : River-dominated Ocean Margin

31 RRP : Rhône River Prodelt

32 SIMPROF : SIMilarity PROFile procedure

33 SSA : Sediment Surface Area

34 THAA : Total Hydrolysable Amino Acids

35 WGMP : West Gironde Mud Patch

36

37

38 **1. INTRODUCTION**

39

40 Continental margins are key areas for the marine component of major biogeochemical
41 cycles, accounting for the mineralization of 50 to 80 % of continental Particulate Organic
42 Carbon (POC) inputs (Aller, 1998; Blair and Aller, 2012; Burdige, 2005). Continental margins

1 impacted by major river freshwater and sediment discharges are defined as River-dominated
2 Ocean Margins (RiOMar). RiOMar provide a large variety of ecosystem services including
3 provisioning (e.g. fisheries), regulating (e.g. carbon mineralization/burial) and supporting (e.g.
4 nutrients cycling, habitats) ones (e.g. Aller, 1998; Levin et al., 2001; Lansard et al., 2009).
5 Their benthic components constitute the main marine primary depositional areas of riverine
6 particule inputs (Burdige, 2005; McKee et al., 2004) and it is estimated that RiOMar account
7 for 40 to 50 % of continental POC burial in continental margins (Blair and Aller, 2012; Burdige,
8 2005; Hedges and Keil, 1995). RiOMar benthic biological compartments (e.g. macrofauna)
9 and biogeochemical fluxes (e.g. mineralization) are largely affected by a variety of natural and
10 anthropogenic disturbances (Aller, 1998; Lansard et al., 2009; Lotze et al., 2006; Rhoads et al.,
11 1985; Tesi et al., 2007; Ulses et al., 2008; Worm et al., 2006).

12
13 The impact of both sediment discharges and associated organic matter inputs on the
14 structuration of benthic macrofauna communities and biogeochemical functioning of the
15 sediment-water interface has been documented for a large variety of RiOMar (Akoumianaki et
16 al., 2013; Aller and Aller, 1986; Aller and Stupakoff, 1996; Alongi et al., 1992; Bonifácio et
17 al., 2014; Harmelin-Vivien et al., 2009; Rhoads et al., 1985; Wheatcroft, 2006). Mostly based
18 on the analysis of macrofauna vertical distribution within the sediment column and X-ray
19 radiographies, Rhoads et al. (1985) first proposed a conceptual model describing the response
20 of benthic macrofauna and surface sediments to the inputs of major rivers. According to this
21 model, macrofauna spatial distribution is mainly controlled by the interaction between: (1) the
22 physical disturbance induced by intense riverine particles inputs, and (2) Particulate Organic
23 Matter (POM) availability. In proximal (i.e., the closest to the river mouth) parts of RiOMar,
24 high and irregular sedimentation rates induce sedimentary instability, precluding the
25 establishment of mature macrobenthic communities. The high turbidity of river plumes also
26 limits primary production in the water column, resulting in mainly low (refractory) POM
27 concentrations in surface sediments. Accordingly, RiOMar proximal areas are characterized by
28 low bioturbation intensities and low mineralization fluxes. Conversely, in distal (i.e., deeper)
29 parts, moderated sedimentation and enhanced primary production in the water column allow
30 for the establishment of mature macrobenthic communities, high bioturbation intensities and
31 mineralization fluxes.

32
33 More recent RiOMar typologies, based on meta-analyses (mostly achieved on tropical
34 and subtropical systems) of geomorphological and biogeochemical processes, highlighted the
35 major effect of local hydrodynamics on RiOMar morphologies and POC mineralization/burial
36 intensities (Blair and Aller, 2012; McKee et al., 2004). This led to a clear distinction between:
37 (1) low-energy systems, with both high sedimentation rates and carbon preservation (later
38 referred as type 1), and (2) high-energy tidal and/or wave systems with both high sediment
39 oxygenation and low carbon preservation rates (later referred as type 2). Lamarque et al. (2021)
40 recently assessed the main environmental factors responsible for the spatial structuration of
41 temperate type 1 and type 2 systems based on the comparison between extensive spatial surveys
42 conducted within the Rhone River Prodelta (RRP; type 1, French Mediterranean coast) and the
43 West Gironde Mud Patch (WGMP; type 2, French Atlantic coast). Their results confirmed the

1 major importance of riverine inputs and local hydrodynamics in the spatial structuration of type
2 1 and type 2 temperate RiOMar, respectively.

3
4 Such a difference in the nature of their main structuring factors could as well affect the
5 spatiotemporal dynamics of these two types of systems. Seasonal changes in both surface
6 sediment characteristics and benthic macrofauna compositions along a depth gradient have
7 already been addressed within the RRP by Bonifácio et al. (2014). Results showed that
8 temporal changes were larger at the most proximal station in relation with changes in river
9 flows. Major changes were observed after a prolonged low-flow period, which allowed for the
10 development of a mature macrobenthic community (otherwise precluded due to high
11 sedimentation rates resulting from POM riverine inputs). Conversely, temporal changes in
12 benthic surface sediment characteristics and benthic macrofauna compositions within the
13 WGMP have only been poorly documented so far. The only available quantitative benthic
14 macrofauna composition data have been collected during a single cruise and at only 3 stations
15 (Massé et al., 2016). Consequently, the explaining factors and the magnitude of spatiotemporal
16 changes in surface sediment characteristics and benthic macrofauna compositions within the
17 WGMP (taken as an example of temperate type 2 RiOMar) are still largely unknown.

18
19 The present study is aiming at filling this gap by: (1) quantifying spatiotemporal
20 changes in surface sediment characteristics and benthic macrofauna compositions within the
21 WGMP over both short (2016-2018) and long (2010/2016-2018) time scales, (2) identifying
22 the main environmental factors explaining these changes, and (3) achieving a comparison with
23 the RRP. Main tackled questions were as follows: (1) What are the relative importance of
24 spatial and temporal changes in both WGMP surface sediment characteristics and benthic
25 macrofauna composition? (2) Are the factors better explaining spatial changes in WGMP
26 sediment profile image characteristics and spatiotemporal changes in benthic macrofauna
27 composition similar? (3) Are the differences between spatiotemporal changes in benthic
28 macrofauna composition in the RRP and the WGMP compatible with their classification as
29 type 1 and type 2 RiOMar?

30 31 32 **2. MATERIALS & METHODS**

33 34 **2.1 WGMP: Sampling area**

35
36 The West Gironde Mud Patch (WGMP) is a 420 km² sedimentary body located in the
37 Bay of Biscay 40 km off the Mouth of the Gironde Estuary, between ca. 30 and 75m depth
38 (Figure 1). This relict paleovalley is the primary depocenter (sedimentation rates between less
39 than 1 and 10 mm.y⁻¹; Dubosq et al., 2021) of fine particles originating from the Gironde
40 Estuary, which has an annual mean water flow of 944 m³.s⁻¹ (Doxaran et al., 2009) with daily
41 flows up to 7,500 m³.s⁻¹ (Constantin et al., 2018). The WGMP is located in a macro-tidal
42 environment with a tidal range from 1.5 to 5m (Jalón-Rojas et al., 2018). The continental shelf
43 off the Mouth of the Gironde Estuary is periodically submitted to strong swells/waves, which
44 can reach maximal amplitudes of 15 m and periods of 15 s (Cirac et al., 2000; Masselink et al.,


1 2016). The sedimentology of the WGMP has been extensively studied based on stratigraphic
 2 sequences, palynological data, X-ray radiographies and radiochronographies (Castaing et al.,
 3 1979, 1982; Castaing and Allen, 1981; Gadel et al., 1997; Jouanneau et al., 1989; Lesueur et
 4 al., 1991, 1996, 2001, 2002; Lesueur and Tastet, 1994; Longère and Dorel, 1970; Parra et al.,
 5 1998; Weber et al., 1991). These surveys have attributed a major role to local hydrodynamics
 6 in controlling the spatial structuration of the WGMP. This paradigm was based on both: (1) the
 7 segmentation between a proximal and a distal part with no modern persistent sedimentation in
 8 the former, and (2) the decreasing frequency of occurrence of vertical erosional sequences
 9 within the sediment column with station depth (Jouanneau et al., 1989; Lesueur et al., 1991,
 10 2001, 2002; Lesueur and Tastet, 1994). It was recently further validated by a synoptic survey
 11 of the spatial distributions of surface sediment and sediment profile image characteristics
 12 (Lamarque et al., 2021).

13
 14

15 2.2. WGMP: Sampling of surface sediment and benthic macrofauna

16

17 Five stations located along a depth gradient (Figure 1A, Table I) were sampled for
 18 surface sediment characteristics and benthic macrofauna during 4 cruises, which took place on
 19 board of the R/V Côtes de la Manche in October 2016 (Deflandre 2016), August 2017
 20 (Deflandre 2017), February 2018 (Deflandre 2018a) and April 2018 (Deflandre, 2018b). This
 21 sampling design was set to document, over the whole WGMP, seasonally contrasted situations
 22 in terms of Gironde Estuary river flows and local hydrodynamics. Only 4 stations were sampled
 23 in February 2018 due to bad meteorological conditions. During each cruise, 4 sediment cores
 24 (10 cm internal diameter) were collected at each station using an Oktopus GmbH® multiple
 25 corer. The upper top 0.5 cm of each core was sliced and immediately frozen (-20°C) on board.
 26 A single core was used to assess sediment grain size, Sediment Surface Area (SSA; Mayer,
 27 1994b), POC concentrations (hereafter POC) and $\delta^{13}\text{C}$. The remaining 3 cores were used to
 28 assess chloropigment and amino acid concentrations. Benthic macrofauna was sampled using
 29 a 0.25 m² Hamon grab (3 replicates), sieved on a 1 mm mesh and fixed with 4 % formalin.



30 **Figure 1. Sampling:** Map showing the delimitation of: (A) the West Gironde Mud Patch
 31 (WGMP) along the French Atlantic Coast together with the locations of the 5 sampled stations.
 32 Stations 1, 3 and 4 had also been sampled in July 2010 by Massé et al. (2016); and (B) the

1 Rhône River Prodelta (RRP) in the Gulf of Lions (NW Mediterranean) together with the
 2 locations of the 5 stations sampled by Bonifácio et al. (2014) in April 2007, May and December
 3 2008, and July 2011.

6 2.3. WGMP: Gironde Estuary water flows

7
 8 Daily water flows of the Gironde Estuary were assessed by summing the flows of the
 9 Garonne and Dordogne Rivers. Data were downloaded at <http://www.hydro.eaufrance.fr>.

12 **Table I.** *Sampling:* Location (WGS84, degrees, and decimal minutes) and depth of the 5
 13 stations (1, 2, 3, 8, 4) sampled in the West Gironde Mud Patch (WGMP) during the present
 14 study and of the 5 stations (A, B, N, C, D) sampled in the Rhône River Prodelta (RRP) by
 15 Bonifácio et al. (2014). (See text for details)

System	Station	Latitude (N)	Longitude (W)	Depth (m)
WGMP	1	45°45.580'	1°31.489'	37
	2	45°43.511'	1°37.773'	47.8
	3	45°41.007'	1°41.545'	56.5
	8	45°38.873'	1°45.777'	64.5
	4	45°36.924'	1°49.712'	71.5
RRP	A	43°18.690'	04°51.042'	24
	B	43°18.013'	04°50.068'	54
	N	43°17.626'	04°47.896'	67
	C	43°16.343'	04°46.565'	76
	D	43°14.917'	04°43.613'	74

34 2.4. WGMP: Local hydrodynamics

35
 36 At each of the 5 sampled stations (Figure 1, Table I), daily values of Bottom Shear
 37 Stress (BSS) were computed from a tridimensional numerical model (Diaz et al., 2020; Grasso
 38 et al., 2018). This model is based on the MARS3D hydrodynamic model (Lazure and Dumas,
 39 2008) and the WAVE WATCH III[®] wave model (Roland and Ardhuin, 2014) with a resolution
 40 of ca. 0.5x0.5 km² over the WGMP. The model integrates realistic hydro-meteorological
 41 forcing (i.e., wind, tide, surge and river flows). BSS were computed following Soulsby's

1 formulation (Soulsby, 1997). Simulations provided hourly outputs from 2010 to 2018 that were
2 daily compiled and 95th percentiles were used to characterize BSS intensities accounting for
3 intense hydrodynamic events.
4

6 **2.5. WGMP: Surface sediment characteristics**

7

8 Grain sizes were measured on aliquots of unreplicated sediment samples using a
9 Malvern[®] Master Sizer laser microgranulometer. Almost all particle-size distributions were
10 unimodal and therefore characterized through their median diameter ($D_{0.5}$). SSA was measured
11 on unreplicated freeze-dried sediment samples, previously degassed overnight at 150°C, using
12 a Gemini[®] VII Surface Area Analyzer (Micromeritics[®] 2390a model) with the multi-point
13 Brunauer-Emmett-Teller method (Mayer, 1994b). POC were assayed using a LECO[®] CS 200
14 analyzer, after 2M HCl overnight decarbonation (Etcheber et al., 1999) of previously freeze-
15 dried unreplicated sediment samples. They were normalized for SSA (i.e., expressed in terms
16 of mass per SSA) since these two parameters were assessed on the same sediment core at each
17 combination of stations*dates.
18

19 Total and Enzymatically Hydrolysable Amino Acids (THAA and EHAA) were
20 analyzed on triplicates for each of the 3 sampled cores. THAA were extracted through acid
21 hydrolysis (6M HCl, 100°C, 24h). EHAA were extracted following the biomimetic approach
22 proposed by Mayer et al. (1995). THAA and EHAA were derivatized to form fluorescent amino
23 compounds, which were separated by reverse phase High-Performance Liquid
24 Chromatography (Agilent[®] 1260 INFINITY) on a Phenomenex[®] Kinetex 5 μ m EVO C18
25 column and detected using a 338 nm excitation wavelength. Their concentrations (hereafter
26 THAA and EHAA) were expressed in terms of mass per mass of sediment Dry Weight (DW)
27 since SSA were not measured on the same sediment cores.
28

29 Chlorophyll-*a* and Phaeophytin-*a* were assayed on unreplicated thawed frozen (-20°C)
30 sediment samples after overnight acetone extraction (90 % final concentration) using a Perkin
31 Elmer[®] LS-55 spectrofluorometer following Neveux & Lantoiné (1993). Their concentrations
32 (hereafter Chl-*a* and Phaeo-*a*) were expressed in terms of mass per mass of sediment DW since
33 SSA were not measured on the same sediment cores.
34

35 Chl-*a*/(Chl-*a* + Phaeo-*a*) ratios (hereafter Chl-*a*/(Chl-*a* + Phaeo-*a*)) were used as a
36 lability index of vegetal biomass (Bonifácio et al., 2014; Pastor et al., 2011) and EHAA/THAA
37 ratios (hereafter EHAA/THAA) were used as a lability index of bulk sedimentary organics
38 (Bonifácio et al., 2014; Grémare et al., 2005; Medernach et al., 2001; Pastor et al., 2011;
39 Wakeham et al., 1997).
40

41 For the analysis of POC isotopic ratio, duplicated freeze-dried sediment samples were
42 decarbonated (1M HCl) and later analyzed using a Thermo Scientific[®] Delta V plus IRMS

1 coupled with a Thermo Scientific® Flash 2000 EA. Raw measurements were converted in usual
2 $\delta^{13}\text{C}$ units (Coplen, 2011).

3 4 5 **2.6. WGMP: benthic macrofauna**

6
7 Macrofauna was sorted, identified to the lowest tractable taxonomic level and counted.
8 Species richness was computed on pooled replicates for each combination of stations and dates.
9 Biomasses (Ash-Free Dry Weights, AFDW) were measured for each individual taxon as
10 ignition loss (450°C, 4 h) except for Ophiuridae for which an allometric regression was used
11 to account for arm loss: $\text{AFDW} = 0.0111 \times (\text{disk diameter})^{2.5268}$ (where AFDW and disk
12 diameter are expressed in g and mm, respectively; N. Lavesque, personal communication).
13 Abundances and biomasses were standardized per m^2 .

14 15 16 **2.7. WGMP: data analysis**

17
18 To identify the most pertinent time scales for assessing the structuring role of river
19 flows and local hydrodynamics on spatiotemporal changes in surface sediment characteristics
20 (i.e., mean: SSA, POC/SSA, Chl-*a*, Phaeo-*a*, Chl-*a*/(Chl-*a*+Phaeo-*a*), THAA, EHAA,
21 EHAA/THAA and $\delta^{13}\text{C}$) and benthic macrofauna compositions, the Pearson correlation
22 coefficients linking their similarity matrices (Euclidean and Bray-Curtis distances,
23 respectively) with similarity matrices based on either integrated river flows or integrated BSS
24 were computed. This was achieved using R version 3.6.1 (R Core Team, 2019) with the
25 additional packages vegan (Oksanen et al., 2019), BBmisc (Bischi et al., 2017) and Hmisc
26 (Harrell, 2021) for river flows and BSS integrated over 1 to 365 days before each of the 4
27 cruises.

28
29 For the same surface sediment characteristics as described above, hierarchical
30 clustering (Euclidean distance and average group linking) and Principal Components Analysis
31 (PCA) were used to define groups of stations*sampling dates and to assess relationships
32 between variables. A SIMilarity PROFile procedure (SIMPROF; Clarke et al., 2008) was used
33 to test for the statistical significance of between-group differences. A DISTance-based Linear
34 Model (DISTLM) was established to assess the potential contributions of river flows and BSS
35 to spatiotemporal changes in surface sediment characteristics. This model was built using the
36 Best selection procedure (Anderson et al., 2008) and the AIC selection criterion (Akaike,
37 1973). It was represented in a multidimensional space through a distance-based Redundancy
38 Analysis (dbRDA; Anderson et al., 2008). All surface sediment analyses were performed on
39 normalized data and river flows and BSS were integrated over both 100- (i.e., seasonal) and
40 365- (i.e., annual) day periods based on results of the above-described correlation analysis.
41 Station depth, Annual Julian Days (i.e., AJD: the number of days since the beginning of the
42 sampling year), and Cumulated Julian Days (i.e., CJD: the number of days since the first day

1 of the October 2016 cruise) were introduced as supplementary variables. The same approach
2 was used for benthic macrofauna abundances (square-root transformed data). However, in this
3 case and because of the use of the Bray-Curtis similarity, a non-Metric Multidimensional
4 Scaling (nMDS) was achieved instead of a PCA. Moreover, an intermediate DISTLM/dbRDA
5 was achieved using macrofauna composition as response variable and normalized surface
6 sediment characteristics as predictor variables. POC, Phaeo-*a* and EHAA were excluded from
7 this last procedure due to their strong correlation with other surface sediment characteristics.
8

9 A comparison was achieved with macrofauna abundance data collected in July 2010 at
10 stations 1, 3 and 4 (i.e., station E, C and W in Massé et al., 2016) using exactly the same
11 methodology (sampling gear and design, sorting procedure). Macrofauna identification was
12 achieved at the same taxonomic resolution by the same Research group as well. Nevertheless,
13 to avoid inconsistencies in taxa identification between the two studies, benthic macrofauna data
14 were degraded at the genus level. The comparison was achieved through hierarchical clustering
15 (square-root transformed data, Bray-Curtis distance and average group linking) and nMDS
16 together with the SIMPROF procedure.
17

18 All multivariate analyses were computed using the PRIMER[®] 6 software package
19 (Clarke and Warwick, 2001) with the PERMANOVA+ add-on (Anderson et al., 2008).
20
21

22 **2.8. Comparison between the WGMP and the RRP**

23

24 Intra-station temporal variabilities (i.e., among sampling dates considering each station
25 individually) in sediment surface characteristics and benthic macrofauna compositions were
26 compared in the WGMP (4 cruises between 2016 and 2018 as described above) and the RRP.
27 The latter was sampled by Bonifácio et al. (2014), using the same approach and methodology
28 as in the present study. More specifically, 5 stations located along a depth gradient from the
29 mouth of the Rhone River, taking into account the preferential direction of its plume (Figure
30 1B; Table I), were sampled during 4 cruises carried out in April 2007, May and December
31 2008, and July 2011 (corresponding to different hydrological regimes of the Rhône River).
32

33 These comparisons were based on the information contained in dissimilarity matrices
34 computed using: (1) D_{0.5}, POC, Chl-*a*, Phaeo-*a*, Chl-*a*/(Chl-*a*+Phaeo-*a*), THAA, EHAA and
35 EHAA/THAA (i.e., all the surface sediment characteristics measured during both studies), and
36 (2) benthic macrofauna abundances. The sums of squared distances from individual points (i.e.,
37 station*sampling date) to their group (i.e., station) centroids were computed for each station
38 (following Anderson, 2001 and Anderson et al., 2008) and used as an indices of intra-station
39 temporal variability after standardization for the number of sampling dates.
40
41

42 **3. RESULTS**


43

44 **3.1. WGMP: Water flows and Bottom Shear Stress over the 2010-2018 period**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

Water flows presented a clear seasonal pattern with winter floods (peaks up to 5 590 $\text{m}^3 \cdot \text{s}^{-1}$ on January 23 2018) and low-flow periods during summer and fall (Figure 2A). Over the whole 2010-2018 period, major events were the January 2018 and to a slightly lower extent the January 2014 floods. Between October 2016 and April 2018, there were clear inter-annual differences in water flows during the high-flow period as indicated by the moderate values recorded during the 2016-2017 as opposed to 2017-2018 winter. The July 2010 cruise (Massé et al., 2016) was achieved immediately after the end of a high-flow period. The October 2016 and August 2017 cruises were achieved during low-flow periods (112 and 67 days after the end of the preceding high-flow period, respectively). Conversely, both the February 2018 and April 2018 cruises took place during a high-flow period, respectively 5 and 87 days after the January 2018 flood.

Although clearly decreasing with depth, Bottom Shear Stress (BSS) showed similar temporal patterns at all 5 stations (Lamarque et al., 2021). Their temporal changes are therefore only presented for station 3. BSS showed a clear seasonal pattern with a succession of peaks induced by winter storms and conversely low values during summer and fall (Figure 2B). Over the whole 2010-2018 period, the major event was the succession of peaks (i.e., up to 3.23 $\text{N} \cdot \text{m}^{-2}$) induced by the repetition of strong storms during the 2013-2014 winter. The July 2010 cruise (Massé et al. 2016) was achieved at the end of a low-BSS period. During the 2016-2018 period, there were slight differences between temporal changes in water flows and BSS since: (1) high-BSS periods generally preceded high-flow periods, and (2) highest BSS (up to 2.40 $\text{N} \cdot \text{m}^{-2}$ on February 3 2017) were recorded during 2016-2017 *versus* 2018 for highest water flows. Nevertheless, both the October 2016 and August 2017 cruises were conducted during low-BSS periods (i.e., $<0.5 \text{ N} \cdot \text{m}^{-2}$) as opposed to the February 2018 and April 2018 cruises, which both took place during the 2017-2018 high-BSS period.



1

2 **Figure 2.** WGMP: Temporal changes in Gironde Estuary mean daily water flows (A) and in
 3 the 95th percentile of Bottom Shear Stress at station 3 (B) between 2010 and 2018. Short-dashed
 4 lines indicate the 4 cruises achieved in 2016-2018 (grey area) and the long-dashed line indicates
 5 the July 2010 cruise (Massé et al., 2016).

6

7

8 **3.2. WGMP: 2016-2018 spatiotemporal changes in surface sediment characteristics**

9

10 $D_{0.5}$ of surface sediments were usually larger (60, 76 and 92 μm in October 2016,
 11 August 2017 and February 2018, respectively) and much more variable (16 μm in April 2018)
 12 at station 1 (Table II, Figure 3A). At all other stations, $D_{0.5}$ were homogeneous (mean values
 13 around 20 μm) and temporally stable. Spatiotemporal changes in $\delta^{13}\text{C}$ were limited (i.e.,
 14 between -25.12 and -23.01 ‰) with the exception of station 1 in October 2016 (-18.11 ‰,
 15 Table II, Figure 3B). Particulate Organic Carbon (POC) clearly increased with station depth
 16 (i.e., from stations 1 to 4), with a maximal value of 1.79 mg.m^{-2} of Sediment Surface Area
 17 (SSA) at station 4 during April 2018 (Table II, Figure 3C). The only exception to this pattern
 18 was February 2018 with a maximal value of 1.58 mg.m^{-2} SSA at station 3. At station 4, POC
 19 was higher in April 2018 than during the 3 other cruises (mean value of 1.44 mg.m^{-2} SSA).
 20 Total Hydrolysable Amino Acids concentrations (THAA) also increased with station depth


1 (Table II, Figure 3D). THAA ranged from 0.51 mg.gDW⁻¹ (station 1 in August 2017) to 4.65
2 mg.gDW⁻¹ (stations 3 and 4 in April 2018). Temporal changes in THAA were lower at station
3 4 (variation coefficient of 5.4 % *versus* a mean of 22.0 % at the 4 other stations). Chlorophyll-
4 *a* concentrations (Chl-*a*) at station 1 were low (i.e., between 0.23 and 0.66 µg.gDW⁻¹ in August
5 2017 and April 2018, respectively) during all cruises, (Table II, Figure 3E). At all other stations,
6 Chl-*a* were much higher in April 2018 (with a maximal value of 5.86 µg.gDW⁻¹ at station 4)
7 than during the 3 other cruises (mean value of 1.09 ± 0.37 µg.gDW⁻¹ for stations 2, 3, 8 and 4).
8 The ratio between Enzymatically and Total Hydrolysable Amino Acids (EHAA/THAA)
9 clearly decreased with station depth (Table II, Figure 3F). EHAA/THAA was maximal (41.0
10 %) at station 1 in February 2018 and minimal (9.1 %) at station 4 in August 2017. Temporal
11 changes were larger at stations 4 and 1 (variation coefficient of 21.0 and 19.2 %, respectively)
12 than at the 3 other stations (mean variation coefficient of 12.7 %).

13

14

15

16




1 **Figure 3.** *WGMP*: Spatiotemporal changes in the (mean) values of surface sediment
 2 characteristics during the 4 cruises achieved between 2016 and 2018: Median grain sizes ($D_{0.5}$;
 3 **A**), $\delta^{13}\text{C}$ (**B**), Particulate Organic Carbon concentrations normalized by SSA (POC; **C**), Total
 4 Hydrolysable Amino Acids concentration (THAA; **D**), Chlorophyll-*a* concentrations (**E**), and
 5 Enzymatically/Total Hydrolysable Amino Acids ratios (EHAA/THAA; **F**). Stations are
 6 ordered according to their depth.

7

8 The Pearson correlation coefficients linking the similarity matrix based on integrated
 9 river flows and those based on either surface sediment characteristics or benthic macrofauna
 10 compositions are shown in Figure 4A. When using surface sediment characteristics, correlation
 11 coefficients first increased with integration period durations, became significant for a 53-day
 12 integration period and reached a maximal value of 0.34 for integration periods between 156

1 and 222 days. They then decreased constantly down to a value of 0.01 for an integration period
 2 of 365 days. When using benthic macrofauna compositions, correlation coefficients remained
 3 insignificant over the whole range of river flows integration periods.



4 **Figure 4.** *WGMP*: Changes in the Pearson correlation coefficient (Rho) linking the similarity
 5 matrices based on either surface sediment characteristics or benthic macrofauna
 6 compositions (data collected between 2016 and 2018) with river flows (A) and Bottom Shear Stress (B)
 7 integrated over 1 to 365-day time periods preceding the 4 cruises. Dotted lines represent 5 %
 8 significance thresholds.


9
 10

11 The Pearson correlation coefficients linking the similarity matrix based on integrated
 12 BSS and those based on either surface sediment characteristics or benthic macrofauna
 13 compositions are shown in Figure 4B. In both cases, they increased with BSS integration
 14 periods and their maximal values were reached around 1-year (i.e., 0.56 and 0.73 at 365 and
 15 351 days for surface sediment characteristics and benthic macrofauna compositions,
 16 respectively). However, the patterns of changes with increasing integration periods clearly
 17 differed. When using surface sediment characteristics, correlation coefficients were always
 18 significant and almost constant for integration periods between 1 to ca. 250 days. Conversely,
 19 when using benthic macrofauna compositions, correlation coefficients were not significant for
 20 integration periods shorter than 65 days. They then presented a relative maximum (0.19 for a

1 90-day integration period) before almost constantly increasing for integration periods longer
2 than 130 days.

3
4 The first two components of the PCA based on surface sediment characteristics
5 accounted for 72.8 % (i.e., 55.4 % and 17.4 %, respectively) of their total variance (Figure 5A).
6 Hierarchical clustering and the associated SIMPROF procedure resulted in the identification
7 of 3 groups and 2 “isolated” stations*dates (Figure 5A). “Isolated” stations*dates consisted of
8 station 2 in February 2018 and station 4 in April 2018. Group I was exclusively composed of
9 stations 1 (all sampling dates). Group II was composed of all remaining stations*dates except
10 for the April 2018 cruise. All the stations sampled during this cruise clustered into group III,
11 except stations 1 (Group I) and 4 (isolated). The first principal component was mostly defined
12 by the opposition between the quantitative and qualitative characteristics of surface
13 sedimentary organics with THAA, EHAA, POC, Chl-*a*, Phaeo-*a* and POC concentrations on
14 one side and EHAA/THAA on the other. The second component was mostly defined by the
15 Chl-*a*/(Chl-*a*+Phaeo-*a*) and to a lesser extent EHAA/THAA (Figure 5B).

16
17 The DISTLM included river flows and BSS integrated over both 100- and 365-day
18 periods (hereafter Flow₁₀₀, Flow₃₆₅, BSS₁₀₀ and BSS₃₆₅) as independent variables. It explained
19 70.7 % of the total variance of surface sediment characteristics. Its representation through the
20 first plane of the dbRDA accounted for 61.6 % of this total variance and showed two main
21 orientations (Figure 5C). The first one, mainly along the first component of the dbRDA,
22 corresponded to the positioning of the stations along the depth gradient and was mainly cued
23 by BSS₃₆₅. The second one, mainly along the second component of the dbRDA, separated the
24 4 cruises and was mainly cued by Flow₁₀₀ and to a lesser extent BSS₁₀₀. Cumulated and Annual
25 Julian Days (CJD and AJD, respectively) correlated significantly ($p < 0.05$) with this second
26 orientation. This correlation was slightly higher for CJD ($r = 0.934$ versus $r = 0.822$).




1 **Figure 5.** *WGMP*: Multivariate analyses of surface sediment characteristics recorded between
2 2016 and 2018. Projection of stations*cruses on the first plane of a Principal Component
3 Analysis (**A**). Figures refer to stations and symbols to cruises. Dotted lines indicate groups of
4 stations*cruses issued from hierarchical clustering. Correlations of the variables with the first
5 two principal components (**B**). Distance-based Redundancy Analysis based on integrated
6 Bottom Shear Stress and river flows (**C**). Station depth (*d-St*), Annual Julian Days (*AJD*) and
7 Cumulated Julian Days (*CJD*) were used as supplementary variables. SSA: Sediment Surface
8 Area; POC: Particulate Organic Carbon; Chl-*a*: Chlorophyll-*a*; Phaeo-*a*: Phaeophytin-*a*;
9 THAA; Total Hydrolysable Amino Acids; EHAA: Enzymatically Hydrolysable Amino Acids;
10 BSS_100 and BSS_365: integrated Bottom Shear Stress over 100 and 365 days; Flow_100 and
11 Flow_365: integrated river flows over 100 and 365 days. (See text for details)

1 **3.3. WGMP: 2016-2018 spatiotemporal changes in benthic macrofauna compositions**

2

3 Overall, 6391 specimens belonging to 146 taxa were collected and identified (123 taxa
 4 after pooling species at the *Ampelisca*, *Glycera* and *Nephtys* genus). Molluscs (mainly
 5 abundant at stations 1 and 2) represented 34 % of total macrofauna abundance, followed by
 6 polychaetes (27.8 %), echinoderms (22 %) and crustaceans (13.4 %).




7 **Figure 6.** WGMP: Spatiotemporal changes in the (mean) values of main univariate benthic
 8 macrofauna characteristics recorded between 2016 and 2018: species richness (A), abundance
 9 (B), and biomass (C). Hatching corresponds to the exceptional presence of: (1) numerous
 10 individuals of a single taxon for abundance barplot and (2), high biomass of a single taxon for
 11 biomass barplot. Stations are ordered according to their depth.

1 Species richness did not show any clear spatiotemporal pattern (Table II, Figure 6A). It
2 seemed higher at station 1, with a maximum of 43 taxa in August 2017. It tended to decrease
3 from stations 1 to 3, where a minimal value of 16 taxa was recorded in October 2016. Species
4 richness then seemed to increase from stations 3 to 4 in August 2017 and April 2018.
5 Equitability tended to be lower at stations 1 and 2 than at stations 3, 8 and 4 (Table II). It was
6 temporally more stable at stations 1 and 2 than at stations 8 and 4, whose variability resulted
7 from low values in April 2018 caused by the exceptionally high abundances of *Ampelisca spp.*
8 and *Hyala vitrea*. Mean macrobenthic abundances were between 882.7 (station 4*April 2018)
9 and 89.3 (station 8*August 2017) individuals per m² (Table II, Figure 6B). Abundances were
10 higher at station 1 and tended to decrease with station depth. Only stations 8*April 2018 and
11 4*April 2018 differed from this general pattern with especially high values due to high
12 abundances of *Ampelisca spp.* (182.7 and 593.3 individuals per m² at station 8 and 4
13 respectively) and *Hyala vitrea* (93.3 and 100.0 individuals per m² at stations 8 and 4
14 respectively). Changes in mean macrobenthic biomasses did not show any clear spatiotemporal
15 pattern (Table II, Figure 6C) and were characterized by the occurrence of 3 outliers featuring
16 high values. The high biomass recorded for station 2*October 2016 resulted from the presence
17 of both a single individual of *Asterias rubens* (2019.0 mgAFDW) and a high biomass of
18 numerous individuals (i.e., 57.3 per m²) of *Abra alba* (910.6 mgAFDW.m⁻²). The high
19 biomasses recorded for station 4*August 2017 and station 4*April 2018 both resulted from the
20 presence of single individuals of *Cereus pedunculatus* in August 2017 (747.0 mgAFDW) and
21 of *Nephrops norvegicus* in April 2018 (2379.2 mgAFDW).

22
23 The nMDS based on 2016-2018 macrofauna abundance data is shown in Figure 7A.
24 The horizontal dimension of the reduced space corresponded to station depth, reflecting the
25 progressive change in benthic macrofauna compositions along this gradient. *Amphiura*
26 *filiformis* and *Kurtiella bidentata* accounted for 20.6 and 40.9 % of total macrofauna abundance
27 at station 1 (43.4 and 10.3 % at station 2). Stations 8 and 4 were, conversely, characterized by
28 the presence of burrowing macrofauna (e.g. *Callianassa subterranea* and *Oestergrenia*
29 *digitata*). For all stations, the vertical dimension of the reduced space mostly separated
30 sampling cruises. The dispersion of the different cruises for each station suggests that between-
31 cruises changes in benthic macrofauna compositions were larger at deep (i.e., stations 3, 4 and
32 8) than at shallow (i.e., stations 1 and 2) stations. The hierarchical clustering further confirmed
33 this pattern with the identification of 5 groups: (I) station 1 in February and April 2018, (II)
34 station 2 during all cruises, (III) station 3 in February and April 2018, (IV) stations 8 and 4 in
35 October 2016, and (V) stations 8 and 4 in August 2017, February 2018 and April 2018.

36 The DISTLM involving surface sediment characteristics as predatory variables
37 included THAA and Chl-*a*. It explained 42.0 % of the total variance of benthic macrofauna
38 compositions. Its representation through the first plane of the dbRDA also accounted for 42.0
39 % of this total variance and showed two main orientations (Figure 7B). The first one, mainly
40 along the first component of the dbRDA, corresponded to station depth and was mainly
41 explained by THAA. The second one, mainly along the second component of the dbRDA,
42 separated stations (except station 1) sampled during April 2018 from all other stations*dates,
43 and was mainly cued by Chl-*a* concentrations. CJD correlated significantly with this second
44 orientation ($r = 0.560$, $p < 0.05$) but not AJD ($r = 0.393$, $p = 0.07$). The DISTLM involving

1 BSS and river flows included BSS₃₆₅, Flow₁₀₀ and Flow₃₆₅ as independent variables. It
 2 explained 53.5 % of the total variance of benthic macrofauna compositions. Its representation
 3 through the first plane of a dbRDA accounted for 49.9 % of this initial variance and here again
 4 showed two main orientations (Figure 7C). The first one, mainly along the first component of the
 5 the dbRDA, corresponded to station depth and was mainly explained by BSS₃₆₅. The second
 6 one, mainly along the second component of the dbRDA, separated the 4 cruises, and was
 7 mainly cued by Flow₁₀₀. The contribution of Flow₃₆₅ was poorly described by the first plane of
 8 the dbRDA. CJD and AJD correlated significantly ($p < 0.05$) with this second orientation. This
 9 correlation was slightly higher for CJD ($r = 0.992$ versus $r = 0.820$).
 10



11 **Figure 7.** WGMP: Multivariate analyses of spatiotemporal changes in benthic macrofauna
 12 compositions recorded between 2016 and 2018. Non-Metric Multidimensional Scaling of
 13 benthic macrofauna compositions data for stations*dates during 2016-2018 cruises (A).
 14 Figures refer to stations and symbols to cruises. Solid lines represent trajectories of stations

1 over time and dotted lines indicate groups of stations*dates issued from hierarchical clustering.
2 Distance-based Redundancy Analysis based on surface sediment characteristics (**B**). Distance-
3 based Redundancy Analysis based on integrated BSS and river flows (**C**). Station depth (*d-St*),
4 Annual Julian Days (*AJD*) and Cumulated Julian Days (*CJD*) were used as supplementary
5 variables. Chl-*a*: Chlorophyll-*a*; THAA; Total Hydrolysable Amino Acids; BSS_365:
6 integrated Bottom Shear Stress over 365 days; Flow_100 and Flow_365: integrated river flows
7 over 100 and 365 days.


1 **Table II.** *WGMP*: Mean values (\pm standard deviations for replicated measures) of surface sediment and main univariate benthic macrofauna
2 characteristics during the 4 cruises achieved between 2016 and 2018. $D_{0.5}$: median grain size, SSA: Sediment Surface Area, POC: Particulate
3 Organic Carbon, Chl-*a*: Chlorophyll-*a*, Phaeo-*a*: Phaeophytin-*a*, THAA: Total Hydrolyzable Amino Acids, EHAA: Enzymatically Hydrolyzable
4 Amino Acids, SR: Species Richness and J' : Pielou's evenness.

Cruise	Station	$D_{0.5}$ (μm)	SSA ($\text{m}^2\cdot\text{gDW}^{-1}$)	POC ($\text{mg}\cdot\text{m}^{-2}$ SSA)	Chl- <i>a</i> ($\mu\text{g}\cdot\text{gDW}^{-1}$)	Phaeo- <i>a</i> ($\mu\text{g}\cdot\text{gDW}^{-1}$)	Chl- <i>a</i> /(Chl- <i>a</i> +Phaeo- <i>a</i>) (%)	THAA ($\text{mg}\cdot\text{gDW}^{-1}$)	EHAA ($\text{mg}\cdot\text{gDW}^{-1}$)	EHAA/THAA (%)	$\delta^{13}\text{C}$ (‰)	Abundance ($\text{ind}\cdot\text{m}^{-2}$)	Biomass ($\text{mgAFDW}\cdot\text{m}^{-2}$)	SR ($\text{taxa}\cdot 0.75\text{m}^{-2}$)	J'
October 2016	1	60.0	5.0	0.97	0.36 \pm 0.10	2.95 \pm 0.27	10.9 \pm 2.0	0.69 \pm 0.06	0.18 \pm 0.05	26.5 \pm 4.5	-18.11 \pm 0.41	544.0 \pm 198.9	1969.2 \pm 2260.5	38	0.73
	2	26.0	6.1	1.14	0.90 \pm 0.36	8.27 \pm 1.05	10.0 \pm 4.6	2.13 \pm 0.13	0.44 \pm 0.06	20.7 \pm 1.8	-23.20 \pm 0.05	410.7 \pm 112.0	4473.4 \pm 4494.1	30	0.69
	3	20.0	8.1	1.42	0.94 \pm 0.02	8.39 \pm 1.11	10.2 \pm 1.3	2.77 \pm 0.10	0.45 \pm 0.02	16.4 \pm 1.1	-23.23 \pm 0.13	141.3 \pm 68.2	474.2 \pm 433.1	16	0.89
	8	23.1	7.5	1.43	0.87 \pm 0.35	6.75 \pm 0.37	11.3 \pm 4.2	3.60 \pm 0.23	0.53 \pm 0.07	14.9 \pm 2.6	-23.16 \pm 0.08	228.0 \pm 28.0	1922.4 \pm 890.6	27	0.88
	4	20.1	10.3	1.49	1.39 \pm 0.47	9.59 \pm 1.69	13.0 \pm 5.8	4.49 \pm 0.27	0.65 \pm 0.12	14.6 \pm 3.2	-23.01 \pm 0.03	92.0 \pm 42.3	796.2 \pm 418.2	18	0.92
August 2017	1	75.8	2.5	1.15	0.23 \pm 0.09	1.46 \pm 0.46	13.4 \pm 2.5	0.51 \pm 0.18	0.16 \pm 0.04	32.2 \pm 3.8	-24.44 \pm 0.35	784.0 \pm 360.6	958.1 \pm 552.5	43	0.67
	2	23.8	7.3	1.27	1.15 \pm 0.20	6.65 \pm 0.88	14.9 \pm 2.7	3.23 \pm 0.29	0.61 \pm 0.02	18.9 \pm 1.4	-24.22 \pm 0.55	294.7 \pm 109.3	1405.8 \pm 259.1	26	0.68
	3	21.2	8.4	1.28	0.87 \pm 0.05	6.07 \pm 1.64	13.1 \pm 3.2	3.97 \pm 0.07	0.66 \pm 0.10	16.5 \pm 2.2	-24.46 \pm 0.10	218.7 \pm 154.6	754.9 \pm 189.9	24	0.67
	8	24.4	7.6	1.40	0.77 \pm 0.29	5.25 \pm 0.57	12.7 \pm 4.0	3.55 \pm 0.10	0.44 \pm 0.01	12.3 \pm 0.5	-24.22 \pm 0.19	89.3 \pm 43.1	1081.0 \pm 1147.4	18	0.92
	4	21.9	9.3	1.50	0.91 \pm 0.18	6.90 \pm 1.28	12.0 \pm 4.0	4.10 \pm 0.19	0.38 \pm 0.02	9.1 \pm 0.5	-24.03 \pm 0.38	176.0 \pm 60.4	2569.4 \pm 1698.6	27	0.92
February 2018	1	92.0	1.9	0.89	0.52 \pm 0.37	2.53 \pm 1.58	16.4 \pm 1.8	0.72 \pm 0.32	0.29 \pm 0.12	41.0 \pm 2.0	-25.12 \pm 0.23	840.0 \pm 342.0	1625.4 \pm 541.0	32	0.62
	2	17.5	14.2	1.16	2.09 \pm 0.72	9.61 \pm 2.47	17.6 \pm 1.5	3.10 \pm 0.16	0.59 \pm 0.14	19.0 \pm 4.2	-24.58 \pm 0.48	269.3 \pm 217.0	1149.6 \pm 1093.7	35	0.81
	3	23.7	6.2	1.58	0.96 \pm 0.19	6.22 \pm 0.63	13.3 \pm 1.3	2.87 \pm 0.43	0.45 \pm 0.06	15.7 \pm 0.3	-24.41 \pm 0.01	174.7 \pm 130.1	1113.3 \pm 1190.6	27	0.85
	4	17.4	12.6	1.32	1.12 \pm 0.17	9.51 \pm 1.86	10.6 \pm 1.0	4.32 \pm 0.43	0.57 \pm 0.07	13.3 \pm 3.0	-23.91 \pm 0.10	162.7 \pm 112.4	1313.4 \pm 1035.7	22	0.88
April 2018	1	16.3	12.9	0.93	0.66 \pm 0.05	3.67 \pm 0.38	15.2 \pm 0.5	0.98 \pm 0.51	0.26 \pm 0.04	29.6 \pm 11.0	-24.29 \pm 0.002	821.3 \pm 615.5	1271.2 \pm 175.0	35	0.62
	2	15.8	11.0	1.44	4.54 \pm 1.96	18.16 \pm 8.17	20.1 \pm 0.6	3.94 \pm 0.67	0.56 \pm 0.07	14.2 \pm 1.4	-24.05 \pm 0.05	314.7 \pm 32.6	1569.5 \pm 1071.7	25	0.67
	3	17.9	7.7	1.49	3.48 \pm 1.98	15.57 \pm 8.18	18.1 \pm 0.6	4.65 \pm 1.44	0.56 \pm 0.11	12.2 \pm 1.2	-24.11 \pm 0.20	134.7 \pm 50.0	306.8 \pm 354.9	25	0.84
	8	16.2	10.0	1.48	3.99 \pm 1.66	18.37 \pm 6.26	17.6 \pm 2.1	4.36 \pm 0.26	0.61 \pm 0.03	14.0 \pm 0.9	-23.69 \pm 0.18	388.0 \pm 69.4	1522.5 \pm 1301.5	28	0.62
	4	17.4	8.9	1.79	5.86 \pm 0.85	29.35 \pm 2.83	16.6 \pm 1.3	4.65 \pm 0.49	0.72 \pm 0.20	15.3 \pm 2.8	-23.74 \pm 0.31	882.7 \pm 202.6	4985.9 \pm 5475.4	33	0.45

1 **3.4. WGMP: 2010/2016-2018 spatiotemporal changes in benthic macrofauna**
2 **compositions**

3
4 The 2010/2016-2018 comparison of univariate benthic macrofauna descriptors
5 highlighted major differences between July 2010 (Massé et al., 2016) and the 2016-2018
6 cruises. Species richness values were in the same order of magnitude in July 2010 (32, 22 and
7 21 taxa at stations 1, 3 and 4, respectively) and in 2016-2018 (means of 37, 23 and 25 taxa at
8 stations 1, 3 and 4, respectively). Conversely, abundances were much higher at station 1 in July
9 2010 than during all 2016-2018 cruises (1628.0 vs 747.3 ± 137.5 ind.m⁻², respectively).
10 Abundances were also higher at station 3 in July 2010 (328.0 vs 167.4 ± 38.5 ind.m⁻²). They
11 conversely tended to be equivalent at station 4 (118.7 in July 2010 vs 155.0 ± 43.4 ind.m⁻²
12 during 2016-2018 cruises) when excluding the numerous *Ampelisca spp.* and *Hyala vitrea*
13 individuals present in April 2018. Biomasses were higher at stations 1 and 3 in July 2010
14 (4696.0 and 3495.7 mgAFDW.m⁻², respectively) than during the 2016-2018 cruises (means of
15 1456.0 ± 437.5 and 662.3 ± 353.0 mgAFDW.m⁻², respectively). They also tended to be
16 equivalent at station 4 between July 2010 (1901.9 mgAFDW.m⁻²) and 2016-2018 cruises
17 (1374.2 ± 436.1 mgAFDW.m⁻²) when excluding the large and rarely present individuals
18 identified at station 4 during August 2017 and April 2018 (see section 3.3).

19
20 The representation of stations*dates through the nMDS based on macrofauna
21 abundance data from the 2010 and 2016-2018 cruises is shown in Figure 8. The horizontal
22 dimension accounted for the positioning of stations along the depth gradient, whereas the
23 vertical dimension separated sampling dates and more specifically July 2010 (Massé et al.,
24 2016) from all the 2016-2018 cruises. Hierarchical clustering confirmed this pattern with the
25 identification of 5 clusters. The first 4 were almost identical to those of the analysis conducted
26 on the sole 2016-2018 data (see above) and group V was only constituted by stations 3 and 4
27 in July 2010. The positioning of stations*dates along the horizontal dimension were similar in
28 2010 and 2016-2018 as opposed to their positioning along the second dimension, which
29 showed major 2010/2016-2018 changes in benthic macrofauna compositions and suggested
30 that these changes were larger at stations 3 and 4 than at station 1.



1
2
3
4
5
6
7
8
9

Figure 9. Comparison between the West Gironde Mud Patch (WGMP) and the Rhône River Prodelta (RRP): Relationship between station depth and intra-station temporal variabilities (standardized squared distances from points to their group centroid) in sediment surface characteristics (A) and benthic macrofauna compositions (B) in both the WGMP (data collected between 2016 and 2018) and the RRP (data collected between 2007 and 2011; Bonifácio et al., 2014). (See text for details)

10
11
12
13
14
15

4. DISCUSSION

4.1. WGMP 2016-2018 spatiotemporal changes

4.1.1. Surface sediment characteristics

1
2 Spatial changes in surface sediment characteristics were larger than temporal ones. Station
3 1 was located in the proximal part of the West Gironde Mud Patch (WGMP). It presented
4 coarser surface sediments during 3 cruises (i.e., October 2016, August 2017 and February
5 2018) than during April 2018. This likely resulted from the transient deposition of coarser
6 sediments during high-energy events (Lesueur et al., 1991, 1996, 2001, 2002; Lesueur and
7 Tastet, 1994). All other 4 stations were located in the distal part. They presented restricted
8 spatial changes in surface sediment granulometry, but conversely a clear increase in bulk
9 organic contents (i.e., Particulate Organic Carbon and Total Hydrolysable Amino Acids) with
10 station depth, which supports previous observations by Lamarque et al. (2021) and Massé et
11 al. (2016). Such a pattern is rather uncommon in River-dominated Ocean Margins (RiOMar;
12 e.g., Bonifacio et al., 2014; Cathalot et al., 2013; Goñi et al., 1998; Gordon et al., 2001; Keil et
13 al., 1997) and has been attributed to a particle sieving process cued by particle density
14 (Lamarque et al., 2021) rather than size (Hedges and Keil, 1995; Keil et al., 1998; Mayer,
15 1994a, b) during the succession of sedimentation/resuspension cycles governing the transfer of
16 particles offshore in RiOMar (Blair and Aller, 2012). The results of the DISTLM/dbRDA
17 analysis support the predominant role of local hydrodynamics as the driving factor of the spatial
18 distribution of surface sediment characteristics in the WGMP as already put forward by
19 Lamarque et al. (2021) based on a synoptic spatial survey. Caution should nevertheless be taken
20 in interpreting this result since river flows did not include any spatial component.

21
22 The importance of the seasonal component in driving 2016-2018 temporal changes in
23 surface sediment characteristics is supported by the results of the DISTLM/dbRDA analysis,
24 showing that temporal changes correlate better with river flows, and to a lesser extent
25 hydrodynamics, integrated over a seasonal (i.e., Flow₁₀₀ and BSS₁₀₀) than a yearly period (i.e.,
26 Flow₃₆₅ and BSS₃₆₅). The most important changes in surface sediment characteristics occurred
27 during spring 2018. A spring increase in chloropigment concentrations has already been
28 reported by Relexans et al. (1992) in the distal part of the WGMP. The seasonality of pelagic
29 primary production in this area of the Bay of Biscay is characterized by the occurrence of a
30 spring phytoplankton bloom (Herbland et al., 1998; Labry et al., 2002). Both fluorescence
31 profiles and satellite image analyses (Copernicus Sentinel data 2018 processed using SNAP
32 software, data available on request to B. Lamarque) confirmed that the period immediately
33 preceding April 2018 was characterized by high Chlorophyll-*a* (Chl-*a*) concentrations (i.e., ca.
34 10 mg.m⁻³) in surface waters overlying the WGMP. The high chloropigment concentrations
35 and Chl-*a*/(Chl-*a*+Phaeo-*a*) recorded in surface sediments during April 2018 thus probably
36 resulted from the sedimentation of a spring bloom.

37 38 39 **4.1.2. Benthic macrofauna**

40
41 Our results clearly show the higher importance of spatial relative to 2016-2018
42 temporal changes in benthic macrofauna composition. Benthic macrofauna compositions
43 changed with station depth. The spatial pattern observed in July 2010 by Massé et al. (2016)
44 was strongly dominated by differences between the proximal (i.e., station 1) and the distal (i.e.,

1 stations 3 and 4) parts. During the present study, stations 1 and 2 were dominated by *Amphiura*
2 *filiformis* and *Kurtiella bidentata*, which is consistent with the presence of the *A. filiformis*-*K.*
3 *bidentata*-*Abra nitida* community in cohesive muddy sands off wave exposed coast (Hiscock,
4 1984; Picton et al., 1994). This corresponds to the A5.351 (“*Amphiura filiformis*, *Kurtiella*
5 *bidentata* and *Abra nitida* in circalittoral sandy mud”) habitat of the EUNIS classification
6 (Bajjouk et al., 2015). Stations 8 and 4 were strongly bioturbated (Lamarque et al., 2021 and
7 unpubl. data) and characterized by the presence of: (1) a large variety of polychaetes, (2)
8 seapens (*Cavernularia pusilla* and *Veretillum cynomorium*) and (3) *Nephrops norvegicus* (G.
9 Bernard, personal observation). They therefore can be related with the EUNIS A5.361
10 (“Seapens and burrowing megafauna in circalittoral fine mud”) habitat (Bajjouk et al., 2015).
11 Our results thus confirm that benthic macrofauna compositions clearly differ between the
12 proximal and distal parts of the WGMP. They however also show the occurrence of a depth
13 gradient in these compositions within the sole distal part.

14
15 Overall, benthic macrofauna abundances tended to decrease with depth, which is
16 conform to the trend observed by Massé et al. (2016). Such a decreasing pattern is not fully
17 consistent with the results of surveys achieved in other RiOMar, where increasing trends in
18 benthic macrofauna abundances with the distance to the river mouth have been most often
19 observed (Akoumianaki et al., 2013; Aller and Aller, 1986; Alongi et al., 1992; Bonifácio et
20 al., 2014; Rhoads et al., 1985) and interpreted based on the Rhoads et al. (1985) model.
21 According to this model, benthic macrofauna communities at the immediate vicinity of river
22 mouths are limited by the sediment instability induced by riverine inputs. During the present
23 study, spatial changes in benthic macrofauna compositions correlated best with BSS₃₆₅ and not
24 significantly with river flows. Although caution should clearly be taken (see above), this
25 suggests that local hydrodynamics is the main factor explaining benthic macrofauna
26 composition in the WGMP. Its predominance over riverine inputs within the proximal part
27 being further supported by the lack of modern sedimentation (Lesueur et al., 2002, 2001;
28 Lesueur and Tastet, 1994). Along the same line, the importance of local hydrodynamics in
29 explaining spatial changes in sediment profile image characteristics was put forward by
30 Lamarque et al. (2021). According to the current RiOMar typology (Blair and Aller, 2012), the
31 WGMP would therefore correspond to a type 2/high energy “bypassed” (*sensu* McKee et al.,
32 2004) system where the impact of local hydrodynamics is dominant relative to the one of
33 riverine inputs.

34
35 2016-2018 temporal changes in benthic macrofauna compositions tended to be larger
36 at deep stations than at the two shallowest ones, with some evidence that these changes included
37 both a seasonal and an inter-annual component. There are several rationales for the occurrence
38 of seasonal changes. First, in the reduced space of the nMDS, individual station trajectories
39 tended to “close” when they were defined based on Annual Julian Days (AJD), which can be
40 considered as indicative of within year/seasonal variability. Second, temporal changes
41 correlated best with Flow₁₀₀, which suggests the link between changes in seasonally integrated
42 river flows and benthic macrofauna compositions. Third, AJD correlated significantly with the
43 second component of the dbRDA, which was indicative of temporal changes. There is some
44 indirect evidence for an inter-annual component to 2016-2018 temporal changes as well since:

1 (1) a common inter-annual trend was observed at stations 1, 8 and 4, and (2) the second
2 component of the dbRDA correlated slightly better with Cumulated Julian Days (CJD;
3 indicative of inter-annual/cumulated changes) than with AJD.
4
5

6 **4.2. WGMP: 2010/2016-2018 comparison of benthic macrofauna compositions**

7

8 The existence of an inter-annual component to temporal changes in benthic macrofauna
9 compositions became even more obvious when July 2010 (Massé et al., 2016) and 2016-2018
10 data were compared. Differences between 2010 and 2016-2018 were indeed much larger than
11 differences observed during 2016-2018. At all 3 sampled stations, differences in benthic
12 macrofauna compositions relative to July 2010 were maximal in October 2016 and, for most
13 of them, changes occurring during the 2016-2018 period tended to reduce those differences,
14 which would be indicative of an ongoing cicatrization process following a major disturbance
15 (Pearson and Rosenberg, 1978) that would have taken place between July 2010 and October
16 2016.
17

18 In the WGMP, local hydrodynamics is a key factor structuring the spatial distributions
19 of benthic macrofauna composition and activity (Lamarque et al., 2021 and above). The
20 frequency of sediment disturbances induced by strong hydrodynamic events have been
21 documented based on the analysis of vertical erosional sequences within the sediment column
22 (Lesueur et al., 2002, 1991; Lesueur and Tastet, 1994). It has been estimated that such
23 disturbances occur every ca. 22 years at 35m depth and only every ca. 80 years at 64m depth
24 (Lesueur et al., 2002). The analysis of the 2010-2018 Bottom Shear Stress time series showed
25 higher values during the 2013-2014 winter when the WGMP experienced higher significant
26 wave heights and a longer total storm duration (i.e., 40 and 300 %, respectively) than the overall
27 means of the 1948-2015 period. Overall, the 2013-2014 winter has been the most energetic
28 during the 1948-2015 period along most of the European Atlantic coast (Masselink et al.,
29 2016). Storm-induced physical disturbance during this uncommon winter is likely to have
30 affected surface sediments, and also possibly, benthic macrofauna compositions over the whole
31 WGMP. In this context, the increase in 2010/2016-2018 temporal changes in benthic
32 macrofauna compositions with station depth may seem counter-intuitive since the intensity of
33 physical disturbance during strong storms is clearly higher at shallower stations. In “normal”
34 conditions (i.e., not during extreme events), these stations are, however, also experiencing
35 tougher hydrodynamics. Moreover, the frequency of extreme events affecting the sediment
36 column and thus potentially benthic macrofauna compositions (Dobbs and Vozarik, 1983;
37 Glémarec, 1978; Rees et al., 1977) also decreases with station depth (Lesueur et al., 2002). One
38 can thus assume that resident benthic macrofauna in the WGMP are less tolerant to physical
39 disturbance at deeper than at shallower stations, thereby accounting for larger changes in
40 benthic macrofauna composition at deeper stations. Based on these elements, our current
41 interpretation is that: (1) the repetition of major storms during the 2013-2014 winter has
42 induced strong physical disturbances of benthic habitats and consequently strong changes in
43 benthic macrofauna compositions over the whole WGMP, and (2) since then temporal changes
44 in benthic macrofauna compositions consist in an ongoing (pluri-annual) cicatrization process

1 superimposed to a seasonal dynamics (Cárcamo et al., 2017). It should nevertheless be stressed
2 that this working hypothesis clearly remains to be further tested through an extension of the
3 WGMP observation period.

6 **4.3. Comparison between the WGMP and the RRP**

8 Intra-station temporal variabilities in surface sediment characteristics were much higher
9 at the immediate vicinity of the Rhone River Mouth than in the proximal area of the WGMP.
10 In the shallowest area of the Rhône River Prodelta (RRP), this variability is mainly resulting
11 from high sedimentation during floods (Bonifácio et al., 2014; Cathalot et al., 2010; Pastor et
12 al., 2018). Mostly during autumn and winter, part of deposited sediments are resuspended
13 initiating a series of resuspension/depositions, which results in the displacement of fine
14 particles offshore (Marion et al., 2010; Ulses et al., 2008). During this transfer, particles are
15 sorted relative to their size and the most labile particulate organic matter component is degraded
16 (Bonifácio et al., 2014), which contributes to reduce temporal variabilities in surface sediment
17 characteristics in deeper areas. Conversely, temporal variabilities in surface sediment
18 characteristics tended to be constant all over the WGMP. In the proximal part, temporal
19 variability is mostly linked to granulometry, whereas in the distal part, it mostly resulted from
20 elevated chloropigment concentrations and ratios in April 2018. Dilution/mixing effects
21 between continental and marine Particulate Organic Matter sources are affecting the whole
22 WGMP as indicated by the almost constancy in surface sediment $\delta^{13}\text{C}$ and previous water
23 column observations by Fontugne and Jouanneau (1987). Conversely, they clearly increase
24 with the distance to the river mouth (and thus station depth) in the RRP (Bourgeois et al., 2011;
25 Cathalot et al., 2013; Lansard et al., 2009; Tesi et al., 2007). Moreover, due to strong local
26 hydrodynamics, the fine particles originating from the Gironde Estuary do not settle in the
27 proximal part of the WGMP (Lesueur et al., 2001), whereas conversely, high sedimentation
28 rates are taking place at the immediate vicinity of the Rhône River Mouth (Miralles et al., 2005;
29 Zuo et al., 1991). Overall, in the proximal part of the WGMP, this contributes to reduce
30 temporal variabilities in surface sediment characteristics possibly induced by seasonal changes
31 in the hydrological regime of the Gironde Estuary.

33 Intra-station temporal variabilities in benthic macrofauna compositions showed
34 opposite spatial patterns in the RRP and the WGMP. In the RRP, temporal variabilities were
35 much higher at the immediate vicinity of the Rhône River due to the shift between high and
36 low sedimentation rates during high- and low-flow periods, respectively (Bonifácio et al.,
37 2014). This pattern largely conforms to the Rhoads et al. (1985) model, which attributes a major
38 role to sedimentation in controlling benthic macrofauna in RiOMar. Conversely, temporal
39 variabilities in benthic macrofauna compositions within the WGMP tended to be lower at the
40 2 shallowest stations than at the 3 deepest ones, which could be attributed a better tolerance of
41 benthic macrofauna to wave-induced disturbance in the shallowest part of the WGMP (see
42 above). Here again, this pinpoints to differences in the relative roles of riverine inputs and local
43 hydrodynamics in explaining spatiotemporal changes within the two systems and thereby
44 supports the current RiOMar typologies (Blair and Aller, 2012; McKee et al., 2004).

5. CONCLUSIONS

The present assessment of 2016-2018 spatiotemporal changes in West Gironde Mud Patch (WGMP) surface sediment characteristics confirms the existence of: (1) a spatial structuration relative to station depth mainly explained by local hydrodynamics, and (2) seasonal changes mainly resulting from the sedimentation of the spring bloom.

Our results also demonstrate the existence of: (1) differences in benthic macrofauna composition between the proximal and distal parts of the WGMP, and (2) a depth gradient in its distal part. Overall, spatial changes in benthic macrofauna composition were best explained by local hydrodynamics integrated over a 1-year period. Benthic macrofauna compositions presented inter-annual changes superimposed over seasonal ones. The marked shift observed between 2010 and 2016-2018 suggests that a major disturbance occurred between these two periods. Our current working hypothesis is that this disturbance was caused by the series of extremely severe winter storms that took place during 2013-2014.

The comparison between the WGMP and Rhône River Prodelta (RRP) shows major differences in the control of temporal variabilities in both surface sediment characteristics and benthic macrofauna compositions within their proximal parts. In the RRP, these variabilities are associated with changes in sedimentation rates in relation with the Rhône River hydrological regime, whereas in the WGMP they result from transient particle deposition induced by intense hydrodynamic events.

Our results further support the characterization of the WGMP as temperate type 2 (i.e., high-energy system) RiOMar. They highlight the different mechanisms involved in the control of surface sediment characteristics and benthic macrofauna compositions in the WGMP and the RRP and are in line with current RiOMar typologies derived from meta-analyses mainly achieved on tropical and subtropical systems (Blair and Aller, 2012; McKee et al., 2004).

Author Contributions: Conceptualization, B.L., B.D., and A.G.; Data curation, B.L.; Formal analysis, B.L. and A.G.; Funding acquisition, B.D. and A.G.; Investigation, B.L., B.D., S.S., G.B., N.D., and A.G.; Methodology, B.L., B.D., S.S., M.D. (Mélanie Diaz), F.G. (Florent Grasso), A.S. and A.G.; Resources, B.L., B.D., S.S., G.B., N.D., M.D. (Mélanie Diaz), N.L., F.G. (Frédéric Garabetian), F.G. (Florent Grasso), A.S., S.R., A.R.-R., M.-A.C., D.P. and M.D. (Martin Danilo); Validation, B.L., B.D., and A.G.; Visualization, B.L. and A.G.; Writing—original draft, B.L. and A.G.; Writing—review and editing, B.L., B.D., S.S., G.B., N.D., F.G. (Frédéric Garabetian), F.G. (Florent Grasso), S.R., A.R.-R. and A.G. All authors have read and agreed to the published version of the manuscript.

Acknowledgements

1 This work is part of the PhD thesis of Bastien Lamarque (Bordeaux University). Bastien
2 Lamarque was partly supported by a doctoral grant from the French “Ministère de
3 l’Enseignement Supérieur, de la Recherche et de l’Innovation”. This work was supported by:
4 (1) the JERICO-NEXT project (European Union’s Horizon 2020 Research and Innovation
5 program under grant agreement no. 654410), (2) the VOG project (LEFE-CYBER and
6 EC2CO-PNEC), and (3) the MAGMA project (COTE cluster of Excellence ANR-10-LABX-
7 45). It also benefited from additional fundings allocated by the Conseil Régional Nouvelle-
8 Aquitaine and the Office Français de la Biodiversité. Operations at sea were funded by the
9 French Oceanographic Fleet. The authors wish to thank the crew of the R/V Côtes de la Manche
10 for their help during field sampling, Christophe Fontanier, Marie Claire Perello, Pascal Lebleu
11 and Hervé Derriennic for their help during sampling and analyses.
12
13

14 **References**

- 15 Akaike, H., 1973. Information theory as an extension of the maximum likelihood principle, in:
16 Csaki, F., Petrov, B.N. (Eds.), Proceedings, 2nd International Symposium on Information
17 Theory. Budapest, pp. 267–281.
- 18 Akoumianaki, I., Papaspyrou, S., Kormas, K.A., Nicolaidou, A., 2013. Environmental
19 variation and macrofauna response in a coastal area influenced by land runoff. *Estuar.
20 Coast. Shelf Sci.* 132, 34–44. <https://doi.org/10.1016/j.ecss.2012.04.009>
- 21 Aller, J.Y., Aller, R.C., 1986. General characteristics of benthic faunas on the Amazon inner
22 continental shelf with comparison to the shelf off the Changjiang River, East China Sea.
23 *Cont. Shelf Res.* 6, 291–310. [https://doi.org/10.1016/0278-4343\(86\)90065-8](https://doi.org/10.1016/0278-4343(86)90065-8)
- 24 Aller, J.Y., Stupakoff, I., 1996. The distribution and seasonal characteristics of benthic
25 communities on the Amazon shelf as indicators of physical processes. *Cont. Shelf Res.*
26 16, 717–751. [https://doi.org/10.1016/0278-4343\(96\)88778-4](https://doi.org/10.1016/0278-4343(96)88778-4)
- 27 Aller, R.C., 1998. Mobile deltaic and continental shelf muds as suboxic, fluidized bed reactors.
28 *Mar. Chem.* 61, 143–155. [https://doi.org/10.1016/S0304-4203\(98\)00024-3](https://doi.org/10.1016/S0304-4203(98)00024-3)
- 29 Alongi, D.M., Christoffersen, P., Tirendi, F., Robertson, A.I., 1992. The influence of
30 freshwater and material export on sedimentary facies and benthic processes within the Fly
31 Delta and adjacent Gulf of Papua (Papua New Guinea). *Cont. Shelf Res.* 12, 287–326.
32 [https://doi.org/10.1016/0278-4343\(92\)90033-G](https://doi.org/10.1016/0278-4343(92)90033-G)
- 33 Anderson, M., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA + for PRIMER user manual.
- 34 Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance.
35 *Austral Ecol.* 26, 32–46. [https://doi.org/https://doi.org/10.1111/j.1442-
36 9993.2001.01070.pp.x](https://doi.org/https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x)
- 37 Bajjouk, T., Guillaumont, B., Michez, N., Thouin, B., Croguennec, C., Populus, J., Louvel-
38 Glaser, J., Gaudillat, V., Chevalier, C., Tourolle, J., Hamon, D., 2015. Classification
39 EUNIS, Système d’information européen sur la nature : Traduction française des habitats
40 benthiques des Régions Atlantique et Méditerranée. Vol. 2. Habitats subtidaux &
41 complexes d’habitats. <https://doi.org/https://archimer.ifremer.fr/doc/00271/38223/>

- 1 Bischl, B., Lang, M., Bossek, J., Horn, D., Richter, J., Surmann, D., 2017. BBmisc:
2 Miscellaneous Helper Functions for B. Bischl. R Packag. version 1-1-1.
- 3 Blair, N.E., Aller, R.C., 2012. The Fate of Terrestrial Organic Carbon in the Marine
4 Environment. *Ann. Rev. Mar. Sci.* 4, 401–423. [https://doi.org/10.1146/annurev-marine-](https://doi.org/10.1146/annurev-marine-120709-142717)
5 [120709-142717](https://doi.org/10.1146/annurev-marine-120709-142717)
- 6 Bonifácio, P., Bourgeois, S., Labrune, C., Amouroux, J.M., Escoubeyrou, K., Buscail, R.,
7 Romero-Ramirez, A., Lantoine, F., Vétion, G., Bichon, S., Desmalades, M., Rivière, B.,
8 Deflandre, B., Grémare, A., 2014. Spatiotemporal changes in surface sediment
9 characteristics and benthic macrofauna composition off the Rhône River in relation to its
10 hydrological regime. *Estuar. Coast. Shelf Sci.* 151, 196–209.
11 <https://doi.org/10.1016/j.ecss.2014.10.011>
- 12 Bourgeois, S., Pruski, A.M., Sun, M.Y., Buscail, R., Lantoine, F., Kerhervé, P., Vétion, G.,
13 Rivière, B., Charles, F., 2011. Distribution and lability of land-derived organic matter in
14 the surface sediments of the Rhône prodelta and the adjacent shelf (Mediterranean Sea,
15 France): A multi proxy study. *Biogeosciences* 8, 3107–3125. [https://doi.org/10.5194/bg-](https://doi.org/10.5194/bg-8-3107-2011)
16 [8-3107-2011](https://doi.org/10.5194/bg-8-3107-2011)
- 17 Burdige, D.J., 2005. Burial of terrestrial organic matter in marine sediments: A re-assessment.
18 *Global Biogeochem. Cycles* 19, 1–7. <https://doi.org/10.1029/2004GB002368>
- 19 Cárcamo, P.J., Hernández-Miranda, E., Veas, R., Quiñones, R.A., 2017. Macrofaunal
20 community structure in Bahía Concepción (Chile) before and after the 8.8 Mw Maule
21 mega-earthquake and tsunami. *Mar. Environ. Res.* 130, 233–247.
22 <https://doi.org/10.1016/j.marenvres.2017.07.022>
- 23 Castaing, P., Allen, G., Houdart, M., Moign, Y., 1979. Étude par télédétection de la dispersion
24 en mer des eaux estuariennes issues de la Gironde et du Pertuis de Maumusson. *Oceanol.*
25 *Acta* 2, 459–468. <https://doi.org/https://archimer.ifremer.fr/doc/00122/23361/>
- 26 Castaing, P., Allen, G.P., 1981. Mechanisms controlling seaward escape of suspended sediment
27 from the Gironde: A macrotidal estuary in France. *Mar. Geol.* 40, 101–118.
28 [https://doi.org/10.1016/0025-3227\(81\)90045-1](https://doi.org/10.1016/0025-3227(81)90045-1)
- 29 Castaing, P., Philipps, I., Weber, O., 1982. Répartition et dispersion des suspensions dans les
30 eaux du plateau continental aquitain. *Oceanol. Acta* 5, 85–96.
31 <https://doi.org/https://archimer.ifremer.fr/doc/00121/23193/>
- 32 Cathalot, C., Rabouille, C., Pastor, L., Deflandre, B., Viollier, E., Buscail, R., Grémare, A.,
33 Treignier, C., Pruski, A., 2010. Temporal variability of carbon recycling in coastal
34 sediments influenced by rivers: Assessing the impact of flood inputs in the Rhône River
35 prodelta. *Biogeosciences* 7, 1187–1205. <https://doi.org/10.5194/bg-7-1187-2010>
- 36 Cathalot, C., Rabouille, C., Tisnérat-Laborde, N., Toussaint, F., Kerhervé, P., Buscail, R.,
37 Loftis, K., Sun, M.Y., Tronczynski, J., Azoury, S., Lansard, B., Treignier, C., Pastor, L.,
38 Tesi, T., 2013. The fate of river organic carbon in coastal areas: A study in the Rhône
39 River delta using multiple isotopic ($\delta^{13}\text{C}$, $\delta^{14}\text{C}$) and organic tracers. *Geochim.*
40 *Cosmochim. Acta* 118, 33–55. <https://doi.org/10.1016/j.gca.2013.05.001>

- 1 Cirac, P., Berne, S., Castaing, P., Weber, O., 2000. Processus de mise en place et d'évolution
2 de la couverture sédimentaire superficielle de la plate-forme nord-aquitaine. *Oceanol.*
3 *Acta* 23, 663–686. [https://doi.org/10.1016/s0399-1784\(00\)00110-9](https://doi.org/10.1016/s0399-1784(00)00110-9)
- 4 Clarke, K.R., Somerfield, P.J., Gorley, R.N., 2008. Testing of null hypotheses in exploratory
5 community analyses: similarity profiles and biota-environment linkage. *J. Exp. Mar. Bio.*
6 *Ecol.* 366, 56–69. <https://doi.org/10.1016/j.jembe.2008.07.009>
- 7 Clarke, K.R., Warwick, R.M., 2001. *Change in Marine Communities: An Approach to*
8 *Statistical Analysis and Interpretation*, 2nd edition. PRIMER-E, Plymouth.
- 9 Constantin, S., Doxaran, D., Derkacheva, A., Novoa, S., Lavigne, H., 2018. Multi-temporal
10 dynamics of suspended particulate matter in a macro-tidal river Plume (the Gironde) as
11 observed by satellite data. *Estuar. Coast. Shelf Sci.* 202, 172–184.
12 <https://doi.org/10.1016/j.ecss.2018.01.004>
- 13 Coplen, T.B., 2011. Guidelines and recommended terms for expression of stable- isotope-ratio
14 and gas-ratio measurement results. *Rapid Commun. Mass Spectrom* 25, 2538–2560.
15 <https://doi.org/10.1002/rcm.5129>
- 16 Deflandre, B., 2018a. JERICOBENT-3 croise, RV Côtes De La Manche.
17 <https://doi.org/10.17600/18000469>
- 18 Deflandre, B., 2018b. JERICOBENT-4 croise, RV Côtes De La Manche.
19 <https://doi.org/10.17600/18000470>
- 20 Deflandre, B., 2017. JERICOBENT-2 croise, RV Côtes De La Manche.
21 <https://doi.org/10.17600/17011000>
- 22 Deflandre, B., 2016. JERICOBENT-1 croise, RV Côtes De La Manche.
23 <https://doi.org/10.17600/16010400>
- 24 Diaz, M., Grasso, F., Le Hir, P., Sottolichio, A., Caillaud, M., Thouvenin, B., 2020. Modeling
25 Mud and Sand Transfers Between a Macrotidal Estuary and the Continental Shelf:
26 Influence of the Sediment Transport Parameterization. *J. Geophys. Res. Ocean.* 125.
27 <https://doi.org/10.1029/2019JC015643>
- 28 Dobbs, F.C., Vozarik, J.M., 1983. Immediate effects of a storm on coastal infauna. *Mar. Ecol.*
29 *Prog. Ser.* 11, 273–279. <https://doi.org/https://doi.org/10.3354/meps011273>
- 30 Doxaran, D., Froidefond, J.M., Castaing, P., Babin, M., 2009. Dynamics of the turbidity
31 maximum zone in a macrotidal estuary (the Gironde, France): Observations from field
32 and MODIS satellite data. *Estuar. Coast. Shelf Sci.* 81, 321–332.
33 <https://doi.org/10.1016/j.ecss.2008.11.013>
- 34 Dubosq, N., Schmidt, S., Walsh, J.P., Grémare, A., Gillet, H., Lebleu, P., Poirier, D., Perello,
35 M.C., Lamarque, B., Deflandre, B., 2021. A first assessment of organic carbon burial in
36 the West Gironde Mud Patch (Bay of Biscay). *Cont. Shelf Res.* 221.
37 <https://doi.org/10.1016/j.csr.2021.104419>
- 38 Etcheber, H., Relexans, J.C., Beliard, M., Weber, O., Buscail, R., Heussner, S., 1999.
39 *Distribution and quality of sedimentary organic matter on the Aquitanian margin (Bay of*

- 1 Biscay). *Deep. Res. Part II* 46, 2249–2288. <https://doi.org/10.1016/S0967->
2 0645(99)00062-4
- 3 Fontugne, M.R., Jouanneau, J.M., 1987. Modulation of the particulate organic carbon flux to
4 the ocean by a macrotidal estuary: Evidence from measurements of carbon isotopes in
5 organic matter from the Gironde system. *Estuar. Coast. Shelf Sci.* 24, 377–387.
6 [https://doi.org/10.1016/0272-7714\(87\)90057-6](https://doi.org/10.1016/0272-7714(87)90057-6)
- 7 Gadel, F., Jouanneau, J.M., Weber, O., Serve, L., Comellas, L., 1997. Traceurs organiques dans
8 les dépôts de la vasière Ouest-Gironde (Golfe de Gascogne). *Oceanol. Acta* 20, 687–695.
9 <https://doi.org/https://archimer.ifremer.fr/doc/00093/20426/>
- 10 Glémarec, M., 1978. Problèmes d'écologie dynamique et de succession en baie de Concarneau.
11 *Vie Milieu* 1–20.
- 12 Goñi, M.A., Ruttenberg, K.C., Eglinton, T.I., 1998. A reassessment of the sources and
13 importance of land-derived organic matter in surface sediments from the Gulf of Mexico.
14 *Geochim. Cosmochim. Acta* 62, 3055–3075. <https://doi.org/10.1016/S0016->
15 7037(98)00217-8
- 16 Gordon, E.S., Goñi, M.A., Roberts, Q.N., Kineke, G.C., Allison, M.A., 2001. Organic matter
17 distribution and accumulation on the inner Louisiana shelf west of the Atchafalaya River.
18 *Cont. Shelf Res.* 21, 1691–1721. [https://doi.org/10.1016/S0278-4343\(01\)00021-8](https://doi.org/10.1016/S0278-4343(01)00021-8)
- 19 Grasso, F., Verney, R., Le Hir, P., Thouvenin, B., Schulz, E., Kervella, Y., Khojasteh Pour
20 Fard, I., Lemoine, J.P., Dumas, F., Garnier, V., 2018. Suspended Sediment Dynamics in
21 the Macrotidal Seine Estuary (France): 1. Numerical Modeling of Turbidity Maximum
22 Dynamics. *J. Geophys. Res. Ocean.* 123, 558–577. <https://doi.org/10.1002/2017JC013185>
- 23 Grémare, A., Gutiérrez, D., Anschutz, P., Amouroux, J.M., Deflandre, B., Vétion, G., 2005.
24 Spatio-temporal changes in totally and enzymatically hydrolyzable amino acids of
25 superficial sediments from three contrasted areas. *Prog. Oceanogr.* 65, 89–111.
26 <https://doi.org/10.1016/j.pocean.2005.02.016>
- 27 Harmelin-Vivien, M.L., Bănar, D., Dierking, J., Hermand, R., Letourneur, Y., Salen-Picard,
28 C., 2009. Linking benthic biodiversity to the functioning of coastal ecosystems subjected
29 to river runoff (NW Mediterranean). *Anim. Biodivers. Conserv.* 32, 135–145.
30 <https://doi.org/https://doi.org/10.32800/abc.2009.32.0135>
- 31 Harrell, F.E., 2021. Hmisc: Harrell Miscellaneous. R Packag. version 4-5-0.
- 32 Hedges, J.I., Keil, R.G., 1995. Sedimentary organic matter preservation: an assessment and
33 speculative synthesis. *Mar. Chem.* 49, 81–115. <https://doi.org/10.1016/0304->
34 4203(95)00008-F
- 35 Herbland, A., Delmas, D., Laborde, P., Sautour, B., Artigas, F., 1998. Phytoplankton spring
36 bloom of the Gironde plume waters in the Bay of Biscay: Early phosphorus limitation and
37 food-web consequences. *Oceanol. Acta* 21, 279–291. <https://doi.org/10.1016/S0399->
38 1784(98)80015-7
- 39 Hiscock, K., 1984. Rocky shore surveys of the Isles of Scilly. March 27th to April 1st and July
40 7th to 15th 1983. Peterbrgh. Nat. Conserv. Council. CSD Rep. 509.

- 1 Jalón-Rojas, I., Sottolichio, A., Hanquiez, V., Fort, A., Schmidt, S., 2018. To what extent
2 multidecadal changes in morphology and fluvial discharge impact tide in a convergent
3 (turbid) tidal river. *J. Geophys. Res. Ocean.* 123, 3241–3258.
4 <https://doi.org/10.1002/2017JC013466>
- 5 Jouanneau, J.M., Weber, O., Latouche, C., Vernet, J.P., Dominik, J., 1989. Erosion, non-
6 deposition and sedimentary processes through a sedimentological and radioisotopic study
7 of surficial deposits from the “Ouest-Gironde vasière” (Bay of Biscay). *Cont. Shelf Res.*
8 9, 325–342. [https://doi.org/10.1016/0278-4343\(89\)90037-X](https://doi.org/10.1016/0278-4343(89)90037-X)
- 9 Keil, R.G., Mayer, L.M., Quay, P.D., Richey, J.E., Hedges, J.I., 1997. Loss of organic matter
10 from riverine particles in deltas. *Geochim. Cosmochim. Acta* 61, 1507–1511.
11 [https://doi.org/10.1016/S0016-7037\(97\)00044-6](https://doi.org/10.1016/S0016-7037(97)00044-6)
- 12 Keil, R.G., Tsamakis, E., Giddings, J.C., Hedges, J.I., 1998. Biochemical distributions (amino
13 acids, neutral sugars, and lignin phenols) among size-classes of modern marine sediments
14 from the Washington coast. *Geochim. Cosmochim. Acta* 62, 1347–1364.
15 [https://doi.org/10.1016/S0016-7037\(98\)00080-5](https://doi.org/10.1016/S0016-7037(98)00080-5)
- 16 Labry, C., Herbland, A., Delmas, D., 2002. The role of phosphorus on planktonic production
17 of the Gironde plume waters in the Bay of Biscay. *J. Plankton Res.* 24, 97–117.
18 <https://doi.org/10.1093/plankt/24.2.97>
- 19 Lamarque, B., Deflandre, B., Dalto, A.G., Schmidt, S., Romero-Ramirez, A., Garabetian, F.,
20 Dubosq, N., Diaz, M., Grasso, F., Sottolichio, A., Bernard, G., Gillet, H., Cordier, M.A.,
21 Poirier, D., Lebleu, P., Derriennic, H., Danilo, M., Tenório, M.M.B., Grémare, A., 2021.
22 Spatial distributions of surface sedimentary organics and sediment profile image
23 characteristics in a high-energy temperate marine riomar: The west gironde mud patch. *J.*
24 *Mar. Sci. Eng.* 9, 242. <https://doi.org/10.3390/jmse9030242>
- 25 Lansard, B., Rabouille, C., Denis, L., Grenz, C., 2009. Benthic remineralization at the land-
26 ocean interface: A case study of the Rhône River (NW Mediterranean Sea). *Estuar. Coast.*
27 *Shelf Sci.* 81, 544–554. <https://doi.org/10.1016/j.ecss.2008.11.025>
- 28 Lazure, P., Dumas, F., 2008. An external-internal mode coupling for a 3D hydrodynamical
29 model for applications at regional scale (MARS). *Adv. Water Resour.* 31, 233–250.
30 <https://doi.org/10.1016/j.advwatres.2007.06.010>
- 31 Lesueur, P., Jouanneau, J.M., Boust, D., Tastet, J.P., Weber, O., 2001. Sedimentation rates and
32 fluxes in the continental shelf mud fields in the Bay of Biscay (France). *Cont. Shelf Res.*
33 21, 1383–1401. [https://doi.org/10.1016/S0278-4343\(01\)00004-8](https://doi.org/10.1016/S0278-4343(01)00004-8)
- 34 Lesueur, P., Tastet, J.P., Weber, O., Sinko, J.A., 1991. Modèle faciologique d’un corps
35 sédimentaire pélagique de plate-forme la vasière Ouest-Gironde (France). *Oceanol. Acta*
36 11, 143–153. <https://doi.org/https://archimer.ifremer.fr/doc/00267/37871/>
- 37 Lesueur, P., Tastet, J.P., 1994. Facies, internal structures and sequences of modern Gironde-
38 derived muds on the Aquitaine inner shelf, France. *Mar. Geol.* 120, 267–290.
39 [https://doi.org/10.1016/0025-3227\(94\)90062-0](https://doi.org/10.1016/0025-3227(94)90062-0)
- 40 Lesueur, P., Tastet, J.P., Marambat, L., 1996. Shelf mud fields formation within historical

- 1 times: Examples from offshore the Gironde estuary, France. *Cont. Shelf Res.* 16, 1849–
2 1870. [https://doi.org/10.1016/0278-4343\(96\)00013-1](https://doi.org/10.1016/0278-4343(96)00013-1)
- 3 Lesueur, P., Tastet, J.P., Weber, O., 2002. Origin and morphosedimentary evolution of fine-
4 grained modern continental shelf deposits: The Gironde mud fields (Bay of Biscay,
5 France). *Sedimentology* 49, 1299–1320. <https://doi.org/10.1046/j.1365-3091.2002.00498.x>
- 7 Levin, L.A., Boesch, D.F., Covich, A., Dahm, C., Erséus, C., Ewel, K.C., Kneib, R.T.,
8 Moldenke, A., Palmer, M.A., Snelgrove, P., Strayer, D., Weslawski, J.M., 2001. The
9 function of marine critical transition zones and the importance of sediment biodiversity.
10 *Ecosystems* 4, 430–451. <https://doi.org/10.1007/s10021-001-0021-4>
- 11 Longère, P., Dorel, D., 1970. Etude des sédiments meubles de la vasière de la Gironde et des
12 régions avoisinantes. *Rev. des Trav. l'Institut des Pêches Marit.* 34, 233–256.
- 13 Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell,
14 S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion, Degradation, and
15 Recovery Potential of Estuaries and Coastal Seas. *Science.* 312, 1806–1809.
16 <https://doi.org/10.1126/science.1128035>
- 17 Marion, C., Dufois, F., Arnaud, M., Vella, C., 2010. In situ record of sedimentary processes
18 near the Rhône River mouth during winter events (Gulf of Lions, Mediterranean Sea).
19 *Cont. Shelf Res.* 30, 1095–1107. <https://doi.org/10.1016/j.csr.2010.02.015>
- 20 Massé, C., Meisterhans, G., Deflandre, B., Bachelet, G., Bourasseau, L., Bichon, S., Ciutat, A.,
21 Jude-Lemeilleur, F., Lavesque, N., Raymond, N., Grémare, A., Garabetian, F., 2016.
22 Bacterial and macrofaunal communities in the sediments of the West Gironde Mud Patch,
23 Bay of Biscay (France). *Estuar. Coast. Shelf Sci.* 179, 189–200.
24 <https://doi.org/10.1016/j.ecss.2016.01.011>
- 25 Masselink, G., Castelle, B., Scott, T., Dodet, G., Suanez, S., Jackson, D., Floc'h, F., 2016.
26 Extreme wave activity during 2013/2014 winter and morphological impacts along the
27 Atlantic coast of Europe. *Geophys. Res. Lett.* 43, 2135–2143.
28 <https://doi.org/10.1002/2015GL067492>.Received
- 29 Mayer, L.M., 1994a. Relationships between mineral surfaces and organic carbon
30 concentrations in soils and sediments. *Chem. Geol.* 114, 347–363.
31 [https://doi.org/10.1016/0009-2541\(94\)90063-9](https://doi.org/10.1016/0009-2541(94)90063-9)
- 32 Mayer, L.M., 1994b. Surface area control of organic carbon accumulation in continental shelf
33 sediments. *Geochim. Cosmochim. Acta* 58, 1271–1284.
34 [https://doi.org/https://doi.org/10.1016/0016-7037\(94\)90381-6](https://doi.org/https://doi.org/10.1016/0016-7037(94)90381-6)
- 35 Mayer, L.M., Linda L., S., Sawyer, T., Plante, C.J., Jumars, P.A., Sel, R.L., 1995. Bioavailable
36 amino acids in sediments: A biomimetic, kinetics based approach. *Limnol. Oceanogr.* 40,
37 511–520. <https://doi.org/10.4319/lo.1995.40.3.0511>
- 38 McKee, B.A., Aller, R.C., Allison, M.A., Bianchi, T.S., Kineke, G.C., 2004. Transport and
39 transformation of dissolved and particulate materials on continental margins influenced
40 by major rivers: Benthic boundary layer and seabed processes. *Cont. Shelf Res.* 24, 899–

- 1 926. <https://doi.org/10.1016/j.csr.2004.02.009>
- 2 Medernach, L., Grémare, A., Amoureux, J.M., Colomines, J.C., Vétion, G., 2001. Temporal
3 changes in the amino acid contents of particulate organic matter sedimenting in the Bay
4 of Banyuls-sur-Mer (northwestern Mediterranean). *Mar. Ecol. Prog. Ser.* 214, 55–65.
5 <https://doi.org/10.3354/meps214055>
- 6 Miralles, J., Radakovitch, O., Aloisi, J.C., 2005. ²¹⁰Pb sedimentation rates from the
7 Northwestern Mediterranean margin. *Mar. Geol.* 216, 155–167.
8 <https://doi.org/10.1016/j.margeo.2005.02.020>
- 9 Neveux, J., Lantoiné, F., 1993. Spectrofluorometric assay of chlorophylls and phaeopigments
10 using the least squares approximation technique. *Deep. Res. Part I* 40, 1747–1765.
11 [https://doi.org/10.1016/0967-0637\(93\)90030-7](https://doi.org/10.1016/0967-0637(93)90030-7)
- 12 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R.,
13 O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.,
14 2019. *vegan: Community Ecology Package*. R Packag. version 2-5-6.
- 15 Parra, M., Castaing, P., Jouanneau, J.M., Grousset, F., Latouche, C., 1998. Nd-Sr isotopic
16 composition of present-day sediments from the Gironde Estuary, its draining basins and
17 the WestGironde mud patch (SW France). *Cont. Shelf Res.* 19, 135–150.
18 [https://doi.org/10.1016/S0278-4343\(98\)00083-1](https://doi.org/10.1016/S0278-4343(98)00083-1)
- 19 Pastor, L., Deflandre, B., Viollier, E., Cathalot, C., Metzger, E., Rabouille, C., Escoubeyrou,
20 K., Lloret, E., Pruski, A.M., Vétion, G., Desmalades, M., Buscail, R., Grémare, A., 2011.
21 Influence of the organic matter composition on benthic oxygen demand in the Rhône
22 River prodelta (NW Mediterranean Sea). *Cont. Shelf Res.* 31, 1008–1019.
23 <https://doi.org/10.1016/j.csr.2011.03.007>
- 24 Pastor, L., Rabouille, C., Metzger, E., Thibault de Chanvalon, A., Viollier, E., Deflandre, B.,
25 2018. Transient early diagenetic processes in Rhône prodelta sediments revealed in
26 contrasting flood events. *Cont. Shelf Res.* 166, 65–76.
27 <https://doi.org/10.1016/j.csr.2018.07.005>
- 28 Pearson, T.H., Rosenberg, R., 1978. Macrobenthos succession in relation to organic enrichment
29 and pollution of the marine environment. *Oceanogr. Mar. Biol. Annu. Rev.* 16, 229–331.
- 30 Picton, B.E., Emblow, C.S., Morrow, C.C., Sides, E.M., Costello, M.J., 1994. Marine
31 communities of the Mulroy Bay and Lough Swill area, north-west Ireland, with an
32 assessment of their nature conservation importance. *Environ. Sci. Unit, Trinity Coll. (f.*
33 *Surv. Report)*.
- 34 R Core Team, 2019. *R: A Language and Environment for Statistical Computing*. R Foundation
35 for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- 36 Rees, E.I.S., Nicholaidou, A., Laskaridou, P., 1977. The Effects of Storms on the Dynamics of
37 Shallow Water Benthic Associations. *Biol. Benthic Org.* 465–474.
38 <https://doi.org/10.1016/b978-0-08-021378-1.50052-x>
- 39 Relexans, J.C., Lin, R.G., Castel, J., Etcheber, H., Laborde, P., 1992. Response of biota to
40 sedimentary organic matter quality of the west Gironde mud patch, Bay of Biscay

- 1 (France). *Oceanol. Acta* 15, 639–649.
2 <https://doi.org/https://archimer.ifremer.fr/doc/00101/21231/>
- 3 Rhoads, D.C., Boesch, D.F., Zhican, T., Fengshan, X., Liqiang, H., Nilsen, K.J., 1985.
4 Macrobenthos and sedimentary facies on the Changjiang delta platform and adjacent
5 continental shelf, East China Sea. *Cont. Shelf Res.* 4, 189–213.
6 [https://doi.org/10.1016/0278-4343\(85\)90029-9](https://doi.org/10.1016/0278-4343(85)90029-9)
- 7 Roland, A., Ardhuin, F., 2014. On the developments of spectral wave models: Numerics and
8 parameterizations for the coastal ocean. *Ocean Dyn.* 64, 833–846.
9 <https://doi.org/10.1007/s10236-014-0711-z>
- 10 Soulsby, R., 1997. *Dynamics of Marine Sands: A Manual for Practical Applications*, Thomas
11 Tel. ed.
- 12 Tesi, T., Miserocchi, S., Goñi, M.A., Langone, L., 2007. Source, transport and fate of terrestrial
13 organic carbon on the western Mediterranean Sea, Gulf of Lions, France. *Mar. Chem.* 105,
14 101–117. <https://doi.org/10.1016/j.marchem.2007.01.005>
- 15 Ulses, C., Estournel, C., Durrieu de Madron, X., Palanques, A., 2008. Suspended sediment
16 transport in the Gulf of Lions (NW Mediterranean): Impact of extreme storms and floods.
17 *Cont. Shelf Res.* 28, 2048–2070. <https://doi.org/10.1016/j.csr.2008.01.015>
- 18 Wakeham, S.G., Lee, C., Hedges, J.I., Hernes, P.J., Peterson, M.L., 1997. Molecular indicators
19 of diagenetic status in marine organic matter. *Geochim. Cosmochim. Acta* 61, 5363–5369.
20 <https://doi.org/10.5833/jjgs.33.70>
- 21 Weber, O., Jouanneau, J.M., Ruch, P., Mirmand, M., 1991. Grain-size relationship between
22 suspended matter originating in the Gironde estuary and shelf mud-patch deposits. *Mar.*
23 *Geol.* 96, 159–165. [https://doi.org/10.1016/0025-3227\(91\)90213-N](https://doi.org/10.1016/0025-3227(91)90213-N)
- 24 Wheatcroft, R.A., 2006. Time-series measurements of macrobenthos abundance and sediment
25 bioturbation intensity on a flood-dominated shelf. *Prog. Oceanogr.* 71, 88–122.
26 <https://doi.org/10.1016/j.pcean.2006.06.002>
- 27 Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Jackson, J.B.C.,
28 Lotze, H.K., Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson,
29 R., 2006. Impacts of biodiversity loss on ocean ecosystem services. *Science* 314, 787–
30 790. <https://doi.org/10.1126/science.1132294>
- 31 Zuo, Z., Eisma, D., Berger, G.W., 1991. Determination of sediment accumulation and mixing
32 rates in the Gulf of Lions, Mediterranean Sea. *Oceanol. Acta* 14, 253–262.
33 <https://doi.org/https://archimer.ifremer.fr/doc/00101/21255/>