



HAL
open science

Existence of minimizers for the Dirac-Fock model of crystals

Isabelle Catto, Long Meng, Eric Paturel, Eric Séré

► **To cite this version:**

Isabelle Catto, Long Meng, Eric Paturel, Eric Séré. Existence of minimizers for the Dirac-Fock model of crystals. *Archive for Rational Mechanics and Analysis*, 2024, 248 (issue 4, article 63), 10.1007/s00205-024-01988-8 . hal-03871041v5

HAL Id: hal-03871041

<https://hal.science/hal-03871041v5>

Submitted on 30 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Existence of minimizers for the Dirac–Fock model of crystals

Isabelle Catto* Long Meng† Éric Paturel‡ Éric Séré§

Abstract

Whereas many different models exist in the mathematical and physical literature for ground states of non-relativistic crystals, the relativistic case has been much less studied and we are not aware of any mathematical result on a fully relativistic treatment of crystals. In this paper, we introduce a mean-field relativistic energy for crystals in terms of periodic density matrices. This model is inspired both from a recent definition of the Dirac–Fock ground state for atoms and molecules, due to one of us, and from the non-relativistic Hartree–Fock model for crystals. We prove the existence of a ground state when the number of electrons per cell is not too large.

1 Introduction

For solids with heavy atoms, relativistic shifts may affect the bonding properties and the optical properties. It is shown in [37] that the yellow color of gold is a result of relativistic effects. Furthermore, by studying the relativistic band structure in solids, it is shown in [11, 12] that the relativistic shifts of the $5d$ bands relative to the $s - p$ bands in gold change the main interband edge by more than 1 eV.

A natural way to build quantum models for the crystal phase is to consider the so-called thermodynamic limit of quantum molecular models. Roughly speaking it consists in considering a finite but large piece of an (infinite and neutral) crystal. The thermodynamic law predicts that the ground state energy of the obtained large neutral molecule is proportional to the volume of this finite piece (which turns out to be also proportional to the total number of particles composing the molecule). The energy for the whole crystal is then identified with the limit—if it exists—of the energy per unit volume (or equivalently per particle) of the large molecule when the size of the considered piece goes to infinity. This method was applied successfully by different authors for several well-known models from quantum chemistry [9, 8, 6, 32]—see also [7] for a review—but always for non-relativistic crystals.

The Dirac–Fock model (DF) was introduced in atomic physics by Swirls [41] in 1935. It is widely used in relativistic quantum chemistry, and gives numerical results on atoms and molecules in excellent agreement with experimental data [13, 20, 30]. Its relation with QED was investigated by Mittleman [35]. Mittleman’s approach was studied mathematically in [2–4, 18, 34]. To our knowledge the Dirac–Fock model has not been extended to crystals: there exist fully relativistic treatments of crystals in the physics literature, but they use the Kohn–Sham approach (see [15, 28] and the references therein).

The first rigorous existence results for the atomic and molecular Dirac–Fock equations were obtained in [16, 36]. Compared to the non-relativistic models, the situation is different: the existence of bound states has only been proved when the total number of protons does not exceed 124, for the physical value $\alpha \approx 1/137$ of the fine structure constant. Moreover, the Dirac–Fock energy functional is strongly indefinite and the notion of ground state has to be handled very carefully [16]. These difficulties exclude a thermodynamic limit approach to derive the Dirac–Fock model for crystals.

In [17] it was shown that certain solutions of the (relativistic) Dirac–Fock equations converge towards the energy-minimizing solutions of the (non-relativistic) Hartree–Fock equations when the speed of light tends to infinity. This validates *a posteriori* the notions of ground state solutions and ground state energy for the Dirac–Fock equations. In the approach of [17], the multi-electronic state

*ISABELLE CATTO, CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, UNIVERSITÉ PSL, CNRS, 75016 PARIS, FRANCE
E-mail address: catto@ceremade.dauphine.fr

†LONG MENG, CERMICS, ÉCOLE DES PONTS PARISTECH, 6 AND 8 AV. PASCAL, 77455 MARNE-LA-VALLÉE, FRANCE
E-mail address: long.meng@enpc.fr

‡ÉRIC PATUREL, LABORATOIRE DE MATHÉMATIQUES J. LERAY (UMR CNRS 6629), UNIVERSITÉ DE NANTES, 44322 NANTES, FRANCE
E-mail address: eric.paturel@univ-nantes.fr

§ÉRIC SÉRÉ, CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, UNIVERSITÉ PSL, CNRS, 75016 PARIS, FRANCE
E-mail address: sere@ceremade.dauphine.fr

is modeled by a Slater determinant of mono-electronic wavefunctions. On the other hand, Huber and Siedentop use a density matrix formulation and a fixed-point iteration to define and construct ground states of the Dirac–Fock model [27]. Unfortunately their assumptions do not cover the physical value of α . Recently, in [39] one of us gave a new definition and an existence proof for the ground state of the Dirac–Fock model in atoms and molecules, under assumptions covering the physical value of α , thanks to a density matrix formulation and a retraction technique combined with a minimization principle. Inspired by this work and by the analysis of the periodic Hartree–Fock model due to Le Bris, Lions, and one of us [8], we propose a definition for the ground state of the Dirac–Fock model for crystals which is a relativistic analogue of Lieb’s variational principle for the Hartree–Fock model [1, 31], and we prove the existence of minimizers. In addition, we show that these minimizers solve a self-consistent equation. Our method can be used to calculate the ground state of neutral crystals with at most 17 electrons per cell. However, some estimates used in this paper are not optimal, and we strongly believe that this limiting bound can be improved.

The minimization problem under consideration in this paper combines several difficulties related to compactness issues. The Dirac operator, hence the Dirac–Fock energy functional, is not bounded from below and the kinetic energy term is of the same order as the Coulomb-type potential energy terms, a standard feature of Coulomb–Dirac–Fock type models. Our proof of existence of minimizers for crystals is neither a straightforward adaptation of the one for atoms and molecules in [39] nor of the one for crystals in Hartree–Fock theory in [8]: a major issue arises from the compactness of the density matrices and of the self-consistent operators in the momentum variable ξ , resulting from the Bloch decomposition of the space. Compactness in the momentum variable is crucial to deal with the (non-linear) exchange term in the DF periodic functional and with the nonlinear constraint ensuring that the electrons lie in the positive spectral subspace of the self-consistent periodic Dirac–Fock operator. Our results rely on a careful analysis of the periodic exchange potential. In passing, we have corrected some wrong estimates on the exchange term in [8] and improved the regularity results therein (see Appendix B). Furthermore, we provide an asymptotically optimal constant for the Hardy inequality associated with the periodic Coulomb potential that is new in the literature, as far as we know.

In addition, compared with existing results for crystals’ ground state energy, such as the Hartree–Fock model [8], we provide a new method to prove the existence of minimizers for crystals: based on the spectral analysis of the self-consistent periodic DF operator, we build minimizing sequences that feature both a uniform dependence with respect to the momentum ξ and a better regularity in the space variables, and we rely on it to improve the relative compactness of subsequences in the periodic energy space.

Before ending this section, let us mention the Bogoliubov–Dirac–Fock (BDF) model proposed by Chaix and Iracane in [10] as an alternative to Dirac–Fock for heavy atoms and molecules, and later studied in a series of mathematical works (see the review paper [25] and references therein, see also the more recent works [21, 22, 24]). Compared with DF, the BDF model has several advantages: the corresponding energy is bounded from below, so the notion of ground state becomes straightforward; vacuum polarization effects are taken into account; the derivation of the model as a mean-field approximation of no-photon QED is more convincing. However the mathematical definition of the BDF energy involves a rather complex functional framework as well as an ultraviolet regularization, and a renormalization procedure is needed to interpret the equations. Thus, in the present work we restrict ourselves to the conceptually simpler DF model, and the study of relativistic crystals in the BDF approximation is left for future research.

2 General setting of the model and main result

2.1 Preliminaries and functional framework

Throughout the paper, we choose units for which $m = c = \hbar = 1$, where m is the mass of the electron, c the speed of light and \hbar the Planck constant. For the sake of simplicity, we only consider the case of a cubic crystal with a single point-like nucleus per unit cell, which is located at the center of the cell. The reader should however keep in mind that the general case could be handled as well. Let $\ell > 0$ denote the length of the elementary cell $Q_\ell = (-\frac{\ell}{2}, \frac{\ell}{2}]^3$. The nuclei with positive charge z are treated as classical particles with infinite mass that are located at each point of the lattice $\ell\mathbb{Z}^3$. The electrons are treated quantum mechanically through a periodic density matrix. The electronic density is modeled by a Q_ℓ -periodic function whose L^1 norm over the elementary cell equals the “number of

electrons" q (the electrons' charge per cell is equal to $-q$). When $q = z$, electrical neutrality per cell is ensured.

In this periodic setting, the Q_ℓ -periodic Coulomb potential G_ℓ resulting from a distribution of point particles of charge 1 that are periodically located at the centers of the cubic cells of the lattice is defined, up to a constant, by

$$-\Delta G_\ell = 4\pi \left[-\frac{1}{\ell^3} + \sum_{k \in \mathbb{Z}^3} \delta_{\ell k} \right]. \quad (2.1)$$

By convention, we choose G_ℓ such that

$$\int_{Q_\ell} G_\ell dx = 0. \quad (2.2)$$

The function G_ℓ is actually the Green function of the periodic Laplace operator on Q_ℓ . The Fourier series of G_ℓ writes

$$G_\ell(x) = \frac{1}{\pi\ell} \sum_{p \in \mathbb{Z}^3 \setminus \{0\}} \frac{e^{\frac{2i\pi}{\ell} p \cdot x}}{|p|^2}, \quad \text{for every } x \in \mathbb{R}^3. \quad (2.3)$$

Under the convention (2.2), the periodic Coulomb potential changes sign, but is bounded from below (see Lemma A.1 in Appendix A).

Remark 2.1. *The size ℓ of the unit cell does not play a specific role here. It is however involved in the study of the Hardy-type inequalities for the periodic Coulomb potential (see Section 4.1). When ℓ goes to infinity, one expects to recover the Dirac–Fock model for atoms.*

The free Dirac operator is defined by $D^0 = -i \sum_{r=1}^3 \alpha_r \partial_r + \beta$, with 4×4 complex matrices $\alpha_1, \alpha_2, \alpha_3$ and β , whose standard forms are $\beta = \begin{pmatrix} 1_2 & 0 \\ 0 & -1_2 \end{pmatrix}$, $\alpha_r = \begin{pmatrix} 0 & \sigma_r \\ \sigma_r & 0 \end{pmatrix}$ where 1_2 is the 2×2 identity matrix and the σ_r 's, for $r \in \{1, 2, 3\}$, are the well-known 2×2 Pauli matrices $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.

The operator D^0 acts on 4–spinors; that is, on functions from \mathbb{R}^3 to \mathbb{C}^4 . It is self-adjoint in $L^2(\mathbb{R}^3; \mathbb{C}^4)$, with domain $H^1(\mathbb{R}^3; \mathbb{C}^4)$ and form domain $H^{1/2}(\mathbb{R}^3; \mathbb{C}^4)$ (denoted by L^2 , H^1 and $H^{1/2}$ in the following, when there is no ambiguity). Its spectrum is $\sigma(D^0) = (-\infty, -1] \cup [+1, +\infty)$. Following the notation in [16, 36], we denote by Λ^+ and $\Lambda^- = 1_{L^2} - \Lambda^+$ respectively the two orthogonal projectors on $L^2(\mathbb{R}^3; \mathbb{C}^4)$ corresponding to the positive and negative eigenspaces of D^0 ; that is

$$\begin{cases} D^0 \Lambda^+ = \Lambda^+ D^0 = \Lambda^+ \sqrt{1 - \Delta} = \sqrt{1 - \Delta} \Lambda^+; \\ D^0 \Lambda^- = \Lambda^- D^0 = -\Lambda^- \sqrt{1 - \Delta} = -\sqrt{1 - \Delta} \Lambda^-. \end{cases}$$

According to the Floquet theory [38], the underlying Hilbert space $L^2(\mathbb{R}^3; \mathbb{C}^4)$ is unitarily equivalent to $L^2(Q_\ell^*) \otimes L^2(Q_\ell; \mathbb{C}^4)$, where $Q_\ell^* = [-\frac{\pi}{\ell}, \frac{\pi}{\ell}]^3$ is the so-called reciprocal cell of the lattice, with volume $|Q_\ell^*| = (2\pi)^3/\ell^3$ (in the physics literature Q_ℓ^* is known as the first Brillouin zone). The Floquet unitary transform $U : L^2(\mathbb{R}^3; \mathbb{C}^4) \rightarrow L^2(Q_\ell^*) \otimes L^2(Q_\ell; \mathbb{C}^4)$ is given by

$$(U\phi)_\xi = \sum_{k \in \mathbb{Z}^3} e^{-i\ell k \cdot \xi} \phi(\cdot + \ell k) \quad (2.4)$$

for every $\xi \in Q_\ell^*$ and ϕ in $L^2(\mathbb{R}^3; \mathbb{C}^4)$. For every $\xi \in Q_\ell^*$, the function $(U\phi)_\xi$ belongs to the space

$$L_\xi^2(Q_\ell; \mathbb{C}^4) = \{ \psi \in L_{\text{loc}}^2(\mathbb{R}^3; \mathbb{C}^4) \mid e^{-i\xi \cdot x} \psi \text{ is } Q_\ell\text{-periodic} \},$$

which will be denoted by L_ξ^2 in the sequel. Functions ψ of this form are called Bloch waves or Q_ℓ -quasi-periodic functions with quasi-momentum $\xi \in Q_\ell^*$. They satisfy

$$\psi(\cdot + \ell k) = e^{i\ell k \cdot \xi} \psi(\cdot), \quad \text{for every } k \in \mathbb{Z}^3.$$

For any function $\phi_\xi \in L^2_\xi$, using the definition of Fourier series expansion for Q_ℓ -periodic functions, we write

$$\phi_\xi(x) = \sum_{k \in \mathbb{Z}^3} \widehat{\phi}_\xi(k) e^{i(\frac{2\pi}{\ell}k + \xi) \cdot x}, \quad \text{a.e. } x \in \mathbb{R}^3, \quad (2.5)$$

with coefficients

$$\widehat{\phi}_\xi(k) = \frac{1}{\ell^3} \int_{Q_\ell} \phi_\xi(y) e^{-i(\frac{2\pi}{\ell}k + \xi) \cdot y} dy \in \mathbb{C}^4.$$

The Hilbert space L^2_ξ is endowed with the norm

$$\|\phi\|_{L^2_\xi} := \left(\ell^3 \sum_{k \in \mathbb{Z}^3} |\widehat{\phi}_\xi(k)|^2 \right)^{1/2} = \left(\int_{Q_\ell} |\phi_\xi(x)|^2 dx \right)^{1/2} = \|\phi_\xi\|_{L^2(Q_\ell)}.$$

Here, and in the whole paper, we use the same notation $|\cdot|$ for the canonical Euclidean norm in \mathbb{R}^n , \mathbb{C}^n or $\mathcal{M}_n(\mathbb{C})$. When applied to self-adjoint operators, $|T|$ means the absolute value of T .

For every real number s , we also define

$$H^s_\xi(Q_\ell; \mathbb{C}^4) := L^2_\xi(Q_\ell; \mathbb{C}^4) \cap H^s_{\text{loc}}(\mathbb{R}^3; \mathbb{C}^4)$$

endowed with the norm

$$\|\phi_\xi\|_{H^s_\xi} = \left(\ell^3 \sum_{k \in \mathbb{Z}^3} \left(1 + \left| \frac{2\pi}{\ell}k + \xi \right|^2 \right)^s |\widehat{\phi}_\xi(k)|^2 \right)^{1/2}.$$

To simplify the notation, we simply write here and below H^s_ξ when there is no ambiguity.

Operators \mathcal{L} on $L^2(\mathbb{R}^3; \mathbb{C}^4)$ that commute with the translations of $\ell\mathbb{Z}^3$ can be decomposed accordingly into a direct integral of operators \mathcal{L}_ξ acting on L^2_ξ and defined by

$$\mathcal{L}_\xi(U\phi)_\xi = (U\mathcal{L}\phi)_\xi \text{ for every } \phi \in L^2(\mathbb{R}^3; \mathbb{C}^4), \text{ a.e. } \xi \in Q_\ell^* \quad (2.6)$$

(see [38] for more details). We use the notation $\mathcal{L} = \int_{Q_\ell^*}^\oplus \mathcal{L}_\xi d\xi$, with the shorthand \int_Ω for $\frac{1}{|\Omega|} \int_\Omega$, to refer to this decomposition. In particular, for the free Dirac operator D^0 we have

$$D^0 = \int_{Q_\ell^*}^\oplus D_\xi d\xi, \quad (2.7)$$

where the D_ξ 's are self-adjoint operators on L^2_ξ with domains H^1_ξ and form-domains $H^{1/2}_\xi$. Note that $D^2_\xi = 1 - \Delta_\xi$, where $-\Delta = \int_{Q_\ell^*}^\oplus -\Delta_\xi d\xi$. For every function $\phi_\xi \in H^1_\xi$, the operator D_ξ is also defined by

$$D_\xi \phi_\xi(x) = \sum_{k \in \mathbb{Z}^3} \left[\sum_{r=1}^3 \left(\frac{2\pi}{\ell} k_r + \xi_r \right) \cdot \alpha_r + \beta \right] \widehat{\phi}_\xi(k) e^{i(\frac{2\pi}{\ell}k + \xi) \cdot x}.$$

In particular,

$$(\phi_\xi, |D_\xi| \phi_\xi)_{L^2_\xi} = \ell^3 \sum_{k \in \mathbb{Z}^3} \sqrt{1 + \left| \xi + \frac{2\pi}{\ell}k \right|^2} |\widehat{\phi}_\xi(k)|^2. \quad (2.8)$$

For every $\xi \in Q_\ell^*$, the positive spectrum of D_ξ is composed of a non-decreasing sequence of real eigenvalues $(d_j^+(\xi))_{j \geq 1}$ counted with multiplicity. Each function $\xi \mapsto d_j^+(\xi)$ is continuous and Q_ℓ^* -periodic, and one has $d_j^+(Q_\ell^*) \in [c_*(j), c^*(j)]$ with

$$c_*(j) := \min_{\xi \in Q_\ell^*} d_j^+(\xi) \quad \text{and} \quad c^*(j) := \max_{\xi \in Q_\ell^*} d_j^+(\xi). \quad (2.9)$$

Note that

$$c_*(j) \geq 1, \quad \lim_{j \rightarrow +\infty} c_*(j) = +\infty.$$

In the same manner, the negative spectrum of D_ξ is composed of the non-increasing sequence of real eigenvalues $d_j^-(\xi) = -d_j^+(\xi)$. Finally, one has

$$\bigcup_{\xi \in Q_\ell^*} \sigma(D_\xi) = \bigcup_{j \geq 1} [-c^*(j), -c_*(j)] \cup [c_*(j), c^*(j)] = \sigma(D^0) = (-\infty, -1] \cup [+1, +\infty). \quad (2.10)$$

As in the Hartree–Fock model for crystals [8], the electrons will be modeled by an operator on $L^2(\mathbb{R}^3; \mathbb{C}^4)$, called the one-particle density matrix, that reflects their periodic distribution in the nuclei lattice.

We now introduce various functional spaces for linear operators on $L^2(Q_\ell; \mathbb{C}^4)$ and for operators on $L^2(\mathbb{R}^3; \mathbb{C}^4)$ that commute with translations. Let $\mathcal{B}(E)$ be the set of bounded operators from a Banach space E to itself. We use the shorthand $\mathcal{B}(L_\xi^2)$ for $\mathcal{B}(L_\xi^2(Q_\ell; \mathbb{C}^4))$. The space of bounded operators on $\int_{Q_\ell^*}^\oplus L_\xi^2 d\xi = L^2(Q_\ell^*) \otimes L^2(Q_\ell; \mathbb{C}^4)$ which commute with the translations of $\ell\mathbb{Z}^3$ is denoted by Y . It is isomorphic to $L^\infty(Q_\ell^*; \mathcal{B}(L_\xi^2))$. Moreover, for every $h = \int_{Q_\ell^*}^\oplus h_\xi d\xi \in Y$,

$$\|h\|_Y := \operatorname{ess\,sup}_{\xi \in Q_\ell^*} \|h_\xi\|_{\mathcal{B}(L_\xi^2)} = \|h\|_{\mathcal{B}(L^2(\mathbb{R}^3; \mathbb{C}^4))}$$

(see [38, Theorem XIII.83]). For $s \in [1, \infty)$ and $\xi \in Q_\ell^*$, we define

$$\mathfrak{S}_s(\xi) := \left\{ h_\xi \in \mathcal{B}(L_\xi^2) \mid \operatorname{Tr}_{L_\xi^2}(|h_\xi|^s) < \infty \right\}$$

endowed with the norm

$$\|h_\xi\|_{\mathfrak{S}_s(\xi)} := \left(\operatorname{Tr}_{L_\xi^2}(|h_\xi|^s) \right)^{1/s}.$$

We denote by $\mathfrak{S}_\infty(\xi)$ the space of compact operators on L_ξ^2 , endowed with the norm inherited from $\|\cdot\|_{\mathcal{B}(L_\xi^2)}$. Similarly, for $t \in [1, +\infty]$, we define

$$\mathfrak{S}_{s,t} := \left\{ h = \int_{Q_\ell^*}^\oplus h_\xi d\xi \mid h_\xi \in \mathfrak{S}_s(\xi) \text{ a.e. } \xi \in Q_\ell^*, \|h_\xi\|_{\mathfrak{S}_s(\xi)} \in L^t(Q_\ell^*) \right\} \quad (2.11)$$

endowed with the norm

$$\|h\|_{\mathfrak{S}_{s,t}} := \left(\int_{Q_\ell^*} \|h_\xi\|_{\mathfrak{S}_s(\xi)}^t d\xi \right)^{1/t} \quad \text{for } 1 \leq t < +\infty \quad (2.12)$$

and

$$\|h\|_{\mathfrak{S}_{s,\infty}} := \operatorname{ess\,sup}_{\xi \in Q_\ell^*} \|h_\xi\|_{\mathfrak{S}_s(\xi)}. \quad (2.13)$$

In particular $\mathfrak{S}_{\infty,\infty} = L^\infty(Q_\ell^*; \mathfrak{S}_\infty(L_\xi^2)) \subset Y$ is endowed with the norm of Y .

In the sequel of this paper, we work with periodic one-particle density matrices belonging to subspaces $\mathfrak{S}_{1,p}$ of $\mathfrak{S}_{1,1}$, for $1 \leq p \leq +\infty$. On such spaces, we can define the trace per unit cell as

$$\widetilde{\operatorname{Tr}}_{L^2}(h) := \int_{Q_\ell^*} \operatorname{Tr}_{L_\xi^2}(h_\xi) d\xi.$$

Here, $\operatorname{Tr}_{L_\xi^2}$ means the usual trace of operators on the Hilbert space $L_\xi^2(Q_\ell; \mathbb{C}^4)$. It coincides with the trace of the operator with kernel $\operatorname{Tr}_4(h_\xi(\cdot, \cdot))$ on $L_\xi^2(Q_\ell; \mathbb{C})$ with Tr_4 standing for the trace of a 4×4 matrix. The $\widetilde{}$ reminds us that γ is not trace-class on $L^2(\mathbb{R}^3)$.

The trace per unit cell allows to define duality pairings between spaces $\mathfrak{S}_{s,t}$ using the classical duality properties in Schatten's spaces [40]. More precisely, if (s, s') and (t, t') are in $[1, +\infty]^2$ with $1/s + 1/s' = 1$ and $1/t + 1/t' = 1$, then one can define a duality pairing $\langle \cdot, \cdot \rangle$ between $\mathfrak{S}_{s,t}$ and $\mathfrak{S}_{s',t'}$ as follows. For $h \in \mathfrak{S}_{s,t}$ and $h' \in \mathfrak{S}_{s',t'}$, the product hh' is in $\mathfrak{S}_{1,1}$ and one sets

$$\langle h, h' \rangle := \widetilde{\operatorname{Tr}}_{L^2}[hh'].$$

One has

$$|\langle h, h' \rangle| \leq \|hh'\|_{\mathfrak{S}_{1,1}} \leq \|h\|_{\mathfrak{S}_{s,t}} \|h'\|_{\mathfrak{S}_{s',t'}}.$$

We also define

$$X^\alpha(\xi) = \left\{ h \in \mathcal{B}(L_\xi^2) \mid |D_\xi|^{\alpha/2} h_\xi |D_\xi|^{\alpha/2} \in \mathfrak{S}_1(\xi) \right\}$$

endowed with the norm

$$\|h_\xi\|_{X^\alpha(\xi)} = \left\| |D_\xi|^{\alpha/2} h_\xi |D_\xi|^{\alpha/2} \right\|_{\mathfrak{S}_1(\xi)}$$

and

$$X_t^\alpha := \left\{ h = \int_{Q_\ell^*}^\oplus h_\xi d\xi \mid h_\xi \in \mathfrak{S}_1(\xi) \text{ a.e. } \xi \in Q_\ell^*, \left\| |D_\xi|^{\alpha/2} h_\xi |D_\xi|^{\alpha/2} \right\|_{\mathfrak{S}_1(\xi)} \in L^t(Q_\ell^*) \right\}$$

endowed with the norm

$$\|h\|_{X_t^\alpha} := \left\| |D^0|^{\alpha/2} h |D^0|^{\alpha/2} \right\|_{\mathfrak{S}_{1,t}}.$$

For any two functional spaces A and B the norm of the intersected space is defined by

$$\|\gamma\|_{A \cap B} := \max\{\|\gamma\|_A, \|\gamma\|_B\}, \quad \forall \gamma \in A \cap B.$$

For future convenience, we use the notation $X(\xi)$ for $X^1(\xi)$. We also set $X := X_1^1$ and

$$Z := \{\gamma \in X \cap Y \mid \gamma^* = \gamma\}.$$

We endow Z with the norm inherited from $X \cap Y$, that is, we take

$$\|\gamma\|_Z := \max\{\|\gamma\|_X, \|\gamma\|_Y\}, \quad \forall \gamma \in Z.$$

With this norm, Z is a Banach space. The functional spaces $\mathfrak{S}_{1,1}$, X , Y and Z will play an essential role in the whole paper, while the functional space $\mathfrak{S}_{1,\infty}$ and its subspace X_∞^2 are mainly used in Section 6. In addition, we will also use the functional space $\mathfrak{S}_{\infty,1}$ in Section 6 since $\mathfrak{S}_{1,\infty}$ is its dual space.

Definition 2.2 (Periodic one-particle density matrices). *We denote by Γ the following set of Q_ℓ -periodic one-particle density matrices:*

$$\Gamma := \{\gamma \in X \mid \gamma^* = \gamma, \quad 0 \leq \gamma \leq 1_{L^2(\mathbb{R}^3)}\} \subset Z.$$

Remark 2.3. *For $\gamma \in \Gamma$ and for almost every ξ in Q_ℓ^* , the operator γ_ξ is compact on L_ξ^2 and admits a complete set of eigenfunctions $(u_n(\xi, \cdot))_{n \geq 1}$ in L_ξ^2 (actually lying in $H_\xi^{1/2}$), corresponding to a non-decreasing sequence of eigenvalues $0 \leq \mu_n(\xi) \leq 1$ (counted with their multiplicity). This is expressed as*

$$\gamma_\xi = \sum_{n \geq 1} \mu_n(\xi) |u_n(\xi, \cdot)\rangle \langle u_n(\xi, \cdot)|, \quad \langle u_n(\xi, \cdot), u_m(\xi, \cdot) \rangle_{L_\xi^2} = \delta_{n,m} \quad (2.14)$$

where $|u\rangle \langle u|$ denotes the projector onto the vector space spanned by the function u . Equivalently, for almost every ξ in Q_ℓ^* and for any $(x, y) \in \mathbb{R}^3 \times \mathbb{R}^3$, the Hilbert-Schmidt kernel writes

$$\gamma_\xi(x, y) = \sum_{n \geq 1} \mu_n(\xi) u_n(\xi, x) u_n^*(\xi, y). \quad (2.15)$$

In the above equation, $u_n(\xi, \cdot)$ is a column vector with four coefficients and the superscript $*$ refers to transposition composed with complex conjugation of the coefficients. Thus, $\gamma_\xi(x, y)$ is a 4×4 complex matrix, and for every function $\varphi \in L_\xi^2$,

$$(\gamma_\xi \varphi)(x) = \int_{Q_\ell} \gamma_\xi(x, y) \varphi(y) dy = \sum_{n \geq 1} \mu_n(\xi) u_n(\xi, x) \int_{Q_\ell} u_n^*(\xi, y) \varphi(y) dy.$$

By definition of the trace of an operator,

$$\text{Tr}_{L_\xi^2}(\gamma_\xi) = \sum_{n \geq 1} \mu_n(\xi).$$

Definition 2.4 (Integral kernel and electronic density). *Let γ belong to Γ . Then we can define in a unique way an integral kernel $\gamma(\cdot, \cdot) \in L^2(Q_\ell \times \mathbb{R}^3) \cap L^2(\mathbb{R}^3 \times Q_\ell)$ with $\gamma(\cdot + k, \cdot + k) = \gamma(\cdot, \cdot)$ for any $k \in \mathbb{Z}^3$ and a Q_ℓ -periodic density ρ_γ associated to γ by*

$$\gamma(x, y) = \int_{Q_\ell^*} \gamma_\xi(x, y) d\xi \quad (2.16)$$

and

$$\rho_\gamma(x) = \int_{Q_\ell^*} \text{Tr}_4 \gamma_\xi(x, x) d\xi. \quad (2.17)$$

The function ρ_γ is non-negative and belongs to $L^1(Q_\ell; \mathbb{R})$. Indeed, using the decomposition (2.15), we have

$$\rho_\gamma(x) = \int_{Q_\ell^*} \sum_{n=1}^{\infty} \mu_n(\xi) |u_n(\xi, x)|^2 d\xi \quad (2.18)$$

and

$$\int_{Q_\ell} \rho_\gamma(x) dx = \int_{Q_\ell^*} \sum_{n=1}^{\infty} \mu_n(\xi) d\xi = \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}(\gamma_\xi) d\xi.$$

In the physical setting we are interested in, the value of the above integral is the number of electrons per cell.

By the Cauchy–Schwarz inequality, it is easily checked that

$$|\gamma(x, y)|^2 \leq \rho_\gamma(x) \rho_\gamma(y), \quad \text{a.e. } x, y \in \mathbb{R}^3. \quad (2.19)$$

Note that, when h is a Q_ℓ -periodic trace-class operator but is not necessarily a positive operator, we still may define ρ_h with the help of (2.17), but (2.19) becomes $|h(x, y)|^2 \leq \rho_{|h|}(x) \rho_{|h|}(y)$ where $|h| = \sqrt{h^* h}$.

We can now introduce the periodic Dirac–Fock functional.

2.2 The periodic Dirac–Fock model

For $\gamma \in Z$, we define the periodic Dirac–Fock functional

$$\begin{aligned} \mathcal{E}^{DF}(\gamma) &= \int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [D_\xi \gamma_\xi] d\xi - \alpha z \int_{Q_\ell} G_\ell(x) \rho_\gamma(x) dx \\ &\quad + \frac{\alpha}{2} \iint_{Q_\ell \times Q_\ell} \rho_\gamma(x) G_\ell(x - y) \rho_\gamma(y) dx dy \\ &\quad - \frac{\alpha}{2} \iint_{Q_\ell^* \times Q_\ell^*} d\xi d\xi' \iint_{Q_\ell \times Q_\ell} \text{Tr}_4 [\gamma_\xi(x, y) \gamma_{\xi'}(y, x)] W_\ell^\infty(\xi - \xi', x - y) dx dy. \end{aligned}$$

This functional is well-defined on Z (see Remark 4.9 below). In the above definition of the energy functional, the so-called fine structure constant α is a dimensionless positive constant (the physical value is approximately 1/137). Note that $D_\xi \gamma_\xi$ is not a trace-class operator, so $\text{Tr}_{L_\xi^2} [D_\xi \gamma_\xi]$ is not really a trace, it is just a notation for the rigorous mathematical object $\text{Tr}_{L_\xi^2} [|D_\xi|^{1/2} \gamma_\xi |D_\xi|^{1/2} \text{sign}(D_\xi)]$. We will make this abuse of notation throughout the paper.

The last term in (2.20) is called the “exchange term”. The potential W_ℓ^∞ that enters its definition is given by

$$W_\ell^\infty(\eta, x) = \sum_{k \in \mathbb{Z}^3} \frac{e^{i\ell k \cdot \eta}}{|x + \ell k|} = \frac{4\pi}{\ell^3} \sum_{k \in \mathbb{Z}^3} \frac{1}{|\frac{2\pi k}{\ell} - \eta|^2} e^{i(\frac{2\pi k}{\ell} - \eta) \cdot x} \quad (2.20)$$

(see [8] for a formal derivation of the exchange term from its analogue for molecules). It is Q_ℓ^* -periodic with respect to η and quasi-periodic with quasi-momentum η with respect to x . For every $\gamma \in Z$, we now define the mean-field periodic Dirac operator

$$D_\gamma = \int_{Q_\ell^*}^\oplus D_{\gamma, \xi} d\xi \quad \text{with} \quad D_{\gamma, \xi} := D_\xi - \alpha z G_\ell + \alpha V_{\gamma, \xi}$$

where

$$V_{\gamma,\xi} = \rho_\gamma * G_\ell - W_{\gamma,\xi}. \quad (2.21)$$

Here,

$$\rho_\gamma * G_\ell(x) = \int_{Q_\ell} G_\ell(y-x) \rho_\gamma(y) dy = \widetilde{\text{Tr}}_{L^2}[G_\ell(\cdot-x) \gamma] \quad (2.22)$$

and

$$W_{\gamma,\xi} \psi_\xi(x) = \int_{Q_\ell^*} d\xi' \int_{Q_\ell} W_\ell^\infty(\xi' - \xi, x-y) \gamma_{\xi'}(x,y) \psi_\xi(y) dy. \quad (2.23)$$

(In (2.22) we keep the notation $\cdot * \cdot$ for the convolution of periodic functions on Q_ℓ .)

Let us explain the relation between \mathcal{E}^{DF} and D_γ . The periodic DF energy may be rewritten

$$\mathcal{E}^{DF}(\gamma) = \widetilde{\text{Tr}}_{L^2}[(\mathcal{D}^0 - \alpha G) \gamma + \frac{\alpha}{2} V_\gamma \gamma].$$

It is smooth on Z , and its differential at $\gamma \in Z$ is the linear form

$$d\mathcal{E}^{DF}(\gamma) : Z \ni h \mapsto \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[D_{\gamma,\xi} h_\xi] d\xi = \widetilde{\text{Tr}}_{L^2}[D_\gamma h].$$

We introduce the following set of periodic density matrices :

$$\Gamma_q := \{\gamma \in \Gamma \mid \|\gamma\|_{\mathfrak{S}_{1,1}} = q\}$$

and

$$\Gamma_{\leq q} := \{\gamma \in \Gamma \mid \|\gamma\|_{\mathfrak{S}_{1,1}} \leq q\}.$$

Here q is a positive real number. The elements of Γ_q (resp. $\Gamma_{\leq q}$) are Dirac–Fock density matrices with particle number per unit cell equal to q (resp. at most q).

Our goal is to define the ground state despite the fact that the energy functional \mathcal{E}^{DF} is strongly indefinite on $\Gamma_{\leq q}$, due to the unboundedness of the Dirac operator D^0 .

2.3 Ground state energy and main result

We follow Dirac’s interpretation of the negative energy states of Dirac–Fock models: Such states are supposed to be occupied by virtual electrons that form the Dirac sea. Therefore, by the Pauli exclusion principle, the states of physical electrons are orthogonal to all the negative energy states. The ground-energy and state should thus be defined on the positive spectral subspaces of the corresponding Dirac–Fock operator. Let

$$P_\gamma^\pm := \int_{Q_\ell^*}^\oplus P_{\gamma,\xi}^\pm d\xi \quad \text{with} \quad P_{\gamma,\xi}^\pm := 1_{\mathbb{R}_\pm}(D_{\gamma,\xi}).$$

Note that by definition $P_{0,\xi}^\pm = 1_{\mathbb{R}_\pm}(D_\xi - \alpha z G_\ell)$. We define the set

$$\Gamma_q^+ := \{\gamma \in \Gamma_q \mid \gamma = P_\gamma^+ \gamma P_\gamma^+\} \quad (2.24)$$

and the ground state energy

$$I_q := \inf_{\gamma \in \Gamma_q^+} \mathcal{E}^{DF}(\gamma). \quad (2.25)$$

We need the following assumption.

Assumption 2.5. *Let $q^+ := \max\{q, 1\}$ and $\kappa := \alpha(C_{Gz} + C'_{EE} q^+)$. We also introduce the positive constants $e_0 := (1 - \kappa)^{-1} c^*([q])$ and $a := \frac{\alpha}{2} C_{EE} (1 - \kappa)^{-1/2} \lambda_0^{-1/2}$ (well-defined if $\kappa < 1$). Here we have used the standard notation $[q] := \min\{m \in \mathbb{N} \mid m \geq q\}$ and $c^*(\cdot)$ is given by formula (2.9).*

We demand that

1. $\kappa < 1 - \frac{\alpha}{2} C_{EE} q^+$;
2. $2a \sqrt{\max\{(1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} e_0 q; 1\}} q^+ < 1$.

The positive constants C_G , C_{EE} , C'_{EE} and λ_0 are defined respectively in Lemmas 4.1, 4.7 and 4.11 below.

Our main result is the following.

Theorem 2.6 (Existence of a ground state). *When α , q , z and ℓ satisfy Assumption 2.5, there exists $\gamma_* \in \Gamma_q^+$ such that*

$$\mathcal{E}^{DF}(\gamma_*) = I_q = \min_{\gamma \in \Gamma_q^+} \mathcal{E}^{DF}(\gamma). \quad (2.26)$$

Besides, γ_* solves the following nonlinear self-consistent equation

$$\gamma = 1_{[0, \nu)}(D_\gamma) + \delta \quad (2.27)$$

where $0 \leq \delta \leq 1_{\{\nu\}}(D_\gamma)$ and $\lambda_0 \leq \nu \leq e_0$, with e_0 being defined in Assumption 2.5, and $\lambda_0 \geq 1 - \kappa > 0$ in Lemma 4.11.

Remark 2.7 (Projectors). *According to [1, 19, 31] any ground state of the Hartree–Fock model (both for the molecules and crystals) is a projector. However we do not know whether the ground states of Dirac–Fock model are projectors in general.*

Remark 2.8. *In Solid State Physics, the length of the unit cell is about a few Ångströms. In our system of units, $\hbar = m = c = 1$, thus $\alpha \approx \frac{1}{137}$ and $\ell \approx 1000$. Under the condition $q = z$ for electrical neutrality, Assumption 2.5 is satisfied for $q \leq 17$. The proof is detailed in Appendix D. Our estimates are far from optimal: The ideas of this paper are expected to apply to higher values of q .*

3 Sketch of proof of Theorem 2.6

We are convinced that the constraint set Γ_q^+ is not convex, and we are not able to prove that it is closed for the weak-* topology of Z . This is the source of considerable difficulties. Mimicking [39], we shall use a retraction technique as for the Dirac–Fock model for atoms and molecules. This imposes to search the ground state in the set $\Gamma_{\leq q}^+$ defined by

$$\Gamma_{\leq q}^+ := \{\gamma \in \Gamma_{\leq q} \mid \gamma = P_\gamma^+ \gamma P_\gamma^+\}.$$

However, under the above constraint, the minimizers may not be situated in Γ_q^+ . To overcome this problem, we next subtract a penalization term $\epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma)$ from the DF energy functional, for some parameter $\epsilon_P > 0$ to be chosen later, and we first study a minimization problem for the penalized functional with relaxed constraint. We introduce the infimum

$$J_{\leq q} := \inf_{\gamma \in \Gamma_{\leq q}^+} \left[\mathcal{E}^{DF}(\gamma) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma) \right]. \quad (3.1)$$

If this infimum is attained at some $\gamma_* \in \Gamma_{\leq q}^+$, γ_* will be called a *minimizer for $J_{\leq q}$* . We are going to see that for a suitably chosen value of ϵ_P , $J_{\leq q}$ is attained and that every minimizer for $J_{\leq q}$ lies in Γ_q^+ , thus is a minimizer for I_q .

For the study of the penalized problem $J_{\leq q}$, we need an analogue of Assumption 2.5:

Assumption 3.1. *Let $q^+ = \max\{q; 1\}$, $\kappa := \alpha(C_G z + C'_{EE} q^+)$ and $a := \frac{\alpha}{2} C_{EE} (1 - \kappa)^{-1/2} \lambda_0^{-1/2}$ (well-defined if $\kappa < 1$). We assume that*

1. $\kappa < 1 - \frac{\alpha}{2} C_{EE} q^+$;
2. $2a \sqrt{\max\{(1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} \epsilon_P q; 1\} q^+} < 1$.

The relation between assumptions 2.5 and 3.1 is given by the following lemma:

Lemma 3.2 (Choice of ϵ_P). *Assume that Assumption 2.5 on q and z holds. Then, there is a constant $\epsilon_P > e_0$ such that Assumption 3.1 is satisfied.*

Proof. One just needs to take $\epsilon_P = e_0 + \varepsilon$ with ε positive and small enough. □

We now state an existence result:

Theorem 3.3 (Existence of a minimizer for the penalized problem). *We suppose that Assumption 3.1 on q, z, ϵ_P holds and recall the notation $e_0 := (1 - \kappa)^{-1} c^*([q])$. If $\epsilon_P > e_0$, then there exists $\gamma_* \in \Gamma_{\leq q}^+$ such that*

$$\mathcal{E}^{DF}(\gamma_*) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma_*) = J_{\leq q}. \quad (3.2)$$

Besides, $\widetilde{\text{Tr}}_{L^2}(\gamma_*) = \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}(\gamma_{*,\xi}) d\xi = q$ and γ_* solves the following nonlinear self-consistent equation

$$\gamma = 1_{[0,\nu)}(D_\gamma) + \delta \quad (3.3)$$

where $0 \leq \delta \leq 1_{\{\nu\}}(D_\gamma)$ and $\nu \in [\lambda_0, e_0]$ is the Lagrange multiplier due to the charge constraint $\text{Tr}_{L^2}(\gamma) \leq q$.

Theorem 2.6 is a direct consequence of Lemma 3.2 and Theorem 3.3. Indeed, if Assumption 2.5 on q, z holds, Lemma 3.2 guarantees the existence of ϵ_P such that the assumptions of Theorem 3.3 are satisfied. Then this theorem provides a minimizer γ_* for $J_{\leq q}$ which lies in Γ_q^+ , hence $J_{\leq q} + \epsilon_P q = \mathcal{E}^{DF}(\gamma_*) \geq I_q$. On the other hand, for each $\gamma \in \Gamma_q^+$ one has the inequality $\mathcal{E}^{DF}(\gamma) \geq J_{\leq q} + \epsilon_P q$, so $I_q \geq J_{\leq q} + \epsilon_P q$. As a consequence, we get $\mathcal{E}^{DF}(\gamma_*) = J_{\leq q} + \epsilon_P q = I_q$ which is the same as (2.26). Moreover γ_* satisfies (3.3) which is the same as (2.27).

Therefore, in the sequel of this paper we focus on the proof of Theorem 3.3. Before going further, we explain the difficulties we face and the strategy we adopt to solve them, by comparing with the Hartree–Fock case [8]. The method used in [8] is based on some properties of the Schrödinger operator $-\Delta$:

1. This operator is non-negative. Hence the Hartree–Fock model for crystals is well-defined and the kinetic energy is weakly lower semi-continuous w.r.t. the density matrix ;
2. The exchange potential W_ℓ^∞ is rather easily controlled by the Schrödinger operator $-\Delta$.

In [8], these properties allow to deduce bounds on the minimizing sequence of density matrices w.r.t. the ξ, x and y variables, and to pass to the limit in the different terms of the energy functional, in particular in the exchange term which is the most intricate one. In the proof, the strong convergence of the density matrix kernels $\gamma_n(x, y) = \int_{Q_\ell^*} \gamma_{n,\xi}(x, y) d\xi$ plays an important role. In addition, the charge constraint in the periodic Hartree–Fock model is linear with respect to the density, and there is no possible loss of charge in passing to the limit.

In the Dirac–Fock model for crystals, two additional difficulties occur. First of all, the Dirac operator does not control the potential energy terms, which are of the same order. Secondly, the convergence of the nonlinear constraint $\int_{Q_\ell^*} P_{\gamma,\xi}^+ \gamma_\xi d\xi = \int_{Q_\ell^*} \gamma_\xi d\xi$ requires stronger compactness properties of the sequence of density matrices with respect to the ξ variable (of course, this second difficulty does not exist for the Dirac–Fock model of atoms and molecules, since in that case the ξ variable is absent). Therefore the proof of existence of minimizers in the periodic Hartree–Fock setting cannot be applied *mutatis mutandis*. The functional space Z is natural to give a sense to the energy functional and to the constraints, but the weak convergence of minimizing sequences in Z is not sufficient to deal with the exchange term and the non-linear constraints. The whole paper (except Section 5 about the retraction) is devoted to solving the difficulties arising from the ξ variable.

Strategy for the proof of Theorem 3.3. Our strategy rather relies on the spectral analysis of the periodic Dirac–Fock operator, which is new, to our knowledge, for the proof of existence of minimizers in the periodic case. Thanks to this spectral analysis, in Lemma 4.15 together with Lemma 5.1 (see also Remark 4.16), we can prove that every minimizer for $J_{\leq q}$ actually lies in $\mathfrak{S}_{1,\infty}$, and is situated in $\overline{\mathfrak{B}}$, where

$$\overline{\mathfrak{B}} := \{ \gamma \in Z \mid \|\gamma\|_{\mathfrak{S}_{1,\infty}} \leq j_1 \} \quad (3.4)$$

and j_1 is an integer defined in Subsection 4.2.

The key point in the proof of the existence of minimizers for $J_{\leq q}$ is that for any minimizing sequence (γ_n) for $J_{\leq q}$, we are able to construct another minimizing sequence $(\tilde{\gamma}_n)$ satisfying the same regularity estimate as the minimizers, that is, $\tilde{\gamma}_n \in \overline{\mathfrak{B}}$. This is the content of Lemma 6.2. This estimate helps considerably to solve the problem of passing to the limit in the constraint $\int_{Q_\ell^*} P_{\gamma,\xi}^+ \gamma_\xi P_{\gamma,\xi}^+ d\xi = \int_{Q_\ell^*} \gamma_\xi d\xi$.

Organization of the paper. The next sections are devoted to the proof of Theorem 3.3. Our paper is organized as follows.

In Section 4, we collect some fundamental estimates on the potentials G_ℓ and W_ℓ^∞ , that ensure in particular that the DF periodic energy functional is well-defined and smooth on Z . In Subsection 4.2, we study the spectral properties of the Dirac–Fock operators $D_{\gamma,\xi}$ for every $\xi \in Q_\ell^*$. Relying on them, we study in Subsection 4.3 the properties of minimizers for a linearized Dirac–Fock energy. Finally, we collect the first estimates on minimizing sequences for $J_{\leq q}$.

In Section 5, we study the Euler-Lagrange equation associated to $J_{\leq q}$. We conclude that each minimizer for $J_{\leq q}$ is in Γ_q^+ and solves a self-consistent equation (it is a ground state of its own mean-field Hamiltonian) and that minimizing sequences are approximate solutions. In Hartree–Fock type models for molecules [33] or crystals [19], it is a standard fact that the approximate minimizers are also approximate ground states of their mean-field Hamiltonian, and the proof relies on the convexity of the constraint set. However, in the Dirac–Fock model (both for molecules and crystals), the constraint set $\Gamma_{\leq q}^+$ does not seem to be convex. By using a retraction technique, a similar difficulty was recently overcome by one of us for the Dirac–Fock model of molecules [39]. Adapting the method of [39], we define a set \mathcal{V} which is relatively open in $\Gamma_{\leq q}$ for the norm of Z , and we build a regular map $\theta : \overline{\mathcal{V}} \rightarrow \overline{\mathcal{V}} \cap \Gamma_{\leq q}^+$ such that $\theta(\gamma) = \gamma$, $\forall \gamma \in \overline{\mathcal{V}} \cap \Gamma_{\leq q}^+$. Here, $\overline{\mathcal{V}}$ is the closure of \mathcal{V} in Z . Under our assumptions, there exist minimizing sequences for $J_{\leq q}$ lying in $\mathcal{V} \cap \Gamma_{\leq q}^+$, hence the equality

$$J_{\leq q} = \inf_{\gamma \in \mathcal{V}} \left\{ \mathcal{E}^{DF}(\theta(\gamma)) - \epsilon_P \widetilde{\text{Tr}}_{L^2}[\theta(\gamma)] \right\}$$

which allows us to prove that the terms of minimizing sequences are approximate ground states of their mean-field Hamiltonian.

Then, in Section 6, we build modified minimizing sequences lying in $\overline{\mathbf{B}}$. Finally, we prove the convergence of such sequences to a minimizer for $J_{\leq q}$, and this ends the proof of Theorem 3.3.

Assumption 2.5 involves optimal constants in Hardy-type inequalities introduced in Subsection 4.1. Therefore, in Appendix A–C, we prove Lemma 4.1, Lemma 4.5 and Lemma 4.7 respectively. Finally, in Appendix D, we calculate the maximum number of electrons per cell allowed by the model, relying on approximate values of the constants obtained in Appendices A–C.

4 Fundamental estimates

In this section, we give Hardy-type inequalities for the periodic Coulomb potential and provide estimates on the interaction potential between electrons in crystals. Then we study the spectrum of the periodic self-consistent Dirac–Fock operators. Finally, using this spectral analysis, we derive properties of the minimizers for a linearized problem, and *a priori* bounds on minimizing sequences for $J_{\leq q}$.

4.1 Hardy-type estimates on the periodic Coulomb potential

First of all, and this is a major difference with the usual Coulomb potential $\frac{1}{|x|}$ in \mathbb{R}^3 , the periodic Coulomb potential G_ℓ may not be positive, since it is defined up to constant, but it is bounded from below (see Lemma A.1 in Appendix A). Nevertheless, the operator of convolution with G_ℓ is positive on $L^2(Q_\ell)$ in virtue of (2.3). Moreover, we have the following Hardy-type estimates concerning the operator of multiplication by the periodic potential G_ℓ .

Lemma 4.1 (Hardy-type inequalities for the periodic Coulomb potential). *There exist positive constants $C_H = C_H(\ell) > 0$ that only depends on ℓ and such that*

$$G_\ell \leq |G_\ell| \leq C_H |D^0| \tag{4.1}$$

in the sense of operators on $L^2(Q_\ell^) \otimes L^2(Q_\ell; \mathbb{C}^4)$.*

Moreover, there exists a positive constant $C_G = C_G(\ell)$ with $C_G \geq C_H$ that only depends on ℓ and such that

$$\|G_\ell |D^0|^{-1}\|_Y = C_G. \tag{4.2}$$

Remark 4.2. In (4.1), the inequality $A \leq B$ is equivalent to : For almost every $\xi \in Q_\ell^*$, $A_\xi \leq B_\xi$ in the sense of operators on L_ξ^2 .

Remark 4.3. The constant $C_G(\ell)$ is estimated in (A.4) in Appendix A below. While it is far from optimal when ℓ is small, it converges to 2 when ℓ goes to infinity; that is, to the value of the optimal constant for the Coulomb potential on the whole space. By interpolation,

$$C_H \leq C_G. \quad (4.3)$$

Therefore, (4.1) holds with C_H being replaced by C_G . However, C_H is expected to converge to $\pi/2$ as ℓ goes to infinity; that is, to the best constant in the Kato–Herbst inequality on the whole space [26, 29].

A by-product of Lemma 4.1 is the following.

Corollary 4.4 (Estimates on the direct term). *For any $\gamma \in X$, we have*

$$\|\rho_\gamma * G_\ell\|_Y \leq C_H \|\gamma\|_X \quad (4.4)$$

and

$$\|(\rho_\gamma * G_\ell) |D^0|^{-1}\|_Y \leq C_G \|\gamma\|_{\mathfrak{S}_{1,1}}. \quad (4.5)$$

Proof. For every $x \in \mathbb{R}^3$ and $\gamma \in X$, we have

$$\begin{aligned} |\rho_\gamma * G_\ell(x)| &= \left| \widetilde{\text{Tr}}_{L^2} [G_\ell(x - \cdot) \gamma] \right| \\ &= \left| \widetilde{\text{Tr}}_{L^2} [|D^0|^{-1/2} G_\ell(x - \cdot) |D^0|^{-1/2} |D^0|^{1/2} \gamma |D^0|^{1/2}] \right| \\ &\leq \| |D^0|^{-1/2} G_\ell(x - \cdot) |D^0|^{-1/2} \|_Y \| |D^0|^{1/2} \gamma |D^0|^{1/2} \|_{\mathfrak{S}_{1,1}} \leq C_H \|\gamma\|_X. \end{aligned}$$

Indeed, the bound (4.1) in Lemma 4.1 yields

$$\left\| |G_\ell(\cdot - x)|^{1/2} |D^0|^{-1/2} \right\|_Y \leq (C_H)^{1/2}$$

uniformly in x .

We now turn to the proof of (4.5). For every $\xi \in Q_\ell^*$ and φ_ξ in L_ξ^2 , we have

$$\begin{aligned} \|(\rho_\gamma * G_\ell) |D_\xi|^{-1} \varphi_\xi\|_{L_\xi^2} &\leq \int_{Q_\ell} |\rho_\gamma(x)| \| |G_\ell(\cdot - x)| |D_\xi|^{-1} \varphi_\xi \|_{L_\xi^2} dx \\ &\leq \sup_{x \in \mathbb{R}^3} \| |G_\ell(\cdot - x)| |D_\xi|^{-1} \varphi_\xi \|_{L_\xi^2} \int_{Q_\ell} |\rho_\gamma(x)| dx \leq C_G \|\gamma\|_{\mathfrak{S}_{1,1}} \|\varphi_\xi\|_{L_\xi^2}. \end{aligned} \quad (4.6)$$

In (4.6), we have used the bound (4.2) of Lemma 4.1 and the obvious fact that it remains true for $G_\ell(\cdot - x)$ for any $x \in \mathbb{R}^3$. □

Now, we consider the operator $W_\gamma := \int_{Q_\ell^*}^\oplus W_{\gamma,\xi} d\xi$ which enters the definition of the exchange term. The operators $W_{\gamma,\xi}$ have been defined in Formula (2.23), which involves the integral kernel W_ℓ^∞ given in (2.20). We can separate the singularities of W_ℓ^∞ with respect to $\eta \in 2Q_\ell^*$ and $x \in 2Q_\ell$ as follows

$$W_\ell^\infty(\eta, x) = W_{\geq m, \ell}^\infty(\eta, x) + W_{< m, \ell}^\infty(\eta, x), \quad \forall m \in \mathbb{N}, m \geq 2, \quad (4.7)$$

with

$$W_{\geq m, \ell}^\infty(\eta, x) = \frac{4\pi}{\ell^3} \sum_{\substack{|k|_\infty \geq m \\ k \in \mathbb{Z}^3}} \frac{1}{\left| \frac{2\pi k}{\ell} - \eta \right|^2} e^{i\left(\frac{2\pi k}{\ell} - \eta\right) \cdot x}$$

and

$$W_{< m, \ell}^\infty(\eta, x) = \frac{4\pi}{\ell^3} \sum_{\substack{|k|_\infty < m \\ k \in \mathbb{Z}^3}} \frac{1}{\left| \frac{2\pi k}{\ell} - \eta \right|^2} e^{i\left(\frac{2\pi k}{\ell} - \eta\right) \cdot x}$$

where $|k|_\infty := \max\{|k_1|; |k_2|; |k_3|\}$. It is easy to see that the singularity of $W_{< m, \ell}^\infty$ behaves like $\frac{1}{|\eta|^2}$. We will show in Appendix B that the singularity of $W_{\geq m, \ell}^\infty$ behaves like $\frac{1}{|x|}$ or equivalently $G_\ell(x)$, and we will obtain the following estimates on the operator W_γ .

Lemma 4.5 (Estimates on W_γ). *If $\gamma \in Z$, then $W_\gamma \in Y$ and there exist positive constants $C_W = C_W(\ell)$, $C'_W = C'_W(\ell)$, $C''_W = C''_W(\ell)$ that only depend on ℓ , such that*

$$\|W_\gamma\|_Y \leq C_W \|\gamma\|_Z \quad \text{if } \gamma \in Z, \quad (4.8)$$

$$\|W_\gamma\|_Y \leq C''_W (\|\gamma\|_X + \|\gamma\|_{\mathfrak{S}_{1,\infty}}^{3/4} \|\gamma\|_{\mathfrak{S}_{1,1}}^{1/4}) \quad \text{if } \gamma \in X \cap \mathfrak{S}_{1,\infty}, \quad (4.9)$$

$$\|W_\gamma |D^0|^{-1}\|_Y \leq C'_W \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \quad \text{if } \gamma \in \mathfrak{S}_{1,1} \cap Y. \quad (4.10)$$

Remark 4.6. *The constants C_W , C'_W and C''_W are estimated in (B.24) in Appendix B.*

Gathering together Lemmas 4.1, 4.5 and Corollary 4.4, we can get some rough estimates on the self-consistent potential $V_{\gamma,\xi}$ defined in (2.21). In Appendix C we obtain much better estimates by a careful study of the structure of the operator $V_\gamma = \int_{Q_\ell^*}^\oplus V_{\gamma,\xi} d\xi$:

Lemma 4.7 (Estimates on V_γ). *There exist positive constants $C_{EE} = C_{EE}(\ell) > 0$ and $C'_{EE} = C'_{EE}(\ell) > 0$ that only depend on ℓ and such that, for every $\gamma \in Z$,*

$$\|V_\gamma\|_Y \leq C_{EE} \|\gamma\|_Z \quad (4.11)$$

and

$$\|V_\gamma |D^0|^{-1}\|_Y \leq C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}. \quad (4.12)$$

For every $\xi \in Q_\ell^*$ and any $\psi_\xi \in H_\xi^{1/2}$,

$$\left| (\psi_\xi, V_{\gamma,\xi} \psi_\xi)_{L_\xi^2} \right| \leq C_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \|\psi_\xi\|_{H_\xi^{1/2}}^2. \quad (4.13)$$

Furthermore, if $\gamma \geq 0$, for any $\psi \in L_\xi^2$,

$$-C''_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \|\psi_\xi\|_{L_\xi^2}^2 \leq (\psi_\xi, V_{\gamma,\xi} \psi_\xi)_{L_\xi^2}. \quad (4.14)$$

Remark 4.8. *The constants C_{EE} , C'_{EE} and C''_{EE} are estimated in (C.7), (C.5) and (C.8) of Appendix C respectively.*

Remark 4.9. *Using Lemmas 4.1 and 4.7, it is easily checked that $Z \ni \gamma \mapsto \mathcal{E}(\gamma)$ is well-defined.*

4.2 Spectral properties of the mean-field Dirac–Fock operator

Recall that $\kappa := \alpha (C_G z + C'_{EE} q^+)$. We start with the following.

Lemma 4.10. *Let $\gamma \in Z$. We assume that $C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} < 1/\alpha$, then $D_{\gamma,\xi}$ is a self-adjoint operator on L_ξ^2 with domain H_ξ^1 and form-domain $H_\xi^{1/2}$. In addition, the following holds*

$$\left\| |D_\gamma|^{1/2} |D^0|^{-1/2} \right\|_Y \leq (1 + \alpha (C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}))^{1/2} \quad (4.15)$$

and

$$\left\| |D^0|^{1/2} |D_\gamma|^{-1/2} \right\|_Y \leq (1 - \alpha (C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}))^{-1/2}. \quad (4.16)$$

In particular, if $\gamma \in \Gamma_{\leq q}$ and $\kappa < 1$, we have

$$(1 - \kappa) |D^0| \leq |D_\gamma| \leq (1 + \kappa) |D^0|. \quad (4.17)$$

Proof. Recall $q^+ = \max\{1; q\}$. By Lemmas 4.1 and 4.7, we obtain

$$\|(-\alpha z G_\ell + \alpha V_\gamma) |D^0|^{-1}\|_Y \leq \alpha (C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}). \quad (4.18)$$

In particular, D_γ is self-adjoint on $\int_{Q_\ell^*}^\oplus H_\xi^1 d\xi$ by the Rellich–Kato theorem if $C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} < 1/\alpha$ (see [38, Theorem XIII-85]). Let now $\xi \in Q_\ell^*$ and $u_\xi \in H_\xi^1(Q_\ell)$. We have

$$\|D_{\gamma,\xi} u_\xi\|_{L_\xi^2} \leq (1 + \alpha C_G z + \alpha C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}) \|D_\xi u_\xi\|_{L_\xi^2}, \quad (4.19)$$

which implies (4.15). On the other hand,

$$\begin{aligned} \|D_\xi u_\xi\|_{L_\xi^2} &\leq \|(D_{\gamma,\xi} - D_\xi) u_\xi\|_{L_\xi^2} + \|D_{\gamma,\xi} u_\xi\|_{L_\xi^2} \\ &\leq \alpha (C_G z + C'_{EE} q^+) \|D_\xi u_\xi\|_{L_\xi^2} + \|D_{\gamma,\xi} u_\xi\|_{L_\xi^2}. \end{aligned}$$

Hence,

$$\|D_\xi u_\xi\|_{L_\xi^2} \leq (1 - \alpha(C_G z + C'_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y}))^{-1} \|D_{\gamma,\xi} u_\xi\|_{L_\xi^2} \quad (4.20)$$

which implies (4.16). Since $\gamma \in \Gamma_{\leq q}$, $\|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \leq q^+$. Thus (4.19) and (4.20) together give (4.17). This concludes the proof. \square

As a consequence of (4.20), we deduce that the spectrum of D_γ (and of any $D_{\gamma,\xi}$) is included in $\mathbb{R} \setminus [-1 + \kappa; 1 - \kappa]$. In order to allow for as many electrons as possible per cell, we need a more accurate estimate on the bottom of $\sigma(|D_\gamma|)$.

Lemma 4.11 (Further properties of the bottom of the spectrum of $|D_\gamma|$). *Let $\gamma \in \Gamma_{\leq q}$. Then*

$$\inf \sigma(|D_\gamma|) \geq \lambda_0 \geq 1 - \kappa,$$

with $\lambda_0 := 1 - \alpha \max\{C_H z + C''_{EE} q^+, \frac{C_0}{\ell} z + C_{EE} q^+\}$, the constant C_0 being defined in Lemma A.1 in Appendix A below.

Proof. Let $\psi_\xi^+ = \Lambda_\xi^+ \psi_\xi$ and $\psi_\xi^- = \Lambda_\xi^- \psi_\xi$. Notice that $D_{\gamma,\xi} = D_\xi - \alpha z G_\ell + \alpha V_{\gamma,\xi}$ and $V_{\gamma,\xi}$ satisfies (4.13) and (4.14). Now, combining with (A.1) in Appendix A, we have

$$\left(\psi_\xi^+, D_{\gamma,\xi} \psi_\xi^+ \right)_{H_\xi^{1/2} \times H_\xi^{-1/2}} \geq (1 - \alpha(C_H z + C''_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y})) \|\psi_\xi^+\|_{H_\xi^{1/2}}^2$$

and

$$- \left(\psi_\xi^-, D_{\gamma,\xi} \psi_\xi^- \right)_{H_\xi^{1/2} \times H_\xi^{-1/2}} \geq \left(1 - \alpha \left(\frac{C_0}{\ell} z + C_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \right) \right) \|\psi_\xi^-\|_{H_\xi^{1/2}}^2.$$

We get

$$\begin{aligned} \|\psi_\xi\|_{H_\xi^{1/2}} \|D_{\gamma,\xi} \psi\|_{H_\xi^{-1/2}} &\geq \Re \left(\psi_\xi^+ - \psi_\xi^-, D_{\gamma,\xi} \psi_\xi \right)_{H_\xi^{1/2} \times H_\xi^{-1/2}} \\ &= \left(\psi_\xi^+, D_{\gamma,\xi} \psi_\xi^+ \right)_{H_\xi^{1/2} \times H_\xi^{-1/2}} - \left(\psi_\xi^-, D_{\gamma,\xi} \psi_\xi^- \right)_{H_\xi^{1/2} \times H_\xi^{-1/2}} \geq \lambda_0 \|\psi_\xi\|_{H_\xi^{1/2}}^2. \end{aligned}$$

\square

Further spectral properties of the self-consistent operator D_γ are collected in the following.

Lemma 4.12 (Properties of positive eigenvalues of $D_{\gamma,\xi}$). *Assume that $\kappa < 1$ and let $\gamma \in \Gamma_{\leq q}$. We denote by $\lambda_j(\xi)$, for $j \geq 1$, the j -th positive eigenvalue (counted with multiplicity) of the mean-field operator $D_{\gamma,\xi}$. Then $\lambda_j(\xi)$ is situated in the interval $[c_*(j)(1 - \kappa), c^*(j)(1 - \kappa)^{-1}]$ where $c^*(j)$ and $c_*(j)$ are the constants of Formula (2.9).*

In addition, every eigenfunction $u_{j,\xi}$ associated to $\lambda_j(\xi)$ lies in H_ξ^1 and satisfies

$$\|u_{j,\xi}\|_{H_\xi^1} \leq c^*(j) (1 - \kappa)^{-2} \|u_{j,\xi}\|_{L_\xi^2}. \quad (4.21)$$

Proof. We rely on a variational characterization of eigenvalues in spectral gaps (see [14] and references therein). Let

$$\Lambda_\xi^+ := 1_{\mathbb{R}^+}(D_\xi) = \frac{1}{2} + \frac{D_\xi}{2|D_\xi|}$$

and

$$\Lambda_\xi^- := 1_{\mathbb{R}^-}(D_\xi) = \frac{1}{2} - \frac{D_\xi}{2|D_\xi|}.$$

One has $\Lambda_\xi^\pm H_\xi^1 \subset H_\xi^1$, that is, the domain H_ξ^1 of the self-adjoint operator $D_{\gamma,\xi}$ satisfies Condition (H1) of [14]. To each integer $j \geq 0$ we associate the min-max level

$$\hat{\lambda}_j(\xi) := \inf_{\substack{V \text{ subspace of } \Lambda_\xi^+ H_\xi^1 \\ \dim V = j}} \sup_{u_\xi \in (V \oplus \Lambda_\xi^- H_\xi^1) \setminus \{0\}} \frac{(D_{\gamma,\xi} u_\xi, u_\xi)}{\|u_\xi\|_{L_\xi^2}^2}. \quad (4.22)$$

Let $u_\xi \in (V \oplus \Lambda_\xi^- H_\xi^1) \setminus \{0\}$. We write $u_\xi = u_\xi^+ + u_\xi^-$ with

$$u_\xi^+ = \Lambda_\xi^+ u_\xi \in V, \quad u_\xi^- = \Lambda_\xi^- u_\xi \in \Lambda_\xi^- H_\xi^1.$$

By definition of Λ_ξ^\pm ,

$$(D_\xi u_\xi^+, u_\xi^+) = (|D_\xi| u_\xi^+, u_\xi^+), \quad (D_\xi u_\xi^-, u_\xi^-) = -(|D_\xi| u_\xi^-, u_\xi^-) \text{ and } (D_\xi u_\xi^+, u_\xi^-) = 0.$$

Therefore,

$$\begin{aligned} (D_{\gamma,\xi} u_\xi, u_\xi) &= (D_\xi u_\xi, u_\xi) + ((D_{\gamma,\xi} - D_\xi) u_\xi, u_\xi) \\ &= (|D_\xi| u_\xi^+, u_\xi^+) - (|D_\xi| u_\xi^-, u_\xi^-) + ((D_{\gamma,\xi} - D_\xi) u_\xi^+, u_\xi^+) + ((D_{\gamma,\xi} - D_\xi) u_\xi^-, u_\xi^-) \\ &\quad + 2\Re((D_{\gamma,\xi} - D_\xi) u_\xi^+, u_\xi^-). \end{aligned} \quad (4.23)$$

We observe that for $j \geq 1$,

$$\hat{\lambda}_j(\xi) \geq \inf_{\substack{V \text{ subspace of } \Lambda_\xi^+ H_\xi^1 \\ \dim V = j}} \sup_{u_\xi^+ \in V \setminus \{0\}} \frac{(D_{\gamma,\xi} u_\xi^+, u_\xi^+)}{\|u_\xi^+\|_{L_\xi^2}^2}.$$

By (4.23), (4.2) in Lemma 4.1 and (4.12) in Lemma 4.7, for any $u_\xi^+ \in \Lambda_\xi^+ H_\xi^1$,

$$(D_{\gamma,\xi} u_\xi^+, u_\xi^+) = (|D_\xi| u_\xi^+, u_\xi^+) + ((-\alpha z G_\ell + \alpha V_{\gamma,\xi}) u_\xi^+, u_\xi^+) \geq (1 - \kappa) (|D_\xi| u_\xi^+, u_\xi^+).$$

Thus,

$$\begin{aligned} \hat{\lambda}_j(\xi) &\geq (1 - \kappa) \inf_{\substack{V \text{ subspace of } \Lambda_\xi^+ H_\xi^1 \\ \dim V = j}} \sup_{u_\xi^+ \in V \setminus \{0\}} \frac{(|D_\xi| u_\xi^+, u_\xi^+)}{\|u_\xi^+\|_{L_\xi^2}^2} \\ &= (1 - \kappa) d_j^+(\xi) \geq (1 - \kappa) c_*(j). \end{aligned}$$

In particular, $\hat{\lambda}_1(\xi) \geq (1 - \kappa) c_*(1) > 0$. On the other hand, $(D_{\gamma,\xi} u_\xi^-, u_\xi^-) \leq -(1 - \kappa) (|D_\xi| u_\xi^-, u_\xi^-) \leq 0$ for every $u_\xi^- \in \Lambda_\xi^- H_\xi^1$, whenever $\kappa < 1$. This implies that $\hat{\lambda}_0(\xi) \leq 0$, so Conditions (H2) and (H3) of [14] are satisfied, and we can conclude that

$$\lambda_j(\xi) = \hat{\lambda}_j(\xi), \quad \forall j \geq 1, \quad (4.24)$$

hence the lower bound $\lambda_j(\xi) \geq (1 - \kappa) c_*(j)$.

For the upper bound, we proceed as follows. Equations (4.18) and (4.23) together yield

$$\begin{aligned} (D_{\gamma,\xi} u_\xi, u_\xi) &= (|D_\xi| u_\xi^+, u_\xi^+) + ((-\alpha z G_\ell + \alpha V_{\gamma,\xi}) u_\xi^+, u_\xi^+) + 2\Re((-\alpha z G_\ell + \alpha V_{\gamma,\xi}) u_\xi^+, u_\xi^-) \\ &\quad + ((D_{\gamma,\xi} - D_\xi) u_\xi^-, u_\xi^-) - (|D_\xi| u_\xi^-, u_\xi^-) \\ &\leq (1 + \kappa) (|D_\xi| u_\xi^+, u_\xi^+) - (1 - \kappa) (|D_\xi| u_\xi^-, u_\xi^-) + 2\kappa \| |D_\xi|^{1/2} u_\xi^+ \|_{L_\xi^2} \| |D_\xi|^{1/2} u_\xi^- \|_{L_\xi^2} \\ &= (1 + \kappa) \| |D_\xi|^{1/2} u_\xi^+ \|_{L_\xi^2}^2 + 2\kappa \| |D_\xi|^{1/2} u_\xi^+ \|_{L_\xi^2} \| |D_\xi|^{1/2} u_\xi^- \|_{L_\xi^2} - (1 - \kappa) \| |D_\xi|^{1/2} u_\xi^- \|_{L_\xi^2}^2 \\ &\leq (1 - \kappa)^{-1} \| |D_\xi|^{1/2} u_\xi^+ \|_{L_\xi^2}^2 \end{aligned}$$

by Young's inequality. Moreover, $\|u_\xi^+\|_{L_\xi^2} \leq \|u_\xi\|_{L_\xi^2}$, so, recalling (4.22) and (4.24) we see that

$$\begin{aligned} \lambda_j(\xi) &\leq (1 - \kappa)^{-1} \inf_{\substack{V \text{ subspace of } \Lambda_\xi^+ H_\xi^1 \\ \dim V = j}} \sup_{u_\xi^+ \in V \setminus \{0\}} \frac{(|D_\xi| u_\xi^+, u_\xi^+)}{\|u_\xi^+\|_{L_\xi^2}^2} \\ &= (1 - \kappa)^{-1} d_j^+(\xi) \leq (1 - \kappa)^{-1} c^*(j). \end{aligned}$$

Finally, using (4.20) in Lemma 4.10, we obtain

$$\lambda_j(\xi) \|u_{j,\xi}\|_{L_\xi^2} = \|D_{\gamma,\xi} u_{j,\xi}\|_{L_\xi^2} \geq (1 - \kappa) \|D_\xi u_{j,\xi}\|_{L_\xi^2} = (1 - \kappa) \|u_{j,\xi}\|_{H_\xi^2},$$

hence (4.21). The lemma is thus proved. \square

Recall that $\overline{\mathbf{B}} := \{\gamma \in Z \mid \|\gamma\|_{\mathfrak{S}_{1,\infty}} \leq j_1\}$ where j_1 is an integer that has not been defined yet. In the rest of this paper, assuming that $\kappa < 1$ and recalling our notation $e_0 := (1 - \kappa)^{-1} c^*([q])$, we take

$$j_1 := \min\{j \geq 0 \mid (1 - \kappa)c_*(j + 1) > e_0\}. \quad (4.25)$$

This integer is well-defined, since $\lim_{j \rightarrow \infty} c_*(j) = +\infty$. We also introduce the energy level

$$e_1 := (1 - \kappa)c_*(j_1 + 1). \quad (4.26)$$

By construction, one has $0 < e_0 < e_1$ and $j_1 \geq [q]$. Moreover, Lemma 4.12 has an immediate consequence, which will be very useful in the sequel.

Corollary 4.13. *Assuming that $\kappa < 1$, with the above notation, for every γ in $\Gamma_{\leq q}$:*

- The projector $1_{[0,e_0]}(D_{\gamma,\xi})$ has rank at least $[q]$ for a.e. $\xi \in Q_\ell^*$.
- The projector $1_{[0,e_1]}(D_{\gamma,\xi})$ has rank at most j_1 for a.e. $\xi \in Q_\ell^*$.
- If $\gamma' \in Z$ and $0 \leq \gamma' \leq 1_{[0,e_1]}(D_\gamma)$, then $\gamma' \in \overline{\mathbf{B}}$.

We end this subsection with the following proposition.

Proposition 4.14. *Assume that $\kappa < 1$. Let $\gamma, \gamma' \in \Gamma_{\leq q}$ such that*

$$0 \leq \gamma' \leq 1_{[0,\nu]}(D_\gamma)$$

with $\nu > 0$. Then,

$$\|\gamma'\|_Z \leq \max\{(1 - \kappa)^{-1} q \nu; 1\}.$$

Proof. By Lemma 4.10, we have

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[D_{\gamma,\xi} \gamma'_\xi] d\xi = \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[|D_{\gamma,\xi}| \gamma'_\xi] d\xi \geq (1 - \kappa) \|\gamma'\|_X.$$

Since $\gamma' \in \Gamma_{\leq q}$, we obtain

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[D_{\gamma,\xi} \gamma'_\xi] d\xi \leq \nu \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[\gamma'_\xi] d\xi \leq q \nu.$$

Then $\|\gamma'\|_X \leq (1 - \kappa)^{-1} q \nu$. We deduce the desired bound since $\|\gamma'\|_Y \leq 1$. \square

4.3 Properties of the minimizers for a linearized problem

The following lemma will be used in the next sections.

Lemma 4.15. *Let $g \in \Gamma_{\leq q}$ be given, and assume $\kappa < 1$. Then for each $\epsilon_P > 0$, the minimization problem*

$$\inf_{\substack{\gamma \in \Gamma_{\leq q} \\ \gamma = P_g^+ \gamma P_g^+}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \epsilon_P)\gamma_\xi] d\xi$$

admits a minimizer. Every minimizer \hat{g} is of the form $\hat{g} = \int_{Q_\ell^}^\oplus 1_{[0,\nu]}(D_{g,\xi}) d\xi + \hat{\delta}$, with $0 \leq \hat{\delta} \leq \int_{Q_\ell^*}^\oplus 1_{\{\nu\}}(D_{g,\xi}) d\xi$ for some $\nu \in (0, \min(\epsilon_P, e_0)]$, and one has $\hat{g} \in \overline{\mathbf{B}}$.*

If $\epsilon_P > e_0$, the set of minimizers is independent of ϵ_P , and every minimizer satisfies $\widetilde{\text{Tr}}_{L^2}(\hat{g}) = q$ and $\nu \geq \lambda_0$.

Proof. The proof is inspired of [5]. For any $\xi \in Q_\ell^*$ we can choose an orthonormal eigenbasis $\{\psi_j(\xi, \cdot)\}_{j \geq 1}$ of $D_{g,\xi} P_{g,\xi}^+$, such that

$$D_{g,\xi} P_{g,\xi}^+ = \sum_{j \geq 1} \lambda_j(\xi) |\psi_j(\xi)\rangle \langle \psi_j(\xi)|,$$

with $\lambda_j(\xi) \geq 0$. According to Lemma 4.12, for every $\xi \in Q_\ell^*$ and for every $j \geq 1$

$$(1 - \kappa) c_*(j) \leq \lambda_j(\xi) \leq (1 - \kappa)^{-1} c^*(j).$$

Let us introduce as in [5, 19] the non-decreasing function

$$C : \mathbb{R}_+ \ni s \mapsto \frac{\ell^3}{(2\pi)^3} \sum_{j \geq 1} |\{\xi \in Q_\ell^* \mid 0 \leq \lambda_j(\xi) \leq s\}|.$$

By Lemma 4.12, for $0 \leq s < (1 - \kappa)c_*([q])$ one has $C(t) \leq [q] - 1 < q$. On the other hand, $C(e_0) \geq [q] \geq q$. Thus, there exists ν_1 with

$$1 - \kappa \leq (1 - \kappa)c_*([q]) \leq \nu_1 \leq e_0 \tag{4.27}$$

such that

$$\lim_{s \rightarrow \nu_1^-} C(s) \leq q \leq \lim_{s \rightarrow \nu_1^+} C(s). \tag{4.28}$$

Equivalently,

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[1_{[0,\nu_1]}(D_{g,\xi})] d\xi \leq q$$

and

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[1_{[0,\nu_1]}(D_{g,\xi})] d\xi \geq q.$$

Therefore, there exists $0 \leq \tilde{\delta} \leq \int_{Q_\ell^*}^\oplus 1_{\{\nu_1\}}(D_{g,\xi}) d\xi$, such that the density matrix

$$\tilde{g} := \int_{Q_\ell^*}^\oplus 1_{[0,\nu_1]}(D_{g,\xi}) d\xi + \tilde{\delta}$$

satisfies

$$\widetilde{\text{Tr}}_{L^2}(\tilde{g}) = q.$$

We first consider the case $\nu_1 < \epsilon_P$. For any $\gamma \in \Gamma_{\leq q}$ with $\gamma = P_g^+ \gamma P_g^+$, we write

$$\begin{aligned} & \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \epsilon_P)(\gamma_\xi - \tilde{g}_\xi)] d\xi \\ &= \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \nu_1)(\gamma_\xi - \tilde{g}_\xi)] d\xi + \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(\nu_1 - \epsilon_P)(\gamma_\xi - \tilde{g}_\xi)] d\xi \\ &= \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \nu_1)(\gamma_\xi - \tilde{g}_\xi)] d\xi + (\nu_1 - \epsilon_P) \left(\widetilde{\text{Tr}}_{L^2}(\gamma) - q \right) \end{aligned}$$

$$\geq \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \nu_1)(\gamma_\xi - \tilde{g}_\xi)] d\xi. \quad (4.29)$$

On the other hand, we have

$$\begin{aligned} & \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \nu_1)(\gamma_\xi - \tilde{g}_\xi)] d\xi \\ &= \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{g,\xi} - \nu_1)(\gamma_\xi - 1_{[0,\nu_1]}(D_{g,\xi}))] d\xi \\ &= \int_{Q_\ell^*} \sum_{\lambda_j(\xi) < \nu_1} |\lambda_j(\xi) - \nu_1| |(\gamma_\xi \psi_j(\xi), \psi_j(\xi)) - 1| d\xi \\ &\quad + \int_{Q_\ell^*} \sum_{\lambda_j(\xi) > \nu_1} (\lambda_j(\xi) - \nu_1) (\gamma_\xi \psi_j(\xi), \psi_j(\xi)) d\xi \geq 0, \end{aligned} \quad (4.30)$$

since $0 \leq \gamma \leq 1_{L^2(\mathbb{R}^3)}$. Thus \tilde{g} is a minimizer. Conversely, if \hat{g} is a minimizer, then it must be of the form $f_{Q_\ell^*}^\oplus 1_{[0,\nu]}(D_{g,\xi}) d\xi + \hat{\delta}$ with $0 \leq \hat{\delta} \leq f_{Q_\ell^*}^\oplus 1_{\{\nu_1\}}(D_{g,\xi}) d\xi$, $\nu = \nu_1 \in [\lambda_0, \min\{\epsilon_P, e_0\}]$ and $\widetilde{\text{Tr}}_{L^2}(\hat{g}) = q$ since all inequalities in (4.29) and (4.30) above have to be equalities for $g = \hat{g}$.

For the case $\epsilon_P \leq \nu_1$, we prove that $g' := f_{Q_\ell^*}^\oplus \text{Tr}_{L_\xi^2}[1_{[0,\epsilon_P]}(D_{g,\xi})] d\xi$ is a minimizer with $\widetilde{\text{Tr}}_{L^2}(g') \leq q$, thanks to a modified version of (4.30) with ν_1 (resp. \tilde{g}) being replaced by ϵ_P (resp. g'). As in the previous case, every minimizer \hat{g} satisfies $\hat{g} = f_{Q_\ell^*}^\oplus 1_{[0,\epsilon_P]}(D_{g,\xi}) d\xi + \hat{\delta}$, with $0 \leq \hat{\delta} \leq f_{Q_\ell^*}^\oplus 1_{\{\epsilon_P\}}(D_{g,\xi}) d\xi$. Note that in the case $\epsilon_P \leq \nu_1$, the inequality $\widetilde{\text{Tr}}_{L^2}(\hat{g}) \leq q$ automatically holds for any such \hat{g} .

In both cases, thanks to (4.27) we have $\nu = \min\{\nu_1, \epsilon_P\} \leq e_0 < e_1$, hence $0 \leq \hat{g} \leq 1_{[0,e_1]}(D_g)$. Thus, Corollary 4.13 implies that $\hat{g} \in \overline{\mathbf{B}}$. \square

Remark 4.16. *Actually, it follows from Corollary 4.13 that for every minimizer \hat{g} and a.e. $\xi \in Q_\ell^*$, one has $\text{Rank}(\hat{g}_\xi) \leq j_1$.*

4.4 First properties of minimizing sequences for $J_{\leq q}$

We introduce the sublevel set

$$\mathcal{S} := \left\{ \gamma \in \Gamma_{\leq q}^+ \mid \mathcal{E}^{DF}(\gamma) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma) \leq 0 \right\}. \quad (4.31)$$

Note that the operator 0 belongs to $\Gamma_{\leq q}^+$ and satisfies $\mathcal{E}^{DF}(0) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(0) = 0$. Thus,

$$J_{\leq q} = \inf_{\gamma \in \mathcal{S}} \left[\mathcal{E}^{DF}(\gamma) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma) \right] \quad (4.32)$$

and from now on we will only consider minimizing sequences for $J_{\leq q}$ lying in the sublevel set \mathcal{S} . These sequences satisfy *a priori* estimates gathered in the following lemma.

Lemma 4.17 (Boundedness of \mathcal{S}). *Assume that $\kappa < 1$. If $\kappa < 1 - \frac{\alpha}{2} C_{EE} q^+$, then, for every $\gamma \in \mathcal{S}$,*

$$\|\gamma\|_Z \leq \max \left\{ (1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} \epsilon_P q; 1 \right\} \quad (4.33)$$

and

$$\max \left\{ \|\gamma\|_{D^0}^{1/2} \|\gamma\|_{\mathfrak{S}_{1,1}}; \|\gamma\|_Y \right\} \leq \sqrt{\max \left\{ (1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} \epsilon_P q; 1 \right\}} q^+. \quad (4.34)$$

Proof. As $D_\gamma \gamma = |D_\gamma| \gamma$ for any $\gamma \in \Gamma_{\leq q}^+$, we get, by (4.13) and (4.17),

$$\begin{aligned} \mathcal{E}^{DF}(\gamma) - \epsilon_P \widetilde{\text{Tr}}_{L^2}(\gamma) &= \widetilde{\text{Tr}}_{L^2}[(D_\gamma - \epsilon_P - \frac{\alpha}{2} V_\gamma) \gamma] = \widetilde{\text{Tr}}_{L^2}[(|D_\gamma| - \epsilon_P - \frac{\alpha}{2} V_\gamma) \gamma] \\ &\geq \widetilde{\text{Tr}}_{L^2}[((1 - \kappa) |D^0| - \epsilon_P - \frac{\alpha}{2} V_\gamma) \gamma] \\ &\geq (1 - \kappa) \|\gamma\|_X - \frac{\alpha}{2} C_{EE} \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \|\gamma\|_X - \epsilon_P \|\gamma\|_{\mathfrak{S}_{1,1}} \\ &\geq (1 - \kappa - \frac{\alpha}{2} C_{EE} q^+) \|\gamma\|_X - \epsilon_P q. \end{aligned}$$

Hence, for any $\gamma \in \mathcal{S}$,

$$(1 - \kappa - \frac{\alpha}{2} C_{EE} q^+) \|\gamma\|_X - \epsilon_P q \leq 0.$$

Whenever $1 - \kappa - \frac{\alpha}{2} C_{EE} q^+ > 0$, (4.33) holds since $\|\gamma\|_Y \leq 1$.

The estimate (4.34) follows from Hölder's inequality and the fact that $\gamma \geq 0$; namely

$$\|\gamma |D^0|^{1/2}\|_{\mathfrak{S}_{1,1}} \leq \|\gamma^{1/2}\|_{\mathfrak{S}_{2,2}} \| |D^0|^{1/2} \|_{\mathfrak{S}_{2,2}} \leq \|\gamma\|_{\mathfrak{S}_{1,1}}^{1/2} \|\gamma\|_X^{1/2}.$$

□

5 Approximation by a linearized problem

The aim of this section is to show the link between a minimizing sequence $(\gamma_n)_{n \geq 1}$ in \mathcal{S} and the linearized Dirac–Fock problem introduced in Lemma 4.15.

Proposition 5.1 (Link with the linearized problem). *Under Assumption 3.1, let $(\gamma_n)_{n \geq 1} \in \mathcal{S}^{\mathbb{N}^*}$ be a minimizing sequence for $J_{\leq q}$. Then, as n goes to infinity,*

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma_n, \xi} - \epsilon_P) \gamma_{n, \xi}] d\xi - \inf_{\substack{\gamma \in \Gamma_{\leq q}^+ \\ \gamma = P_{\gamma_n}^+ \gamma P_{\gamma_n}^+}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma, \xi} - \epsilon_P) \gamma_\xi] d\xi \rightarrow 0. \quad (5.1)$$

This property is used in Lemma 6.2 below to build a new minimizing sequence with further regularity, and it is also used at the end of Section 6 to show some properties of the minimizers for $J_{\leq q}$.

As mentioned at the end of Section 3, the main difficulty in the proof of Proposition 5.1 is to deal with the nonconvex constraint set $\Gamma_{\leq q}^+$. To do so, we adapt to our setting a retraction technique introduced in [39]. We are going to build an open subset \mathcal{U} of Z stable under the continuous map $T : \gamma \mapsto P_\gamma^+ \gamma P_\gamma^+$ and such that the sequence $(T^p)_{p \geq 1}$ converges uniformly on $\overline{\mathcal{U}}$ to a surjective map $\theta : \overline{\mathcal{U}} \rightarrow \overline{\mathcal{U}} \cap \text{Fix}(T)$. Here, $\overline{\mathcal{U}}$ is the closure of \mathcal{U} in Z and $\text{Fix}(T)$ is the set of fixed points of T . The map θ will be uniformly continuous and such that $\theta \circ \theta = \theta$. Following a classical terminology we call it a retraction of $\overline{\mathcal{U}}$ onto $\overline{\mathcal{U}} \cap \text{Fix}(T)$. The restriction of θ to \mathcal{U} will be of class C^1 , the differential map $d\theta$ being itself uniformly continuous and bounded from \mathcal{U} to the space $\mathcal{B}(Z)$ of bounded linear operators on Z .

Then we will consider the subset $\mathcal{V} := \mathcal{U} \cap \Gamma_{\leq q}^+$, which is relatively open in $\Gamma_{\leq q}^+$ for the topology of Z . We will see that $\theta(\overline{\mathcal{V}}) \subset \overline{\mathcal{V}} \cap \Gamma_{\leq q}^+$ and $\overline{\mathcal{V}} \cap \text{Fix}(\theta) = \overline{\mathcal{V}} \cap \Gamma_{\leq q}^+$, so θ may be considered as a retraction of $\overline{\mathcal{V}}$ onto $\overline{\mathcal{V}} \cap \Gamma_{\leq q}^+$. Under our assumptions, we will prove the inclusion $\mathcal{S} \subset \mathcal{V}$ which implies, in combination with (4.32), the equality

$$J_{\leq q} = \inf_{\gamma \in \mathcal{V}} \left\{ \mathcal{E}^{DF}(\theta(\gamma)) - \epsilon_P \widetilde{\text{Tr}}_{L^2}[\theta(\gamma)] \right\}.$$

It will even turn out that \mathcal{U} is a uniform neighborhood of \mathcal{S} in Z , that is,

$$\mathcal{S} + B_Z(0, \rho) \subset \mathcal{U}$$

for some positive constant ρ . This property, combined with a formula for the differential of θ , will allow us to prove Proposition 5.1.

Before giving the definition of \mathcal{U} , we take $r > 0$ very small, and we introduce the following set, as in [39]:

$$\Gamma_{\leq q, r} := \{ \gamma \in Z \mid \text{dist}_{\mathfrak{S}_{1,1} \cap Y}(\gamma, \Gamma_{\leq q}) < r \}.$$

Then analogously to Lemmas 4.10 and 4.11, we have for any $\gamma \in \Gamma_{\leq q, r}$,

$$(1 - \kappa_r) |D^0| \leq |D_\gamma| \leq (1 + \kappa_r) |D^0| \quad (5.2)$$

and

$$\inf \sigma(|D_\gamma|) \geq \lambda_{0, r} \geq 1 - \kappa_r, \quad (5.3)$$

where $\kappa_r := \alpha (C_G z + C'_{EE}(q^+ + 2r))$ and

$$\lambda_{0,r} := 1 - \alpha \max \left\{ C_{HZ} + C_{EE}r + C''_{EE}(q^+ + r); \frac{C_0}{\ell} z + C_{EE}(q^+ + r) \right\}.$$

Definition 5.2 (Admissible set for the retraction). *Assume that $\kappa_r < 1$. Let $a_r := \frac{\alpha}{2} C_{EE} (1 - \kappa_r)^{-1/2} \lambda_{0,r}^{-1/2}$. Given $0 < \tau < \frac{1}{2a_r}$, let $M := \max \left\{ \frac{2+a_r(q^++r)}{2}; \frac{1}{1-2a_r\tau} \right\}$. We then define*

$$\mathcal{U} := \left\{ \gamma \in \Gamma_{\leq q,r} \mid \max \{ \|\gamma\|_{D^0}^{1/2} \| \mathfrak{S}_{1,1} \}; \|\gamma\|_Y \} + M \|T(\gamma) - \gamma\|_Z < \tau \right\}.$$

We have the following result.

Proposition 5.3 (Existence and differentiability of the retraction). *Take $\kappa_r, a_r, \tau, \mathcal{U}$ as in Definition 5.2. Let $k := 2a_r\tau$ and $\mathcal{V} := \mathcal{U} \cap \Gamma_{\leq q}^+$. Then the sequence of iterated maps $(T^p)_{p \geq 1}$ converges uniformly on $\overline{\mathcal{U}}$ to a limit θ with $\theta(\overline{\mathcal{U}}) = \text{Fix}(T) \cap \overline{\mathcal{U}}$, $\theta(\overline{\mathcal{V}}) = \Gamma_{\leq q}^+ \cap \overline{\mathcal{V}}$ and $\theta \circ \theta = \theta$. We have the estimate*

$$\forall \gamma \in \overline{\mathcal{U}}, \|\theta(\gamma) - T^p(\gamma)\|_Z \leq \frac{k^p}{1-k} \|T(\gamma) - \gamma\|_Z. \quad (5.4)$$

Moreover $\theta \in C^{1,\text{unif}}(\mathcal{U}, Z)$ and $d\theta(T^p)$ converges uniformly to $d\theta$ on \mathcal{U} .

In this way we obtain a continuous retraction θ of $\overline{\mathcal{U}}$ onto $\Gamma_{\leq q}^+ \cap \overline{\mathcal{U}}$ whose restriction to \mathcal{U} is of class $C^{1,\text{unif}}$. This map and its differential are bounded and uniformly continuous on \mathcal{U} .

For any $\gamma \in \text{Fix}(T) \cap \mathcal{U}$ and $\xi \in Q_\ell^*$, the linear operator $h \mapsto d\theta_\xi(\gamma)h$ satisfies

$$P_{\gamma,\xi}^+ d\theta_\xi(\gamma)h P_{\gamma,\xi}^+ = P_{\gamma,\xi}^+ h_\xi P_{\gamma,\xi}^+ \quad \text{and} \quad P_{\gamma,\xi}^- d\theta_\xi(\gamma)h P_{\gamma,\xi}^- = 0,$$

where $\theta(\gamma) = \int_{Q_\ell^*}^\oplus \theta_\xi(\gamma) d\xi$, according to the Floquet-Bloch decomposition. In other words, the splitting $L_\xi^2 = P_{\gamma,\xi}^+ L_\xi^2 \oplus P_{\gamma,\xi}^- L_\xi^2$ gives a block decomposition of $d\theta_\xi(\gamma)h$ of the form

$$d\theta_\xi(\gamma)h = \begin{pmatrix} P_{\gamma,\xi}^+ h_\xi P_{\gamma,\xi}^+ & b_{\gamma,\xi}(h)^* \\ b_{\gamma,\xi}(h) & 0 \end{pmatrix}. \quad (5.5)$$

The proof of Proposition 5.3 is postponed to the end of this section.

To apply Proposition 5.3 to the proof of Proposition 5.1, we need to find $\tau \in \left(0, \frac{1}{2a_r}\right)$ such that \mathcal{U} is a uniform neighborhood of \mathcal{S} . From Lemma 4.17 and the definition of \mathcal{U} , we can observe that if

$$\tau > \sqrt{\max \left\{ (1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} \epsilon_P q; 1 \right\} q^+}.$$

then there is $\rho > 0$ such that for every $\gamma \in \mathcal{S}$, one has the inclusion $B_Z(\gamma, \rho) \subset \mathcal{U}$. Thus, we have the following.

Lemma 5.4. *Assume that $\kappa < 1 - \frac{\alpha}{2} C_{EE} q^+$, and let a_r be as above. Assume in addition that*

$$2a_r \sqrt{\max \left\{ (1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} \epsilon_P q; 1 \right\} q^+} < 1.$$

Then there exist $\tau \in \left(0, \frac{1}{2a_r}\right)$ and $\rho > 0$ such that $\mathcal{S} + B_Z(0, \rho) \subset \mathcal{U}$.

We are now in a position to prove the main result of this section.

Proof of Proposition 5.1 (as a consequence of Proposition 5.3). Under Assumption 3.1, we may choose $r > 0$ so small that the assumptions of Lemma 5.4 hold true. Then we may take $\tau \in \left(0, \frac{1}{2a_r}\right)$ and $\rho > 0$ satisfying the conclusion of this lemma. To prove (5.1), we argue by contradiction. Otherwise, there would be an $\epsilon_0 > 0$ such that, for n large enough,

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma_{n,\xi}} - \epsilon_P) \gamma_{n,\xi}] d\xi \geq \inf_{\substack{\gamma \in \Gamma_{\leq q} \\ \gamma = P_{\gamma_n}^+ \gamma}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma_{n,\xi}} - \epsilon_P) \gamma_\xi] d\xi + \epsilon_0.$$

By Lemma 4.15, there exists an operator $\widehat{\gamma}_n \in \Gamma_{\leq q}$ which solves the following minimization problem:

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma_n, \xi} - \epsilon_P) \widehat{\gamma}_{n, \xi}] d\xi = \min_{\substack{\gamma \in \Gamma_{\leq q} \\ \gamma = P_{\gamma_n}^+ \gamma}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2} [(D_{\gamma_n, \xi} - \epsilon_P) \gamma_\xi] d\xi.$$

From Lemma 4.15 and Proposition 4.14, $\|\widehat{\gamma}_n\|_Z$ is uniformly bounded. So according to Corollary 5.4, there is $\sigma > 0$ such that for any n large enough and any $s \in [0, \sigma]$, $(1-s)\widehat{\gamma}_n + s\widehat{\gamma}_n \in \Gamma_{\leq q} \cap B_Z(\gamma_n, \rho) \subset \mathcal{V}$. Then, from Proposition 5.3, the function $f_n : [0, \sigma] \ni s \mapsto (\mathcal{E}^{DF} - \epsilon_P \widetilde{\text{Tr}}_{L^2})(\theta[(1-s)\gamma_n + s\widehat{\gamma}_n])$ is of class C^1 , and the sequence of derivatives (f'_n) is equicontinuous on $[0, \sigma]$. From (5.5), we infer

$$f'_n(0) = \widetilde{\text{Tr}}_{L^2} [(D_{\gamma_n} - \epsilon_P)(\widehat{\gamma}_n - \gamma_n)] \leq -\frac{\epsilon_0}{2}.$$

So there is $0 < s_0 < \sigma$ independent of n such that for any $s \in [0, s_0]$ we have $f'_n(s) \leq -\frac{\epsilon_0}{4}$. Hence,

$$(\mathcal{E}^{DF} - \epsilon_P \widetilde{\text{Tr}}_{L^2})(\theta[(1-s_0)\gamma_n + s_0\widehat{\gamma}_n]) = f_n(s_0) \leq f_n(0) - \frac{\epsilon_0 s_0}{4} = (\mathcal{E}^{DF} - \epsilon_P \widetilde{\text{Tr}}_{L^2})(\gamma_n) - \frac{\epsilon_0 s_0}{4}.$$

Recalling that $(\mathcal{E}^{DF} - \epsilon_P \widetilde{\text{Tr}}_{L^2})(\gamma_n)$ converges to $J_{\leq q}$, we conclude that $(\mathcal{E}^{DF} - \epsilon_P \widetilde{\text{Tr}}_{L^2})(\theta[(1-s_0)\gamma_n + s_0\widehat{\gamma}_n]) < J_{\leq q}$ when n is large enough. This contradicts the definition of $J_{\leq q}$, since $\theta[(1-s)\gamma_n + s\widehat{\gamma}_n] \in \Gamma_{\leq q}^+$. Hence the proposition. \square

It remains to prove Proposition 5.3, but before that, we need some preliminary results. Recall that $P_0^+ = 1_{\mathbb{R}^+}(D^0 - \alpha z G_\ell)$. We have the following lemma.

Lemma 5.5. *Take κ_r, a_r as in Definition 5.2 and consider the map*

$$Q : \gamma \mapsto P_\gamma^+ - P_0^+,$$

that is, $Q(\gamma) := \int_{Q_\ell^*}^\oplus Q_\xi(\gamma) d\xi$ with $Q_\xi(\gamma) := P_{\gamma, \xi}^+ - P_{0, \xi}^+$.

Then Q is in $C^{1, \text{lip}}(\Gamma_{\leq q, r}; |D^0|^{-1/2} Y)$ and we have the estimates

$$\forall \gamma \in \Gamma_{\leq q, r}, \quad \forall h \in Z, \quad \||D^0|^{1/2} dQ(\gamma)h\|_Y \leq a_r \|h\|_Z \quad (5.6)$$

and

$$\forall \gamma, \gamma' \in \Gamma_{\leq q, r}, \quad \||D^0|^{1/2} [dQ(\gamma)h - dQ(\gamma')h]\|_Y \leq K \|\gamma - \gamma'\|_Z \|h\|_Z, \quad (5.7)$$

where K is a positive constant depending only on κ_r which remains bounded when κ_r stays away from 1.

Remark 5.6. In (5.6), if $\gamma \in \Gamma_{\leq q}$ one can replace a_r by the constant a introduced in Assumption 2.5, since $\lim_{r \rightarrow 0} a_r = a$.

Proof of Lemma 5.5. By Lemma 4.10, for every $\xi \in Q_\ell^*$ and for all $\gamma \in \Gamma_{\leq q, r}$, $D_{\gamma, \xi}$ is a self-adjoint operator, and 0 is in its resolvent set. Then by Taylor's formula [29, Chapter VI.5, Lemma 5.6] or [23], we have

$$P_{\gamma, \xi}^\pm = \frac{1}{2} \pm \frac{1}{2\pi} \int_{-\infty}^{+\infty} (D_{\gamma, \xi} - iz)^{-1} dz = \frac{1}{2} \pm \frac{1}{\pi} \int_0^{+\infty} D_{\gamma, \xi} (|D_{\gamma, \xi}|^2 + z^2)^{-1} dz \quad (5.8)$$

and, by the second resolvent identity,

$$Q_\xi(\gamma) = -\frac{\alpha}{2\pi} \int_{-\infty}^{+\infty} (D_{\gamma, \xi} - iz)^{-1} V_{\gamma, \xi} (D_{0, \xi} - iz)^{-1} dz.$$

Hence, for every $h \in Z$, we deduce from (5.8) and the second resolvent formula again, that

$$dQ_\xi(\gamma)h = dP_{\gamma, \xi}^+ h = -\frac{\alpha}{2\pi} \int_{-\infty}^{+\infty} (D_{\gamma, \xi} - iz)^{-1} V_{h, \xi} (D_{\gamma, \xi} - iz)^{-1} dz. \quad (5.9)$$

Besides, for any $u_\xi \in L_\xi^2(Q_\ell)$ and any $\gamma \in \Gamma_{\leq q, r}$, we have

$$\int_{-\infty}^{+\infty} \left(u_\xi, |D_{\gamma, \xi}|^{1/2} (|D_{\gamma, \xi}|^2 + |z|^2)^{-1} |D_{\gamma, \xi}|^{1/2} u_\xi \right)_{L_\xi^2} dz = \pi \|u_\xi\|_{L_\xi^2}^2. \quad (5.10)$$

We infer from (5.3) that

$$\| |D_\gamma|^{-1} \|_Y \leq \lambda_{0,r}^{-1}. \quad (5.11)$$

Thus, gathering (5.9), (5.10), (5.11) with Lemma 4.7, for any $\phi_\xi, \psi_\xi \in L_\xi^2$ we may write

$$\begin{aligned} & \left| (\psi_\xi, |D_\xi|^{1/2} (dQ_\xi(\gamma)h)\phi_\xi)_{L_\xi^2} \right| \\ &= \frac{\alpha}{2\pi} \left| \int_{-\infty}^{+\infty} (\psi_\xi, |D_\xi|^{1/2} (D_{\gamma,\xi} - iz)^{-1} V_{h,\xi} (D_{\gamma,\xi} - iz)^{-1} \phi_\xi)_{L_\xi^2} dz \right| \\ &\leq \frac{\alpha}{2\pi} \|V_{h,\xi}\|_{\mathcal{B}(L_\xi^2)} \left(\int_{-\infty}^{+\infty} \|(D_{\gamma,\xi} - iz)^{-1} |D_\xi|^{1/2} \psi_\xi\|_{L_\xi^2}^2 dz \right)^{1/2} \left(\int_{-\infty}^{+\infty} \|(D_{\gamma,\xi} - iz)^{-1} \phi_\xi\|_{L_\xi^2}^2 dz \right)^{1/2} \\ &\leq \frac{\alpha}{2} \|V_{h,\xi}\|_{\mathcal{B}(L_\xi^2)} \| |D_\xi|^{1/2} |D_{\gamma,\xi}|^{-1/2} \|_{\mathcal{B}(L_\xi^2)} \| |D_{\gamma,\xi}|^{-1/2} \|_{\mathcal{B}(L_\xi^2)} \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2} \\ &\leq \frac{\alpha}{2} C_{EE} (1 - \kappa_r)^{-1/2} \lambda_{0,r}^{-1/2} \|h\|_Z \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2} \end{aligned} \quad (5.12)$$

hence we obtain (5.6); namely,

$$\| |D^0|^{1/2} dQ(\gamma)h \|_Y \leq \frac{\alpha}{2} C_{EE} (1 - \kappa_r)^{-1/2} \lambda_{0,r}^{-1/2} \|h\|_Z.$$

For the second inequality, we write

$$\begin{aligned} dQ_\xi(\gamma)h - dQ_\xi(\gamma')h &= -\frac{\alpha^2}{2\pi} \int_{-\infty}^{+\infty} (D_{\gamma,\xi} - iz)^{-1} V_{\gamma'-\gamma,\xi} (D_{\gamma',\xi} - iz)^{-1} V_{h,\xi} (D_{\gamma,\xi} - iz)^{-1} dz \\ &\quad - \frac{\alpha^2}{2\pi} \int_{-\infty}^{+\infty} (D_{\gamma',\xi} - iz)^{-1} V_{h,\xi} (D_{\gamma,\xi} - iz)^{-1} V_{\gamma'-\gamma,\xi} (D_{\gamma',\xi} - iz)^{-1} dz. \end{aligned}$$

Proceeding as above, we get (5.7). The fact that $Q \in C^{1,\text{lip}}(\Gamma_{\leq q,r}; |D^0|^{-1/2} Y)$ follows from (5.6) and (5.7). \square

Lemma 5.7. *Take κ_r, a_r as in Definition 5.2. Then the map $T : \gamma \rightarrow P_\gamma^+ \gamma P_\gamma^+$ is well-defined and of class $\mathcal{C}^{1,1}$ on $\Gamma_{\leq q,r}$ with values in $\Gamma_{\leq q,r} \subset Z$. Moreover, for any $\gamma \in \Gamma_{\leq q,r}$,*

$$\begin{aligned} \|T^2(\gamma) - T(\gamma)\|_Z &\leq 2a_r \left(\max\{\|T(\gamma)\|_{\mathfrak{S}_{1,1}} \|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|T(\gamma)\|_Y\} \right. \\ &\quad \left. + \frac{a_r(q^+ + r)}{2} \|\gamma - T(\gamma)\|_Z \right) \|T(\gamma) - \gamma\|_Z. \end{aligned} \quad (5.13)$$

Moreover, there are two positive constants $C_{\kappa,r}, L_{\kappa,r}$ such that

$$\forall \gamma \in \Gamma_{\leq q,r}, \quad \|dT(\gamma)\|_{\mathcal{B}(Z)} \leq C_{\kappa,r} \left(1 + \max\{\|\gamma\|_{\mathfrak{S}_{1,1}} \|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|\gamma\|_Y\} \right) \quad (5.14)$$

and

$$\forall \gamma, \gamma' \in \Gamma_{\leq q,r}, \quad \|dT(\gamma') - dT(\gamma)\|_{\mathcal{B}(Z)} \leq L_{\kappa,r} \left(1 + \max\{\|\gamma\|_{\mathfrak{S}_{1,1}} \|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|\gamma\|_Y\} \right) \|\gamma' - \gamma\|_Z. \quad (5.15)$$

Proof. If $\gamma \in \Gamma_{\leq q,r}$, there is $\gamma_0 \in \Gamma_{\leq q,r}$ such that $\|\gamma - \gamma_0\|_{\mathfrak{S}_{1,1} \cap Y} < r$. Then $P_\gamma^+ \gamma_0 P_\gamma^+ \in \Gamma_{\leq q}$, $T(\gamma) \in Z$ and

$$\|T(\gamma) - P_\gamma^+ \gamma_0 P_\gamma^+\|_{\mathfrak{S}_{1,1} \cap Y} = \|P_\gamma^+ (\gamma - \gamma_0) P_\gamma^+\|_{\mathfrak{S}_{1,1} \cap Y} \leq \|\gamma - \gamma_0\|_{\mathfrak{S}_{1,1} \cap Y} < r,$$

so $T(\gamma) \in \Gamma_{\leq q,r}$.

Let $\gamma, \gamma' \in \Gamma_{\leq q,r}$. Then $P_\gamma^+ - P_{\gamma'}^+$ can be written as

$$P_\gamma^+ - P_{\gamma'}^+ = \int_0^1 dQ(\gamma' + t(\gamma - \gamma')) (\gamma - \gamma') dt.$$

From (5.6),

$$\| |D^0|^{1/2} (P_\gamma^+ - P_{\gamma'}^+) \|_Y \leq a_r \|\gamma - \gamma'\|_Z.$$

For the estimate (5.13), we write

$$\begin{aligned} T^2(\gamma) - T(\gamma) &= (P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma) \left(P_{T(\gamma)}^+ - P_\gamma^+ + P_\gamma^+ \right) + P_\gamma^+T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+) \\ &= (P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma) + T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+) + (P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+). \end{aligned}$$

Then

$$\begin{aligned} \|T^2(\gamma) - T(\gamma)\|_Z &\leq \|(P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma)\|_{X \cap Y} \\ &\quad + \|T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+)\|_{X \cap Y} + \|(P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+)\|_Z. \end{aligned}$$

We have

$$\|T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+)\|_{X \cap Y} \leq \| |D^0|^{1/2}(P_{T(\gamma)}^+ - P_\gamma^+) \|_Y \max\{\|T(\gamma)|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|T(\gamma)\|_Y\}$$

and

$$\|(P_{T(\gamma)}^+ - P_\gamma^+)T(\gamma)(P_{T(\gamma)}^+ - P_\gamma^+)\|_Z \leq \| |D^0|^{1/2}(P_{T(\gamma)}^+ - P_\gamma^+) \|_Y^2 \|T(\gamma)\|_{\mathfrak{S}_{1,1 \cap Y}}.$$

Notice that $\|T(\gamma)\|_{\mathfrak{S}_{1,1 \cap Y}} \leq \|\gamma\|_{\mathfrak{S}_{1,1 \cap Y}} \leq q^+ + r$. Gathering together these estimates, we obtain (5.13).

We now turn to the proof of (5.14) and (5.15). From Lemma 5.5, T is in $C^1(\Gamma_{\leq q,r})$ with

$$dT(\gamma)h = (dQ_\gamma h)\gamma P_\gamma + P_\gamma \gamma (dQ_\gamma h) + P_\gamma h P_\gamma.$$

Notice that for any $\gamma \in \Gamma_{\leq q,r}$,

$$\| |D^0|^{1/2} P_\gamma^+ |D^0|^{-1/2} \|_Y \leq (1 - \kappa_r)^{-1/2} \| |D_\gamma|^{1/2} P_\gamma^+ |D^0|^{-1/2} \|_Y \leq \frac{(1 + \kappa_r)^{1/2}}{(1 - \kappa_r)^{1/2}}. \quad (5.16)$$

Thus, using (5.6), one finds a constant $C_{\kappa,r}$ such that for any $h \in Z$,

$$\begin{aligned} \|dT(\gamma)h\|_Z &\leq \max \left\{ 2 \| |D^0|^{1/2}(dQ_\gamma h) \|_Y \|\gamma\|_{\mathfrak{S}_{1,1}} \| |D^0|^{1/2} \|_{\mathfrak{S}_{1,1}} \| |D^0|^{1/2} P_\gamma^+ |D^0|^{-1/2} \|_Y + \| |D^0|^{1/2} P_\gamma^+ |D^0|^{-1/2} \|_Y^2 \|h\|_X; \right. \\ &\quad \left. 2 \|\gamma\|_Y \|dQ_\gamma h\|_Y + \|h\|_Y \right\} \\ &\leq C_{\kappa,r} \left(1 + \max \{ \|\gamma\|_{\mathfrak{S}_{1,1}} \| |D^0|^{1/2} \|_{\mathfrak{S}_{1,1}}; \|\gamma\|_Y \} \right) \|h\|_Z, \end{aligned}$$

so (5.14) is proved. Finally, for the term $dT(\gamma') - dT(\gamma)$, we have

$$\begin{aligned} dT_\xi(\gamma')h - dT_\xi(\gamma)h &= (dQ_{\gamma,\xi} h)\gamma_\xi P_{\gamma,\xi} + P_{\gamma,\xi} \gamma_\xi (dQ_{\gamma,\xi} h) + P_{\gamma,\xi} h_\xi P_{\gamma,\xi} \\ &\quad - (dQ_{\gamma',\xi} h)\gamma'_\xi P_{\gamma',\xi} - P_{\gamma',\xi} \gamma'_\xi (dQ_{\gamma',\xi} h) - P_{\gamma',\xi} h_\xi P_{\gamma',\xi}. \end{aligned}$$

Proceeding in the same way as for (5.14), we can get (5.15). \square

We now show that \mathcal{U} and T satisfy all the assumptions in [39, Proposition 2.2] in the Banach space Z .

Proposition 5.8. *Let $\kappa_r, a_r, \tau, \mathcal{U}$ be as in Definition 5.2. Then T is in $C^0(\overline{\mathcal{U}}) \cap C^{1, \text{lip}}(\mathcal{U}, Z)$. Moreover $T(\mathcal{U}) \subset \mathcal{U}$ and the following estimates are satisfied:*

$$\sup_{\gamma \in \mathcal{U}} \|dT(\gamma)\|_{\mathcal{B}(Z)} < \infty, \quad \sup_{\gamma \in \mathcal{U}} \|T(\gamma) - \gamma\|_Z < \infty$$

and

$$\forall \gamma \in \mathcal{U}, \quad \|T^2(\gamma) - T(\gamma)\|_Z \leq k \|T(\gamma) - \gamma\|_Z$$

with $k := 2a_r \tau < 1$.

Proof. For any $\gamma \in \mathcal{U}$, we have

$$\|T(\gamma)|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}} \leq \|\gamma|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}} + \|(\gamma - T(\gamma))|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}} \leq \|\gamma|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}} + \|\gamma - T(\gamma)\|_X$$

and

$$\|T(\gamma)\|_Y \leq \|\gamma\|_Y.$$

As a result, as $M \geq \frac{2+a_r(q^++r)}{2}$, (5.13) implies that

$$\|T^2(\gamma) - T(\gamma)\|_Z \leq k\|T(\gamma) - \gamma\|_Z.$$

Moreover, using the inequality $M \geq \frac{1}{1-2a_r\tau}$, we get

$$\begin{aligned} & \max \{ \|T(\gamma)|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|T(\gamma)\|_Y \} + M\|T^2(\gamma) - T(\gamma)\|_Z \\ & \leq \max \{ \|\gamma|D^0|^{1/2}\|_{\mathfrak{S}_{1,1}}; \|\gamma\|_Y \} + (1 + Mk)\|T(\gamma) - \gamma\|_Z < \tau, \end{aligned}$$

so $T(\gamma) \in \mathcal{U}$.

The fact that $\sup_{\gamma \in \mathcal{U}} \|dT(\gamma)\|_{\mathcal{B}(Z)} < \infty$ and that dT is Lipschitz continuous on \mathcal{U} follows from (5.14) and (5.15). Besides, for $\gamma \in \mathcal{U}$, we have $\|T(\gamma) - \gamma\|_Z < \frac{\tau}{M}$. This ends the proof of Proposition 5.8. \square

We can now prove Proposition 5.3, which implies Proposition 5.1, as we have already seen.

Proof of Proposition 5.3. By Proposition 5.8, we may apply [39, Proposition 2.2] to our map T and our open set \mathcal{U} in the Banach space Z . This allows us to construct $\theta \in C^0(\overline{\mathcal{U}}, Z) \cap C^{1,\text{unif}}(\mathcal{U}, Z)$ with the properties $\theta(\overline{\mathcal{U}}) = \text{Fix}(T) \cap \overline{\mathcal{U}}$, $\theta \circ \theta = \theta$ and the convergence estimate (5.4). By our definition of T , we have $T(\Gamma_{\leq q}^+) \subset \Gamma_{\leq q}^+$ and $\Gamma_{\leq q}^+ = \text{Fix}(T) \cap \Gamma_{\leq q}^+$, hence the additional property $\theta(\overline{\mathcal{V}}) = \Gamma_{\leq q}^+ \cap \overline{\mathcal{V}}$. The proof of (5.5) is exactly the same as in [39, Theorem 2.10]. This ends the proof of Proposition 5.3. \square

6 Proof of Theorem 3.3

Throughout this section, we assume that Assumption 3.1 is satisfied and that $\epsilon_P > e_0$. Let $(\gamma_n)_{n \geq 1}$ be a minimizing sequence for $J_{\leq q}$ lying in \mathcal{S} . According to Lemma 4.17, this sequence is uniformly bounded in Z . We split $(\gamma_n)_{n \geq 1}$ into two parts: $(\tilde{\gamma}_n)_{n \geq 1}$ and $(\gamma_n - \tilde{\gamma}_n)_{n \geq 1}$ where, for each n ,

$$\tilde{\gamma}_n := p_n \gamma_n p_n \quad \text{with} \quad p_n := 1_{[0, e_1]}(D\gamma_n) \quad (6.1)$$

where e_1 has been defined in Formula (4.26). Thanks to Corollary 4.13, for almost every $\xi \in Q_\ell^*$ the rank of $p_{n,\xi}$, and therefore of $\tilde{\gamma}_{n,\xi}$, is at most j_1 , so that $\tilde{\gamma}_n \in \overline{\mathbf{B}}$.

Actually, we prove in Lemma 6.1 that, for each $n \geq 1$, $\tilde{\gamma}_n \in X_\infty^2$ whereas $\gamma_n \in X$; roughly speaking, we reach a $L^\infty(Q_\ell^*; H_\xi^1(Q_\ell))$ regularity instead of a $L^2(Q_\ell^*; H_\xi^{1/2}(Q_\ell))$ regularity for the associated eigenfunctions. Moreover $\tilde{\gamma}_n$ is close to γ_n in X (Lemma 6.2), so $(\tilde{\gamma}_n)_{n \geq 1}$ is a modified minimizing sequence with higher regularity than $(\gamma_n)_{n \geq 1}$.

The structure of the proof of Theorem 3.3 is as follows. In Subsection 6.1, we will show that $\|\gamma_n - \tilde{\gamma}_n\|_X \rightarrow 0$ when n goes to infinity. In Subsection 6.2, we first study the convergence of the kernels of $(W_{\tilde{\gamma}_n, \xi})_{n \geq 1}$. Then we deduce the strong convergence of $(V_{\tilde{\gamma}_n, \xi})_{n \geq 1}$. As a result, $\|P_{\gamma_n}^+ - P_{\tilde{\gamma}_n}^+\|_Y \rightarrow 0$. On the other hand, for any $\gamma \in \Gamma_{\leq q}$, we also have $\|(P_{\gamma_n}^+ - P_{\tilde{\gamma}_n}^+)\gamma\|_{\mathfrak{S}_{1,1}} \leq C\|\tilde{\gamma}_n - \gamma_n\|_X^{1/2}\|\gamma\|_Z \rightarrow 0$. Hence in Subsection 6.3, we can pass to the limit in the energy and in the constraints.

6.1 Decomposition of minimizing sequences

We start with some regularity bounds on $\tilde{\gamma}_n$ that will be needed in the sequel.

Lemma 6.1. *The sequence $(\tilde{\gamma}_n)_{n \geq 1}$ and the sequence of kernels $(\tilde{\gamma}_{n,\xi}(\cdot, \cdot))_{n \geq 1}$ are uniformly bounded in X_∞^2 and in $L^\infty(Q_\ell^*; H^1(Q_\ell \times Q_\ell))$, respectively. More precisely, we have the following, for every $n \geq 1$ and for almost every ξ in Q_ℓ^* ,*

$$\|\tilde{\gamma}_n\|_{X_\infty^2} \leq j_1(1 - \kappa)^{-4} c^*(j_1)^2 \quad (6.2)$$

and

$$\|\tilde{\gamma}_{n,\xi}(\cdot, \cdot)\|_{H^1(Q_\ell \times Q_\ell)} \leq 2j_1(1 - \kappa)^{-4} c^*(j_1)^2. \quad (6.3)$$

Proof. We first prove that $\|p_n\|_{X_\infty^2}$ is bounded. Let $(u_{n,j}(\xi))_{j \geq 1}$ be the normalized eigenfunctions of the operator $D_{\gamma_n, \xi}$ with the corresponding eigenvalues $\lambda_{n,j}(\xi)$ counted with multiplicity. Hence,

$$p_{n,\xi} = \sum_{j=1}^{+\infty} \delta_{n,j}(\xi) |u_{n,j}(\xi)\rangle \langle u_{n,j}(\xi)|$$

with $\delta_{n,j}(\xi) = 1$ if $0 \leq \lambda_{n,j}(\xi) < e_1$ and $\delta_{n,j}(\xi) = 0$ otherwise.

By Corollary 4.13, we know that $|\{j \in \mathbb{N}^* \mid \delta_{n,j}(\xi) = 1\}| \leq j_1$. By (4.21), for any eigenfunction $u_{n,j}(\xi)$, we have $\delta_{n,j}(\xi) \|u_{n,j}(\xi)\|_{H_\xi^1(Q_\ell)}^2 \leq (1 - \kappa)^{-4} c^*(j_1)^2$, for every $\xi \in Q_\ell^*$. Now,

$$\|p_{n,\xi}\|_{X^2(\xi)} = \sum_{j=1}^{j_1} \delta_{n,k}(\xi) \|u_{n,k}(\xi)\|_{H_\xi^1}^2 \leq j_1 \sup_{j \geq 1} \delta_{n,j}(\xi) \|u_{n,j}(\xi)\|_{H_\xi^1}^2.$$

Hence,

$$\|p_n\|_{X_\infty^2} \leq j_1(1 - \kappa)^{-4} c^*(j_1)^2.$$

Since $p_n = p_n^2$, $\tilde{\gamma}_n = p_n \tilde{\gamma}_n p_n$ and $0 \leq \tilde{\gamma}_n \leq 1_{L^2(\mathbb{R}^3)}$, we have

$$\begin{aligned} \|\tilde{\gamma}_n\|_{X_\infty^2} &= \|D^0 |p_n \tilde{\gamma}_n p_n| D^0\|_{\mathfrak{S}_{1,\infty}} \leq \|\tilde{\gamma}_n\|_Y \|p_n\|_{X_\infty^2} \\ &\leq \|p_n\|_{X_\infty^2} \leq j_1(1 - \kappa)^{-4} c^*(j_1)^2. \end{aligned}$$

In terms of kernels, this implies that

$$\|D_{\xi,x} |\tilde{\gamma}_{n,\xi}(\cdot, \cdot)|\|_{L^2(Q_\ell \times Q_\ell)} = \|D_\xi |\tilde{\gamma}_{n,\xi}\|_{\mathfrak{S}_2(\xi)} \leq \|\tilde{\gamma}_{n,\xi}\|_{X^2(\xi)} \leq j_1(1 - \kappa)^{-4} c^*(j_1)^2,$$

the same holding for $|D_{\xi,y} |\tilde{\gamma}_{n,\xi}(\cdot, \cdot)|$. Thus, $\tilde{\gamma}_{n,\xi}(\cdot, \cdot) \in L^\infty(Q_\ell^*; H^1(Q_\ell \times Q_\ell))$ and (6.3) holds. \square

We begin the proof of Theorem 3.3 by showing the following result as in the case of molecules [39, Lemma 3.4].

Lemma 6.2. *Under Assumption 3.1, whenever $\epsilon_P > e_0$, for any minimizing sequence $(\gamma_n)_{n \geq 1}$ of (3.1) in $\Gamma_{\leq q}^+$ we have*

$$\widetilde{\text{Tr}}_{L^2}[\gamma_n] \rightarrow q \quad \text{and} \quad \|\gamma_n - \tilde{\gamma}_n\|_X \rightarrow 0.$$

Proof. According to Proposition 5.1, any minimizing sequence $(\gamma_n)_{n \geq 1}$ in $\Gamma_{\leq q}^+$ satisfies (5.1); namely

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{\gamma_n, \xi} - \epsilon_P)\gamma_n, \xi] d\xi - \inf_{\substack{\gamma \in \Gamma_{\leq q}^+ \\ \gamma = P_{\gamma_n}^+ \gamma P_{\gamma_n}^+}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{\gamma, \xi} - \epsilon_P)\gamma_\xi] d\xi \rightarrow 0.$$

By Lemma 4.15, for every n , minimizers of the above minimization problem are of the form $\gamma'_n := \int_{Q_\ell^*}^\oplus 1_{[0, \nu_n]}(D_{\gamma_n, \xi}) d\xi + \delta_n$ for some $\nu_n \in [\lambda_0, e_0]$ and some $0 \leq \delta_n \leq \int_{Q_\ell^*}^\oplus 1_{\nu_n}(D_{\gamma_n, \xi}) d\xi$ such that $\widetilde{\text{Tr}}_{L^2}(\gamma'_n) = q$. In particular, $\limsup_{n \rightarrow +\infty} \nu_n < \epsilon_P$, for every n . We define

$$\pi_n := \int_{Q_\ell^*}^\oplus 1_{[e_1, \infty)}(D_{\gamma_n, \xi}) d\xi, \quad \pi'_n := \int_{Q_\ell^*}^\oplus 1_{(\nu_n, e_1)}(D_{\gamma_n, \xi}) d\xi, \quad \pi''_n := \int_{Q_\ell^*}^\oplus 1_{[0, \nu_n]}(D_{\gamma_n, \xi}) d\xi.$$

We can write $p_n = \pi'_n + \pi''_n$ and we observe that $\gamma'_n = \pi''_n \gamma'_n \pi''_n$. Proceeding as for (4.29) and (4.30), and since $\gamma_n \in \Gamma_{\leq q}^+$ we have

$$\begin{aligned} & \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \epsilon_P)\gamma_n] - \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \epsilon_P)\gamma'_n] \\ &= \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \nu_n)\pi_n \gamma_n \pi_n] + \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \nu_n)\pi'_n \gamma_n \pi'_n] \\ & \quad + \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \nu_n)(\pi''_n \gamma_n \pi''_n - 1_{[0, \nu_n]}(D_{\gamma_n}))] + (\epsilon_P - \nu_n) \left(q - \widetilde{\text{Tr}}_{L^2}(\gamma_n) \right). \end{aligned}$$

We observe that the four terms in the right-hand side of the above equation are non-negative whereas, from Proposition 5.1, their sum goes to 0 as n goes to infinity. Therefore,

$$\widetilde{\text{Tr}}_{L^2}[\gamma_n] \rightarrow q \text{ and } \widetilde{\text{Tr}}_{L^2}[(D_{\gamma_n} - \nu_n)\pi_n \gamma_n \pi_n] \rightarrow 0,$$

since $\liminf_{n \rightarrow +\infty} (\epsilon_P - \nu_n) \geq \epsilon_P - e_0 > 0$. But $\pi_n(D_{\gamma_n} - \nu_n)\pi_n \geq (e_1 - \nu_n)\pi_n \geq (e_1 - e_0)\pi_n$ and $\pi_n(D_{\gamma_n} - \nu_n)\pi_n \geq \pi_n(|D_{\gamma_n}| - e_0)\pi_n$. Thus, taking a convex combination of these two estimates leads to

$$\frac{e_1}{e_1 - e_0} \pi_n(D_{\gamma_n} - \nu_n)\pi_n \geq \pi_n |D_{\gamma_n}| \pi_n.$$

Hence

$$\|\pi_n \gamma_n \pi_n\|_X = \widetilde{\text{Tr}}_{L^2}[\pi_n |D^0| \pi_n \gamma_n] \leq (1 - \kappa)^{-1} \widetilde{\text{Tr}}_{L^2}[\pi_n |D_{\gamma_n}| \pi_n \gamma_n],$$

and the right-hand side goes to 0 by (4.17). It remains to study the limit of $h_n := \pi_n \gamma_n p_n$ as n goes to infinity. Since $(\gamma_n)^2 \leq \gamma_n$, we have

$$(\pi_n \gamma_n \pi_n)^2 + h_n h_n^* = \pi_n (\gamma_n)^2 \pi_n \leq \pi_n \gamma_n \pi_n.$$

Hence

$$\widetilde{\text{Tr}}_{L^2}(|D_{\gamma_n}|^{1/2} h_n h_n^* |D_{\gamma_n}|^{1/2}) \rightarrow 0.$$

In other words, $\| |D_{\gamma_n}|^{1/2} h_n \|_{\mathfrak{S}_{2,2}} \rightarrow 0$. Taking any operator A in Y , by the Cauchy-Schwarz inequality,

$$\begin{aligned} \left| \widetilde{\text{Tr}}_{L^2}[A |D_{\gamma_n}|^{1/2} h_n^* |D_{\gamma_n}|^{1/2}] \right| &= \left| \widetilde{\text{Tr}}_{L^2}[A |D_{\gamma_n}|^{1/2} p_n h_n^* |D_{\gamma_n}|^{1/2}] \right| \\ &\leq \|A |D_{\gamma_n}|^{1/2} p_n\|_{\mathfrak{S}_{2,2}} \| |D_{\gamma_n}|^{1/2} h_n \|_{\mathfrak{S}_{2,2}}. \end{aligned} \quad (6.4)$$

We have already seen that $p_{n,\xi}$ has rank at most j_1 . Therefore,

$$\|A |D_{\gamma_n}|^{1/2} p_n\|_{\mathfrak{S}_{2,2}} \leq \| |D_{\gamma_n}|^{1/2} p_n \|_{\mathfrak{S}_{2,2}} \|A\|_Y \leq j_1 e_1 \|A\|_Y,$$

so we deduce immediately from (6.4) that

$$\left\| |D_{\gamma_n}|^{1/2} h_n |D_{\gamma_n}|^{1/2} \right\|_{\mathfrak{S}_{1,1}} \rightarrow 0,$$

since A is arbitrary. Hence, thanks to (4.15), $\|h_n\|_X = \| |D^0|^{1/2} h_n |D^0|^{1/2} \|_{\mathfrak{S}_{1,1}} \rightarrow 0$. Finally, we obtain that $\|\gamma_n - \tilde{\gamma}_n\|_X \leq \|\pi_n \gamma_n \pi_n\|_X + 2 \|h_n\|_X \rightarrow 0$. \square

By Lemma 6.1, up to the extraction of a subsequence, there is γ_* in $X_\infty^2 \cap Y$, such that

$$\tilde{\gamma}_n \xrightarrow{*} \gamma_* \quad \text{for the weak-}^* \text{ convergence in } X_\infty^2 \cap Y, \quad (6.5)$$

since X_∞^2 is a subspace of $\mathfrak{S}_{1,\infty}$ which is the dual space of $\mathfrak{S}_{\infty,1}$ and Y is the dual space of $\mathfrak{S}_{1,1}$. We immediately get the following.

Lemma 6.3 (Strong convergence of the density). *The sequence $\rho_{\tilde{\gamma}_n}^{1/2}$ converges strongly to $\rho_{\gamma_*}^{1/2}$ in $H^s(Q_\ell)$ with $0 \leq s < 1$, thus in $L^p(Q_\ell)$ for every $1 \leq p < 6$. In particular, whenever $\epsilon_P > e_0$, we have $\int_{Q_\ell} \rho_{\gamma_*} dx = q$.*

Proof. The proof of the strong convergence of $\rho_{\tilde{\gamma}_n}^{1/2}$ to $\rho_{\gamma_*}^{1/2}$ in $L^p(Q_\ell)$ for every $1 \leq p < 6$ is the same as in [8, p. 730] and relies on the fact that $\tilde{\gamma}_n \in X_\infty^2$ (see the proofs of [8, Eqs. (4.51) and (4.55)]). When $\epsilon_P > e_0$, by Lemma 6.2, $\tilde{\gamma}_n - \gamma_n$ converges to 0 in X , thus in $\mathfrak{S}_{1,1}$, whereas $\widetilde{\text{Tr}}_{L^2}[\gamma_n]$ converges to q . Thus, $\widetilde{\text{Tr}}_{L^2}[\gamma_*] = q$. \square

6.2 Convergence of $(P_{\gamma_n}^+)_{n \geq 1}$

We now study the differences between $P_{\gamma_n}^+$ and $P_{\tilde{\gamma}_n}^+$ and between $P_{\tilde{\gamma}_n}^+$ and $P_{\gamma_*}^+$ separately. We do not know whether $\gamma_n - \tilde{\gamma}_n$ goes to 0 in Y and we do not even know whether $\tilde{\gamma}_n - \gamma_*$ goes to 0 in $\mathfrak{S}_{1,1}$, so we cannot rely on the continuity of the map Q introduced in Lemma 5.5. The proof is therefore more involved than in [39].

Convergence of $(P_{\tilde{\gamma}_n}^+ - P_{\gamma_*}^+)_{n \geq 1}$

The main result of this section is Corollary 6.8 which states that the sequence $(P_{\tilde{\gamma}_n}^+ - P_{\gamma_*}^+)_{n \geq 1}$ converges strongly in Y .

Recall that

$$W_\gamma = W_{\geq m, \gamma} + W_{< m, \gamma}, \quad \forall m \in \mathbb{N}, m \geq 2$$

where for $\xi \in Q_\ell^*$ and $x, y \in Q_\ell$, the kernels of $W_{\geq m, \gamma, \xi}$ and $W_{< m, \gamma, \xi}$ are respectively

$$W_{\geq m, \gamma, \xi}(x, y) := \int_{Q_\ell^*} W_{\geq m, \ell}^\infty(\xi' - \xi, x - y) \gamma_{\xi'}(x, y) d\xi' \quad (6.6)$$

and

$$W_{< m, \gamma, \xi}(x, y) := \int_{Q_\ell^*} W_{< m, \ell}^\infty(\xi' - \xi, x - y) \gamma_{\xi'}(x, y) d\xi'. \quad (6.7)$$

We first prove the following.

Lemma 6.4. *The two sequences of operators $(W_{< m, \tilde{\gamma}_n})_n$ and $(W_{\geq m, \tilde{\gamma}_n})_n$ are bounded in $\mathfrak{S}_{2, \infty}$. Thus, up to the extraction of a subsequence, we may assume that*

$$W_{< m, \tilde{\gamma}_n} \xrightarrow{*} W_{< m, \gamma_*} \quad \text{and} \quad W_{\geq m, \tilde{\gamma}_n} \xrightarrow{*} W_{\geq m, \gamma_*} \quad \text{in} \quad \mathfrak{S}_{2, \infty}. \quad (6.8)$$

Note that saying that the operator $A = \int_{Q_\ell^*}^\oplus A_\xi d\xi$ belongs to $\mathfrak{S}_{2, p}$ with $1 \leq p \leq +\infty$ is equivalent to saying that the family of kernels $\xi \mapsto A_\xi(\cdot, \cdot)$ is in $L^p(Q_\ell^*; L^2(Q_\ell \times Q_\ell))$.

Proof. The first claim follows from (B.23) and (B.17) in Appendix B, thanks to (6.2). Let $g = \int_{Q_\ell^*}^\oplus g_\xi d\xi \in \mathfrak{S}_{2,1}$. From (B.23), we know that $W_{< m, g} \in \mathfrak{S}_{2,1}$. Then, by (6.5), and since $\mathfrak{S}_{2,1}^* = \mathfrak{S}_{2, \infty}$, we get

$$\widetilde{\text{Tr}}_{L^2}[g W_{< m, \tilde{\gamma}_n}] = \widetilde{\text{Tr}}_{L^2}[W_{< m, g}^* \tilde{\gamma}_n] \rightarrow \widetilde{\text{Tr}}_{L^2}[W_{< m, g}^* \gamma_*] = \widetilde{\text{Tr}}_{L^2}[g W_{< m, \gamma_*}].$$

The argument is similar for $W_{\geq m, \tilde{\gamma}_n}$. Notice that by (B.17), $W_{\geq m, \tilde{\gamma}_n} \in \mathfrak{S}_{2,1}$. Then we write

$$\widetilde{\text{Tr}}_{L^2}[W_{\geq m, \tilde{\gamma}_n} g] = \widetilde{\text{Tr}}_{L^2}[W_{\geq m, g} \tilde{\gamma}_n] = \widetilde{\text{Tr}}_{L^2}[|D^0|^{-1} W_{\geq m, g} |D^0|^{-1} |D^0| \tilde{\gamma}_n |D^0|],$$

with $|D^0| \tilde{\gamma}_n |D^0|$ converging to $|D^0| \gamma_* |D^0|$ for the weak-* topology of $\mathfrak{S}_{1, \infty}$ thanks to (6.5). Therefore, it remains to show that the operator $|D^0|^{-1} W_{\geq m, g} |D^0|^{-1}$ belongs to $\mathfrak{S}_{\infty, 1}$ whenever $g \in \mathfrak{S}_{2,1}$. Actually, we prove that $W_{\geq m, g} |D^0|^{-1} \in Y$, which gives $|D^0|^{-1} W_{\geq m, g} |D^0|^{-1} \in \mathfrak{S}_{\infty, \infty}$ since $\| |D^0|^{-1} \|_{\mathfrak{S}_{\infty, \infty}} \leq 1$. We conclude since $\mathfrak{S}_{\infty, \infty} \subset \mathfrak{S}_{\infty, 1}$. We use Proposition B.1 on $W_{\geq m, \ell}^\infty$ and focus on the singularity introduced by the potential G_ℓ , the difference being easy to deal with. For every $\xi \in Q_\ell^*$ and every ψ_ξ and ϕ_ξ in $L_\xi^2(Q_\ell)$ with $\|\psi_\xi\|_{L_\xi^2} = 1$, we have

$$\begin{aligned} & \left| \iint_{Q_\ell \times Q_\ell} \int_{Q_\ell^*} \psi_\xi^*(x) G_\ell(x - y) g_{\xi'}(x, y) |D_\xi|^{-1} \phi_\xi(y) dx dy d\xi' \right| \\ & \leq \int_{Q_\ell^*} d\xi' \left(\iint_{Q_\ell \times Q_\ell} |g_{\xi'}(x, y)|^2 dx dy \right)^{1/2} \left(\iint_{Q_\ell \times Q_\ell} |\psi_\xi(x)|^2 |G_\ell(x - y)|^2 \| |D_\xi|^{-1} \phi_\xi(y) \|^2 dx dy \right)^{1/2} \\ & \leq C_G \|g\|_{\mathfrak{S}_{2,1}} \|\phi_\xi\|_{L_\xi^2} \|\psi_\xi\|_{L_\xi^2}, \end{aligned}$$

thanks to (4.2). Hence the result. \square

We are now proving stronger convergence results in (6.8), by improving the bounds on the kernels of $(W_{\tilde{\gamma}_n, \xi})_n$.

Lemma 6.5 (Convergence of the sequence $(W_{\geq m, \tilde{\gamma}_n})_{n \geq 1}$). *The sequence of kernels $(W_{\geq m, \tilde{\gamma}_n, \xi})_n$ is bounded in $W^{1, \infty}(Q_\ell^*; L^2(Q_\ell \times Q_\ell))$. Therefore, up to the extraction of a subsequence, we have*

$$\| |D^0|^{-1/2} W_{\geq m, \tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_{\mathfrak{S}_{2, \infty}} \rightarrow 0.$$

Proof. Let us start with the boundedness of the sequence. We already proved in Lemma 6.4 that $(W_{\geq m, \tilde{\gamma}_n})_n$ is bounded in $\mathfrak{S}_{2, \infty}$, which follows from (B.17). Let us now check that the sequence of norms $\| \nabla_\xi W_{\geq m, \tilde{\gamma}_n, \xi}(\cdot, \cdot) \|_{L^\infty(Q_\ell^*; L^2(Q_\ell \times Q_\ell))}$ is also bounded. Thanks to (B.5), for every $\xi \in Q_\ell^*$,

$$\| \nabla_\xi W_{\geq m, \tilde{\gamma}_n, \xi}(\cdot, \cdot) \|_{L^2(Q_\ell \times Q_\ell)} \leq C \int_{Q_\ell^*} \|\tilde{\gamma}_{n, \xi'}\|_{\mathfrak{S}_2(\xi')} d\xi' \leq C \|\tilde{\gamma}_n\|_{\mathfrak{S}_{2, \infty}},$$

and we conclude with the help of (6.2). Therefore, the kernels $|D_\xi|^{-1/2} W_{\geq m, \tilde{\gamma}_n, \xi} |D_\xi|^{-1/2}(\cdot, \cdot)$ are bounded in $W^{1, \infty}(Q_\ell^*; H^{1/2}(Q_\ell \times Q_\ell))$. Thus, according to the Rellich–Kondrachov and the Arzelà–Ascoli theorems, up to the extraction of a subsequence,

$$|D_\xi|^{-1/2} W_{\geq m, \tilde{\gamma}_n, \xi} |D_\xi|^{-1/2}(\cdot, \cdot) \rightarrow |D_\xi|^{-1/2} W_{\geq m, \tilde{\gamma}_*, \xi} |D_\xi|^{-1/2}(\cdot, \cdot) \quad \text{in} \quad L^\infty(Q_\ell^*; L^2(Q_\ell \times Q_\ell))$$

which yields $\| |D^0|^{-1/2} W_{\geq m, \tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_{\mathfrak{S}_{2, \infty}} \rightarrow 0$, whence the lemma. \square

Lemma 6.6 (Convergence of the sequence $(W_{< m, \tilde{\gamma}_n})_{n \geq 1}$). *The sequence of kernels $(W_{< m, \tilde{\gamma}_n, \xi})_n$ is bounded in $C^{0, \mu}(Q_\ell^*; H_\xi^1(Q_\ell \times Q_\ell))$ for any $\mu \in (0, 1)$. In particular, up to the extraction of a subsequence,*

$$\| W_{< m, \tilde{\gamma}_n - \gamma_*} \|_{\mathfrak{S}_{2, \infty}} \rightarrow 0.$$

Proof. Let $\mu \in (0, 1)$. We first show the uniform boundedness of $W_{< m, \tilde{\gamma}_n, \xi}$ in $C^{0, \mu}(Q_\ell^*; H^1(Q_\ell \times Q_\ell))$. It is based on Lemma 6.1, particularly Eq.(6.3). In the formulas below, C is a positive constant which depends only on m, ℓ and μ . Recall that

$$W_{< m, \ell}^\infty(\eta, z) = \frac{4\pi}{\ell^3} \sum_{\substack{|k|_\infty < m \\ k \in \mathbb{Z}^3}} \frac{1}{|\frac{2\pi}{\ell}k - \eta|^2} e^{i(\frac{2\pi}{\ell}k - \eta) \cdot z}.$$

Thus,

$$\begin{aligned} & \| W_{< m, \tilde{\gamma}_n, \eta} - W_{< m, \tilde{\gamma}_n, \eta'} \|_{H^1(Q_\ell \times Q_\ell)} \\ & \leq \frac{4\pi}{\ell^3} \sum_{\substack{|k|_\infty \leq m-1 \\ k \in \mathbb{Z}^3}} \left\| \int_{Q_\ell^*} \frac{e^{i(\frac{2\pi}{\ell}k - (\eta - \xi')) \cdot (x-y)}}{|\frac{2\pi}{\ell}k - (\eta - \xi')|^2} \tilde{\gamma}_{n, \xi'} d\xi' - \int_{Q_\ell^*} \frac{e^{i(\frac{2\pi}{\ell}k - (\eta' - \xi')) \cdot (x-y)}}{|\frac{2\pi}{\ell}k - (\eta' - \xi')|^2} \tilde{\gamma}_{n, \xi'} d\xi' \right\|_{H^1(Q_\ell \times Q_\ell)}. \end{aligned}$$

For each term on the right-hand side, we have

$$\begin{aligned} & \left\| \int_{Q_\ell^*} \frac{e^{i(\frac{2\pi}{\ell}k - (\eta - \xi')) \cdot (x-y)}}{|\frac{2\pi}{\ell}k - (\eta - \xi')|^2} \tilde{\gamma}_{n, \xi'} d\xi' - \int_{Q_\ell^*} \frac{e^{i(\frac{2\pi}{\ell}k - (\eta' - \xi')) \cdot (x-y)}}{|\frac{2\pi}{\ell}k - (\eta' - \xi')|^2} \tilde{\gamma}_{n, \xi'} d\xi' \right\|_{H^1(Q_\ell \times Q_\ell)} \\ & \leq \int_{Q_\ell^*} \left| \frac{1}{|\frac{2\pi}{\ell}k - (\eta' - \xi')|^2} - \frac{1}{|\frac{2\pi}{\ell}k - (\eta - \xi')|^2} \right| \left\| e^{i(\frac{2\pi}{\ell}k - (\eta - \xi')) \cdot (x-y)} \tilde{\gamma}_{n, \xi'} \right\|_{H_\eta^1(Q_\ell \times Q_\ell)} d\xi' \\ & \quad + \left\| \int_{Q_\ell^*} \frac{(e^{i(\frac{2\pi}{\ell}k - (\eta - \xi')) \cdot (x-y)} - e^{i(\frac{2\pi}{\ell}k - (\eta' - \xi')) \cdot (x-y)})}{|\frac{2\pi}{\ell}k - (\eta' - \xi')|^2} \tilde{\gamma}_{n, \xi'} d\xi' \right\|_{H^1(Q_\ell \times Q_\ell)}. \end{aligned}$$

As $\eta, \xi' \in Q_\ell^*$, according to (6.3) we get

$$\left\| e^{i(\frac{2\pi}{\ell}k - (\eta - \xi')) \cdot (x-y)} \tilde{\gamma}_{n, \xi'} \right\|_{H_\eta^1(Q_\ell \times Q_\ell)} \leq C \|\tilde{\gamma}_{n, \xi'}\|_{H_{\xi'}^1(Q_\ell \times Q_\ell)} \leq 2C j_1 (1 - \kappa)^{-4} c^*(j_1)^2.$$

By the Hölder continuity of the function $\eta \mapsto \int_{Q_\ell^*} \frac{d\xi'}{|\eta - \xi'|^2}$,

$$\begin{aligned} & \int_{Q_\ell^*} \left| \frac{1}{\left| \frac{2\pi}{\ell}k - (\eta' - \xi') \right|^2} - \frac{1}{\left| \frac{2\pi}{\ell}k - (\eta - \xi') \right|^2} \right| \left\| e^{i\left(\frac{2\pi}{\ell}k - (\eta - \xi')\right) \cdot (x-y)} \tilde{\gamma}_{n, \xi'} \right\|_{H_\eta^1(Q_\ell \times Q_\ell)} d\xi' \\ & \leq 2Cj_1(1 - \kappa)^{-4} c^*(j_1)^2 |\eta - \eta'|^\mu. \end{aligned}$$

For the last term, note that

$$|e^{-i\eta \cdot z} - e^{-i\eta' \cdot z}| \leq \|\nabla_\eta e^{i\eta \cdot z}\|_{L^\infty(Q_\ell^* \times 2Q_\ell)} |\eta - \eta'| \leq C|\eta - \eta'|$$

and $|\nabla_z(e^{-i\eta \cdot z} - e^{-i\eta' \cdot z})| \leq C|\eta - \eta'|$. We get

$$\begin{aligned} & \left\| \int_{Q_\ell^*} \frac{e^{i\left(\frac{2\pi}{\ell}k + \xi'\right) \cdot (x-y)} (e^{-i\eta' \cdot (x-y)} - e^{-i\eta \cdot (x-y)})}{\left| \frac{2\pi}{\ell}k - (\eta' - \xi') \right|^2} \tilde{\gamma}_{n, \xi'} d\xi' \right\|_{H^1(Q_\ell \times Q_\ell)} \\ & \leq 2Cj_1(1 - \kappa)^{-4} c^*(j_1)^2 |\eta - \eta'|. \end{aligned}$$

We finally get

$$\|W_{< m, \tilde{\gamma}_n, \xi}\|_{C^{0, \mu}(Q_\ell^*; H_\xi^1(Q_\ell \times Q_\ell))} \leq 2Cj_1(1 - \kappa)^{-4} c^*(j_1)^2.$$

Lastly, thanks to the Arzelà–Ascoli theorem and the Rellich–Kondrachov theorem, the sequence of kernels converges strongly in $L^\infty(Q_\ell^*; L^2(Q_\ell \times Q_\ell))$, up to the extraction of a subsequence, and the limit is the operator $W_{< m, \gamma_*}$ thanks to (6.8). This concludes the proof of the lemma. \square

Then we have the following.

Lemma 6.7 (Strong convergence of the electron–electron interaction). *As n goes to infinity, we have*

$$\||D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2}\|_Y \rightarrow 0.$$

Proof. As $V_\gamma = G_\ell * \rho_\gamma - W_\gamma$, we have

$$\begin{aligned} & \||D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2}\|_Y \\ & \leq \||D^0|^{-1/2} G_\ell * (\rho_{\tilde{\gamma}_n} - \rho_{\gamma_*}) |D^0|^{-1/2}\|_Y + \||D^0|^{-1/2} (W_{\tilde{\gamma}_n} - W_{\gamma_*}) |D^0|^{-1/2}\|_Y. \end{aligned}$$

Notice that, from Lemma 6.3, we infer $\rho_{\tilde{\gamma}_n} \rightarrow \rho_{\gamma_*}$ in $L^2(Q_\ell)$. This, together with the fact that $G_\ell \in L^2(Q_\ell)$, yield

$$\|G_\ell * (\rho_{\tilde{\gamma}_n} - \rho_{\gamma_*})\|_{L^\infty(Q_\ell)} \rightarrow 0.$$

Then, using $|D^0|^{-1} \leq 1$, we infer

$$\||D^0|^{-1/2} G_\ell * (\rho_{\tilde{\gamma}_n} - \rho_{\gamma_*}) |D^0|^{-1/2}\|_Y \rightarrow 0. \quad (6.9)$$

We consider now the second term that we split into two parts. We already proved in Lemma 6.5 that $\||D^0|^{-1/2} W_{\geq m, \tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2}\|_{\mathfrak{S}_{2, \infty}} \rightarrow 0$, which implies the strong convergence in Y . The proof for the other term is even simpler, since

$$\||D^0|^{-1/2} W_{< m, \tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2}\|_{\mathfrak{S}_{2, \infty}} \leq \|W_{< m, \tilde{\gamma}_n - \gamma_*}\|_{\mathfrak{S}_{2, \infty}},$$

for $\||D^0|^{-1/2}\|_Y \leq 1$. We conclude by Lemma 6.6. Finally, we infer

$$\||D^0|^{-1/2} W_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2}\|_Y \rightarrow 0. \quad (6.10)$$

The lemma follows gathering together (6.9) and (6.10). \square

As a result, we have the following.

Corollary 6.8 (Strong convergence of the spectral projectors). *As n goes to infinity, we have*

$$\|P_{\gamma_*}^+ - P_{\tilde{\gamma}_n}^+\|_Y \rightarrow 0.$$

Proof. For any ϕ_ξ and $\psi_\xi \in L_\xi^2$, by (5.8) and the second resolvent identity, we obtain

$$\begin{aligned}
& \left| \left(\psi_\xi, (P_{\gamma_*, \xi}^+ - P_{\tilde{\gamma}_n, \xi}^+) \phi_\xi \right) \right| \\
& \leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} \left| \left(\psi_\xi, (D_{\gamma_*, \xi} - iz)^{-1} V_{\tilde{\gamma}_n - \gamma_*, \xi} (D_{\tilde{\gamma}_n, \xi} - iz)^{-1} \phi_\xi \right)_{L_\xi^2} \right| dz \\
& \leq \frac{1}{2\pi} \| |D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_Y \\
& \quad \times \left(\int_{-\infty}^{+\infty} \| (D_{\gamma_*, \xi} - iz)^{-1} |D_\xi|^{1/2} \psi_\xi \|_{L_\xi^2}^2 dz \right)^{1/2} \left(\int_{-\infty}^{+\infty} \| (D_{\gamma_*, \xi} - iz)^{-1} |D_\xi|^{1/2} \phi_\xi \|_{L_\xi^2}^2 dz \right)^{1/2} \\
& \leq \frac{1}{2} (1 - \kappa)^{-1} \| |D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_Y \|\phi_\xi\|_{L_\xi^2} \|\psi_\xi\|_{L_\xi^2}, \tag{6.11}
\end{aligned}$$

in virtue of (4.16). Therefore,

$$\|P_{\gamma_*}^+ - P_{\tilde{\gamma}_n}^+\|_Y \leq \frac{1}{2} (1 - \kappa)^{-1} \| |D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_Y.$$

The right-hand side goes to 0 by Lemma 6.7. \square

Relationship between $P_{\tilde{\gamma}_n}^+$ and $P_{\gamma_n}^+$

It remains to analyze the behaviour of $P_{\gamma_n}^+ - P_{\tilde{\gamma}_n}^+$, as n goes to infinity. Our main result is the following.

Lemma 6.9 (Relationship between $P_{\tilde{\gamma}_n}^+$ and $P_{\gamma_n}^+$). *For every $\gamma \in Z$ and for some $C > 0$ independent of n and Z , it holds that*

$$\|(P_{\gamma_n}^+ - P_{\tilde{\gamma}_n}^+) \gamma\|_{\mathfrak{S}_{1,1}} \leq C \|\gamma_n - \tilde{\gamma}_n\|_X^{1/2} \|\gamma\|_Z.$$

Proof. We first observe that $\|G * (\rho_{\gamma_n} - \rho_{\tilde{\gamma}_n})\|_Y \rightarrow 0$, thanks to Lemma 6.2 and (4.4) in Lemma 4.4. Next, thanks to the bound (B.10), the sequence of operators $(W_{\geq m, \gamma_n - \tilde{\gamma}_n})_n$ converges strongly to 0 in Y . Therefore, proceeding as for (5.12), we have

$$\frac{1}{2\pi} \left\| \int_{-\infty}^{+\infty} (D_{\gamma_n} - iz)^{-1} (G * (\rho_{\gamma_n} - \rho_{\tilde{\gamma}_n}) - W_{\geq m, \tilde{\gamma}_n - \gamma_n}) (D_{\tilde{\gamma}_n} - iz)^{-1} dz \right\|_Y \leq C \|\gamma_n - \tilde{\gamma}_n\|_X,$$

with the right-hand side going to 0. We now focus on the term involving $W_{< m, \gamma_n - \tilde{\gamma}_n}$. For every $\gamma \in Z$, proceeding as for (6.11), we have

$$\begin{aligned}
& \frac{1}{2\pi} \left\| \int_{-\infty}^{+\infty} (D_{\gamma_n} - iz)^{-1} W_{< m, \tilde{\gamma}_n - \gamma_n} (D_{\tilde{\gamma}_n} - iz)^{-1} \gamma dz \right\|_{\mathfrak{S}_{1,1}} \\
& \leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} \| (D_{\gamma_n} - iz)^{-1} W_{< m, \tilde{\gamma}_n - \gamma_n} (D_{\tilde{\gamma}_n} - iz)^{-1} \gamma \|_{\mathfrak{S}_{1,1}} dz \\
& \leq C \|W_{< m, \tilde{\gamma}_n - \gamma_n}\|_{\mathfrak{S}_{2,2}} \|\gamma\|_{\mathfrak{S}_{2,2}} \int_{-\infty}^{+\infty} \| (D_{\gamma_n} - iz)^{-1} \|_Y \| (D_{\tilde{\gamma}_n} - iz)^{-1} \|_Y dz \\
& \leq C \|\gamma_n - \tilde{\gamma}_n\|_{\mathfrak{S}_{2,2}} \|\gamma\|_{\mathfrak{S}_{2,2}} \\
& \leq C \|\gamma_n - \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}}^{1/2} \|\gamma\|_Z
\end{aligned}$$

thanks to (B.23) and in virtue of $\|\gamma_n - \tilde{\gamma}_n\|_Y \leq 2$. \square

6.3 Existence and properties of minimizers for $J_{\leq q}$

The existence of minimizers for $J_{\leq q}$ now follows by passing to the limit in the constraint and in the energy. The proof is separated into the following two lemmas.

Lemma 6.10. *The limit γ_* lies in Γ_q^+ .*

Proof. As

$$\tilde{\gamma}_n \xrightarrow{*} \gamma_* \quad \text{in } X_\infty^2 \cap Y,$$

we get

$$\|\gamma_*\|_Y \leq \liminf_{n \rightarrow \infty} \|\tilde{\gamma}_n\|_Y \leq 1$$

and

$$\|\gamma_*\|_{X_\infty^2} \leq \liminf_{n \rightarrow \infty} \|\tilde{\gamma}_n\|_{X_\infty^2} \leq j_1(1 - \kappa)^{-4} c^*(j_1)^2.$$

Thus, $\gamma_* \in \Gamma$. Since $\epsilon_P > e_0$, $\widetilde{\text{Tr}}_{L^2}[\gamma_*] = q$ thanks to Lemma 6.3.

To complete the proof, it remains to show that $P_{\gamma_*}^+ \gamma_* = \gamma_*$. From Lemma 6.9 and since $\|\tilde{\gamma}_n - \gamma_n\|_X \rightarrow 0$ (Lemma 6.2), we first prove that

$$\|P_{\tilde{\gamma}_n}^+ \tilde{\gamma}_n - \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}} \rightarrow 0. \quad (6.12)$$

Indeed, since $P_{\gamma_n}^+ \gamma_n = \gamma_n$, we have

$$\|P_{\tilde{\gamma}_n}^+ \tilde{\gamma}_n - \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}} \leq \|(P_{\tilde{\gamma}_n}^+ - P_{\gamma_n}^+) \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}} + \|P_{\gamma_n}^+ (\tilde{\gamma}_n - \gamma_n)\|_{\mathfrak{S}_{1,1}} + \|\gamma_n - \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}},$$

using also $\|P_{\gamma_n}^+\|_Y \leq 1$. Then, the right-hand side goes to 0. Next, by Corollary 6.8,

$$\|(P_{\tilde{\gamma}_n}^+ - P_{\gamma_*}^+) \tilde{\gamma}_n\|_{\mathfrak{S}_{1,1}} \rightarrow 0. \quad (6.13)$$

Let $g \in Y$. Let us show that

$$\widetilde{\text{Tr}}_{L^2}[(P_{\gamma_*}^+ \gamma_* - \gamma_*) g] = 0.$$

Notice that

$$\begin{aligned} \left| \widetilde{\text{Tr}}_{L^2}[(P_{\gamma_*}^+ \gamma_* - \gamma_*) g] \right| &\leq \left| \widetilde{\text{Tr}}_{L^2}[(P_{\tilde{\gamma}_n}^+ - P_{\gamma_*}^+) \tilde{\gamma}_n g] \right| + \left| \widetilde{\text{Tr}}_{L^2}[(P_{\tilde{\gamma}_n}^+ \tilde{\gamma}_n - \tilde{\gamma}_n) g] \right| \\ &\quad + \left| \widetilde{\text{Tr}}_{L^2}[P_{\gamma_*}^+ (\gamma_* - \tilde{\gamma}_n) g] \right| + \left| \widetilde{\text{Tr}}_{L^2}[(\tilde{\gamma}_n - \gamma_*) g] \right|. \end{aligned}$$

The first two terms in the right-hand side goes to 0 thanks to (6.12) and (6.13). For the last two terms, we use the weak-* convergence of $\tilde{\gamma}_n$ to γ_* in X_∞^2 and the fact that $|D^0|^{-1}g|D^0|^{-1}$ and $|D^0|^{-1}P_{\gamma_*}^+g|D^0|^{-1}$ both lie in $\mathfrak{S}_{\infty,1}$. Hence $\gamma_* \in \Gamma_q^+$. This concludes the proof. \square

Lemma 6.11. *We have*

$$\lim_{n \rightarrow +\infty} \left(\mathcal{E}^{DF}(\gamma_n) - \epsilon_P \widetilde{\text{Tr}}_{L^2}[\gamma_n] \right) = \mathcal{E}^{DF}(\gamma_*) - \epsilon_P \widetilde{\text{Tr}}_{L^2}[\gamma_*]. \quad (6.14)$$

Therefore, γ_* is a minimizer of $J_{\leq q}$.

Proof. For the kinetic energy term, we write

$$\widetilde{\text{Tr}}_{L^2}[D^0(\gamma_n - \gamma_*)] = \widetilde{\text{Tr}}_{L^2}[D^0(\gamma_n - \tilde{\gamma}_n)] + \widetilde{\text{Tr}}_{L^2}[D^0(\tilde{\gamma}_n - \gamma_*)].$$

Since $\|\gamma_n - \tilde{\gamma}_n\|_X \rightarrow 0$ thanks to Lemma 6.2, $\widetilde{\text{Tr}}_{L^2}[D^0(\gamma_n - \tilde{\gamma}_n)] \rightarrow 0$. On the other hand, by definition of the weak-* convergence in X_∞^2 , $|D^0|(\tilde{\gamma}_n - \gamma_*)|D^0|$ converges to 0 for the weak-* topology in $\mathfrak{S}_{1,\infty}$. As $|D^0|^{-1}D^0|D^0|^{-1} \in \mathfrak{S}_{\infty,1}$, we obtain that

$$\widetilde{\text{Tr}}_{L^2}[D^0(\tilde{\gamma}_n - \gamma_*)] = \widetilde{\text{Tr}}_{L^2}[|D^0|(\tilde{\gamma}_n - \gamma_*)|D^0||D^0|^{-1}D^0|D^0|^{-1}] \rightarrow 0,$$

hence

$$\widetilde{\text{Tr}}_{L^2}[D^0(\gamma_n - \gamma_*)] \rightarrow 0.$$

In addition, thanks again to Lemma 6.3,

$$\epsilon_P \widetilde{\text{Tr}}_{L^2}[\tilde{\gamma}_n - \gamma_*] \rightarrow 0.$$

The proof for the attractive potential is similar. We start with

$$\widetilde{\text{Tr}}_{L^2}[G_\ell(\gamma_n - \gamma_*)] = \widetilde{\text{Tr}}_{L^2}[G_\ell(\gamma_n - \tilde{\gamma}_n)] + \widetilde{\text{Tr}}_{L^2}[G_\ell(\tilde{\gamma}_n - \gamma_*)].$$

The second term goes to 0 as n goes to infinity since $\rho_{\tilde{\gamma}_n}$ converges to ρ_{γ_*} in $L^2(Q_\ell)$ and $G \in L^2(Q_\ell)$ (Lemma 6.3). For the first term, we use the fact that

$$\left| \widetilde{\text{Tr}}_{L^2}[G_\ell(\gamma_n - \tilde{\gamma}_n)] \right| \leq C_H \|\gamma_n - \tilde{\gamma}_n\|_X \rightarrow 0,$$

and we conclude by Lemma 6.2.

For the repulsive potential, using the fact that $\widetilde{\text{Tr}}_{L^2}[V_{\tilde{\gamma}} \gamma'] = \widetilde{\text{Tr}}_{L^2}[V_{\gamma'} \tilde{\gamma}]$ whenever γ and γ' are in Z , we have

$$\begin{aligned} & \left| \widetilde{\text{Tr}}_{L^2}[V_{\gamma_n} \gamma_n - V_{\gamma_*} \gamma_*] \right| \\ & \leq \left| \widetilde{\text{Tr}}_{L^2}[V_{\gamma_n} \gamma_n - V_{\tilde{\gamma}_n} \tilde{\gamma}_n] \right| + \left| \widetilde{\text{Tr}}_{L^2}[V_{\tilde{\gamma}_n} \tilde{\gamma}_n - V_{\gamma_*} \gamma_*] \right| \\ & = \left| \widetilde{\text{Tr}}_{L^2}[V_{\tilde{\gamma}_n + \gamma_n}(\gamma_n - \tilde{\gamma}_n)] \right| + \left| \widetilde{\text{Tr}}_{L^2}[V_{\tilde{\gamma}_n - \gamma_*}(\tilde{\gamma}_n + \gamma_*)] \right| \\ & \leq C_{EE} (\|\tilde{\gamma}_n\|_Z + \|\gamma_n\|_Z) \|\tilde{\gamma}_n - \gamma_n\|_X + \| |D^0|^{-1/2} V_{\tilde{\gamma}_n - \gamma_*} |D^0|^{-1/2} \|_Y (\|\tilde{\gamma}_n\|_X + \|\gamma_*\|_X) \end{aligned}$$

using the bound (4.11) in Lemma 4.7. Finally, the right-hand side in the above string of inequalities goes to 0, according to Lemmas 6.2 and 6.7. The lemma follows. \square

Since γ_* is a minimizer of $J_{\leq q}$, we may apply Proposition 5.1 to the constant minimizing sequence equal to γ_* , and we obtain that

$$\int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{\gamma_*, \xi} - \epsilon_P) \gamma_{*, \xi}] d\xi = \inf_{\substack{\gamma \in \Gamma_{\leq q} \\ \gamma = P_{\gamma_*}^+ \gamma}} \int_{Q_\ell^*} \text{Tr}_{L_\xi^2}[(D_{\gamma_*, \xi} - \epsilon_P) \gamma_\xi] d\xi.$$

We may then apply Lemma 4.15 to conclude that $\gamma_* = f_{Q_\ell^*}^\oplus 1_{[0, \nu)}(D_{\gamma_*, \xi}) d\xi + \delta$ with $0 \leq \delta \leq 1_{\{\nu\}}(D_{\gamma_*})$ for some $\nu \in [\lambda_0, e_0]$ that is independent of ϵ_P . This ends the proof of Theorem 3.3. Thus, Theorem 2.6 holds true.

Acknowledgments. The authors are grateful to the referees for a careful reading of the manuscript and for remarks and suggestions that greatly improved the quality of this paper. L.M. acknowledges support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement No. 810367).

Declarations.

Conflict of interest The authors declare no conflict of interest.

Data Availability Data sharing is not applicable to this article as it has no associated data.

A Proof of Lemma 4.1

It suffices to prove (4.2). By interpolation, we can choose $C_H = C_G$.

To deal with (4.2), the idea is to decompose the potential G_ℓ on Q_ℓ into two parts, namely $\frac{1}{|x|}$ and $G_\ell - \frac{1}{|x|}$. The first term can be treated as the Hardy inequality on Q_ℓ , whereas the second is bounded. We begin with the second term and prove the following.

Lemma A.1. *There is a constant $C_0 > 0$ independent of ℓ such that*

$$\sup_{x \in Q_\ell} \left| G_\ell(x) - \frac{1}{|x|} \right| \leq \frac{C_0}{\ell}.$$

This implies that, for any $x \in \mathbb{R}^3$,

$$G_\ell(x) \geq -\frac{C_0}{\ell}. \tag{A.1}$$

In particular, we have

$$C_0 \leq \inf_{0 < R < \frac{1}{2}} \left(\frac{3}{2R} + \frac{2\pi R^2}{5} + \frac{3}{4\pi^2 R^3} \min \left\{ \frac{4\pi R^3}{3}; 1 - \frac{4\pi R^3}{3} \right\}^{1/2} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2} \right).$$

Proof. As $G_1(x) = \ell G_\ell(\ell x)$, it suffices to consider the case $\ell = 1$. Let $f(x) = G_1(x) - \frac{1}{|x|}$. Equation (2.1) yields

$$-\Delta f = 4\pi \left(-1 + \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \delta_k \right).$$

Let $B(z, R)$ be a ball of center z and radius R chosen such that $(\bigcup_{z \in Q_1} B(z, R)) \cap (\mathbb{Z}^3 \setminus \{0\}) = \emptyset$. Obviously, we can assume $0 < R < 1/2$. On the one hand, by the divergence theorem, for $0 \leq r \leq R$ and $z \in Q_1$ we obtain

$$\frac{1}{4\pi} \frac{d}{dr} \int_{\mathbb{S}^2} f(z + r\omega) d\omega = \frac{1}{4\pi r^2} \int_{\partial B(z,r)} \frac{\partial f(s)}{\partial n} ds = \frac{1}{4\pi r^2} \int_{B(z,r)} \Delta_x f(x) dx \quad (\text{A.2})$$

with \mathbb{S}^2 denoting the unit sphere. For any $z \in Q_1$,

$$\frac{1}{4\pi r^2} \int_{B(z,r)} \Delta_x f dx = \frac{1}{r^2} \int_{B(z,r)} 1 dx = \frac{4\pi}{3} r,$$

where the first equation holds since

$$\left(\bigcup_{z \in Q_1} B(z, r) \right) \cap (\mathbb{Z}^3 \setminus \{0\}) = \emptyset, \quad \text{for } 0 \leq r \leq R.$$

Therefore, integrating (A.2) with respect to r ,

$$\int_{\mathbb{S}^2} f(z + r\omega) d\omega - 4\pi f(z) = \frac{8\pi^2}{3} r^2.$$

Since $\int_{B(z,R)} f(x) dx = \int_0^R r^2 \left(\int_{\mathbb{S}^2} f(z + r\omega) d\omega \right) dr$, integration over $[0, R]$ leads to

$$|f(z)| \leq \frac{3}{4\pi R^3} \left| \int_{B(z,R)} f(x) dx \right| + \frac{2\pi R^2}{5} \leq \frac{3}{4\pi R^3} \left| \int_{B(z,R)} G_1(x) dx \right| + \frac{3}{4\pi R^3} \left| \int_{B(z,R)} \frac{1}{|x|} dx \right| + \frac{2\pi R^2}{5}.$$

On the other hand,

$$\begin{aligned} \left| \int_{B(z,R)} G_1(x) dx \right| &\leq |B(z, R)|^{1/2} \|G_1\|_{L^2(B(z,R))} \\ &\leq \left(\frac{4\pi R^3}{3} \right)^{1/2} \|G_1\|_{L^2(Q_1)} = \frac{1}{\pi} \left(\frac{4\pi R^3}{3} \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2}. \end{aligned}$$

Using (2.2) and the periodicity of G_1 , we also have

$$\begin{aligned} \left| \int_{B(z,R)} G_1(x) dx \right| &= \left| \int_{(z+Q_1) \setminus B(z,R)} G_1(x) dx \right| \\ &\leq \frac{1}{\pi} \left(1 - \frac{4\pi R^3}{3} \right)^{1/2} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2}. \end{aligned}$$

Thus,

$$\left| \int_{B(z,R)} G_1(x) dx \right| \leq \frac{1}{\pi} \min \left\{ \left(\frac{4\pi R^3}{3} \right)^{1/2}, \left(1 - \frac{4\pi R^3}{3} \right)^{1/2} \right\} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2}.$$

Furthermore,

$$\left| \int_{B(z,R)} \frac{1}{|x|} dx \right| \leq \int_{B(0,R)} \frac{1}{|x|} dx = 2\pi R^2.$$

Therefore, the bound holds for $\ell = 1$ and any $0 < R < \frac{1}{2}$ with

$$C_0 \leq \frac{3}{2R} + \frac{2\pi R^2}{5} + \frac{3}{4\pi^2 R^3} \min \left\{ \frac{4\pi R^3}{3}; 1 - \frac{4\pi R^3}{3} \right\}^{1/2} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2}.$$

□

We now consider the Hardy inequality on Q_ℓ for the potential $\frac{1}{|x|}$.

Lemma A.2. *Let $u \in H^1(Q_\ell)$, then*

$$\left\| \frac{u}{|x|} \right\|_{L^2(Q_\ell)}^2 \leq \frac{4\ell + 12}{\ell} \|\nabla u\|_{L^2(Q_\ell)}^2 + \frac{24 + 12\ell}{\ell^2} \|u\|_{L^2(Q_\ell)}^2.$$

Proof. We start with the relationship:

$$0 \leq \int_{Q_\ell} \left| \nabla u + \frac{xu}{2|x|^2} \right|^2 dx.$$

Thus,

$$0 \leq \int_{Q_\ell} |\nabla u|^2 dx + \frac{1}{4} \int_{Q_\ell} \frac{|u|^2}{|x|^2} dx + \frac{1}{2} \int_{Q_\ell} \frac{\nabla |u|^2 \cdot x}{|x|^2} dx.$$

By the divergence theorem for $\int_{Q_\ell} \nabla \cdot \left(\frac{|u|^2 x}{|x|^2} \right) dx$, we obtain

$$\int_{Q_\ell} \frac{\nabla |u|^2 \cdot x}{|x|^2} dx = \int_{\partial Q_\ell} \frac{\vec{n}x |u|^2}{|x|^2} dx - \int_{Q_\ell} \frac{|u|^2}{|x|^2} dx$$

where \vec{n} is the outward pointing unit normal at each point on the boundary ∂Q_ℓ . To end this proof, it suffices to estimate $\int_{\partial Q_\ell} \frac{\vec{n}x |u|^2}{|x|^2}$.

Let

$$A^{2,3}(x_1) = \int_{(-\frac{\ell}{2}, \frac{\ell}{2}]^2} |u|^2(x_1, x_2, x_3) dx_2 dx_3.$$

As $|\vec{n} \cdot x| = \frac{\ell}{2}$ and $|x| \geq \frac{\ell}{2}$ for any $x \in \partial Q_\ell$, we have

$$\begin{aligned} \left| \int_{\partial Q_\ell} \frac{\vec{n}x |u|^2}{|x|^2} dx \right| &\leq \frac{2}{\ell} \int_{\partial Q_\ell} |u|^2 dx \\ &= \frac{2}{\ell} \left(A^{2,3} \left(-\frac{\ell}{2} \right) + A^{2,3} \left(\frac{\ell}{2} \right) + A^{1,3} \left(-\frac{\ell}{2} \right) + A^{1,3} \left(\frac{\ell}{2} \right) + A^{1,2} \left(-\frac{\ell}{2} \right) + A^{1,2} \left(\frac{\ell}{2} \right) \right). \end{aligned} \quad (\text{A.3})$$

Let $x_1^{(0)} \in (-\frac{\ell}{2}, \frac{\ell}{2}]$ such that

$$A^{2,3}(x_1^{(0)}) \leq \int_{(-\frac{\ell}{2}, \frac{\ell}{2}]^2} A^{2,3}(x_1) dx_1 = \frac{1}{\ell} \int_{(-\frac{\ell}{2}, \frac{\ell}{2}]^2} A^{2,3}(x_1) dx_1 = \frac{1}{\ell} \int_{Q_\ell} |u|^2 dx.$$

Then we have

$$\begin{aligned} A^{2,3} \left(-\frac{\ell}{2} \right) + A^{2,3} \left(\frac{\ell}{2} \right) &= \left[\int_{x_1^{(0)}}^{-\frac{\ell}{2}} - \int_{-\frac{\ell}{2}}^{x_1^{(0)}} \right] \frac{d}{dx_1} A^{2,3} dx_1 + 2A^{2,3}(x_1^{(0)}) \\ &\leq 2A^{2,3}(x_1^{(0)}) + \int_{(-\frac{\ell}{2}, \frac{\ell}{2}]^2} \left| \frac{d}{dx_1} A^{2,3} \right| dx_1. \end{aligned}$$

As

$$\int_{(-\frac{\ell}{2}, \frac{\ell}{2}]^2} \left| \frac{d}{dx_1} A^{2,3}(x_1) \right| dx_1 \leq 2 \int_{Q_\ell} |u| \left| \frac{\partial}{\partial x_1} u \right| \leq 2 \|u\|_{L^2(Q_\ell)} \|\nabla u\|_{L^2(Q_\ell)},$$

we get

$$A^{2,3} \left(-\frac{\ell}{2}\right) + A^{2,3} \left(\frac{\ell}{2}\right) \leq \frac{2}{\ell} \|u\|_{L^2(Q_\ell)}^2 + 2 \|u\|_{L^2(Q_\ell)} \|\nabla u\|_{L^2(Q_\ell)}.$$

Inserting this into (A.3), we can conclude that

$$0 \leq \|\nabla u\|_{L^2(Q_\ell)}^2 - \frac{1}{4} \left\| \frac{u}{|x|} \right\|_{L^2(Q_\ell)}^2 + \frac{6}{\ell^2} \|u\|_{L^2(Q_\ell)}^2 + \frac{6}{\ell} \|u\|_{L^2(Q_\ell)} \|\nabla u\|_{L^2(Q_\ell)}.$$

As a result, by the Cauchy–Schwarz inequality

$$\left\| \frac{u}{|x|} \right\|_{L^2(Q_\ell)}^2 \leq \frac{4\ell + 12}{\ell} \|\nabla u\|_{L^2(Q_\ell)}^2 + \frac{24 + 12\ell}{\ell^2} \|u\|_{L^2(Q_\ell)}^2.$$

□

Actually, Lemma A.2 implies that for any $\xi \in Q_\ell^*$,

$$\begin{aligned} \||x|^{-1} u_\xi\|_{L_\xi^2} &\leq \left(\max \left\{ \frac{24 + 12\ell}{\ell^2}; \frac{4\ell + 12}{\ell} \right\} \right) \|(1 - \Delta_\xi)^{1/2} u_\xi\|_{L_\xi^2} \\ &= 2 \max \left\{ \sqrt{\frac{6}{\ell^2} + \frac{3}{\ell}}; \sqrt{1 + \frac{3}{\ell}} \right\} \|u_\xi\|_{H_\xi^1}. \end{aligned}$$

Thus combining Lemmas A.1 and A.2, we know that

$$\begin{aligned} \|G_\ell u_\xi\|_{L_\xi^2} &\leq \frac{C_0}{\ell} \|u_\xi\|_{L_\xi^2} + 2 \max \left\{ \sqrt{\frac{6}{\ell^2} + \frac{3}{\ell}}; \sqrt{1 + \frac{3}{\ell}} \right\} \|u_\xi\|_{H_\xi^1} \\ &\leq C_G \|u_\xi\|_{H_\xi^1}. \end{aligned}$$

with

$$C_G := 2 \left(1 + \frac{C_0}{\ell} \right) \max \left\{ \sqrt{1 + \frac{3}{\ell}}; \sqrt{\frac{3}{\ell} + \frac{6}{\ell^2}} \right\}. \quad (\text{A.4})$$

We now turn to the estimates on the operator W_γ .

B Proof of Lemma 4.5

We first study the properties of W_ℓ^∞ , then we prove Lemma 4.5. In passing, we correct wrong estimates in [8]. Indeed, contrary to what was claimed, the function $W_\infty(\eta, x) - e^{-i\eta \cdot x} G(x) - 4\pi \frac{e^{-i\eta \cdot x}}{|\eta|^2}$ introduced in [8, p. 745, Eq. (5.3)] is not harmonic in $(1 + \epsilon)Q_\ell$ with respect to the x -variable. Therefore, some arguments need to be modified.

Properties of W_ℓ^∞

Recall that $W_\ell^\infty = W_{\geq m, \ell}^\infty + W_{< m, \ell}^\infty$ is given by (4.7). We are going to prove the Hardy type inequalities for $W_{\geq m, \ell}^\infty$. A natural idea is to compare it with the potential G_ℓ .

Proposition B.1 (Singularities for the potential $W_{\geq m, \ell}^\infty$). *For every $m \geq 2$, there exists a positive constant $C_{\geq m}$ such that, for any $\ell > 0$, we have*

$$\sup_{\substack{\eta \in 2Q_\ell^* \\ x \in Q_\ell}} \left| W_{\geq m, \ell}^\infty(\eta, x) - G_\ell(x) \right| \leq \frac{C_{\geq m}}{\ell} \quad (\text{B.1})$$

with

$$C_{\geq m} \leq \inf_{0 < R < 1/2} \left\{ \frac{\sqrt{3}}{2(\pi R)^{3/2}} \frac{m^2 + 2}{(m-1)^2} \left(\sum_{|k|_\infty \geq m} \frac{1}{|k|^4} \right)^{1/2} + \frac{2\pi[(2m-1)^3 + 1]R^2}{5} \right\}$$

$$+ \frac{3}{4\pi^2 R^3} \min \left\{ \frac{4\pi R^3}{3}; 1 - \frac{4\pi R^3}{3} \right\}^{1/2} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2} \Bigg\}.$$

Proof. The proof is similar to Lemma A.1. Notice that

$$W_\ell^\infty(\eta, x) = \lambda W_{\lambda\ell}^\infty \left(\frac{\eta}{\lambda}, \lambda x \right), \quad \eta \in \mathbb{R}^3, x \in \mathbb{R}^3.$$

We therefore take $\ell = 1$. Observe, from (2.20), that

$$-\Delta_z W_1^\infty(\eta, x) = 4\pi \sum_{k \in \mathbb{Z}^3} e^{-i\eta \cdot k} \delta_k(x).$$

Let $f(\eta, x) = W_{\geq m, 1}^\infty(\eta, x) - G_1(x)$. Then

$$-\Delta_x f(\eta, x) = 4\pi \sum_{\substack{k \neq 0 \\ k \in \mathbb{Z}^3}} (e^{-i\eta \cdot k} - 1) \delta_k(x) + 4\pi - 4\pi \sum_{\substack{|k|_\infty < m \\ k \in \mathbb{Z}^3}} e^{i(2\pi k - \eta) \cdot x}.$$

Let $B(z, R)$ be a ball of center z and radius R chosen such that $(\bigcup_{z \in Q_1} B(z, R)) \cap (\mathbb{Z}^3 \setminus \{0\}) = \emptyset$. Obviously, we can assume $0 < R < 1/2$. Analogously to (A.2), for $0 \leq r \leq R$ and $z \in Q_1$ we obtain

$$\frac{d}{dr} \left(\frac{1}{4\pi r^2} \int_{\partial B(z, r)} f(\eta, s) ds \right) = \frac{1}{4\pi r^2} \int_{B(z, r)} \Delta_x f(\eta, x) dx. \quad (\text{B.2})$$

On the one hand, for any $z \in Q_1$,

$$\frac{1}{4\pi r^2} \left| \int_{B(z, r)} \Delta_x f dx \right| = \frac{1}{r^2} \left| \int_{B(z, r)} \left(1 - \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} e^{i(2\pi k - \eta) \cdot x} \right) dx \right| \leq \frac{4\pi [(2m-1)^3 + 1]}{3} r,$$

where the first equality holds since

$$\left(\bigcup_{z \in Q_1} B(z, r) \right) \cap (\mathbb{Z}^3 \setminus \{0\}) = \emptyset, \quad \text{for } 0 \leq r \leq R.$$

Therefore, integrating (B.2) with respect to r ,

$$-\frac{8\pi^2 [(2m-1)^3 + 1]}{3} r^2 \leq \int_{\mathbb{S}^2} f(\eta, z + r\omega) d\omega - 4\pi f(\eta, z) \leq \frac{8\pi^2 [(2m-1)^3 + 1]}{3} r^2.$$

Then integration over $[0, R]$ leads to

$$|f(\eta, z)| \leq \frac{3}{4\pi R^3} \left| \int_{B(z, R)} f(\eta, x) dx \right| + \frac{2\pi [(2m-1)^3 + 1] R^2}{5}.$$

On the other hand,

$$\left| \int_{B(z, R)} G_1(x) dx \right| \leq \frac{1}{\pi} \min \left\{ \left(\frac{4\pi R^3}{3} \right)^{1/2}, \left(1 - \frac{4\pi R^3}{3} \right)^{1/2} \right\} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2}.$$

Furthermore, according to the quasi-periodicity of $W_{\geq m, 1}^\infty$ with respect to $z \in \mathbb{R}^3$, for any $\eta \in 2Q_1^*$,

$$\begin{aligned} \left| \int_{B(z, R)} W_{\geq m, 1}^\infty(\eta, x) dx \right| &\leq |B(z, R)|^{1/2} \|W_{\geq m, 1}^\infty\|_{L^2(B(z, R))} \leq \left(\frac{4\pi R^3}{3} \right)^{1/2} \|W_{\geq m, 1}^\infty\|_{L^2(Q_1)} \\ &\leq 4\pi \left(\frac{4\pi R^3}{3} \right)^{1/2} \left(\sum_{|k|_\infty \geq m} \frac{1}{|2\pi k - \eta|^4} \right)^{1/2} \end{aligned}$$

$$\begin{aligned}
&\leq 4\pi \left(\frac{4\pi R^3}{3} \right)^{1/2} \sup_{\substack{|k|_\infty \geq m \\ \eta \in 2Q_1^*}} \frac{|2\pi k|^2}{|2\pi k - \eta|^2} \left(\sum_{|k|_\infty \geq m} \frac{1}{|2\pi k|^4} \right)^{1/2} \\
&= \left(\frac{4\pi R^3}{3} \right)^{1/2} \frac{m^2 + 2}{\pi(m-1)^2} \left(\sum_{|k|_\infty \geq m} \frac{1}{|k|^4} \right)^{1/2}.
\end{aligned}$$

Therefore, the bound (B.1) holds for $\ell = 1$ with

$$\begin{aligned}
C_{\geq m} &\leq \frac{\sqrt{3}}{2(\pi R)^{3/2}} \frac{m^2 + 2}{(m-1)^2} \left(\sum_{|k|_\infty \geq m} \frac{1}{|k|^4} \right)^{1/2} + \frac{2\pi[(2m-1)^3 + 1]R^2}{5} \\
&\quad + \frac{3}{4\pi^2 R^3} \min \left\{ \left(\frac{4\pi R^3}{3} \right)^{1/2}, \left(1 - \frac{4\pi R^3}{3} \right)^{1/2} \right\} \left(\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \right)^{1/2},
\end{aligned}$$

for any $0 < R < \frac{1}{2}$. The corresponding result for any $\ell > 0$ follows immediately by a scaling argument. \square

We can immediately conclude from Lemma 4.1 and Proposition B.1 the following.

Corollary B.2 (Hardy-type inequalities for the potential $W_{\geq m, \ell}^\infty$). *For $m \geq 2$, we have*

$$\| |W_{\geq m, \ell}^\infty|^{1/2} |D_\xi|^{-1/2} \|_Y \leq \left(C_H + \frac{C_{\geq m}}{\ell} \right) \quad (\text{B.3})$$

and

$$\| W_{\geq m, \ell}^\infty |D_\xi|^{-1} \|_Y \leq \left(C_G + \frac{C_{\geq m}}{\ell} \right). \quad (\text{B.4})$$

We also have the following estimate on $W_{\geq m, \ell}^\infty$.

Lemma B.3. *Let $m \geq 2$. There is a positive constant $C = C(\ell, m)$ such that*

$$\sup_{\eta \in 2Q_\ell^*} \|\nabla_\eta W_{\geq m, \ell}^\infty(\eta, \cdot)\|_{L^\infty(Q_\ell)} \leq C. \quad (\text{B.5})$$

Proof. Take $\ell = 1$ for simplicity. Note that

$$-\Delta_x \nabla_\eta W_{\geq m, 1}^\infty(\eta, x) = -4\pi \sum_{k \in \mathbb{Z}^3 \setminus \{0\}} i k e^{-i\eta \cdot k} \delta_k(x) + 4\pi \sum_{\substack{|k|_\infty < m \\ k \in \mathbb{Z}^3}} i x e^{i(2\pi k - \eta) \cdot x},$$

from which we obtain

$$|\Delta_x \nabla_\eta W_{\geq m, 1}^\infty(\eta, x)| \leq C$$

for any $\eta \in 2Q_\ell^*$ and $x \in Q_\ell$. Following the proof of Lemma B.1, we know

$$|\nabla_\eta W_{\geq m, 1}^\infty(\eta, x)| \leq C.$$

The corresponding result for any $\ell > 0$ follows immediately by a scaling argument as for Lemma B.1. \square

Estimates for the exchange term

We consider now the exchange term. Let $\psi_\xi \in H_\xi^{1/2}$. As

$$\|W_{\gamma, \xi} \psi_\xi\|_{L_\xi^2} = \sup_{\phi_\xi \in L_\xi^2, \|\phi_\xi\|_{L_\xi^2} = 1} |(\phi_\xi, W_{\gamma, \xi} \psi_\xi)|, \quad (\text{B.6})$$

we only need to study the inner product $(W_{\gamma, \xi} \psi_\xi, \phi_\xi)$. For $m \geq 2$, $\eta \in \mathbb{R}^3$ and $z \in \mathbb{R}^3$,

$$W_\ell^\infty(\eta, z) = W_{\geq m, \ell}^\infty(\eta, z) + W_{< m, \ell}^\infty(\eta, z). \quad (\text{B.7})$$

For the term that carries all singularities in the x variable (i.e., $W_{\geq m, \ell}^\infty$), we use the decomposition (2.14) and Corollary B.2. Let $\gamma \in X$ with $\gamma^* = \gamma$. For any $\xi \in Q_\ell^*$ we have

$$|D_\xi|^{1/2} \gamma_\xi |D_\xi|^{1/2} = \sum_{n \geq 1} \lambda_n(\xi) |v_n(\xi, \cdot)\rangle \langle v_n(\xi, \cdot)| \quad (\text{B.8})$$

with $(v_n(\xi, \cdot), v_m(\xi, \cdot))_{L_\xi^2} = \delta_{m, n}$ and $\|\gamma\|_X = \int_{Q_\ell^*} \sum_{n \geq 1} |\lambda_n(\xi)| d\xi$. Hence

$$\gamma_\xi = \sum_{n \geq 1} \lambda_n(\xi) |u_n(\xi, \cdot)\rangle \langle u_n(\xi, \cdot)|$$

with $u_n(\xi, \cdot) = |D_\xi|^{-1/2} v_n(\xi, \cdot)$. Now, we have

$$\begin{aligned} & \left| \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} W_{\geq m, \ell}^\infty(\xi - \xi', x - y) \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy \right| \\ & \leq \int_{Q_\ell^*} d\xi' \sum_{n \geq 1} |\lambda_n(\xi')| \int_{Q_\ell \times Q_\ell} |W_{\geq m, \ell}^\infty(\xi - \xi', x - y)| |u_n(\xi', x)| |u_n(\xi', y)| |\psi_\xi(y)| |\phi_\xi(x)| dx dy \\ & \leq \int_{Q_\ell^*} d\xi' \sum_{n \geq 1} |\lambda_n(\xi')| \left(\int_{Q_\ell} |\psi_\xi(y)|^2 dy \int_{Q_\ell} |W_{\geq m, \ell}^\infty(\xi - \xi', x - y)| |u_n(\xi', x)|^2 dx \right)^{1/2} \\ & \quad \times \left(\int_{Q_\ell} |\phi_\xi(x)|^2 dx \int_{Q_\ell} |W_{\geq m, \ell}^\infty(\xi - \xi', x - y)| |u_n(\xi', y)|^2 dy \right)^{1/2} \\ & \leq \left(\int_{Q_\ell^*} \left(C_H + \frac{C_{\geq m}}{\ell} \right) \sum_{n \geq 1} |\lambda_n(\xi')| \| |D_{\xi'}|^{1/2} u_n(\xi', \cdot) \|_{L^2(\xi')}^2 d\xi' \right) \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2} \\ & \leq \left(C_H + \frac{C_{\geq m}}{\ell} \right) \|\gamma\|_X \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2}, \end{aligned} \quad (\text{B.9})$$

with the help of (B.3). Thus, for every $\gamma \in X$ with $\gamma = \gamma^*$, we have

$$\|W_{\geq m, \gamma}\|_Y \leq \left(C_H + \frac{C_{\geq m}}{\ell} \right) \|\gamma\|_X. \quad (\text{B.10})$$

Using the Cauchy–Schwarz inequality and (B.4), we can also argue as follows whenever $\gamma \in \mathfrak{S}_{1,1}$ with $\gamma^* = \gamma$:

$$\begin{aligned} & \left| \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} W_{\geq m, \ell}^\infty(\xi - \xi', x - y) \phi_\xi^*(x) \gamma_{\xi'}(x, y) (|D_\xi|^{-1} \psi_\xi)(y) dx dy \right| \\ & \leq \int_{Q_\ell^*} \left(\iint_{Q_\ell \times Q_\ell} \rho_{|\gamma_{\xi'}|}(y) |\phi_\xi(x)|^2 dx dy \right)^{1/2} \left(\iint_{Q_\ell \times Q_\ell} \rho_{|\gamma_{\xi'}|}(x) |W_{\geq m, \ell}^\infty(\xi - \xi', x - y)|^2 \||D_\xi|^{-1} \psi_\xi(y)|^2 dx dy \right)^{1/2} d\xi' \\ & \leq \left(C_G + \frac{C_{\geq m}}{\ell} \right) \|\gamma\|_{\mathfrak{S}_{1,1}} \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2}. \end{aligned} \quad (\text{B.11})$$

Thus, for every $\gamma \in \mathfrak{S}_{1,1}$ with $\gamma^* = \gamma$, since $|D^0|^{-1} \in \mathfrak{S}_{\infty, \infty}$, we have

$$\|W_{\geq m, \gamma} |D^0|^{-1}\|_Y \leq \left(C_G + \frac{C_{\geq m}}{\ell} \right) \|\gamma\|_{\mathfrak{S}_{1,1}}. \quad (\text{B.12})$$

Additionally, when $\gamma \in X_1^2$ with $\gamma = \gamma^*$ (which includes the case $\gamma \in X_\infty^2$), one can prove that $W_{\geq m, \gamma}$ belongs to $\mathfrak{S}_{2, \infty}$. Recall that $\|W_{\geq m, \gamma}\|_{\mathfrak{S}_{2, p}} = \|W_{\geq m, \gamma}(\cdot, \cdot)\|_{L^p(Q_\ell^*; L^2(Q_\ell \times Q_\ell))}$. The main ingredient is that $\sqrt{\rho_{|\gamma|}} \in H^1(Q_\ell)$ when $\gamma \in X_1^2$. Indeed, analogously to (B.8), for any $\gamma \in X_1^2$, we have

$$|D_\xi| \gamma_\xi |D_\xi| = \sum_{n \geq 1} \mu_n(\xi) |v_n(\xi, \cdot)\rangle \langle v_n(\xi, \cdot)| \in \mathfrak{S}_{1, p} \quad (\text{B.13})$$

with $(v_n(\xi, \cdot), v_m(\xi, \cdot))_{L^2_\xi} = \delta_{m,n}$ and $\|\gamma\|_{X_p^2} = \left(\int_{Q_\ell^*} (\sum_{n \geq 1} |\mu_n(\xi)|)^p d\xi \right)^{1/p}$. Hence

$$\gamma_\xi = \sum_{n \geq 1} \mu_n(\xi) |u_n(\xi, \cdot)\rangle \langle u_n(\xi, \cdot)|$$

with $u_n(\xi, \cdot) = |D_\xi|^{-1} v_n(\xi, \cdot) \in H_\xi^1(Q_\ell)$, from which we deduce, adapting the proof in [8, Eq. (4.42)], that

$$\|\sqrt{\rho_{|\gamma|}}\|_{H^1(Q_\ell)} \leq \|\gamma\|_{X_1^2}^{1/2}, \quad (\text{B.14})$$

since $\rho_{|\gamma|} = \int_{Q_\ell^*} \sum_{n \geq 1} |\mu_n(\xi)| |u_n(\xi, \cdot)|^2 d\xi$. For every $\xi \in Q_\ell^*$, the kernel of $W_{\geq m, \gamma, \xi}$ writes

$$W_{\geq m, \gamma, \xi}(x, y) = \int_{Q_\ell^*} W_{\geq m, \ell}^\infty(\xi' - \xi, x - y) \gamma_{\xi'}(x, y) d\xi'. \quad (\text{B.15})$$

We now use the bound (B.1) on $W_{\geq m, \ell}^\infty$ and split the kernel $W_{\geq m, \gamma, \xi}$ in two terms :

$$\begin{aligned} W_{\geq m, \gamma, \xi}(x, y) &= \int_{Q_\ell^*} G_\ell(x - y) \gamma_{\xi'}(x, y) d\xi' + \int_{Q_\ell^*} f(\xi' - \xi, x - y) \gamma_{\xi'}(x, y) d\xi' \\ &= G_\ell(x - y) \gamma(x, y) + \int_{Q_\ell^*} f(\xi' - \xi, x - y) \gamma_{\xi'}(x, y) d\xi', \end{aligned} \quad (\text{B.16})$$

with $f \in L^\infty(2Q_\ell^* \times 2Q_\ell)$. The second term in (B.16) lies in $L^\infty(Q_\ell^*; L^2(Q_\ell \times Q_\ell))$ as soon as $\gamma \in \mathfrak{S}_{2,1}$. Therefore, the delicate contribution comes from the term involving the potential G_ℓ that we bound from above by using (2.19):

$$\begin{aligned} \iint_{Q_\ell \times Q_\ell} |G_\ell(x - y)|^2 \rho_{|\gamma|}(x) \rho_{|\gamma|}(y) dx dy &\leq \| |G_\ell|^2 \|_{L^1(Q_\ell)} \cdot \|\rho\|_{L^3(Q_\ell)} \|\rho_{|\gamma|}\|_{L^{3/2}(Q_\ell)} \\ &\leq \|G_\ell\|_{L^2(Q_\ell)}^2 \|\sqrt{\rho_{|\gamma|}}\|_{L^6(Q_\ell)}^2 \|\sqrt{\rho_{|\gamma|}}\|_{L^3(Q_\ell)}^2 \\ &\leq C \|G_\ell\|_{L^2(Q_\ell)}^2 \|\sqrt{\rho_{|\gamma|}}\|_{H^1(Q_\ell)}^4. \end{aligned}$$

Finally, using (B.14), we have

$$\|W_{\geq m, \gamma}\|_{\mathfrak{S}_{2,\infty}} \leq C \|\gamma\|_{X_1^2}, \quad (\text{B.17})$$

where C is a positive constant that depends only on m and ℓ .

We now study the contribution of the term involving $W_{< m, \ell}^\infty$, that carries the singularities in the η variable. For this part of the proof, we only need that $\gamma \in Y$. Then, we may write

$$\begin{aligned} &\left| \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} W_{< m, \ell}^\infty(\xi - \xi', x - y) \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy \right| \\ &\leq \left| \int_{Q_\ell^*} \iint_{Q_\ell \times Q_\ell} \frac{4\pi}{\ell^3} \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \frac{e^{-i(\xi' - \xi - \frac{2\pi k}{\ell}) \cdot (x-y)}}{|\xi' - \xi - \frac{2\pi k}{\ell}|^2} \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy d\xi' \right| \\ &\leq \frac{4\pi}{\ell^3} \left| \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{Q_\ell^* + \frac{2k\pi}{\ell}} \iint_{Q_\ell \times Q_\ell} \frac{e^{-i(\xi' - \xi) \cdot (x-y)}}{|\xi' - \xi|^2} \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy d\xi' \right| \\ &\leq \frac{4\pi}{\ell^3} \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{Q_\ell^* + \frac{2k\pi}{\ell}} \frac{d\xi'}{|\xi' - \xi|^2} \left| \left(e^{i(\xi' - \xi) \cdot (\cdot)} \phi_\xi(\cdot), \gamma_{\xi'} e^{i(\xi' - \xi) \cdot (\cdot)} \psi_\xi(\cdot) \right)_{L^2} \right| \\ &\leq C_{\leq m, \ell} \operatorname{ess\,sup}_{\xi' \in Q_\ell^*} \|\gamma_{\xi'}\|_{\mathcal{B}(L_\xi^2)} \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2} \\ &= C_{\leq m, \ell} \|\gamma\|_Y \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2} \end{aligned} \quad (\text{B.18})$$

with

$$C_{\leq m, \ell} = \frac{4\pi}{\ell^3} \sup_{\xi \in Q_\ell^*} \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq (m-1)}} \int_{Q_\ell^* + \frac{2k\pi}{\ell}} \frac{d\xi'}{|\xi' - \xi|^2} = \frac{(2m-1)}{2\pi\ell} \int_{[-1,1]^3} \frac{d\xi'}{|\xi'|^2}. \quad (\text{B.19})$$

Thus, for every $\gamma \in Y$, $W_{< m, \gamma} \in Y$ and

$$\|W_{< m, \gamma}\|_Y \leq C_{\leq m, \ell} \|\gamma\|_Y. \quad (\text{B.20})$$

We now make a further assumption on γ ; namely, $\gamma \in \mathfrak{S}_{1, \infty}$. (Actually, we need $\gamma \in \mathfrak{S}_{1, 4}$.) We first observe that

$$\gamma_{\xi' + \frac{2k\pi}{\ell}}(x, y) = e^{\frac{2ik\pi}{\ell} \cdot (x-y)} \gamma_{\xi'}(x, y) \quad \text{for every } \xi' \in Q_\ell^*, k \in \mathbb{Z}^3 \text{ and } x, y \in \mathbb{R}^3.$$

In particular, $\rho_{\gamma_{\xi' + \frac{2k\pi}{\ell}}} = \rho_{\gamma_{\xi'}}$ for every $\xi' \in Q_\ell^*$ and $k \in \mathbb{Z}^3$, and the function of $\xi \mapsto \text{Tr}_{L_\xi^2}(\gamma_\xi)$ is Q_ℓ^* -periodic. Next, we write

$$\begin{aligned} & \left| \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} \frac{4\pi}{\ell^3} \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \frac{e^{-i(\xi' - \xi - \frac{2\pi k}{\ell}) \cdot (x-y)}}{|\xi' - \xi - \frac{2\pi k}{\ell}|^2} \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy \right| \\ & \leq \frac{4\pi}{\ell^3} \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{Q_\ell^* + \frac{2k\pi}{\ell}} \frac{d\xi'}{|\xi' - \xi|^2} \iint_{Q_\ell \times Q_\ell} \rho_{|\gamma_{\xi'}|}^{1/2}(x) \rho_{|\gamma_{\xi'}|}^{1/2}(y) |\psi_\xi(y)| |\phi_\xi^*(x)| dx dy \\ & = \frac{1}{2\pi^2} \int_{(2m-1)Q_\ell^*} \frac{d\xi'}{|\xi' - \xi|^2} \iint_{Q_\ell \times Q_\ell} \rho_{|\gamma_{\xi'}|}^{1/2}(x) \rho_{|\gamma_{\xi'}|}^{1/2}(y) |\psi_\xi(y)| |\phi_\xi(x)| dx dy \\ & \leq \frac{1}{2\pi^2} \left(\int_{(2m-1)Q_\ell^*} \frac{\|\gamma_{\xi'}\|_{\mathfrak{S}_1(\xi')}}{|\xi' - \xi|^2} d\xi' \right) \|\psi_\xi\|_{L_\xi^2} \|\phi_\xi\|_{L_\xi^2}, \end{aligned} \quad (\text{B.21})$$

where the last estimate follows from the Cauchy–Schwarz inequality. Here and below we use the fact that

$$\sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{Q_\ell^*} f(\xi' - \frac{2\pi k}{\ell}) d\xi' = \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{\frac{2\pi k}{\ell} + Q_\ell^*} f(\xi') d\xi' = \frac{\ell^3}{(2\pi)^3} \int_{(2m-1)Q_\ell^*} f(\xi') d\xi',$$

since $(Q_\ell^* + 2\pi k/\ell) \cap (Q_\ell^* + 2\pi k'/\ell) = \emptyset$ whenever $k, k' \in \mathbb{Z}^3$ with $k \neq k'$. We focus on the quantity inside the brackets in the last inequality. By Hölder's inequality, for $\gamma \in \mathfrak{S}_{1, \infty}$, and for some constant $C'_{\leq m, \ell}$, we obtain

$$\begin{aligned} \|W_{< m, \gamma}\|_Y & \leq \frac{1}{2\pi^2} \text{ess sup}_{\xi \in Q_\ell^*} \int_{(2m-1)Q_\ell^*} \frac{\|\gamma_{\xi'}\|_{\mathfrak{S}_1(\xi')}}{|\xi' - \xi|^2} d\xi' \\ & \leq C \text{ess sup}_{\xi \in Q_\ell^*} \left(\int_{(2m-1)Q_\ell^*} \frac{d\xi'}{|\xi' - \xi|^{8/3}} \right)^{3/4} \left(\int_{(2m-1)Q_\ell^*} \|\gamma_{\xi'}\|_{\mathfrak{S}_1(\xi')}^4 d\xi' \right)^{1/4} \\ & = (2m-1)^{3/4} C \left(\int_{(2m-1)Q_\ell^*} \frac{d\xi'}{|\xi' - \xi|^{8/3}} \right)^{3/4} \|\gamma\|_{\mathfrak{S}_{1, 4}} \\ & \leq C'_{\leq m, \ell} \|\gamma\|_{\mathfrak{S}_{1, \infty}}^{3/4} \|\gamma\|_{\mathfrak{S}_{1, 1}}^{1/4}. \end{aligned} \quad (\text{B.22})$$

Following the lines of the proof of (B.21) and (B.22), we obtain, for every $\xi \in Q_\ell^*$,

$$\|W_{< m, \gamma, \xi}(\cdot, \cdot)\|_{L^2(Q_\ell \times Q_\ell)} \leq C \sum_{\substack{k \in \mathbb{Z}^3 \\ |k|_\infty \leq m-1}} \int_{Q_\ell^*} \frac{\|\gamma_{\xi'}(\cdot, \cdot)\|_{L^2(Q_\ell \times Q_\ell)}}{|\xi' - \xi - \frac{2\pi k}{\ell}|^2} d\xi' = C \int_{(2m-1)Q_\ell^*} \frac{\|\gamma_{\xi'}\|_{\mathfrak{S}_2(\xi')}}{|\xi' - \xi|^2} d\xi',$$

and, by Young's convolution inequality, the right-hand side belongs to $L^p(Q_\ell^*)$ as soon as $\gamma \in \mathfrak{S}_{2,p}$ with $1 \leq p \leq +\infty$ (which is guaranteed whenever $\gamma \in \mathfrak{S}_{1,p}$), where C is a positive constant which depends only on m and ℓ . In particular, for every $1 \leq p \leq +\infty$,

$$\|W_{<m,\gamma}\|_{\mathfrak{S}_{2,p}} \leq C \|\gamma\|_{\mathfrak{S}_{2,p}}. \quad (\text{B.23})$$

Since $\|\gamma\|_{\mathfrak{S}_{1,1}} \leq \|\gamma\|_X$ and $|D^0|^{-1/2} \leq 1$, the statement of the lemma follows : From (B.9) and (B.18), we obtain (4.8); from (B.9) and (B.22), we obtain (4.9); from (B.11) and (B.18), we obtain (4.10). More precisely,

$$C_W = C_H + C_\ell, \quad C'_W = C_G + C_\ell, \quad C''_W = C_H + C'_\ell, \quad (\text{B.24})$$

with

$$C_\ell := \inf_{\substack{m \in \mathbb{N} \\ m \geq 2}} \left(\frac{C_{\geq m}}{\ell} + C_{\leq m,\ell} \right), \quad C'_\ell := \inf_{\substack{m \in \mathbb{N} \\ m \geq 2}} \left(\frac{C_{\geq m}}{\ell} + C'_{\leq m,\ell} \right). \quad (\text{B.25})$$

C Proof of Lemma 4.7

Analogously to (B.6), we have

$$\|V_{\gamma,\xi}\psi_\xi\|_{L^2_\xi} = \sup_{\phi_\xi \in L^2_\xi, \|\phi_\xi\|_{L^2_\xi}=1} |(\phi_\xi, V_{\gamma,\xi}\psi_\xi)|. \quad (\text{C.1})$$

We can rewrite as $W_\ell^\infty = W_{<m,\ell}^\infty + G_\ell + (W_{\geq m,\ell}^\infty - G_\ell)$. According to Proposition B.1 and (B.18), the terms associated to $W_{<m,\ell}^\infty$ and $(W_{\geq m,\ell}^\infty - G_\ell)$ are easily bounded. The aim of this section is to get a better estimate on the following term :

$$\iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \rho_{\gamma'}(y) \phi_\xi^*(x) \psi_\xi(x) dx dy - \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \phi_\xi^*(x) \gamma_{\xi'}(x,y) \psi_\xi(y) dx dy.$$

From now on, for any function $f \in L^2(Q_\ell, \mathbb{C}^4)$, we denote $f := (f^\alpha)_{1 \leq \alpha \leq 4}$. We use the decomposition (2.14) for $\gamma \in \mathfrak{S}_{1,1} \cap Y$ such that $\gamma = \gamma^*$. Then as $G(x) = G(-x)$, for almost every $\xi \in Q_\ell^*$, we may write

$$\begin{aligned} & \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left[\rho_{\gamma'}(y) \phi_\xi^*(x) \psi_\xi(x) - \phi_\xi^*(x) \gamma_{\xi'}(x,y) \psi_\xi(y) \right] dx dy \\ &= \sum_{n \geq 1} \lambda_n(\xi') \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(|u_n(\xi', y)|^2 \phi_\xi^*(x) \psi_\xi(x) - \phi_\xi^*(x) u_n(\xi', x) u_n^*(\xi', y) \psi_\xi(y) \right) dx dy \\ &= \frac{1}{2} \sum_{n \geq 1} \sum_{1 \leq \alpha, \beta \leq 4} \lambda_n(\xi') \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(u_n^\alpha(\xi', y) \phi_\xi^\beta(x) - \phi_\xi^\alpha(y) u_n^\beta(\xi', x) \right)^* \\ & \quad \times \left(u_n^\alpha(\xi', y) \psi_\xi^\beta(x) - \psi_\xi^\alpha(y) u_n^\beta(\xi', x) \right) dx dy. \end{aligned} \quad (\text{C.2})$$

Estimate for (4.12). By Lemma A.1, we have

$$\begin{aligned} & \left| \sum_{1 \leq \alpha, \beta \leq 4} \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(u_n^\alpha(\xi', y) \phi_\xi^\beta(x) - \phi_\xi^\alpha(y) u_n^\beta(\xi', x) \right)^* \left(u_n^\alpha(\xi', y) \psi_\xi^\beta(x) - \psi_\xi^\alpha(y) u_n^\beta(\xi', x) \right) dx dy \right| \\ & \leq \left(\sum_{1 \leq \alpha, \beta \leq 4} \iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)|^2 \left| u_n^\alpha(\xi', y) \psi_\xi^\beta(x) - u_n^\beta(\xi', x) \psi_\xi^\alpha(y) \right|^2 dx dy \right)^{1/2} \\ & \quad \times \left(\sum_{1 \leq \alpha, \beta \leq 4} \iint_{Q_\ell \times Q_\ell} \left| u_n^\alpha(\xi', y) \phi_\xi^\beta(x) - u_n^\beta(\xi', x) \phi_\xi^\alpha(y) \right|^2 dx dy \right)^{1/2}. \end{aligned} \quad (\text{C.3})$$

Thus according to the Cauchy–Schwarz inequality, we have

$$\sum_{1 \leq \alpha, \beta \leq 4} \left| \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(u_n^\alpha(\xi', y) \phi_\xi^\beta(x) - \phi_\xi^\alpha(y) u_n^\beta(\xi', x) \right)^* \left(u_n^\alpha(\xi', y) \psi_\xi^\beta(x) - \psi_\xi^\alpha(y) u_n^\beta(\xi', x) \right) dx dy \right|$$

$$\begin{aligned} &\leq 2 \left(\sum_{1 \leq \alpha, \beta \leq 4} \iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)|^2 |u_n(\xi', y)|^2 |\psi_\xi(x)|^2 dx dy \right)^{1/2} \left(\iint_{Q_\ell \times Q_\ell} |u_n^*(\xi', y)|^2 |\phi_\xi^*(x)|^2 dx dy \right)^{1/2} \\ &\leq 2C_G \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2}. \end{aligned}$$

Substituting this inequality into (C.2) and using the decomposition (2.15), we get

$$\begin{aligned} &\left| \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(\rho_{\gamma_{\xi'}} \phi_\xi^*(x) \psi_\xi(x) - \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) \right) dx dy \right| \\ &\leq C_G \sum_{n \geq 1} |\lambda_n(\xi')| \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2} = C_G \|\gamma_{\xi'}\|_{\mathfrak{S}_1(\xi')} \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2}, \end{aligned}$$

from which we get

$$\begin{aligned} &\left| \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \rho_\gamma(y) \phi_\xi^*(x) \psi_\xi(x) dx dy - \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy \right| \\ &\leq C_G \|\gamma\|_{\mathfrak{S}_{1,1}} \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2}. \end{aligned} \quad (\text{C.4})$$

Combining (C.4) with Proposition B.1 and (B.18), we get for any $\phi_\xi \in L_\xi^2$ and $\psi_\xi \in H_\xi^1$,

$$|(\phi_\xi, V_{\gamma, \xi} \psi_\xi)| \leq (C_G + C_\ell) \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2},$$

hence (4.12) with

$$C'_{EE} := C_G + C_\ell \quad (\text{C.5})$$

with C_ℓ given in (B.25).

Estimate for (4.11). As $\gamma \in Z$, we use the decomposition (B.8) for γ_ξ . Analogously to (C.3), we also have

$$\begin{aligned} &\left| \sum_{1 \leq \alpha, \beta \leq 4} \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \left(u_n^\alpha(\xi', y) \phi_\xi^\beta(x) - \phi_\xi^\alpha(y) u_n^\beta(\xi', x) \right)^* \left(u_n^\alpha(\xi', y) \psi_\xi^\beta(x) - \psi_\xi^\alpha(y) u_n^\beta(\xi', x) \right) dx dy \right| \\ &\leq 2 \left(\iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)| |u_n(\xi', y)|^2 |\psi_\xi(x)|^2 dx dy \right)^{1/2} \left(\iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)| |u_n^*(\xi', y)|^2 |\phi_\xi^*(x)|^2 dx dy \right)^{1/2} \end{aligned}$$

from which by the decomposition (B.8) we get

$$\begin{aligned} &\left| \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \rho_\gamma(y) \phi_\xi^*(x) \psi_\xi(x) dx dy - \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \phi_\xi^*(x) \gamma_{\xi'}(x, y) \psi_\xi(y) dx dy \right| \\ &\leq \left(\int_{Q_{\xi'}} d\xi' \sum_{n \geq 1} |\lambda_n(\xi')| \iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)| |u_n(\xi', y)|^2 |\psi_\xi(x)|^2 dx dy \right)^{1/2} \\ &\quad \times \left(\int_{Q_{\xi'}} d\xi' \sum_{n \geq 1} |\lambda_n(\xi')| \iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)| |u_n(\xi', y)|^2 |\phi_\xi^*(x)|^2 dx dy \right)^{1/2} \\ &\leq C_H \|\gamma\|_X \|\phi_\xi\|_{L_\xi^2} \|\psi_\xi\|_{L_\xi^2} \end{aligned} \quad (\text{C.6})$$

where the last inequality holds by using Lemma 4.1.

Combining (C.6) with Proposition B.1 and estimate (B.18), we get for any $\phi_\xi \in L_\xi^2$ and $\psi_\xi \in H_\xi^1$,

$$|(\phi_\xi, V_{\gamma, \xi} \psi_\xi)| \leq (C_H + C_\ell) \|\gamma\|_Z \|\phi_\xi\|_{L_\xi^2} \|D_\xi |\psi_\xi|\|_{L_\xi^2},$$

hence (4.12) and

$$C_{EE} := C_H + C_\ell. \quad (\text{C.7})$$

Estimate for (4.13). Combining with Proposition B.1 and estimate (B.18), analogously to (C.6) it can be derived directly from:

$$\begin{aligned} & \left| \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \rho_\gamma(y) \psi_\xi^*(x) \psi_\xi(x) dx dy - \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \psi_\xi^*(x) \gamma_{\xi'}(x,y) \psi_\xi(y) dx dy \right| \\ & \leq \left(\int_{Q_{\xi'}} d\xi' \sum_{n \geq 1} |\lambda_n(\xi')| \iint_{Q_\ell \times Q_\ell} |G_\ell(x-y)| |u_n(\xi', y)|^2 |\psi_\xi(x)|^2 dx dy \right) \\ & \leq C_H \|\gamma\|_{\mathfrak{S}_{1,1}} \| |D_\xi|^{1/2} \psi_\xi \|_{L_\xi^2}^2 \end{aligned}$$

using the decomposition (2.14) for γ_ξ . Hence (4.13) and C_{EE} .

Estimate for (4.14). Notice that $|\gamma_{\xi'}(x,y)| \leq \rho_{\gamma_{\xi'}}(x)^{1/2} \rho_{\gamma_{\xi'}}(y)^{1/2}$ since $\gamma \geq 0$. Thus, according to Lemma A.1 and the Cauchy-Schwarz inequality,

$$\begin{aligned} & \int_{Q_\ell^*} d\xi' \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \psi_\xi^*(x) \gamma_{\xi'}(x,y) \psi_\xi(y) dx dy - \iint_{Q_\ell \times Q_\ell} G_\ell(x-y) \rho_\gamma(x) |\psi_\xi(y)|^2 dx dy \\ & \leq \iint_{Q_\ell \times Q_\ell} (|G_\ell(x-y)| - G_\ell(x-y)) \rho_\gamma(y) |\psi_\xi(x)|^2 dx dy \leq \frac{2C_0}{\ell} \|\gamma\|_{\mathfrak{S}_{1,1}} \|\psi\|_{L_\xi^2}^2. \end{aligned}$$

Combining with Proposition B.1 and (B.18), we get

$$(\psi_\xi, V_{\gamma,\xi} \psi_\xi) \geq - \left(\frac{C_0}{\ell} + C_\ell \right) \|\gamma\|_{\mathfrak{S}_{1,1} \cap Y} \|\psi_\xi\|_{L_\xi^2}^2,$$

hence (4.13) and

$$C_{EE}'' = \frac{2C_0}{\ell} + C_\ell. \quad (\text{C.8})$$

D Numerical results about constants

In this section, we will show the numerical results about the constants used in Remark 2.8 under the condition $\ell = 1000$. Next, we show that Assumption 2.5 is satisfied for $q \leq 17$ for the neutral systems.

We compute numerically the value of the bound of the potential $G_\ell - \frac{1}{|x|}$. First of all, we calculate

$$\sum_{k \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|k|^4} \approx 16.512.$$

Thus, $C_0 \approx 5.019$ and we can choose $C_H = C_G \approx 2.011$. Concerning the estimates involving the potential W_ℓ , we set $m = 2$. When $R \approx \frac{1}{2}$,

$$|C_{\geq 2}| \leq 20.912, \quad C_{\leq 2,1000} \approx 0.010.$$

Thus, we get $C_W \approx 2.042$, and $C'_W \approx 2.042$. Then, $C_{EE} \approx 2.052$, $C'_{EE} \approx 2.052$ and $C''_{EE} \approx 0.041$.

Finally, we estimate $c^*([q])$ which is given by (2.9). Let $u_{p,\xi}(x) = e^{i(\frac{2\pi}{\ell} p + \xi) \cdot x}$ with $p \in \mathbb{Z}^3$. Then $(u_{p,\xi})_{p \in \mathbb{Z}^3}$ is an orthogonal basis on $L_\xi^2(Q_\ell)$. Obviously, $(\Lambda^+ u_{p,\xi})_p$ is also an orthogonal basis on $L_\xi^2(Q_\ell)$. Let

$$V_{[q]} = \text{Span}\{\Lambda^+ u_{p,\xi}(x) \mid p = (j, 0, 0), j \in \{1, \dots, [q]\}\}.$$

Then

$$c^*([q]) \leq \sup_{\xi \in Q_\ell^*} \sup_{u_\xi^+ \in V_{[q]}} \frac{\| |D_\xi|^{1/2} u_\xi^+ \|_{L_\xi^2}^2}{\|u_\xi^+\|_{L_\xi^2}^2} \leq \sqrt{1 + \frac{4\pi^2([q] + 1)^2}{\ell^2}}.$$

Now we can check Assumption 2.5 for $z = q = 17$. The calculation leads to $a \approx 0.010$ and $c^*(17) \leq 1.006$. Thus, we have

- $\kappa + \frac{\alpha}{2} C_{EE} q^+ \approx 0.631 < 1$,
- $2a \sqrt{\max\{(1 - \kappa - \frac{\alpha}{2} C_{EE} q^+)^{-1} (1 - \kappa)^{-1} c^*([q]) q; 1\}} q^+ \leq 0.973 < 1$.

Consequently, Assumption 2.5 is satisfied for $z = q \leq 17$ whenever $\ell = 1000$.

Bibliography

- [1] V. Bach. “Error bound for the Hartree–Fock energy of atoms and molecules”. In: *Commun. Math. Phys.* 147.3 (1992), pp. 527–548.
- [2] J.-M. Barbaroux, M. J. Esteban, and É. Séré. “Some connections between Dirac–Fock and electron-positron Hartree–Fock”. In: *Ann. Henri Poincaré* 6.1 (2005), pp. 85–102.
- [3] J.-M. Barbaroux, W. Farkas, B. Helffer, and H. Siedentop. “On the Hartree–Fock equations of the electron-positron field”. In: *Commun. Math. Phys.* 255.1 (2005), pp. 131–159.
- [4] J.-M. Barbaroux, B. Helffer, and H. Siedentop. “Remarks on the Mittleman max-min variational method for the electron-positron field”. In: *J. Phys. A Math. Gen.* 39.1 (2006), pp. 85–98.
- [5] É. Cancès, A. Deleurence, and M. Lewin. “A new approach to the modeling of local defects in crystals: the reduced Hartree–Fock case”. In: *Commun. Math. Phys.* 281.1 (2008), pp. 129–177.
- [6] I. Catto, C. Le Bris, and P.-L. Lions. *The mathematical theory of thermodynamic limits: Thomas–Fermi type models*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1998, pp. xiv+277. ISBN: 0-19-850161-7.
- [7] I. Catto, C. Le Bris, and P.-L. Lions. “Recent mathematical results on the quantum modeling of crystals”. In: *Mathematical models and methods for ab initio quantum chemistry*. Vol. 74. Lecture Notes in Chem. Springer, Berlin, 2000, pp. 95–119.
- [8] I. Catto, C. Le Bris, and P.-L. Lions. “On the thermodynamic limit for Hartree–Fock type models”. In: *Ann. Inst. Henri Poincaré C Anal. Non Linéaire* 18.6 (2001), pp. 687–760.
- [9] I. Catto, C. Le Bris, and P.-L. Lions. “On some periodic Hartree-type models for crystals”. In: *Ann. Inst. Henri Poincaré C Anal. Non Linéaire* 19.2 (2002), pp. 143–190.
- [10] P. Chaix and D. Iracane. “From quantum electrodynamics to mean field theory: I. The Bogoliubov–Dirac–Fock formalism”. In: *J. Phys. B* 22 (1989), pp. 3791–3814.
- [11] N. E. Christensen. “Relativistic band structure calculations”. In: *Int. J. Quantum Chem.* 25.1 (1984), pp. 233–261.
- [12] N. E. Christensen and B. O. Seraphin. “Relativistic Band Calculation and the Optical Properties of Gold”. In: *Phys. Rev. B* 4.10 (1971), pp. 3321–3344.
- [13] J.-P. Desclaux. “Relativistic Dirac–Fock expectation values for atoms with $Z = 1$ to $Z = 120$ ”. In: *Atomic Data Nucl. Data Tables* 12.4 (1973), pp. 311–406.
- [14] J. Dolbeault, M. J. Esteban, and É. Séré. “Distinguished self-adjoint extension and eigenvalues of operators with gaps. Application to Dirac–Coulomb operators”. In: *J. Spectr. Theory* 13.2 (2023), pp. 491–524.
- [15] H. Eschrig, M. Richter, and I. Opahle. “Chapter 12 - Relativistic Solid State Calculations”. In: *Relativistic Electronic Structure Theory*. Ed. by Peter Schwerdtfeger. Vol. 14. Theoretical and Computational Chemistry. Elsevier, 2004, pp. 723–776.
- [16] M. J. Esteban and É. Séré. “Solutions of the Dirac–Fock equations for atoms and molecules”. In: *Commun. Math. Phys.* 203.3 (1999), pp. 499–530.
- [17] M. J. Esteban and É. Séré. “Nonrelativistic limit of the Dirac–Fock equations”. In: *Ann. Henri Poincaré* 2.5 (2001), pp. 941–961.
- [18] M. J. Esteban and É. Séré. “A max-min principle for the ground state of the Dirac–Fock functional”. In: *Mathematical results in quantum mechanics (Taxco, 2001)*. Vol. 307. Contemp. Math. American Math Mathematical Society Providence, 2002, pp. 135–141.
- [19] M. Ghimenti and M. Lewin. “Properties of periodic Hartree–Fock minimizers”. In: *Calc. Var. Partial Differ. Equ.* 35.1 (2009), pp. 39–56.
- [20] I. P. Grant. “Relativistic calculation of atomic structures”. In: *Adv. Phys.* 19.82 (1970), pp. 747–811.
- [21] P. Gravejat, M. Lewin, and É. Séré. “Ground state and charge renormalization in a nonlinear model of relativistic atoms”. In: *Commun. Math. Phys.* 286.1 (2009), pp. 179–215.
- [22] P. Gravejat, M. Lewin, and É. Séré. “Renormalization and asymptotic expansion of Dirac’s polarized vacuum”. In: *Commun. Math. Phys.* 306.1 (2011), pp. 1–33.

- [23] M. Griesemer, R. T. Lewis, and H. Siedentop. “A minimax principle for eigenvalues in spectral gaps: Dirac operators with Coulomb potentials”. In: *Doc. Math.* 4 (1999), pp. 275–283.
- [24] C. Hainzl, M. Lewin, and É. Séré. “Existence of Atoms and Molecules in the Mean-Field Approximation of No-Photon Quantum Electrodynamics”. In: *Arch. Ration. Mech. Anal.* 192.3 (2009), pp. 453–499.
- [25] C. Hainzl, M. Lewin, É. Séré, and J. P. Solovej. “Minimization method for relativistic electrons in a mean-field approximation of quantum electrodynamics”. In: *Phys. Rev. A* 76 (2007), p. 052104.
- [26] I. W. Herbst. “Spectral theory of the operator $(p^2 + m^2)^{1/2} - Ze^2/r$ ”. In: *Commun. Math. Phys.* 53.3 (1977), pp. 285–294.
- [27] M. Huber and H. Siedentop. “Solutions of the Dirac–Fock equations and the energy of the electron-positron field”. In: *Arch. Ration. Mech. Anal.* 184.1 (2007), pp. 1–22.
- [28] M. Kadec, M. Repisky, and K. Ruud. “All-electron fully relativistic Kohn–Sham theory for solids based on the Dirac–Coulomb Hamiltonian and Gaussian-type functions”. In: *Phys. Rev. B* 99.20 (2019), p. 205103.
- [29] T. Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Reprint of the 1980 edition. Springer, Berlin, 1995, pp. xxii+619.
- [30] Y. K. Kim. “Relativistic self-consistent-field theory for closed-shell atoms”. In: *Phys. Rev.* 154 (1 Feb. 1967), pp. 17–39.
- [31] E. H. Lieb. “Variational principle for many-fermion systems”. In: *Phys. Rev. Lett.* 46.7 (1981), pp. 457–459.
- [32] E. H. Lieb and B. Simon. “The Thomas–Fermi theory of atoms, molecules and solids”. In: *Adv. Math.* 23.1 (1977), pp. 22–116.
- [33] P.-L. Lions. “Solutions of Hartree–Fock equations for Coulomb systems”. In: *Commun. Math. Phys.* 109.1 (1987), pp. 33–97.
- [34] L. Meng. “A rigorous justification of the Mittleman’s approach to the Dirac–Fock model”. In: *Calc. Var. Partial Differ. Equ.* 63.2 (2024).
- [35] M. H. Mittleman. “Theory of relativistic effects on atoms: Configuration-space Hamiltonian”. In: *Phys. Rev. A* 24.3 (1981), pp. 1167–1175.
- [36] É. Paturel. “Solutions of the Dirac–Fock equations without projector”. In: *Ann. Henri Poincaré* 1.6 (2000), pp. 1123–1157.
- [37] P. Pyykko and J.-P. Desclaux. “Relativity and the periodic system of elements”. In: *Acc. Chem. Res.* 12.8 (1979), pp. 276–281.
- [38] M. Reed and B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich, Publishers], New York, 1978, pp. xv+396.
- [39] É. Séré. “A new definition of the Dirac–Fock ground state”. In: *Commun. Math. Phys.* 404.3 (2023), pp. 1275–1307.
- [40] B. Simon. *Trace ideals and their applications*. Vol. 120. American Mathematical Society, Providence, 2005.
- [41] B. Swirles. “The relativistic self-consistent field”. In: *Proc. R. Soci. Lond. Ser. A - Math. Phys. Sci.* 152.877 (1935), pp. 625–649.