



HAL
open science

Effects of a Cognitive Schema Account on the Stigma of Schizophrenia: A Study in a French University Student Sample

Charlotte Devoisin-Lagarde, Bastien Trémolière, Elodie Charbonnier, Serge Caparos

► To cite this version:

Charlotte Devoisin-Lagarde, Bastien Trémolière, Elodie Charbonnier, Serge Caparos. Effects of a Cognitive Schema Account on the Stigma of Schizophrenia: A Study in a French University Student Sample. *American Journal of Psychology*, 2022, 135 (3), pp.287 - 300. 10.5406/19398298.135.3.03 . hal-03871380v2

HAL Id: hal-03871380

<https://hal.science/hal-03871380v2>

Submitted on 25 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of a cognitive schema account on the stigma of schizophrenia: An exploratory study in a French university student sample

Charlotte Devoisin-Lagarde ^a

^a UNIV. NIMES, Student in Master of Clinical Psychology and CBT, F- 30021 Nîmes Cedex 1, France

Bastien Trémolière ^b

^b UNIV. NIMES, APSY-V, F-30021 Nîmes Cedex 1, France

Serge Caparos ^{c,d}

^c Université Paris 8, Laboratoire DysCo, 2 rue de la Liberté, 93526, Saint Denis

^d Institut Universitaire de France

Elodie Charbonnier ^b

^b UNIV. NIMES, APSY-V, F-30021 Nîmes Cedex 1, France

This is the PrePrint version. It may not exactly replicate the final published version of the article. It is not the copy of record. Published version of the article may be found at this link:
<https://doi.org/10.5406/19398298.135.3.03>

Abstract

Background: Amongst people with psychiatric disorders, people experiencing schizophrenia are among those who are victims of high levels of stigmatization.

Aims: The present study explored how educational strategies emphasizing different types of etiological accounts of schizophrenia may help reduce the level of stigma towards the pathology.

Methods: A total of 378 students were randomly assigned to one of four groups. Three experimental groups were presented with a vignette which introduced one out of three different etiological accounts of schizophrenia (Cognitive distortion, Cognitive Schema, or Biogenetics). The fourth group was presented with a vignette unrelated to schizophrenia. The participants completed questionnaires which measured their attitudes, empathic concern and social distance towards people with schizophrenia, before and after reading the vignette.

Results: The "Cognitive Schema" etiological account, which emphasized maladaptive cognitive schemas, was the only one which led to a significant decrease in stigma. The latter decrease was mostly driven by an increase in the level of empathic concern towards people with schizophrenia.

Conclusions: The results of the present study suggest that using cognitive-schema cues may be a promising avenue to reduce the stigmatization of schizophrenia. Future research is warranted which will explore the many factors jointly that explain schizophrenia for use in anti-stigma campaigns.

Keywords: Stigma, causal attributions, empathic concern, maladaptive cognitive schema

1. Introduction

About one in four people will suffer from a psychiatric disorder in their lifetime (World Health Organization, 2001). Not only do individuals with psychiatric disorders have to deal with their symptoms, but they must also cope with the stigmatization that they generally suffer from (Corrigan & Watson, 2002a). Stigmatization is a process of exclusion leading an individual, on the basis of a given characteristic that does not conform to a social norm, to be identified as abnormal or deviant, and reduced to this characteristic (Crocker & Major, 1989; Goffman, 1963).

Levels of stigmatization differ a lot, and are mainly function of the diagnosis (Parle, 2012): among all psychiatric disorders, people experiencing schizophrenia are victims of high levels of rejection and stigmatization (Angermeyer & Schulze, 2001; Castillo et al., 2008; Pescosolido et al., 2010). They are commonly perceived by lay people as dangerous, incompetent, and unpredictable (Angermeyer & Dietrich, 2006; Angermeyer & Matschinger, 2004; Corrigan et al., 2001; Crisp et al., 2000; Dietrich et al., 2004; Graves et al., 2005; Link et al., 1999). Schizophrenia is frequently and intimately associated with the term "violence" and is perceived as the "model of madness" by society (Castillo et al., 2008). Consistent with this observation, some of the symptoms of schizophrenia, such as inappropriate affect and the propensity to talk aloud to oneself, are manifest indicators of a psychiatric disorder and may lead to stigmatizing reactions (see Corrigan, 2000).

Stigmatization is likely to have detrimental consequences for people with schizophrenia. The consequences of stigmatization can be observed either within the psychological sphere (Berge and Ranney, 2005), the social and professional sphere (Daumerie et al., 2012), the family sphere (González-Torres et al., 2007), or the sphere of medical care (Demyttenaere et al., 2004; Fung et al., 2008). In the specific case of France, recent studies

have confirmed that the levels of stigmatization of schizophrenia are high (Angermeyer et al., 2013; Durand-Zaleski et al., 2012; Lampropoulos et al., 2017).

According to attribution theory (Corrigan, 2000; Weiner et al., 1988; Weiner, 1995), an important factor that may contribute to the stigmatization of a disorder is the etiological account that people attribute to this disorder. Specifically, level of control (i.e., the extent to which people can actively act on their disease) and stability of the disorder (i.e., the extent to which the disorder is maintained over time) are thought key elements of stigma. Anti-stigma campaigns which have emphasized biogenetic or psychosocial factors to account for the development of schizophrenia, have had mixed effects on measured levels of stigma (see the meta-analyses of Kvaale et al., 2013a; Kvaale et al., 2013b). Endorsing biological factors as the cause of schizophrenia was associated with a greater desire for social distance, i.e., the desire not to interact with them (Angermeyer & Matschinger, 2005; Dietrich et al., 2004). Biogenetic explanations may reinforce feelings of dangerousness and therefore the desire for social distance (Corrigan and Watson, 2004; Mehta & Farina, 1997; Phelan, 2005; Phelan et al., 2006). Biogenetic causal beliefs are also positively related to prejudice and fear (Read et al., 2006). It is possible that biogenetic accounts promote the idea that the person may be considered to have little control over his/her behavior, which may lead to expectations of higher levels of unpredictable behaviors and may increase fear responses in lay people (Angermeyer & Matschinger, 2003).

Likewise, psychosocial accounts of mental disorders (e.g., accounts such as stress, death of a loved one, insults at school) have generally shown mixed effects on the level of stigma (see the systematic review and meta-analysis of Schomerus et al., 2012). Stress-related accounts of mental disorders have sometimes been shown to be associated with greater tolerance of the disorders (Martin et al., 2000; Schnittker, 2008; Van 't Veer et al., 2006), but

not in all studies (Bag et al., 2006; Jorm & Griffiths, 2008; Martin et al., 2007). Similar mixed results have been observed with early-stressor accounts (Dietrich et al., 2004; Schnittker, 2008; Van 't Veer et al., 2006). According to Schomerus et al. (2014), lay people may infer from some psychosocial accounts that people experiencing schizophrenia are responsible for their disorder (e.g., with accounts such as the use of psychoactive substances), and this may increase stigma as a result. On the other hand, other psychosocial accounts (e.g., childhood sexual abuse) may reduce the perceived level of personal responsibility and, consequently, improve social acceptance. However, psychosocial explanations, such as childhood sexual abuse, may drive lay people to consider schizophrenia as incurable, and this may increase stigma as a result.

In an attempt to reduce stigma, the use of cognitive accounts of schizophrenia have been largely overlooked. It appears that, so far, such accounts have been mobilized mainly in the context of depression (Botha & Dozois, 2015). Yet, cognitive factors – for instance cognitive distortion factors and maladaptive cognitive schemas – are known to hold an important role in the development of symptoms in schizophrenia (Bortolon et al., 2013; Sundag et al., 2016). Early maladaptive schemas develop during childhood, via interpersonal interactions, and form a template that guides the interpretation of later experience. They are extremely stable, considered to be more unconditional, automatic, and to increase the risk of developing a mental disorder (Schmidt et al., 1995). Cognitive models of schizophrenia suggest that exposure to negative life events during childhood (e.g., school harassment) promotes the development of maladaptive cognitive schemas, including a threatening worldview, external attribution of negative events and experiences, which may favor the development of schizophrenia, and more specifically, paranoid delusions (Bentall et al., 2009; Garety et al., 2001; Howes and Murray, 2014). Early maladaptive schemas are higher people experiencing

schizophrenia (Boyda et al., 2018; Giblin et al., 2004; Khosravani et al., 2019; Sundag et al., 2016). Negative schemas ease up the development of delusions and hallucinations, including a decrease in performance at reality tests (Beck & Rector, 2005; Birchwood et al., 2000; Garety et al., 2001).

Cognitive distortions refer to erroneous interpretations of situations (e.g., minimizing positive accomplishments, personalizing neutral events; Beck, 1979). Cognitive distortions are important early risk factors for psychosis (de Paula et al., 2015; Gawęda et al., 2018). More precisely, an exaggerated tendency to pay attention to threat (Gawęda et al., 2015; Prochwicz & Klosowska, 2017), external attribution bias (So et al., 2015), jumping to conclusions (Ross et al., 2015; McLean et al., 2017) or bias against disconfirmatory evidence (Zawadzki et al., 2012) are important factors driving delusions. In the same way, deficits in recognizing the source of information plays a central role in hallucinations (see the meta-analysis of Waters et al., 2012).

Because cognitive distortions may be perceived by lay people as controllable (Cook & Wang, 2011; Goldstein & Rosselli, 2003), accounts based on such factor may *increase* public stigma. On the other hand, maladaptive cognitive schemas emerge during childhood, through experiences of harmful events (e.g., critical, abusive or neglectful events which do not meet the needs of the child) (Young et al., 2003). Thus, cognitive schema accounts emphasize the importance of external factors as causes of the disorder, they support the idea that patients have a reduced control over the disorder and, as a result, cannot be responsible for their pathology (Schomerus et al., 2014). Cognitive schema accounts may therefore *reduce* public stigma. Table 1 summarizes the characteristics of the Biogenetic, Cognitive Distortion, and Cognitive schema accounts.

Insert Table 1 here

The present research explored how lay people's beliefs regarding the causes of schizophrenia shape the stigmatization of people experiencing this disorder. To this end, we exposed a sample of French university students to three different etiological accounts of the pathology (Cognitive Distortion, Cognitive Schema, or Biogenetic explanation), and we measured the effect of this exposition on their level of expressed stigma (stigmatizing attitude, desire for social distance, and empathic concern), compared to a control group. We expected the largest decrease in stigma in participants exposed to the Cognitive schema account, a lower or no decrease in participants exposed to the Cognitive Distortion and Genetic accounts, and no decrease in the control group.

2. Method

2.1. Participants

An a priori power analysis was conducted using Gpower (Erdfelder et al., 1996). The analysis indicated that a sample size of 332 participants (83 participants per group) would be sufficient to detect a significant interaction between our text conditions and moment of measurement (pre/post text exposure), with a power of .95 and an alpha of .05, based on a small effect size of 0.05 (η_p^2). Anticipating drop out and exclusion issues, a total of 415 students were recruited from an undergraduate introductory psychology course from [omitted for masked review] University, and randomly assigned to one of the four conditions of our experimental manipulation (Cognitive Distortion, Cognitive Schema, Biogenetic, or Control condition). The participants completed the questionnaire in the lecture hall before lecture started. Thirty-seven participants did not complete the questionnaire and were excluded from the analyses, leaving a final sample of 378 participants (263 females, 2 undefined) aged

between 16 and 58 ($M_{\text{age}} = 19.90$ years; $SD_{\text{age}} = 4.68$ years), assigned to the following conditions: Cognitive Distortion condition ($N = 99$), Cognitive Schema condition ($N = 106$), Biogenetic condition ($N = 107$), or Control condition ($N = 66$).

2.2. Compliance with Ethical Standards

All participants signed a consent form before participation and were informed that their answers would remain anonymous and that their participation was voluntary and could be withdrawn at any time. All the data were stored on a secure university computer. We respected the ethical principles of psychologists and the code of conduct of the American Psychological Association (American Psychological Association, 2002).

2.3. Measures

In the absence of validated tools in French language for our variables of interest, all scales used were translated in French via a back-and-forth translation procedure. Firstly, the English version was translated into French by an English native speaker specialized in this research field. Then, a French native speaker translated this document back into English. The two translations (English to French and French to English) were compared in order to ensure that they were both faithful to the original and that the items in the French version were clear.

2.3.1. Familiarity with schizophrenia and mental illness

To assess people's familiarity with schizophrenia, we used a French translation of the Familiarity with mental illness scale (Corrigan et al., 2003). This 7-item questionnaire measures participants' familiarity with schizophrenia through everyday encounters with the pathology

(e.g., interaction in the workplace, presence of people experiencing the pathology in the entourage or at home). Participants answered using a dichotomous "Yes"/"No" response to each item. Scores ranged from 0 to 7 and higher scores indicated higher familiarity with schizophrenia. We also included two additional questions in order to assess participants' general familiarity with mental illnesses: "Do you have a diagnosis of mental disorder" and "Do you have a member of your family with a diagnosis of mental disorder". Both questions were answered using a dichotomous "Yes"/"No" response.

2.3.2. Attitudes

We used a French translation of the Brief Version of the Attribution Questionnaire (AQ-9) (Corrigan et al., 2002) in order to assess attitudes towards people experiencing schizophrenia. Each item from the attitude questionnaire assesses a reaction towards people with schizophrenia, exploring the domains of blame, anger, pity, help, dangerousness, fear, avoidance, segregation, and coercion (e.g., How angry do persons with schizophrenia make you feel?). Participants gave their answer using 9-point Likert scales (ranging from 1: "Not at all" to 9: "Very much"). Scores ranged from 9 to 81, and higher scores indicated more negative attitudes towards people with schizophrenia. For the sake of comparability with the other measures, we subsequently reversed the questionnaire score and transformed it into a percentage, leading to a percentage score of positive attitude (from 0 to 100). The scale showed acceptable internal consistency (Cronbach's α at T0 = .71 and T1 = .73).

2.3.3. Social Distance

In order to evaluate social distance, we used a French translation of the Social Distance Items (Link et al., 1987), which is the most used tool for assessing social stigma of

schizophrenia (Lampropoulos et al., 2018). It includes seven items displaying social situations (e.g., renting a room) and participants are asked to indicate the extent to which they would be willing to accept to share these situations with people experiencing schizophrenia, using a 4-point Likert scale ranging from 1 ("Definitely willing") to 4 ("Definitely unwilling"). Scores ranged from 7 to 28, and higher scores indicated a preference for greater social distance. Scores were then reversed and transformed into a percentage, leading to a percentage score of social proximity (from 0 to 100). The scale showed very good internal consistency (Cronbach's α at T0 = .86 and T1 = .89).

2.3.4. Empathic concern

In order to measure empathic concern, we used a French translation of the emotional response scale (Batson, 1987; Batson, 1991). Participants were instructed to indicate how much they felt each of six emotions (sympathy, soft-hearted, warm, compassionate, tender and moved) towards individuals experiencing schizophrenia, using a 7-point Likert scale ranging from 1 ("Not at all") to 7 ("Very much"). Scores ranged from 6 to 42, and higher scores indicated greater empathic concern. Scores were then transformed into a percentage score of empathic concern (from 0 to 100). The scale showed very good internal consistency (Cronbach's α at T0 = .87 and T1 = .90).

2.4. Causal explanations of schizophrenia

Four texts were developed in French language and presented to the participants. Three of the texts detailed a specific factor identified as the main cause for the development of schizophrenia: biogenetics, cognitive distortion, or cognitive schema. As it was the case in the vignettes designed by Botha and Dozois (Botha & Dozois, 2015), these three texts were

presented in the form of a (fictitious) scientific article, recently published in the "Journal of Behavioral and Cognitive Therapy" (for the Cognitive Schema and Cognitive Distortion conditions) or the "Journal of Molecular Biology and Genetics" (for the Biogenetics condition) by a fictitious researcher, known worldwide and specialized in the field of schizophrenia. In the article, the researcher emphasized that all the data collected over the past thirty years indicate that schizophrenia is caused by factors related to biogenetics/cognitive distortion/cognitive schema (depending on the condition). All three texts had identical structures and similar contents, except for the target sentences regarding the cause of schizophrenia, that differed according to each condition (see Appendix A1 for the full version of the texts). A fourth text, unrelated to schizophrenia and presenting a topic related to quantum physics, was used in the control condition.

Procedure

After giving their consent, participants provided their basic sociodemographic data (age, sex) and indicated their familiarity with schizophrenia. Next, all participants read a short description of the daily life of an individual experiencing schizophrenia (the description is displayed in the Appendix), in order to prime a correct and realistic conception of the disorder (see Corrigan et al., 2017). Then, participants completed the questionnaires about attitudes (Corrigan et al., 2002), empathic concern (Batson, 1987; Batson, 1991), and social distance (Link et al., 1987) towards people experiencing schizophrenia. In the next step, they were randomly assigned to one of the four conditions: Biogenetics, Cognitive Distortion, Cognitive Schema or Control, and read the vignette relevant to their condition. Finally, they completed for a second time the questionnaires about attitudes, empathic concern and social distance towards people experiencing schizophrenia.

Following the experiment, participants were informed about the purpose of the study. We took caution to emphasize that each scenario they were presented with displayed only one etiological account (Biogenetics, Cognitive Distortion, Cognitive Schema or Control). Then, they received exhaustive information about causal attributions of schizophrenia and the multiple etiological accounts of the disorder.

2.5. Statistical analysis

We conducted a mixed-design analysis of variance using Measure (positive attitude, social proximity, and empathic concern; within-subject), Text condition (Cognitive Schema, Cognitive Distortion, Biogenetics, Control; between-subject), and Moment of measurement (pre-text measure vs. post-text measure; within-subject) as the independent variables, and percentage score (for attitude, social proximity, and empathic concern) as the dependent measure. In addition, we added familiarity, self-reported psychiatric disorder, and self-reported psychiatric disorder in the participant's family as control variables in the analysis. Data were analyzed using the SPSS software (version 24).

3. Results

Figure 1 displays the detailed results distinguishing the effect of text condition for each of the measures and moment of measurement (see also Table 2 for the full descriptive statistics). A main effect of Measure was observed, $F(1.70, 608.56) = 4.83, p = .012, \eta_p^2 = .013$. Most importantly, with regards to our current purpose, the Moment of measurement x Text condition interaction was significant, $F(3, 358) = 4.42, p = .005, \eta_p^2 = .04$. Before exploring the origin of this interaction with pairwise comparisons, and because Measure did not significantly interact with any of our independent variables, we computed a composite score of *acceptance*

(by averaging the percentage scores of positive attitude, social proximity, and empathic concern). Pairwise comparisons using paired sample *t*-tests showed that only the Cognitive Schema condition yielded an increase in acceptance of schizophrenia between T0 ($M_{\text{acceptance}} = 57.52$; $SD_{\text{acceptance}} = 15.02$) and T1 ($M_{\text{acceptance}} = 60.22$; $SD_{\text{acceptance}} = 15.75$), $t(105) = 4.41$, $p < .001$, Cohen's $d = 0.18$. No significant difference was observed between T0 and T1 in the Cognitive Distortion, Biogenetics, or Control condition (all t s $< .99$, all p s $> .32$; see Table A1, in appendix, for full descriptive statistics).

Insert Table 2 and Figure 1 here

4. Discussion

Amongst people with psychiatric disorders, people experiencing schizophrenia are among those who are victims of high levels of stigmatization (Angermeyer & Schulze, 2001; Castillo et al., 2008; Corrigan & Watson, 2002a; Durand-Zaleski et al., 2012; Graves et al., 2005; Schomerus et al., 2014). A research effort is necessary to identify new strategies that may help reduce the public stigma of schizophrenia, and the associated multiple negative consequences for people experiencing this mental disorder. To date, the few researches that exist on this topic have highlighted mixed effects. Educational interventions using biogenetics or psychosocial accounts of the development of schizophrenia do not always lead to a decrease in stigma, and may even sometimes lead to an increase in stigma (Bag et al., 2006; Jorm & Griffiths, 2008; Kvaale et al., 2013a; Kvaale et al., 2013b; Schomerus et al., 2012). In the context of schizophrenia, educational strategies using cognitive accounts have never been evaluated, yet studies performed in the context of depression have shown these strategies to be promising (Botha and Dozois, 2015). The present study addressed this issue and compared

the effect on public stigma of three different educational strategies, based on cognitive distortion, maladaptive cognitive schema, and biogenetics. In line with our expectations, the information that emphasized the role of maladaptive cognitive schemas in the development of schizophrenia was the only one that led to a significant decrease in stigma. In contrast, our results showed that none of the other two accounts of schizophrenia impacted stigma, and neither did stigma vary in the control condition.

Why should a Cognitive Schema account of schizophrenia be more efficient in reducing stigma? In light of attribution theories (Corrigan, 2000; Weiner et al., 1988; Weiner, 1995), level of stigma may depend on etiological account, specifically on control and stability criteria. Previous studies have shown strong positive relationships between patients' control or responsibility in the development of their particular disorder and public stigma towards the disorder in question (Dietrich et al., 2004; Lincoln et al., 2008; Schomerus et al., 2013). In addition, studies have highlighted a positive link between stability of the disorder (i.e., the impossibility to treat or reduce it) and public stigma (Phelan, 2005; Phelan et al., 2006). Because Cognitive Schema accounts of schizophrenia emphasize the importance of external factors (e.g., childhood experiences, such as humiliations at school) as causes of the disorder, they highlight its low controllability. In addition, Cognitive Schema accounts draw attention to the fact that the joint experience of similar situations experienced in childhood (during which cognitive schema is built) and adulthood (during which cognitive schema is activated) is partly responsible for schizophrenia symptoms, putting light on the unstable nature of the disease over time.

The Cognitive Schema exposure effect was driven mainly by an increase in the level of empathic concern towards people experiencing schizophrenia. According to Batson et al., (2007), level of empathy for another person in need depends on three dimensions: perceiving

that the other is in need, valuing the other's welfare, and adopting the other's perspective. In the Cognitive Schema condition, the example given in the vignette was related to a child being humiliated at school, which is a common situation that the participants may have themselves experienced during childhood. This may have increased their propensity to adopt the other's perspective, and, as a result, empathic concern. This result is all the more important given that much research has evidenced that a high empathic concern is an important factor for prosocial motivation and helping behaviors (Batson et al., 2007; Eisenberg & Miller, 1987). According to this idea, emotional changes may induce cognitive changes (i.e., decrease in negative attitudes) and behavioral changes (decrease in desire for social distance).

We believe that this first step of research may open new avenues for future research on the prevention of stigmatization of schizophrenia. First, the present results shed light on the necessity to raise public awareness of schizophrenia. Educational strategies based on etiological accounts of schizophrenia may be used, and they should emphasize both the low stability and low controllability in the disease. We see a great importance in the necessity to balance these two components in the explanation of schizophrenia, which is consistent with the multidimensional and balanced approach proposed by Corrigan and Watson (2004). If biogenetic-related explanations (associated with low control but high stability) are used, they may be accompanied by information about the efficacy of treatments, in order to reduce the feelings of dangerousness and bad prognosis induced by this kind of account. Second, our results call for the need to evaluate the effectiveness of the prevention methods used, the non-standardization of the different explanations being an important factor which possibly explains their mixed effects on stigma (Lambooy & Saias, 2013).

Our present results, although promising, need to be taken with caution, for several reasons which we list here. First, it is worth noting that our effect is a small one. The significant

pairwise difference that we observed for the Cognitive Schema condition (between T0 and T1) showed only a small effect size (Cohen's $d = .18$). We acknowledge that many other factors (and their interaction) contribute to explain stigma. Second, our research follows an experimental paradigm, and aims to untangle the effects of distinct educational strategies presenting different types of etiological accounts of schizophrenia. As a result, each scenario was designed to emphasize only one cause for schizophrenia. It stands to reason that the etiology of schizophrenia is multifactorial, and that maladaptive cognitive schema cannot be considered the only explanation for schizophrenia. Future research is warranted which will explore the many factors jointly that explain schizophrenia. Third, because cognitive schemas develop during childhood (Young et al., 2003), one may worry about the possibility that Cognitive Schema accounts of schizophrenia may drive the public to blame the social environment, and especially the parents, for the development of schizophrenia in a child. This would be an undesirable side-effect of an anti-stigma campaign. Fourth, although the present results show a promising effect of Cognitive Schema accounts in the short term, they need to be replicated in the long term. Because the three measures capturing stigma were obtained immediately after the anti-stigma treatment, one must refrain from overgeneralizing these findings as well as to consider that the positive effects would persist over time. Finally, our sample consisted of predominantly French female university students who are certainly sensitive to the issue of mental health. Consequently, replications need to be conducted with a more representative population before one may be able to generalize such findings.

References

American Psychological Association. (2002). Ethical Principles of Psychologists and Code of Conduct. *American psychologist*, 57(12), 1060–1073. <https://doi.org/10.1037/0003->

066X.57.12.1060

- Angermeyer, M. C., and Dietrich, S. (2006). Public beliefs about and attitudes towards people with mental illness: a review of population studies. *Acta Psychiatrica Scandinavica*, 113(3), 163–179. <https://doi.org/10.1111/j.1600-0447.2005.00699.x>
- Angermeyer, M. C, and Schulze, B. (2001). Reinforcing stereotypes: how the focus on forensic cases in news reporting may influence public attitudes towards the mentally ill. *International Journal of Law and Psychiatry*, 24(4–5), 469–486.
- Angermeyer, M. C., and Matschinger, H. (2003). Public beliefs about schizophrenia and depression: similarities and differences. *Social psychiatry and psychiatric epidemiology*, 38(9), 526–534. <https://doi.org/10.1007/s00127-003-0676-6>
- Angermeyer, M. C., and Matschinger, H. (2005). Causal beliefs and attitudes to people with schizophrenia. *The British Journal of Psychiatry*, 186(04), 331–334. <https://doi.org/10.1192/bjp.186.4.331>
- Angermeyer, M. C, and Matschinger, H. (2004). The stereotype of schizophrenia and its impact on discrimination against people with schizophrenia: results from a representative survey in Germany. *Schizophrenia bulletin*, 30(4), 1049–1061.
- Angermeyer, M. C, Millier, A., Rémuzat, C., Refaï, T., and Toumi, M. (2013). Attitudes and beliefs of the French public about schizophrenia and major depression: results from a vignette-based population survey. *BMC Psychiatry*, 13(1), 313. <https://doi.org/10.1186/1471-244X-13-313>
- Bag, B., Yilmaz, S., and Kirpinar, I. (2006). Factors influencing social distance from people with schizophrenia. *International Journal of Clinical Practice*, 60(3), 289–294. <https://doi.org/10.1111/j.1368-5031.2006.00743.x>
- Batson, C. D. (1987). Prosocial motivation: Is it ever truly altruistic? In L. Berkowitz (Ed.),

Advances in experimental social psychology (pp. 65–122). New York: Academic Press.

Batson, C. Daniel, Eklund, J. H., Chermok, V. L., Hoyt, J. L., and Ortiz, B. G. (2007). An additional antecedent of empathic concern: Valuing the welfare of the person in need. *Journal of personality and social psychology*, 93(1), 65–74.
<https://doi.org/10.1037/0022-3514.93.1.65>

Batson, C.D. (1991). *The altruism question: Toward a social-psychological answer*. Hillsdale, NJ: Erlbaum .

Beck, A. T. (1979). *Cognitive therapy of depression*. Guilford Press.

Beck, A. T., and Rector, N. A. (2005). Cognitive approaches to schizophrenia: theory and therapy. *Annual Review of Clinical Psychology*, 1, 577-606

Bentall, R. P., Rowse, G., Shryane, N., Kinderman, P., Howard, R., Blackwood, N., ... and Corcoran, R. (2009). The cognitive and affective structure of paranoid delusions: a transdiagnostic investigation of patients with schizophrenia spectrum disorders and depression. *Archives of general psychiatry*, 66(3), 236-247

Berge, M., and Ranney, M. (2005). Self-esteem and stigma among persons with schizophrenia: implications for mental health. *Care Management Journals*, 6(3), 139–144. <https://doi.org/10.1891/CMAJ.6.3.139>

Birchwood, M., Meaden, A., Trower, P., Gilbert, P., and Plaistow, J. (2000). The power and omnipotence of voices: Subordination and entrapment by voices and significant others. *Psychological medicine*, 30(2), 337-344

Boyda, D., McFeeters, D., Dhingra, K., and Rhoden, L. (2018). Childhood maltreatment and psychotic experiences: Exploring the specificity of early maladaptive schemas. *Journal of clinical psychology*, 74(12), 2287-2301.

Bortolon, C., Capdevielle, D., Boulenger, J.-P., Gely-Nargeot, M.-C., and Raffard, S. (2013).

Early maladaptive schemas predict positive symptomatology in schizophrenia: A cross-sectional study. *Psychiatry Research*, 209(3), 361–366.

<https://doi.org/10.1016/j.psychres.2013.03.018>

Botha, F. B., and Dozois, D. J. A. (2015). The influence of emphasizing psychological causes of depression on public stigma. *Canadian Journal of Behavioural Science*, 47(4), 313–320. <https://doi.org/10.1037/a0039611>

Castillo, M. C., Lannoy, V., Seznec, J. C., Januel, D., and Petitjean, F. (2008). Étude des représentations sociales de la schizophrénie dans la population générale et dans une population de patients schizophrènes. *L'Évolution Psychiatrique*, 73(4), 615–628.

<https://doi.org/10.1016/J.EVOPSY.2007.06.006>

Cook, T. M., and Wang, J. (2011). Causation beliefs and stigma against depression: Results from a population-based study. *Journal of Affective Disorders*, 133(1–2), 86–92.

<https://doi.org/10.1016/J.JAD.2011.03.030>

Corrigan, P. W., Schmidt, A., Bink, A. B., Nieweglowski, K., Al-Khouja, M. A., Qin, S., and Discont, S. (2017). Changing public stigma with continuum beliefs. *Journal of Mental Health*, 26(5), 411–418. <https://doi.org/10.1080/09638237.2016.1207224>

Corrigan, P. W., and Watson, A. C. (2002a). Understanding the impact of stigma on people with mental illness. *World Psychiatry*, 1(1), 16–20.

Corrigan, P. W., & Watson, A. C. (2002b). The paradox of self-stigma and mental illness. *Clinical Psychology: Science and Practice*, 9(1), 35–53.

Corrigan, P. W., and Watson, A. C. (2004). At issue: Stop the stigma: call mental illness a brain disease. *Schizophrenia Bulletin*, 30(3), 477–479.

Corrigan, P. W., Markowitz, F. E., Watson, A., Rowan, D., and Kubiak, M. A. (2003). An attribution model of public discrimination towards persons with mental illness. *Journal*

of health and Social Behavior, 44(2), 162–179.

- Corrigan, P. W. (2000). Mental health stigma as social attribution: Implications for research methods and attitude change. *Clinical psychology: science and practice, 7(1), 48–67.*
- Corrigan, P. W, Green, A., and Lundin, R. (2001). Familiarity with and social distance from people who have serious mental illness. *Psychiatric services, 52(7), 953–958.*
- Corrigan, P. W, Rowan, D., Green, A., and Lundin, R. (2002). Challenging two mental illness stigmas: Personal responsibility and dangerousness. *Schizophrenia bulletin, 28(2), 293–309.*
- Crisp, A. H., Gelder, M. G., Rix, S., Meltzer, H. I., and Rowlands, O. J. (2000). Stigmatisation of people with mental illnesses. *The British journal of psychiatry, 177, 4–7.*
- Crocker, J., & Major, B. (1989). Social stigma and self-esteem: The self-protective properties of stigma. *Psychological review, 96(4), 608-630.*
- Daumerie, N., Vasseur Bacle, S., Giordana, J.-Y., Bourdais Mannone, C., Caria, A., and Roelandt, J.-L. (2012). Discrimination perceived by people with a diagnosis of schizophrenic disorders. International study of Discrimination and stiGma Outcomes (INDIGO): French results. *L'Encephale, 38(3), 224–231.*
<https://doi.org/10.1016/j.encep.2011.06.007>
- Demyttenaere, K., Bruffaerts, R., Posada-Villa, J., Gasquet, I., Kovess, V., Lepine, J. P., ... WHO World Mental Health Survey Consortium. (2004). Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *JAMA, 291(21), 2581–2590.*
<https://doi.org/10.1001/jama.291.21.2581>

- de Paula, A. L. D., Hallak, J. E. C., Maia-de-Oliveira, J. P., Bressan, R. A., and Machado-de-Sousa, J. P. (2015). Cognition in at-risk mental states for psychosis. *Neuroscience & Biobehavioral Reviews*, *57*, 199-208
- Dietrich, S., Beck, M., Bujantugs, B., Kenzine, D., Matschinger, H., and Angermeyer, M. C. (2004). The relationship between public causal beliefs and social distance toward mentally ill people. *Australian & New Zealand Journal of Psychiatry*, *38*(5), 348–354. <https://doi.org/10.1080/j.1440-1614.2004.01363.x>
- Durand-Zaleski, I., Scott, J., Rouillon, F., and Leboyer, M. (2012). A first national survey of knowledge, attitudes and behaviours towards schizophrenia, bipolar disorders and autism in France. *BMC Psychiatry*, *12*(1), 128.
- Eisenberg, N., and Miller, P. A. (1987). The relation of empathy to prosocial and related behaviors. *Psychological bulletin*, *101*(1), 91–119.
- Erdfelder, E., Faul, F., and Buchner, A. (1996). GPOWER: A general power analysis program. *Behavior research methods, instruments, & computers*, *28*(1), 1–11. <https://doi.org/10.3758/BF03203630>
- Fung, K. M. T., Tsang, H. W. H., and Corrigan, P. W. (2008). Self-stigma of people with schizophrenia as predictor of their adherence to psychosocial treatment. *Psychiatric rehabilitation journal*, *32*(2), 95–104. <https://doi.org/10.2975/32.2.2008.95.104>
- Garety, P. A., Kuipers, E., Fowler, D., Freeman, D., and Bebbington, P. E. (2001). A cognitive model of the positive symptoms of psychosis. *Psychological Medicine*, *31*(2), 189-195.
- Gawęda, Ł., Prochwicz, K., and Cella, M. (2015). Cognitive biases mediate the relationship between temperament and character and psychotic-like experiences in healthy adults. *Psychiatry research*, *225*(1-2), 50-57.

Gawęda, Ł., Prochwicz, K., Krężolek, M., Kłosowska, J., Staszkiwicz, M., and Moritz, S.

(2018). Self-reported cognitive distortions in the psychosis continuum: A Polish 18-item version of the Davos Assessment of Cognitive Biases Scale (DACOBS-18). *Schizophrenia research, 192*, 317-326.

Giblin, S., Clare, L., Livingston, G., and Howard, R. (2004). Psychosocial correlates of late-onset psychosis: life experiences, cognitive schemas, and attitudes to ageing.

International Journal of geriatric psychiatry, 19(7), 611-623

Goffman, E. (1963). *Stigma: Notes on the management of spoiled identity*. New York: Simon & Schuster, Inc.

Goldstein, B., and Rosselli, F. (2003). Etiological paradigms of depression: The relationship between perceived causes, empowerment, treatment preferences, and stigma. *Journal of Mental Health, 12(6)*, 551–563. <https://doi.org/10.1080/09638230310001627919>

González-Torres, M. A., Oraa, R., Arístegui, M., Fernández-Rivas, A., and Guimon, J. (2007).

Stigma and discrimination towards people with schizophrenia and their family members. *Social psychiatry and psychiatric epidemiology, 42(1)*, 14–23.

<https://doi.org/10.1007/s00127-006-0126-3>

Graves, R. E., Cassisi, J. E., and Penn, D. L. (2005). Psychophysiological evaluation of stigma towards schizophrenia. *Schizophrenia research, 76(2–3)*, 317–327.

<https://doi.org/10.1016/j.schres.2005.02.003>

Howes, O. D., and Murray, R. M. (2014). Schizophrenia: an integrated sociodevelopmental-cognitive model. *The Lancet, 383(9929)*, 1677-1687.

Jorm, A. F., and Griffiths, K. M. (2008). The public's stigmatizing attitudes towards people with mental disorders: how important are biomedical conceptualizations? *Acta Psychiatrica Scandinavica, 118(4)*, 315–321. <https://doi.org/10.1111/j.1600->

<https://doi.org/10.1111/j.1600->

0447.2008.01251.x

- Khosravani, V., Mohammadzadeh, A., and Oskouyi, L. S. (2019). Early maladaptive schemas in patients with schizophrenia and non-patients with high and low schizotypal traits and their differences based on depression severity. *Comprehensive psychiatry*, *88*, 1-8
- Kvaale, E. P., Gottdiener, W. H., and Haslam, N. (2013)a. Biogenetic explanations and stigma: A meta-analytic review of associations among laypeople. *Social science & medicine*, *96*, 95–103. <https://doi.org/10.1016/J.SOCSCIMED.2013.07.017>
- Kvaale, E. P., Haslam, N., and Gottdiener, W. H. (2013)b. The ‘side effects’ of medicalization: A meta-analytic review of how biogenetic explanations affect stigma. *Clinical psychology review*, *33*(6), 782–794. <https://doi.org/10.1016/J.CPR.2013.06.002>
- Lambooy, B., and Saïas, T. (2013). Réduire la stigmatisation des personnes souffrant de troubles psychiques par une campagne de communication ? Une synthèse de la littérature. *Annales Médico-psychologiques*, *171*(2), 77–82. <https://doi.org/10.1016/J.AMP.2010.11.022>
- Lampropoulos, D., Fonte, D., and Apostolidis, T. (2018). La stigmatisation sociale des personnes vivant avec la schizophrénie : une revue systématique de la littérature. *L'Évolution Psychiatrique*. <https://doi.org/10.1016/j.evopsy.2018.09.002>
- Lampropoulos, D., Wolman, A., and Apostolidis, T. (2017). Analyzing the presentation and the stigma of schizophrenia in French newspapers. *Social psychiatry and psychiatric epidemiology*, *52*(12), 1541–1547. <https://doi.org/10.1007/s00127-017-1455-0>
- Lincoln, T. M., Arens, E., Berger, C., and Rief, W. (2008). Can antistigma campaigns be improved? A test of the impact of biogenetic Vs psychosocial causal explanations on implicit and explicit attitudes to schizophrenia. *Schizophrenia Bulletin*, *34*(5), 984–994. <https://doi.org/10.1093/schbul/sbm131>

- Link, B. G., Cullen, F. T., Frank, J., and Wozniak, J. F. (1987). The social rejection of former mental patients: Understanding why labels matter. *American journal of Sociology*, 92(6), 1461–1500. <https://doi.org/10.1086/228672>
- Link, B., Phelan, J. C., Bresnahan, M., Stueve, A., and Pescosolido, B. A. (1999). Public conceptions of mental illness: labels, causes, dangerousness, and social distance. *American journal of public health*, 89(9), 1328–1333. <https://doi.org/10.2105/AJPH.89.9.1328>
- Martin, J. K., Pescosolido, B. A., Olafsdottir, S., and Mcleod, J. D. (2007). The construction of fear: americans' preferences for social distance from children and adolescents with mental health problems. *Journal of Health and Social Behavior*, 48(1), 50–67. <https://doi.org/10.1177/002214650704800104>
- Martin, J. K., Pescosolido, B. A., and Tuch, S. A. (2000). Of fear and loathing: The role of “disturbing behavior,” labels, and causal attributions in shaping public attitudes toward people with mental illness. *Journal of health and social behavior*, 41(2), 208. <https://doi.org/10.2307/2676306>
- McLean, B. F., Mattiske, J. K., and Balzan, R. P. (2017). Association of the jumping to conclusions and evidence integration biases with delusions in psychosis: a detailed meta-analysis. *Schizophrenia Bulletin*, 43(2), 344-354
- Mehta, S., and Farina, A. (1997). Is being “sick” really better? Effect of the disease view of mental disorder on stigma. *Journal of Social and Clinical psychology*, 16(4), 405–419. <https://doi.org/10.1521/jscp.1997.16.4.405>
- Parle, S. (2012). How does stigma affect people with mental illness? *Nursing times*, 108(28), 12–14.
- Pescosolido, B. A., Martin, J. K., Long, J. S., Medina, T. R., Phelan, J. C., and Link, B. G. (2010).

- “A disease like any other”? A decade of change in public reactions to schizophrenia, depression, and alcohol dependence. *American Journal of Psychiatry*, 167(11), 1321–1330. <https://doi.org/10.1176/appi.ajp.2010.09121743>
- Phelan, J. (2005). Geneticization of deviant behavior and consequences for stigma: The case of mental illness. *Journal of health and social behavior*, 46(4), 307–322.
- Phelan, J. C., Yang, L. H., and Cruz-Rojas, R. (2006). Effects of attributing serious mental illnesses to genetic causes on orientations to treatment. *Psychiatric Services*, 57(3), 382–387. <https://doi.org/10.1176/appi.ps.57.3.382>
- Prochwicz, K., and Kłosowska, J. (2017). Attentional focus moderates the relationship between attention to threat bias and delusion-like experiences in healthy adults. *European Psychiatry*, 39, 27-32.
- Read, J., Haslam, N., Sayce, L., and Davies, E. (2006). Prejudice and schizophrenia: a review of the “mental illness is an illness like any other” approach. *Acta Psychiatrica Scandinavica*, 114(5), 303–318. <https://doi.org/10.1111/j.1600-0447.2006.00824.x>
- Read, J., and Harré, N. (2001). The role of biological and genetic causal beliefs in the stigmatisation of “mental patients.” *Journal of mental health*, 10(2), 223–235.
- Ross, R. M., McKay, R., Coltheart, M., and Langdon, R. (2015). Jumping to conclusions about the beads task? A meta-analysis of delusional ideation and data-gathering. *Schizophrenia bulletin*, 41(5), 1183-1191.
- Schnittker, J. (2008). An uncertain revolution: Why the rise of a genetic model of mental illness has not increased tolerance. *Social science & medicine*, 67(9), 1370–1381. <https://doi.org/10.1016/j.socscimed.2008.07.007>
- Schomerus, G., Matschinger, H., and Angermeyer, M. C. (2014). Causal beliefs of the public and social acceptance of persons with mental illness: a comparative analysis of

schizophrenia, depression and alcohol dependence. *Psychological medicine*, 44(02), 303–314. <https://doi.org/10.1017/S003329171300072X>

Schomerus, G., Schwahn, C., Holzinger, A., Corrigan, P. W., Grabe, H. J., Carta, M. G., and Angermeyer, M. C. (2012). Evolution of public attitudes about mental illness: a systematic review and meta-analysis. *Acta Psychiatrica Scandinavica*, 125(6), 440–452. <https://doi.org/10.1111/j.1600-0447.2012.01826.x>

Schomerus, G., Matschinger, H., and Angermeyer, M. C. (2013). Continuum beliefs and stigmatizing attitudes towards persons with schizophrenia, depression and alcohol dependence. *Psychiatry Research*, 209(3), 665–669. <https://doi.org/10.1016/j.psychres.2013.02.006c>

Schmidt, N. B., Joiner, T. E., Young, J. E., & Telch, M. J. (1995). The schema questionnaire: Investigation of psychometric properties and the hierarchical structure of a measure of maladaptive schemas. *Cognitive therapy and research*, 19(3), 295-321.

So, S. H. W., Tang, V., and Leung, P. W. L. (2015). Dimensions of delusions and attribution biases along the continuum of psychosis. *PloS one*, 10(12), e0144558

Sundag, J., Ascone, L., de Matos Marques, A., Moritz, S., and Lincoln, T. M. (2016). Elucidating the role of early maladaptive schemas for psychotic symptomatology. *Psychiatry Research*, 238, 53–59. <https://doi.org/10.1016/j.psychres.2016.02.008>

Van 't Veer, J. T. B., Kraan, H. F., Drosseart, S. H. C., and Modde, J. M. (2006). Determinants that shape public attitudes towards the mentally ill. *Social psychiatry and psychiatric epidemiology*, 41(4), 310–317. <https://doi.org/10.1007/s00127-005-0015-1>

Waters, F., Woodward, T., Allen, P., Aleman, A., and Sommer, I. (2010). Self-recognition deficits in schizophrenia patients with auditory hallucinations: a meta-analysis of the literature. *Schizophrenia bulletin*, 38(4), 741-750.

<https://doi.org/10.1093/schbul/sbq144>

Weiner, B., Perry, R., and Magnusson, J. (1988). An attributional analysis of reactions to stigmas. *Journal of personality and social psychology*, 55(5), 738.

Weiner, Bernard. (1995). *Judgments of responsibility : a foundation for a theory of social conduct*. Guilford Press.

World Health Organization. (2001). *The World Health Report 2001 - Mental health: new understanding, new hope*. World Health Organization.

Young, J. E., Klosko, J. S., and Weishaar, M. E. (2003). *Schema therapy : a practitioner's guide*. Guilford Press.

Zawadzki, J. A., Woodward, T. S., Sokolowski, H. M., Boon, H. S., Wong, A. H. C., and Menon, M. (2012). Cognitive factors associated with subclinical delusional ideation in the general population. *Psychiatry Research*, 197(3), 345-349.

Table 1. Characteristics emphasized in the biogenetic, cognitive distortion, and cognitive schema accounts of Schizophrenia in terms of control of etiology and stability, and the resulting predictions formulated in the present study.

		Biogenetic account	Cognitive Distortion account	Cognitive Schema account
Etiological account	Control of etiology of the disorder	Low	High	Low
	Stability of etiology of the disorder over time	High	Low	Low
Hypotheses from the present research regarding stigma		Increased	Increased	Decreased

Table 2. Mean (and SD) of positive attitude, empathic concern and social proximity regarding schizophrenia at T0 and T1, as a function of text condition.

Text Condition	Measure	T0	T1	Difference (T1-T0)
Cognitive Schema	<i>Empathic Concern</i>	54.45 (20.68)	58.38 (21.43)	3.93
	<i>Positive Attitude</i>	71.72 (14.44)	73.55 (13.87)	1.83

Effects of a cognitive schema account on the stigma of schizophrenia

	<i>Social Proximity</i>	46.41 (22.50)	48.74 (24.54)	2.33
Genetic	<i>Empathic</i>			-0.63
	<i>Concern</i>	60.99 (19.89)	60.36 (21.59)	
	<i>Positive Attitude</i>	73.06 (12.79)	74.02 (14.06)	0.96
	<i>Social Proximity</i>	49.93 (20.54)	51.27 (22.28)	1.34
Cognitive Distortion	<i>Empathic</i>			0.48
	<i>Concern</i>	54.11 (17.98)	54.59 (20.30)	
	<i>Positive Attitude</i>	71.05 (12.78)	71.49 (13.82)	0.44
	<i>Social Proximity</i>	45.50 (20.75)	45.74 (21.94)	0.24
Control	<i>Empathic</i>			-1.18
	<i>Concern</i>	52.61 (17.22)	51.43 (18.50)	
	<i>Positive Attitude</i>	73.76 (12.72)	74.73 (12.92)	0.97
	<i>Social Proximity</i>	50.29 (21.48)	50.36 (23.16)	0.07

Effects of a cognitive schema account on the stigma of schizophrenia




Figure 1. Differential scores (T1-T0) of positive attitude, empathic concern and social proximity regarding schizophrenia, as a function of text condition. The stars indicate a statistical difference between T0 and T1 (* $p < .05$; ** $p < .01$; *** $p < .001$). Error bars indicate standard error of the mean.

Appendix

A1. Short description of the daily life of an individual with schizophrenia

Please read carefully the text below. Later, we will ask you a few questions regarding the content of this text.

"Some people may be affected by a serious mental illness called schizophrenia. These people can go to university, work in the field of justice and live in a self-contained apartment. They do not always present symptoms, but when they do, they can sometimes hear voices, or think that the CIA is watching them. Aside from these specific symptoms of schizophrenia, they can also feel very anxious, which can cause them to miss work on some days. People with schizophrenia are often hospitalized and take medication for their illness."

Control condition

Please carefully read the article below. Later, we will ask you a few questions regarding the content of the article. This article was published in the August-28 edition of this year's "Journal of the American Academy of Quantum Physics". The "Journal of the American Academy of Quantum Physics" is a journal that covers the latest news about quantum physics. It is highly respected in its academic field.

"Fission is largely the result of electrostatic factors." This is the first conclusion of an article recently published in the Journal of the American Academy of Quantum Physics. The author of this article is Dr Morel Edouard from Paris-Sorbonne University, who is renowned worldwide for his expertise in quantum physics. Dr Morel and his collaborators have investigated the factors that may be involved in the appearance of antineutrons in the fission process. They concluded: "Taking into account all the data gathered over the past 30 years, we can conclude that fission is the phenomenon by which the nucleus of a heavy atom (nucleus that contains many nucleons, such as uranium and plutonium nuclei) is divided into several lighter nuclides, usually two nuclides. This reaction results in the emission of neutrons and antineutrons (usually two or three), and a very important release of energy." These results have been praised by many researchers working in quantum physics, more specifically on fission, who now base their work on the results obtained by Dr Morel and his collaborators."

Biogenetic condition

Please carefully read the article below. Later, we will ask you a few questions regarding the content of the article. This article was published in the August-28 edition of this year's "Journal of Molecular Biology and Genetics". The "Journal of Molecular Biology and Genetics" is a journal that covers the latest news about genetics and the improvement of the well-being of the individual. It is highly respected in its academic field.

"Schizophrenia is largely the result of genetic factors." This is the first conclusion of an article recently published in the "Journal of Molecular Biology and Genetics". The author of this article is Dr Morel Edouard of Paris-Sorbonne University, who is renowned worldwide for his expertise in schizophrenia. Dr Morel and his collaborators have investigated the factors that may be involved in the onset of symptoms of schizophrenia. They concluded: "Taking into account all the data gathered over the past 30 years, we can conclude that genetic factors, linked to an excessive amount of neurotransmitters in the brain, a reduction of the volume of the grey matter, and abnormalities in some cortical circuits (all genetically predisposed), are the main causes of schizophrenia. For example, an individual with a parent experiencing schizophrenia has 30% risk of developing schizophrenia symptoms in their life." These results have been praised by many researchers working on schizophrenia, who now base their work on the results obtained by Dr Morel and his collaborators.

Cognitive distortion condition

Please carefully read the article below. Later, we will ask you a few questions regarding the content of the article. This article was published in the August-28 edition of this year's "Journal of Behavioral and Cognitive Therapy". The "Journal of Behavioral and Cognitive Therapy" is a journal that covers the latest news about psychopathology and cognitive and behavioral therapy. It is highly respected in its academic field.

"Schizophrenia is largely the result of cognitive factors." This is the first conclusion of an article recently published in the "Journal of Behavioral and Cognitive Therapy". The author of this article is Dr Morel Edouard of Paris-Sorbonne University, who is renowned worldwide for its expertise in schizophrenia. Dr Morel and collaborators have investigated factors that may be involved in the onset of symptoms of schizophrenia. They concluded: "Considering all the data gathered over the past 30 years, we can conclude that reasoning errors largely influence the onset of schizophrenia symptoms. One example of reasoning error may be that if someone blames you, it means that everyone blames you, which is called over-generalization. When this type of thoughts develops over time and multiplies, the symptoms of schizophrenia may appear." These results have been praised by many researchers working on schizophrenia, who now base their work on the results obtained by Dr Morel and his collaborators.

Cognitive Schema Condition

Please carefully read the article below. Later, we will ask you a few questions regarding the content of the article. This article was published in the August-28 edition of this year's "Journal of Behavioral and Cognitive Therapy". The "Journal of Behavioral and Cognitive Therapy" is a journal that covers the latest news about psychopathology and cognitive and behavioral therapy. It is highly respected in its academic field.

"Schizophrenia is largely the result of cognitive factors." This is the first conclusion of an article recently published in the "Journal of Behavioral and Cognitive Therapy". The author of this article is Dr Morel Edouard of Paris-Sorbonne University, who is renowned worldwide for its expertise in schizophrenia. Dr Morel and collaborators have investigated factors that may be involved in the onset of symptoms of schizophrenia. They concluded: "Considering all the data

gathered over the past 30 years, we can conclude that dysfunctional cognitive schemas, in other words errors of interpretation, which have their origin in events experienced during childhood, are the primary factors influencing the symptoms of schizophrenia. We can take the example of a child who is humiliated by his schoolmates. When this type of behavior is repeated in adulthood, the dysfunctional schema of mistrust built during childhood can then activate and generate very strong negative thoughts and emotions favoring the emergence of erroneous ideas that take the form of delusions. These results have been praised by many researchers working on schizophrenia, who now base their work on the results obtained by Dr Morel and his collaborators.