



Tropospheric Ozone Alters the Chemical Signal Emitted by an Emblematic Plant of the Mediterranean Region: The True Lavender (*Lavandula angustifolia* Mill.)

Candice Dubuisson, Florence Nicolè, Bruno Buatois, Martine Hossaert-Mckey, Magali Proffit

► To cite this version:

Candice Dubuisson, Florence Nicolè, Bruno Buatois, Martine Hossaert-Mckey, Magali Proffit. Tropospheric Ozone Alters the Chemical Signal Emitted by an Emblematic Plant of the Mediterranean Region: The True Lavender (*Lavandula angustifolia* Mill.). *Frontiers in Ecology and Evolution*, 2022, 10, 10.3389/fevo.2022.795588 . hal-03871592

HAL Id: hal-03871592

<https://hal.science/hal-03871592>

Submitted on 25 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tropospheric Ozone Alters the Chemical Signal Emitted by an Emblematic Plant of the Mediterranean Region: The True Lavender (*Lavandula angustifolia* Mill.)

Candice Dubuisson^{1*}, Florence Nicolè², Bruno Buatois¹, Martine Hossaert-McKey¹ and Magali Proffit¹

¹ Centre d'Écologie Fonctionnelle et Évolutive (CEFE), Université de Montpellier, CNRS, EPHE, IRD, Montpellier, France,

² Université de Lyon, UJM-Saint-Etienne, CNRS, LBVpam UMR 5079, Saint-Etienne, France

OPEN ACCESS

Edited by:

Robbie Girling,
University of Reading,
United Kingdom

Reviewed by:

Leif Abrell,
University of Arizona, United States
Tao Li,
Sichuan University, China

*Correspondence:

Candice Dubuisson
candice.dubuisson@cefe.cnrs.fr

Specialty section:

This article was submitted to
Chemical Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 15 October 2021

Accepted: 15 February 2022

Published: 15 March 2022

Citation:

Dubuisson C, Nicolè F, Buatois B, Hossaert-McKey M and Proffit M (2022) Tropospheric Ozone Alters the Chemical Signal Emitted by an Emblematic Plant of the Mediterranean Region: The True Lavender (*Lavandula angustifolia* Mill.). *Front. Ecol. Evol.* 10:795588. doi: 10.3389/fevo.2022.795588

Among air pollutants, tropospheric ozone (O_3) is one of the most stressful for organisms due to its strong oxidative potential. For instance, high ozone concentration ($[O_3]$) has the potential to affect (i) the emission of volatile organic compounds (VOCs) by plants and (ii) the lifetime of these VOCs in the atmosphere, and consequently disturb crucial signals in the interactions between plants and other organisms. However, despite the determinant role of VOCs emitted by flowers for pollinator attraction, a very limited number of studies have investigated the impact of O_3 on floral VOCs. In this study, we investigated the effect of high $[O_3]$ episodes on the VOCs emitted by a flowering Mediterranean plant: the true lavender (*Lavandula angustifolia* Mill., Lamiaceae). To do so, in controlled conditions, we exposed (i) the entire plant to high but realistic $[O_3]$ (200 ppb for 5 h) and (ii) only the VOCs emitted by lavender to increasing $[O_3]$ (0, 40, 80, 120, and 200 ppb). We sampled VOCs of lavender in both conditions and analyzed them by Gas Chromatography-Mass Spectrometry in order to qualify and quantify the flowering lavender's emissions and the reaction of VOCs with O_3 in the atmosphere. Our results showed that exposure to high $[O_3]$ during a short period (5 h) did not affect the emission of VOCs by flowering lavender. Incidentally, we also showed that the chemical signal varied in quantities and proportions over the day. Moreover, we showed that after their emission by the plant, composition of the VOCs changed quantitatively and qualitatively in an atmosphere containing $[O_3]$ naturally observed nowadays. Quantities of several of the major terpenes emitted by lavender decreased drastically during O_3 exposure, whereas concentrations of some VOCs increased, such as carbonyls and carboxylic acids, which are probably reaction products of terpenes with O_3 . Exposure to high $[O_3]$ thus directly affected the proportions of VOCs in the

atmosphere. Because pollinators generally use a blend of VOCs in particular proportions as a signal to localize flowers, the numerous pollinators of lavender may experience difficulty in recognizing specific floral odors during frequent and moderate $[O_3]$ episodes in the Mediterranean region.

Keywords: VOCs, flowers, ozone, pollution, lavender, Mediterranean ecosystems

INTRODUCTION

Since the beginning of the industrial era, anthropic activities have impacted ecosystems and their functioning (Barnosky et al., 2012). The main consequences are habitat fragmentation, climate change and the pollution of soils, water, and the atmosphere. Atmospheric pollution consists in increased concentrations of primary pollutants, such as carbon dioxide or nitrogen oxides (NO_x), released mainly by industry and agriculture. These pollutants are involved in complex chemical reactions in the troposphere that lead to the formation of secondary pollutants (Jenkin and Clemitshaw, 2000). Among these, tropospheric ozone (O_3) is one of the most important pollutants in terms of its contribution to global warming (Sicard et al., 2017; Yeung et al., 2019). Ozone is formed by the reaction between UV light and NO_x , the latter generally released by human activities in cities, but O_3 tends to accumulate in the countryside, because of complex processes involving wind, altitude and NO_x concentration. Thus, ozone concentration ($[O_3]$) varies in space and, as its formation is correlated with UV light, it also varies in time, with $[O_3]$ being generally highest during mid-day in summer (Cape, 2008; Sicard et al., 2017; Young et al., 2018). Currently, in many Northern-hemisphere countries, the mean daily $[O_3]$ reaches a minimum of 40 parts per billion (ppb) with numerous O_3 pollution episodes during summer. Between 1900 and 2000, the mean $[O_3]$ already increased by 30 ppb in the Northern hemisphere (Vingarzan, 2004; Ashmore, 2005) and some atmospheric chemistry models predict a further increase of about 5 ppb on average at the horizon of 2050 considering changes in climate with no control on the precursors source (Fowler et al., 2008). Due to its strong oxidative potential, O_3 is very stressful for organisms and causes damage to human health, notably respiratory problems (Gryparis et al., 2004; Nuvolone et al., 2018). In plants, physiological injuries have been reported, such as leaf lesions (Long and Naidu, 2002), but also decreased vegetative growth and reproductive output (Black et al., 2007), or the inhibition of activities of PEPc and Rubisco carboxylases and of photosynthetic pigments (chlorophylls a and b) (Leitao et al., 2007). As a consequence, European directives (Directive 2008/50/EC, 2008 of the European parliament and of the council of 21 May 2008 on ambient air quality and cleaner air for Europe) have set an alert threshold of 120 ppb of exposure to O_3 over 1 h for human health, beyond which emergency sanitary measures must be taken. For plants, the target value is set to $18,000 \mu g m^{-3} h^{-1}$ (900 ppb) of AOT40 (Accumulated Ozone over a Threshold of 40 ppb between 8:00 and 20:00 from April to August) in order to prevent physical and physiological injuries. Surprisingly, these thresholds do not take into account the effect of O_3 on other characteristics of plants, such as the emission of volatile organic

compounds (VOCs). However, VOCs are the foundation of plants' chemical communication with their environment. For instance, plants produce them to communicate with other plants (Ueda et al., 2012), to protect themselves against pathogens, to repel herbivores and also to attract pollinators, which are crucial for plant reproduction (Dudareva et al., 2013; Abbas et al., 2017). Several studies have investigated the effect of O_3 on the emission of VOCs by vegetative parts of plants (see reviews by Yuan et al., 2009; Holopainen and Gershenzon, 2010; Loreto and Schnitzler, 2010). These studies hypothesize that O_3 exposure induces emission of terpenes by leaves (to an extent varying among species) because their high antioxidant potential could protect the plants from oxidative stress (Peñuelas and Staudt, 2010; Pinto et al., 2010; Bourtsoukidis et al., 2012). In addition, studies have been conducted on the effect of O_3 on the lifetime of VOCs in the atmosphere (Atkinson and Arey, 2003a,b) and on the attraction of associated herbivorous insects (Fuentes et al., 2013; Li et al., 2016). Compared to plant-herbivore interactions, studies of the effect of O_3 on signal recognition in plant-pollinator interactions are limited.

Only in the last 10 years have investigations been conducted on the impact of O_3 on the emission of floral VOCs and on their lifetime in the atmosphere. However, flowers are well-known to emit VOCs to attract pollinators and ensure the plant's reproduction, and their VOCs play a major role in structuring plant-pollinator interactions (Raguso, 2008b; Dudareva et al., 2013; Burkle and Runyon, 2019; Kantsa et al., 2019). In fact, as for the vegetative parts of the plant, exposure to increased O_3 levels could modify the emission of some floral VOCs (Saunier and Blande, 2019) and/or reduce the lifetime of VOCs in the atmosphere, including those attractive to pollinators (McFrederick et al., 2009; Farré-Armengol et al., 2016). Because reaction time of the different VOCs with O_3 is extremely variable (Atkinson and Arey, 2003a), during periods with elevated O_3 levels significant changes in the proportions of VOCs are expected as soon as VOCs are emitted in the atmosphere. As the proportions of VOCs are crucial for the specific recognition of the floral chemical signal by pollinators (Raguso, 2008a; Proffit et al., 2020), these changes can reduce pollinator attraction to their host plants (Farré-Armengol et al., 2016). McFrederick et al. (2008) and Fuentes et al. (2016) showed by modeling under different scenarios of O_3 pollution that the scent plume downwind of floral sources could be degraded and as a result increase honeybee foraging time. More recently, an experimental study conducted in the field reported a reduction in the number of pollinating insects of several families and their rate of flower visits in an environment enriched in O_3 ($90 \text{ ppb} < [O_3] < 120 \text{ ppb}$) (Ryalls et al., 2022). Authors of this study concluded that these changes were caused by the reactions of floral VOCs with O_3 resulting in

an alteration of the pollinators' perceptions of the chemical signal emitted by the plants.

In the Mediterranean region, background $[O_3]$ is high and O_3 episodes, i.e., short periods with high $[O_3]$, are frequent in summer due to high levels of both sunlight and human activities. For instance, in summer 2003 in the French Mediterranean region, a peak of 200 ppb O_3 was registered (Vautard et al., 2005). Several Mediterranean flowering plants bloom during this same period. For instance, the true lavender (*Lavandula angustifolia* Mill.) is omnipresent in Mediterranean landscapes and flowers from the beginning of June to the end of July, depending on the altitude. It is a nectariferous plant that attracts a large number of pollinators and associated insects (e.g., predators) (Benachour, 2017). The censuses that we carried out on various fields of cultivated lavender show up to 140 different species of arthropods, including numerous species of bumblebees, solitary bees, butterflies, honeybees and Diptera (Nicolè, pers. comm.) (Li et al., 2019). It grows as a wild species but is also widely cultivated for its essential oil and for the production of honey by beekeepers (Barbier, 1963), creating a real olfactory landscape in Provence (McGregor, 1976). Moreover, behavioral experiments in controlled conditions, showed that essential oil extracts of *L. angustifolia* can act as plant signals, for example by significantly attracting honeybees (*Apis mellifera*) or repelling other insects (Li et al., 2019; Ganassi et al., 2020). The VOCs emitted by *L. angustifolia* inflorescences are well-known and include 50–90 molecules, mostly terpenes, dominated by linalyl acetate, linalool, 1,8-cineole, camphor, and borneol (Li et al., 2019; Stierlin et al., 2020; Héral et al., 2021). The reaction constants of some of these VOCs with O_3 have already been described, e.g., linalool 1,8-cineole or camphor (Atkinson and Arey, 2003b). The results of these studies strongly suggest that during O_3 episodes, the VOCs emitted from flowering lavender should react at different speed with O_3 , reducing differently their concentration, thereby affecting their relative proportions.

In this study we aimed to test to what extent exposure to high but realistic O_3 levels can alter (i) lavender emissions of VOCs and (ii) the concentration and proportions of lavender VOCs once they are emitted in the atmosphere. To do so, in order to simulate a high $[O_3]$ episode in controlled conditions, we exposed flowering lavenders to 200 ppb for 5 h, collected VOCs surrounding the plant by dynamic headspace, and analyzed them by Gas Chromatography-Mass Spectrometry (GC-MS). In order to characterize the effect of O_3 on the VOCs themselves, we isolated lavender VOCs in a reactor, exposed them to different $[O_3]$ (0, 40, 80, 120, and 200 ppb) for 30 min, collected the blend of VOCs in the reactor and analyzed them by GC-MS.

MATERIALS AND METHODS

Study System


The true lavender, *Lavandula angustifolia* Miller (Lamiaceae), is a plant from the Mediterranean region (Upson et al., 2004; Passalacqua et al., 2017). Here we used clones of *L. angustifolia* cv. "Diva" from the Centre Régionalisé Interprofessionnel d'Expérimentation en Plantes à Parfum, Aromatiques et

Médicinales (CRIEPPAM, Manosque, France). One-year-old plants were grown outdoors in individual pots in an NPK (nitrogen, phosphorus, potassium) 15 :15 :15 soil, from July 2018 at the Terrain d'Expériences, a technical platform of the Laboratoire d'Expertises Centre Méditerranéen Environnement et Biodiversité (LabEx CeMEB) located at the Centre d'Écologie Fonctionnelle et Évolutive (CEFE, Montpellier, France). In June 2020, we exposed 3-year-old plants to high O_3 levels (experiment 1) and in June 2021, we used 4-year-old plants to expose only the VOCs to different O_3 levels (experiment 2). The emissions of controlled plants were compared and proved to be similar between 3- and 4-year-old plants. In each experiment we used plants at the same flowering stage, when at least half of the flowers in the inflorescences were open.

Experiment 1: Volatile Organic Compounds Sampling of Flowering Lavender Exposed to High O_3 Level

Lavender plants were exposed to O_3 in greenhouses at the Terrain d'Expériences in which the temperature was controlled and fixed at 25°C in the day and 15°C during the night under natural light conditions (15 h/9 h day/night periods).

In order to expose the entire flowering plant to O_3 and monitor the VOCs, a dynamic enclosure system was designed. Plants were individually enclosed in a 42 L inert and odorless PTFE (polytetrafluoroethylene) frame closed by a 50 μ m thick PTFE film (Figure 1A). Air was pushed using a PTFE pump (KNF, N816.1.2K.18®, Germany) through a charcoal filter into the chambers at 6 L min⁻¹. A PTFE fan ensured air mixing in the chambers. Air flow rates were controlled by flowmeters and all tubing lines were made of PTFE. Plants were bagged at least 12 h before the treatment phase to allow plants to acclimate to the enclosure. Exposure was always conducted on pairs of plants: the first plant was exposed to 0 ppb O_3 and represented the control and the other plant was maintained at 200 ppb O_3 using an O_3 generator-analyzer (UV photometric Ozone Analyser with a generator option, Model 49i, Thermo Fisher Scientific, Franklin, MA, United States) for 5 h between 10:00 and 15:00, corresponding to the time of greatest pollinator activity in the fields (Nicolè, pers. comm.). For both plants, VOCs contained in the chambers were collected four times: (t0) before starting to ozonate the treated plant (0 ppb of O_3 in both chambers); (t1) 3 h after the beginning of generation of O_3 for the ozonated plant; (t2) 5 h after the beginning of generation of O_3 for the ozonated plant; and (t3) between 30 min and 1 h after O_3 generation was stopped (Figure 1B). VOCs were collected by pushing the air out of the chambers at 200 mL min⁻¹ through an adsorbent steel cartridge acting as VOCs trap filled with 40 mg of Carbotrap and 80 mg of Tenax phases (Sigma-Aldrich, Munich, Germany). We collected VOCs of each plant of the pair for 10 min following the same protocol. In addition, before introducing the plant into the chambers, to assess background scent level in the chambers, we collected, using the same method, blank samples, i.e., samples of air within empty chambers. Indeed, in semi-controlled experiments with whole plants, despite the fact that the air injected into the sampling chambers is purified with



activated charcoal, exogenous compounds, which are considered as contaminants, are always present in the samples, and therefore must be removed from the plant samples (see methods in Palmqvist et al., 2021; Rupp et al., 2021). At the end of each O₃ exposure in the first experiment, inflorescences and leaves were cut from the plant and placed for 4 days at 40°C in an oven to measure the dry biomass. Ten pairs of plants were used, representing 20 individuals in total for this first experiment.

Experiment 2: Sampling of Flowering Lavender Volatile Organic Compounds Exposed to Different [O₃]

In order to evaluate the effect of O₃ exposure on VOCs only, after their emission by the plant, we sequentially but randomly exposed lavender VOCs to 40, 80, 120, or 200 ppb of O₃ (Figure 2A). Based on the results of experiment 1, experiments were conducted between 11:00 and 18:00 because variations of VOC estimated concentrations and proportions over time were limited during this period (see results section “Experiment 1: Effects of Exposure to High O₃ Level on the Emission of VOCs by Flowering Lavender”). The same device described in experiment 1 was used to enclose the flowering lavender plant. This time, the set-up was installed in a room where the temperature (26.36 ± 0.041°C) and light were controlled (artificial light 400–700 nm SpectraBulb, LED E27, FloraLed, France) to better

characterize the effect of O₃ on lavender VOCs and avoid temperature- and light-related odor variations. Air purified by a charcoal filter was pushed into the chamber at 2 L min⁻¹. The air charged with VOCs present in the chamber was sent into a glass reactor of 2 L (Hei-Vap Core model, Heidolph Instruments GmbH & Co. KG, Regensburg, Germany) at a flow rate of 400 mL min⁻¹ using the same O₃ generator/analyzer as in the experiment 1 that analyzed at 1.4 L min⁻¹ the [O₃] at the outlet of the reactor (Figure 2A). This very same device generated 40, 80, 120, and 200 ppb at a flow rate of 1.8 L min⁻¹ at the inlet of the reactor. Respectively, 20, 50, 80, and 110 ppb were measured at the outlet of the reactor by a second O₃ analyser (Model 106-L Ozone monitor, 2B Technologies, Boulder, CO, United States) at 850 mL min⁻¹ (Figure 2A). The difference in [O₃] between the inlet and the outlet of the reactor confirmed that O₃ was reacting with VOCs in the reactor (loss of ~50% of the [O₃], but a loss of only ~25% when only O₃ was present in the reactor).

Each O₃ phase lasted 30 min and was always preceded by a control exposure phase of 30 min (0 ppb of O₃ in the reactor, Figure 2B). Plants were bagged at least 4 h before the treatment phase. VOCs in the reactor were collected for 10 min using the adsorbent traps containing Carbotrap/Tenax and analyzed by GC-MS for control (20 samples) and O₃ conditions (20 samples: N = 5 per [O₃] > 0 ppb). This experiment was conducted over 5 days with a different lavender plant every day (N = 5 plants for this second experiment). Each day, plant VOCs were exposed to




FIGURE 2 | Schematic diagram of **(A)** the experimental set-up presenting the exposure of flowering lavender volatile organic compounds (VOCs) to different $[O_3]$ and **(B)** timeline of the exposure for experiment 2. We present the temporal scheme showing the 4-hour control period at the beginning followed by alternating ozone periods (red) and control periods (purple). Each period was 30 min and VOC samples were collected through adsorbent cartridges (gray bars in panel **A**) at the end of each period during 10 min (including controls and ozone exposure).

the five $[O_3]$ and the order of the different concentrations used was randomized among the days.

Chemical Analyses

All chemical analyses were performed at the Plateforme d'Analyses Chimiques en Ecologie (PACE technical platform of the Labex CeMEB).

Analyses by GC-MS were conducted using the method detailed in Proffit et al. (2020). A gas chromatograph (GC, Trace 1310, Thermo Fisher Scientific, Milan, Italy) coupled to a mass spectrometer (ISQ QD Single Quadrupole, Thermo Fisher Scientific, Milan, Italy) was used with an Optima 5-MS capillary column (30 m, 0.25 mm ID, 0.25 μm film thickness, Machery-Nagel, Düren, Germany). Adsorbent traps were handled with a Multi Purpose Sampler (Gerstell, Mülheim, Germany) and desorbed with a double stage desorption system, composed of a Thermal Desorption Unit (TDU) and a Cold Injection System (CIS) (Gerstell, Mülheim, Germany). First, the injector was splitless with a temperature of 250°C on the CIS trap cooled at -80°C by liquid nitrogen. Then, the CIS trap was heated to 250°C with a 1:4 split ratio to inject the compounds into the column. We used helium at $1 \text{ mL} \cdot \text{min}^{-1}$ as a carrier gas. Oven temperature was held at 40°C for 3 min, increased from 40 to 220°C at a rate of $5^\circ\text{C} \cdot \text{min}^{-1}$ and from 220 to 250°C at $10^\circ\text{C} \cdot \text{min}^{-1}$, and finally held for 2 min at 250°C . The temperatures of the transfer line and the ion source of the mass

spectrometer were 250 and 200°C , respectively. The acquisition was at 70 eV ionization energy, from 38 to 350 m/z. XcaliburTM 266 software (Thermo Fisher Scientific, Milan, Italy) was used for data processing. Retention times of a series of n-alkanes (Alkanes standard solution, Sigma Aldrich, Munich, Germany) were used to convert retention times into retention indices. Compounds were then identified based on matching of the mass spectra using Total Ion Current mass chromatogram (TIC) with a database (NIST 2007 MS library, Wiley 9th edition), and on confirmation by comparison of calculated retention index (RI) with the Adams (2007) library and, for some VOCs, with matching of their mass spectra and retention time with those of synthetic compounds using the same analytic method. Volatiles detected only in the empty chamber or reactor (blank controls) were considered contaminants and were removed from the samples.

In order to evaluate the concentrations of the different VOCs, peak areas of VOCs were integrated using TIC and calibration factors were calculated with the same spectral method. To do so, we prepared a stock solution in dichloromethane (Pestisnorm, >99.8%) of four major VOCs of lavender, each of them at a mean concentration of $9.66 \pm 0.8 \mu\text{g}/\mu\text{L}$: linalyl acetate (Sigma-Aldrich, Analytical standard), camphor (Fluka, >99%), 1,8-cineole (Sigma-Aldrich, ~99%) and linalool (Fluka, >99%). In order to have calibration in the concentration range of the VOCs detected in lavender samples, dilutions of 1:1,000, 1:400, and 1:200 of this stock solution were prepared in

dichloromethane. For each of the solutions, 1 μL was injected in the adsorbent trap, with three replicates per solution, and analyzed by GC-MS using the same method as for the lavender samples. Calibration factors were obtained for each VOC using the slope of the regression between their peak area (integrated from TIC) and their concentration in the three replicates of the four dilutions, setting the intercept to zero. Because we do not have a reference standard for all VOCs of the lavender samples and we do not have clear expectations about which ones represent a signal for interacting species, we used the mean calibration factor of these four compounds to estimate the concentrations of all compounds present in our samples.

For experiment 1, estimated concentration (ng/min/g of dry biomass) of each VOC was calculated. For experiment 2, in order to evaluate the quantitative variation of the different VOCs after O_3 exposure compared to the corresponding control (0 ppb) we calculated a relative index of variation following the formula:

$$\text{Variation index of VOC}_{[\text{O}_3]i} = \frac{(\text{area of VOC}_{[\text{O}_3]i} + 1) - (\text{area of VOC}_{[\text{O}_3]0} + 1)}{(\text{area of VOC}_{[\text{O}_3]i} + 1) + (\text{area of VOC}_{[\text{O}_3]0} + 1)}$$

“area of $\text{VOC}_{[\text{O}_3]i}$ ” stands for the peak area of a given compound at a given $[\text{O}_3]$ i (i : from 0 to 200 ppb), “area of $\text{VOC}_{[\text{O}_3]0}$ ” stands for the peak area of a given compound in the sample of the corresponding control (0 ppb of O_3). The variation index varied between -1 and 1 .

Statistical Analyses

All the data visualization and analyses were performed in R (v. 3.6.2; R Development Core Team)¹ and required the following packages: *ade4* (Thioulouse et al., 2018), *car* (Fox and Weisberg, 2019), *vegan* (Oksanen et al., 2020), *mixOmics* (Rohart et al., 2017), *Hotelling* (Curran, 2018), *RVAideMemoire* (Hervé, 2021), *nlme* (Pinheiro et al., 2021), *contrast* (O’Callaghan et al., 2020), and *corplot* (Wei and Simko, 2021).

Experiment 1: Effects of Exposure to High O_3 Level on the Emission of Volatile Organic Compounds by Flowering Lavender

Estimated concentrations (ng/min/g of dry biomass) and relative proportions in the total bouquet of each compound were calculated and set into two distinct data tables.

To assess the effect of O_3 on the concentration of each compound, but also on the total estimated concentration of all VOCs, we fitted linear mixed models on their log-transformed quantity with treatment (control or ozone), time (t_0 , t_1 , t_2 , and t_3) and their interaction as fixed factors and the pair of plants as random factor. Analyses of deviance allowed testing the significance of effects of these factors on concentration of each VOC. Then, pairwise comparisons were made between modalities of the factors using contrasts.

If significant variations of VOC estimated concentrations were detected between the control and treated plants, we also evaluated the effect of O_3 on the total estimated concentration of VOCs

of lavender plants with a linear model with treatment, time and their interaction as fixed factors and the pair of plants as random factor. Data were log-transformed prior to the analysis to normalize the distribution of the residuals. An analysis of deviance was used in order to test the significance of effects of these factors on the plants’ total emission. Then, contrasts were conducted between modalities of the factors to reveal true differences.

In addition, variations in the relative proportions of VOCs in the overall odor bouquet were tested using partial redundancy analyses (pRDA) (Hervé et al., 2018). This multivariate analysis consisted of two steps. Firstly, a multivariate linear regression on the relative proportions of VOCs with treatment, time and their interaction, was performed with pair of plants as a random factor. Secondly, a constrained Principal Component Analysis (PCA) on the fitted values of the regression model was conducted in order to evaluate the variation in the relative proportions of VOCs that was explained by the two factors and their interaction. In addition, a permutation test was performed to assess the significance of the effects of the fixed factors. Finally, pairwise comparisons were conducted between the modalities of the factors and p -values were adjusted with the “false discovery rate” (fdr) method (Benjamini and Hochberg, 1995).

Experiment 2: Effects of Different $[\text{O}_3]$ on Flowering Lavender Volatile Organic Compounds

To assess the effect of $[\text{O}_3]$ (0, 40, 80, 120, and 200 ppb) on the variation index of each compound, we fitted linear models with $[\text{O}_3]$ as fixed factor and the plant as random factor. An analysis of deviance was conducted to test the significance of the effect of $[\text{O}_3]$ on the variation index. Then, pairwise comparisons were made between modalities of this factor using contrast.

In order to highlight possible relationships between the different VOCs in terms of degradation-formation with O_3 , we performed a correlation matrix between each pair of compounds based on Pearson’s correlation test with scaled peak areas of VOCs ozonated at all $[\text{O}_3]$.

The effect of $[\text{O}_3]$ on the relative proportions of VOCs in the reactor was tested using a pRDA, with plants as a random factor. A permutation test was performed to assess the global effect of $[\text{O}_3]$ but also pairwise comparisons between modalities of the factor; p -values were adjusted with the “fdr” method (Benjamini and Hochberg, 1995).

RESULTS

Experiment 1: Effects of Exposure to High O_3 Level on the Emission of Volatile Organic Compounds by Flowering Lavender

In the volatile emission of flowering lavender, we detected a total of 34 VOCs, seven of which were present only as traces (see Table 1 and Supplementary Figure 1). As expected,

¹<http://www.R-project.org>

TABLE 1 | Effects of the O₃ treatment and time on the estimated concentrations of the different VOCs detected in experiment 1. For each modality of the different factors, see **Figure 1**.

			Ozone treatment													
			Control N=10🌿				Ozone N=10👉				Results of the linear models					
Compounds	CalcRI	LitRI	t0	t1	t2	t3	t0	t1	t2	t3	Treatment		Time		Treatment * Time	
											df = 1		df = 3		df = 3	
											x²	p	x²	p	x²	p
Monoterpenes																
α-thujene	933	924	trace	trace	trace	trace	trace	trace	trace	trace						
α-pinene*	940	932	1.54 ± 0.50 ^a	0.55 ± 0.14 ^b	0.51 ± 0.10 ^b	0.63 ± 0.16 ^b	1.03 ± 0.11 ^a	0.47 ± 0.17 ^{bc}	0.13 ± 0.04 ^c	0.57 ± 0.15 ^b	3.77	0.05	34.21	<0.001	2.30	0.51
Camphene	946	946	0.67 ± 0.48	0.08 ± 0.08	0.32 ± 0.20	0.41 ± 0.23	0.05 ± 0.04	0.49 ± 0.49	0.06 ± 0.06	0.17 ± 0.07	1.82	0.18	0.78	0.85	3.18	0.37
Sabinene	979	969	trace	trace	trace	trace	trace	trace	trace	trace			trace			
Myrcene	995	988	0.62 ± 0.27	0.74 ± 0.40	1.08 ± 0.33	0.94 ± 0.41	0.98 ± 0.76	2.19 ± 1.94	0.33 ± 0.17	0.78 ± 0.30	0.50	0.48	0.39	0.94	1.67	0.64
p-cymene	1024	1020	0.12 ± 0.03 ^a	0.24 ± 0.06 ^{ab}	0.40 ± 0.09 ^b	0.49 ± 0.12 ^b	0.22 ± 0.12 ^a	0.27 ± 0.12 ^a	0.30 ± 0.13 ^a	0.37 ± 0.11 ^a	0.37	0.54	10.27	0.02	2.17	0.54
o-cymene	1027	1022	2.60 ± 1.70	0.87 ± 0.26	1.18 ± 0.30	1.41 ± 0.36	1.05 ± 0.42	1.04 ± 0.34	0.99 ± 0.36	1.16 ± 0.35	0.86	0.35	1.18	0.76	0.77	0.86
Limonene*	1035	1024	19.42 ± 15.29	2.32 ± 1.62	0.60 ± 0.39	0.34 ± 0.23	4.39 ± 2.16	1.84 ± 1.39	1.84 ± 0.99	1.51 ± 1.02	0.26	0.61	4.89	0.18	0.50	0.92
1,8-cineole*	1035	1026	0.26 ± 0.19	0.93 ± 0.54	1.58 ± 0.95	1.96 ± 1.01	0.05 ± 0.05	2.72 ± 2.72	0.56 ± 0.43	0.96 ± 0.53	2.56	0.11	7.01	0.07	0.71	0.87
(Z)-β-ocimene	1041	1032	0.18 ± 0.06 ^a	0.27 ± 0.11 ^a	0.71 ± 0.18 ^b	0.76 ± 0.11 ^b	0.41 ± 0.18 ^a	0.12 ± 0.12 ^{ac}	0.03 ± 0.03 ^c	0.65 ± 0.22 ^a	8.22	0.004	25.37	<0.001	16.92	<0.001
(E)-β-ocimene	1051	1044	0.32 ± 0.10 ^a	0.58 ± 0.26 ^{ac}	1.36 ± 0.39 ^b	1.33 ± 0.25 ^b	0.81 ± 0.35 ^a	0.19 ± 0.19 ^c	0.03 ± 0.03 ^c	1.78 ± 0.76 ^b	5.85	0.02	26.76	<0.001	18.97	<0.001
Linalool*	1106	1095	0.51 ± 0.14 ^a	0.41 ± 0.12 ^a	0.8 ± 0.19 ^a	0.81 ± 0.14 ^a	0.6 ± 0.18 ^a	0.46 ± 0.26 ^{ab}	0.09 ± 0.05 ^b	1.14 ± 0.43 ^a	2.35	0.13	12.56	0.006	9.72	0.02
Allo-ocimene	1137	1128	0.06 ± 0.04	0.15 ± 0.12	0.27 ± 0.12	0.16 ± 0.07	0.07 ± 0.05	0.25 ± 0.19	0.22 ± 0.17	0.09 ± 0.06	0.12	0.73	2.39	0.50	0.78	0.85
Camphor*	1151	1141	2.95 ± 2.63	0.98 ± 0.61	1.08 ± 0.91	1.54 ± 0.93	0.35 ± 0.18	0.19 ± 0.1	0.69 ± 0.62	0.86 ± 0.67	3.04	0.08	1.30	0.73	0.25	0.97
Borneol	1175	1165	0.91 ± 0.34	0.54 ± 0.12	0.59 ± 0.10	0.63 ± 0.12	0.69 ± 0.24	1.81 ± 1.23	0.61 ± 0.15	1.32 ± 0.55	1.03	0.31	1.28	0.73	2.57	0.46
Terpinen-4-ol	1185	1174	2.61 ± 2.49	0.08 ± 0.03	0.11 ± 0.04	0.12 ± 0.03	0.26 ± 0.15	0.08 ± 0.04	0.03 ± 0.02	0.17 ± 0.07	0.68	0.41	5.26	0.15	1.52	0.68
Linalyl acetate*	1260	1254	1.66 ± 0.43 ^a	2.70 ± 0.93 ^{ab}	3.38 ± 0.98 ^{ab}	3.29 ± 0.65 ^b	3.02 ± 1.52 ^a	1.41 ± 0.97 ^c	0.29 ± 0.18 ^c	3.77 ± 1.34 ^a	12.16	<0.001	15.56	0.001	21.07	<0.001
Lavandulyl acetate	1278	1288	trace	trace	trace	trace	trace	trace	trace	trace						
Bornyl acetate	1278	1287	trace	trace	trace	trace	trace	trace	trace	trace						
Sesquiterpenes																
α-santalene*	1433	1416	0.10 ± 0.03 ^a	0.18 ± 0.04 ^{ab}	0.25 ± 0.06 ^{ab}	0.32 ± 0.07 ^b	0.26 ± 0.10 ^a	0.00 ± 0.00 ^c	0.00 ± 0.00 ^c	0.30 ± 0.10 ^a	5.35	0.02	18.78	<0.001	17.59	<0.001
(E)-β-caryophyllene*	1435	1417	0.33 ± 0.07 ^a	1.09 ± 0.71 ^a	0.54 ± 0.15 ^a	0.64 ± 0.12 ^a	0.48 ± 0.16 ^a	0.00 ± 0.00 ^b	0.00 ± 0.00 ^b	0.44 ± 0.19 ^a	14.14	<0.001	4.92	0.18	11.61	<0.001
Aliphatics																
(Z)-hex-3-enyl acetate	1010	1004	0.01 ± 0.00	0.01 ± 0.00	0.03 ± 0.02	0.005 ± 0.001	0.14 ± 0.13	0.03 ± 0.02	0.01 ± 0.01	0.01 ± 0.00	1.27	0.26	2.04	0.56	3.14	0.37
2-ethylhexanol	1034	1035	3.25 ± 3.05	0.37 ± 0.25	0.07 ± 0.03	0.15 ± 0.11	0.68 ± 0.62	0.42 ± 0.30	0.68 ± 0.53	0.05 ± 0.04	0.008	0.93	2.93	0.40	1.92	0.59
Nonanal*	1108	1104	0.31 ± 0.23	0.06 ± 0.03	0.07 ± 0.03	0.08 ± 0.03	0.13 ± 0.23	0.05 ± 0.03	0.06 ± 0.03	0.08 ± 0.04	0.50	0.48	4.54	0.21	0.93	0.82
Oct-1-en-3-yl acetate	1115	1110	0.08 ± 0.04 ^a	0.33 ± 0.12 ^{ab}	0.48 ± 0.38 ^b	0.62 ± 0.17 ^b	0.28 ± 0.16 ^a	0.48 ± 0.38 ^a	0.32 ± 0.16 ^a	0.54 ± 0.21 ^a	0.17	0.68	9.49	0.02	2.53	0.47
Octanoic acid	1188	1167	trace	trace	trace	trace	trace	trace	trace	trace			trace			
Decanal*	1207	1203	0.004 ± 0.002	0.05 ± 0.03	0.004 ± 0.002	0.01 ± 0.003	0.01 ± 0.00	0.02 ± 0.01	0.02 ± 0.01	0.01 ± 0.00	0.006	0.94	6.73	0.08	3.11	0.38
Dodecanal*	1411	1408	0.66 ± 0.27 ^a	0.94 ± 0.16 ^a	0.53 ± 0.14 ^a	0.66 ± 0.04 ^a	0.68 ± 0.12 ^a	2.03 ± 0.68 ^b	1.13 ± 0.34 ^{bc}	0.74 ± 0.21 ^{ac}	5.45	0.02	13.47	0.004	3.86	0.28
Dodecanol*	1478	1469	0.91 ± 0.48	0.44 ± 0.07	0.4 ± 0.08	0.38 ± 0.08	0.27 ± 0.06	1.33 ± 0.87	0.38 ± 0.22	0.34 ± 0.08	0.21	0.64	3.72	0.29	5.31	0.15
Pentadecane*	1502	1500	0.07 ± 0.06	0.008 ± 0.008	0.04 ± 0.02	0.02 ± 0.01	0.03 ± 0.02	0.03 ± 0.02	0.09 ± 0.04	0.10 ± 0.04	1.66	0.50	2.99	0.18	3.83	0.95

(Continued)

TABLE 1 | (Continued)

Compounds	CalcRI	LitRI	Control N=10				Ozone treatment				Results of the linear models							
			t0	t1	t2	t3	t0	t1	t2	t3	Treatment		Time		Treatment * Time			
											df = 1	x ²	p	x ²	p	df = 3	x ²	p
Monoterpenes																		
Irregular terpenes																		
Geranyl acetone	1455	1453	0.30 ± 0.09	0.37 ± 0.11	0.38 ± 0.11	0.37 ± 0.12	0.22 ± 0.09	0.64 ± 0.37	0.39 ± 0.10	0.46 ± 0.10	0.07	0.78	2.29	0.52	0.82	0.84		
Benzenoids																		
Benzaldehyde*	964	952	trace	trace	trace	trace	trace	trace	trace	trace			trace					
Acetophenone	1072	1059	trace	trace	trace	trace	trace	trace	trace	trace			trace					
Other																		
Unknown compound	906		1.44 ± 1.40	0.55 ± 0.45	0.30 ± 0.16	0.19 ± 0.12	0.27 ± 0.22	0.28 ± 0.22	0.46 ± 0.39	0.24 ± 0.19	0.29	0.59	0.52	0.91	0.53	0.91		
Total			41.89 ± 23.39 ^a	15.85 ± 3.63 ^a	17.06 ± 3.53 ^a	18.26 ± 3.02 ^a	17.42 ± 5.55 ^{ab}	18.86 ± 7.91 ^{ab}	9.74 ± 2.40 ^b	18.6 ± 4.69 ^a	4.38	0.04	4.72	0.19	1.68	0.64		
We present the means ± standard error in ng/min/g of dry biomass per individual. In addition, results of the linear models are presented for each VOC. Bold numbers correspond to statistically significant <i>p</i> -values (<i>p</i> < 0.05) and modalities that are associated with different letters are significantly different. CalcRI means "Calculated Retention Index", LitRI means "Literature Retention Index" from Adams (2007); "trace" means that VOCs were present in the chromatogram but not quantifiable. The purple plant corresponds to lavenders that were not subject to O ₃ exposure and the red plant corresponds to lavenders that were exposed to 200 ppb O ₃ . Mean dry weight of lavender was 2.66 ± 0.71 g.																		
*Identification based on authentic standards (see Supplementary Figure 1).																		
VOCs collection time; df, degrees of freedom.																		

monoterpenes were the family of compounds most represented in terms of concentrations and number of compounds.


Volatile Organic Compounds Estimated Concentrations and Proportions in Control Conditions

The control conditions (0 ppb) provided information about the expected chemical profile of *L. angustifolia* in normal conditions without ozone exposure. For all control samples (*N* = 40), limonene was the major compound (control sample mean ± SE: 24.4 ± 15.49% of the total of VOCs emitted and 22.68 ± 6.88 ng/min/g of dry biomass), owing to its high estimated concentration at t0 (19.42 ± 15.29 ng/min/g of dry biomass), followed by linalyl acetate (11.9 ± 6.7% of the total VOCs emitted by control plants and 2.76 ± 0.58 ng/min/g of dry biomass), which dominated the odor bouquet of all the control samples except at t0. Camphor (7.05 ± 0.74%), o-cymene (6.53 ± 0.74%) and 1,8-cineole (5.1 ± 3.38%) were the other most abundant compounds, each of them being present at around 1.45 ± 0.18 ng/min/g of dry biomass on average (Table 1). Each of the other VOCs were present at less than 6% on average of the total blend.

Effects of Ozone Treatment and Time of Day on Volatile Organic Compounds Estimated Concentrations

Both O₃ treatment and time of day affected the estimated concentrations of emitted VOCs. Of the 34 VOCs in the chamber, concentrations of 10 VOCs were significantly affected by one or both of these factors (linear models, at least one significant *p*-value, i.e., *p* < 0.02, per VOC, Table 1). Significant decreases in the estimated concentrations of monoterpenes [α -pinene, (Z) and (E)- β -ocimene, linalool, linalyl acetate, and p-cymene] and of two sesquiterpenes [α -santalene and (E)- β -caryophyllene] were observed when the plants were exposed to 200 ppb O₃ for 3 h or 5 h (*p* ≤ 0.02) (Table 1). The concentration of only one compound, dodecanal, increased significantly when plants were exposed to 200 ppb O₃ for 3 h ("ozone:t1") compared to the corresponding control ("control:t1") (*p* = 0.02). There were no significant differences between the estimated concentrations of different VOCs of lavender plants that after a 5 h exposure were no longer ozonated ("ozone:t3"), and the corresponding control plants ("control:t3").

Incidentally, this experiment also showed, apparently for the first time, that emission of VOCs by flowering lavender varied over the course of the day. Concentration of α -pinene decreased over the day, while estimated concentrations of p-cymene, 1,8-cineole, (Z) and (E)- β -ocimene, linalyl acetate, oct-1-en-3-yl acetate and α -santalene increased (Table 1). The total concentration of VOCs emitted by flowering lavender was affected only by the O₃ treatment (*p* = 0.04) and not by the time of the day or the interaction of these two factors (*p* = 0.19 and *p* = 0.64, respectively, Table 1). Pairwise comparisons revealed only one significant difference: the total VOCs concentration of lavenders ozonated for 5 h ("ozone:t2") was significantly lower than in the corresponding control ("control:t2").



Effects of Ozone Treatment and Time of Day on the Relative Proportions of Volatile Organic Compounds

Because estimated concentrations of different VOCs were affected both by O_3 exposure and by time of day, their relative proportions also changed during O_3 exposure. The O_3 treatment (0 vs. 200 ppb ozone), time of day (t0, t1, t2, and t3) and their interaction together explained 20% of this variability in relative proportions of VOCs, as shown by the multivariate linear regression analysis (Figure 3). The random factor “pair of plants” explained 4.9% of the variation. The relative proportions of VOCs varied significantly between O_3 treatments (permutation test: $F_{1, 71} = 4.53$, $p = 0.001$) and among times of day (permutation test: $F_{3, 71} = 3.04$, $p = 0.001$), and the interaction of the two factors was also significant (permutation test: $F_{3, 71} = 1.74$, $p = 0.008$). Regarding diurnal variation, pairwise comparisons detected significant differences in relative proportions of VOCs emitted by lavender plants from the morning to the afternoon (from t0 to t3) between the control groups (Figure 3C). Moreover, the relative proportions of VOCs in chambers with plants exposed to O_3 for 3 h (“ozone:t1”) and 5 h (“ozone:t2”) were significantly different from those in all other groups (respectively $p < 0.012$ and $p < 0.002$). The pRDA score plot from the constrained PCA (Figure 3A) shows the significant effects of ozone treatment,

time of day and their interaction on the relative proportions of VOCs. A total of 82% of the constrained variance is explained by the first and the second components of the PCA (67.50 and 14.44%, respectively). As previously shown by the pairwise comparisons, three groups appear on this PCA. First, the effect of time of day can be observed on the second component, where samples collected in the morning and mid-day (t0 and t1) were completely separated from those collected in the afternoon (t2 and t3). Second, the effect of the O_3 treatment is clearly observed in the first component. Samples collected when lavender plants were ozonated for 3 and 5 h (“ozone:t1” and “ozone:t2”) were positively correlated with the first component, in contrast to those collected in control chambers and with plants that were no longer ozonated (“ozone:t3”), which were negatively correlated with the first component.

The correlation circle of the constrained PCA (Figure 3B) highlights the VOCs correlated individually with the first or the second component at the threshold of 30% and shows which compounds were affected by the different experimental conditions. Linalool, (Z)- and (E)- β -ocimene, (E)- β -caryophyllene, α -santalene, and linalyl acetate were associated with control plants and with previously exposed plants (“ozone:t3”). In contrast, dodecanol, p-cymene, decanal,




FIGURE 4 | Relative proportions of volatile organic compounds (VOCs) emitted by flowering lavenders in experiment 1 exposed to 0 ppb O_3 for 5 h (right) vs. proportions of VOCs emitted by flowering lavenders exposed to 200 ppb O_3 for 5 h (left). Mean $\pm$ standard error per individual of the relative proportions of VOCs emitted by 10 plants without O_3 in the chamber and by 10 plants exposed to 200 ppb O_3 for 5 h.

dodecanal, pentadecane, an unknown compound and 2-ethylhexanol were positively correlated with the first component, suggesting that their proportions were higher in plants that sustained exposure to 200 ppb of O_3 for 3 and 5 h (“ozone:t1” and “ozone:t2”). In fact, compared to control plants, in plants ozonated at 200 ppb for 5 h (“ozone:t2”), linalyl acetate was no longer the major compound in the scent bouquet, being replaced by dodecanal and limonene (**Figure 4**). Regarding diurnal variation, geranyl acetone, 1,8-cineole and oct-1-en-3-yl acetate were positively correlated with the second component, indicating that their proportions were higher in odor bouquets of plants sampled in the afternoon, whereas α -pinene was present in higher proportion in odor bouquets of plants sampled in the morning.

Experiment 2: Effects of Different $[O_3]$ on Flowering Lavender Volatile Organic Compounds

In this experiment, plants were not ozonated. Only the VOCs, after their emission by the plant, were exposed to O_3 . We detected 35 VOCs in the reactor chamber, of which 23 could be quantified and 12 were present only as traces.

Volatile Organic Compounds Proportions in Control Conditions

In control conditions (0 ppb in the reactor), among the VOCs detected in the reactor chamber, terpenes were still the most represented family (43.94% of the total blend), with linalyl acetate being the main monoterpene, comprising $21.04 \pm 15.22\%$ of the total blend (**Supplementary Table 1**).

Effects of $[O_3]$ on the Quantities of Volatile Organic Compounds in the Reactor

The $[O_3]$ in the reactor significantly affected the variation index of 16 VOCs among the 23 VOCs quantified in the chemical blend (**Figure 5**). The remaining seven VOCs did not vary significantly with increasing $[O_3]$: borneol ($x^2 = 4.92$, $p = 0.3$), camphene ($x^2 = 2.48$, $p = 0.65$), camphor ($x^2 = 1$, $p = 0.91$), α -cymene ($x^2 = 5.88$, $p = 0.21$), p-cymene ($x^2 = 8.41$, $p = 0.08$), an unknown compound ($x^2 = 4.8$, $p = 0.31$) and undecanal ($x^2 = 6.09$, $p = 0.19$). Linear models revealed a significant effect of the $[O_3]$ on the variation index of VOCs. Pairwise comparisons highlighted once again significant decreases in the quantities of monoterpenes [(Z)- and (E)- β -ocimene, linalool, linalyl acetate, limonene, myrcene, α -pinene, 1,8-cineole, and bornyl acetate], of one aliphatic compound (1-octen-3-ol acetate) and of some sesquiterpenes [(E)- β -caryophyllene, α -santalene, and caryophyllene oxide] when VOCs were ozonated (**Figure 5A**), while estimated concentrations of dodecanal and decanal increased significantly (**Figure 5B**).

Quantities of decanal, dodecanal, the unknown compound and geranyl acetone were significantly negatively correlated with quantities of some monoterpenes and sesquiterpenes of the bouquets [mostly linalyl acetate, bornyl acetate, borneol, 1-octen-3-ol acetate, (Z) and (E)- β -ocimene] (**Figure 6**; $r < 0.4$, $p < 0.05$). These results confirm the assumption that different VOCs emitted by plants react differentially with O_3 and may form new VOCs in the odor bouquet.

Effects of $[O_3]$ on the Proportions of Volatile Organic Compounds in the Reactor

The $[O_3]$ in the reactor had a significant effect on variation of the relative proportions of VOCs (permutation test: $F_{4, 16} = 5.03$,




FIGURE 5 | Effects of exposure to increasing $[O_3]$ in experiment 2 on the volatile organic compounds (VOCs) variation index (mean $\pm$ standard error). Panel (A) lists VOCs whose variation index significantly decreased with $[O_3]$, (B) panel lists VOCs whose variation index increased significantly with increasing $[O_3]$. Purple, yellow, orange, red, and brown colors correspond, respectively to an $[O_3]$ exposure of 0, 40, 80, 120, and 200 ppb. For each VOC, results of the linear models are presented and values of the variation index with different letters indicate statistically significant difference in pairwise comparisons ($p < 0.05$).

$p = 0.001$). Twenty-nine percent of the total variation of relative proportions of VOCs was explained by the $[O_3]$ (0, 40, 80, 120, and 200 ppb) in the redundancy analysis. Forty-eight percent of this variation was explained by the random factor (plants). Pairwise comparisons showed that the relative proportions of VOCs were significantly different between the non-ozonated VOCs (0 ppb) and all the ozonated VOCs, irrespective of the $[O_3]$ applied ($p < 0.04$). Among the ozonated samples, the proportions of different VOCs were not significantly different among treatments ($p > 0.08$), except between the 40 and 200 ppb treatments ($p = 0.04$).

The significant effect of $[O_3]$ on the relative proportions of VOCs in the reactor is clearly illustrated in the pRDA score plot from the constrained PCA (Figure 7A). A total of 96% of the constrained variance was explained by the first and second components of the PCA (92 and 4%, respectively). The effect of $[O_3]$ can be observed on the first component, where ozonated samples were positively correlated with this component, in contrast to non-ozonated controls, which were negatively correlated. On the correlation circle of the constrained PCA, camphor, decanal, dodecanal, o-cymene, p-cymene, camphene, borneol, 1,8-cineole, geranyl acetone, α -pinene, and unknown compound were positively correlated with the first component (Figure 7B): their relative proportions were higher in the ozonated odor bouquets than in control ones (Supplementary Table 1). On the contrary, (Z)- and (E)- β -ocimene, α -santalene, (E)- β -caryophyllene, myrcene, linalool, and caryophyllene oxide were negatively correlated with the first component; their

proportions were higher in the control odor bouquet than in ozonated ones (Supplementary Table 1). The second component of the pRDA clearly separates the chemical profiles as a function of the value of increasing $[O_3]$. There was a gradient from VOCs exposed to 40 ppb with a positive score on the axis to VOCs exposed to 200 ppb with a negative score on the axis. Undecanal and 1-octen-3-ol acetate were negatively correlated with the second component and were associated with samples ozonated at high $[O_3]$ (at 80 ppb and higher values) (Figure 7B).

DISCUSSION

This study expands our knowledge of VOCs emission in *Lavandula angustifolia* and also provides important insights on the impact of high $[O_3]$ episodes on chemical signals emitted by flowers.

In this study, we created a new experimental device allowing plants to be exposed to different $[O_3]$ during several hours under controlled conditions (light, ventilation, temperature and humidity). Our results on the control emissions attest to the validity of our method. In our study, the chemical composition of *L. angustifolia* cv. "Diva" included more than 30 VOCs, dominated by oxygenated monoterpenes (linalyl acetate, camphor, 1,8-cineole), other monoterpenes (myrcene, limonene, linalool, and o- and p-cymene) and sesquiterpenes [α -santalene and (E)- β -caryophyllene]. The chemical composition of our plants was similar to what has been reported in a recent




FIGURE 6 | Visualization of the pairwise correlation matrix of the concentrations-like (peak areas) of all flowering lavender compounds during volatile organic compounds (VOCs) O_3 exposure in experiment 2. Red symbols correspond to negative correlation coefficients, blue symbols correspond to positive correlation coefficients. The size of circles represents the magnitude of the value of the correlation coefficient (r). No circles means that the correlation coefficient was not significant at the threshold of 95% based on Pearson's test.

review of the compounds already identified in other studies of *L. angustifolia* cv. “Diva” (Héral et al., 2021), except that we found a smaller number of compounds. This might be explained by the fact that in all these studies headspace collection of flowering lavender VOCs was performed immediately, or at least shortly after enclosure of the inflorescences, or VOCs were sampled in liquid extractions, thereby including some other compounds stored in the plant but not emitted by flowers (Ormeño et al., 2011). In contrast, in our study, in order to avoid measuring a handling effect (pressure on the glandular trichomes) in the chemical data coming from the release of VOCs stored in the handled tissues (Li et al., 2018), we chose to wait for 12 h after manipulating the plants to start the collection of VOCs. Even if VOCs concentrations were not calculated with full precisions, interestingly, this study highlights, to our knowledge for the first time, a variation in lavender emissions over the course of the day. Proportions and estimated concentrations of monoterpenes changed significantly from morning to afternoon: α -pinene, p-cymene, 1,8-cineole, (Z) and (E)- β -ocimene, linalyl acetate,

and α -santalene increased significantly from morning to noon. A recent study documented diurnal variation in quantities of chemicals stored in different structures of *L. angustifolia* with the highest amount of compounds stored in flowers harvested at 15:00 (Yildirim et al., 2019). Daily emission rhythms in VOCs emission have already been shown in other plants (Zeng et al., 2017). This change in the emissions of plant VOCs could be caused by diurnal changes in abiotic conditions, e.g., light and temperature (Owen et al., 2002; Lerda and Gray, 2003), or may be due to an internal circadian rhythm (Fenske and Imaizumi, 2016; Picazo-Aragón et al., 2020). Since the temperature was controlled in our study, we can assume that VOCs that varied significantly in time in control conditions are compounds that are part of the daily emission rhythm or else compounds whose emissions are light-dependent. As the frequency of visitation of lavender pollinators varies over the course of the day (Nicolè, pers. comm.), it would be interesting to test if the rhythm of emission of lavender volatiles coincides with variations in pollinator foraging periods.




FIGURE 7 | Partial Redundancy Analysis (pRDA) score plot from the constrained Principal Component Analysis of the relative proportions of flowering lavender volatile organic compounds (VOCs) sampled in the reactor as a function of the [O₃] during experiment 2 (A) and the associated correlation circle (B). Purple, yellow, orange, red, and brown colors correspond, respectively, to an exposure of 0, 40, 80, 120, and 200 ppb O₃ and symbols indicate whether the VOCs in the reactor were exposed to 0 ppb O₃ (circles) or increasing [O₃] (triangles). Groups with different letters next to the barycenters show statistically significant differences based on the permutation test assessing the pairwise comparisons at the threshold of 95%.

When we exposed the whole plant to 200 ppb O₃ for 5 h, we detected a significant difference in proportions of VOCs in the chamber between the ozonated and the control lavenders. However, no difference was found between these plants when all the O₃ was removed from the chamber (VOC collection ~45 min after O₃ exposure). These results, combined with the observed decrease in estimated concentrations of terpenes during O₃ exposure, suggest that the changes detected are due to reaction of O₃ with the VOCs in the chamber, rather than to changes in the emission of VOCs by lavenders. This suggests that either short-term O₃ exposure had no effect on VOC emission by lavender, or our sampling time after O₃ exposure was too short to detect a possible response in the emission of lavender VOCs. Indeed, specific organs such as trichomes can be a barrier to O₃ exposure and could thus protect *L. angustifolia* from short but high [O₃] episodes, as has been shown for other plants (Li et al., 2018). Because *L. angustifolia* stores VOCs in trichomes (Guittou et al., 2010) and the time between the production and the emission of stored compounds is longer than for non-stored VOCs, the effect of short-term exposure to O₃ on VOCs emission from lavender might be observed after a longer periods of time than those studied here. For instance, tobacco plants exposed to 170 ppb O₃ for 5 h showed a significant increase in emission of some monoterpenes by leaves 24 h after the exposure (Heiden et al., 1999). Compared to studies conducted on leaf VOCs, studies investigating the effect of O₃ on floral VOCs are limited. Interestingly, it has been reported in some

species of Brassicaceae that emission of some floral VOCs, such as indole and methyl salicylate, can be triggered after 5 days under high [O₃] (Saunier and Blande, 2019). In our study we aimed to explore the immediate response of the plant to a short-term O₃ exposure but it would thus be interesting to test the effect on *L. angustifolia* of a longer period of O₃ exposure, for example from 24 h.

The direct effect of O₃ on the VOCs was clearly confirmed by our experiment testing the result of exposure in the reactor chamber to different [O₃] of the VOCs only, after their emission by the plant. Interestingly, in both experiments, when the whole plant was exposed to O₃ (experiment 1) and when only VOCs were ozonated in the reactor (experiment 2), we detected a significant decrease in the estimated concentrations of terpenes such as linalool, linalyl acetate, α-pinene, and (Z)- and (E)-β-ocimene, (E)-β-caryophyllene and α-santalene, whereas other monoterpenes such as 1,8-cineole, camphor and borneol were relatively more stable during exposure to O₃. In addition, when only the VOCs were ozonated there was a significant decrease in the estimated concentrations of myrcene, geranyl acetone, 1-octen-3-ol-acetate and bornyl acetate. For all these VOCs, estimated concentration decreased linearly with increasing [O₃]. On the contrary, for caryophyllene oxide, the estimated concentration did not seem to follow a monotonous decrease with [O₃]: it decreased significantly at 40, 120, and 200 ppb O₃ but not at 80 ppb. This pattern of variation could be explained by the limited occurrence of this VOC in the blend emitted by

flowering lavender and particularly the absence of this compound in most control samples and samples ozonated at 80 ppb. For the compounds detected in the volatile profile of lavender in this study, the detailed reaction kinetics with O₃ have been described for only two, linalool (McFrederick et al., 2008; Bernard et al., 2012) and (E)- β -caryophyllene (Winterhalter et al., 2009; Jokinen et al., 2016). Because the oxidation of some terpenes by O₃ can produce the same secondary compounds in some cases (McFrederick et al., 2008; Winterhalter et al., 2009; Bernard et al., 2012; Jokinen et al., 2016), accurate determination of the reaction kinetics of VOCs with O₃ can only be approached by performing experimental oxidation studies with a single VOC. Therefore, we can only speculate on the possible degradation-formation of VOCs under O₃ exposure. Besides, we found significant negative correlations between the estimated concentrations of decanal, dodecanal and the unknown compound, on the one hand, and estimated concentrations of monoterpenes [mainly (Z)- and (E)-ocimene, bornyl acetate, borneol, 1-octen-3-ol-acetate, linalyl acetate, and α -pinene], on the other under O₃ exposure (even at [O₃] as low as 40 ppb). These results are in accordance with chemical kinetics studies on the gas-phase terpene oxidation with O₃ (Calogirou et al., 1999; Atkinson and Arey, 2003b), which report carbonyls and carboxylic acids (and sometimes alcohols, epoxides, esters, nitrates, or peroxy nitrates) to be the most prevalent products resulting from these reactions.

As expected, this study highlights different reactivities of some specific lavender VOCs with O₃, suggesting that O₃ exposure not only alters concentrations, but also changes the proportions of compounds in the odor bouquet. Similar results were found in a study investigating the degradation of floral scent volatiles from *Brassica nigra* by reaction with O₃ (at 80 and 120 ppb) over a distance gradient (Farré-Armengol et al., 2016). This latter study showed that supplementing air with O₃ led to a reduction of the concentrations of floral volatiles in air and a change in the proportions with increasing distance from the volatile source, at [O₃] as low as 80 ppb. In that study, attraction tests with *Bombus terrestris* revealed that attractiveness of floral scent to pollinators was reduced after exposure to 120 ppb O₃. In the present study, we found an alteration in the reactor of the VOCs emitted by flowering lavender at even lower [O₃] (40 ppb).

As in most studies exploring the effect of ozone exposure on VOCs and their consequences for plant-insect chemical communication (Fuentes et al., 2013; Farré-Armengol et al., 2016; Li et al., 2016), some conditions of our experiments do not represent the realistic probability of encounter of O₃ with VOCs in normal atmosphere (although the [O₃] we tested were within a realistic range). Nevertheless, our results are alarming. In fact, some carbonyls and carboxylic acids—decanal, dodecanol and dodecanal—appeared to be in higher concentration and/or proportion when the VOCs of lavender were ozonated. These compounds could be detected by pollinators' antennae and could also affect their behavior. For instance, electroantennographic tests showed that bees responded less to high quantities of decanal than to lower quantities (Mas et al., 2018). This study suggested that high concentration of decanal has a repellent effect on bees but behavioral tests are now needed to address this direct effect on this pollinator attraction.

Our findings warn of the strong impact of such modifications of floral chemical signals on the communication of plants with their biotic environment, particularly pollinating insects. Pollination is a fundamental service to natural and agricultural ecosystems (Klein et al., 2007; Winfree et al., 2011). Unfortunately, limited research has been conducted on the impact of O₃ on the chemical communication of plants in the context of plant-pollinator interactions, particularly in the field. Because *L. angustifolia* attracts different groups of insects (Li et al., 2019), which use different signals to find their host plant olfactory cues being of prime importance (Junker and Parachnowitsch, 2015), it would be of great interest to compare the recognition of non-ozonated and ozonated lavender plants by different pollinators. Differences in pollinator sensitivity to variation in plant VOCs induced by high [O₃] could consequently affect the composition of insects visiting lavender plants.

CONCLUSION

We successfully demonstrate that the reactivity with O₃ of major and ubiquitous compounds emitted by lavender plants leads to the formation of new compounds, including carbonyls and carboxylic acids, even at [O₃] as low as 40 ppb, an [O₃] frequently observed in summer. These modifications of individual compounds lead to variation in the proportions of VOCs within the overall bouquet. In contrast, we do not show here an immediate plant response to an O₃ pollution episode affecting the emission of floral VOCs. A general issue to be addressed is whether pollinators might thereby be unable to recognize the O₃-degraded scent. Here we underline the importance of studying the effects of O₃ pollution on plant-insect and especially plant-pollinator interactions in the Mediterranean region. With an increasing number of studies investigating this fundamental issue, thresholds set by European directives on chronic and episodic pollution levels may need to be reviewed and revised to values that take into account scientific studies of the impact of ozone on plant-pollinator interactions.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

CD, MP, and MH-M designed the experiment. CD performed the experiments and analyzed the data. CD and BB identified the chemical compounds. CD, FN, and MP wrote the first version of the article. All authors read and approved the final version.

FUNDING

This work was partly funded by the Montpellier Université d'Excellence (MUSE, ANR under the Investissements d'avenir

program, reference ANR-16-IDEX-0006), International Research Project (IRP)-CNRS-MOST, and the French National Research Program for Environmental and Occupational Health of ANSES (reference 2018/1/138).

ACKNOWLEDGMENTS

We thank the PACE and the TE (LabEx CeMEB, the French National Research Agency under the Investissements d'avenir) for hosting our experiments. We thank the CRIEPPAM for providing the lavender plants and Doyle McKey (a native

speaker) for English proofreading and scientific advices on the manuscript. We also thank two reviewers for their comments on the manuscript, and Michael Staudt, Elena Ormeño-Lafuente, Henri Wortham for their valuable technical advices.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fevo.2022.795588/full#supplementary-material>

REFERENCES

- Abbas, F., Ke, Y., Yu, R., Yue, Y., Amanullah, S., Jahangir, M. M., et al. (2017). Volatile terpenoids: multiple functions, biosynthesis, modulation and manipulation by genetic engineering. *Planta* 246, 803–816. doi: 10.1007/s00425-017-2749-x
- Adams, R. P. (2007). *Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy*, 4th Edn. Carol Stream, IL: Allured publishing corporation.
- Ashmore, M. R. (2005). Assessing the future global impacts of ozone on vegetation. *Plant Cell Environ.* 28, 949–964. doi: 10.1111/j.1365-3040.2005.01341.x
- Atkinson, R., and Arey, J. (2003a). Atmospheric degradation of volatile organic compounds. *Chem. Rev.* 103, 4605–4638. doi: 10.1021/cr0206420
- Atkinson, R., and Arey, J. (2003b). Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review. *Atmos. Environ.* 37, 197–219. doi: 10.1016/S1352-2310(03)00391-1
- Barbier, E. (1963). Les lavandes et l'apiculture dans le sud-est de la France. *Ann. Abeille* 6, 85–159. doi: 10.1051/apido:19630201
- Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., et al. (2012). Approaching a state shift in Earth's biosphere. *Nature* 486, 52–58. doi: 10.1038/nature11018
- Benachour, K. (2017). Insect visitors of lavender (*Lavandula officinalis* L.): comparison of quantitative and qualitative interactions of the plant with its main pollinators. *Afr. Entomol.* 25, 435–444. doi: 10.4001/003.025.0435
- Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* 57, 289–300. doi: 10.1111/j.2517-6161.1995.tb02031.x
- Bernard, F., Daële, V., and Mellouki, A. (2012). Studies of the gas phase reactions of linalool, 6-methyl-5-hepten-2-ol and 3-methyl-1-penten-3-ol with O₃ and OH radicals. *J. Phys. Chem. A* 116, 6113–6126. doi: 10.1021/jp211355d
- Black, V. J., Stewart, C. A., Roberts, J. A., and Black, C. R. (2007). Ozone affects gas exchange, growth and reproductive development in *Brassica campestris* (Wisconsin Fast Plants). *New Phytol.* 176, 150–163. doi: 10.1111/j.1469-8137.2007.02163.x
- Bourtsoukidis, E., Bonn, B., Dittmann, A., Hakola, H., Hellén, H., and Jacobi, S. (2012). Ozone stress as a driving force of sesquiterpene emissions: a suggested parameterisation. *Biogeosciences* 9, 4337–4352. doi: 10.5194/bg-9-4337-2012
- Burkle, L. A., and Runyon, J. B. (2019). Floral volatiles structure plant–pollinator interactions in a diverse community across the growing season. *Funct. Ecol.* 33, 2116–2129. doi: 10.1111/1365-2435.13424
- Calogirou, A., Larsen, B. R., and Kotzias, D. (1999). Gas-phase terpene oxidation products: a review. *Atmos. Environ.* 33, 1423–1439. doi: 10.1016/S1352-2310(98)00277-5
- Cape, J. N. (2008). Surface ozone concentrations and ecosystem health: past trends and a guide to future projections. *Sci. Total Environ.* 400, 257–269. doi: 10.1016/j.scitotenv.2008.06.025
- Curran, J. (2018). *Hotelling: Hotelling's T² Test and Variants*. Available online at: <https://cran.r-project.org/package=Hotelling> (accessed December 22, 2021).
- Directive 2008/50/EC (2008). *Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on Ambient Air Quality and Cleaner Air for Europe*. Luxembourg: Publications office of the European Union.
- Dudareva, N., Klempien, A., Muhlemann, J. K., and Kaplan, I. (2013). Biosynthesis, function and metabolic engineering of plant volatile organic compounds. *New Phytol.* 198, 16–32. doi: 10.1111/nph.12145
- Farré-Armengol, G., Peñuelas, J., Li, T., Yli-Pirilä, P., Filella, I., Llusia, J., et al. (2016). Ozone degrades floral scent and reduces pollinator attraction to flowers. *New Phytol.* 209, 152–160. doi: 10.1111/nph.13620
- Fenske, M. P., and Imaizumi, T. (2016). Circadian rhythms in floral scent emission. *Front. Plant Sci.* 7:462. doi: 10.3389/fpls.2016.00462
- Fowler, D., Amann, M., Anderson, R., Ashmore, M., Cox, P., Depledge, M., et al. (2008). *Ground-Level Ozone in the 21st Century: Future Trends, Impacts and Policy Implications*. London: The Royal Society.
- Fox, J., and Weisberg, S. (2019). *An {R} Companion to Applied Regression*, 3rd Edn. Thousand Oaks CA: Sage.
- Fuentes, J. D., Chamecki, M., Roulston, T., Chen, B., and Pratt, K. R. (2016). Air pollutants degrade floral scents and increase insect foraging times. *Atmos. Environ.* 141, 361–374. doi: 10.1016/j.atmosenv.2016.07.002
- Fuentes, J. D., Roulston, T. H., and Zenker, J. (2013). Ozone impedes the ability of a herbivore to find its host. *Environ. Res. Lett.* 8:014048. doi: 10.1088/1748-9326/8/1/014048
- Ganassi, S., Cascone, P., Di Domenico, C., Pistillo, M., Formisano, G., Giorgini, M., et al. (2020). Electrophysiological and behavioural response of *Phlaenus spumarius* to essential oils and aromatic plants. *Sci. Rep.* 10:3114. doi: 10.1038/s41598-020-59835-1
- Gryparis, A., Forsberg, B., Katsouyanni, K., Analitis, A., Touloumi, G., Schwartz, J., et al. (2004). Acute effects of ozone on mortality from the “Air pollution and health: a European approach” project. *ATS J.* 170, 1080–1087. doi: 10.1164/rccm.200403-333oc
- Guittion, Y., Nicolé, F., Moja, S., Valot, N., Legrand, S., Jullien, F., et al. (2010). Differential accumulation of volatile terpene and terpene synthase mRNAs during lavender (*Lavandula angustifolia* and *L. x intermedia*) inflorescence development. *Physiol. Plant.* 138, 150–163. doi: 10.1111/j.1399-3054.2009.01315.x
- Heiden, A. C., Hoffmann, T., Kahl, J., Kley, D., Klockow, D., Langebartels, C., et al. (1999). Emission of volatile organic compounds from ozone-exposed plants. *Ecol. Appl.* 9, 1160–1167. doi: 10.1890/1051-0761(1999)009[1160:eovocf]2.0.co;2
- Héral, B., Stierlin, É., Fernandez, X., and Michel, T. (2021). Phytochemicals from the genus *Lavandula*: a review. *Phytochem. Rev.* 20, 751–771. doi: 10.1007/s11101-020-09719-z
- Hervé, M. (2021). *RVAideMemoire: Testing and Plotting Procedures for Biostatistics*. Available online at: <https://cran.r-project.org/package=RVAideMemoire> (accessed December 22, 2021).
- Hervé, M. R., Nicolé, F., and Lê Cao, K. A. (2018). Multivariate analysis of multiple datasets: a practical guide for chemical ecology. *J. Chem. Ecol.* 44, 215–234. doi: 10.1007/s10886-018-0932-6
- Holopainen, J. K., and Gershenzon, J. (2010). Multiple stress factors and the emission of plant VOCs. *Trends Plant Sci.* 15, 176–184. doi: 10.1016/j.tplants.2010.01.006
- Jenkin, M. E., and Clemitshaw, K. C. (2000). Ozone and other secondary photochemical pollutants: chemical processes governing their formation in the planetary boundary layer. *Atmos. Environ.* 34, 2499–2527. doi: 10.1016/S1474-8177(02)80014-6

- Jokinen, T., Kausiala, O., Garmash, O., Peräkylä, O., Junninen, H., Schobesberger, S., et al. (2016). Production of highly oxidized organic compounds from ozonolysis of β -caryophyllene: laboratory and field measurements. *Boreal Environ. Res.* 21, 262–273.
- Junker, R. R., and Parachnowitsch, A. L. (2015). Working towards a holistic view on flower traits-how floral scents mediate plant-animal interactions in concert with other floral characters. *J. Indian Inst. Sci.* 95, 43–67.
- Kantsa, A., Raguso, R. A., Lekkas, T., Kalantzi, O.-I., and Petanidou, T. (2019). Floral volatiles and visitors: a meta-network of associations in a natural community. *J. Ecol.* 107, 2574–2586. doi: 10.1111/1365-2745.13197
- Klein, A.-M., Vaissire, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., et al. (2007). Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. Biol. Sci.* 274, 303–313. doi: 10.1098/rspb.2006.3721
- Leitao, L., Bethenod, O., and Biolley, J. P. (2007). The impact of ozone on juvenile maize (*Zea mays* L.) plant photosynthesis: effects on vegetative biomass, pigmentation, and carboxylases (PEPC and Rubisco). *Plant Biol.* 9, 478–488. doi: 10.1055/s-2007-964942
- Lerdau, M., and Gray, D. (2003). Ecology and evolution of light-dependent and light-independent phytochemical volatile organic carbon. *New Phytol.* 157, 199–211. doi: 10.1046/j.1469-8137.2003.00673.x
- Li, H., Li, J., Dong, Y., Hao, H., Ling, Z., Bai, H., et al. (2019). Time-series transcriptome provides insights into the gene regulation network involved in the volatile terpenoid metabolism during the flower development of lavender. *BMC Plant Biol.* 19:313. doi: 10.1186/s12870-019-1908-6
- Li, S., Tosens, T., Harley, P. C., Jiang, Y., Kanagendran, A., Grosberg, M., et al. (2018). Glandular trichomes as a barrier against atmospheric oxidative stress: relationships with ozone uptake, leaf damage, and emission of LOX products across a diverse set of species. *Plant Cell Environ.* 41, 1263–1277. doi: 10.1111/pce.13128
- Li, T., Blande, J. D., and Holopainen, J. K. (2016). Atmospheric transformation of plant volatiles disrupts host plant finding. *Sci. Rep.* 6:33851. doi: 10.1038/srep33851
- Long, S. P., and Naidu, S. L. (2002). “Effects of oxidants at the biochemical, cell and physiological levels, with particular reference to ozone,” in *Air Pollution and Plant Life*, eds J. N. B. Bell and M. Treshow (West Sussex: John Wiley & Sons, Ltd), 69–88.
- Loreto, F., and Schnitzler, J. P. (2010). Abiotic stresses and induced BVOCs. *Trends Plant Sci.* 15, 154–166. doi: 10.1016/j.tplants.2009.12.006
- Mas, F., Harper, A., Horner, R., Welsh, T., Jaksons, P., and Suckling, D. M. (2018). The importance of key floral bioactive compounds to honey bees for the detection and attraction of hybrid vegetable crops and increased seed yield. *J. Sci. Food Agric.* 98, 4445–4453. doi: 10.1002/jsfa.8967
- McFrederick, Q. S., Fuentes, J. D., Roulston, T., Kathilankal, J. C., and Lerdau, M. (2009). Effects of air pollution on biogenic volatiles and ecological interactions. *Oecologia* 160, 411–420. doi: 10.1007/s00442-009-1318-9
- McFrederick, Q. S., Kathilankal, J. C., and Fuentes, J. D. (2008). Air pollution modifies floral scent trails. *Atmos. Environ.* 42, 2336–2348. doi: 10.1016/j.atmosenv.2007.12.033
- McGregor, E. S. (1976). “Crops dependent upon or benefited by insect pollination,” in *Insect Pollination of Cultivated Crop Plants* (Washington, DC: Agricultural Research Service United States Department of Agriculture), 238. doi: 10.5860/choice.37-2165
- Nuvolone, D., Petri, D., and Voller, F. (2018). The effects of ozone on human health. *Environ. Sci. Pollut. Res.* 25, 8074–8088. doi: 10.1007/s11356-017-9239-3
- O’Callaghan, A., Kuhn, M., Weston, S., Wing, J., Forester, J., and Thaler, T. (2020). *contrast: A Collection of Contrast Methods*. Available online at: <https://cran.r-project.org/package=contrast> (accessed December 22, 2021).
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2020). *vegan: Community Ecology Package. R Package Version 2.5-7*. Available online at: <https://CRAN.R-project.org/package=vegan> (accessed March 01, 2022).
- Ormeño, E., Goldstein, A., and Niinemets, Ü. (2011). Extracting and trapping biogenic volatile organic compounds stored in plant species. *Trends Anal. Chem.* 30, 978–989. doi: 10.1016/j.trac.2011.04.006
- Owen, S. M., Harley, P., Guenther, A., and Hewitt, C. N. (2002). Light dependency of VOC emissions from selected Mediterranean plant species. *Atmos. Environ.* 36, 3147–3159. doi: 10.1016/s1352-2310(02)00235-2
- Palmqvist, B., Brazeau, H. A., and Parachnowitsch, A. L. (2021). Differences in floral scent and petal reflectance between diploid and tetraploid *Chamerion angustifolium*. *Front. Ecol. Evol.* 9:734128. doi: 10.3389/fevo.2021.734128
- Passalacqua, N. G., Tundis, R., and Upson, T. M. (2017). A new species of *Lavandula* sect. *Lavandula* (Lamiaceae) and review of species boundaries in *Lavandula angustifolia*. *Phytotaxa* 292, 161–170. doi: 10.11646/phytotaxa.292.2.3
- Peñuelas, J., and Staudt, M. (2010). BVOCs and global change. *Trends Plant Sci.* 15, 133–144. doi: 10.1016/j.tplants.2009.12.005
- Picazo-Aragón, J., Terrab, A., and Balao, F. (2020). Plant volatile organic compounds evolution: transcriptional regulation, epigenetics and polyploidy. *Int. J. Mol. Sci.* 21:8956. doi: 10.3390/ijms21238956
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and R Core Team (2021). *nlme: Linear and Nonlinear Mixed Effects Models*. Available online at: <https://cran.r-project.org/package=nlme>
- Pinto, D. M., Blande, J. D., Souza, S. R., Nerg, A. M., and Holopainen, J. K. (2010). Plant volatile organic compounds (VOCs) in ozone (O₃) polluted atmospheres: the ecological effects. *J. Chem. Ecol.* 36, 22–34. doi: 10.1007/s10886-009-9732-3
- Proffitt, M., Lapeyre, B., Buatois, B., Deng, X. X., Arnal, P., Gouzerh, F., et al. (2020). Chemical signal is in the blend: bases of plant-pollinator encounter in a highly specialized interaction. *Sci. Rep.* 10:10071. doi: 10.1038/s41598-020-66655-w
- Raguso, R. A. (2008b). Wake up and smell the roses: the ecology and evolution of floral scent. *Annu. Rev. Ecol. Syst.* 39, 549–569. doi: 10.1146/annurev.ecolsys.38.091206.095601
- Raguso, R. A. (2008a). Start making scents: the challenge of integrating chemistry into pollination ecology. *Entomol. Exp. Appl.* 128, 196–207. doi: 10.1111/j.1570-7458.2008.00683.x
- Rohart, F., Gautier, B., Singh, A., and Lê Cao, K. A. (2017). mixOmics: an R package for ‘omics feature selection and multiple data integration. *PLoS Comput. Biol.* 13:e1005752. doi: 10.1371/journal.pcbi.1005752
- Rupp, T., Oelschlägel, B., Rabitsch, K., Mahfoud, H., Wenke, T., Disney, R. H. L., et al. (2021). Flowers of deceptive *Aristolochia microstoma* are pollinated by phorid flies and emit volatiles known from invertebrate carrion. *Front. Ecol. Evol.* 9:658441. doi: 10.3389/fevo.2021.658441
- Ryalls, J. M. W., Langford, B., Mullinger, N. J., Bromfield, L. M., Nemitz, E., Pfrang, C., et al. (2022). Anthropogenic air pollutants reduce insect-mediated pollination services. *Environ. Pollut.* 297:118847. doi: 10.1016/j.envpol.2022.118847
- Saunier, A., and Blande, J. D. (2019). The effect of elevated ozone on floral chemistry of *Brassicaceae* species. *Environ. Pollut.* 255:113257. doi: 10.1016/j.envpol.2019.113257
- Sicard, P., Anav, A., De Marco, A., and Paoletti, E. (2017). Projected global ground-level ozone impacts on vegetation under different emission and climate scenarios. *Atmos. Chem. Phys.* 17, 12177–12196. doi: 10.5194/acp-17-12177-2017
- Stierlin, É., Nicolé, F., Costes, T., Fernandez, X., and Michel, T. (2020). Metabolomic study of volatile compounds emitted by lavender grown under open-field conditions: a potential approach to investigate the yellow decline disease. *Metabolomics* 16:31. doi: 10.1007/s11306-020-01654-6
- Thioulouse, J., Dray, S., Dufour, A., Siberchicot, A., Jombart, T., and Pavoine, S. (2018). *Multivariate Analysis of Ecological Data with ade4*. New York, NY: Springer. doi: 10.1007/978-1-4939-8850-1
- Ueda, H., Kikuta, Y., and Matsuda, K. (2012). Plant communication. Mediated by individual or blended VOCs? *Plant Signal. Behav.* 7, 222–226. doi: 10.4161/psb.18765
- Upson, T. M., Andrews, S., and Harriott, G. (2004). *The Genus Lavandula*. Portland, OR: Timber Press.
- Vautard, R., Honoré, C., Beekmann, M., and Rouil, L. (2005). Simulation of ozone during the August 2003 heat wave and emission control scenarios. *Atmos. Environ.* 39, 2957–2967. doi: 10.1016/j.atmosenv.2005.01.039
- Vingarzan, R. (2004). A review of surface ozone background levels and trends. *Atmos. Environ.* 38, 3431–3442. doi: 10.1016/j.atmosenv.2004.03.030
- Wei, T., and Simko, V. (2021). *R Package “corrplot”: Visualization of a Correlation Matrix*. Available online at: <https://github.com/taiyun/corrplot> (accessed December 22, 2021).

- Winfree, R., Gross, B. J., and Kremen, C. (2011). Valuing pollination services to agriculture. *Ecol. Econ.* 71, 80–88. doi: 10.1016/j.ecolecon.2011.08.001
- Winterhalter, R., Herrmann, F., Kanawati, B., Nguyen, T. L., Peeters, J., Vereecken, L., et al. (2009). The gas-phase ozonolysis of β -caryophyllene (C₁₅H₂₄). Part I: an experimental study. *Phys. Chem. Chem. Phys.* 11, 4152–4172. doi: 10.1039/b817824k
- Yeung, L. Y., Murray, L. T., Martinerie, P., Witrant, E., Hu, H., Banerjee, A., et al. (2019). Isotopic constraint on the twentieth-century increase in tropospheric ozone. *Nature* 570, 224–227. doi: 10.1038/s41586-019-1277-1
- Yildirim, M., Sarihan, E., Kil, H., and Khawar, K. (2019). Diurnal and nocturnal variability of essential oil content and components of *Lavandula angustifolia* Mill. (Lavender). *Mustafa Kemal Univ. J. Agric. Sci.* 24, 268–278.
- Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M. Y., et al. (2018). Tropospheric ozone assessment report: assessment of global-scale model performance for global and regional ozone distributions, variability, and trends. *Elementa* 6:10. doi: 10.1525/elementa.265
- Yuan, J. S., Himanen, S. J., Holopainen, J. K., Chen, F., and Stewart, C. N. (2009). Smelling global climate change: mitigation of function for plant volatile organic compounds. *Trends Ecol. Evol.* 24, 323–331. doi: 10.1016/j.tree.2009.01.012
- Zeng, L., Wang, X., Kang, M., Dong, F., and Yang, Z. (2017). Regulation of the rhythmic emission of plant volatiles by the circadian clock. *Int. J. Mol. Sci.* 18:2408. doi: 10.3390/ijms18112408

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Dubuisson, Nicolé, Buatois, Hossaert-McKey and Proffit. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.