



Superlocalization reveals long-range synchronization of vibrating soliton molecules

Said Hamdi, Aurélien Coillet, Benoit Cluzel, Philippe Grelu, Pierre Colman

► To cite this version:

Said Hamdi, Aurélien Coillet, Benoit Cluzel, Philippe Grelu, Pierre Colman. Superlocalization reveals long-range synchronization of vibrating soliton molecules. *Physical Review Letters*, 2022, 128 (21), pp.213902. 10.1103/PhysRevLett.128.213902 . hal-03872036

HAL Id: hal-03872036

<https://hal.science/hal-03872036>

Submitted on 25 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Superlocalization reveals long-range synchronization of vibrating soliton molecules

Said Hamdi, Aurélien Coillet, Benoit Cluzel, Philippe Grelu, and Pierre Colman ^{*1}

¹Laboratoire Interdisciplinaire Carnot de Bourgogne UMR CNRS 6303,
Université Bourgogne-Franche-Comté, 9 av. Savary, 21000 Dijon, France*

(Dated: January 24, 2022)

We implement a super-localization method in the time domain that allows the observation of the external motion of soliton molecules in a fiber ring cavity laser with unprecedented accuracy. In particular, we demonstrate the synchronization of two oscillating soliton molecules separated by several nanoseconds, with inter-molecules oscillations following the same pattern as the intramolecular motion of the individual molecules. These experimental findings indicate an interplay between the different interaction mechanisms that coexist inside the laser cavity, despite their very different characteristic ranges, timescales, strengths, and physical origins.

Keywords: Point Spread Function Deconvolution, Soliton Molecule, HyperResolution, Long range interaction, Multiscale Interplay

INTRODUCTION

In recent years, the dynamics of optical soliton molecules in ultrafast lasers has attracted increasing attention [1]. Soliton molecules consist in solitons forming a bound state owing to their strong interaction [2]. Sometimes, these bound states exhibit periodic internal motions similar to the vibrations of molecules in chemistry. Previous real-time experiments highlighted different oscillatory behaviors, such as vibrations, pure phase, or anharmonic oscillations [3–5]. The specific pattern depends on the laser parameters, namely the existence of a limit cycle attractor [6], combined with the laser noise [7] and other experimental perturbations [8–10]. Beyond their fundamental appeal, soliton molecules could be involved in harmonic mode-locking or in optical data manipulation [11]. Therefore, experiments investigate now soliton molecule dynamics for a larger number of interacting solitons. In a significant number of cases, solitons are not distributed in a regular train of pulses, but form a supra-molecule or molecular complex composed of several, often identical, molecules [10, 12, 13]. This raises the open question of the interplay between the mechanisms responsible for the dynamics governing each molecule, and for the macro organization of the molecular complexes.

Many experiments rely on the dispersive Fourier-transform (DFT) technique [14], which allows a single-shot real-time recording of the spectra over successive cavity roundtrips at multi-MHz frame rates. After numerical processing, DFT spectra yield the dynamics of relative timing and phase between temporally separated pulses. As a major limitation, the observation windows T_{obs} is limited to pulse separations typically below a hundred of picoseconds [8–10, 12]. Such practical limitation is bound on one side by the speed of the detection electronics and on another side by the magnitude of the dispersive line used for pulse stretching. Indeed, the need to avoid the overlap of the DFT

traces for two subsequent pulses prevents the use of a dispersive line of arbitrary length. To set typical orders of magnitudes, considering a electronic bandwidth of ~ 10 GHz and a repetition rate of ~ 10 MHz for ~ 200 fs pulses yields approximatively $T_{\text{obs}} \sim 100$ ps. Moreover, whereas the DFT recording can retrieve the internal motion of a given soliton molecule, it does not provide an accurate information regarding the global motion of the molecule round the laser cavity. Instead, the direct observation - i.e. without pulse stretching - using a GHz-bandwidth oscilloscope can provide a temporal resolution only down to a few tens of picoseconds. Such a timing precision is not enough to observe the weak timing fluctuations that are expected to take place within the supra-molecule. Indeed the strongest mechanisms have the shortest range of interaction [1], while long range interactions are much weaker [7, 15, 16]. Consequently observing the latter would require an acute measurements precision, about on par with the resolution provided by the DFT, but with a much larger observation windows.

To do so we have implemented on the timing channel that records the undispersed optical signal a super-localization procedure akin to what is done in fluorescence spectroscopy [17–19]. Thus we got a timing resolution down to 0.14 ps, 100 times the native sampling resolution. Subtle changes of the molecules cruising velocity round the laser cavity can now be observed. We applied this new technique to the co-evolution of two soliton-pair molecules [5], which are separated by $\simeq 7.6$ ns, one third of the roundtrip time (23.0 ns, FSR = 43.4 MHz). For a clear demonstration of the possibilities offered by combining short and long time scales measurement, two soliton-pair molecules make indeed the simplest multi-molecular system.

INTERNAL MOTION: TWIN MOLECULES

The fiber ring laser cavity under study is composed of 1 meter of erbium-doped fiber closed by 3 meters of SMF fiber (see Figure 1-a)). Nonlinear polarization evolution (NPE) in the fibers provides a virtual saturable absorber effect for mode locking: a polarization beam-splitter (PBS) converts NPE into amplitude modulation and provides as well a laser output port. The first channel of a 80 GSamples/s oscilloscope records the DFT spectrum after an accumulated a chromatic dispersion of -49 ps/nm. It yields the information about the internal motion for each molecule. The second channel records the direct pulsed laser output to track down the global motion of the molecules round the cavity. The direct detection reveals the presence of two sets of pulses and the DFT signal confirms that both sets are actually stable pairs of solitons separated by 5.57 ± 0.01 ps.




FIG. 1. a) Experimental setup; PBS: polarization beam splitter; DCF: 545 m of dispersion-compensating fiber (-90.15 ps/nm/km); Pump: 980-nm laser diode; WDM: pump-signal multiplexer. Single-shot spectra in color scale for b) the leading and c) the trailing soliton molecule. d) Temporal autocorrelation traces after Fourier-transform of c). e) Internal vibration motion for each molecule. Blue (resp. red) line stands for the leading (resp. trailing) molecule. They both exhibits an average soliton separation of 5.57 ps.

The internal vibration of each soliton molecule is ob-

tained by Fourier transform of the corresponding DFT signal, as illustrated by Figure 1-b,c,d). We see in Figure 1-e) that the two molecules both follow a common periodic oscillation with an amplitude of 92 ± 6 fs and a periodicity of 143.5 round trips (RTs). The correlation coefficient between the two trajectories is 0.89 for a relative delay of 87.2 RTs. This strong correlation means that the two oscillations belong to the same family, minus some slight phase drifts - as discussed later in this article. Since the existence and vibrations of soliton molecules are fixed by a common dissipative attractor [6], it is not surprising that the two pairs exhibit similar features. However, the two molecules are not isolated physical objects. Instead, they are likely to interact, as the existence of weak long-range (ns) interactions has been established [7, 16, 20]. Therefore, one could wonder whether the two molecules following a common vibration pattern would benefit from an additional synchronization mechanism. In theory, several distinct vibration patterns may coexist [21?, 22].

EXTERNAL MOTION: POINT SPREAD FUNCTION (PSF) DECONVOLUTION




FIG. 2. a) Timing trace displaying one intensity peak per molecule and per round-trip (RT). b) Reconstruction of the point spread function (PSF) for M1: the random interleaved sampling (RIS) effect between the cavity RT time and the oscilloscope's sampling clock is shown for a few timing pulses. Dashed-black: reconstructed PSF after averaging over 4221 RTs. c) Evolution of the temporal separation between the two molecules (M1, M2) with sub 20-fs resolution. The average inter-molecular separation is $\langle \tau_{ext0} \rangle = 7.58$ ns.

To investigate this matter, we examined further the timing pulse signal. To improve the native 12.5 ps (80 GS/s) sampling resolution, we super-localize the pulses using the point spread response function (PSF) of the detection link. Such a deconvolution technique is widely used in fluorescence microscopy to achieve spatial super-resolution [17–19]. In a nutshell, by superposing together all the pulses acquired at each round-trip, it is possible to take advantage of the aliasing between the cavity free spectral range and the oscilloscope’s internal sampling clock to perform a random interleaved sampling (RIS) and retrieve the shape of PSF with at least a tenfold improvement in the temporal resolution (Figure 2-b)). Note that the PSF is defined using a self-consistent procedure so that a perfectly recurring and stable signal is not required here [23], and pulse jitter can be observed. Once the high resolution PSF is obtained, the time of passage of each pulse can be determined with a precision down to 140 fs. The residual uncertainty is due to a mixture between the oscilloscope’s and laser intrinsic jitters. More details regarding the implementation of the PSF deconvolution can be found in Appendix A. A major difference with fluorescence microscopy is that RIS is possible and more events are recorded, resulting in a better improvement.

The advance or delay for each molecule compared to the average round trip time is shown in Figure 2-c). The most striking feature is that the distance between the two molecules oscillates, following a pattern very similar to the internal motion: same periodicity and nearly the same amplitude (106 ± 18 fs versus 92 ± 6 fs). Similarly to the internal motions, this external oscillation has its own phase offset.

In order to better understand the specificity of this inter-molecular oscillation, we compared in Figure 3 the dynamics of the dual-molecule system with that of a single-molecule, recorded from the same laser setup. Both dynamics show nearly the same internal oscillatory motion (Figure 3-a,d)). However, the two-molecule system exhibits higher fluctuations (Figure 3-b,e)). Using the Parseval’s theorem, most of the external motion (81%) of the two-molecule systems can be explained by an oscillation tuned to the internal vibration state. On the contrary for the single molecule system, the trajectory around the cavity is mainly composed of noise fluctuations which are not related to the internal vibration. The remaining 20% contribution synchronized to the internal oscillation remain hypothetical: a small influence of the molecule dynamics on its own velocity; or a periodic exchange of energy between the two solitons that would cause an apparent motion of its center of mass[24, 25]. Indeed, a vibrating soliton molecule is not a simple two-parameter oscillators [25]. However the comparative analysis of the single molecule systems rules out this




FIG. 3. a) Internal vibrational motion within the soliton molecule M1 composing the two-molecule system. b) Variation of the inter-molecular distance with respect to the round trip number. c) Fourier transform of b): the dashed line indicates the period of the internal vibration shown in a). d-f) Same as a-c, but typical results for a single-molecule system.

phenomenon as being the cause of the external motion observed for the two-molecule systems.

LONG-RANGE INTERACTION AND SYNCHRONIZATION

Clues regarding any interaction between the different oscillators can be inferred by examining how the oscillators respond to fluctuations. Indeed, since the oscillators are separated by several nanoseconds, they do not experience exactly the same random noise. Hence, fully independent and non-interacting oscillators would slowly drift from each other following a random walk pattern.

For slow and weak fluctuations (adiabatic regime), the motion $\tau(t)$ of a harmonic oscillator around its equilibrium position τ_0 can be described as $\tau(t) = \tau_0(t) + A(t) \cos(\omega t + \phi(t))$, where τ_0 , A , and ϕ are the equilibrium position, the oscillation amplitude, and the phase offset, respectively. These parameters fluctuate under the action of noise and drift of the lasing conditions. Therefore, it is possible to get a finer description of the dynamics of the oscillators by monitoring the evolution of the latter three parameters, as shown in Figure 4.

Firstly, the oscillation amplitudes for the internal motions are more stable than for the external relative motion (Figure 4-a). Noting the small temporal extension of each soliton molecule (< 7 ps here), the moderate amplitude excursions may result from a stronger stabilizing interaction. Indeed, dissipative solitons are subjected to phase-sensitive short-range interactions




FIG. 4. a) Instantaneous oscillation amplitude for molecule M1 (A_1), M2 (A_2), and for the inter molecular oscillation (A_{ext}). b) Similar to a) but regarding the relative phase of the oscillation. Black-dashed : result of eq.(1). c) Evolution of the average soliton separation for M1 (blue) and M2 (red), and of the distance between the two molecules (yellow).

[26, 27]. This strong binding attenuates the impact of noise. In addition, the impact of external noise depends on the separation between the oscillator's constituents: it filters out noise at lower frequency that then impacts the molecule as a whole (global drift) but not its internal motion. The weaker interaction and greater sensitivity to low frequency noise explain the faster drift observed for the external oscillator, as displayed on Figure 4-c). That said, considering its spatial extension, the external oscillator has a relative drift proportionally much smaller than that of the oscillating molecules.

Regarding the link between the two molecules, hence the external motion, long-range (nanosecond) interactions are very difficult to model precisely in fiber lasers, because they can combine physical effects such as gain depletion and recovery, electrostriction, and random walk on a noise floor [7, 15, 16, 20]. In contrast with short range interactions, long range effects are much weaker. In addition, they are sensitive to the optical intensity but not to the relative phase between the dissipative solitons. For close-by solitons, the relative optical dephasing is an important features that controls the attraction and repulsion, hence the oscillation of the molecule. A priori there does not exist any mechanism that would control the relative optical phase offset between the very distant two molecules. However a crucial point here is that the internal motion of the molecule is an oscillator that possesses its own phase, which is of different nature than the optical phase while still being closely related to it. Therefore, it is still possible that long-range interactions become sensitive to the phase of the soliton molecule oscillators. Some indications about the possible synchronization between the three oscillators, can be found in

Figure 4-b) that features the evolution of the relative phase offset $\phi(RT) - \phi(0)$. As the two molecules circulate round the cavity, their respective internal motions experience different phase fluctuations and drift away significantly from each other. When it comes to the external motion, we see that its phase evolution is similar to the one that would experience a simple spring-like oscillator when excited at both ends by two oscillatory forces that would derive from the internal motion. Basically, the phase evolution of such a spring oscillator would then satisfy:

$$\begin{aligned} \cos(\omega t + \phi_{spring}(t) + \delta) = \\ \cos(\omega t + \phi_1(t)) + \cos(\omega t + \phi_2(t)) \end{aligned} \quad (1)$$

Where $\phi_1(t)$ and $\phi_2(t)$ are the relative phases of the two driving forces, which we assume here to be the relative phases of the internal motion for each molecule. δ is a constant phase offset that accounts for a delayed response of the oscillator to the external driving, taken here as $\delta \simeq 20^\circ$ (equivalent to an offset of 8 RTs), which yields the dashed line in Figure 4-b). Therefore, the present data analysis indicates that the external oscillation is most likely driven by the combination of the internal motions of the soliton molecules.

DISCUSSION AND CONCLUSION

The above analysis is limited by the number of recorded roundtrips (4221 RTs), itself limited by the oscilloscope memory, so that we cannot give a definite statement regarding the strength of the synchronization between the internal and external motions of the soliton molecules. Nevertheless, the progress of our argument in that direction is made possible by the super-localization techniques and refined data analysis which shed new lights on the complex nonlinear dynamics of multiple soliton molecules. Therefore, by presenting such tools and approach, our article should stimulate further experiments aimed at exploring the dynamics of soliton molecular complexes with an acute precision. We have also analyzed a second experimental data set, namely consisting in another set of oscillating soliton-pair molecules, which is presented in Appendix B. Endowed with similar but not identical features, it confirms the existence of an external oscillation relative to the separation between the soliton molecules. It also leaves the possibility for a synchronization between the external oscillator and the internal ones.

At this point, it is essential to discuss into more details the soliton molecules dynamics. Occurrences of multiples soliton molecules systems have been previously reported in the literature [12, 28]. Studies focus either on the binding mechanisms controlling the equilibrium position between the solitons [9, 10, 28];

or on the internal molecules' dynamics and their possible interaction [12]. A second distinction concerns the type of interactions that are involved. Indeed they can be sensitive or not to the optical phase, hence have different repulsive and attractive landscapes. Therefore depending on the interactions, completely different dynamics are to be expected. In a nutshell, multi-molecules systems are characterized by at least two main features: the macroscopic organization of the molecules; their dynamic. It is indeed important to distinguish between on one side the adiabatic adaptation of the equilibrium positions to an external change of the laser parameters [9]; and on the other side the intrinsic dynamics that occurs at constant laser parameters. Only the latter can be considered as a true oscillator. Thus far, the intrinsic limit of the DFT technics, as discussed in the introduction, limited the studies to situation where the supramolecular cohesion is governed by the same interactions as for the individual molecules binding: short range interactions. Moreover these are strong coherent interactions which lead to a perfect locking of the molecules [12]. In this article, the long range organization of the molecules, which are separated by precisely one third of cavity round trip, is well ascribed by acoustic and optomechanical interactions [29, 30]. These interactions are of different nature than a direct soliton interaction as they are not sensitive to the optical phase. Nevertheless we showed that they may be sensitive to the oscillator's phase and create thus a link between the internal motions. This synchronization effect is very weak, in particular with regard to the laser noise, but it can accumulate over numerous round trips [30]; and hence lead to noticeable effects: the two molecules have the same oscillation pattern. The direct observation of the said synchronization mechanism requires however an extremely precise timing with sub pico-second resolution.

To conclude, we have shown that soliton molecules influence each other over long-range. Despite being separated by several nano-seconds, compact picosecond soliton molecules cannot be considered as independent. Interestingly, long range interactions do not only determine the relative position of the molecules, but can also serve as media to exchange dynamical information by coupling the internal degree of the molecules together. Despite the fact that such possible synchronization will remain weak and cannot give rise to a strict locking, it is an open question whether different molecules could share the same vibration properties and can thus drive each other resonantly. Obviously, the mechanisms involved here would also be present in harmonically mode-locked fiber lasers [10, 31].

The Point Spread Function deconvolution that we

implemented is critical in order to unveil such subtle interactions, as this requires a very precise timing resolution below 300 fs. This technique provides new possibilities to study the dynamics of multi-pulse systems in fiber ring cavity lasers [10]. As a pure numerical post-measurement processing, this deconvolution technique is easy to implement. Note that the final timing resolution after deconvolution depends on the quality of the experimental implementation. As a practical example, we managed in this paper a two order of magnitude improvement of the resolution.

Concerning the specific topic of soliton molecule analysis, performing an instantaneous phase and amplitude analysis – as shown in Figure 4 – is essential to describe with accuracy the vibration patterns. In particular it provides information about the fluctuations, the latter being directly related to the laser noise. Therefore, fluctuations of the molecule vibration reveal the intrinsic properties of the limit-cycle attractor in response to the environment noise. Thus far, description of soliton molecules has been mostly qualitative, with scarce information regarding the stability of the molecules' motion. We hope the present work will serve as incentive for more thorough and quantitative analysis of soliton molecules' vibration patterns.

This work was supported by the French “Investissements d’Avenir” program / project ISITE-BFC (contract ANR-15-IDEX-0003), and by the Agence Nationale de la Recherche (ANR) project “CoMuSim” (contract ANR-17-CE24-0010-01).

The authors declare no conflicts of interest.

APPENDIX A: Determination of the Point Spread Function; and deconvolution procedure

The precise determination of the time of passage is limited by three factors: (i) the finite sampling rate of the oscilloscope, (ii) the measurement noise, and (iii) the presence of a background noise. In the paper, the first factor limits a priori the temporal resolution to 12.5 ps (sampling rate is 80 GS/s). The simplest approach to improve the resolution would be to define the time of passage as a weighted momentum of the whole timing trace. However, the use of a first order momentum is intrinsically limited by the fluctuations of background noise because the timing pulse is short - its spans over a few sampling points, between 2 and 4 - so that the numerous extra data points that constitute the background have a great influence. In turn, an higher order weighted momentum dramatically reduces the background's contribution, but increases the impact of measurement noise. In contrast, the matching of the timing trace with regards to a well deter-

mined Point Spread Function can remove both background fluctuation and measurement noise.

The fitting of the timing pulse against the PSF is a simple task as it is just a matter of translating and scaling the nominal PSF so that it best matches the few data points that constitute the timing trace. The main difficulty is to determine the PSF with enough accuracy.

PSF determination and metrics

Providing one knows with enough accuracy the time of passage $\tau_0(N)$ of each molecule for each round trip N , it is then possible to superpose all the timing trace on top of each other. This operation is equivalent to performing a random interleaved sampling (RIS) because the sampling rate of the oscilloscope and the cavity round trip time are not matched. A weighted Savitzky-Golay filter (i.e. noise reduction through a polynomial regression of the raw data over a sliding windows) is then performed to retrieve the high resolution PSF. In details we set the amplitude $Y(t)$ of the PSF at a given time t as the result of the weighted least square error (W-LSE) fitting of a polynom of 4 order over a time windows of 4 ps span centered around t . Considering the overall much slow response of the detection link, the choice of the polynom's order and of the windows' width have little influence on the final result. The weight that is set on each data point is determined by the error given by a prior non-weighted LSE fitting. The weighted fitting has the advantage to decreases the impact of spurious points and oddly behaving timing pulses. All in all, the PSF is defined with a 10-fold temporal resolution and about 4-bit Y resolution improvement compared to the oscilloscope's specifications. Note that the temporal random interleaved sampling (RIS) has no equivalent in optical microscopy where only noise reduction is possible. As a result the increase of super-localization are smaller than what we achieve here. There is clearly a trade-off between the increase in the temporal resolution and the reduction of the Y measurement noise. Thus far we have not checked what would be the best trade-off.

If knowledge of the PSF allows the precise determination of $\tau_0(N)$, a (good) estimation of $\tau_0(N)$ is at the same time required in order to reconstruct the PSF. The PSF and $\tau_0(N)$ must then be defined using a self-consistent procedure. The measured $\tau_0^{\text{Meas}}(N)$ is composed of several elements :

$$\begin{aligned}\tau_0^{\text{Meas}}(N) &= N \tau_{RT} + \delta^{\text{Meas}}(N) \\ &= N \tau_{RT} + \delta^{\text{slow}}(N) + \delta^{\text{p2p}}(N) + \delta^{\text{error}}(N)\end{aligned}\quad (2)$$

τ_{RT} is the average round trip time, N the round trip number, while the other terms represent the variations

of the time of passage with respect to a perfectly periodic circulation. $\delta^{\text{slow}}(N)$, $\delta^{\text{p2p}}(N)$, and $\delta^{\text{error}}(N)$ correspond respectively to the slow (about 143.5 RTs period here) evolution of the molecule around the laser cavity, the pulse to pulse jitter, and the timing error that we make. The standard deviation of δ^{Meas} taken over all the round trips ($n = \infty$) is then

$$\sigma_{\text{Meas}}^2(n) = \sigma_{\text{slow}}^2(n) + \sigma_{\text{p2p}}^2(n) + \sigma_{\text{error}}^2(n) \quad (3)$$

$\sigma_{\text{slow}}^2(n = \infty)$ dominates over $\sigma_{\text{p2p}}^2(n = \infty)$. However if we consider a number of n consecutive round trips much shorter than characteristic slow evolution of the soliton molecule then $\sigma_{\text{slow}}^2(n) \approx 0$. Considering the molecules oscillate over 143.5 RTs, we chose $n = 8$. The expected value of the standard deviation over all the possible n consecutive bin, reads then:

$$\begin{aligned}\langle \sigma_{\text{Meas}}^2(8) \rangle &\approx \langle \sigma_{\text{p2p}}^2(8) \rangle + \langle \sigma_{\text{error}}^2(8) \rangle \\ &\approx C^{te} + \langle \sigma_{\text{error}}^2(8) \rangle\end{aligned}\quad (4)$$

We define Eq. 4 as our metric. It is minimal when δ^{Error} is minimal - ideally null.

First pass : Initial evaluation of $\tau_0^{\text{Meas}}(N)$

The initial evaluation of $\tau_0^{\text{Meas}}(N)$ ($\delta^{\text{Meas}}(N)$) is given by the computation of the momentum of the timing pulse to the power k : $\tau_0^{\text{Meas}}(N) = \int_N t.Y(t)^k.dt$. k is chosen so that our metrics is minimized as seen in Figure 5. The resulting $\tau_0^{\text{Meas}}(N)$ is shown in Figure 6-a). This simple approach allows the determination of the time of passage with an accuracy of 0.18 ps. Using this first estimate of $\tau_0^{\text{Meas}}(N)$, we compute its corresponding PSF and perform a deconvolution on the timing signal. This first deconvolution reduces the timing accuracy down to 0.13 ps. The resulting variation of time of passage δ^{Meas} is shown in Figure 6-b). If the motion of the soliton molecule round the laser cavity appears now more clearly that in Figure 6-a), large pulse to pulse fluctuations remain. We ascribe these fluctuations to the fact that the PSF that we used is not the exact one but a rough approximation of it. Indeed initial any error on $\tau_0^{\text{Meas}}(N)$ results in an inaccurate RIS sampling, hence an inaccurate determination of the PSF trace, hence a subsequent inaccurate PSF deconvolution.

Subsequent pass(es) : Self-consistent iteration(s)

Considering that rapid fluctuations in Figure 6-b) are mostly artefacts caused by an inaccurate determination of the PSF, they can be filter out (red line Figure 6-b) to result in a better new approximation of $\delta^{\text{Meas}}(N)$. Similarly to what has been done for the first pass, we can now redefine the PSF with better




FIG. 5. Black line: $\langle \sigma_{\text{Meas}}^2(8) \rangle$ versus the power exponent k used in the initial definition of $\tau_0^{\text{Meas}}(N) = \int_N t.Y(t)^k .dt$. Red square: $\langle \sigma_{\text{Meas}}^2(8) \rangle$ after the first pass of super-localization.

accuracy and deconvolve the timing trace once again. The result of this second pass is shown in Figure 6-c). The metrics $\langle \sigma_{\text{Meas}}^2(8) \rangle$ tells the numerical timing uncertainty has been reduced down to 0.02 ps.

This procedure can be repeated until convergence. In our case we did not observe any notable improvement of the metrics by performing a third pass. We see that the final $\delta^{\text{Meas}}(N)$ clearly exhibit distinct temporal features and that most of the spurious peaks seen in Figure 6-b) have been removed. The remaining fluctuations (≈ 100 fs) are much larger than the metrics (20 fs). The small metrics indicate that the errors related to the determination of the time of passage have been reduced to a minimum, the remaining fluctuation correspond then real experimental jitter: laser intrinsic jitter, oscilloscope's clock jitter.




FIG. 6. a) $\delta^{\text{Meas}}(N)$ evaluated using the momentum method with $k = 3.2$ ($\langle \sigma_{\text{Meas}}^2(8) \rangle = 0.18$ ps). b) Super-localization of $\delta^{\text{Meas}}(N)$: first pass ($\langle \sigma_{\text{Meas}}^2(8) \rangle = 0.13$ ps). Red line is the filtered function used as initial guess for the second pass. c) $\delta^{\text{Meas}}(N)$ obtained after the second - and last - pass ($\langle \sigma_{\text{Meas}}^2(8) \rangle = 0.02$ ps).

Hypothesis associated to the deconvolution procedure

The procedure we just described relies on two major assumptions.

First we assume that the shape of the PSF does not change with the internal state of the molecules. Indeed the molecules are about 6 ps long and they undergo little changes: the total power is stable to a few percent and the internal vibration amplitude of only few hundreds of fs. Consequently, these small changes cannot be recorded by the detection link: electronic residual jitter, 80 GS/s sampling rate, oscilloscope with a 40 GHz electronic bandwidth, optical photodiode with a cut-off around 35 GHz, etc. A posteriori this first hypothesis appears to hold. If not, one would have need to short out timing pulses depending of the molecule internal state, and construct one PSF for each internal state. This mitigation plan would nevertheless reduce the number of timing traces used for each PSF, hence reducing the quality of the latter.

The second hypothesis concerns the possibility to define a metrics, hence that the pulse to pulse jitter occurring over one Round-trip is of small amplitude; and that the largest fluctuations are of a much slower dynamics so that it can be simply filtered out through the proper choice of the number of round trips n taken to define the metrics $\langle \sigma_{\text{Meas}}^2(n) \rangle$. Considering that the PSF and $\tau_0^{\text{Meas}}(N)$ are determined using a self-consistent procedure, is it indeed important to define a reliable convergence criteria.

APPENDIX B: Second set of soliton molecules

Here we present the data corresponding to the observation of a second set of soliton-pair molecules using the same experimental setup (Figure 1-a)) and recorded subsequently to the case presented in the main text.

Compared to the first data set, the vibration period is significantly larger (495.6 RTs versus 143.5 RTs), and the two soliton molecules are slightly more compact (temporal extension of 4.6 ps instead of 5.6 ps), see Figure 7. A direct consequence of the longer vibration period is that the molecules can be observed during only 5 oscillation periods. Like in the first data set, the separation between the two molecules is close to 1/3 of the cavity round trip time. These similar conditions allow us to compare the two dynamics.

By monitoring the evolution of the three oscillators' parameters in Figure 8 and comparing with Figure 4, we can see that the second system is more stable and exhibits less fluctuations. We ascribe this to the overall slower dynamics of the systems, the period of oscillation being 3 times larger. That said, we see in Figure 4-c) that the average inter-molecular distance




FIG. 7. a) Motions of the molecule M1 (blue) and M2 (red). b) Variations of the inter-molecular distance. c) Internal vibration of the molecules (M1=Blue; M2=red).




FIG. 8. a) Evolution of the instantaneous vibration amplitude for the intra-molecular oscillations (M1=Blue; M2=red) and for the inter-molecular vibration (yellow) b) Same as a) but regarding the instantaneous phase of the vibration patterns. Dashed-black is the results of Eq.(1). c) Evolution of the average soliton separation for M1 (blue) and M2 (red), and of the distance between the two molecules (yellow).

experiences more drift. Note that due to the long oscillation period, the oscillator's are observed here during only 5 cycles, so that their parameters may not be accurately extracted for the few first and last hundreds of round trips (border effect in the numerical treatment). Finally in Figure 4-a), we see that the amplitude of motion for the external oscillator can be significantly different from the amplitudes of the internal motions (the oscillators' amplitudes were almost equals for the case presented in the main text). This

second set of data confirms the findings regarding the first set. It also provides some further indications regarding the possible relations between the oscillators' parameters: if the two internal motions are very similar to each other, the external oscillation can have a different amplitude and drifts. The limitation in our present investigations comes from the limited memory of the oscilloscope that prevented us to observe the evolution of the molecules over a larger number of round trips.

* * Corresponding author: pierre.colman@u-bourgogne.fr

- [1] P. Grelu and J. M. Soto-Crespo, Temporal soliton molecules in mode-locked lasers: Collisions, pulsations, and vibrations, in *Dissipative Solitons: From Optics to Biology and Medicine* (Springer Berlin Heidelberg, Berlin, Heidelberg, 2008) pp. 1–37.
- [2] M. Stratmann, T. Pagel, and F. Mitschke, Experimental observation of temporal soliton molecules, *PRL* 95, 143902 (2005).
- [3] K. Krupa, K. Nithyanandan, U. Andral, P. Tchofodinda, and P. Grelu, Real-time observation of internal motion within ultrafast dissipative optical soliton molecules, *Phys. Rev. Lett.* 118, 243901 (2017).
- [4] G. Herink, F. Kurtz, B. Jalali, D. R. Solli, and C. Ropers, Real-time spectral interferometry probes the internal dynamics of femtosecond soliton molecules, *Science* 356, 50 (2017).
- [5] S. Hamdi, A. Coillet, and P. Grelu, Real-time characterization of optical soliton molecule dynamics in an ultrafast thulium fiber laser, *Opt. Lett.* 43, 4965 (2018).
- [6] P. Grelu and N. Akhmediev, Dissipative solitons for mode-locked lasers, *Nature Photonics* 6, 84 (2012).
- [7] R. Weill, A. Bekker, V. Smulakovsky, B. Fischer, and O. Gat, Noise-mediated casimir-like pulse interaction mechanism in lasers, *Optica* 3, 189 (2016).
- [8] L. Nimmesgern, C. Beckh, H. Kempf, A. Leitenstorfer, and G. Herink, Soliton molecules in femtosecond fiber lasers: universal binding mechanism and direct electronic control, *Optica* 8, 1334 (2021).
- [9] M. Zhou, J. He, C. Li, Y.-G. Liu, Y. Yue, R. He, S. Chen, L. Zhang, L. Zhu, K. Zhu, K. Chang, and Z. Wang, Oscillatory self-organization dynamics between soliton molecules induced by gain fluctuation, *Optics Express* 29, 16362 (2021).
- [10] W. He, M. Pang, D. H. Yeh, J. Huang, C. R. Menyuk, and P. S. J. Russell, Formation of optical supramolecular structures in a fibre laser by tailoring long-range soliton interactions, *Nature Communications* 10, 10.1038/s41467-019-13746-6 (2019).
- [11] N. Akhmediev, J. M. Soto-Crespo, and G. Town, Pulsating solitons, chaotic solitons, period doubling, pulse coexistence in mode-locked lasers: Complex ginzburg-landau equation approach, *PRE* 63, 056602 (2001).
- [12] Z. Q. Wang, K. Nithyanandan, A. Coillet, P. Tchofodinda, and P. Grelu, Optical soliton molecular com-

- plexes in a passively mode-locked fibre laser, *Nature Communications* 10, 830 (2019).
- [13] Y. Luo, R. Xia, P. P. Shum, W. Ni, Y. Liu, H. Q. Lam, Q. Sun, X. Tang, and L. Zhao, Real-time dynamics of soliton triplets in fiber lasers, *Photon. Res.* 8, 884 (2020).
 - [14] K. Goda and B. Jalali, Dispersive fourier transformation for fast continuous single shot measurements, *Nature Photonics* 7, 102 (2013).
 - [15] J. K. Jang, M. Erkintalo, S. G. Murdoch, and S. Coen, Ultraweak long-range interactions of solitons observed over astronomical distances, *Nature Photonics* 7, 657 (2013).
 - [16] A. N. Pilipetskii, E. A. Golovchenko, and C. R. Menyuk, Acoustic effect in passively mode-locked fiber ring lasers, *Optics Letters* 20, 907 (1995).
 - [17] G. H. Patterson, Fluorescence microscopy below the diffraction limit, *Seminars in Cell & Developmental Biology* 20, 886 (2009).
 - [18] X. Michalet, T. D. Lacoste, and S. Weiss, Ultrahigh-resolution colocalization of spectrally separable point-like fluorescent probes, *Methods* 25, 87 (2001).
 - [19] R. E. Thompson, D. R. Larson, and W. W. Larson, Webb, Precise nanometer localization analysis for individual fluorescent probes, *Biophysical Journal* 82, 2775 (2002).
 - [20] J. N. Kutz, B. C. Collings, K. Bergman, and W. H. Knox, Stabilized pulse spacing in soliton lasers due to gain depletion and recovery, *IEEE Journal of Quantum Electronics* 34, 1749 (1998).
 - [21] T. Schreiber, B. Ortay, J. Limpert, and A. Tünnermann, On the study of pulse evolution in ultra-short pulse mode-locked fiber lasers by numerical simulations, *Opt. Express* 15, 8252 (2007).
 - [22] A. Zavyalov, R. Iliev, O. Egorov, and F. Lederer, Discrete family of dissipative soliton pairs in mode-locked fiber lasers, *Phys. Rev. A* 79, 053841 (2009).
 - [23] E. Lucas, M. Karpov, H. Guo, M. L. Gorodetsky, and T. J. Kippenberg, Breathing dissipative solitons in optical microresonators, *Nature Communications* 8, 10.1038/s41467-017-00719-w (2017).
 - [24] J. Igbonacho, K. Nithyanandan, K. Krupa, P. T. Dinda, P. Grelu, and A. B. Moubissi, Dynamics of distorted and undistorted soliton molecules in a mode-locked fiber laser, *Phys. Rev. A* 99, 063824 (2019).
 - [25] P. Colman, A. Coillet, S. Hamdi, P. Tchofo-Dinda, and P. Grelu, Higher dimensional oscillations of soliton molecules in ultrafast fiber laser, in *Conference on Laser and Electro Optics (CLEO Europe)*, pp. EF-8.2.
 - [26] B. A. Malomed, Bound solitons in the nonlinear schrödinger-ginzburg-landau equation, *Physical Review A* 44, 6954 (1991).
 - [27] J. M. Soto-Crespo, N. Akhmediev, P. Grelu, and F. Belhache, Quantized separations of phase-locked soliton pairs in fiber lasers, *Opt. Lett.* 28, 1757 (2003).
 - [28] Y. Zhou, Y.-X. Ren, J. Shi, H. Mao, and K. K. Y. Wong, Buildup and dissociation dynamics of dissipative optical soliton molecules, *Optica* 7, 965 (2020).
 - [29] M. Erkintalo, K. Luo, J. K. Jang, S. Coen, and S. G. Murdoch, Bunching of temporal cavity solitons via forward brillouin scattering, *New Journal of Physics* 17, 115009 (2015).
 - [30] Y. Jaouën and L. du Mouza, Transverse brillouin effect produced by electrostriction in optical fibers and its impact on soliton transmission systems, *J. of Stat. Mech.* 7 (3), 141.
 - [31] X. Liu and M. Pang, Revealing the buildup dynamics of harmonic mode-locking states in ultrafast lasers, *Laser & Photonics Reviews* 13, 1800333 (2019).