



HAL
open science

Propagation of viral genomes by replicating ammonia-oxidising archaea during soil nitrification

Sungeun Lee, Ella Sieradzki, Graeme Nicol, Christina Hazard

► To cite this version:

Sungeun Lee, Ella Sieradzki, Graeme Nicol, Christina Hazard. Propagation of viral genomes by replicating ammonia-oxidising archaea during soil nitrification. *The International Society of Microbiological Ecology Journal*, In press, 10.1038/s41396-022-01341-5 . hal-03872694

HAL Id: hal-03872694

<https://hal.science/hal-03872694>

Submitted on 25 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BRIEF COMMUNICATION OPEN



Propagation of viral genomes by replicating ammonia-oxidising archaea during soil nitrification

Sungeun Lee¹, Ella T. Sieradzki¹, Graeme W. Nicol¹✉ and Christina Hazard¹✉

© The Author(s) 2022

Ammonia-oxidising archaea (AOA) are a ubiquitous component of microbial communities and dominate the first stage of nitrification in some soils. While we are beginning to understand soil virus dynamics, we have no knowledge of the composition or activity of those infecting nitrifiers or their potential to influence processes. This study aimed to characterise viruses having infected autotrophic AOA in two nitrifying soils of contrasting pH by following transfer of assimilated CO₂-derived ¹³C from host to virus via DNA stable-isotope probing and metagenomic analysis. Incorporation of ¹³C into low GC mol% AOA and virus genomes increased DNA buoyant density in CsCl gradients but resulted in co-migration with dominant non-enriched high GC mol% genomes, reducing sequencing depth and contig assembly. We therefore developed a hybrid approach where AOA and virus genomes were assembled from low buoyant density DNA with subsequent mapping of ¹³C isotopically enriched high buoyant density DNA reads to identify activity of AOA. Metagenome-assembled genomes were different between the two soils and represented a broad diversity of active populations. Sixty-four AOA-infecting viral operational taxonomic units (vOTUs) were identified with no clear relatedness to previously characterised prokaryote viruses. These vOTUs were also distinct between soils, with 42% enriched in ¹³C derived from hosts. The majority were predicted as capable of lysogeny and auxiliary metabolic genes included an AOA-specific multicopper oxidase suggesting infection may augment copper uptake essential for central metabolic functioning. These findings indicate virus infection of AOA may be a frequent process during nitrification with potential to influence host physiology and activity.

The ISME Journal; <https://doi.org/10.1038/s41396-022-01341-5>

Microbially mediated oxidation of ammonia to nitrate during nitrification is a central component of the global nitrogen (N) cycle. It is also responsible for major losses of applied fertiliser N in soil, generating atmospheric pollution via direct and indirect production of nitrous oxide (N₂O) as well as nitrate (NO₃⁻) pollution of groundwater [1]. Autotrophic ammonia-oxidising archaea (AOA) of the class *Nitrososphaeria* are a ubiquitous component of soil microbial communities and often dominate ammonia oxidation and nitrification-associated N₂O emissions when ammonia is supplied at low rates via organic matter mineralisation [2], slow-release fertilisers [3] or in acidic soils [4]. Integrated temperate viruses (proviruses) and other virus-associated protein-encoding genes are found in most AOA genomes suggesting frequent interaction (see Supplementary Text). While viruses infecting marine AOA have been characterised through metagenomic approaches [5] and cultivation [6], those infecting soil AOA or other nitrifier groups are currently uncharacterised.

Virus infection can influence biogeochemical cycling by augmenting host activity or causing cell mortality and subsequent release of nutrients [7]. Recent advances have demonstrated that soil virus communities are dynamic in a wide range of soils [e.g. 8, 9] and augmenting virus loads modulate C and N fluxes [10, 11]. Nevertheless, identifying active interactions with specific

populations or functional groups in soil remains challenging due to structural complexity and the vast diversity of hosts and viruses. Recent use of stable-isotope approaches has investigated whole community host-virus dynamics [12, 13] or interactions between individual host-virus populations specific to a functional process and substrate [14]. The aim of this study was to utilise the latter approach with ¹³CO₂-based DNA-SIP to focus on nitrification-associated interactions and to test the hypothesis that viruses are a dynamic component of soil AOA activity.


HYBRID ANALYSIS OF GC MOL% FRACTIONATION AND ¹³C-DNA-SIP FOR IDENTIFYING ACTIVE AOA HOSTS AND VIRUSES

Microcosms were established (see Supplementary Text) with two sandy loam soils of contrasting pH (4.5 or 7.5), which were characterised previously as containing distinct AOA communities [15]. As ammonia from hydrolysed urea is used by AOA in these and other soils [1, 3], nitrification was fuelled by addition of urea (two pulses of 100 µg N g⁻¹ soil (dry weight)) to stimulate AOA growth and potential enrichment of AOA-infecting viruses. Microcosms were incubated at 25 °C for 30 days (Supplementary Fig. 1) with a 5% ¹²C- or ¹³C-CO₂ headspace to target viruses of autotrophs via transfer of assimilated carbon. Extracted DNA was subjected to isopycnic centrifugation in CsCl gradients, in which

¹Univ Lyon, CNRS, INSA Lyon, Université Claude Bernard Lyon 1, Ecole Centrale de Lyon, Ampère, UMR5005, 69134 Ecully, France. ✉email: graeme.nicol@ec-lyon.fr; christina.hazard@ec-lyon.fr

Received: 26 August 2022 Revised: 2 November 2022 Accepted: 4 November 2022

Published online: 21 November 2022



DNA migrates as a function of GC mol% and isotopic enrichment, before fractionation and analysis of genomic DNA of different buoyant densities [14]. Quantification of AOA genomic DNA distribution across CsCl gradients demonstrated growth of AOA populations (Fig. 1A; Supplementary Fig. 2) and genomic DNA in high buoyant density (HBD) fractions ($>1.719 \text{ g ml}^{-1}$) was then

pooled for each replicate microcosm and metagenomes sequenced (Supplementary Table 1). While ^{13}C -incorporation clearly separated DNA from growing vs non-growing AOA populations, ^{13}C -enriched AOA genomes (average contig GC mol% content of 38%) co-migrated within the CsCl gradient with more abundant community ^{12}C genomes with higher GC mol%

Fig. 1 Contigs and MAGs from low GC mol% DNA and in situ growth of AOA in pH 4.5 and 7.5 soils determined from hybrid analysis of ^{12}C - and ^{13}C -enriched DNA. **A** Distribution of total prokaryotic and AOA genomes in CsCl gradients determined from the relative abundance of 16S rRNA genes and *amoA* genes, respectively. Example profiles are shown for pH 4.5 soil only (see Supplementary Fig. 2 for pH 7.5 profiles). Vertical error bars are the standard error of the mean relative abundance and horizontal bars (mostly smaller than the symbol size) the standard error of the mean buoyant density of individual fractions from three independent CsCl gradients, each representing an individual microcosm. Genomic DNA in fractions highlighted in blue or pink areas were pooled for each replicate microcosm for metagenomic sequencing. **B** Distribution of assembled contigs (≥ 5 kb) from metagenomic libraries prepared from LBD or HBD DNA in sequence coverage vs. GC mol% plots. Contigs binned into ten medium- and high-quality nitrifier MAGs and predicted viruses of nitrifiers (≥ 10 kb) are highlighted, with numbers in parenthesis beside each MAG identifier giving estimated completeness and contamination (%). **C** Mean and standard error of the relative proportion of metagenomic reads from ^{12}C - and ^{13}C -enriched HBD DNA mapped onto contigs of nine AOA and one *Nitrobacter* MAGs. A significantly greater relative proportion of reads in ^{13}C -derived libraries are indicated with * ($p < 0.05$, two-sample Student's t-test or Welsch's t-test when variances were not homogenous). GTDB genus [16] and *amoA* gene lineage-affiliation [17] are given.

(>57%), reducing AOA host and virus sequencing depth, contig assembly and binning. Consequently, only one medium-quality AOA metagenome-assembled genome (MAG) was recovered from HBD DNA and no AOA-associated virus contigs were identified amongst 290 viral operational taxonomic units (vOTUs) ≥ 10 kb (Fig. 1B).

To facilitate analysis of low GC mol% genomes while also identifying activity, we developed a hybrid approach combining metagenomic analysis of low buoyant density (LBD) ^{12}C DNA with read-mapping of sequenced HBD DNA (Fig. 1; Supplementary Fig. 2). The AOA and virus genomes were assembled from metagenomic libraries of ^{12}C LBD DNA only ($< 1.699 \text{ g ml}^{-1}$) representing both dormant and active populations. Individual sequence reads from HBD DNA libraries from both ^{12}C - and ^{13}C -incubations were then mapped to identify activity with a significantly greater relative abundance in ^{13}C -incubated samples demonstrating isotope incorporation. For taxonomically defined contigs of hosts or host-linked viruses, in comparison to HBD, assemblies from LBD libraries increased the relative proportion of contigs from the AOA-containing *Thermoproteota* phylum (GTDB taxonomy; *Nitrososphaerota* in ICNP taxonomy and *Thaumarchaeota* in NCBI taxonomy) and *Thermoproteota*-infecting virus contigs from 0.3 to 2.8% and 0 to 22.3%, respectively (Supplementary Fig. 3). In LBD metagenomes, 123 medium- and high-quality MAGs from both soils were assigned taxonomy (Supplementary Table 2) including nine AOA MAGs representing six genera (GTDB classification [16]) within the *Nitrososphaerales* and *Nitrosopumilales* orders and one nitrite-oxidising bacterium (NOB) *Nitrobacter* MAG (MAG 15) (Fig. 1B; Supplementary Fig. 4), contributing to the oxidation of nitrite (NO_2^-) to NO_3^- during the second step of canonical nitrification. Mapping of HBD sequence reads revealed that eight of the ten MAGs were derived from active autotrophic populations (Fig. 1C). Consistent with previous studies characterising the distribution of soil AOA from different taxa [17], this included four *Nitrosotalea* genus populations (MAGs 9, 61, 77 and 107) from pH 4.5 soil and three *Nitrososphaera* genus populations (MAGs 12, 52 and 96) from pH 7.5 soil. No evidence of autotrophic growth was observed for populations represented by a further two MAGs within the *Nitrososphaerales*. Both are from genera with no cultivated representatives with MAG 112 from genus TA-21 (*amoA*-gene lineage NS- β -2 [17]) and MAG 63 (*amoA*-gene lineage NS- δ -2.1) representing a novel genus (designated here as CS63). The latter belongs to the *amoA* gene-defined NS- δ lineage that while widely found in soil, is rarely implicated in contributing to ammonia oxidation. Although these two MAGs may simply be from AOA populations not active under the incubation conditions used, they are also consistent with microorganisms possessing physiologies distinct from cultivated AOA.


AOA VIRUSES

After predicting contigs of virus origin using established bioinformatic tools, those from AOA-infecting viruses were identified with a custom database of hallmark genes from AOA proviruses and

analysis of other shared homologues (see Supplementary Text). Specifically, proviruses were identified in 184 (87%) of 212 AOA genome sequences downloaded from GTDB. Capsid, terminase, portal and integrase protein sequences from these proviruses were subsequently used as search queries. While previous analysis of 12 total soil metagenome or virome libraries from the same soil samples did not enable identification of any AOA-associated virus contigs (average of 152 million quality filtered reads and 1,947 contigs ≥ 10 kb per library) [14], this targeted approach using LBD DNA from nitrifying microcosms yielded 64 contigs predicted as derived from AOA-infecting viruses and each representing a unique vOTU. In addition, viruses infecting other nitrifier populations were identified with five and three vOTUs derived from viruses infecting ammonia-oxidising bacteria (AOB) and NOB, respectively. Virus and host genome GC mol% correlate [18] and AOA-infecting virus contigs possessed a similar average content of 36.3% which was considerably lower than averages of 46.6% and 67.0% of all viral contigs recovered from LBD and HBD metagenomes, respectively. As with their hosts, AOA virus communities were distinct between the two pH soils (Fig. 2A) as also observed previously for total virus communities [19] with 29 (45%) and 34 (53%) vOTUs significantly more relatively abundant in pH 4.5 and 7.5 LBD libraries compared to the other soil, respectively. Comparison of the relative abundance of HBD sequence reads from both ^{12}C and ^{13}C libraries revealed that 27 contigs from both soils (42%) were enriched in ^{13}C and thus confirming infection during incubation and transfer of C from AOA hosts to viruses.

Soil AOA viruses were distinct from characterised archaeal and bacterial viruses. Gene-sharing network analysis using vConTACT v2.0 [20] did not identify any associations with reference sequences or recently described viruses of marine AOA [6] and genome-wide sequence comparisons [21] placed 60 of the 64 soil AOA viral contigs within a unique proteomic grouping (Fig. 2B). The majority were predicted as capable of lysogeny with 55 encoding an integrase. While it can be difficult to determine virulent versus temperate capability from partial genomes, only 6.8 and 8.9% of all virus contigs from LBD and HBD metagenomes, respectively, possessed a predicted integrase. Although suggesting a higher proportion of AOA viruses may be temperate, it should be noted that this could be a consequence of using AOA provirus-associated genes to identify viral contigs, biasing towards identification of those from temperate viruses. These results also highlight that while DNA-SIP of total metagenomes enables identification of ^{13}C incorporation into virus genomes, it is potentially limited in facilitating confident bioinformatic prediction of free temperate virus particles versus proviruses present from a previous infection of a currently active host. While the integration status of a temperate virus can be predicted bioinformatically, using targeted "viromes" metagenomes of filtered virus particles will provide greater confidence of identifying genomes from free virus particles.

Eleven contigs contained large terminase (TerL) sub-unit gene sequences and were similar to those recovered from other predicted AOA virus or provirus genomes only (e-value $< 10^{-5}$)



(Fig. 2C). In addition to viral hallmark genes encoding portal, endonuclease, tail and capsid proteins, contig 053596 contained a type 1 multicopper oxidase (MCO1) homologue found exclusively in AOA genomes [22] (Fig. 2D). Copper (Cu) is essential for ammonia oxidation and electron transport in AOA and MCO1

is upregulated under Cu limitation and potentially involved in Cu sequestration from the environment [23]. This virus-encoded MCO1 homologue therefore represents an auxiliary metabolic gene that may contribute to a core process in AOA physiology.

Fig. 2 Identification and determination of activity of viruses infecting AOA in pH 4.5 and 7.5 soils using a hybrid analysis of ^{12}C - and ^{13}C -enriched DNA. **A** Heatmap showing the relative abundance (reads per kb, ln transformed) of 64 vOTUs identified in LBD DNA libraries from pH 4.5 and 7.5 soil ($\geq 1\times$ coverage, $\geq 75\%$ contig breadth). To identify activity, reads from ^{12}C - and ^{13}C -enriched HBD DNA were mapped onto LBD-derived contigs with a significantly greater relative proportion of reads in ^{13}C -derived libraries indicated with * ($p < 0.05$, two-sample Student's t-test or Welsch's t-test when variances were not homogenous). **B** Proteomic tree showing genome-wide sequence similarities between AOA viral contigs and 1,981 curated virus genomes. Values at dotted circles represent a distance metric based on normalized tBLASTx scores plotted on a log scale. **C** Maximum likelihood phylogenetic analysis of derived large sub-unit terminase (TerL) protein sequences (336 unambiguously aligned positions, LG substitution model) identified in this and other metagenomic studies or in proviruses of cultivated AOA. NCBI accession numbers are given in parenthesis. Circles at nodes represent percentage bootstrap support from 1000 replicates, scale bar denotes an estimated 0.05 changes per position and colour-coding describes the comparative relative abundance in each soil. **D** Genetic map of AOA-infecting virus-derived contig 053596 containing a type 1 multicopper oxidase (MCO1) gene. Maximum likelihood phylogenetic tree describes placement of MCO1 within the AOA MCO/NirK family of cultivated AOA following designations of Kerou et al. [22] (206 unambiguously aligned positions, LG substitution model, invariant and gamma distributed sites). NCBI accession numbers are given in parenthesis. Circles at nodes represent percentage bootstrap support from 1000 replicates and the scale bar denotes an estimated 0.05 changes per position.

SUMMARY

These results from two soils indicate that virus infection of AOA during nitrification is likely a common and dynamic process, with the potential to influence nitrification rates in soil. Although viruses were only identified from contigs representing partial viral genomes, analysis of individual marker genes and genome-wide similarity comparisons revealed that those infecting AOA are highly divergent from known viruses. While one *Nitrobacter* MAG was obtained and both AOB and NOB viruses were also predicted, the hybrid approach developed in this study targeted low GC mol% DNA and was specifically tailored for the analysis of AOA populations and their viruses. A broader analysis of all viruses infecting nitrifying prokaryotic populations would therefore likely require a more comprehensive sequencing effort of all fractions throughout the CsCl gradient using quantitative SIP metagenomics [24] or alternative approaches, such as virus-targeted metagenomic viromes.

DATA AVAILABILITY

Metagenome sequence data are deposited in NCBI's GenBank under BioProject accession nos. PRJNA621418-PRJNA621429 (^{13}C data) and PRJNA868779 (^{12}C data) with MAGs available under accession numbers SAMN30436125- SAMN30436246.

REFERENCES

- Prosser JI, Hink L, Gubry-Rangin C, Nicol GW. Nitrous oxide production by ammonia oxidizers: Physiological diversity, niche differentiation and potential mitigation strategies. *Glob Chang Biol.* 2020;26:103–18.
- Huang L, Chakrabarti S, Cooper J, Perez A, John SM, Daroub SH, et al. Ammonia-oxidizing archaea are integral to nitrogen cycling in a highly fertile agricultural soil. *ISME Commun.* 2021;1:19.
- Hink L, Gubry-Rangin C, Nicol GW, Prosser JI. The consequences of niche and physiological differentiation of archaeal and bacterial ammonia oxidisers for nitrous oxide emissions. *ISME J.* 2018;12:1084–93.
- Li Y, Chapman SJ, Nicol GW, Yao H. Nitrification and nitrifiers in acidic soils. *Soil Biol Biochem.* 2018;116:290–301.
- Ahlgren NA, Fuchsman CA, Rocap G, Fuhrman JA. Discovery of several novel, widespread, and ecologically distinct marine Thaumarchaeota viruses that encode *amoC* nitrification genes. *ISME J* 2019;13:618–31.
- Kim J-G, Kim S-J, Cvirkaite-Krupovic V, Yu W-J, Gwak J-H, López-Pérez M, et al. Spindle-shaped viruses infect marine ammonia-oxidizing thaumarchaea. *Proc Natl Acad Sci USA.* 2019;116:15645–50.
- Emerson JB. Soil Viruses: A New Hope. *mSystems* 2019;4:e00120–19.
- Santos-Medellin C, Estera-Molina K, Yuan M, Pett-Ridge J, Firestone MK, Emerson JB. Spatial turnover of soil viral populations and genotypes overlain by cohesive responses to moisture in grasslands. *Proc Natl Acad Sci USA.* 2022;119:e2209132119.
- Wu R, Davison MR, Gao Y, Nicora CD, Mcdermott JE, Burnum-Johnson KE, et al. Moisture modulates soil reservoirs of active DNA and RNA viruses. *Commun Biol.* 2021;4:992.
- Braga LPP, Spor A, Kot W, Breuil M-C, Hansen LH, Setubal JC, et al. Impact of phages on soil bacterial communities and nitrogen availability under different assembly scenarios. *Microbiome* 2020;8:52.

- Albright MBN, Gallegos-Graves LV, Feeser KL, Montoya K, Emerson JB, Shakya M, et al. Experimental evidence for the impact of soil viruses on carbon cycling during surface plant litter decomposition. *ISME Commun.* 2022;2:24.
- Starr EP, Shi S, Blazewicz SJ, Koch BJ, Probst AJ, Hungate BA, et al. Stable-isotope-informed, genome-resolved metagenomics uncovers potential cross-kingdom interactions in rhizosphere soil. *mSphere* 2021;6:e0008521.
- Trubl G, Kimbrel JA, Lique-Gonzalez J, Nuccio EE, Weber PK, Pett-Ridge J, et al. Active virus-host interactions at sub-freezing temperatures in Arctic peat soil. *Microbiome* 2021;9:208.
- Lee S, Sieradzki ET, Nicolas AM, Walker RL, Firestone MK, Hazard C, et al. Methane-derived carbon flows into host-virus networks at different trophic levels in soil. *Proc Natl Acad Sci USA.* 2021;118:e2105124118.
- Nicol GW, Leininger S, Schleper C, Prosser JI. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environ Microbiol.* 2008;10:2966–78.
- Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 2020;36:1925–7.
- Alves RJE, Minh BQ, Ulrich T, von Haesele A, Schleper C. Unifying the global phylogeny and environmental distribution of ammonia-oxidising archaea based on *amoA* genes. *Nat Commun.* 2018;9:1517.
- Cardinale DJ, Duffy S. Single-stranded genomic architecture constrains optimal codon usage. *Bacteriophage* 2011;1:219–24.
- Lee S, Sorensen JW, Walker RL, Emerson JB, Nicol GW, Hazard C. Soil pH influences the structure of virus communities at local and global scales. *Soil Biol Biochem.* 2022;166:108569.
- Jang HB, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al. Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by gene-sharing networks. *Nat Biotechnol.* 2019;37:632–9.
- Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, Goto S. ViPTree: the viral proteomic tree server. *Bioinformatics* 2017;33:2379–80.
- Kerou M, Offre P, Valleder L, Abby SS, Melcher M, Nagler M, et al. Proteomics and comparative genomics of *Nitrososphaera viennensis* reveal the core genome and adaptations of archaeal ammonia oxidizers. *Proc Natl Acad Sci Usa* 2016;113:7937–46.
- Reyes C, Hodgskiss LH, Kerou M, Pribansnig T, Abby SS, Bayer B, et al. Genome wide transcriptomic analysis of the soil ammonia oxidizing archaeon *Nitrososphaera viennensis* upon exposure to copper limitation. *ISME J* 2020;14:2659–74.
- Sieradzki ET, Greenlon A, Nicolas AM, Firestone MK, Pett-Ridge J, Blazewicz SJ, et al. Functional succession of actively growing soil microorganisms during rewetting is shaped by precipitation history. *bioRxiv.* 2022; 2022.06.28.498032.

ACKNOWLEDGEMENTS

This work was funded by an AXA Research Chair in Microbial Ecology awarded to GWN, Agence Nationale de la Recherche grant 'CONSERVE' (ANR-22-CE02-0006-01) awarded to GWN and CH, and an EU Horizon Europe MSCA Postdoctoral Fellowship 'DIVOBIS' (101064472) awarded to ETS. Sequencing data generated under JGI Community Science Program Proposal No. 503702 awarded to GWN and CH was conducted by the US DOE JGI, a DOE Office of Science User Facility, and supported by the Office of Science of the US DOE under Contract No. DE-AC02-05CH11231. We acknowledge use of the Newton high-performance computing cluster at PMCS21, École Centrale de Lyon. The authors would like to thank Dr. Willm Martens-Habbena (University of Florida) for continued discussions on soil AOA ecophysiology and taxonomy, Dr. Melina Kerou (University of Vienna) for guidance on AOA MCO diversity, Dr. Gary Trubl for comments on the manuscript and Dr. Robin Walker (SRUC, Aberdeen) for access to the much-lamented Craibstone Estate Woodlands Field long term experiment.

AUTHOR CONTRIBUTIONS

SL, GWN and CH conceived and designed the experiment with additional input from ETS. SL performed all laboratory work and analysed the data with input from GWN in phylogenetic analyses. SL, GWN and CH drafted the manuscript text and figures with all authors contributing to the final paper.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41396-022-01341-5>.

Correspondence and requests for materials should be addressed to Graeme W. Nicol or Christina Hazard.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022