



**HAL**  
open science

# Cyclic Diaryl $\lambda^3$ -Bromanes: A Rapid Access to Molecular Complexity via Cycloaddition Reactions

Matteo Lanzi, Racha Abed, Ali Abdine, Maxime de Abreu, Joanna Wencel-Delord

► **To cite this version:**

Matteo Lanzi, Racha Abed, Ali Abdine, Maxime de Abreu, Joanna Wencel-Delord. Cyclic Diaryl  $\lambda^3$ -Bromanes: A Rapid Access to Molecular Complexity via Cycloaddition Reactions. *Organic Letters*, 2021, 23 (23), pp.9047-9052. 10.1021/acs.orglett.1c03278 . hal-03874364

**HAL Id: hal-03874364**

**<https://hal.science/hal-03874364>**

Submitted on 28 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Cyclic Diaryl $\lambda^3$ -Bromanes: a Rapid Access to Molecular Complexity via Cycloaddition Reactions

Matteo Lanzi, Racha Abed Ali Abdine, ‡ Maxime De Abreu, ‡ and Joanna Wencel-Delord\*

Laboratoire d'Innovation Moléculaire et Applications (UMR CNRS 7042), Université de Strasbourg/Université de Haute Alsace, ECPM, 25 rue Becquerel, 67087 Strasbourg (France)

Supporting Information Placeholder



**ABSTRACT:** Biaryls have widespread applications in organic synthesis. However, sequentially polysubstituted biaryls are underdeveloped due to their challenging preparation. Herein, we report the synthesis of unsymmetric 2,3,2',3',4-substituted biaryls *via* pericyclic reactions of cyclic diaryl  $\lambda^3$ -bromanes. The functional groups tolerance and atom economy allow access to molecular complexity in a single reaction step. Continuous flow protocol has been designed for the scale-up of the reaction, while post-functionalizations have been developed taking advantage of the residual bromide.


The direct access to molecular complexity under sustainable, simple, and mild reaction conditions is a critical challenge of modern organic chemistry. Biaryl compounds are ubiquitous motifs in many fields of chemistry, including medicinal chemistry, agrochemistry, material science, and ligand design. A straightforward synthesis of highly decorated biaryls thus remains an important challenge.<sup>1</sup> In particular, dissymmetrical 2,3,2',3',4-substituted biaryls are poorly explored motifs due to the arduous synthesis *via* standard cross-coupling reactions which require densely decorated precursors. Albeit scarcely investigated, a general and straightforward alternative approach based on a modular functionalization *via* pericyclic reactions of a finely designed biaryl benzyne precursor represents an attractive solution to this synthetic issue (Scheme 1a).<sup>2</sup> However, multiple drawbacks hamper the state-of-the-art of the aryne-based approach (Scheme 1b): 1) lengthy synthetic routes for the preparation of aryne precursors; 2) the limited atom economy resulting from the aryne generation with the removal of leaving groups (-OTf and -TMS; -X and -OR; Aux-I) which translate into significant waste formation and 3) the limited functional group tolerance as the base- and fluorine-sensitive moieties are generally incompatible.<sup>3-5</sup> Accordingly, the implementation of the pericyclic reactions to rapidly access dissymmetrical 2,3,2',3',4-substituted biaryls diversity requests design of original biaryl aryne precursors, activated under mild reaction conditions and in presence of a weak base.

Despite being known since the 19th century and albeit their unique reactivity, hypervalent bromane (III) reagents

have received much less attention compared to the corresponding iodine compounds.<sup>6</sup>

The lack of general and simple synthetic routes deliver-


**Scheme 1: a) Common aryne precursors; b) Aryne strategy to *ortho*-, *ortho*-, *meta*-substituted biaryls, c) Cyclic diaryl  $\lambda^3$ -bromanes as aryne precursors in pericyclic approach towards *ortho*-, *ortho*-, *meta*-substituted**



ing these unique scaffolds has drastically hampered the development of bromane (III) chemistry,<sup>7</sup> and thus the rare advances in this field have been reported only recently.<sup>8</sup> Our group has latterly established a simple, safe, and scalable protocol for the preparation of cyclic diaryl  $\lambda^3$ -bromanes in good to excellent yields.<sup>9</sup> The superior electron-withdrawing effect and the nucleofugality of hypervalent bromane (III) compared to the iodine (III) counterpart (approximately  $10^6$  times higher), confer them a complementary reactivity.<sup>10</sup> Encouraged by the disclosed reactivity of cyclic diaryl  $\lambda^3$ -bromanes towards the construction of carbon-oxygen and carbon-nitrogen bonds, we speculate that they could represent a new family of biaryl coupling partners for pericyclic reactions (Scheme 1c). Moreover, cyclic hypervalent  $\lambda^3$ -bromane products carry a residual functional bromine atom which opens additional perspectives for further post-functionalizations, while warranting excellent atom economy of the overall process. Thus unique reactivity provides a new strategy for the synthesis of valuable and unprecedented highly decorated dissymmetric 2,3,2',3',4'-substituted biaryls, helicenes<sup>11</sup> and four consequently substituted aromatic units.<sup>4e,12</sup>

Our investigation begun by studying the reaction between the well-known cyclic diaryl  $\lambda^3$ -iodanes and furan **2a** (Table 1, entry 2), but no product formation was observed neither by GC-MS nor <sup>1</sup>HNMR (see Supporting Information). Remarkably, while using cyclic diaryl  $\lambda^3$ -bromane **1a-OTf** and **2a**, in a presence of cesium carbonate and DCM, the desired product **3aa** was afforded in 92% yield after 16 hours at room temperature (entry 1), unambiguously illustrating the unique reactivity of these re-

**Table 1. Optimization of the Diels-Alder Reaction with cyclic diaryl hypervalent bromanes<sup>a</sup>**



entry	deviation from standard conditions	yield (%) <sup>b</sup>
1	none	92% <sup>c</sup>
2	$\lambda^3$ -I-OTf/-OMs instead of <b>1a-OTf</b>	- <sup>d</sup>
3	no Cs <sub>2</sub> CO <sub>3</sub>	- <sup>d</sup>
4	<i>t</i> BuONa / K <sub>2</sub> CO <sub>3</sub> instead to Cs <sub>2</sub> CO <sub>3</sub>	59% / 23%
5	CH <sub>3</sub> CN / H <sub>2</sub> O instead to CH <sub>2</sub> Cl <sub>2</sub>	69% / 13%


<sup>a</sup> Reaction conditions:  $\lambda^3$ -X-Y (0.1 mmol), furan (1.2 equiv), r.t. 16h; <sup>b</sup> <sup>1</sup>HNMR yields using CH<sub>2</sub>Br<sub>2</sub> as internal standard; <sup>c</sup> Isolated yield; <sup>d</sup> 5 equiv furan.

gents.

No reactivity was observed in the absence of Cs<sub>2</sub>CO<sub>3</sub> (entry 3). Further investigation showed that the nature of the base holds a key role. Despite being a stronger base, *t*BuONa provided **3aa** in lower yield, while K<sub>2</sub>CO<sub>3</sub> performed poorly (entry 4). Finally, the role of a chlorinated polar solvent, namely CH<sub>2</sub>Cl<sub>2</sub>, was confirmed (entry 5). Surprising, albeit in low yield, the reaction occurred in

water media.<sup>13</sup> With the optimal reaction conditions in hand, the scope of this cycloaddition was explored, focusing on various cyclic diaryl  $\lambda^3$ -bromanes **1a-1** and aromatic cyclic Diels-Alder partners **2b-2d** (Scheme 2). The symmetric bromane **1b**, presenting inductive electron-donating methyl groups provided the desired product **3ba** in good yield (69%). Regioselective functionalizations took place in presence of asymmetric substrates, such as **1c**, **1d**, and **1e**, where the enhanced reactivity of one aryl moieties translated into the formation of a unique products in good yields. The introduction of the methyl group in *ortho*-position directs the desired transformation onto one aromatic ring, thus promoting the formation of **3fa** in moderate yield (51%). Bifunctional dissymmetric  $\lambda^3$ -bromanes bearing methyl ester and chlorine afforded **3ga** and **3ha** in 59% and 98% yields respectively. Finally, mono-substituted cyclic diaryl  $\lambda^3$ -bromanes were tested. A weak electron-donating methyl group, **1i**, unsurprisingly leads to a 1:1 mixture of products **3ia** and **3ia'**, while the enhanced electron-withdrawing character of methyl ester **1j**, chloro **1k** and trifluoromethyl **1l** enabled the preferential formation of one regioisomer with good selectivity. Moreover, substituted furans, **2b** and **2c** and N-protected pyrrole **2d** were well tolerated under our reaction conditions, affording the desired products in good yields. Moreover, the dissymmetric bromane **1h** reacted smoothly with **2d** offering the highly functionalized bromo, chloro cycloadduct **3hd** in 84% yield. Targeting a rapid generation of a broad range of diverse molecules, we have challenged the  $\lambda^3$ -bromanes with other arynophiles. Various azides have been successfully coupled with **1a-OTf**, furnishing highly decorated bromo benzotriazole derivatives. Simple benzyl azides **2e** and substituted **2f - 2i** supplied the desired regioisomeric **3** and **3'** in moderate to good yields, which were straightforwardly separated *via* column providing the pure products. The single-crystal X-ray analysis of **3af** confirmed the triazole structure and the regioselectivity. Remarkably, thanks to the fluorine-free conditions, the silyl-protected phenols **3ai** and **3ai'** could be obtained in excellent yields, highlighting the complementarity of this protocol. The cholesterol-de derived **2j** and the cyclohexyl azide **2k** react smoothly under the standard reaction conditions. Diazomethane **2l** (TMSCHN<sub>2</sub>) also proved to be an excellent arynophile, yielding **3al** and **3cl** in 92% and 50% yields respectively, albeit as a mixture of two cycloadducts. In accordance with the aryne distorted and steric models, a moderate regiocontrol has been observed in the 1,3-dipolar cycloaddition where the nucleophilic nitrogen reacts with the more distorted alkyne terminus position furnishing the less steric demanding product as major isomer.<sup>14</sup> While Diels-Alder reactions involving aryne have been largely investigated,<sup>15</sup> the [2+2] reactions have received much less attention despite the interest of the resulting peculiar structures.<sup>16</sup> Rewardingly, the reaction of **1a-OTf** with 2,3-dihydrofuran **2m** and 2,3-dihydropyran **2n** conducted to the selective formation of **3am** and **3an** in excellent yields, furnishing a single regioisomer with a ratio up to 14.3:1. Moreover, strained cycloalkenes, namely norbornene **2o** and norbornadiene **2p**, provided the desired cyclobutene products **3ao** and **3ap** with good yields. Interestingly, an elusive [2+2+2] mechanism has been observed within **1a-OTf** and **2p** yielding the uncommon

### Scheme 3: Scope of the reaction




**3ap'** in 17% yield. The acrylate **2q** showed sufficient reactivity under our reaction conditions. Asymmetric  $\lambda^3$ -bromanes **1h**, **1c** and **1g** provided the highly decorated biaryls **3hm**, **3cn** and **3go** in excellent yields. Finally, structurally condensed polycyclic aromatic compounds, valuable motifs in material science, were targeted. We investigated the reaction of **1a-OTf** with the  $\beta$ -bromostyrenes **2r**. After a minor reoptimization of the reaction conditions, a cascade [4+2] cycloaddition/base-mediated aromatization occurred smoothly, at room temperature, in dioxane yielding the *endo*-product **3ar** as a single regioisomer. Electron-rich bromo styryls showed higher reactivity, delivering **3as** and **3ay** respectively in 85% and 58% yields. The single-crystal X-ray analysis of **3as** confirmed the formation of the *endo*-product. Electron withdrawing groups, such as trifluoromethyl, impact the reaction outcome, delivering **3at** in a decreased yield.

On the other hand, halogenated aromatic rings **2u**, **2v**, **2w** and **2x**, offered synthetically useful bifunctional products in moderate to high yields. The electron-rich, protected bromo caffeic acid derivative **2z** turned out to be suitable for the synthesis of highly decorated phenyl


phenanthrene **3az**, providing a functional dibromo compound. Sterically demanding substrate **2A** furnished the conformationally stable atropoisomers **3aA**. Finally, when dissymmetric  $\lambda^3$ -bromane **1f** was tested, a selective functionalization occurred, yielding **3fs**. Remarkably, the excellent regioselectivity reached in all of these cases, providing the sterically hindered *endo*-regioisomers as unique product, outcompetes the literature precedents.<sup>17</sup>

### Scheme 2: Synthesis of 3aa via flow-chemistry approach



Encouraged by the high versatility observed using cyclic diaryl  $\lambda^3$ -bromanes, we endeavored on developing a flow protocol for this reaction (Scheme 3). Despite clear advantages of flow chemistry approach as scalability, safety

### Scheme 4: Synthesis of functional biaryl compounds from 3aa, 3ar, 3as and 3au



and reduction of the reaction time, implementation of this process into the aryne chemistry remains rare and the chemically difficult.<sup>18</sup> The difficulty in our case lays in the use of insoluble Cs<sub>2</sub>CO<sub>3</sub>, however, attempts to replace it with organic bases failed. (see Supporting Information). Accordingly, the flow-reaction was performed in a biphasic system, which involved a water solution of Cs<sub>2</sub>CO<sub>3</sub> and a solution of **1a-OTf** and **2a** in CH<sub>3</sub>CN. The optimal flow-reaction conditions enabled the synthesis of 516 mg of **3aa** (66% yield) within 22 min of residence time at 60 °C. Interestingly, the substrate **1a-OTf** is stable under these reaction conditions and the unreacted portion was recovered *via* precipitation with Et<sub>2</sub>O from the reaction steam. The conspicuous reduction of the reaction time and the use of green solvents make this approach a suitable alternative to the batch protocol, especially for further reaction scale-up.

Beyond the mild reaction conditions for the aryne generation and the high functional group tolerance demonstrated, the cycloadduct products feature a bromine atom which can serve as an additional handle for the incorporation of molecular complexity (Scheme 4). Initially, the robustness of our procedures was confirmed by performing the reactions on 1 or 2 mmol scales, furnishing **3aa**, **3ar**, **3as** and **3au** in multi milligram scale. The functionalization of the **3aa** through lithium-mediated bromine exchange provided the corresponding carboxylic acid **4**, aldehyde **5**, and the new phosphine **6** in 64%, 93% and 76% yields respectively. Moreover, a standard deoxygenation protocol generated **7** in 74% yield. Also, bromo-biphenyl phenanthrenes **3ar**, **3as** and **3au** are synthetically interesting scaffolds and the residual bromine enables access to highly adorned structures. Pd-catalyzed cross-couplings, such as Sonogashira and Suzuki reactions granted the introduction of alkyne **8** and aryl **9** moieties with useful yields. Furthermore, a Buchwald-Hartwig Pd-catalyzed couplings delivered **10** and the sterically challenging **11** diaryl amines, in good yields. A lithium exchange protocol using **3ar**, **3as** and **3au**, led to the versatile aldehydes **12**, **13** and **14**. A subsequent Wittig

reaction involving the aldehyde **13** provided the stereoisomers *E*-**15** and *Z*-**15** in good yield (70% + 17% respectively), which were easily separated by flash chromatography. To further illustrate the utility of the bromine moiety, the synthesis of screw-shaped helicene was envisioned. A standard Corey-Fuchs alkyne synthesis using the aldehyde **14** furnished **17**. Then, a Pt-catalyzed cyclization afforded the desired [5]-helicene **18** in 12% overall yield.

In conclusion, we demonstrated herein the peculiar versatility, chemoselectivity and atom economy of cyclic diaryl λ<sup>3</sup>-bromanes as biaryl aryne precursors in cycloaddition reactions for the synthesis of highly decorated dissymmetric 2,3,2',3',4'-substituted biaryl motifs. The mild aryne generation conditions, namely carbonate base and room temperature, allowed a wide aryne compatibility, together with great functional group tolerance. Robustness of these aryne precursors guaranteed an efficient biphasic flow protocol for the pericyclic reaction further improving the sustainability. High molecular complexity *via* multiple C-C and C-N bonds formation was attained in a single reaction step, while subsequent functionalizations further expand access to a plethora of complex functional compounds.

### ASSOCIATED CONTENT

#### Supporting Information

Experimental procedures, synthesis of reagents, copies of NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

### AUTHOR INFORMATION

#### Corresponding Author

\* wenceldelord@unistra.fr

#### Author Contributions

‡These authors contributed equally.



## Notes

The authors declare no conflict of interest.

## ACKNOWLEDGMENT

We thank the CNRS (Centre National de la Recherche Scientifique), the "Ministère de l'Éducation Nationale et de la Recherche" (France) for financial support. J.W.D., M.L. and M.D.A. are very grateful to European Commission for the ERC-Starting Grant "AICHIMIE" N°949804. We are grateful to Dr. Lydia Karmazin, Dr. Corinne Bailly and Dr. Nathalie Gruber for X-ray diffraction analysis (Service de Radiocristallographie, CNRS-Université de Strasbourg, Strasbourg, France). Authors are also very grateful to Dr. Morgan Donnard, Dr. Armen Panossian and Dr. Frédéric R. Leroux for helpful discussions on flow-chemistry.

## REFERENCES

- (1) (a) Wu, P.; Nielsen, T. E.; Clausen, M. H. Small-molecule kinase inhibitors: an analysis of FDA-approved drugs. *Drug Discovery Today* **2016**, *21*, 5–10. (b) Mullard, A. 2020 FDA Drug Approvals. *Nat Rev Drug Discov* **2021**, *20*, 85–90. (c) Bhutani, P.; Joshi, G.; Raja, N.; Bachhav, N.; Rajanna, P. K.; Bhutani, H.; Paul, A. T.; Kumar, R. U.S. FDA Approved Drugs from 2015–June 2020: A Perspective. *J. Med. Chem.* **2021**, *64*, 2339–2381. (d) Jhulki, S.; Ghosh, A.; Chow, T. J.; Moorthy, J. N. Twisted Biaryl-Amines as Novel Host Materials for Green-Emissive Phosphorescent Organic Light-Emitting Diodes (PhOLEDs). *RSC Adv.* **2015**, *5*, 101169–101176. (e) Esmaeli, A. A.; Moradi, A.; Mohammadi, H. K. Regioselective Synthesis of Highly-Substituted Biaryls by Reaction of Vinyl Malononitriles with Acetylenic Esters. *Tetrahedron* **2010**, *66*, 3575–3578.
- (2) (a) Ständera, M.; Häfliger, R.; Gershoni-Poranne, R.; Stanger, A.; Jeschke, G.; van Beek, J. D.; Bertschi, L.; Schlüter, A. D. Evidence for Fully Conjugated Double-Stranded Cycles. *Chem. Eur. J.* **2011**, *17*, 12163–12174. (b) Diemer, V.; Garcia, J. S.; Leroux, F. R.; Colobert, F. Aryne-Mediated Fluorination: Synthesis of Fluorinated Biaryls via a Sequential Desilylation–Halide Elimination–Fluoride Addition Process. *J. Fluorine Chem.* **2012**, *134*, 146–155. (c) Neudorff, W. D.; Schulte, N.; Lentz, D.; Schlüter, A. D. 8,9-Didehydrofluoranthenes as Building Blocks for the Synthesis of Extended Polycyclic Aromatic Hydrocarbons (PAHs). *Org. Lett.* **2001**, *3*, 3115–3118. (d) Akai, S.; Ikawa, T.; Takayanagi, S.; Morikawa, Y.; Mohri, S.; Tsubakiyama, M.; Egi, M.; Wada, Y.; Kita, Y. Synthesis of Biaryl Compounds through Three-Component Assembly: Ambidentate Effect of the Tert-Butyldimethylsilyl Group for Regioselective Diels-Alder and Hiyama Coupling Reactions. *Angew. Chem. Int. Ed.* **2008**, *47*, 7673–7676. (e) Moreira, B.; Muraca, A.; Raminelli, C. Synthesis of Silylbiaryl Triflates by Chemoselective Suzuki Reaction. *Synthesis* **2016**, *49*, 1093–1102. (f) Asgari, P.; Dakarapu, U. S.; Nguyen, H. H.; Jeon, J. Aryne Cycloaddition Reactions of Benzodioxasilines as Aryne Precursors Generated by Catalytic Reductive Ortho-C-H Silylation of Phenols with Traceless Acetal Directing Groups. *Tetrahedron* **2017**, *73*, 4052–4061. (g) Wei, Y.-L.; Dauvergne, G.; Rodriguez, J.; Coquerel, Y. Enantiospecific Generation and Trapping Reactions of Aryne Atropisomers. *J. Am. Chem. Soc.* **2020**, *142*, 16921–16925. (h) Werz, D. B.; Biju, A. T. Uncovering the Neglected Similarities of Arynes and Donor–Acceptor Cyclopropanes. *Angew. Chem. Int. Ed.* **2020**, *59*, 3385–3398.
- (3) (a) Shi, J.; Li, L.; Li, Y. O-Silylaryl Triflates: A Journey of Kobayashi Aryne Precursors. *Chem. Rev.* **2021**, *121*, 3892–4022. (b) Sarmah, M.; Sharma, A.; Gogoi, P. Exploration of Kobayashi's Aryne Precursor: A Potent Reactive Platform for the Synthesis of Polycyclic Aromatic Hydrocarbons. *Org. Biomol. Chem.* **2021**, *19*, 722–737. (c) Idiris, F. I. M.; Jones, C. R. Recent Advances in Fluoride-Free Aryne Generation from Arene Precursors. *Org. Biomol. Chem.* **2017**, *15*, 9044–9056. (d) Roy, T.; Biju, A. T. Recent Advances in Molecular Rearrangements Involving Aryne Intermediates. *Chem. Commun.* **2018**, *54*, 2580–2594.
- (4) (a) Sundalam, S. K.; Nilova, A.; Seidl, T. L.; Stuart, D. R. A Selective C–H Deprotonation Strategy to Access Functionalized Arynes by Using Hypervalent Iodine. *Angew. Chem. Int. Ed.* **2016**, *55*, 8431–8434. (b) Chen, H.; Han, J.; Wang, L. Diels–Alder Cycloadditions of N-Arylpyrroles via Aryne Intermediates Using Diaryliodonium Salts. *Beilstein J. Org. Chem.* **2018**, *14*, 354–363. (c) Zhang, Z.; Wu, X.; Han, J.; Wu, W.; Wang, L. Direct Arylation of Tertiary Amines via Aryne Intermediates Using Diaryliodonium Salts. *Tetrahedron Letters* **2018**, *59*, 1737–1741. (d) Nilova, A.; Metzke, B.; Stuart, D. R. Aryl(TMP)Iodonium Tosylate Reagents as a Strategic Entry Point to Diverse Aryl Intermediates: Selective Access to Arynes. *Org. Lett.* **2021**, *23*, 4813–4817. (e) Nilova, A.; Sibbald, P. A.; Valente, E. J.; González-Montiel, G. A.; Richardson, H. C.; Brown, K. S.; Cheong, P. H.; Stuart, D. R. Regioselective Synthesis of 1,2,3,4-Tetrasubstituted Arenes by Vicinal Functionalization of Arynes Derived from Aryl(Mes)Iodonium Salts. *Chem. Eur. J.* **2021**, *27*, 7168–7175.
- (5) (a) Heinz, B.; Djukanovic, D.; Filipponi, P.; Martin, B.; Paraghiosoff, K.; Knochel, P. Regioselective Difunctionalization of Pyridines via 3,4-Pyridynes. *Chem. Sci.* **2021**, *12*, 6143–6147. (b) Lin, W.; Sapountzis, I.; Knochel, P. Preparation of Functionalized Aryl Magnesium Reagents by the Addition of Magnesium Aryl Thiolates and Amides to Arynes. *Angew. Chem. Int. Ed.* **2005**, *44*, 4258–4261. (c) Sapountzis, I.; Lin, W.; Fischer, M.; Knochel, P. Preparation of Polyfunctional Arynes via 2-Magnesiated Diaryl Sulfonates. *Angew. Chem. Int. Ed.* **2004**, *43*, 4364–4366.
- (6) (a) Wirth, T. Hypervalent Iodine Chemistry in Synthesis: Scope and New Directions. *Angew. Chem. Int. Ed.* **2005**, *44*, 3656–3665. (c) Ochiai, M. Hypervalent Aryl-, Alkynyl-, and Alkenyl- $\lambda^3$ -Bromanes. *Synlett* **2009**, *2009*, 159–173. (d) Bhojgude, S. S.; Thangaraj, M.; Suresh, E.; Biju, A. T. *Org. Lett.* **2014**, *16*, 3576–3579.
- (7) Ochiai, M.; Miyamoto, K.; Kaneaki, T.; Hayashi, S.; Nakanishi, W. Highly Regioselective Amination of Unactivated Alkanes by Hypervalent Sulfonylimino- $\lambda^3$ -Bromane. *Science* **2011**, *332*, 448–451.
- (8) (a) Yoshida, Y.; Ishikawa, S.; Mino, T.; Sakamoto, M. Bromonium Salts: Diaryl- $\lambda^3$ -Bromanes as Halogen-Bonding Organocatalysts. *Chem. Commun.* **2021**, *57*, 2519–2522. (b) Sokolovs, I.; Mohebbati, N.; Francke, R.; Suna, E. Electrochemical Generation of Hypervalent Bromine(III) Compounds. *Angew. Chem. Int. Ed.* **2021**, *60*, 15832–15837. (c) Miyamoto, K.; Saito, M.; Tsuji, S.; Takagi, T.; Shiro, M.; Uchiyama, M.; Ochiai, M. Benchtop-Stable Hypervalent Bromine(III) Compounds: Versatile Strategy and Platform for Air- and Moisture-Stable  $\lambda^3$ -Bromanes. *J. Am. Chem. Soc.* **2021**, *143*, 9327–9331.
- (9) Lanzi, M.; Dherbassy, Q.; Wencel-Delord, J. Cyclic Diaryl  $\lambda^3$ -Bromanes as Original Aryne Precursors. *Angew. Chem. Int. Ed.* **2021**, *60*, 14852–14857.
- (10) (a) Tolstaya, T. P.; Demkina, I. I.; Grushin, V. V.; Vanchikov, A. N. Nucleophilic Substitution in Diphenylbromonium and Diphenyl-chloronium Salts. *J. Org. Chem. USSR* **1989**, *25*, 2305–2310. (b) Nakajima, M.; Miyamoto, K.; Hirano, K.; Uchiyama, M. Diaryl- $\lambda^3$ -chloranes: versatile synthesis and unique reactivity as aryl cation equivalent. *J. Am. Chem. Soc.* **2019**, *141*, 6499–6503.
- (11) Shen, Y.; Chen, C.-F. Helicenes: Synthesis and Applications. *Chem. Rev.* **2012**, *112*, 1463–1535.
- (12) Biju, A. T. *Modern Aryne Chemistry*, Wiley, **2021**.
- (13) Yoshimura, A.; Fuchs, J. M.; Middleton, K. R.; Maskavev, A. V.; Rohde, G. T.; Saito, A.; Postnikov, P. S.; Yusubov, M. S.; Nemykin, V. N.; Zhdankin, V. V. Pseudocyclic Arylbenziodoxaboroles: Efficient Benzyne Precursors Triggered by Water at Room Temperature. *Chem. Eur. J.* **2017**, *23*, 16738–16742.
- (14) (a) Medina, J. M.; Mackey, J. L.; Garg, N. K.; Houk, K. N. The Role of Aryne Distortions, Steric Effects, and Charges in Regioselectivities of Aryne Reactions. *J. Am. Chem. Soc.* **2014**, *136*, 15798–15805. (b) Bronner, S. M.; Mackey, J. L.; Houk, K. N.; Garg, N. K. Steric Effects Compete with Aryne Distortion To Control

Regioselectivities of Nucleophilic Additions to 3-Silylarynes. *J. Am. Chem. Soc.* **2012**, *134*, 13966–13969.

(15) Bhojgude, S. S.; Bhunia, A.; Biju, A. T. Employing Arynes in Diels–Alder Reactions and Transition-Metal-Free Multicomponent Coupling and Arylation Reactions. *Acc. Chem. Res.* **2016**, *49*, 1658–1670.

(16) (a) Gilmore, C. D.; Allan, K. M.; Stoltz, B. M. Orthogonal Synthesis of Indolines and Isoquinolines via Aryne Annulation. *J. Am. Chem. Soc.* **2008**, *130*, 1558–1559. (b) Chen, Z.; Han, X.; Liang, J.-H.; Yin, J.; Yu, G.-A.; Liu, S.-H. *Chin. Chem. Lett.* **2014**, *25*, 1535–1539. (c) Yedulla, V. R.; Pradhan, P.; Yang, L.; Lakshman, M. K. Cycloaddition of Arynes and Cyclic Enol Ethers as a Platform for Access to Stereochemically Defined 1,2-Disubstituted Benzocyclobutenes: Cycloaddition of Arynes and Cyclic Enol Ethers. *Eur. J. Org. Chem.* **2015**, *2015*, 750–764.

(17) (a) Singh, V.; Verma, R. S.; Khatana, A. K.; Tiwari, B. Construction of Phenanthrenes and Chrysenes from  $\beta$ -

Bromovinylarenes via Aryne Diels–Alder Reaction/Aromatization. *J. Org. Chem.* **2019**, *84*, 14161–14167. (b) Bhojgude, S. S.; Bhunia, A.; Gonnade, R. G.; Biju, A. T. Efficient Synthesis of 9-Aryldihydrophenanthrenes by a Cascade Reaction Involving Arynes and Styrenes. *Org. Lett.* **2014**, *16*, 676–679.

(18) (a) García-Lacuna, J.; Domínguez, G.; Pérez-Castells, J. Flow Chemistry for Cycloaddition Reactions. *ChemSusChem* **2020**, *13*, 5138–5163. (b) Guidi, M.; Seeberger, P. H.; Gilmore, K. How to Approach Flow Chemistry. *Chem. Soc. Rev.* **2020**, *49*, 8910–8932. (c) Michel, B.; Greaney, M. F. Continuous-Flow Synthesis of Trimethylsilylphenyl Perfluorosulfonate Benzyne Precursors. *Org. Lett.* **2014**, *16*, 2684–2687. (d) Kleoff, M.; Boeser, L.; Baranyi, L.; Heretsch, P. Scalable Synthesis of Benzotriazoles via [3+2] Cycloaddition of Azides and Arynes in Flow. *Eur. J. Org. Chem.* **2021**, *2021*, 979–982. (17)

---