



HAL
open science

Full discretization of time dependent convection-diffusion-reaction equation coupled with the Darcy system

Nancy Chalhoub, Pascal Omnes, Toni Sayah, Rebecca El Zahlaniyeh

► **To cite this version:**

Nancy Chalhoub, Pascal Omnes, Toni Sayah, Rebecca El Zahlaniyeh. Full discretization of time dependent convection-diffusion-reaction equation coupled with the Darcy system. *Calcolo*, 2020, 57 (1), pp.4. 10.1007/s10092-019-0352-1 . hal-03878148

HAL Id: hal-03878148

<https://hal.science/hal-03878148v1>

Submitted on 29 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

FULL DISCRETIZATION OF TIME DEPENDENT CONVECTION-DIFFUSION-REACTION EQUATION COUPLED WITH THE DARCY SYSTEM.

NANCY CHALHOUB[†], PASCAL OMNES^{‡*}, TONI SAYAH[†], AND REBECCA EL ZAHLANIYEH^{†*}

ABSTRACT. In this article, we study the time dependent convection-diffusion-reaction equation coupled with the Darcy equation. We propose and analyze two numerical schemes based on finite element methods for the discretization in space and the implicit Euler method for the discretization in time. An optimal *a priori* error estimate is then derived for each numerical scheme. Finally, we present some numerical experiments that confirm the theoretical accuracy of the discretization.

KEYWORDS. Darcy's equations; convection-diffusion-reaction equation; finite element method; *a priori* error estimates.

1. INTRODUCTION.

Let Ω be a connected bounded open set in \mathbb{R}^d , $d = 2, 3$, with a Lipschitz-continuous boundary $\Gamma = \partial\Omega$, and let $[0, T]$ be an interval of \mathbb{R} . In this work, we study the concentration distribution of a fluid in a porous medium modelled by a time dependent convection-diffusion-reaction equation coupled with Darcy's law. The system of equations is

$$(P) \left\{ \begin{array}{ll} \nu(C(\mathbf{x}, t))\mathbf{u}(\mathbf{x}, t) + \nabla p(\mathbf{x}, t) & = \mathbf{f}(\mathbf{x}, t, C(\mathbf{x}, t)) \quad \text{in } \Omega \times]0, T[, \\ (\operatorname{div} \mathbf{u})(\mathbf{x}, t) & = 0 \quad \text{in } \Omega \times]0, T[, \\ \frac{\partial C}{\partial t}(\mathbf{x}, t) - \alpha \Delta C(\mathbf{x}, t) + (\mathbf{u}(\mathbf{x}, t) \cdot \nabla C)(\mathbf{x}, t) + r_0 C(\mathbf{x}, t) & = g(\mathbf{x}, t) \quad \text{in } \Omega \times]0, T[, \\ (\mathbf{u} \cdot \mathbf{n})(\mathbf{x}, t) & = 0 \quad \text{on } \Gamma \times [0, T], \\ C(\mathbf{x}, t) & = 0 \quad \text{on } \Gamma \times [0, T], \\ C(\mathbf{x}, 0) & = 0 \quad \text{in } \Omega, \end{array} \right.$$

where \mathbf{n} is the unit outward normal vector on Γ . The unknowns are the velocity \mathbf{u} , the pressure p and the concentration C of the fluid. The function \mathbf{f} represents a force density that depends on the concentration C and the function g represents an external concentration source. The viscosity ν also depends on the concentration C but the diffusion coefficient α and the parameter r_0 are positive constants. To simplify, a homogeneous Dirichlet boundary condition is prescribed on the concentration C , but the present analysis easily extends to a non homogeneous boundary condition.

The existence of a weak solution of (P) is established in [12, 8]. As far as its numerical approximation is concerned, problem (P) was treated for example in [16, 17]. In these works, the authors used the semi-implicit Euler method for the time discretization and the Raviart-Thomas $H(\operatorname{div}, \Omega)$ finite element

November 29, 2022.

[†] Laboratoire de Mathématiques et Applications, Unité de recherche Mathématiques et Modélisation, Faculté des Sciences, Université Saint-Joseph, B.P 11-514 Riad El Solh, Beyrouth 1107 2050, Liban.

E-mails: nancy.chalhoub@usj.edu.lb, rebecca.zahlaniyeh@net.usj.edu.lb, toni.sayah@usj.edu.lb.

[‡] DEN-Service de thermohydraulique et de mécanique des fluides (STMF), CEA, Université Paris-Saclay, F-91191, Gif-sur-Yvette, France.

E-mail : pascal.omnes@cea.fr.

* Université Paris 13, LAGA, CNRS UMR 7539, Institut Galilée, 99 Avenue J.B. Clément F-93430 Villetaneuse Cedex, France.

E-mail: elzahlaniyeh@math.univ-paris13.fr.

method for the space discretization of the velocity/pressure unknowns. They established an *a priori* error estimate that is valid in 2-dimensions and place themselves in a somewhat restrictive framework, where Ω is a square, discretized by square cells. Moreover, in order to derive the estimate, the authors imposed multiple conditions on the space and time steps, which lead to a time step $\Delta t = o(h)$ for the lowest-order Raviart-Thomas scheme. The heat equation coupled with the Navier-Stokes system has been treated by many works (see for instance Bernardi, Métivet and Pernaud-Thomas [4], Deteix, Jendoubi and Yakoubi [11], or Gaultier and Lezaun [13]). The stationary coupling of Darcy's system with the heat equation where the viscosity is constant but the exterior force depends on the temperature (like in the model proposed by Boussinesq [7]) was analyzed by Bernardi, Yacoubi and Maarouf [6] and discretized with a spectral method. The same stationary system but coupled by a nonlinear viscosity depending on the temperature is studied by Bernadi et al. in [5], where they propose and analyze two numerical schemes based on finite element methods. In [20, 19], the authors discretize a problem similar to (P) using Raviart-Thomas elements methods for the discretization in space. For the discretization in time, Rivière and Walkington in [20] used the Discontinuous Galerkin (DG) method and Li *et al.* in [19] used the Interior Penalty Discontinuous Galerkin (IPDG) method. These two works do not establish *a priori* error estimates but rather prove the convergence of the schemes using compactness results for functions that may be discontinuous in time. Vassilev and Yotov coupled in [23] the non-stationary Stokes-Darcy equation with the time dependent Transport equation, and established an *a priori* error estimate.

In this work, we study two types of discrete schemes for the full discretization of Problem (P) in time and space and for both, we prove existence and uniqueness and derive optimal *a priori* error estimates for the solutions. The first scheme is the lowest order Raviart-Thomas scheme for the velocity/pressure unknowns, for which we extend and improve the results of [16, 17], first by considering general domains covered by simplicial meshes, then by extending the proof to the three-dimensional case.

The second scheme uses the \mathbb{P}_1 -bubble / \mathbb{P}_1 scheme for the velocity/pressure unknowns, which is known as the "mini-element" in the Stokes context [2]. Both schemes use the \mathbb{P}_1 scheme for the concentration unknown. Finally, we perform several numerical tests to validate the theoretical results. They show that, in certain circumstances, the second scheme, although of higher numerical complexity, may have a better accuracy/complexity ratio.

The outline of the paper is as follows:

- In Section 2, we introduce some notations and functional spaces that are useful for the study of the problem.
- In section 3, we introduce two variational formulations.
- Section 4 is devoted to the study of two numerical schemes and the establishment of an *a priori* error estimation under regularity assumptions of the exact solutions.
- Some numerical experiments are presented in Section 5.

2. PRELIMINARIES

In this section, we recall the main notations and results which we use later on. We introduce the Sobolev space

$$W^{m,r}(\Omega)^d = \{ \mathbf{v} \in L^r(\Omega)^d; \partial^k \mathbf{v} \in L^r(\Omega)^d, \forall |k| \leq m \},$$

where $k = \{k_1, \dots, k_d\}$ is a vector of non negative integers, such that $|k| = k_1 + \dots + k_d$ and

$$\partial^k \mathbf{v} = \frac{\partial^{|k|} \mathbf{v}}{\partial^{k_1} x_1 \dots \partial^{k_d} x_d}.$$

This space is equipped with the semi-norm

$$|\mathbf{v}|_{W^{m,r}(\Omega)^d} = \left(\sum_{|k|=m} \int_{\Omega} |\partial^k \mathbf{v}|^r d\mathbf{x} \right)^{\frac{1}{r}},$$

and is a Banach space for the norm

$$\| \mathbf{v} \|_{W^{m,r}(\Omega)^d} = \left(\sum_{l=0}^m \int_{\Omega} |\mathbf{v}|_{W^{l,r}(\Omega)^d}^r d\mathbf{x} \right)^{\frac{1}{r}}.$$

When $r = 2$, this space is the Hilbert space $H^m(\Omega)^d$. In particular, we consider the following spaces

$$H_0^1(\Omega)^d = \{ \mathbf{v} \in H^1(\Omega)^d; \mathbf{v}|_{\partial\Omega} = 0 \},$$

and its dual $H^{-1}(\Omega)^d$.

We shall also introduce

$$L_0^2(\Omega) = \left\{ q \in L^2(\Omega); \int_{\Omega} q(\mathbf{x}) d\mathbf{x} = 0 \right\}.$$

We define the following scalar product in $L^2(\Omega)$:

$$(v, w) = \int_{\Omega} v(\mathbf{x})w(\mathbf{x})d\mathbf{x}, \quad \forall v, w \in L^2(\Omega).$$

We recall the following Poincaré and Sobolev inequalities:

Lemma 2.1. *For any $p \geq 1$ when $d = 1$ or 2 , or $1 \leq p \leq \frac{2d}{d-2}$ when $d \geq 3$, there exist two positive constants S_p and S_p^0 such that*

$$\forall \mathbf{v} \in H_0^1(\Omega)^d, \quad \| \mathbf{v} \|_{L^p(\Omega)^d} \leq S_p^0 \| \mathbf{v} \|_{H_0^1(\Omega)^d},$$

and

$$\forall \mathbf{v} \in H^1(\Omega)^d, \quad \| \mathbf{v} \|_{L^p(\Omega)^d} \leq S_p \| \mathbf{v} \|_{H^1(\Omega)^d}.$$

We shall also use the following continuous embedding:

$$\forall q > d, \quad W^{1,q}(\Omega) \hookrightarrow L^\infty(\Omega). \quad (2.1)$$

We recall the standard spaces for Darcy's equations

$$H(\operatorname{div}, \Omega) = \{ \mathbf{v} \in L^2(\Omega)^d; \operatorname{div} \mathbf{v} \in L^2(\Omega) \}, \quad (2.2)$$

$$H_0(\operatorname{div}, \Omega) = \{ \mathbf{v} \in H(\operatorname{div}, \Omega); (\mathbf{v} \cdot \mathbf{n})|_{\Gamma} = 0 \}, \quad (2.3)$$

and

$$\mathcal{V} = \{ \mathbf{v} \in H_0(\operatorname{div}, \Omega); \operatorname{div} \mathbf{v} = 0 \}, \quad (2.4)$$

equipped with the norm

$$\| \mathbf{v} \|_{H(\operatorname{div}, \Omega)}^2 = \| \mathbf{v} \|_{L^2(\Omega)^d}^2 + \| \operatorname{div} \mathbf{v} \|_{L^2(\Omega)}^2. \quad (2.5)$$

Finally, we recall the inf-sup condition between $L_0^2(\Omega)$ and $H_0(\operatorname{div}, \Omega)$,

$$\inf_{q \in L_0^2(\Omega)} \sup_{\mathbf{v} \in H_0(\operatorname{div}, \Omega)} \frac{\int_{\Omega} (\operatorname{div} \mathbf{v}) q d\mathbf{x}}{\| \mathbf{v} \|_{H(\operatorname{div}, \Omega)} \| q \|_{L^2(\Omega)}} \geq \beta, \quad (2.6)$$

with a constant $\beta > 0$, and the inf-sup condition between $H^1(\Omega) \cap L_0^2(\Omega)$ and $L^2(\Omega)^d$,

$$\inf_{q \in H^1(\Omega) \cap L_0^2(\Omega)} \sup_{\mathbf{v} \in L^2(\Omega)^d} \frac{\int_{\Omega} \mathbf{v} \cdot \nabla q d\mathbf{x}}{\| \mathbf{v} \|_{L^2(\Omega)^d} \| q \|_{H^1(\Omega)}} \geq 1. \quad (2.7)$$

Condition (2.6) follows immediately by solving a Laplace equation in Ω with a Neumann boundary condition on Γ , and condition (2.7) by choosing $\mathbf{v} = \nabla q$. As usual, for handling time-dependent problems, it is convenient to consider functions defined on a time interval $]a, b[$ with values in a separable functional space W equipped with a norm $\| \cdot \|_W$. Then, for any $r \geq 1$, we introduce the space

$$L^r(a, b; W) = \left\{ f \text{ measurable on }]a, b[; \int_a^b \| f(t) \|_W^r dt < \infty \right\};$$

equipped with the norm

$$\|f\|_{L^r(a,b;W)} = \left(\int_a^b \|f(t)\|_W^r dt \right)^{\frac{1}{r}}.$$

If $r = \infty$, then

$$L^\infty(a,b;W) = \left\{ f \text{ measurable on }]a,b[; \sup_{t \in [a,b]} \|f(t)\|_W < \infty \right\}.$$

Remark 2.2. $L^r(0,T;W)$ is a Banach space if W is a Banach space.

In addition, we define $C^j(0,T;W)$ as the space of functions C^j in time with values in W .

Remark 2.3. Let a and b be two real numbers.

(1) For any positive real number ε , we have

$$ab \leq \frac{1}{2\varepsilon}a^2 + \frac{1}{2}\varepsilon b^2. \quad (2.8)$$

(2) We also have

$$a(a-b) = \frac{1}{2}a^2 - \frac{1}{2}b^2 + \frac{1}{2}(a-b)^2. \quad (2.9)$$

3. VARIATIONAL FORMULATIONS

In this section, we start by writing a variational formulation of problem (P). Next, we prove the existence and the uniqueness of the solution. We assume that the data of the problem verify the following assumptions:

Assumption 3.1. We assume that the data \mathbf{f} , g and ν verify:

(1) \mathbf{f} can be written as follows:

$$\mathbf{f}(\mathbf{x}, t, C) = \mathbf{f}_0(\mathbf{x}, t) + \mathbf{f}_1(\mathbf{x}, C), \quad (3.1)$$

where $\mathbf{f}_0 \in L^\infty(0,T;L^2(\Omega)^d)$ and $\mathbf{f}_1(C)$ is (uniformly in \mathbf{x}) $c_{\mathbf{f}_1}^*$ -Lipschitz with respect to its second variable with values in \mathbb{R}^d . In addition, we suppose that

$$\forall \mathbf{x} \in \Omega, \forall \xi \in \mathbb{R}, |\mathbf{f}_1(\mathbf{x}, \xi)| \leq c_{\mathbf{f}_1} |\xi|, \quad (3.2)$$

where $c_{\mathbf{f}_1}$ is a positive constant.

(2) $g \in L^2(0,T,L^2(\Omega))$.

(3) ν is λ -Lipschitz on \mathbb{R} and there exist two strictly positive constants ν_1 and ν_2 such that, for any $\theta \in \mathbb{R}$

$$\nu_1 \leq \nu(\theta) \leq \nu_2. \quad (3.3)$$

There are two possible choices of spaces for Darcy's velocity and pressure (\mathbf{u}, p) . The first choice is $L^\infty(0,T;H_0(\text{div}, \Omega)) \times L^\infty(0,T;L_0^2(\Omega))$; it corresponds to a mixed formulation and is analyzed in this section. The second choice is $L^\infty(0,T;L^2(\Omega)^d) \times L^\infty(0,T;(H^1(\Omega)) \cap L_0^2(\Omega))$; it leads to an alternative formulation equivalent to the first. In both cases, the concentration C is in $L^2(0,T;H_0^1(\Omega))$. Then, whereas there is no difficulty in setting Darcy's system in variational form, a variational formulation of the concentration equation is not that obvious. Indeed, the convection term $\mathbf{u} \cdot \nabla C$ cannot be tested by an $H^1(\Omega)$ function, since it is only in $L^1(\Omega)$. Therefore, we choose the test functions in $H_0^1(\Omega) \cap L^\infty(\Omega)$.

where c is positive constant independent of h and m .

Proof. It is clear that the first equation of Problem $(V_{h,1})$ has a unique solution (\mathbf{u}_h^n, p_h^n) as a consequence of the coerciveness of the corresponding bilinear form on $\mathcal{W}_{h,1} \times \mathcal{W}_{h,1}$ and the inf-sup condition (4.11). Thus, knowing $\mathbf{u}_h^n \in \mathcal{W}_{h,1}$ and $C_h^{n-1} \in X_h$, the third equation of Problem $(V_{h,1})$ also admits a unique solution $C_h^n \in X_h$. Therefore, by taking $\mathbf{v}_h = \mathbf{u}_h^n$ in the first equation we get (4.13), and $S_h = C_h^n$ in the third equation of Problem $(V_{h,1})$ we get, using the Cauchy-Schwarz inequality, Lemma 2.1 and Remark 2.3

$$\frac{1}{2}(\|C_h^n\|_{L^2(\Omega)}^2 - \|C_h^{n-1}\|_{L^2(\Omega)}^2) + \frac{\alpha}{2}\tau\|C_h^n\|_{H^1(\Omega)}^2 + \tau r_0\|C_h^n\|_{L^2(\Omega)}^2 \leq \frac{(S_2^0)^2}{2\alpha}\tau\|g^n\|_{L^2(\Omega)}^2. \quad (4.15)$$

We sum over $n = 1, \dots, m$ and we obtain (4.14).

4.2. Second discrete scheme. Let K be an element of \mathcal{T}_h with vertices a_i , $1 \leq i \leq d+1$, and corresponding barycentric coordinates λ_i . We denote by $b_K \in \mathbb{P}_{d+1}(K)$ the basic bubble function

$$b_K(\mathbf{x}) = \lambda_1(\mathbf{x}) \dots \lambda_{d+1}(\mathbf{x}). \quad (4.16)$$

We observe that $b_K(\mathbf{x}) = 0$ on ∂K and that $b_K(\mathbf{x}) > 0$ in the interior of K .

Let $(\mathcal{W}_{h,2}, M_{h,2})$ be a pair of discrete spaces approximating $L^2(\Omega)^d \times (H^1(\Omega) \cap L_0^2(\Omega))$ defined by

$$\mathcal{W}_{h,2} = \{\mathbf{v}_h \in (C^0(\bar{\Omega}))^d; \forall K \in \mathcal{T}_h, \mathbf{v}_h|_K \in \mathcal{P}(K)^d\}, \quad (4.17)$$

$$\tilde{M}_h = \{q_h \in C^0(\bar{\Omega}); \forall K \in \mathcal{T}_h, q_h|_K \in \mathbb{P}_1(K)\} \quad \text{and} \quad M_{h,2} = \tilde{M}_h \cap L_0^2(\Omega), \quad (4.18)$$

where

$$\mathcal{P}(K) = \mathbb{P}_1(K) \oplus \text{Vect}\{b_K\},$$

is the space associated to the discretisation in space by the ‘‘mini-élément’’ introduced by Arnold *et al.* in [2].

Let $\mathcal{V}_{h,2}$ be the kernel of the divergence in $\mathcal{W}_{h,2}$,

$$\mathcal{V}_{h,2} = \{\mathbf{v}_h \in \mathcal{W}_{h,2}; \forall q_h \in M_{h,2}, \int_{\Omega} \mathbf{v}_h \cdot \nabla q_h \, d\mathbf{x} = 0\}. \quad (4.19)$$

We shall use a variant of R_h denoted by \mathcal{F}_h which is constructed in [5, p. 336] and has the following properties:

$$\forall \mathbf{v} \in H^1(\Omega)^d, \|\mathbf{v} - \mathcal{F}_h(\mathbf{v})\|_{L^2(K)^d} \leq Ch\|\mathbf{v}\|_{H^1(\Delta_K)^d}, \quad (4.20)$$

and $\mathcal{F}_h(\mathbf{v}) \in \mathcal{V}_{h,2}$ when $\text{div } \mathbf{v} = 0$.

Regarding the pressure, since Z_h coincides with \tilde{M}_h , an easy modification of R_h yields an operator r_h in $\mathcal{L}(H^1(\Omega); \tilde{M}_h)$ and in $\mathcal{L}(H^1(\Omega) \cap L_0^2(\Omega); M_{h,2})$ (see for instance Abboud, Girault and Sayah [1]), satisfying (4.2). We approximate problem (V_a) by the following discrete scheme:

$$(V_{h,2}) \left\{ \begin{array}{l} \text{Having } C_h^{n-1} \in X_h, \text{ Find } (\mathbf{u}_h^n, p_h^n) \in \mathcal{W}_{h,2} \times M_{h,2} \text{ such as} \\ \forall \mathbf{v}_h \in \mathcal{W}_{h,2}, \int_{\Omega} \nu(C_h^{n-1})\mathbf{u}_h^n \cdot \mathbf{v}_h \, d\mathbf{x} + \int_{\Omega} \nabla p_h^n \cdot \mathbf{v}_h \, d\mathbf{x} = \int_{\Omega} \mathbf{f}^n(C_h^{n-1}) \cdot \mathbf{v}_h \, d\mathbf{x}, \\ \forall q_h \in M_{h,2}, \int_{\Omega} \nabla q_h \cdot \mathbf{u}_h^n \, d\mathbf{x} = 0, \\ \text{Having } C_h^{n-1} \in X_h, \text{ Find } C_h^n \in X_h \text{ such that} \\ \forall S_h \in X_h, \int_{\Omega} \frac{C_h^n - C_h^{n-1}}{\tau} S_h \, d\mathbf{x} + \alpha \int_{\Omega} \nabla C_h^n \cdot \nabla S_h \, d\mathbf{x} + \int_{\Omega} (\mathbf{u}_h^n \cdot \nabla C_h^n) S_h \, d\mathbf{x} \\ \quad + \frac{1}{2} \int_{\Omega} \text{div}(\mathbf{u}_h^n) C_h^n S_h \, d\mathbf{x} + r_0 \int_{\Omega} C_h^n S_h \, d\mathbf{x} = \int_{\Omega} g^n S_h \, d\mathbf{x}. \end{array} \right.$$

where as usual, the second nonlinear term in the last equation is added to compensate the fact that $\text{div } \mathbf{u}_h^n \neq 0$. It is well-known that Green’s formula and the functions regularity imply that

$$\int_{\Omega} (\mathbf{u}_h^n \cdot \nabla C_h^n) S_h \, d\mathbf{x} + \frac{1}{2} \int_{\Omega} (\text{div } \mathbf{u}_h^n) C_h^n S_h \, d\mathbf{x} = \frac{1}{2} \left(\int_{\Omega} (\mathbf{u}_h^n \cdot \nabla C_h^n) S_h \, d\mathbf{x} - \int_{\Omega} (\mathbf{u}_h^n \cdot \nabla S_h) C_h^n \, d\mathbf{x} \right), \quad (4.21)$$

so that the nonlinear term is antisymmetric. One of the key points for studying $(V_{h,2})$ is the discrete inf-sup condition satisfied by the pair of spaces $(\mathcal{W}_{h,2}, M_{h,2})$ (see for instance [5]):

$$\forall q_h \in M_{h,2}, \sup_{\mathbf{v}_h \in \mathcal{W}_{h,2}} \frac{\int_{\Omega} \nabla q_h \cdot \mathbf{v}_h \, d\mathbf{x}}{\|\mathbf{v}_h\|_{L^2(\Omega)^d}} \geq \beta_2 |q_h|_{H^1(\Omega)}, \quad (4.22)$$

with a constant $\beta_2 > 0$ independent of h .

By following the same steps of the proof of Theorem (4.1), we deduce the following theorem:

Theorem 4.2 (Existence and uniqueness of the solution of $(V_{h,2})$). *At each time step n and for a given $C_h^{n-1} \in X_h$, Problem $(V_{h,2})$ has a unique solution $(\mathbf{u}_h^n, p_h^n, C_h^n) \in \mathcal{W}_{h,2} \times M_{h,2} \times X_h$ which verifies, for $m = 1, \dots, N$, the following bounds*

$$\|\mathbf{u}_h^m\|_{L^2(\Omega)^d} \leq \frac{1}{\nu_1} (\|\mathbf{f}_0\|_{L^\infty(0,T;L^2(\Omega)^d)} + c_{\mathbf{f}_1} \|C_h^{m-1}\|_{L^2(\Omega)}) \quad (4.23)$$

and

$$\|C_h^m\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^m \tau |C_h^n|_{H^1(\Omega)}^2 + 2r_0 \sum_{n=1}^m \tau \|C_h^n\|_{L^2(\Omega)}^2 \leq \frac{(S_2^0)^2}{\alpha} \|g\|_{L^2(0,T;L^2(\Omega))}^2, \quad (4.24)$$

where c is positive constant independent of h and m .

5. A PRIORI ERROR ESTIMATE

In this section, we establish the *a priori* estimates corresponding to the proposed numerical schemes. We begin by establishing the error estimates corresponding to the velocity and the pressure, and then we will establish those corresponding to the concentration for both schemes $(V_{h,1})$ and $(V_{h,2})$.

In all the rest of the paper, we denote by $\mathbf{u}^n = \mathbf{u}(t_n)$, $p^n = p(t_n)$ and $C^n = C(t_n)$.

Theorem 5.1. *Let (\mathbf{u}, p, C) be the solution of Problem (V) and $(\mathbf{u}_h^n, p_h^n, C_h^n)$ be the solution of Problem $(V_{h,1})$. If $\mathbf{u} \in L^\infty(0, T; H^1(\Omega)^d) \cap L^\infty(0, T; L^\infty(\Omega)^d)$, $\frac{\partial \mathbf{u}}{\partial t} \in L^2(0, T; L^4(\Omega)^d)$, $p \in L^\infty(0, T; H^1(\Omega))$, $C \in L^\infty(0, T; W^{2,4}(\Omega))$ and $\frac{\partial C}{\partial t} \in L^2(0, T; W^{1,4}(\Omega))$ and under Assumption 3.1, there exist positive constants c, c_1, c', c'' depending on \mathbf{u} and α such that,*

$$\|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} \leq \frac{1}{\nu_1} (ch + (c_{\mathbf{f}_1}^* + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)}) \|C^n - C_h^{n-1}\|_{L^2(\Omega)}), \quad (5.1)$$

$$\|p^n - p_h^n\|_{L^2(\Omega)} \leq c'h + \frac{\nu_2}{\beta_1} \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} + \frac{1}{\beta_1} (c_{\mathbf{f}_1}^* + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)}) \|C^n - C_h^{n-1}\|_{L^2(\Omega)}, \quad (5.2)$$

and

$$\begin{aligned} \sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 + \sum_{n=1}^N \|(C_h^n - C^n) - (C_h^{n-1} - C^{n-1})\|_{L^2(\Omega)}^2 \\ + r_0 \sum_{n=1}^N \tau \|C_h^n - C^n\|_{L^2(\Omega)}^2 \leq c''(h^2 + \tau^2) + c_1 \sum_{n=1}^N \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2. \end{aligned} \quad (5.3)$$

Proof. Let (\mathbf{u}, p, C) and $(\mathbf{u}_h^n, p_h^n, C_h^n)$ solve respectively (V) and $(V_{h,1})$. We shall prove first (5.1), next (5.2), and finally (5.3).

We start by estimating the error on the velocity approximation. By taking the difference between the first equations of (V) for $t = t_n$ and $(V_{h,1})$ and testing with $\mathbf{v} = \mathbf{v}_h \in \mathcal{V}_{h,1}$, we obtain

$$\begin{aligned} \int_{\Omega} \nu(C_h^{n-1}(\mathbf{x})) (\mathbf{u}^n - \mathbf{u}_h^n)(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) \, d\mathbf{x} &= \int_{\Omega} (\mathbf{f}_1(C^n(\mathbf{x})) - \mathbf{f}_1(C_h^{n-1}(\mathbf{x}))) \cdot \mathbf{v}_h(\mathbf{x}) \, d\mathbf{x} \\ &+ \int_{\Omega} (\nu(C_h^{n-1}(\mathbf{x})) - \nu(C^n(\mathbf{x}))) \mathbf{u}^n(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) \, d\mathbf{x}. \end{aligned} \quad (5.4)$$

By inserting $\xi_h^1 \mathbf{u}^n$, taking $\mathbf{v}_h = \xi_h^1 \mathbf{u}^n - \mathbf{u}_h^n$, using the triangle inequality and the properties of ξ_h^1 , and using the properties of \mathbf{f}_1 and ν , we obtain

$$\nu_1 \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} \leq ch + c_{\mathbf{f}_1}^* \|C^n - C_h^{n-1}\|_{L^2(\Omega)} + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)} \|C^n - C_h^{n-1}\|_{L^2(\Omega)}.$$

Hence, we deduce (5.1).

To prove the error estimate on the pressure, we take the difference between the first equations of (V) (for $t = t_n$) and $(V_{h,1})$, insert $\rho_h(p^n)$, test with \mathbf{v}_h in $\mathcal{W}_{h,1}$, and obtain

$$\begin{aligned} \int_{\Omega} (\rho_h(p^n) - p_h^n)(\mathbf{x}) \operatorname{div} \mathbf{v}_h(\mathbf{x}) d\mathbf{x} &= \int_{\Omega} (\rho_h(p^n) - p^n)(\mathbf{x}) \operatorname{div} \mathbf{v}_h(\mathbf{x}) d\mathbf{x} + \int_{\Omega} \nu(C_h^{n-1})(\mathbf{u}^n - \mathbf{u}_h^n)(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} \\ &+ \int_{\Omega} (\nu(C^n(\mathbf{x})) - \nu(C_h^{n-1}(\mathbf{x}))) \mathbf{u}^n(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} \\ &- \int_{\Omega} (\mathbf{f}_1(C^n(\mathbf{x})) - \mathbf{f}_1(C_h^{n-1}(\mathbf{x}))) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x}. \end{aligned} \quad (5.5)$$

It follows from the inf-sup condition (4.11) that there exists \mathbf{v}_h in $\mathcal{W}_{h,1}$ such that

$$\operatorname{div} \mathbf{v}_h = \rho_h(p^n) - p_h^n \quad \text{and} \quad \|\mathbf{v}_h\|_{H(\operatorname{div},\Omega)} \leq \frac{1}{\beta_1} \|\rho_h(p^n) - p_h^n\|_{L^2(\Omega)}.$$

With this \mathbf{v}_h , (5.5) implies (5.2) by using the properties of ρ_h .

Let us now focus on (5.3). We choose the test function $S_h = r_h^n = C_h^n - R_h C^n$ in the third equation of $(V_{h,1})$ and multiply it by the time step τ . Then, we subtract the third equation of (V) integrated over $[t_{n-1}; t_n]$. We obtain, using the definition of g^n by (4.12):

$$\begin{aligned} &\left((C_h^n - C_h^{n-1}) - (C^n - C^{n-1}), r_h^n \right) + \alpha \left(\tau \nabla C_h^n - \int_{t_{n-1}}^{t_n} \nabla C(t) dt, \nabla r_h^n \right) \\ &+ \left(\tau \mathbf{u}_h^n \cdot \nabla C_h^n - \int_{t_{n-1}}^{t_n} \mathbf{u}(t) \cdot \nabla C(t) dt, r_h^n \right) + r_0 \left(\tau C_h^n - \int_{t_{n-1}}^{t_n} C(t) dt, r_h^n \right) = 0. \end{aligned} \quad (5.6)$$

The first term in the left-hand side of (5.6) can be bounded, by inserting $R_h C^n$ and $R_h C^{n-1}$ and using (2.9). We obtain:

$$\begin{aligned} \left((C_h^n - C_h^{n-1}) - (C^n - C^{n-1}), r_h^n \right) &= \frac{1}{2} \|r_h^n\|_{L^2(\Omega)}^2 - \frac{1}{2} \|r_h^{n-1}\|_{L^2(\Omega)}^2 + \frac{1}{2} \|r_h^n - r_h^{n-1}\|_{L^2(\Omega)}^2 \\ &+ \left((C^{n-1} - R_h C^{n-1}) - (C^n - R_h C^n), r_h^n \right). \end{aligned} \quad (5.7)$$

The last term of the previous equality can be bounded as follows, for any $\xi_1 > 0$, thanks to (2.8)

$$\begin{aligned} \left| \left((C^{n-1} - R_h C^{n-1}) - (C^n - R_h C^n), r_h^n \right) \right| &= \left| - \left(\int_{t_{n-1}}^{t_n} (R_h C' - C')(t) dt, r_h^n \right) \right| \\ &\leq \frac{c^2 h^2}{2\xi_1} \int_{t_{n-1}}^{t_n} \|C'(t)\|_{H^1(\Omega)}^2 dt + \frac{\tau \xi_1}{2} \|r_h^n\|_{L^2(\Omega)}^2 \\ &\leq \frac{c^2 h^2}{2\xi_1} \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; H^1(\Omega))}^2 + (S_2^0)^2 \frac{\tau \xi_1}{2} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.8)$$

By choosing $\xi_1 = \frac{\alpha}{25(S_2^0)^2}$, we obtain

$$\left| \left((C^{n-1} - R_h C^{n-1}) - (C^n - R_h C^n), r_h^n \right) \right| \leq c_1 h^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; H^1(\Omega))}^2 + \frac{\alpha}{50} \tau |r_h^n|_{1,\Omega}^2, \quad (5.9)$$

where c_1 is positive constant independent of h and τ .

The second term of the left-hand side of (5.6) can be bounded, by inserting $\tau(\nabla R_h C^n, \nabla r_h^n)$ and

$\int_{t_{n-1}}^{t_n} (\nabla R_h C(t), \nabla r_h^n) dt$. We have:

$$\begin{aligned} \alpha \left(\tau \nabla C_h^n - \int_{t_{n-1}}^{t_n} \nabla C(t) dt, \nabla r_h^n \right) &= \alpha \tau |r_h^n|_{1,\Omega}^2 + \alpha \int_{t_{n-1}}^{t_n} (\nabla (R_h C^n - R_h C(t)), \nabla r_h^n) dt \\ &\quad + \alpha \int_{t_{n-1}}^{t_n} (\nabla (R_h C(t) - C(t)), \nabla r_h^n) dt. \end{aligned} \quad (5.10)$$

Since $C^n - C(t) = \int_t^{t_n} C'(s) ds$, then we have by using the stability of the operator R_h in $H_0^1(\Omega)$, the Fubini theorem, the Cauchy-Schwarz inequality and (2.8) for any $\xi_2 > 0$:

$$\begin{aligned} \left| \alpha \int_{t_{n-1}}^{t_n} (\nabla (R_h C^n - R_h C(t)), \nabla r_h^n) dt \right| &\leq \alpha \int_{t_{n-1}}^{t_n} \int_t^{t_n} |R_h C'(s)|_{1,\Omega} |r_h^n|_{1,\Omega} ds dt \\ &\leq \alpha c |r_h^n|_{1,\Omega} \int_{t_{n-1}}^{t_n} |C'(s)|_{1,\Omega} (s - t_{n-1}) ds \\ &\leq \frac{\alpha c^2 \tau^2 \xi_2}{2\sqrt{3}} \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; H^1(\Omega))}^2 + \frac{\tau \alpha}{2\sqrt{3} \xi_2} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.11)$$

In addition, we apply the Cauchy-Schwarz inequality, (2.8) and (4.2) to obtain for any $\xi_3 > 0$:

$$\begin{aligned} \left| \alpha \int_{t_{n-1}}^{t_n} (\nabla (R_h C(t) - C(t)), \nabla r_h^n) dt \right| &\leq \frac{\alpha \xi_3}{2} \int_{t_{n-1}}^{t_n} |(R_h C - C)(t)|_{1,\Omega}^2 dt + \frac{\tau \alpha}{2\xi_3} |r_h^n|_{1,\Omega}^2 \\ &\leq \frac{\alpha c^2 h^2 \xi_3}{2} \|C\|_{L^2(t_{n-1}, t_n; H^2(\Omega))}^2 + \frac{\tau \alpha}{2\xi_3} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.12)$$

By choosing $\xi_2 = 25/\sqrt{3}$ and $\xi_3 = 25$, we obtain, using the properties of the operator R_h and (5.10), (5.11) and (5.12)

$$\begin{aligned} \left| \alpha \left(\tau \nabla C_h^n - \int_{t_{n-1}}^{t_n} \nabla C(t) dt, \nabla r_h^n \right) - \alpha \tau |r_h^n|_{1,\Omega}^2 \right| &\leq c_1 h^2 \|C\|_{L^2(t_{n-1}, t_n; H^2(\Omega))}^2 \\ &\quad + c_2 \tau^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; H^1(\Omega))}^2 + \frac{\alpha}{25} \tau |r_h^n|_{1,\Omega}^2, \end{aligned} \quad (5.13)$$

where c_1 and c_2 are positive constants independent of h and τ .

Let the third term in the left-hand side of (5.6) be denoted by

$$b = \left(\tau \mathbf{u}_h^n \cdot \nabla C_h^n - \int_{t_{n-1}}^{t_n} \mathbf{u}(t) \cdot \nabla C(t) dt, r_h^n \right). \quad (5.14)$$

We insert $\left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla C(t) dt, r_h^n \right)$ and $\tau(\mathbf{u}_h^n \cdot \nabla R_h C^n, r_h^n)$ to get by noticing that $(\mathbf{u}_h^n \cdot \nabla r_h^n, r_h^n) = 0$:

$$b = b_1 + b_2 = \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (R_h C^n - C(t)) dt, r_h^n \right) + \left(\int_{t_{n-1}}^{t_n} (\mathbf{u}_h^n - \mathbf{u}(t)) \cdot \nabla C(t) dt, r_h^n \right). \quad (5.15)$$

We insert $\pm \tau(\mathbf{u}_h^n \cdot \nabla C^n, r_h^n)$ in b_1 and we get:

$$b_1 = \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (R_h C^n - C^n) dt, r_h^n \right) + \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (C^n - C(t)) dt, r_h^n \right). \quad (5.16)$$

Using the L^2 - L^4 - L^4 generalized Cauchy-Schwarz inequality, (4.2) and Lemma 2.1, we obtain, for any $\xi_4 > 0$:

$$\begin{aligned} \left| \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (R_h C^n - C^n) dt, r_h^n \right) \right| &\leq \tau \|\mathbf{u}_h^n\|_{L^2(\Omega)^d} \|R_h C^n - C^n\|_{W^{1,4}(\Omega)} \|r_h^n\|_{L^4(\Omega)} \\ &\leq c \tau h \|\mathbf{u}_h^n\|_{L^2(\Omega)^d} \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))} \|r_h^n\|_{L^4(\Omega)} \\ &\leq \frac{c^2 \xi_4 h^2 \tau}{2} \|\mathbf{u}_h^n\|_{L^2(\Omega)^d}^2 \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))}^2 + \frac{\tau (S_4^0)^2}{2\xi_4} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.17)$$

By choosing $\xi_4 = 25(S_4^0)^2/\alpha$, we obtain by using (4.13):

$$\left| \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (R_h C^n - C^n) dt, r_h^n \right) \right| \leq c_3 h^2 \tau \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))}^2 + \frac{\alpha}{50} \tau |r_h^n|_{1,\Omega}^2. \quad (5.18)$$

By treating as above, we have for any $\xi_5 > 0$:

$$\begin{aligned} \left| \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla (C^n - C(t)) dt, r_h^n \right) \right| &= \left| \left(\int_{t_{n-1}}^{t_n} \int_t^{t_n} \mathbf{u}_h^n \cdot \nabla C'(s) ds dt, r_h^n \right) \right| \\ &= \left| \left(\int_{t_{n-1}}^{t_n} \int_s^{t_{n-1}} \mathbf{u}_h^n \cdot \nabla C'(s) dt ds, r_h^n \right) \right| \\ &\leq \|\mathbf{u}_h^n\|_{L^2(\Omega)^d} \|r_h^n\|_{L^4(\Omega)} \int_{t_{n-1}}^{t_n} \|\nabla C'(s)\|_{L^4(\Omega)} (s - t_{n-1}) ds \\ &\leq \frac{\tau^2 \xi_5}{6} \|\mathbf{u}_h^n\|_{L^2(\Omega)^d}^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; W^{1,4}(\Omega))}^2 + \frac{\tau (S_4^0)^2}{2\xi_5} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.19)$$

We deduce by regrouping (5.16), (5.18) and (5.19) (for $\xi_5 = 25(S_4^0)^2/\alpha$) that

$$|b_1| \leq c_3 \tau h^2 \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))}^2 + c_4 \tau^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; W^{1,4}(\Omega))}^2 + \frac{\alpha}{25} \tau |r_h^n|_{1,\Omega}^2. \quad (5.20)$$

We insert \mathbf{u}^n in b_2 and we get, using the L^2 - L^4 - L^4 inequality

$$\begin{aligned} |b_2| &= \left| \left(\int_{t_{n-1}}^{t_n} (\mathbf{u}_h^n - \mathbf{u}^n) \cdot \nabla C(t) dt, r_h^n \right) + \left(\int_{t_{n-1}}^{t_n} \int_t^{t_n} \mathbf{u}'(s) \cdot \nabla C(t) ds dt, r_h^n \right) \right| \\ &\leq \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d} \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))} \|r_h^n\|_{L^4(\Omega)} \\ &\quad + \tau^{3/2} \|C\|_{L^\infty(0,T;H^1(\Omega))} \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^4(\Omega)^d)} \|r_h^n\|_{L^4(\Omega)}. \end{aligned} \quad (5.21)$$

By using (2.8) and Lemma 2.1 and taking $\xi = 25(S_4^0)^2/\alpha$, we obtain:

$$\begin{aligned} |b_2| &\leq c_{51} \tau \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))}^2 \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 \\ &\quad + c_{52} \tau^2 \|C\|_{L^\infty(0,T;H^1(\Omega))}^2 \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^4(\Omega)^d)}^2 + \frac{\alpha}{25} \tau |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.22)$$

Finally, we combine (5.14), (5.20) and (5.22), and we deduce that

$$\begin{aligned} \left| \left(\tau \mathbf{u}_h \cdot \nabla C_h^n - \int_{t_{n-1}}^{t_n} \mathbf{u} \cdot \nabla C(t) dt, r_h^n \right) \right| &\leq c_3 \tau h^2 \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))}^2 + c_4 \tau^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; W^{1,4}(\Omega))}^2 \\ &\quad + c_{51} \tau \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))}^2 \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 \\ &\quad + c_{52} \tau^2 \|C\|_{L^\infty(0,T;H^1(\Omega))}^2 \left\| \frac{\partial \mathbf{u}}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^4(\Omega)^d)}^2 \\ &\quad + \frac{2\alpha}{25} \tau |r_h^n|_{1,\Omega}^2, \end{aligned} \quad (5.23)$$

where c_3 , c_4 , c_{51} and c_{52} are positive constants independent of h and τ .

The last term in the left-hand side of (5.6) can be bounded, by inserting $\pm(\tau R_h C^n, r_h^n)$ and $\pm\tau(C^n, r_h^n)$

$$\left(\tau C_h^n - \int_{t_{n-1}}^{t_n} C(t) dt, r_h^n \right) = \tau(C_h^n - R_h C^n, r_h^n) + \tau(R_h C^n - C^n, r_h^n) + \left(\int_{t_{n-1}}^{t_n} (C^n - C(t)) dt, r_h^n \right)$$

so that

$$\left| \left(\tau C_h^n - \int_{t_{n-1}}^{t_n} C(t) dt, r_h^n \right) - \tau \|r_h^n\|_{L^2(\Omega)}^2 \right| \leq \tau \left| (R_h C^n - C^n, r_h^n) \right| + \left| \left(\int_{t_{n-1}}^{t_n} (C^n - C(t)) dt, r_h^n \right) \right|. \quad (5.24)$$

Moreover, using the approximation properties of R_h , we have:

$$\begin{aligned} \tau \left| (R_h C^n - C^n, r_h^n) \right| &\leq \tau \|R_h C^n - C^n\|_{L^2(\Omega)} \|r_h^n\|_{L^2(\Omega)} \\ &\leq c\tau h \|r_h^n\|_{L^2(\Omega)} \|C\|_{L^\infty(0,T;H^1(\Omega))} \\ &\leq \frac{c^2 h^2 \tau \xi_6}{2} \|C\|_{L^\infty(0,T;H^1(\Omega))}^2 + \frac{\tau (S_2^0)^2}{2\xi_6} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.25)$$

Furthermore,

$$\begin{aligned} \left| \left(\int_{t_{n-1}}^{t_n} (C^n - C(t)) dt, r_h^n \right) \right| &= \left| \left(\int_{t_{n-1}}^{t_n} \int_t^{t_n} C'(s) ds dt, r_h^n \right) \right| \\ &\leq \frac{\tau^{3/2}}{\sqrt{3}} \|r_h^n\|_{L^2(\Omega)} \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^2(\Omega))} \\ &\leq \frac{\xi_7 \tau^2}{6} \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^2(\Omega))}^2 + \frac{\tau (S_2^0)^2}{2\xi_7} |r_h^n|_{1,\Omega}^2. \end{aligned} \quad (5.26)$$

Next, we combine (5.24), (5.25) and (5.26), and we choose $\xi_6 = \xi_7 = \frac{25(S_2^0)^2}{\alpha}$ to obtain:

$$\left| \left(\tau C_h^n - \int_{t_{n-1}}^{t_n} C(t) dt, r_h^n \right) - \tau \|r_h^n\|_{L^2(\Omega)}^2 \right| \leq c_6 \tau h^2 \|C\|_{L^\infty(0,T;H^1(\Omega))}^2 + c_7 \tau^2 \left\| \frac{\partial C}{\partial t} \right\|_{L^2(t_{n-1}, t_n; L^2(\Omega))}^2 + \frac{\alpha}{25} \tau |r_h^n|_{1,\Omega}^2, \quad (5.27)$$

where c_6 and c_7 are positive constants independent of τ and h .

Now we use (5.6), (5.7), (5.9), (5.13), (5.23) and (5.27) and we sum over n from 1 to $m \leq N$. This leads to

$$\begin{aligned} \frac{1}{2} \|r_h^m\|_{L^2(\Omega)}^2 + \frac{1}{2} \sum_{n=1}^m \|r_h^n - r_h^{n-1}\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^m \tau |r_h^n|_{1,\Omega}^2 + r_0 \sum_{n=1}^m \tau \|r_h^n\|_{L^2(\Omega)}^2 &\leq \\ c(h^2 + \tau^2) + c' \sum_{n=1}^m \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 + \frac{9\alpha}{50} \sum_{n=1}^m \tau |r_h^n|_{1,\Omega}^2. \end{aligned}$$

Finally, we obtain

$$\begin{aligned} \sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 + \sum_{n=1}^N \|(C_h^n - C^n) - (C_h^{n-1} - C^{n-1})\|_{L^2(\Omega)}^2 \\ + r_0 \sum_{n=1}^N \tau \|C_h^n - C^n\|_{L^2(\Omega)}^2 \leq c(h^2 + \tau^2) + c' \sum_{n=1}^N \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2, \end{aligned}$$

where c and c' are positive constants independent of h and τ .

Remark 5.2. If the viscosity ν is constant independent of C , it suffices to take $\mathbf{u} \in L^\infty(0, T; H^1(\Omega)^d)$ in Theorem 5.1, as in this case, Equation (5.4) becomes

$$\int_{\Omega} \nu (\mathbf{u}^n - \mathbf{u}_h^n)(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} = \int_{\Omega} (\mathbf{f}_1(C^n(\mathbf{x})) - \mathbf{f}_1(C_h^{n-1}(\mathbf{x}))) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x}. \quad (5.28)$$

Corollary 5.3. Under the assumptions of Theorem 5.1, we have the a priori error estimates corresponding to $(V_{h,1})$:

$$\begin{aligned} \sup_{0 \leq n \leq N} \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} &\leq c(h + \tau), \\ \sup_{0 \leq n \leq N} \|p^n - p_h^n\|_{L^2(\Omega)} &\leq c'(h + \tau), \\ \sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 &\leq c''(h^2 + \tau^2), \end{aligned} \quad (5.29)$$

where c, c' and c'' are independent of h and τ .

Proof. We first consider relation (5.1). By inserting C^{n-1} and using the triangle inequality, we have

$$\sum_{n=1}^N \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 \leq c \left(\sum_{n=1}^N \tau \|C_h^{n-1} - C^{n-1}\|_{L^2(\Omega)}^2 + \sum_{n=1}^N \tau \int_{\Omega} \left| \int_{t_{n-1}}^{t_n} C'(s) ds \right|^2 dx \right).$$

We then deduce from (5.3) the following relation

$$\sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 \leq c(h^2 + \tau^2) + c' \sum_{n=1}^N \tau \|C_h^{n-1} - C^{n-1}\|_{L^2(\Omega)}^2. \quad (5.30)$$

Finally, the following inequality follows from the discrete Grönwall Lemma

$$\sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 \leq c''(h^2 + \tau^2). \quad (5.31)$$

It is crucial to note that in the right-hand side of (5.30), $\sum_{n=1}^N \tau$ is bounded by the final time T which is a quantity that does not depend on N ; thus the constant c'' that appears in (5.31) does not depend on τ .

Inequality (5.31) and Inequalities (5.1) and (5.2) yield the desired bounds in (5.29).

Theorem 5.4. *Let (\mathbf{u}, p, C) be the solution of Problem (V_a) and $(\mathbf{u}_h^n, p_h^n, C_h^n)$ be the solution of Problem $(V_{h,2})$. If $\mathbf{u} \in L^\infty(0, T; H^1(\Omega)^d) \cap L^\infty(0, T; L^\infty(\Omega)^d)$, $p \in L^\infty(0, T; H^2(\Omega))$, $C \in L^\infty(0, T; W^{2,4}(\Omega))$ and $\frac{\partial C}{\partial t} \in L^2(0, T; W^{1,4}(\Omega))$ and under Assumption 3.1, there exist positive constants c, c_1, c', c'' depending on \mathbf{u} and α such that,*

$$\|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} \leq \frac{1}{\nu_1} (ch + (c_{\mathbf{f}_1}^* + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)}) \|C^n - C_h^{n-1}\|_{L^2(\Omega)}), \quad (5.32)$$

$$|p^n - p_h^n|_{1,\Omega} \leq c'h + \frac{\nu_2}{\beta_2} \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} + \frac{1}{\beta_2} (c_{\mathbf{f}_1}^* + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)}) \|C^n - C_h^{n-1}\|_{L^2(\Omega)}, \quad (5.33)$$

and

$$\begin{aligned} \sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 + \sum_{n=1}^N \|(C_h^n - C^n) - (C_h^{n-1} - C^{n-1})\|_{L^2(\Omega)}^2 \\ + r_0 \sum_{n=1}^N \tau \|C_h^n - C^n\|_{L^2(\Omega)}^2 \leq c''(h^2 + \tau^2) + c_1 \sum_{n=1}^m \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2. \end{aligned} \quad (5.34)$$

Proof. Let (\mathbf{u}, p, C) and $(\mathbf{u}_h^n, p_h^n, C_h^n)$ solve respectively (V_a) and $(V_{h,2})$. We shall prove first (5.32), next (5.33), and finally (5.34).

Let us estimate the velocity error. By taking the difference between the first equations of (V_a) and $(V_{h,2})$ and testing with $\mathbf{v} = \mathbf{v}_h \in \mathcal{V}_{h,2}$, we obtain

$$\begin{aligned} \int_{\Omega} \nu(C_h^{n-1}(\mathbf{x})) (\mathbf{u}^n - \mathbf{u}_h^n)(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) dx &= \int_{\Omega} (\mathbf{f}_1(C^n(\mathbf{x})) - \mathbf{f}_1(C_h^{n-1}(\mathbf{x}))) \cdot \mathbf{v}_h(\mathbf{x}) dx \\ &+ \int_{\Omega} (\nu(C_h^{n-1}(\mathbf{x})) - \nu(C^n(\mathbf{x}))) \mathbf{u}^n(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) dx \\ &- \int_{\Omega} \nabla(p^n - r_h(p^n)) \cdot \mathbf{v}_h dx. \end{aligned} \quad (5.35)$$

By inserting $\mathcal{F}_h \mathbf{u}^n$, choosing $\mathbf{v}_h = (\mathcal{F}_h \mathbf{u}^n - \mathbf{u}_h^n)$ which belongs to $\mathcal{V}_{h,2}$ using the triangle inequality and the approximation properties of \mathcal{F}_h and r_h and the bounds on ν , we obtain (5.32).

To prove the error estimate on the pressure, we take the difference between the first equations of (V_a) and $(V_{h,2})$, insert $\nabla r_h(p^n)$, test with \mathbf{v}_h in $\mathcal{W}_{h,2}$, and we obtain

$$\begin{aligned}
\int_{\Omega} \nabla(p_h^n - r_h(p^n))(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} &= \int_{\Omega} \nabla(p^n - r_h(p^n))(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} \\
&+ \int_{\Omega} \nu(C_h^{n-1})(\mathbf{u}^n - \mathbf{u}_h^n)(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} \\
&+ \int_{\Omega} (\nu(C^n(\mathbf{x})) - \nu(C_h^{n-1})(\mathbf{x})) \mathbf{u}^n(\mathbf{x}) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x} \\
&- \int_{\Omega} (\mathbf{f}_1(C^n(\mathbf{x})) - \mathbf{f}_1(C_h^{n-1}(\mathbf{x}))) \cdot \mathbf{v}_h(\mathbf{x}) d\mathbf{x}.
\end{aligned} \tag{5.36}$$

It follows from the inf-sup condition (4.22) and by applying the Cauchy-Schwarz inequality to the right-hand side of (5.36) and Assumption 3.1 that

$$|p_h^n - r_h(p^n)|_{1,\Omega} \leq \frac{1}{\beta_2} \left[|p^n - r_h(p^n)|_{1,\Omega} + \nu_2 \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} + (c_{\mathbf{f}_1}^* + \lambda \|\mathbf{u}\|_{L^\infty(0,T;L^\infty(\Omega)^d)}) \|C^n - C_h^{n-1}\|_{L^2(\Omega)} \right]. \tag{5.37}$$

Inserting $r_h(p^n)$ in $|p^n - r_h(p^n)|_{1,\Omega}$ and using a triangular inequality and (5.37) implies (5.33) by using the properties of r_h .

Now we prove (5.34). We choose the test function $S_h = r_h^n = C_h^n - R_h C^n$ in the third equation of $(V_{h,2})$ and multiply it by the time step τ . Then, we subtract the third equation of (V_a) integrated over $[t_{n-1}, t_n]$. We obtain:

$$\begin{aligned}
&\left((C_h^n - C_h^{n-1}) - (C^n - C^{n-1}), r_h^n \right) + \alpha \left(\tau \nabla C_h^n - \int_{t_{n-1}}^{t_n} \nabla C(t) dt, \nabla r_h^n \right) \\
&+ \left(\tau \mathbf{u}_h^n \cdot \nabla C_h^n - \int_{t_{n-1}}^{t_n} \mathbf{u}(t) \cdot \nabla C(t) dt, r_h^n \right) + \frac{\tau}{2} (\operatorname{div} \mathbf{u}_h^n C_h^n, r_h^n) + r_0 \left(\tau C_h^n - \int_{t_{n-1}}^{t_n} C(t) dt, r_h^n \right) = 0.
\end{aligned} \tag{5.38}$$

All the terms of (5.38) can be treated as the proof of Theorem 5.1 (see (5.9), (5.13), (5.27)) except the non-linear one denoted by b_n .

By inserting $\pm \tau(\mathbf{u}_h^n \cdot \nabla R_h C^n, r_h^n)$, $\pm \frac{\tau}{2}(\operatorname{div} \mathbf{u}_h^n R_h C^n, r_h^n)$, b_n becomes:

$$\begin{aligned}
b_n &= \tau(\mathbf{u}_h^n \cdot \nabla(C_h^n - R_h C^n), r_h^n) + \tau(\mathbf{u}_h^n \cdot \nabla R_h C^n, r_h^n) - \left(\int_{t_{n-1}}^{t_n} \mathbf{u}(t) \cdot \nabla C(t) dt, r_h^n \right) \\
&+ \frac{\tau}{2} (\operatorname{div} \mathbf{u}_h^n (C_h^n - R_h C^n), r_h^n) + \frac{\tau}{2} (\operatorname{div} \mathbf{u}_h^n R_h C^n, r_h^n).
\end{aligned}$$

Then, we insert $\pm \tau(\mathbf{u}_h^n \cdot \nabla C^n, r_h^n)$, $\pm \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla C(t) dt, r_h^n \right)$ and $\pm \frac{\tau}{2}(\operatorname{div} \mathbf{u}_h^n C^n, r_h^n)$ to get by noticing that $\tau(\mathbf{u}_h^n \cdot \nabla(C_h^n - R_h C^n), r_h^n) + \frac{\tau}{2}(\operatorname{div} \mathbf{u}_h^n (C_h^n - R_h C^n), r_h^n) = 0$:

$$\begin{aligned}
b_n &= \tau(\mathbf{u}_h^n \cdot \nabla(R_h C^n - C^n), r_h^n) + \left(\int_{t_{n-1}}^{t_n} \mathbf{u}_h^n \cdot \nabla(C^n - C(t)) dt, r_h^n \right) \\
&+ \int_{t_{n-1}}^{t_n} ((\mathbf{u}_h^n - \mathbf{u}(t)) \cdot \nabla C(t) dt, r_h^n) \\
&+ \frac{\tau}{2} (\operatorname{div} \mathbf{u}_h^n (R_h C^n - C^n), r_h^n) + \frac{\tau}{2} (\operatorname{div} \mathbf{u}_h^n C^n, r_h^n).
\end{aligned} \tag{5.39}$$

The sum of the first two terms in the right-hand side of (5.39) is exactly the expression of b_1 in (5.16). The third term in the right-hand side of (5.39) is exactly b_2 as defined in (5.15). Therefore, these terms can be treated exactly as in the proof of theorem 5.1. The last two terms, denoted below as $b_{n,4}$ and $b_{n,5}$

can be treated as follows:

$$\begin{aligned}
b_{n,4} &= \frac{\tau}{2}(\operatorname{div} \mathbf{u}_h^n (R_h C^n - C^n), r_h^n) \\
&= -\frac{\tau}{2}(\mathbf{u}_h^n \cdot \nabla (R_h C^n - C^n), r_h^n) - \frac{\tau}{2}(\mathbf{u}_h^n \cdot \nabla r_h^n, (R_h C^n - C^n)) \\
&= b_{n,4}^1 + b_{n,4}^2,
\end{aligned}$$

where, using Lemma 2.1, (2.8) and (4.2), we have, for any $\bar{\xi}_1 > 0$

$$\begin{aligned}
|b_{n,4}^1| &\leq \frac{\tau}{2} \|\mathbf{u}_h^n\|_{L^2(\Omega)^d} \|\nabla (R_h C^n - C^n)\|_{L^4(\Omega)} \|r_h^n\|_{L^4(\Omega)} \\
&\leq \frac{ch^2 \tau \bar{\xi}_1}{8} \|\mathbf{u}_h^n\|_{L^2(\Omega)^d}^2 \|C\|_{L^\infty(0,T;W^{2,4}(\Omega))}^2 + \frac{\tau(S_4^0)^2}{2\bar{\xi}_1} |r_h^n|_{1,\Omega}^2
\end{aligned}$$

and where, using (4.2) and (4.3) with $d \leq 3$ we have, for any $\bar{\xi}_2 > 0$

$$\begin{aligned}
|b_{n,4}^2| &\leq \frac{\tau}{2} \|\mathbf{u}_h^n\|_{L^4(\Omega)^d} \|\nabla r_h^n\|_{L^2(\Omega)} \|R_h C^n - C^n\|_{L^4(\Omega)} \\
&\leq \frac{ch^2}{2} \tau \|\mathbf{u}_h^n\|_{L^2(\Omega)^d} \times h^{-\frac{d}{4}} |r_h^n|_{1,\Omega} \|C^n\|_{W^{2,4}(\Omega)} \\
&\leq \frac{c\bar{\xi}_2}{8} h^{\frac{8-d}{2}} \tau \|\mathbf{u}_h^n\|_{L^2(\Omega)^d}^2 \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))}^2 + \frac{\tau}{2\bar{\xi}_2} |r_h^n|_{1,\Omega}^2.
\end{aligned}$$

Note that in the bounds of $|b_{n,4}^1|$ and $|b_{n,4}^2|$, the $L^2(\Omega)$ norm of \mathbf{u}_h^n can be bounded by a constant depending only on the data of the problem through Theorem 4.2 using both (4.23) and (4.24). Note also that $\frac{8-d}{2} \geq 2$ for $d \leq 3$, which is sufficient for our proof.

Moreover, $b_{n,5}$ can be decomposed in the same way and treated as follows (where we use that $\operatorname{div} \mathbf{u}^n = 0$):

$$\begin{aligned}
b_{n,5} &= \frac{\tau}{2}(\operatorname{div} (\mathbf{u}_h^n - \mathbf{u}^n) C^n, r_h^n) \\
&= -\frac{\tau}{2}((\mathbf{u}_h^n - \mathbf{u}^n) \cdot \nabla C^n, r_h^n) - \frac{\tau}{2}((\mathbf{u}_h^n - \mathbf{u}^n) \cdot \nabla r_h^n, C^n) \\
&= b_{n,5}^1 + b_{n,5}^2.
\end{aligned}$$

Using Lemma 2.1 and (2.8) we have, for any $\bar{\xi}_3 > 0$

$$\begin{aligned}
|b_{n,5}^1| &= \left| \frac{\tau}{2}((\mathbf{u}_h^n - \mathbf{u}^n) \cdot \nabla C^n, r_h^n) \right| \\
&\leq \frac{\tau}{2} \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d} \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))} \|r_h^n\|_{L^4(\Omega)} \\
&\leq \frac{\tau \bar{\xi}_3}{8} \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 \|C\|_{L^\infty(0,T;W^{1,4}(\Omega))}^2 + \frac{\tau(S_4^0)^2}{2\bar{\xi}_3} |r_h^n|_{1,\Omega}^2
\end{aligned}$$

and for any $\bar{\xi}_4 > 0$

$$\begin{aligned}
|b_{n,5}^2| &= \left| \frac{\tau}{2}((\mathbf{u}_h^n - \mathbf{u}^n) \cdot \nabla r_h^n, C^n) \right| \\
&\leq \frac{\tau}{2} \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d} \|C^n\|_{L^\infty(\Omega)} |r_h^n|_{1,\Omega} \\
&\leq \frac{\tau \bar{\xi}_4}{8} \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2 \|C\|_{L^\infty([0,T] \times \Omega)}^2 + \frac{\tau}{2\bar{\xi}_4} |r_h^n|_{1,\Omega}^2.
\end{aligned}$$

By using the above bounds with a suitable choice of $\bar{\xi}_i, i = 1, \dots, 4$, and summing over n from 1 to $m \leq N$, we get

$$\frac{1}{2} \|r_h^m\|_{L^2(\Omega)}^2 + \frac{1}{2} \sum_{n=1}^m \|r_h^n - r_h^{n-1}\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^m \tau |r_h^n|_{1,\Omega}^2 + r_0 \sum_{n=1}^m \tau \|r_h^n\|_{L^2(\Omega)}^2 \leq c(h^2 + \tau^2) + c \sum_{n=1}^m \tau \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^d}^2,$$

and we deduce finally (5.34).

Corollary 5.5. *Under the assumption of Theorem 5.4, we have the a priori error estimates corresponding to $(V_{h,2})$:*

$$\begin{aligned} \sup_{0 \leq n \leq N} \|\mathbf{u}^n - \mathbf{u}_h^n\|_{L^2(\Omega)^d} &\leq c(h + \tau), \\ \sup_{0 \leq n \leq N} |p^n - p_h^n|_{H^1(\Omega)} &\leq c'(h + \tau), \\ \sup_{0 \leq n \leq N} \|C_h^n - C^n\|_{L^2(\Omega)}^2 + \alpha \sum_{n=1}^N \tau |C_h^n - C^n|_{1,\Omega}^2 &\leq c''(h^2 + \tau^2), \end{aligned} \quad (5.40)$$

where c, c' and c'' are independent of h and τ .

6. NUMERICAL RESULTS

To validate the theoretical results, we perform several numerical simulations using Freefem++ (see [18]). We consider a square domain $\Omega =]0, 1[^2$. Each edge is divided into N equal segments so that Ω is divided into $2N^2$ triangles. For the numerical tests, we consider $\alpha = 1$, $r_0 = 1$, $\mathbf{f}_1(C) = C + 1$ and $\nu(C) = \sin(C) + 2$. We choose the right-hand sides \mathbf{f}_0 and g so that the exact solution is given by $(\mathbf{u}, p, C) = (e^{-t/4} \text{curl} \psi, p, C)$ where ψ, p and C are defined by

$$\begin{aligned} \psi(x, y) &= e^{-100((x-\frac{1}{2})^2 + (y-\frac{1}{2})^2)}, \\ p(x, y, t) &= (t + 1) \cos(\pi x) \cos(\pi y), \\ C(x, y, t) &= \sin t x^2(x-1)^2 y^2(y-1)^2. \end{aligned} \quad (6.1)$$

We define the following total relative error between the exact and numerical solutions :

$$err = \left(\frac{\sup_{1 \leq n \leq N} \|\mathbf{u}_h^n - \mathbf{u}^n\|_{L^2(\Omega)^2}^2 + \sup_{1 \leq n \leq N} |p_h^n - p^n|_{L^2(\Omega)}^2 + \sum_{n=1}^N \tau |C_h^n - C^n|_{H^1(\Omega)}^2}{\sup_{1 \leq n \leq N} \|\mathbf{u}^n\|_{L^2(\Omega)^2}^2 + \sup_{1 \leq n \leq N} |p^n|_{L^2(\Omega)}^2 + \sum_{n=1}^N \tau |C^n|_{H^1(\Omega)}^2} \right)^{1/2}. \quad (6.2)$$

We test the algorithms for N ranging from 60 to 120, by a step of 10 with $T = 1$, $h = \frac{1}{N}$ and $\tau = h$. Moreover, in order to be able to compare the accuracy of the schemes on similar number of unknowns and non-zero entries in the matrices associated to the linear systems, we also use $N = 30$ for the second scheme and $N = 240$ for the first one.




FIGURE 1. Pressure error with respect to the mesh size.




FIGURE 2. Velocity error with respect to the mesh size.

Figures 1, 2, and 3 show, in logarithmic scale, the curves of the errors on the pressure, the velocity and the concentration, respectively, according to the mesh step h for the first and second schemes $(V_{h,1})$ and




FIGURE 3. Concentration error with respect to the mesh size.




FIGURE 4. Total relative error with respect to the mesh size.

($V_{h,2}$). Figure 4 represents, in logarithmic scale, the curve of the total relative error according to the mesh step h for the first and second discrete schemes ($V_{h,1}$) and ($V_{h,2}$). Their respective slopes are 1.0013 and 1.0142 which is in conformity with the theoretical order of convergence of both schemes. We also note that the second scheme, which uses higher order polynomials for the pressure than the first scheme, yields a second order accuracy on this variable.




FIGURE 5. Total relative error with respect to the number of unknowns.




FIGURE 6. Total relative error with respect to the number of non-zero coefficients in the matrices.

In order to include a more complete comparison of the accuracy of the two schemes with respect to their numerical complexity, we show on Figures 4, 5 and 6 their total relative errors, not only with respect to the mesh size, but also with respect to the number of unknowns and with respect to the number of non-zero coefficients in the associated matrices. In all three figures, the curve of the second scheme is below the one of the first scheme, which means that, for a given complexity, the second scheme is more accurate than the first one for this particular example. It is probable that this advantage is linked to the high stiffness of the exact solution, which is better captured by the second scheme which uses higher-order polynomials.

REFERENCES

- [1] H. ABBOUD, V. GIRAULT AND T. SAYAH, A SECOND ORDER ACCURACY IN TIME FOR A FULL DISCRETIZED TIME-DEPENDENT NAVIER-STOKES EQUATIONS BY A TWO-GRID SCHEME. *Numer. Math.*, **114**, (2009), pp. 189-231.
- [2] D. ARNOLD, F. BREZZI AND M. FORTIN, A STABLE FINITE ELEMENT FOR THE STOKES EQUATIONS, *Calcolo* **21**, (1984), pp. 337-344.
- [3] C. BERNARDI AND V. GIRAULT, A LOCAL REGULARISATION OPERATION FOR TRIANGULAR AND QUADRILATERAL FINITE ELEMENTS. *SIAM J. Numer. Anal.*, **35**, (1998), pp. 1893-1916.
- [4] C. BERNARDI, B. MÉTIVET AND B. PERNAUD-THOMAS, COUPLAGE DES ÉQUATIONS DE NAVIER-STOKES ET DE LA CHALEUR: LE MODÈLE ET SON APPROXIMATION PAR ÉLÉMENTS FINIS. *RAIRO Modél. Math. Anal. Numér.*, **29(7)**, (1995), pp. 871-921.
- [5] C. BERNARDI, S. DIB, V. GIRAULT, F. HECHT, F. MURAT, T. SAYAH, FINITE ELEMENT METHOD FOR DARCY'S PROBLEM COUPLED WITH THE HEAT EQUATION. *Numer. Math.*, **139 (2)**, (2018), pp. 315-348.
- [6] C. BERNARDI, D. YAKOUBI AND S. MAAROUF, SPECTRAL DISCRETIZATION OF DARCY'S EQUATIONS COUPLED WITH THE HEAT EQUATION. *IMA Journal of Numerical Analysis*, (2015), pp. 1-24.
- [7] J. BOUSSINESQ, THÉORIE ANALYTIQUE DE LA CHALEUR, *Volume 2 of Lecture Notes in Mathematics. Gauthier-Villars, Paris*, (1903).
- [8] Z. CHEN AND R. EWING, MATHEMATICAL ANALYSIS FOR RESERVOIR MODELS. *SIAM J. Math. Anal.*, **30**, (1999), pp. 431-453.
- [9] P. G. CIARLET, BASIC ERROR ESTIMATES FOR ELLIPTIC PROBLEMS. *Handbook of Numerical Analysis, Finite Element Methods (Part I)*, **II**, North-Holland, (1991), pp. 17-343.
- [10] P. CLÉMENT, APPROXIMATION BY FINITE ELEMENT FUNCTIONS USING LOCAL REGULARIZATION. *RAIRO Anal. Numér.*, (1975), pp. 77-84.
- [11] J. DETEIX, A. JENDOUBI AND D. YAKOUBI, A COUPLED PREDICTION SCHEME FOR SOLVING THE NAVIER-STOKES AND CONVECTION-DIFFUSION EQUATIONS. *SIAM J. Numer. Anal.*, **52(5)**, (2014), pp. 2415-2439.
- [12] X. FENG, ON EXISTENCE AND UNIQUENESS RESULTS FOR A COUPLED SYSTEM MODELING MISCIBLE DISPLACEMENT IN POROUS MEDIA. *J. Math. Anal. Appl.*, **194**, (1995), pp. 883-910.
- [13] M. GAULTIER AND M. LEZAUN, ÉQUATIONS DE NAVIER-STOKES COUPLÉES À DES ÉQUATIONS DE LA CHALEUR: RÉOLUTION PAR UNE MÉTHODE DE POINT FIXE EN DIMENSION INFINIE. *Ann. Sci. Math. Québec*, **13(1)**, (1989), pp. 1-17.
- [14] V. GIRAULT & P.-A. RAVIART, FINITE ELEMENT METHODS FOR THE NAVIER-STOKES EQUATIONS. THEORY AND ALGORITHMS, in *Springer Series in Computational Mathematics 5*, Springer-Verlag, Berlin, (1986).
- [15] V. GIRAULT, J. L. LIONS, TWO-GRID FINITE-ELEMENT SCHEMES FOR THE STEADY NAVIER-STOKES PROBLEM IN POLYHEDRA. *Portugaliae Mathematica, Nova Série* **58(1)**, (2001), pp. 25-57.
- [16] J. DOUGLAS, R. E. EWING AND M. F. WHEELER, THE APPROXIMATION OF THE PRESSURE BY A MIXED METHOD IN THE SIMULATION OF MISCIBLE DISPLACEMENT, *RAIRO Analyse numérique*, tome 17, (1983), pp. 17-33.
- [17] J. DOUGLAS, R. E. EWING AND M. F. WHEELER, A TIME-DISCRETIZATION PROCEDURE FOR A MIXED FINITE ELEMENT APPROXIMATION OF MISCIBLE DISPLACEMENT IN POROUS MEDIA, *RAIRO - Analyse numérique*, tome 17, (1983), pp. 249-265.
- [18] F. HECHT, NEW DEVELOPMENT IN FREEFEM++, *Journal of Numerical Mathematics*, **20**, (2012), pp. 251-266.
- [19] J. LI, B. RIVIÈRE AND N.J. WALKINGTON, CONVERGENCE OF A HIGH ORDER METHOD IN TIME AND SPACE FOR THE MISCIBLE DISPLACEMENT EQUATIONS. *ESAIM: Mathematical Modelling and Numerical Analysis*, **49**, (2015) pp. 953-976.
- [20] B. RIVIÈRE AND N.J. WALKINGTON, CONVERGENCE OF A DISCONTINUOUS GALERKIN METHOD FOR THE MISCIBLE DISPLACEMENT EQUATION UNDER LOW REGULARITY. *SIAM J. Numer. Anal.*, **49(3)**, (2011) pp. 1085-1110.
- [21] J. E. ROBERTS AND J.-M. THOMAS, MIXED AND HYBRID METHODS. *Handbook of Numerical Analysis, Finite Element Methods (Part I)*, **II**, North-Holland, (1991), pp. 523-637.
- [22] L.R. SCOTT AND S. ZHANG, FINITE ELEMENT INTERPOLATION OF NONSMOOTH FUNCTIONS SATISFYING BOUNDARY CONDITIONS. *Math. Comp.*, **54**, (1990), pp. 483-493.
- [23] D. VASSILEV AND I. YOTOV, COUPLING STOKES-DARCY FLOW WITH TRANSPORT. *SIAM J. Sci. Comput.*, **31 (5)**, (2009), pp. 3661-3684.