



HAL
open science

Permutation-invariant log-Euclidean geometries on full-rank correlation matrices

Yann Thanwerdas

► **To cite this version:**

Yann Thanwerdas. Permutation-invariant log-Euclidean geometries on full-rank correlation matrices. 2022. hal-03878729v1

HAL Id: hal-03878729

<https://hal.science/hal-03878729v1>

Preprint submitted on 30 Nov 2022 (v1), last revised 24 Nov 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PERMUTATION-INVARIANT LOG-EUCLIDEAN GEOMETRIES ON FULL-RANK CORRELATION MATRICES *

YANN THANWERDAS[†]

Abstract. There is a growing interest in defining specific tools on correlation matrices which depart from those suited to SPD matrices. Several geometries have been defined on the open elliptope of full-rank correlation matrices: some are permutation-invariant, some others are log-Euclidean, i.e. diffeomorphic to a Euclidean space. In this work, we merge these two properties by defining the families of off-log metrics and log-scaled metrics. Firstly, we prove that the recently introduced off-log bijection is a diffeomorphism, allowing to pullback (permutation-invariant) inner products. We introduce the “cor-inverse” involution on the open elliptope which can be seen as analogous to the inversion of SPD matrices. We show that off-log metrics are not inverse-consistent. That is why secondly, we define the log-scaling diffeomorphism between the open elliptope and the vector space of symmetric matrices with null row sums. This map is based on the congruence action of positive diagonal matrices on SPD matrices, more precisely on the existence and uniqueness of a “scaling”, i.e. an SPD matrix with unit row sums within an orbit. Thanks to this multiplicative approach, log-scaled metrics are inverse-consistent. We provide the main Riemannian operations in closed form for the two families modulo the computation of the respective bijections.

Key words. SPD matrices, elliptope, correlation matrices, log-Euclidean metric, permutation-invariant, cor-inversion, off-log metric, log-scaled metric, quotient-affine metric

AMS subject classifications. 15B48, 15A63, 53B20, 53C22, 58D17, 53-08, 53A04, 62H20.

1. Introduction. In many domains such as Diffusion Tensor Imaging, Brain-Computer Interfaces, brain connectomes or radar signals, the data are time series which are often represented by their covariance matrices. They encode the dependence between the variables and the scale of intensity of these variables. Many Riemannian geometries were proposed to compute with covariance matrices with more natural tools than the Euclidean ones. The use of the affine-invariant metric was shown to outperform many results based on Euclidean metrics such as fiber reconstruction in DTI [29], movement classification in BCI [5] or detection of brain functional connectivity [35]. Shortly after, the log-Euclidean metric [4, 12] was shown to be a more efficient alternative to the affine-invariant metric with similar results. The Bures-Wasserstein metric was also proposed to deal with low-rank matrices since the two previous ones are only defined on the space $\text{Sym}^+(n)$ of Symmetric Positive Definite (SPD) matrices. All these metrics belong to the wide families of kernel metrics [15] and $O(n)$ -invariant metrics [33]. Non $O(n)$ -invariant metrics were also proposed such as the Cholesky [36, 14], log-Euclidean-Cholesky [23] and log-Cholesky [24] metrics.

In the previously cited domains and in other ones such as phylogenetic trees [13] or Gaussian graphical networks [22, 11], another possible and natural choice to represent the data is the correlation matrix instead of the covariance matrix. However, the geometries of correlation matrices have been much less studied. Hence they are often considered as covariance matrices on which one can use the classical tools. Nevertheless, these tools are not adapted to correlation matrices, at least the $O(n)$ -invariant ones. Indeed, firstly, the manifold of full-rank correlation matrices is not stable by the congruence action of the orthogonal group so this action has no sense for them. Secondly, it is not a totally geodesic submanifold for neither of the noted $O(n)$ -invariant metrics on SPD matrices, except the Euclidean metric. This motivates the study of

*Preprint, February 19th, 2022.

[†]Université Côte d’Azur and Inria, Epione Project Team (yann.thanwerdas@inria.fr).

intrinsic geometries of correlation matrices. Such geometries could also have a great impact on applications with covariance matrices since they would provide product metrics with one part on diagonal matrices and the other part on correlation matrices. It would thus allow to decouple the scale of the variables from the dependence between the variables.

Among the geometries proposed on correlation matrices, one involves a surjection from a product of spheres [30, 19]. It is an orbit space [1], this construction is quite analogous to the Bures-Wasserstein geometry of covariance matrices. However, to our knowledge, it has not been precisely described yet. A metric space structure called the Hilbert geometry relies on the convexity of the set [27]. Among the Riemannian structures, the recently introduced quotient-affine metric is obtained by taking the quotient of the affine-invariant metric under the congruence action of positive diagonal matrices $\star : (\Delta, \Sigma) \in \text{Diag}^+(n) \times \text{Sym}^+(n) \mapsto \Delta \Sigma \Delta \in \text{Sym}^+(n)$ [9, 32]. Indeed, full-rank correlation matrices can be seen as the orbits of this action so the space $\text{Cor}^+(n)$ of such matrices is the quotient manifold $\text{Sym}^+(n)/\text{Diag}^+(n)$ and any invariant metric on $\text{Sym}^+(n) \simeq \text{Cor}^+(n) \times \text{Diag}^+(n)$ descends to a Riemannian metric on $\text{Cor}^+(n)$. These constructions have the common property to be invariant under permutations. It means that the statistical analysis is invariant under reordering the variables, which can be a relevant hypothesis when the order is arbitrary. When the order of the variables is meaningfully chosen depending on the application (e.g. for auto-correlation matrices), other Riemannian metrics that are not permutation-invariant can be considered. The metrics proposed in [34] provide a Hadamard structure or even a vector space structure, which are very convenient for computing with correlation matrices.

Given this short survey on geometries of correlation matrices, there is an obvious gap to fill in: no permutation-invariant log-Euclidean metrics have been derived yet on the space $\text{Cor}^+(n)$ of full-rank correlation matrices. By *log-Euclidean*, we mean the pullback of an inner product on a vector space \mathcal{V} by a diffeomorphism referred to as a logarithm and denoted $\text{Log} : \text{Cor}^+(n) \rightarrow \mathcal{V}$. In this work, we propose two approaches to define such a diffeomorphism. The first one is based on a recent bijective parametrization of full-rank correlation matrices by the space $\text{LT}^0(n)$ of lower triangular matrices with null diagonal introduced by Archakov and Hansen [3]. The second one is entirely new. We rephrase their framework to present the two approaches in a similar way to facilitate the comparison between them and the comprehension of the second one by analogy with the first one. These two methods are respectively summarized in Tables 1.1 and 1.2 and explained below.

1.1. Results and organization of the paper. In the remainder of this section, we introduce the necessary notations. We also define a natural involution on the open ellipsope called the cor-inversion, which allows to define a notion of inverse-consistency for Riemannian metrics.

In [3], Archakov and Hansen show the claim in Table 1.1, that is for all symmetric matrix S , there exists a unique diagonal matrix $D = \mathcal{D}(S)$ such that $\exp(D + S)$ is a full-rank correlation matrix. Thus it defines a surjective map $\pi : S \in \text{Sym}(n) \mapsto \exp \circ (\mathcal{D}(S) + S) \in \text{Cor}^+(n)$ which is equivariant under permutations, and a bijective map $L \in \text{LT}^0(n) \mapsto \pi(L + L^\top) \in \text{Cor}^+(n)$. We astutely replace the space $\text{LT}^0(n)$ by the space $\mathcal{V} = \text{Hol}(n)$ of symmetric matrices with null diagonal (which is of same dimension) so that the restriction $\text{Exp} = \pi|_{\mathcal{V}} : \mathcal{V} \rightarrow \text{Cor}^+(n)$ is also equivariant under permutations. Note that $\text{Hol}(n)$ is actually the tangent space of $\text{Cor}^+(n)$. Moreover, we trivially observe that π is invariant by the additive action of a diagonal matrix. Our contribution is to show that the bijection Exp is a smooth diffeomorphism, to

	Additive approach: off-log diffeomorphism
Action	$+$: $\begin{cases} \text{Diag}(n) \times \text{Sym}(n) & \longrightarrow \text{Sym}(n) \\ (D, S) & \longmapsto D + S \end{cases}$
Claim	$\forall S \in \text{Sym}(n), \exists! D := \mathcal{D}(S) \in \text{Diag}(n) :$ $\exp(D + S) \in \text{Cor}^+(n)$
Status	Claim proved by Archakov and Hansen [3]
Diagram	$ \begin{array}{ccc} & & \text{Cor}^+(n) \\ & \swarrow \text{Log} = \text{Off} \circ \log & \nearrow \\ & \text{Hol}(n) & \xrightarrow{\text{Exp}} \\ & \searrow & \uparrow \text{exp} \circ (\mathcal{D} + \text{Id}_{\text{Sym}(n)}) \\ & \text{Sym}(n) & \xrightarrow{\mathcal{D}} \text{Diag}(n) \end{array} $
Outline	Section 2.1. Definition of \mathcal{D} Section 2.2. Definition of pullback metrics

Table 1.1: Additive approach to define permutation-invariant log-Euclidean metrics on $\text{Cor}^+(n)$, based on [3].

	Multiplicative approach: cor-exp diffeomorphism
Action	\star : $\begin{cases} \text{Diag}^+(n) \times \text{Sym}^+(n) & \longrightarrow \text{Sym}^+(n) \\ (\Delta, \Sigma) & \longmapsto \Delta \Sigma \Delta \end{cases}$
Claim	$\forall \Sigma \in \text{Sym}^+(n), \exists! \Delta := \mathcal{D}^*(\Sigma) \in \text{Diag}^+(n) :$ $\log(\Delta \star \Sigma) \in \mathcal{V}^*$
Status	Claim proved for $\mathcal{V}^* = \text{Row}_0(n)$ in Section 3
Diagram	$ \begin{array}{ccc} \text{Cor}^+(n) & \hookrightarrow \text{Sym}^+(n) & \xrightarrow{\mathcal{D}^*} \text{Diag}^+(n) \\ & \searrow \text{Log}^* & \downarrow \log \circ (\mathcal{D}^* \star \text{Id}_{\text{Sym}^+(n)}) \\ & & \mathcal{V}^* \\ & \swarrow \text{Exp}^* = \text{Cor} \circ \exp & \nearrow \end{array} $
Outline	Section 3. Def. of \mathcal{D}^* (choice of appropriate \mathcal{V}^*) Section 4. Definition of pullback metrics

Table 1.2: Multiplicative approach to define permutation-invariant log-Euclidean metrics on $\text{Cor}^+(n)$. This is new. The vector space \mathcal{V}^* is stable by permutations and satisfies $\text{Sym}(n) = \mathcal{V}^* \oplus \text{Diag}(n)$.

define by pullback the family of *off-log metrics* and to provide all the Riemannian operations in closed form modulo the computation of \mathcal{D} . This additive approach is summarized in Table 1.1 and exposed in Section 2.

Our second approach consists in inverting the roles played by the vector spaces $\text{Sym}(n) = \mathcal{V} \oplus \text{Diag}(n)$ and the manifolds $\text{Sym}^+(n) = \text{Cor}^+(n) \times \text{Diag}^+(n)$, as well as the matrix exponential and the matrix logarithm, and especially to replace the additive action of $\text{Diag}(n)$ on $\text{Sym}(n)$ by the congruence action $\star : (\Delta, \Sigma) \in \text{Diag}^+(n) \times \text{Sym}^+(n) \mapsto \Delta \Sigma \Delta \in \text{Sym}^+(n)$. In this work, we find a vector space \mathcal{V}^* such that for all $\Sigma \in \text{Sym}^+(n)$, there exists a unique $\Delta = \mathcal{D}^*(\Sigma) \in \text{Diag}^+(n)$ such that $\log(\Delta \Sigma \Delta) \in \mathcal{V}^*$. This allows to define the surjective map $\pi^* : \Sigma \in \text{Sym}^+(n) \mapsto \log(\mathcal{D}^*(\Sigma) \star \Sigma) \in \mathcal{V}^*$ and the bijective map $\text{Log}^* = \pi^*|_{\text{Cor}^+(n)} : \text{Cor}^+(n) \longrightarrow \mathcal{V}^*$ which are equivariant under permutations. This multiplicative approach is summarized in Table 1.2 and exposed in Section 3. One major advantage of this multiplicative approach is that it intrinsically respects the structure of correlation matrices since $\text{Cor}^+(n) = \text{Sym}^+(n)/\text{Diag}^+(n)$, contrarily to the additive approach. The main con-

sequence is the compatibility with the inversion, i.e. $\pi^*(C^{-1}) = -\pi^*(C)$ for all $C \in \text{Cor}^+(n)$.

More precisely, we try to prove the claim with $\mathcal{V}^* = \text{Hol}(n)$ and $\mathcal{V}^* = \text{Row}_0(n)$, that are the vector spaces of symmetric matrices respectively with null diagonal and null row sums. With the first choice, we only manage to prove the existence. We actually prove that the uniqueness would imply the uniqueness of the Riemannian logarithm at identity of the quotient-affine metric mentioned above, which is an open problem. This is a secondary contribution that relates two problems on full-rank correlation matrices. In contrast, we prove the claim with $\mathcal{V}^* = \text{Row}_0(n)$. Indeed, we show that $\exp(\text{Row}_0(n)) = \text{Row}_{\mathbf{1}}^+(n)$, where $\text{Row}_{\mathbf{1}}^+(n)$ is the submanifold of SPD matrices with unit row sums. This reduces our question to the famous problem of scaling an SPD matrix to prescribed row sums by congruence of a positive diagonal matrix: for all SPD matrix Σ , does there exist a positive diagonal matrix $\Delta = \mathcal{D}^*(\Sigma)$ such that $\Delta\Sigma\Delta \in \text{Row}_{\mathbf{1}}^+(n)$. The answer is yes [26, 18] so the claim is true.

In Section 4, similarly to the additive approach, we prove that the bijection Log^* is a smooth diffeomorphism and we define by pushforward the family of *log-scaled metrics*. Contrarily to off-log metrics, they are inverse-consistent. We provide all the Riemannian operations in closed form modulo the computation of \mathcal{D}^* , that is the computation of the scaling of an SPD matrix. We conclude in Section 5.

1.2. Notations.

1.2.1. Matrices.

Tables 1.3 and 1.4 summarize our notations for matrix spaces. We also use the following constant and linear operators on vectors and matrices:

- $\mathbf{1} \in \mathbb{R}^n$ is the vector with all entries equal to 1;
- $\text{diag} : \mathbb{R}^n \rightarrow \text{Diag}(n)$ makes a diagonal matrix from a vector;
- $\text{sum} : \mathbb{R}^n \rightarrow \mathbb{R}$ computes the sum of the entries of a vector;
- $\text{Diag} : \text{Mat}(n) \rightarrow \text{Diag}(n)$ extracts the diagonal matrix from a matrix;
- $\text{Off} : \text{Mat}(n) \rightarrow \ker \text{Diag}$ subtracts the diagonal matrix from a matrix;
- $\text{Sum} : \text{Mat}(n) \rightarrow \mathbb{R}$ computes the sum of entries of a matrix;
- $\bullet : \text{Mat}(n) \times \text{Mat}(n) \rightarrow \text{Mat}(n)$ computes the Schur (entry-wise) product of matrices.

Squared of size n	$\text{Mat}(n) = \{M = [M_{ij}]_{1 \leq i, j \leq n} M_{ij} \in \mathbb{R}\}$
Skew-symmetric	$\text{Skew}(n) = \{Y \in \text{Mat}(n) Y^\top = -Y\}$
Symmetric	$\text{Sym}(n) = \{X \in \text{Mat}(n) X^\top = X\}$
Diagonal	$\text{Diag}(n) = \{\text{Diag}(X) X \in \text{Mat}(n)\}$
Symmetric hollow (null diagonal)	$\text{Hol}(n) = \{X \in \text{Sym}(n) \text{Diag}(X) = 0\}$
Symmetric null-row-sum	$\text{Row}_0(n) = \{X \in \text{Sym}(n) X\mathbf{1} = 0\}$

Table 1.3: Matrix vector spaces

Invertible	$\text{GL}(n) = \{A \in \text{Mat}(n) \det(A) \neq 0\}$
Orthogonal	$\text{O}(n) = \{A \in \text{GL}(n) AA^\top = I_n\}$
Symmetric Positive Definite	$\text{Sym}^+(n) = \{AA^\top A \in \text{GL}(n)\}$
Positive diagonal	$\text{Diag}^+(n) = \text{Sym}^+(n) \cap \text{Diag}(n)$
Full-rank correlation	$\text{Cor}^+(n) = \{C \in \text{Sym}^+(n) \text{Diag}(C) = I_n\}$
SPD unit-row-sum	$\text{Row}_{\mathbf{1}}^+(n) = \{\Sigma \in \text{Sym}^+(n) \Sigma\mathbf{1} = \mathbf{1}\}$

Table 1.4: Matrix manifolds

We recall the definition of the congruence action $\star : (A, M) \in \text{GL}(n) \times \text{Mat}(n) \mapsto AMA^\top \in \text{Mat}(n)$. Since the permutation group $\mathfrak{S}(n)$ is a subgroup of the orthogonal group via the canonical injection $\sigma \in \mathfrak{S}(n) \mapsto P_\sigma = [\delta_{i, \sigma(j)}]_{1 \leq i, j \leq n} \in \text{O}(n)$, the permutation action $(\sigma, M) \in \mathfrak{S}(n) \times \text{Mat}(n) \mapsto \sigma \cdot M = P_\sigma M P_\sigma^\top \in \text{Mat}(n)$ is the congruence action of the subgroup $\mathfrak{S}(n)$.

We recall the definition of the matrix exponential map $\exp : M \in \text{Mat}(n) \rightarrow \sum_{k=0}^{+\infty} \frac{1}{k!} M^k \in \text{GL}(n)$ which is a smooth map. Its restriction to symmetric matrices is a smooth diffeomorphism onto SPD matrices, $\exp : \text{Sym}(n) \rightarrow \text{Sym}^+(n)$. The symmetric matrix logarithm is its smooth inverse, $\log : \text{Sym}^+(n) \rightarrow \text{Sym}(n)$. The computation of \exp , \log and their differentials is particularly simple modulo eigenvalue decomposition. Given $\Sigma = PDP^\top \in \text{Sym}^+(n)$, $X = Q\Delta Q^\top, Y \in \text{Sym}(n)$ where $P, Q \in \text{O}(n)$, $D \in \text{Diag}^+(n)$ and $\Delta \in \text{Diag}(n)$:

$$(1.1) \quad \exp(X) = Q \exp(\Delta) Q^\top,$$

$$(1.2) \quad \log(\Sigma) = P \log(D) P^\top,$$

$$(1.3) \quad d_X \exp(Y) = Q \left([\exp^{[1]}(\delta_i, \delta_j)]_{1 \leq i, j \leq n} \bullet (Q^\top Y Q) \right) Q^\top,$$

$$(1.4) \quad d_\Sigma \log(Y) = P \left([\log^{[1]}(d_i, d_j)]_{1 \leq i, j \leq n} \bullet (P^\top Y P) \right) P^\top,$$

where $f^{[1]}(x, y) = \begin{cases} \frac{f(x)-f(y)}{x-y} & \text{if } x \neq y \\ f'(x) & \text{if } x = y \end{cases}$ is the first divided difference of $f \in \{\exp, \log\}$ [7]. In other words, the maps $\exp, \log, d\exp, d\log$ are $\text{O}(n)$ -equivariant, and therefore $\mathfrak{S}(n)$ -equivariant.

1.2.2. Correlation matrices. The manifold of full-rank correlation matrices is called the open ellipsope. It is relatively open in $\text{Sym}(n)$, i.e. open in $I_n + \text{Hol}(n)$.

We introduce the following smooth submersion from SPD matrices to full-rank correlation matrices, $\text{Cor} : \Sigma \in \text{Sym}^+(n) \mapsto \text{Diag}(\Sigma)^{-1/2} \Sigma \text{Diag}(\Sigma)^{-1/2} \in \text{Cor}^+(n)$. Given $\Sigma \in \text{Sym}^+(n)$, $X \in \text{Sym}(n)$, denoting $\Delta = \text{Diag}(\Sigma)^{-1/2}$, its differential is:

$$(1.5) \quad d_\Sigma \text{Cor}(X) = \Delta \left[X - \frac{1}{2} (\Delta^2 \text{Diag}(X) \Sigma + \Sigma \text{Diag}(X) \Delta^2) \right] \Delta.$$

We introduce a notation for equicorrelation matrices $C(\rho) = (1 - \rho)I_n + \rho \mathbf{1}\mathbf{1}^\top \in \text{Cor}^+(n)$ where $\rho \in (-\frac{1}{n-1}, 1)$. Given a correlation matrix $C \in \text{Cor}^+(n)$, there exist partitions of n , i.e. sets $I = \{i_1, \dots, i_p\}$ satisfying $i_1, \dots, i_p \geq 1$ and $i_1 + \dots + i_p = n$, partitioning the matrix C into equicorrelation diagonal blocks and constant off-diagonal blocks. The signature of C is the maximum I_C of such sets I with respect to the natural order on partitions of n . We say that C is a block equicorrelation matrix of signature I_C (see Table 1.5). For example, an equicorrelation matrix is a block equicorrelation matrix with signature $\{n\}$. The maps introduced in this paper preserve the signature.

$$C(\rho) = \begin{pmatrix} 1 & \rho & \cdots & \rho \\ \rho & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \rho \\ \rho & \cdots & \rho & 1 \end{pmatrix} \left| \begin{pmatrix} C(\rho_1) & \rho_{12} \mathbf{1}\mathbf{1}^\top & \cdots & \rho_{1p} \mathbf{1}\mathbf{1}^\top \\ \rho_{12} \mathbf{1}\mathbf{1}^\top & C(\rho_2) & \ddots & \vdots \\ \vdots & \ddots & \ddots & \rho_{p-1,p} \mathbf{1}\mathbf{1}^\top \\ \rho_{1p} \mathbf{1}\mathbf{1}^\top & \cdots & \rho_{p-1,p} \mathbf{1}\mathbf{1}^\top & C(\rho_p) \end{pmatrix} \right.$$

Table 1.5: Equicorrelation and block equicorrelation matrices

The elliptope is stable by the permutation action, it is not stable by the congruence action of the orthogonal group $O(n)$. It is not stable by inversion either. However, the correlation matrix of its inverse, namely $\text{Cor}(C^{-1})$, contains the same information as the partial correlation matrix defined below.

DEFINITION 1.1 (Cor-inversion, partial correlation matrix).

- The cor-inversion is the smooth involution $\mathcal{I} : C \in \text{Cor}^+(n) \mapsto \text{Cor}(C^{-1}) \in \text{Cor}^+(n)$.
- The partial correlation matrix of an SPD matrix $\Sigma \in \text{Sym}^+(n)$ is the matrix $\Gamma \in \text{Sym}(n)$ defined by $\Gamma = I_n - \text{Off}(\mathcal{I}(C)) = 2I_n - \mathcal{I}(C)$, where $C = \text{Cor}(\Sigma)$.

The bijective parametrization $C \mapsto \Gamma$ is used in the theory of stationary stochastic processes where the (potentially infinite and complex) matrices are Toeplitz. The set of partial correlation coefficients (along with the common variance) is considered as an alternative “representation of the second-order statistics” [8, Section II.B.5] of the process with respect to the traditional “auto-correlation” (or auto-covariance) function. This characterization is used in signal processing, especially in radar signal processing where the manifold of SPD Toeplitz is traditionally endowed with the Poincaré polydisk geometry [6]. In Gaussian graphical networks, the partial correlation between two variables indicates the correlation between them conditionally to the other variables. Thus the partial correlations are the weights of the arrows in the network [22, 20, 11]. This approach is applied in many domains such as genomics [10, 28] or brain connectomics [25]. Hence the importance of partial correlations confirms that the cor-inversion is a relevant concept.

We consider the cor-inversion as analogous to the matrix inversion for SPD matrices $\text{inv} : \text{Sym}^+(n) \rightarrow \text{Sym}^+(n)$. The cor-inversion commutes with permutations on full-rank correlation matrices as well as the inversion commutes with the congruence by $O(n)$ on SPD matrices. Moreover, for all signature I , the space of block equicorrelation matrices of signature I is stable by the cor-inversion. In addition, in relation to the problem of this paper, we have the following result.

THEOREM 1.2 (Compatibility between the multiplicative approach and the cor-inversion). *Let \mathcal{V}^* be a vector space stable by permutations and such that $\text{Sym}(n) = \mathcal{V}^* \oplus \text{Diag}(n)$. We assume that the claim in Table 1.1 is true, i.e. for all $C \in \text{Cor}^+(n)$, there exists a unique $\Delta \in \text{Diag}^+(n)$ such that $\log(\Delta C \Delta) \in \mathcal{V}^*$. This defines the inverse bijections $\text{Log}^* : C \in \text{Cor}^+(n) \mapsto \log(\Delta C \Delta) \in \mathcal{V}^*$ and $\text{Exp}^* = \text{Cor} \circ \text{exp} : \mathcal{V}^* \rightarrow \text{Cor}^+(n)$. Then we automatically have $\text{Log}^*(\mathcal{I}(C)) = -\text{Log}^*(C)$, i.e. the following diagram commutes.*

$$(1.6) \quad \begin{array}{ccc} \text{Cor}^+(n) & \xrightarrow{\mathcal{I}} & \text{Cor}^+(n) \\ \downarrow \text{Log}^* & & \downarrow \text{Log}^* \\ \mathcal{V}^* & \xrightarrow{-\text{Id}} & \mathcal{V}^* \end{array}$$

Proof. Note that for all $\Sigma \in \text{Sym}^+(n)$, we have $\text{Cor}(\Sigma^{-1}) = \text{Cor}(\text{Cor}(\Sigma)^{-1})$. Indeed, if $\Sigma = DCD$ with $C = \text{Cor}(\Sigma) \in \text{Cor}^+(n)$, then we have $\text{Cor}(\Sigma^{-1}) = \text{Cor}(D^{-1}C^{-1}D^{-1}) = \text{Cor}(C^{-1})$. Therefore, for all $X \in \mathcal{V}^*$ we have $\text{Exp}^*(-X) = \text{Cor}(\text{exp}(-X)) = \text{Cor}(\text{exp}(X)^{-1}) = \text{Cor}(\text{Exp}^*(X)^{-1}) = \mathcal{I}(\text{Exp}^*(X))$. Thus with $C = \text{Exp}^*(X)$, we have $\text{Log}^*(\mathcal{I}(X)) = -\text{Log}^*(C)$. \square

Otherwise said, the multiplicative approach is automatically compatible with the cor-inversion. This is due to the use of the congruence action of positive diagonal

matrices on SPD matrices instead of the additive action of diagonal matrices on symmetric matrices. Indeed, the former is intrinsically related to the definition of a correlation matrix. On the contrary, we can expect that the bijections built via the additive approach are not compatible with the cor-inversion in general.

Thus, if one finds a vector space \mathcal{V}^* satisfying the claim and if the bijections are smooth, the log-Euclidean metrics defined by pullback will automatically be inverse-consistent. This is quite satisfying for log-Euclidean metrics on full-rank correlation matrices in analogy with SPD matrices.

2. Permutation-invariant log-Euclidean metrics via the off-log diffeomorphism. In this section, we rephrase the framework of [3] (Section 2.1) to ease the comprehension of the next sections by analogy. In Section 2.2, we prove that the bijection they define, that we call the off-log bijection and that we denote $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$, is actually a diffeomorphism. It allows to pullback inner products on full-rank correlation matrices. Since the off-log diffeomorphism is equivariant under permutations, we give a characterization of permutation-invariant inner products on $\text{Hol}(n)$ so that their pullbacks provide permutation-invariant log-Euclidean metrics on $\text{Cor}^+(n)$. Then, we detail the Riemannian operations of these metrics. We prove that, as expected, the log-Euclidean metrics such defined are not inverse-consistent with respect to the cor-inversion. In Section 2.3, we simply recall the algorithm of [3] to compute the inverse diffeomorphism $\text{Exp} = \text{Log}^{-1}$, the speed of convergence and the complexity.

2.1. The off-log bijection. Theorem 2.1 states that the claim in Table 1.1 is true. It allows to define the off-log bijection $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$. Theorem 2.2 states some interesting properties of the off-log bijection. These results are due to Archakov and Hansen [3].

THEOREM 2.1 (Definition of \mathcal{D}). [3] *For all $S \in \text{Sym}(n)$, there exists a unique $D \in \text{Diag}(n)$ such that $\exp(D + S) \in \text{Cor}^+(n)$. This allows to define:*

- *the surjective map $\mathcal{D} : S \in \text{Sym}(n) \mapsto D \in \text{Diag}(n)$,*
- *the surjective map $\pi : S \in \text{Sym}(n) \mapsto \exp(\mathcal{D}(S) + S) \in \text{Cor}^+(n)$ which is invariant under the additive group action $+: \text{Diag}(n) \times \text{Sym}(n) \rightarrow \text{Sym}(n)$,*
- *the bijective map $\text{Exp} = \pi|_{\text{Hol}(n)} : \text{Hol}(n) \rightarrow \text{Cor}^+(n)$ (note that $\pi = \text{Exp} \circ \text{Off}$),*
- *the smooth bijective inverse map $\text{Log} = \text{Exp}^{-1} = \text{Off} \circ \log : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ that we call the off-log bijection.*

THEOREM 2.2 (Properties of the off-log bijection). [3]

1. (Equivariance) *Log and Exp are equivariant under permutations.*
2. (Equicorrelation matrix) *$\text{Log}(C(\rho)) = \frac{1}{n} \ln \left(\frac{1+(n-1)\rho}{1-\rho} \right) (\mathbf{1}\mathbf{1}^\top - I_n)$ for all $\rho \in (-\frac{1}{n-1}, 1)$. In dimension $n = 2$, $\text{Log}(C(\rho)) = \begin{bmatrix} 0 & F(\rho) \\ F(\rho) & 0 \end{bmatrix}$ where $F(\rho) = \frac{1}{2} \log(\frac{1+\rho}{1-\rho}) \in \mathbb{R}$ is the Fisher transformation of the correlation coefficient $\rho \in (-1, 1)$.*
3. (Block equicorrelation matrix) *If C is a block equicorrelation matrix of signature $I = \{i_1, \dots, i_p\}$, then $\text{Log}(C)$ is a block symmetric hollow matrix of signature I with multiples of $\mathbf{1}_{i_j}\mathbf{1}_{i_j}^\top - I_{i_j}$ on diagonal blocks and multiples of $\mathbf{1}_{i_j}\mathbf{1}_{i_k}^\top$ on off-diagonal blocks.*
4. (Generalization) *For all $\Delta \in \text{Diag}^+(n)$, for all $S \in \text{Sym}(n)$, there exists a unique $D \in \text{Diag}(n)$ such that $\text{Diag}(\exp(D + S)) = \Delta$.*

Note that Theorem 2.1 is a particular case of Theorem 2.2 item 4 with $\Delta = I_n$.

The result in dimension 2 was stated as a motivation in [3] to use the map $\text{Log} = \text{Off} \circ \log$ in higher dimensions since it gives in dimension 2 a well known transformation of the correlation coefficient. Interestingly, the same coefficient appears in dimension 2 for the quotient-affine metric [32].

By analogy with the symmetric matrix logarithm $\log : \text{Sym}^+(n) \rightarrow \text{Sym}(n)$ satisfying $\log(\Sigma^{-1}) = -\log(\Sigma)$, one could expect that the off-log bijection $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ “commutes with inversion”, i.e. satisfies $\text{Log}(\mathcal{I}(C)) = -\text{Log}(C)$. We show that it is not the case as we argued in the introduction.

THEOREM 2.3 (Incompatibility between cor-inversion and off-log bijection). *Let $n \geq 3$. There exists $C \in \text{Cor}^+(n)$, such that $\text{Log}(\mathcal{I}(C)) \neq -\text{Log}(C)$. Otherwise said, the following diagram does not commute.*

$$(2.1) \quad \begin{array}{ccc} \text{Cor}^+(n) & \xrightarrow{\mathcal{I}} & \text{Cor}^+(n) \\ \downarrow \text{Log} & & \downarrow \text{Log} \\ \text{Hol}(n) & \xrightarrow{-\text{Id}} & \text{Hol}(n) \end{array}$$

Proof. It is easy to see it numerically. For a formal proof, one can look for a matrix $C \in \text{Cor}^+(3)$ such that $\log(C)$ and $\log(\mathcal{I}(C))$ are easy to compute manually. We propose the following example with $x = \frac{1}{\sqrt{7}}$: $C = \begin{bmatrix} 1 & x & -x \\ x & 1 & 0 \\ -x & 0 & 1 \end{bmatrix} = PDP^\top$ with $P = \frac{1}{2} \begin{bmatrix} 0 & \sqrt{2} & \sqrt{2} \\ \sqrt{2} & 1 & -1 \\ \sqrt{2} & -1 & 1 \end{bmatrix}$ and $D = \text{diag}(1, 1 + \frac{\sqrt{2}}{\sqrt{7}}, 1 - \frac{\sqrt{2}}{\sqrt{7}})$. Then $\mathcal{I}(C) = \begin{bmatrix} 1 & -a & a \\ -a & 1 & -a^2 \\ a & -a^2 & 1 \end{bmatrix} = Q\Delta Q^\top$ with $a = \frac{x}{\sqrt{1-x^2}} = \frac{1}{\sqrt{6}}$, $Q = \frac{1}{\sqrt{14}} \begin{bmatrix} 0 & \sqrt{6} & 2\sqrt{2} \\ \sqrt{7} & -2 & \sqrt{3} \\ \sqrt{7} & 2 & -\sqrt{3} \end{bmatrix}$ and $\Delta = \frac{1}{6} \text{diag}(5, 10, 3)$. Thus:

$$\begin{aligned} [\text{Log}(C) + \text{Log}(\mathcal{I}(C))]_{12} &= [\log(C) + \log(\mathcal{I}(C))]_{12} \\ &= [P \log(D) P^\top + Q \log(\Delta) P^\top]_{12} \\ &= \sum_{k=1}^3 (\ln(D_{kk}) P_{1k} P_{2k} + \ln(\Delta_{kk}) Q_{1k} Q_{2k}) \\ &= \frac{1}{2\sqrt{2}} \ln \left(\frac{\sqrt{7} + \sqrt{2}}{\sqrt{7} - \sqrt{2}} \right) + \frac{\sqrt{6}}{7} \ln \left(\frac{3}{10} \right) > 0. \end{aligned}$$

For $n \geq 4$, it suffices to take the block diagonal matrix $\text{Diag}(C, I_{n-3})$. \square

This incompatibility is one of the justifications of the multiplicative approach that we present in Section 3. Still, this bijection $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ remains a very nice tool that allows to define permutation-invariant log-Euclidean metrics on full-rank correlation matrices. Let us show this.

2.2. Permutation-invariant pullback metrics via the off-log diffeomorphism. This section is part of our contributions. We prove that the off-log bijection $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ is actually a diffeomorphism (Section 2.2.1). Then we characterize all permutation-invariant inner products on $\text{Hol}(n)$ (Section 2.2.2) and we pull them back to permutation-invariant log-Euclidean metrics on $\text{Cor}^+(n)$ (Section 2.2.3).

2.2.1. The off-log bijection is a diffeomorphism.

THEOREM 2.4 ($\text{Log} = \text{Off} \circ \log$ is a diffeomorphism). *The off-log bijection $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ is a smooth diffeomorphism. We give the differentials of Log and*

Exp in function of the differentials of the symmetric matrix logarithm and exponential maps log and exp. For all $C \in \text{Cor}^+(n)$ and $S, X, Y \in \text{Hol}(n)$:

$$(2.2) \quad d_C \text{Log}(X) = \text{Off}(d_C \log(X)),$$

$$(2.3) \quad d_S \text{Exp}(Y) = d_{\log(\text{Exp}(S))} \exp(Y - \text{Diag}(\text{Exp}(S)Y)).$$

Proof. It suffices to show that \mathcal{D} is smooth. We use the implicit function theorem with the smooth map $\Phi : (S, D) \in \text{Hol}(n) \times \text{Diag}(n) \mapsto \text{Diag}(\exp(D + S)) - I_n \in \text{Diag}(n)$ which is such that $D = \mathcal{D}(S)$ if and only if $\Phi(S, D) = 0$. We want to show that for $(S, D) \in \Phi^{-1}(0)$, the differential of $\Phi_S : D \in \text{Diag}(n) \mapsto \Phi(S, D) \in \text{Diag}(n)$ is invertible. We introduce the intermediate smooth map $\varphi_S : D \in \text{Diag}(n) \mapsto D - \log \text{Diag} \exp(D + S) \in \text{Diag}(n)$ as in [3] so that $\Phi_S(D) = \exp(D - \varphi_S(D)) - I_n$. It is proved in the Appendix of [3] that for all $D \in \text{Diag}(n)$, the linear map $d_D \varphi_S : \text{Diag}(n) \rightarrow \text{Diag}(n)$ has its eigenvalues in $[0, 1) \subset \mathbb{R}$. For all $(S, D) \in \Phi^{-1}(0)$ and all $\Delta \in \text{Diag}(n)$, we have:

$$\begin{aligned} d_D \Phi_S(\Delta) &= \underbrace{\exp(D - \varphi_S(D))}_{I_n} (\Delta - d_D \varphi_S(\Delta)) \\ &= (\text{Id} - d_D \varphi_S)(\Delta). \end{aligned}$$

Therefore, the linear map $d_D \Phi_S = \text{Id} - d_D \varphi_S$ has its eigenvalues in $(0, 1]$ so it is invertible. Hence, the implicit function $\mathcal{D} : \text{Sym}(n) \rightarrow \text{Diag}(n)$ is smooth so $\pi = \exp \circ (\mathcal{D} + \text{Id}_{\text{Sym}(n)}) : \text{Sym}(n) \rightarrow \text{Cor}^+(n)$ is smooth. Then $\text{Exp} = \pi|_{\text{Hol}(n)} : \text{Hol}(n) \rightarrow \text{Cor}^+(n)$ is smooth and Log is a smooth diffeomorphism.

Since $\text{Log} = \text{Off} \circ \log$ and Off is linear, the differential of Log is clear. Let us compute the differential of f , \mathcal{D} and Exp. Let $S \in \text{Sym}(n)$, $C = \text{Exp}(S) \in \text{Cor}^+(n)$, $D, \Delta \in \text{Diag}(n)$, $X \in \text{Hol}(n)$ and $Y = d_C \text{Log}(X) \in \text{Hol}(n)$.

$$\begin{aligned} d_{(S,D)} \Phi(Y, \Delta) &= \text{Diag}(\exp(D + S)(\Delta + Y)) \\ &= \text{Diag}(\exp(D + S))\Delta + \text{Diag}(\exp(D + S)Y), \\ d_{(S,\mathcal{D}(S))} \Phi(Y, \Delta) &= \Delta + \text{Diag}(CY), \\ \underbrace{d_{(S,\mathcal{D}(S))} \Phi(Y, d_S \mathcal{D}(Y))}_{=d_S(\Phi \circ (\text{Id}, \mathcal{D}))(Y)=0} &= d_S \mathcal{D}(Y) + \text{Diag}(CY), \\ d_S \mathcal{D}(Y) &= -\text{Diag}(CY) = -\text{Diag}(\text{Exp}(S)Y), \\ d_S \text{Exp}(Y) &= d_{\mathcal{D}(S)+S} \exp(d_S \mathcal{D}(Y) + Y) \\ &= d_{\log(C)} \exp(Y - \text{Diag}(\text{Exp}(S)Y)). \quad \square \end{aligned}$$

2.2.2. Permutation-invariant inner products on $\text{Hol}(n)$. The characterization of permutation-invariant inner products on $\text{Hol}(n)$ can be found in [31, Example 3.8].

THEOREM 2.5 (Permutation-invariant inner products on $\text{Hol}(n)$). [31] For $n \geq 4$, permutation-invariant inner products on $\text{Hol}(n)$ are the symmetric bilinear forms associated to the following positive definite quadratic forms defined for $X \in \text{Hol}(n)$:

$$(2.4) \quad q(X) = \alpha \text{tr}(X^2) + \beta \text{Sum}(X^2) + \gamma \text{Sum}(X)^2$$

with $\alpha > 0$, $2\alpha + (n - 2)\beta > 0$ and $\alpha + (n - 1)(\beta + n\gamma) > 0$. For $n = 3$, the permutation-invariant inner products have the same form with $\alpha = 0$, i.e. $q(X) = \beta \text{Sum}(X^2) + \gamma \text{Sum}(X)^2$ with $\beta > 0$ and $\beta + 3\gamma > 0$. For $n = 2$, they have the same form with $\alpha = \beta = 0$, i.e. $q(X) = \gamma \text{Sum}(X)^2$ with $\gamma > 0$.

2.2.3. Pullback metrics via the off-log diffeomorphism.

DEFINITION 2.6 (Off-log metrics). *An off-log metric on $\text{Cor}^+(n)$ is the pullback metric of a permutation-invariant inner product characterized by a quadratic form q as in Theorem 2.5. For all $C \in \text{Cor}^+(n)$ and $X \in T_C \text{Cor}^+(n) = \text{Hol}(n)$, it writes $g_C(X, X) = q(d_C \text{Log}(X))$ where $d_C \text{Log}(X) = \text{Off}(d_C \log(X))$.*

THEOREM 2.7 (Riemannian operations of off-log metrics). *We consider an off-log metric characterized by the quadratic form q . Let $C, C', C_1, \dots, C_k \in \text{Cor}^+(n)$, $X \in \text{Hol}(n)$. The Riemannian operations of this metric are summarized in Table 2.1.*

Exponential map	$\text{Exp}_C(X) = \text{Exp}(\text{Log}(C) + d_C \text{Log}(X))$
Logarithm map	$\text{Log}_C(C') = d_{\text{Log}(C)} \text{Exp}(\text{Log}(C') - \text{Log}(C))$
Geodesic	$\gamma(t) = \text{Exp}((1-t)\text{Log}(C) + t\text{Log}(C'))$
Squared distance	$d(C, C')^2 = q(\text{Log}(C') - \text{Log}(C))$
Fréchet mean	$\bar{C} = \text{Exp}(\frac{1}{k} \sum_{i=1}^k \text{Log}(C_i))$
Curvature	$R = 0$
Parallel transport	$\Pi_{C \rightarrow C'} X = (d_{C'} \text{Log})^{-1}(d_C \text{Log}(X))$

Table 2.1: Riemannian operations of off-log metrics

Beware that the Riemannian exponential and logarithm maps only coincide with the diffeomorphisms $\text{Exp} : \text{Hol}(n) \rightarrow \text{Cor}^+(n)$ and $\text{Log} : \text{Cor}^+(n) \rightarrow \text{Hol}(n)$ at $C = I_n$ introduced in Theorem 2.1. They differ from the symmetric matrix diffeomorphisms $\text{exp} : \text{Sym}(n) \rightarrow \text{Sym}^+(n)$ and $\text{log} : \text{Sym}^+(n) \rightarrow \text{Sym}(n)$.

Therefore, the off-log diffeomorphism provides a closed-form distance between two full-rank correlation matrices (modulo the computation of a symmetric matrix logarithm, i.e. modulo an eigenvalue decomposition). Moreover, all the other Riemannian operations can be computed in closed form modulo the computation of Exp , i.e. the computation of \mathcal{D} . We recall that Archakov and Hansen [3] proved that $\mathcal{D}(S)$ for $S \in \text{Sym}(n)$ can be computed in logarithmic time complexity by defining the following sequence: $D_0 = 0$ and $D_{k+1} = \varphi_S(D_k)$ where $\varphi_S : D \in \text{Diag}(n) \mapsto D - \text{log}(\text{Diag}(\exp(D + S))) \in \text{Diag}(n)$ is a contractant map.

In this section, we recalled the main facts on the off-log parametrization introduced in [3] and we transformed it as a geometric tool to introduce log-Euclidean metrics on full-rank correlation matrices. We also formalized this tool in terms of invariance under a group action and we showed that off-log metrics are not inverse-consistent. In the next sections, we rely on this formalization to introduce the family of log-scaled metrics which are permutation-invariant, log-Euclidean and inverse-consistent.

3. The log-scaling bijection. In this section, we examine two versions of the following conjecture: for all $\Sigma \in \text{Sym}^+(n)$, there exists a unique $\Delta \in \text{Diag}^+(n)$ such that $\text{log}(\Delta \Sigma \Delta) \in \mathcal{V}^*$. This conjecture depends on \mathcal{V}^* , which is a vector space stable by permutations satisfying $\text{Sym}(n) = \mathcal{V}^* \oplus \text{Diag}(n)$. In Section 3.1, we relate the conjecture with $\mathcal{V}^* = \text{Hol}(n)$ to the problem of the quotient-affine logarithm [9, 32]. We prove the existence and we explain why the uniqueness remains difficult to prove. In Section 3.2, we explain why $\mathcal{V}^* = \text{Row}_0(n)$ is a good candidate for the conjecture to be true and in Section 3.3, we prove the conjecture thanks to a result known as

the existence and uniqueness of the *scaling* of SPD matrices [26, 18]. In Section 3.4, we give the properties of our new Euclideanization called the log-scaling bijection.

3.1. Is the conjecture true with $\mathcal{V}^* = \text{Hol}(n)$? Before relating the conjecture with $\mathcal{V}^* = \text{Hol}(n)$ to the problem of existence and uniqueness of the Riemannian logarithm at I_n of the quotient-affine metric, we recall the conjecture and the definition of the quotient-affine metric.

CONJECTURE 3.1 (The result is true with $\mathcal{V}^* = \text{Hol}(n)$). *For all $\Sigma \in \text{Sym}^+(n)$, there exists a unique matrix $\Delta \in \text{Diag}^+(n)$ such that $\log(\Delta\Sigma\Delta) \in \text{Hol}(n)$.*

DEFINITION 3.2 (Quotient-affine metric). *The quotient-affine metric on $\text{Cor}^+(n)$ is the quotient metric of the affine-invariant metric on $\text{Sym}^+(n)$ by the congruence action of $\text{Diag}^+(n)$ [9]. At I_n , the horizontal space is $\mathcal{H}_{I_n}^{\text{QA}} = \text{Hol}(n)$, the quotient-affine metric writes $g_{I_n}^{\text{QA}}(X, X) = \text{tr}(X^2)$ and the exponential map writes $\text{Exp}_{I_n}^{\text{QA}}(X) = \text{Cor}(\exp(X))$ for all $X \in \text{Hol}(n)$ [32].*

THEOREM 3.3 (Existence and equivalence of conjectures). *We define the smooth map $f : \Delta \in \text{Diag}^+(n) \mapsto d^{\text{AI}}(I_n, \Delta C \Delta)^2 = \text{tr}(\log(\Delta C \Delta)^2)$. It gives the affine-invariant squared distance between I_n and all points of the fiber $\text{Cor}^{-1}(C) = \{\Delta C \Delta \in \text{Sym}^+(n) | \Delta \in \text{Diag}^+(n)\}$.*

1. *The smooth map f has a global minimizer.*
2. *For all $\Sigma \in \text{Sym}^+(n)$, there exists $\Delta \in \text{Diag}^+(n)$ such that $\log(\Delta\Sigma\Delta) \in \text{Hol}(n)$.*
3. *The following conjectures are equivalent for all $C \in \text{Cor}^+(n)$.*
 - (i) *There exists a unique $\Delta \in \text{Diag}^+(n)$ such that $\log(\Delta C \Delta) \in \text{Hol}(n)$ (Conjecture 3.1).*
 - (ii) *There exists a unique $X \in \text{Hol}(n)$ such that $\text{Exp}_{I_n}^{\text{QA}}(X) = C$.*
 - (iii) *There exists a unique local minimizer of the smooth map f , which is actually the global minimizer ensured by statement 1.*
4. *The previous conjectures imply the uniqueness of the quotient-affine logarithm at I_n .*

Proof. 1. The smooth map f has a global minimizer because it is coercive [32, cf. HAL version].

2. Hence, there exists $\Delta \in \text{Diag}^+(n)$ such that $f(\Delta) = \min f$. In other words, $\Delta C \Delta$ is “in optimal position” to I_n [16, Definition 2.3]. Thus [16, Theorem 2.4], the geodesic from I_n to $\Delta C \Delta$ is horizontal, i.e. $\text{Log}_{I_n}^{\text{AI}}(\Delta C \Delta) \in \mathcal{H}_{I_n}^{\text{QA}}$, i.e. $\log(\Delta C \Delta) \in \text{Hol}(n)$. For $\Sigma = \text{Diag}(\Sigma)^{1/2} C \text{Diag}(\Sigma)^{1/2}$, it suffices to take $\Delta \text{Diag}(\Sigma)^{-1/2} \in \text{Diag}^+(n)$.

3. Since $\text{Exp}_{I_n}^{\text{QA}} = \text{Cor} \circ \exp$, we clearly have (i) \iff (ii) because $X = \log(\Delta C \Delta)$ and $\Delta = \text{Diag}(\exp(X))^{1/2}$. To prove (i) \iff (iii), let us compute the differential and the Hessian of f . Let $\Delta \in \text{Diag}^+(n)$ and $D, D' \in \text{Diag}(n)$. We denote $E = D\Delta^{-1} \in \text{Diag}(n)$ and $A = \Delta C \Delta = PBP^\top \in \text{Sym}^+(n)$ with $P \in \text{O}(n)$ and $B \in \text{Diag}^+(n)$.

$$\begin{aligned}
d_\Delta f(D) &= 2 \text{tr}(\log(\Delta C \Delta) d_{\Delta C \Delta} \log((DC\Delta + \Delta CD))) \\
&= 2 \text{tr}(\log(A) d_A \log(EA + AE^\top)) \\
&= 2 \text{tr}(\log(B) d_B \log(P^\top EPB + BP^\top E^\top P)) \\
&= 2 \text{tr}(\log(B) B^{-1} (P^\top EPB + BP^\top E^\top P)) \\
&= 4 \text{tr}(\log(A) E) = 4 \text{tr}(\log(\Delta C \Delta) D \Delta^{-1}),
\end{aligned}$$

$$\begin{aligned}
d_\Delta f &= 0 \iff \text{Diag}(\log(\Delta C \Delta)) = 0 \iff \log(\Delta C \Delta) \in \text{Hol}(n), \\
H_\Delta f(D, D') &= 4 \text{tr}(d_A \log(EA + AE^\top) D' \Delta^{-1} - \log(A) D' \Delta^{-1} D \Delta^{-1}) \\
&= 4 \text{tr}(2\Delta^{-2} D D' - \text{Diag}(\log(\Delta C \Delta)) \Delta^{-2} D D'), \\
H_\Delta f(D, D) &= 4 \text{tr}((2I_n - \text{Diag}(\log(\Delta C \Delta))) \Delta^{-2} D^2).
\end{aligned}$$

Hence, if $\Delta \in \text{Diag}^+(n)$ is such that $\log(\Delta C \Delta) \in \text{Hol}(n)$, then the Hessian of f at Δ is positive definite so f has a local minimum at Δ . Thus, if proposition (iii) is true, then Δ has to be the global minimizer so it is unique. Conversely, if f has a local minimum at $\Delta \in \text{Diag}^+(n)$, then $d_\Delta f = 0$ so $\log(\Delta C \Delta) \in \text{Hol}(n)$. Thus, if proposition (i) is true, then Δ is unique. Therefore, assertions (i) and (iii) are equivalent.

4. A quotient-affine logarithm of $C \in \text{Cor}^+(n)$ at I_n is a tangent vector $X \in \text{Hol}(n)$ of *minimal length* such that $\text{Exp}_{I_n}^{\text{QA}}(X) = \text{Cor}(\exp(X)) = C$. Otherwise said, it is a tangent vector $X = \log(\Delta C \Delta)$ where Δ minimizes $f(\Delta) = \text{tr}(X^2) = \|X\|^2$. Thus the uniqueness in the conjectures of statement 3 imply the uniqueness of the quotient-affine logarithm at I_n . \square

Otherwise said, Conjecture 3.1 is stronger than the conjecture stating the uniqueness of the quotient-affine logarithm at I_n . On the one hand, this could provide a new path to prove the latter. However, Conjecture 3.1 seems difficult to prove because the manifold $\exp(\text{Hol}(n))$ is hard to describe in terms of properties on the coefficients of the matrices. Thus it is difficult to determine whether its intersection with the fiber $\text{Diag}^+(n) \star \Sigma = \text{Cor}^{-1}(\text{Cor}(\Sigma))$ is reduced to one point or not. On the other hand, this could also help to show that Conjecture 3.1 is false. Indeed, the quotient-affine metric has both positive and negative curvature so the quotient-affine logarithm might not be unique. Hence, this seems to be a difficult problem.

3.2. Why $\mathcal{V}^* = \text{Row}_0(n)$ seems to be a better choice. Nevertheless, another interesting decomposition of symmetric matrices where each subspace is stable by permutations is given by $\text{Sym}(n) = \text{Row}_0(n) \oplus \text{Diag}(n)$, where $\text{Row}_0(n) = \{S \in \text{Sym}(n) | S\mathbf{1} = 0\}$ is the vector space of symmetric matrices with null row sum. That is why we propose to examine $\mathcal{V}^* = \text{Row}_0(n)$. In the following theorem, we show that $\exp(\text{Row}_0(n))$ has a nice form.

THEOREM 3.4 ($\exp : \text{Row}_0(n) \rightarrow \text{Row}_\mathbb{1}^+(n)$ is a smooth diffeomorphism). *The symmetric matrix logarithm is a smooth diffeomorphism from $\text{Row}_\mathbb{1}^+(n) = \{\Sigma \in \text{Sym}^+(n) | \Sigma\mathbf{1} = \mathbf{1}\}$ onto $\text{Row}_0(n) = \{S \in \text{Sym}(n) | S\mathbf{1} = 0\}$.*

Proof. It is clear that $\exp(\text{Row}_0(n)) \subset \text{Row}_\mathbb{1}^+(n)$ since if $S \in \text{Row}_0(n)$, then $\exp(S)\mathbf{1} = \sum_{k=0}^{\infty} \frac{1}{k!} S^k \mathbf{1} = \mathbf{1}$. Conversely, let $\Sigma \in \text{Row}_\mathbb{1}^+(n)$. Then the Lagrange polynomial $P(X) = \sum_{\lambda \in \text{eig}(\Sigma)} \log(\lambda) \prod_{\mu \in \text{eig}(\Sigma), \mu \neq \lambda} \frac{X+1-\mu}{\lambda-\mu}$ satisfies $P(\Sigma - I_n) = \log \Sigma$. Since $1 \in \text{eig}(\Sigma)$, $P(0) = \log(1) + \sum_{\lambda \neq 1} \log(\lambda) \frac{1-\lambda}{\lambda-1} \prod_{\mu \neq \lambda, 1} \frac{1-\mu}{\lambda-\mu} = 0$. Hence $\log(\Sigma)\mathbf{1} = P(\Sigma - I_n)\mathbf{1} = P(0)\mathbf{1} = 0$. So $\log(\text{Row}_\mathbb{1}^+(n)) \subset \text{Row}_0(n)$. Finally, $\text{Row}_\mathbb{1}^+(n) = \exp(\text{Row}_0(n))$ so $\log : \text{Row}_\mathbb{1}^+(n) \rightarrow \text{Row}_0(n)$ is a smooth diffeomorphism. \square

Hence, the question becomes: for all $\Sigma \in \text{Sym}^+(n)$, does there exist a unique $\Delta \in \text{Diag}^+(n)$ such that $\Delta \Sigma \Delta \in \text{Row}_\mathbb{1}^+(n)$? The answer is yes [26, 18], let us explain why.

3.3. The conjecture is true with $\mathcal{V}^* = \text{Row}_0(n)$. We recall that we denote $\star : \text{Diag}^+(n) \times \text{Sym}^+(n) \rightarrow \text{Sym}^+(n)$ the congruence action of positive diagonal matrices on SPD matrices.

THEOREM 3.5 (Definition of \mathcal{D}^*). *For all $\Sigma \in \text{Sym}^+(n)$, there exists a unique $\Delta \in \text{Diag}^+(n)$ such that $\Delta\Sigma\Delta \in \text{Row}_0^+(n)$ [26, 18] or equivalently $\log(\Delta\Sigma\Delta) \in \text{Row}_0(n)$ by Theorem 3.4. This allows to define:*

- the surjective map $\mathcal{D}^* : \Sigma \in \text{Sym}^+(n) \mapsto \Delta \in \text{Diag}^+(n)$,
- the surjective map $\pi^* : \Sigma \in \text{Sym}^+(n) \mapsto \log(\mathcal{D}^*(\Sigma) \star \Sigma) \in \text{Row}_0(n)$ which is invariant under the congruence group action of $\text{Diag}^+(n)$ on $\text{Sym}^+(n)$,
- the bijective map $\text{Log}^* = \pi^*_{|\text{Cor}^+(n)} : \text{Cor}^+(n) \rightarrow \text{Row}_0(n)$ that we call the log-scaling (note that $\pi^* = \text{Log}^* \circ \text{Cor}$),
- the smooth bijective inverse map $\text{Exp}^* = (\text{Log}^*)^{-1} = \text{Cor} \circ \text{exp} : \text{Row}_0(n) \rightarrow \text{Cor}^+(n)$.

Proof. The existence and uniqueness are due to [26], the uniqueness has been proved differently later in [18].

- The map \mathcal{D}^* is surjective because $\mathcal{D}^*(\Delta^{-2}) = \Delta$ for all $\Delta \in \text{Diag}^+(n)$.
- The map π^* is surjective because if $S \in \text{Row}_0(n)$, then $\pi^*(\text{exp}(S)) = S$.
- The map Log^* is surjective because $\text{Log}^*(\text{Cor}(\text{exp}(S))) = S$ and injective because if $C, C' \in \text{Cor}^+(n)$ are such that $\text{Log}^*(C) = \text{Log}^*(C')$, then $\mathcal{D}^*(C) \star C = \mathcal{D}^*(C') \star C'$ so $C = \text{Cor}(\mathcal{D}^*(C) \star C) = \text{Cor}(\mathcal{D}^*(C') \star C') = C'$.
- We just showed that $(\text{Log}^*)^{-1} = \text{Cor} \circ \text{exp}$ so $\text{Exp}^* = (\text{Log}^*)^{-1}$ is bijective and smooth. \square

3.4. Properties of the log-scaling bijection. Let us give properties of the log-scaling bijection Log^* that are analogous to the properties of off-log bijection Log introduced in Section 2.

THEOREM 3.6 (Properties of the log-scaling bijection).

1. (Equivariance) Log^* and Exp^* are equivariant under permutations.
2. (Equicorrelation) $\text{Log}^*(C(\rho)) = \frac{1}{n} \ln \left(\frac{1+(n-1)\rho}{1-\rho} \right) (\mathbf{1}\mathbf{1}^\top - nI_n)$ for all $\rho \in (-\frac{1}{n-1}, 1)$. In dimension $n = 2$, $\text{Log}^*(C(\rho)) = \begin{pmatrix} -F(\rho) & F(\rho) \\ F(\rho) & -F(\rho) \end{pmatrix}$ where $F(\rho) = \frac{1}{2} \log \left(\frac{1+\rho}{1-\rho} \right) \in \mathbb{R}$ is the Fisher transformation of the correlation coefficient $\rho \in (-1, 1)$.
3. (Block equicorrelation matrix) If C is a block equicorrelation matrix of signature $I = \{i_1, \dots, i_p\}$, then $\text{Log}^*(C)$ is a block symmetric matrix with null row sum of signature I with diagonal blocks of the form $(\alpha_j - \beta_j)I_{i_j} + \beta_j \mathbf{1}_{i_j} \mathbf{1}_{i_j}^\top$ and off-diagonal blocks of the form $\beta_{jk} \mathbf{1}_{i_j} \mathbf{1}_{i_k}^\top$.
4. (Generalization) For all $x \in (\mathbb{R}^+)^n$, for all $\Sigma \in \text{Sym}^+(n)$, there exists a unique $\Delta \in \text{Diag}^+(n)$ such that $\log(\Delta\Sigma\Delta)x = 0$.

Proof. 1. This is clear.

2. Let $C = C(\rho)$. The result is clear for $\rho = 0$ so we assume that $\rho \neq 0$. One easily checks that $\Delta = \sqrt{a}I_n$ with $a = \frac{1}{1+(n-1)\rho}$ satisfies $\Sigma := \Delta C \Delta = aC = (a-b)I_n + b\mathbf{1}\mathbf{1}^\top \in \text{Row}_0^+(n)$ with $b = a\rho = \frac{\rho}{1+(n-1)\rho}$. Indeed, $a + (n-1)b = 1$. Since $\log(\Sigma)$ is a polynomial in Σ , there exists $(\alpha, \beta) \in \mathbb{R}^2$ such that $\log(\Sigma) = (\alpha - \beta)I_n + \beta\mathbf{1}\mathbf{1}^\top$ and $\alpha + (n-1)\beta = 0$. Moreover, $\text{eig}(\Sigma) = \{1; a-b\}$ with $a-b \neq 1$ and $\text{eig}(\log(\Sigma)) = \{0; \alpha - \beta\}$ so $\alpha - \beta = \ln(a-b) = -\ln \left(\frac{1+(n-1)\rho}{1-\rho} \right)$.

Therefore, $n\beta = -(\alpha - \beta) = \ln \left(\frac{1+(n-1)\rho}{1-\rho} \right)$ and:

$$\text{Log}^*(C) = (\alpha - \beta)I_n + \beta\mathbf{1}\mathbf{1}^\top = \beta(\mathbf{1}\mathbf{1}^\top - nI_n) = \frac{1}{n} \ln \left(\frac{1+(n-1)\rho}{1-\rho} \right) (\mathbf{1}\mathbf{1}^\top - nI_n).$$

3. If C is a block equicorrelation matrix of signature $I = \{i_1, \dots, i_p\}$, it is clear

that the matrix Δ is a block diagonal matrix of signature I with scalar blocks because the sums of all the rows belonging to the same interval $[i_j + 1; i_{j+1}]$ are equal. The matrix product preserves the signature and the form of the blocks so the logarithm as well.

4. Similarly to Theorem 3.4, one can prove that given $x \in (\mathbb{R}^+)^n$, the map $\exp : \{S \in \text{Sym}(n) | Sx = 0\} \rightarrow \{\Sigma \in \text{Sym}^+(n) | \Sigma x = x\}$ is a diffeomorphism. We denote $X = \text{diag}(x) \in \text{Diag}^+(n)$. Therefore, for all $\Delta \in \text{Diag}^+(n)$, $\log(\Delta \Sigma \Delta)x = 0$ if and only if $\Delta \Sigma \Delta x = x$ if and only if $X \Delta \Sigma \Delta X \mathbf{1} = X^2 \mathbf{1}$ if and only if $X \Delta \in \text{Diag}^+(n)$ scales Σ onto an SPD matrix with row sums prescribed by $X^2 \mathbf{1}$. Thus the existence and uniqueness are ensured by [26]. \square

The log-scaling bijection has an additional property: it is compatible with the cor-inversion. It is a corollary of Theorem 1.2, the key reason being the use of the congruence action of positive diagonal matrices instead of the additive action of diagonal matrices.

THEOREM 3.7 (Compatibility between inversion and log-scaling bijection). *For all $C \in \text{Cor}^+(n)$, $\text{Log}^*(\mathcal{I}(C)) = -\text{Log}^*(C)$. Otherwise said, the following diagram commutes.*

$$(3.1) \quad \begin{array}{ccc} \text{Cor}^+(n) & \xrightarrow{\mathcal{I}} & \text{Cor}^+(n) \\ \downarrow \text{Log}^* & & \downarrow \text{Log}^* \\ \text{Row}_0(n) & \xrightarrow{-\text{Id}} & \text{Row}_0(n) \end{array}$$

In this section, we proved the existence in the conjecture with $\mathcal{V}^* = \text{Hol}(n)$ and we proved the conjecture for $\mathcal{V}^* = \text{Row}_0(n)$. This provides a bijective map $\text{Log}^* : \text{Cor}^+(n) \rightarrow \text{Row}_0(n)$ called the log-scaling bijection.

4. Permutation-invariant log-Euclidean metrics via the log-scaling bijection. In this section, we use the log-scaling bijection to define log-Euclidean metrics on $\text{Cor}^+(n)$. More precisely, in Section 4.1, we prove that the log-scaling bijection is a diffeomorphism. In Section 4.2, we characterize all permutation-invariant inner products on $\text{Row}_0(n)$. In Section 4.3, we define permutation-invariant log-Euclidean metrics by pullback and we give their geometric properties.

4.1. The cor-exp bijection is a diffeomorphism.

THEOREM 4.1 ($\text{Exp}^* = \text{Cor} \circ \exp$ is a diffeomorphism). *The log-scaling bijection $\text{Log}^* : \text{Cor}^+(n) \rightarrow \text{Row}_0(n)$ is a smooth diffeomorphism. We give the differentials of Log^* and Exp^* in function of the differentials of the symmetric matrix logarithm and exponential maps \log and \exp . For all $C \in \text{Cor}^+(n)$, $S, Y \in \text{Row}_0(n)$ and $X \in \text{Hol}(n)$ such that $\Sigma = \mathcal{D}^*(C) \star C = \exp(S)$:*

$$(4.1) \quad d_S \text{Exp}^*(Y) = \Delta^{-1} \left[d_S \exp(Y) - \frac{1}{2} (\Delta^{-2} \text{Diag}(d_S \exp(Y)) \Sigma + \Sigma \text{Diag}(d_S \exp(Y)) \Delta^{-2}) \right] \Delta^{-1},$$

$$(4.2) \quad d_C \text{Log}^*(X) = d_\Sigma \log \left(\Delta X \Delta + \frac{1}{2} (X^0 \Sigma + \Sigma X^0) \right),$$

where $\Delta = \text{Diag}(\Sigma)^{1/2}$ and $X^0 = -2 \text{diag}((I_n + \Sigma)^{-1} \Delta X \Delta \mathbf{1})$.

Proof. It suffices to show that \mathcal{D}^* is smooth. We apply the implicit function theorem to the smooth function $\Phi^* : (\Sigma, \Delta) \in \text{Sym}^+(n) \times \text{Diag}^+(n) \mapsto \Delta \Sigma \Delta \mathbf{1} - \mathbf{1} \in$

$(\mathbb{R}^+)^n$ which satisfies $\Delta = \mathcal{D}^*(\Sigma)$ if and only if $\Phi^*(\Sigma, \Delta) = 0$. Let us prove that for all $(\Sigma, \Delta) \in (\Phi^*)^{-1}(0)$, the differential of the partial function $\Phi_\Sigma^* : \Delta \in \text{Diag}^+(n) \mapsto \Phi^*(\Sigma, \Delta) \in (\mathbb{R}^+)^n$ is invertible. In the direction $D \in T_\Delta \text{Diag}^+(n) = \text{Diag}(n)$:

$$\begin{aligned} d_\Delta \Phi_\Sigma^*(D) &= D\Sigma\Delta\mathbf{1} + \Delta\Sigma D\mathbf{1} \\ &= D\Delta^{-1}\mathbf{1} + \Delta\Sigma D\mathbf{1} \\ &= \Delta(\Delta^{-2} + \Sigma)D\mathbf{1}. \end{aligned}$$

Since $\Delta(\Delta^{-2} + \Sigma) \in \text{GL}(n)$, the differential is invertible so \mathcal{D}^* is smooth.

Since $\text{Exp}^* = \text{Cor} \circ \exp$, we have $d_S \text{Exp}^*(Y) = d_\Sigma \text{Cor}(d_S \exp(Y))$ with $\Sigma = \exp(S)$. Using $d_\Sigma \text{Cor}(Z) = \Delta^{-1} [Z - \frac{1}{2}(\Delta^{-2} \text{Diag}(Z)\Sigma + \Sigma \text{Diag}(Z)\Delta^{-2})] \Delta^{-1}$ with $\Delta = \text{Diag}(\Sigma)^{1/2}$ and $Z = d_S \exp(Y) \in T_\Sigma \text{Row}_\mathbf{1}^+(n) = \text{Row}_0(n)$, we get the expected result. Now we want to invert the relation $X = d_S \text{Exp}^*(Y)$ to get $Y = d_\Sigma \text{Log}^*(X)$. We use the intermediate matrices $\Sigma \in \text{Row}_\mathbf{1}^+(n)$ and $Z \in \text{Row}_0(n)$ and the relations $X = d_\Sigma \text{Cor}(Z)$ and $Y = d_\Sigma \log(Z)$.

$$\begin{aligned} \Delta X \Delta &= Z - \frac{1}{2}(\Delta^{-2} \text{Diag}(Z)\Sigma + \Sigma \text{Diag}(Z)\Delta^{-2}), \\ \Delta X \Delta \mathbf{1} &= -\frac{1}{2}(\Delta^{-2} \text{Diag}(Z)\Sigma \mathbf{1} + \Sigma \text{Diag}(Z)\Delta^{-2} \mathbf{1}) \\ &= -\frac{1}{2}(I_n + \Sigma)\Delta^{-2} \text{Diag}(Z)\mathbf{1}, \\ \Delta^{-2} \text{Diag}(Z)\mathbf{1} &= -2(I_n + \Sigma)^{-1} \Delta X \Delta \mathbf{1}, \\ \Delta^{-2} \text{Diag}(Z) &= X^0, \\ Z &= \Delta X \Delta + \frac{1}{2}(X^0 \Sigma + \Sigma X^0), \end{aligned}$$

which allows to conclude with $d_\Sigma \text{Log}^*(X) = Y = d_\Sigma \log(Z)$. \square

4.2. Permutation-invariant inner products on $\text{Row}_0(n)$.

THEOREM 4.2 (Permutation-invariant inner products on $\text{Row}_0(n)$). *For $n \geq 4$, permutation-invariant inner products on $\text{Row}_0(n)$ are the symmetric bilinear forms associated to the following positive definite quadratic forms q^* defined for $Y \in \text{Row}_0(n)$:*

$$(4.3) \quad q^*(Y) = \alpha \text{tr}(Y^2) + \delta \text{tr}(\text{Diag}(Y)^2) + \zeta \text{tr}(Y)^2,$$

with $\alpha > 0$, $n\alpha + (n-2)\delta > 0$ and $n\alpha + (n-1)(\delta + n\zeta) > 0$. For $n = 3$, the permutation-invariant inner products have the same form with $\alpha = 0$. For $n = 2$, they have the same form with $\alpha = \delta = 0$.

Proof. We rely on the characterization of permutation-invariant inner products on $\text{Hol}(n)$ and on the equivariant isomorphism $\Theta : \begin{cases} \text{Hol}(n) & \longrightarrow & \text{Row}_0(n) \\ X & \longmapsto & Y = AXA^\top \end{cases}$ with $A = I_n - \frac{1}{n}\mathbf{1}\mathbf{1}^\top$ found in [21], whose inverse isomorphism is given by $X = \Theta^{-1}(Y) = Y - (\mu\mathbf{1}\mathbf{1}^\top + \mathbf{1}\mu^\top)$ with $\mu = \frac{1}{2}\text{Diag}(Y)\mathbf{1}$. Indeed, let $q^* : \text{Row}_0(n) \longrightarrow \mathbb{R}$ be a permutation-invariant quadratic form. Then q^* is positive definite if and only if $q^* \circ \Theta : \text{Hol}(n) \longrightarrow \mathbb{R}$ is a permutation-invariant quadratic form on $\text{Hol}(n)$. Hence, by Theorem 2.5, q^* is of the form $q^*(Y) = \alpha \text{tr}(\Theta^{-1}(Y)^2) + \beta \text{Sum}(\Theta^{-1}(Y)^2) + \gamma \text{Sum}(\Theta^{-1}(Y))^2$ with $\min(\alpha, 2\alpha + (n-2)\beta, \alpha + (n-1)(\beta + n\gamma)) > 0$. We compute

X^2 with $X = \Theta^{-1}(Y)$, and the three terms:

$$\begin{aligned} X^2 &= Y^2 + \frac{1}{2}\text{tr}(Y)(\mu\mathbf{1}^\top + \mathbf{1}\mu^\top) + n\mu\mu^\top + \|\mu\|^2\mathbf{1}\mathbf{1}^\top - Y\mu\mathbf{1}^\top - \mathbf{1}\mu^\top Y, \\ \text{tr}(X^2) &= \text{tr}(Y^2) + \frac{1}{2}\text{tr}(Y)^2 + \frac{n}{2}\text{tr}(\text{Diag}(Y)^2), \\ \text{Sum}(X^2) &= \frac{n}{2}\text{tr}(Y)^2 + \frac{n}{4}\text{tr}(Y)^2 + \frac{n^2}{4}\text{tr}(\text{Diag}(Y)^2), \\ \text{Sum}(X) &= -n\text{tr}(Y), \\ \text{Sum}(X)^2 &= n^2\text{tr}(Y)^2. \end{aligned}$$

Hence $g(Y, Y) = \alpha\text{tr}(Y^2) + \delta\text{tr}(\text{Diag}(Y)^2) + \zeta\text{tr}(Y)^2$ with $\delta = \frac{n}{2}(\alpha + \frac{n}{2}\beta)$ and $\zeta = \frac{\alpha}{2} + \frac{3n}{4}\beta + n^2\gamma$. The inverse relations between coefficients are $\beta = \frac{4}{n^2}\delta - \frac{2}{n}\alpha$ and $\gamma = \frac{1}{n^2}(\zeta - \frac{3}{n}\delta + \alpha)$. Thus $2\alpha + (n-2)\beta = \frac{4}{n^2}(n\alpha + (n-2)\delta)$ and $\alpha + (n-1)(\beta + n\gamma) = \frac{1}{n^2}(n\alpha + (n-1)(\delta + n\zeta))$, which gives the expected positivity condition. \square

4.3. Pullback metrics via the log-scaling diffeomorphism.

DEFINITION 4.3 (Log-scaled metrics). *A log-scaled metric on $\text{Cor}^+(n)$ is the pullback metric of a permutation-invariant inner product characterized by a quadratic form q^* as in Theorem 4.2. For all $C \in \text{Cor}^+(n)$ and $X \in T_C\text{Cor}^+(n) = \text{Hol}(n)$, it writes $g_C^*(X, X) = q^*(d_C\text{Log}^*(X))$ where $d_C\text{Log}^*(X) = d_C\log(X + \frac{1}{2}(X^0C + CX^0))$ with $X^0 = -2\text{diag}((I_n + C)^{-1}X\mathbf{1})$.*

THEOREM 4.4 (Riemannian operations of log-scaled metrics). *We consider a log-scaled metric characterized by the quadratic form q^* . Let $C, C', C_1, \dots, C_n \in \text{Cor}^+(n)$, $X \in \text{Row}_0(n)$. The Riemannian operations of this metric are summarized in Table 4.1. Moreover, the metric is permutation-invariant and inverse-consistent, i.e. it is invariant under the pullback by the cor-inversion $\mathcal{I} : \text{Cor}^+(n) \rightarrow \text{Cor}^+(n)$.*

Exponential map	$\text{Exp}_C^*(X) = \text{Exp}^*(\text{Log}^*(C) + d_C\text{Log}^*(X))$
Logarithm map	$\text{Log}_C^*(C') = d_{\text{Log}^*(C)}\text{Exp}^*(\text{Log}^*(C') - \text{Log}^*(C))$
Geodesic	$\gamma^*(t) = \text{Exp}^*((1-t)\text{Log}^*(C) + t\text{Log}^*(C'))$
Squared distance	$d^*(C, C')^2 = q^*(\text{Log}^*(C') - \text{Log}^*(C))$
Fréchet mean	$\bar{C}^* = \text{Exp}^*(\frac{1}{k}\sum_{i=1}^k \text{Log}^*(C_i))$
Curvature	$R^* = 0$
Parallel transport	$\Pi_{C \rightarrow C'}^* X = (d_{C'}\text{Log}^*)^{-1}(d_C\text{Log}^*(X))$

Table 4.1: Riemannian operations of log-scaled metrics

Beware not to confuse the Riemannian maps $\text{Exp}_C^* : T_C\text{Cor}^+(n) = \text{Hol}(n) \rightarrow \text{Cor}^+(n)$ and $\text{Log}_C^* : \text{Cor}^+(n) \rightarrow T_C\text{Cor}^+(n) = \text{Hol}(n)$ with the diffeomorphisms $\text{Exp}^* : \text{Row}_0(n) \rightarrow \text{Cor}^+(n)$ and $\text{Log}^* : \text{Cor}^+(n) \rightarrow \text{Row}_0(n)$.

As for the off-log metrics, all the Riemannian operations can be computed in closed form modulo the computation of Log^* , i.e. the computation of \mathcal{D}^* , i.e. the computation of the scaling of an SPD matrix. Several algorithms exist to compute the scaling of a matrix to prescribed row sums since it is an important research topic in different scientific communities from linear algebra to probability theory. In general, scaling a matrix $\Sigma \in \text{Mat}(n)$ to $\mu \in \mathbb{R}^n$ means finding positive diagonal matrices $D_1, D_2 \in \text{Diag}(n)$ such that $D_1\Sigma D_2\mathbf{1} = \mu$. The reader may refer to [18, 17, 2] for

surveys on theoretical results and algorithms. The best method to compute the scaling of an SPD matrix $\Sigma \in \text{Sym}^+(n)$ seems to be the one in [2] where the total complexity is $O(m + n^{4/3})$ where m is the number of non-zero entries of Σ .

Since one motivation behind the off-log bijection in [3] was the closed form expression in dimension 2 (cf. Theorem 2.2, item 2), it seems important to note the following result.

THEOREM 4.5 (Coincidence of the metrics in dimension 2). *In dimension 2, up to a positive scaling factor, the quotient-affine metric, the off-log metric and the log-scaled metric coincide. We recall that the Fisher transformation is the increasing map $F : \rho \in (-1, 1) \mapsto \frac{1}{2} \ln\left(\frac{1+\rho}{1-\rho}\right) \in \mathbb{R}^+$. Let $C = C(\rho)$ and $X = \begin{pmatrix} 0 & x \\ x & 0 \end{pmatrix}$ with $\rho \in (-1, 1)$ and $x \in \mathbb{R}$. Then:*

1. (Metric) $g_C(X, X) = \frac{x^2}{(1-\rho^2)^2}$ (up to a scaling factor $\alpha > 0$),
2. (Geodesic) $\gamma(t) = C(\rho(t))$ where $\rho(t) = \frac{\rho_1 \cosh(\lambda t) + \sinh(\lambda t)}{\rho_1 \sinh(\lambda t) + \cosh(\lambda t)}$ with $\lambda = F(\rho_2) - F(\rho_1)$ is monotonic (increasing if $\rho_1 < \rho_2$, decreasing if $\rho_1 > \rho_2$, constant if $\rho_1 = \rho_2$),
3. (Distance) $d(C_1, C_2) = |\lambda| = |F(\rho_2) - F(\rho_1)|$ (up to a scaling factor $\sqrt{\alpha}$).

Proof. It suffices to use the 2nd statement of Theorems 2.2 and 3.6 and the formulae of distances of the off-log metric and the log-scaled metric. For the former, $d(C_1, C_2) = q(F(\rho_2) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} - F(\rho_1) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}) \propto \lambda^2$ and similarly for the ladder. Up to a multiplicative constant, these distances are equal to the quotient-affine distance in dimension 2 [32]. Therefore, the Riemannian metrics coincide up to a constant and the geodesics coincide. The formulae can be found in [34]. \square

We showed in this section that the log-scaling bijection is a diffeomorphism. Therefore it provides a family of permutation-invariant and inverse-consistent log-Euclidean metrics by pulling back permutation-invariant inner products on $\text{Row}_0(n)$. Thus, the Riemannian operations are trivial. As shown in Table 4.1, they are known in closed form modulo the computation of $\text{Log}^* : \text{Cor}^+(n) \rightarrow \text{Row}_0(n)$ or equivalently the computation of $\mathcal{D}^* : \text{Sym}^+(n) \rightarrow \text{Diag}^+(n)$. This amounts to compute the scaling of an SPD matrix so there exist very efficient methods to do it, for example the one in [2].

5. Conclusion. We introduced two families of permutation-invariant log-Euclidean metrics on full-rank correlation matrices. The first family was built via the off-log diffeomorphism whose algebraic properties were introduced in [3]. The second family was built via the log-scaling diffeomorphism thanks to the result on the scaling of symmetric matrices [26]. In addition, the log-scaled metrics are inverse-consistent contrarily to the off-log metrics. These metrics allow to compute with correlation matrices very efficiently since they are flat and the main Riemannian operations are known in closed form modulo the computation of the maps \mathcal{D} and \mathcal{D}^* , which can be done very efficiently.

These permutation-invariant log-Euclidean metrics fill a gap in the study of Riemannian metrics on the open elliptope since they gather the invariance under permutations satisfied by several existing structures [19, 27, 9, 32] and the log-Euclidean geometry of some others [34]. More generally, we tried to introduce a comprehensive formalism on full-rank correlation matrices in terms of stability under the action of permutations and under the cor-inverse involution. This systematic approach allowed to satisfy intrinsically the requirement of inverse-consistency in the construction of

log-Euclidean metrics. We hope that this presentation will help to manipulate correlation matrices as easily as we work with SPD matrices. This approach could also help define appropriate geometries of block equi-correlation matrices of a given signature.

Acknowledgements. This work has been supported by the French government, through the UCAJEDI Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-15-IDEX-01. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant G-Statistics agreement No 786854). The author warmly thank Xavier Pennec for his careful proofreading of this manuscript.

REFERENCES

- [1] D. ALEKSEEVSKY, A. KRIEGL, M. LOSIK, AND P. W. MICHOR, *The Riemannian geometry of orbit spaces. The metric, geodesics, and integrable systems*, Publicationes Mathematicae, 62 (2001), pp. 1–30.
- [2] Z. ALLEN-ZHU, Y. LI, R. OLIVEIRA, AND A. WIGDERSON, *Much faster algorithms for matrix scaling*, in 2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS), Los Alamitos, CA, USA, oct 2017, IEEE Computer Society, pp. 890–901, <https://doi.org/10.1109/FOCS.2017.87>, <https://doi.ieeecomputersociety.org/10.1109/FOCS.2017.87>.
- [3] I. ARCHAKOV AND P. R. HANSEN, *A New Parametrization of Correlation Matrices*, *Econometrica*, 89 (2021), pp. 1699–1715, <https://doi.org/10.3982/ECTA16910>, <https://onlinelibrary.wiley.com/doi/abs/10.3982/ECTA16910>.
- [4] V. ARSIGNY, P. FILLARD, X. PENNEC, AND N. AYACHE, *Log-Euclidean metrics for fast and simple calculus on diffusion tensors*, *Magnetic Resonance in Medicine*, 56 (2006), pp. 411–421, <https://doi.org/10.1002/mrm.20965>, <https://hal.inria.fr/inria-00502678>.
- [5] A. BARACHANT, S. BONNET, M. CONGEDO, AND C. JUTTEN, *Classification of covariance matrices using a Riemannian-based kernel for BCI applications*, *Neurocomputing*, 112 (2013), pp. 172–178, <https://doi.org/10.1016/j.neucom.2012.12.039>, <https://hal.archives-ouvertes.fr/hal-00820475>.
- [6] F. BARBARESCO, *Information Geometry of Covariance Matrix: Cartan-Siegel Homogeneous Bounded Domains, Mostow/Berger Fibration and Fréchet Median*, in *Matrix Information Geometry*, F. Nielsen and R. Bhatia, eds., Springer, Berlin, Heidelberg, 2013, pp. 199–255, https://doi.org/10.1007/978-3-642-30232-9_9, https://doi.org/10.1007/978-3-642-30232-9_9.
- [7] R. BHATIA, *Matrix Analysis*, vol. 169 of Graduate Texts in Mathematics, Springer New York, New York, NY, 1997, <https://doi.org/10.1007/978-1-4612-0653-8>, <http://link.springer.com/10.1007/978-1-4612-0653-8>.
- [8] J. P. BURG, *Maximum Entropy Spectral Analysis*, PhD thesis, Stanford University, 1975.
- [9] P. DAVID AND W. GU, *A Riemannian structure for correlation matrices*, *Operators and Matrices*, 13 (2019), pp. 607–627.
- [10] A. DE LA FUENTE, N. BING, I. HOESCHELE, AND P. MENDES, *Discovery of meaningful associations in genomic data using partial correlation coefficients*, *Bioinformatics*, 20 (2004), pp. 3565–3574, <https://doi.org/10.1093/bioinformatics/bth445>, <https://doi.org/10.1093/bioinformatics/bth445>.
- [11] S. EPSKAMP AND E. I. FRIED, *A tutorial on regularized partial correlation networks*, *Psychological Methods*, 23 (2018), pp. 617–634, <https://doi.org/10.1037/met0000167>. Place: US Publisher: American Psychological Association.
- [12] P. FILLARD, X. PENNEC, V. ARSIGNY, AND N. AYACHE, *Clinical DT-MRI estimation, smoothing, and fiber tracking with log-Euclidean metrics.*, *IEEE Transactions on Medical Imaging*, 26 (2007), pp. 1472–82, <https://doi.org/10.1109/TMI.2007.899173>, <https://hal.inria.fr/inria-00502645>.
- [13] M. K. GARBA, T. M. W. NYE, J. LUEG, AND S. F. HUCKEMANN, *Information geometry for phylogenetic trees*, *Journal of Mathematical Biology*, 82 (2021), pp. 1–39, <https://doi.org/10.1007/s00285-021-01553-x>, <https://doi.org/10.1007/s00285-021-01553-x>.
- [14] I. GRUBIŠIĆ AND R. PIETERSZ, *Efficient rank reduction of correlation matrices*, *Linear Algebra and its Applications*, 422 (2007), pp. 629–653, <https://doi.org/https://doi.org/10.1016/j>.

- laa.2006.11.024, <https://www.sciencedirect.com/science/article/pii/S0024379506005222>.
- [15] F. HIAI AND D. PETZ, *Riemannian metrics on positive definite matrices related to means*, Linear Algebra and its Applications, 430 (2009), pp. 3105–3130, <https://doi.org/https://doi.org/10.1016/j.laa.2009.01.025>, <https://www.sciencedirect.com/science/article/pii/S0024379509000767>.
- [16] S. HUCKEMANN, T. HOTZ, AND A. MUNK, *Intrinsic shape analysis: Geodesic PCA for Riemannian manifolds modulo isometric Lie group actions*, Statistica Sinica, 20 (2010), pp. 1–58.
- [17] M. IDEL, *A review of matrix scaling and Sinkhorn's normal form for matrices and positive maps*, 2016, <https://arxiv.org/abs/1609.06349>.
- [18] C. R. JOHNSON AND R. REAMS, *Scaling of symmetric matrices by positive diagonal congruence*, Linear and Multilinear Algebra, 57 (2009), pp. 123–140, <https://doi.org/10.1080/03081080600872327>, <http://www.tandfonline.com/doi/abs/10.1080/03081080600872327>.
- [19] A. N. KERCHEVAL, *On Rebonato and Jäckel's parametrization method for finding nearest correlation matrices*, International Journal of Pure and Applied Mathematics, 45 (2008), pp. 383–390.
- [20] D. KOLLER AND N. FRIEDMAN, *Probabilistic graphical models: principles and techniques*, Adaptive computation and machine learning, MIT Press, Cambridge, MA, 2009.
- [21] H. KURATA AND R. B. BAPAT, *Moore-Penrose inverse of a hollow symmetric matrix and a pre-distance matrix*, Special Matrices, 4 (2016), <https://doi.org/doi:10.1515/spma-2016-0028>, <https://doi.org/10.1515/spma-2016-0028>.
- [22] S. L. LAURITZEN, *Graphical Models*, Oxford Statistical Science Series, Oxford University Press, Oxford, New York, May 1996.
- [23] P. LI, Q. WANG, H. ZENG, AND L. ZHANG, *Local Log-Euclidean Multivariate Gaussian Descriptor and Its Application to Image Classification*, IEEE Transactions on Pattern Analysis and Machine Intelligence, 39 (2017), pp. 803–817, <https://doi.org/10.1109/TPAMI.2016.2560816>.
- [24] Z. LIN, *Riemannian Geometry of Symmetric Positive Definite Matrices via Cholesky Decomposition*, SIAM Journal on Matrix Analysis and Applications, 40 (2019), pp. 1353–1370, <https://doi.org/10.1137/18M1221084>, <https://epubs.siam.org/doi/10.1137/18M1221084>.
- [25] G. MARRELEC, A. KRAINIK, H. DUFFAU, M. PÉLÉGRINI-ISSAC, S. LEHÉRICY, J. DOYON, AND H. BENALI, *Partial correlation for functional brain interactivity investigation in functional MRI*, NeuroImage, 32 (2006), pp. 228–237, <https://doi.org/10.1016/j.neuroimage.2005.12.057>.
- [26] A. W. MARSHALL AND I. OLKIN, *Scaling of matrices to achieve specified row and column sums*, Numerische Mathematik, 12 (1968), pp. 83–90, <https://doi.org/10.1007/BF02170999>, <https://doi.org/10.1007/BF02170999>.
- [27] F. NIELSEN AND K. SUN, *Clustering in Hilbert's Projective Geometry: The Case Studies of the Probability Simplex and the Elliptope of Correlation Matrices*, in Geometric Structures of Information, F. Nielsen, ed., Springer International Publishing, Cham, 2019, pp. 297–331, https://doi.org/10.1007/978-3-030-02520-5_11, https://doi.org/10.1007/978-3-030-02520-5_11.
- [28] J. PENG, P. WANG, N. ZHOU, AND J. ZHU, *Partial Correlation Estimation by Joint Sparse Regression Models*, Journal of the American Statistical Association, 104 (2009), pp. 735–746, <https://doi.org/10.1198/jasa.2009.0126>, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2770199/>.
- [29] X. PENNEC, P. FILLARD, AND N. AYACHE, *A Riemannian Framework for Tensor Computing*, International Journal of Computer Vision, 66 (2006), pp. 41–66, <https://doi.org/10.1007/s11263-005-3222-z>, <https://doi.org/10.1007/s11263-005-3222-z>.
- [30] R. REBONATO AND P. JAECKEL, *The most general methodology to create a valid correlation matrix for risk management and option pricing purposes*, Journal of Risk, 2 (2001), pp. 17–27, <https://doi.org/10.2139/ssrn.1969689>.
- [31] Y. THANWERDAS, *Riemannian and stratified geometries of covariance and correlation matrices*, PhD thesis, Université Côte d'Azur and Inria, 2022.
- [32] Y. THANWERDAS AND X. PENNEC, *Geodesics and Curvature of the Quotient-Affine Metrics on Full-Rank Correlation Matrices*, in Proceedings of GSI 2021 - 5th conference on Geometric Science of Information, vol. 12829 of Lecture Notes in Computer Science, Paris, France, July 2021, Springer International Publishing, pp. 93–102, <https://hal.archives-ouvertes.fr/hal-03157992>.
- [33] Y. THANWERDAS AND X. PENNEC, *O(n)-invariant Riemannian metrics on SPD matrices*. Preprint, Sept. 2022, <https://hal.archives-ouvertes.fr/hal-03338601>.
- [34] Y. THANWERDAS AND X. PENNEC, *Theoretically and computationally convenient geometries on full-rank correlation matrices*. Preprint, Jan. 2022, <https://hal.archives-ouvertes.fr/>

- [hal-03527072](#).
- [35] G. VAROQUAUX, F. BARONNET, A. KLEINSCHMIDT, P. FILLARD, AND B. THIRION, *Detection of brain functional-connectivity difference in post-stroke patients using group-level covariance modeling*, in Medical Image Computing and Computer Added Intervention, D. Shen, A. Frangi, and G. Szekely, eds., Lecture Notes in Computer Science, Beijing, China, Sept. 2010, Tianzi Jiang, Springer, pp. 200–208, https://doi.org/10.1007/978-3-642-15705-9_25, <https://hal.inria.fr/inria-00512417>.
- [36] Z. WANG, B. C. VEMURI, Y. CHEN, AND T. H. MARECI, *A constrained variational principle for direct estimation and smoothing of the diffusion tensor field from complex dwi*, IEEE Transactions on Medical Imaging, 23 (2004), pp. 930–939, <https://doi.org/10.1109/TMI.2004.831218>.