



Can physiological signals estimation improve fMRI connectivity assessment in anesthetized rats?

Mikael Naveau, Coline Vanschamelhout, Marin Mauboussin, Romaric Saulnier, Palma Pro-Sistiaga, Thomas Freret, Olivier M Etard

► To cite this version:

Mikael Naveau, Coline Vanschamelhout, Marin Mauboussin, Romaric Saulnier, Palma Pro-Sistiaga, et al.. Can physiological signals estimation improve fMRI connectivity assessment in anesthetized rats?. OHBM2022, Jun 2022, Glasgow, United Kingdom. . hal-03878758

HAL Id: hal-03878758

<https://hal.science/hal-03878758>

Submitted on 30 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



COMETE
UMR 1075 UNICAEN / INSERM



Can physiological signals estimation improve fMRI connectivity assessment in anesthetized rats?

Naveau Mikael¹, Vanschamelhout Coline², Mauboussin Marin², Saulnier Romaric¹, Pro Palma¹, Freret Thomas², Etard Olivier²

¹ Normandie Université, UNICAEN, CNRS, INSERM, UAR-3408, US-50 CYCERON, Caen, France

² Normandie Université, UNICAEN, INSERM, COMETE, CYCERON, CHU Caen, Caen, France

Background

Resting-state functional magnetic resonance imaging (fMRI) is a common translational method to evaluate brain functional connectivity (FC) in preclinical neurologic applications. However, unlike human studies, preclinical fMRI studies are usually performed under anesthetic condition. The confounding effect of the anesthesia on fMRI measurements is not clear. Indeed, a controversy exists between those who claim that anesthesia change FC (Paasonen et al., 2018) and other who maintain that FC is preserved (Becq et al., 2020). Our hypothesis is that physiological changes induced by anesthesia could explain this discrepancy. While in human studies, fMRI analysis generally includes cleaning of physiological-noise (Glover et al., 2000; Kassinosopoulos & Mitsis, 2019), preclinical study processing is rather often limited to a low-pass temporal filtering. We aimed first to demonstrate that the global MRI signal of animal brain includes, like in human, a physiological component. Second, that a normalization of this component based on global signal can reduce variability of FC assessment.

Results

Methods





Figure 1: Time-frequency analysis of the physiological and MRI signals. Fourier transforms are computed on a 10s time window with a 5s overlap over the 60min fMRI experiment. Photoplethysmogram (PPG) of the rat-6 is excluded due to sensor failure. Colormap of the power spectrum are the same across animals but scaled independently for each signal type. Blue arrows show some examples of the main frequencies, harmonics and confounds of each physiological signal.




Figure 2: A Symmetric matrices of the standard deviation of the Functional Connectivity (FC) over time for each animal (in rows). Regions of interest are ordered this way : Right Motor Cortex (RMC), Left Motor Cortex (LMC), Right Somatosensory Cortex (RSSC), Left Somatosensory Cortex (LSSC), Right Thalamus (RTh), Left Thalamus (LTh), Right Ventricle (RV), Left Ventricle (LV). **B** Distribution comparison of the standard deviation of the FC across animals (Kruskal-Wallis). For each pair of regions, the five jittered plots correspond to (1) Raw FC, (2) partial-FC with the Global Signal (GS), (3) partial-FC with comp1 of GS, (4) partial-FC with comp1-2 of the GS, (5) partial-FC with comp1-2-3 of the GS. Gray dot means that the distribution of the FC does not differ statistically of the Raw FC distribution ($\alpha=0.05$, Tukey-HSD corrected for multiple comparisons).

Conclusion

We showed that the physiological variability of both respiratory and cardiac cycles is captured by the global brain fMRI signal. When the global signal was used as a confounding factor to compute the FC, the variability across the different stages of the anesthesia is reduced, even if there were strong physiological modifications.

We now have to investigate the best way to correct the fMRI signal for the physiological variation. Our best bet is to use the methods developed for human brain fMRI preprocessing and adapt them for preclinical imaging.

References

- Becq, G. J.-P. C., Habet, T., Collomb, N., Faucher, M., Delon-Martin, C., Coizet, V., Achard, S., & Barbier, E. L. (2020). **Functional connectivity is preserved but reorganized across several anesthetic regimes.** *NeuroImage*, 219, 116945. <https://doi.org/10.1016/j.neuroimage.2020.116945>
- Behzadi, Y., Restom, K., Liu, J., & Liu, T. T. (2007). **A component based noise correction method (CompCor) for BOLD and perfusion based fMRI.** *NeuroImage*, 37(1), 90-101. <https://doi.org/10.1016/j.neuroimage.2007.04.042>
- Glover, G. H., Li, T. Q., & Ress, D. (2000). **Image-based method for retrospective correction of physiological motion effects in fMRI : RETROICOR.** *Magnetic Resonance in Medicine*, 44(1), 162-167. [https://doi.org/10.1002/1522-2594\(200007\)44:1<162::aid-mrm23>3.0.co;2-e](https://doi.org/10.1002/1522-2594(200007)44:1<162::aid-mrm23>3.0.co;2-e)
- Kassinopoulos, M., & Mitsis, G. D. (2019). **Identification of physiological response functions to correct for fluctuations in resting-state fMRI related to heart rate and respiration.** *NeuroImage*, 202, 116150. <https://doi.org/10.1016/j.neuroimage.2019.116150>
- Paasonen, J., Stenroos, P., Salo, R. A., Kiviniemi, V., & Gröhn, O. (2018). **Functional connectivity under six anesthesia protocols and the awake condition in rat brain.** *NeuroImage*, 172, 9-20. <https://doi.org/10.1016/j.neuroimage.2018.01.014>

The French Research ministry and Ethical Comity approved this preclinical study (#28792).

Contact: olivier.etard@unicaen.fr or naveau@cyceron.fr

France Life Imaging fund the inscription fees of M. Naveau to OHBM2022

