

Distribution and accumulation of metals and metalloids in planktonic food webs of the Mediterranean Sea (MERITE-HIPPOCAMPE campaign)

Sandrine Chifflet^{a*}, Nicolas Briant^b, Javier Angel Tesán-Onrubia^a, Noureddine Zaaboub^c, Sirine Armi^c, Olivier Radakovitch^{d,e}, Daniel Bănaru^a, Marc Tedetti^a

^a Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille, France

^b Ifremer, CCEM Contamination Chimique des Écosystèmes Marins, F-44000 Nantes, France

^c Institut National des Sciences et Technologies de la Mer (INSTM), 28, rue 2 mars 1934, Salammbô 2025, Tunisia

^d Aix Marseille Univ., CNRS, IRD, Collège de France, INRAE, CEREGE, 13545 Aix-en-Provence Cedex 4, France

^e IRSN (Institut de Radioprotection et de Sûreté Nucléaire), PSE-ENV/SRTE/LRTA, Saint-Paul-Les-Durance, France

* Corresponding author: sandrine.chifflet@mio.osupytheas.fr; Phone: + 33 4 86 09 05 32;
Fax: + 33 4 91 82 96 41

For submission to Marine Pollution Bulletin Special issue “*Plankton and Contaminants in the Mediterranean Sea: Biological pump and interactions from regional to global approaches*”

as full-length research paper

Revised version

04 November 2022

APPENDIX 1

Data processing: determination of biotic metal concentrations

Metals and metalloids concentrations in samples can be expressed using the mass balance formula (equation 1):

$$[M]_{sample} = a \cdot [Ti]_a + b \cdot [POP]_b \quad (1)$$


where a is the Ti-normalized elemental ratio in geogenic component; b is the P-normalized elemental ratio in biotic component; $[M]_{sample}$ is the elemental concentration in the sample; $[Ti]_a$ is the Ti concentration in geogenic component and $[POP]_b$ is the particulate organic phosphorus concentration in the biotic component. Assuming that P and Ti concentrations in samples are mainly due to the biotic and geogenic components, respectively ($[POP]_{sample} \equiv [POP]_b$; $[Ti]_{sample} \equiv [Ti]_a$), we can turn the mass balance formula into the first-order equation 2:

$$\frac{[M]_{sample}}{[Ti]_{sample}} \equiv S \cdot \frac{[POP]_{sample}}{[Ti]_{sample}} + C \quad (2)$$

where S is the slope of the regression line between M/Ti and POP/Ti in the sample; C is a constant value. Following this basic model, metals and metalloids concentrations in biotic component were evaluated from equation 3:

$$[bM] \equiv b \cdot [POP]_{sample} \quad (3)$$

The slope S was calculated for each station and for each element using the samples in the 7 particle size classes F1-7 (Fig. S1). Detailed of $[M]_{sample}$ and $[POP]_{sample}$ were presented in Table S3.






Figure S1. Linear regressions between $(Zn/Ti)_{sample}$ and $(POP/Ti)_{sample}$ in St1 (northern coastal zone), St11 (offshore zone) and St19 (southern coastal zone). Data were from particulate size classes (F1: 0.8-3; F2: 3-20; F3: 60-200; F4: 200-500; F5: 500-1000; F6: 1000-2000 and F7 > 2000 μm) collected at the deep chlorophyll maximum. Concentrations were expressed in $\mu\text{g/g}$ for Zn and Ti, and in mg/g for POP. At St19, data for F6 and F7 were not available. Zn_{sample} concentrations in F1 at St11 and St19 (red point) were higher than expected and were not included in the linear regression.

APPENDIX 2

Data processing: determination of trophic magnification factors

Due to the complexity and variability of taxonomic groups, planktonic marine food webs were structured in trophic level (TL) according to the following equation ([Kidd et al, 2018](#)):

$$TL_{Fn} = 1.0 + \frac{(\delta^{15}N_{Fn} - \delta^{15}N_{baseline})}{\Delta^{15}N} \quad (1)$$

where TL_{Fn} is the TL of the sample corresponding to the fraction Fn, $\delta^{15}N_{Fn}$ and $\delta^{15}N_{baseline}$ are the $\delta^{15}N$ values of fractions Fn and a baseline, respectively. The fraction F1 (0.8-3 μ m) is assumed to be the baseline fraction with a TL_{F1} defined to 1.0 as TL corresponding to the phytoplankton communities. $\Delta^{15}N$ ($= \delta^{15}N_{prey} - \delta^{15}N_{predator}$) is the isotopic ^{15}N fractionation between prey and predator. For marine food webs, $\Delta^{15}N$ vary from +3.3 to +3.8‰ ([Liu et al., 2019](#)) and a value of +3.4‰ is recommend without precise knowledge on the composition of the trophic structure ([Post, 2002](#)). TL determination were detailed in a companion study ([Tésan-Onrubia et al., 2022](#)) and results shown in Table S4.

Trophic magnification factor (TMF) is most often derived from the slope of the regression of log-transformed element against their corresponding trophic level (TL) and, N isotope ($\delta^{15}N$) is used as proxy to assess the relative trophic position of an organism in marine food webs.

$$\log_{10}(bM) = S \cdot (TL) + C \quad (2)$$

$$TMF = 10^S \quad (3)$$

where bM is the biotic metal concentration, S is the slope of the linear equation 2 and, TL is the trophic level corresponding to the sample (Fig. S2). Specific TMF (equation 3) were calculated for each geographical area (the northern coastal zone, the offshore zone and the southern coastal zone) from their respective data (stations and fractions).






Figure S2. Determination of the linear regressions between the log-transformed of bZn concentrations in relation to the trophic level (TL) in the 3 geographical areas (the northern coastal zone, the offshore zone and the southern coastal zone). The northern coastal, offshore and southern coastal zones included 25, 18 and 12 data from available stations and fractions. Detailed values were presented in Table S4.

References:

- Kidd K.A., Burkhard L.P., Babut M., Borga K., Muir D., Perceval O., Ruedel H., Woodburn K., Embry M.R., 2018. Practical advice for selecting or determining trophic magnification factors for application under the European Union Water Framework Directive. *Integr. Environ. Assess. Manag.*, 15, 266-277.
- Liu J., Cao L., Dou S., 2019. Trophic transfer, biomagnification and risk assessments of four common heavy metals in the food web of Laizhou Bay, the Bohai Sea. *Sci. Tot. Environ.*, 670, 508-522.
- Post DM., 2002. Using stable isotopes to estimate trophic position: Models, methods, and assumptions. *Ecol.*, 83, 703-718.
- Tesán-Onrubia, J.A., Tedetti, M., Carlotti, F., Tenaille, M., Guilloux, L., Pagano, M., Lebreton, B., Guillou, G., Fierro-Gonzalez, P., Guigue, C., Chifflet, S., Garcia, T., Boudriga, I., Belhassen, M., Zouari, A., Bănaru, D. Spatial variations of stable isotope ratios and biochemical content of size-fractionated plankton in the Mediterranean Sea (MERITE-HIPPOCAMPE campaign). *Mar. Pollut. Bull.*, submitted to this special issue.

Table S1. GPS position of sampling stations

<i>Station</i>	Latitude (North)	Longitude (East)
<i>St01</i>	43°3.819'	5°59.080'
<i>St02</i>	42°56.020'	5°58.041'
<i>St03</i>	43°8.150'	5°15.280'
<i>St04</i>	43°14.500'	5°17.500'
<i>St09</i>	41°53.508'	6°19.998'
<i>St10</i>	40°18.632'	7°14.753'
<i>St11</i>	39°7.998'	7°41.010'
<i>St15</i>	36°12.883'	11°7.641'
<i>St17</i>	34°30.113'	11°43.573'
<i>St19</i>	33°51.659'	11°18.509'

Table S2. Detail of the quality control of metals metalloids analyses (certified material for marine plankton, BCR 414; blank filters, MCE; limit of detection, LOD). Mean metals and metalloids concentrations and standard deviation (σ) associated are expressed in $\mu\text{g/g}$. The limit of detection (LOD) are expressed in $\mu\text{g/L}$.

	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	Ti	V	Zn
<i>CMR BCR 414</i>											
Mean ($n = 3$)	8.03	0.36	21.97	29.08	1702.68	-	19.27	-	-	8.71	121.70
σ ($n = 3$)	0.07	0.04	2.62	0.96	160.79	-	0.10	-	-	0.34	19.31
Recovery (%)	117.7	93.9	92.3	98.6	92.0	-	102.5	-	-	107.6	109.1
<i>BLANK MCE filter, 47mm</i>											
Mean $0.8\mu\text{m}$ ($n = 6$)	0.03	0.05	2.21	0.41	10.57	0.33	0.31	0.01	2.65	0.06	8.16
$\sigma 0.8\mu\text{m}$ ($n = 6$)	0.01	0.01	0.64	0.3	3.94	0.09	0.05	0.00	0.29	0.01	5.14
Mean $3\mu\text{m}$ ($n = 3$)	0.03	0.05	1.48	0.48	10.02	0.49	0.34	0.02	2.43	0.06	3.44
$\sigma 3\mu\text{m}$ ($n = 3$)	0.01	0.01	0.07	0.16	2.16	0.01	0.03	0.00	0.01	0.01	0.08
LOD ($\mu\text{g/L}$)	0.01	0.005	0.008	0.01	0.2	0.008	0.2	0.005	0.02	0.008	0.1

Table S3. Detail of total metals and metalloids (M_{sample}) concentrations ($\mu\text{g/g}$) and particulate organic phosphorus ($\text{POP}_{\text{sample}}$) concentrations (mg/g) in samples from the MERITE-HIPPOCAMPE campaign. Particles were collected in the deep chlorophyll maximum, at 10 stations (St1-4, St9-11, St15, St1, St19) and sieved into 7 size fractions (F1: 0.8-3 μm ; F2: 3-20 μm ; F3: 60-200 μm ; F4: 200-500 μm ; F5: 500-1000 μm ; F6: 1000-2000 μm ; F7 > 2000 μm).

Station	Fraction	POP	Ti	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
St01	F1	1.20	54.75	1.85	0.65	0.08	6.80	35.39	0.84	1.22	-	0.27	50.01
St01	F2	1.92	222.40	12.59	1.04	2.98	21.10	2046.29	155.80	13.54	0.28	9.62	54.16
St01	F3	4.67	150.25	14.88	1.40	4.93	8.63	964.05	23.85	6.85	1.24	4.35	148.10
St01	F4	5.32	52.66	10.48	1.15	2.39	8.95	303.00	6.85	4.41	0.99	1.38	126.70
St01	F5	5.94	153.94	20.28	1.10	7.82	10.85	940.75	20.17	8.73	4.28	4.05	197.97
St01	F6	2.48	445.09	23.40	0.53	8.91	11.60	2243.06	66.28	11.13	14.61	12.12	125.54
St01	F7	3.84	75.24	5.19	0.38	0.78	4.76	261.46	4.96	2.73	0.29	0.96	57.85
St02	F1	5.81	178.12	14.04	3.45	14.69	72.41	1151.66	96.98	22.61	0.32	5.67	702.33
St02	F2	1.56	164.06	2.31	0.10	-	10.63	311.84	20.39	5.44	0.33	1.02	35.16
St02	F3	2.45	59.07	10.22	1.18	18.63	20.57	630.02	7.72	9.09	0.94	1.42	216.00
St02	F4	4.31	36.71	13.02	1.24	8.12	13.49	498.32	6.47	6.43	0.70	0.74	193.30
St02	F5	3.78	93.59	10.67	0.64	39.55	13.58	1404.05	13.11	18.07	1.94	1.36	214.78
St02	F6	3.16	132.60	6.38	0.39	78.14	12.81	5336.71	32.82	25.22	1.96	1.54	160.32
St02	F7	3.72	204.53	6.31	1.44	41.74	8.21	1207.48	13.19	13.47	3.43	1.51	242.53
St03	F1	3.05	281.64	7.63	1.51	2.80	39.57	671.96	16.39	10.00	0.13	1.81	74.43
St03	F2	5.15	217.47	13.35	2.66	9.47	53.76	1201.47	74.83	26.41	0.62	3.81	278.15
St03	F3	1.64	79.50	6.68	0.58	12.99	9.69	537.55	7.74	8.89	5.10	2.40	165.67
St03	F4	3.97	54.81	5.90	0.60	5.85	10.42	277.98	5.52	7.41	5.99	1.81	530.69
St03	F5	3.71	61.61	5.92	0.40	7.92	6.82	311.61	6.45	8.03	8.61	2.01	110.16
St03	F6	3.17	-	-	-	-	-	-	-	-	-	-	-
St03	F7	1.74	47.76	11.46	0.21	3.00	5.78	202.60	6.31	5.72	10.24	1.32	60.04
St04	F1	1.48	113.04	16.22	1.38	6.88	28.60	975.87	29.86	7.49	0.10	33.25	221.92
St04	F2	2.26	100.38	13.58	0.05	-	1.86	225.40	9.30	5.29	0.16	20.64	41.83
St04	F3	1.40	32.29	17.27	0.08	3.10	4.07	345.98	5.09	2.25	1.16	44.83	33.63
St04	F4	1.67	91.86	55.40	0.27	5.33	8.37	816.25	11.94	4.57	3.78	181.14	70.51
St04	F5	1.45	197.55	8.78	0.21	18.17	12.55	1483.43	20.42	9.78	12.28	40.97	111.47
St04	F6	-	243.14	7.26	0.13	23.03	16.99	1724.68	29.33	18.73	37.75	43.28	149.42
St04	F7	-	-	-	-	-	-	-	-	-	-	-	-
St09	F1	4.61	272.74	32.77	9.13	12.67	22.59	837.73	20.41	12.21	0.13	4.30	1406.61
St09	F2	2.79	75.47	24.08	1.61	-	9.38	311.79	16.13	5.29	0.11	2.25	45.59
St09	F3	2.70	59.35	9.61	1.27	2.14	5.14	268.96	4.48	3.66	1.23	1.42	124.11
St09	F4	3.19	16.46	13.63	1.76	0.48	9.65	50.21	3.07	6.47	1.90	0.48	149.54
St09	F5	3.11	37.56	12.98	1.88	1.81	9.36	103.79	3.05	6.92	2.01	0.78	197.92
St09	F6	-	37.78	24.25	0.65	0.62	7.50	151.64	2.80	2.17	1.65	1.39	132.38
St09	F7	-	125.60	1.92	0.75	2.18	11.59	76.35	9.63	3.85	0.03	0.69	63.88
St10	F1	1.30	82.28	11.71	1.16	9.97	11.05	159.17	5.06	3.90	0.01	2.89	419.09
St10	F2	4.82	266.50	59.10	5.26	-	54.68	855.84	73.65	21.62	0.49	9.76	190.45

<i>St10</i>	<i>F3</i>	4.61	267.03	12.90	1.64	13.35	9.81	1633.36	31.99	12.47	7.39	7.47	189.76
<i>St10</i>	<i>F4</i>	5.24	133.86	7.40	1.13	4.53	9.14	890.99	15.60	8.66	11.62	4.30	177.28
<i>St10</i>	<i>F5</i>	4.24	186.16	8.87	0.82	5.26	8.76	1151.63	24.55	10.05	22.25	5.91	141.08
<i>St10</i>	<i>F6</i>	4.98	274.68	11.53	0.71	8.39	9.59	1466.68	37.43	14.16	34.95	7.61	139.39
<i>St10</i>	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>St11</i>	<i>F1</i>	1.65	59.70	3.35	0.55	2.01	9.20	185.21	2.98	2.45	0.03	0.78	294.65
<i>St11</i>	<i>F2</i>	10.31	516.78	50.05	4.53	-	71.22	1827.99	154.07	36.26	0.78	13.73	297.99
<i>St11</i>	<i>F3</i>	5.80	191.10	16.74	1.47	5.05	9.32	892.33	16.07	7.50	17.60	6.47	147.16
<i>St11</i>	<i>F4</i>	6.39	103.69	11.07	1.23	1.57	8.17	430.96	7.64	6.12	16.49	9.20	133.56
<i>St11</i>	<i>F5</i>	6.64	417.43	57.16	5.91	11.86	40.09	2013.35	37.76	30.71	82.17	36.96	580.68
<i>St11</i>	<i>F6</i>	6.48	110.36	12.34	1.07	1.88	9.21	508.19	10.11	4.77	16.57	3.94	129.32
<i>St11</i>	<i>F7</i>	4.52	32.72	24.89	1.72	0.27	5.11	51.11	1.72	0.78	1.96	0.55	127.42
<i>St15</i>	<i>F1</i>	0.72	25.49	1.98	1.05	1.24	8.09	22.95	0.14	2.33	-	0.67	176.64
<i>St15</i>	<i>F2</i>	10.52	617.62	42.59	8.37	-	65.05	4624.77	161.33	28.58	0.93	66.75	438.98
<i>St15</i>	<i>F3</i>	3.03	316.03	17.87	0.68	5.98	9.28	1322.87	28.57	8.79	12.65	14.96	148.03
<i>St15</i>	<i>F4</i>	3.83	234.89	9.93	0.84	4.22	7.33	1153.40	15.07	5.80	14.70	10.62	130.82
<i>St15</i>	<i>F5</i>	5.47	146.14	11.45	0.62	2.15	9.14	692.88	13.66	5.03	7.70	7.52	130.38
<i>St15</i>	<i>F6</i>	-	264.53	11.46	0.59	6.16	8.18	1740.47	18.71	5.17	9.72	10.80	108.30
<i>St15</i>	<i>F7</i>	-	59.99	8.77	0.40	1.14	3.26	393.85	5.47	1.47	4.46	3.41	42.61
<i>St17</i>	<i>F1</i>	2.81	240.31	6.42	0.65	2.96	19.25	246.77	1.56	6.30	-	1.18	308.61
<i>St17</i>	<i>F2</i>	2.68	675.56	10.63	0.69	11.71	20.25	4803.63	63.26	11.69	0.32	19.30	50.58
<i>St17</i>	<i>F3</i>	1.14	178.13	31.09	0.16	3.26	5.81	920.72	26.82	6.03	7.76	9.30	29.63
<i>St17</i>	<i>F4</i>	1.52	290.34	34.57	0.28	4.90	5.85	1475.74	34.71	6.74	12.62	11.62	39.70
<i>St17</i>	<i>F5</i>	-	590.41	3.11	0.71	150.45	21.92	2238.08	37.59	158.97	1.90	6.90	192.53
<i>St17</i>	<i>F6</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>St17</i>	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-	-
<i>St19</i>	<i>F1</i>	12.83	294.63	13.09	20.33	-	40.99	1002.86	18.47	17.30	0.33	3.39	777.14
<i>St19</i>	<i>F2</i>	0.20	596.53	9.36	3.41	9.60	14.09	4740.58	119.91	13.42	0.28	16.18	61.76
<i>St19</i>	<i>F3</i>	2.96	278.59	5.83	1.13	8.12	10.08	857.71	20.56	12.57	3.14	3.49	104.96
<i>St19</i>	<i>F4</i>	4.26	116.46	4.30	1.10	2.10	4.52	411.88	8.44	2.12	3.88	1.76	105.83
<i>St19</i>	<i>F5</i>	3.88	251.62	4.79	1.10	13.68	7.02	850.06	17.62	11.90	11.14	3.60	132.35
<i>St19</i>	<i>F6</i>	-	842.93	5.55	0.52	18.68	5.62	3858.54	67.05	8.29	82.28	13.57	67.74
<i>St19</i>	<i>F7</i>	-	542.25	2.35	0.56	118.58	8.92	2170.58	27.20	79.21	1.90	5.48	87.51

Table S4. Detail of biotic metal (bM) concentrations ($\mu\text{g/g}$) according to the 7 size fractions (F1: 0.8-3 μm ; F2: 3-20 μm ; F3: 60-200 μm ; F4: 200-500 μm ; F5: 500-1000 μm ; F6: 1000-2000 μm ; F7 > 2000 μm) and the 10 stations (St1-4. St9-11. St15. St1. St19) of the MERITE-HIPPOCAMPE campaign. TL determination were discussed in a companion study ([Tésan-Onrubia et al., 2022](#)).

Station	Fraction	TL	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
St01	F1	1.0	1.82	0.21	0.08	1.58	35	0.84	0.55	0.00	0.27	25
	F2	1.1	2.92	0.33	2.98	2.53	120	2.75	0.88	0.28	0.52	40
	F3	1.7	7.09	0.80	4.93	6.15	292	6.68	2.13	0.82	1.26	97
	F4	1.9	8.08	0.91	2.39	7.01	303	6.85	2.43	0.93	1.38	111
	F5	2.2	9.02	1.01	7.82	7.82	371	8.49	2.71	1.04	1.60	124
	F6	2.2	3.77	0.42	7.39	3.27	155	3.55	1.13	0.43	0.67	52
	F7	2.5	5.19	0.38	0.78	4.76	240	4.96	1.75	0.29	0.96	58
St02	F1	1.0	14.04	1.74	14.69	17.32	527	19.81	2.36	0.32	2.64	261
	F2	1.6	2.31	0.10	-	4.65	142	5.31	0.63	0.20	0.71	35
	F3	2.0	7.83	0.73	17.34	7.30	222	7.72	0.99	0.31	1.11	110
	F4	2.0	13.02	1.24	8.12	12.84	391	6.47	1.75	0.54	0.74	193
	F5	2.1	10.67	0.64	26.76	11.26	343	12.88	1.53	0.48	1.36	170
	F6	2.1	6.38	0.39	22.40	9.43	287	10.78	1.28	0.40	1.44	142
	F7	2.3	6.31	1.11	26.36	8.21	338	12.69	1.51	0.47	1.51	167
St03	F1	1.0	3.43	0.51	2.80	4.87	135	8.60	1.16	0.13	1.27	74
	F2	1.3	5.80	0.87	9.47	8.23	228	14.52	1.95	0.62	2.15	166
	F3	2.3	1.85	0.28	3.88	2.63	73	4.64	0.62	0.30	0.69	53
	F4	2.5	4.47	0.60	5.85	6.34	176	5.52	1.51	0.71	1.65	128
	F5	2.5	4.17	0.40	7.92	5.92	164	6.45	1.41	0.67	1.54	110
	F6	-	-	-	-	-	-	-	-	-	-	-
	F7	2.2	1.96	0.21	3.00	2.78	77	4.91	0.66	0.31	0.73	56
St04	F1	1.0	8.65	0.04	1.88	2.47	121	2.15	0.89	0.10	33.25	29
	F2	1.5	13.18	0.05	-	1.86	184	3.28	1.36	0.16	20.64	42
	F3	1.8	8.18	0.04	1.78	2.34	114	2.03	0.84	0.11	41.72	27
	F4	1.8	9.71	0.04	2.11	2.78	135	2.41	1.00	0.13	49.51	33
	F5	1.7	8.44	0.04	1.83	2.41	118	2.10	0.87	0.11	40.97	28
	F6	-	-	-	-	-	-	-	-	-	-	-
	F7	-	-	-	-	-	-	-	-	-	-	-
St09	F1	1.3	18.18	2.35	12.67	13.49	241	4.39	3.28	0.13	1.36	245
	F2	1.4	10.98	1.42	-	8.15	145	2.65	1.98	0.11	0.82	46
	F3	2.4	9.61	1.27	2.14	5.14	141	2.57	1.92	1.23	0.79	124
	F4	2.7	12.57	1.63	0.48	9.33	50	3.04	2.27	1.90	0.48	150
	F5	3.2	12.26	1.59	1.81	9.10	104	2.96	2.21	2.01	0.78	165
	F6	-	-	-	-	-	-	-	-	-	-	-
	F7	-	-	-	-	-	-	-	-	-	-	-
St10	F1	1.0	0.60	0.22	0.27	1.94	159	1.46	0.96	0.01	0.97	41
	F2	1.4	2.25	0.81	N/A	7.21	643	5.45	3.56	0.49	3.60	153
	F3	2.7	2.15	0.78	0.95	6.90	615	5.21	3.41	0.62	3.44	147

	<i>F4</i>	2.8	2.44	0.88	1.08	7.83	697	5.91	3.86	0.70	3.91	166
	<i>F5</i>	3.1	1.98	0.72	0.88	6.35	565	4.79	3.13	0.57	3.17	135
	<i>F6</i>	3.6	2.32	0.71	1.03	7.44	663	5.62	3.67	0.67	3.71	139
	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-
<i>St11</i>	<i>F1</i>	1.0	3.35	0.10	0.73	1.62	134	2.21	0.94	0.03	0.56	42
	<i>F2</i>	1.1	50.05	0.64	-	10.15	838	13.83	5.87	0.78	3.48	260
	<i>F3</i>	2.1	16.74	0.36	2.57	5.72	472	7.79	3.31	17.60	1.96	147
	<i>F4</i>	2.2	11.07	0.40	1.57	6.30	431	7.64	3.64	16.49	2.16	134
	<i>F5</i>	2.4	35.55	0.41	2.93	6.54	540	8.90	3.78	25.51	2.24	167
	<i>F6</i>	2.5	12.34	0.40	1.88	6.38	508	8.70	3.69	16.57	2.19	129
	<i>F7</i>	2.8	24.24	0.28	0.27	4.46	51	1.72	0.78	1.96	0.55	114
<i>St15</i>	<i>F1</i>	1.0	0.60	0.39	1.23	0.91	23	0.14	0.67	-	0.12	32
	<i>F2</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F3</i>	2.0	2.53	0.68	5.15	3.82	289	4.19	2.81	6.53	0.50	133
	<i>F4</i>	2.1	3.20	0.84	6.52	4.83	366	5.30	3.56	8.26	0.63	131
	<i>F5</i>	1.8	4.58	0.62	9.32	6.90	523	7.58	5.03	7.70	0.90	130
	<i>F6</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-
<i>St17</i>	<i>F1</i>	1.0	3.66	0.65	3.42	2.99	205	1.56	2.22	-	1.18	93
	<i>F2</i>	1.4	3.49	0.62	3.26	2.86	196	55.31	2.12	0.32	19.30	51
	<i>F3</i>	1.5	1.49	0.16	1.39	1.22	84	23.63	0.91	7.76	9.22	30
	<i>F4</i>	1.7	1.99	0.28	1.86	1.62	111	31.45	1.21	12.62	11.62	40
	<i>F5</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F6</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-
<i>St19</i>	<i>F1</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F2</i>	1.6	0.14	0.03	0.39	0.36	13	0.02	0.08	0.07	0.08	5
	<i>F3</i>	1.8	2.07	0.51	5.79	5.36	185	0.34	1.22	0.96	1.12	70
	<i>F4</i>	1.9	2.97	0.74	2.10	4.52	266	0.48	1.75	1.38	1.61	100
	<i>F5</i>	2.2	2.71	0.67	7.58	7.02	243	0.44	1.59	1.26	1.47	91
	<i>F6</i>	-	-	-	-	-	-	-	-	-	-	-
	<i>F7</i>	-	-	-	-	-	-	-	-	-	-	-

Table S5: Results of ANOVA (A) or Kruskal-Wallis (KW) tests per element to examine the potential variations of bM concentrations across all stations and fractions (n = 55). Significant differences ($p < 0.05$) are in bold.

	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
Test	A	KW	A	A	A	KW	A	A	KW	A
<i>p</i> -Value	0.003	0.008	0.015	0.026	0.005	0.038	<0.0001	0.012	0.086	0.009

Table S6. Detail of bM/POP ratios in planktonic food webs per elements and stations.

	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
<i>North coastal zone</i>										
St01	0.62	0.04	0.79	0.64	31.98	0.74	0.24	0.03	0.16	9.54
St02	0.96	0.06	2.96	1.41	50.34	1.77	0.21	0.03	0.25	19.64
St03	0.47	0.04	1.09	0.78	24.58	1.39	0.20	0.03	0.26	13.96
St04	2.41	0.01	0.63	0.74	45.11	0.82	0.32	0.02	12.74	9.17
<i>Offshore zone</i>										
St09	1.60	0.13	0.72	1.34	23.83	0.54	0.37	0.08	0.16	17.88
St10	0.19	0.05	0.11	0.73	73.03	0.64	0.39	0.02	0.46	13.53
St11	1.37	0.02	0.18	0.48	39.08	0.68	0.27	0.40	0.19	10.56
<i>South coastal zone</i>										
St15	0.35	0.08	1.01	0.61	45.90	0.65	0.49	0.29	0.10	15.02
St17	0.54	0.05	0.73	0.52	40.51	9.37	0.42	0.78	3.84	11.87
St19	0.29	0.05	0.99	0.81	34.70	0.06	0.22	0.07	0.23	10.53

Table S7: Results of ANOVA (A) or Kruskal-Wallis (KW) tests per element to examine the potential variations of bM/POP ratios across all stations and fractions (n = 55). Significant differences (p < 0.05) are in bold.

	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
<i>Test</i>	A	KW	A	KW	A	A	A	A	A	A
<i>p-Value</i>	0.014	0.038	0.035	0.022	0.298	0.018	< 0.0001	< 0.0001	0.112	0.579

Table S8: Results of Student (t) or Mann-Whitney-Wilcoxon (MW) tests per element to examine the potential differences in bM concentrations between geographical areas (n = 25 in the northern coastal zone; n = 18 in the offshore zone; n = 12 in the southern coastal zone). Significant differences ($p < 0.05$) are in bold.

	As	Cd	Cr	Cu	Fe	Mn	Ni	Sb	V	Zn
<i>Differences between the northern coastal zone and the offshore zone</i>										
<i>Test</i>	MW	MW	MW	MW	t	MW	t	MW	MW	MW
<i>p-Value</i>	0.207	0.066	< 0.001	0.190	0.040	0.428	< 0.001	0.015	0.724	0.014
<i>Differences between the northern coastal zone and the southern coastal zone</i>										
<i>Test</i>	t	MW	MW	MW	t	MW	t	MW	MW	MW
<i>p-Value</i>	< 0.001	0.713	0.221	0.049	0.956	0.227	0.082	0.010	0.327	0.554
<i>Differences between the offshore zone and the southern coastal zone</i>										
<i>Test</i>	MW	MW	MW	t	t	MW	t	MW	MW	MW
<i>p-Value</i>	0.015	0.158	0.021	0.002	0.037	0.346	0.005	0.408	0.573	0.001

Table S6: Elemental stoichiometry (bM/POP ratio, mmol/mol) in planktonic food webs including the 7 size fractions (F1-7) from the MERITE-HIPPOCAMPE campaign (Mediterranean Sea) detailed for the 10 stations (St1-4, St9-11, St15, St1, St19).

	As	Cd	Cu	Cr	Fe	Mn	Ni	Sb	V	Zn
<i>Northern coastal zone</i>										
<i>St01</i>	0.62	0.04	0.64	0.79	31.98	0.74	0.24	0.03	0.16	9.54
<i>St02</i>	0.96	0.06	1.41	2.96	50.34	1.77	0.21	0.03	0.25	19.64
<i>St03</i>	0.47	0.04	0.78	1.09	24.58	1.39	0.20	0.03	0.26	13.96
<i>St04</i>	2.41	0.01	0.74	0.63	45.11	0.82	0.32	0.02	12.74	9.17
<i>Offshore zone</i>										
<i>St09</i>	1.60	0.13	1.34	0.72	23.83	0.54	0.37	0.08	0.16	17.88
<i>St10</i>	0.19	0.05	0.73	0.11	73.03	0.64	0.39	0.02	0.46	13.53
<i>St11</i>	1.37	0.02	0.48	0.18	39.08	0.68	0.27	0.40	0.19	10.56
<i>Southern coastal zone</i>										
<i>St15</i>	0.35	0.08	0.61	1.01	45.90	0.65	0.49	0.29	0.10	15.02
<i>St17</i>	0.54	0.05	0.52	0.73	40.51	9.37	0.42	0.78	3.84	11.87
<i>St19</i>	0.29	0.05	0.81	0.99	34.70	0.06	0.22	0.07	0.23	10.53