



HAL
open science

State sums for some super quantum link invariants

Louis-Hadrien Robert, Emmanuel Wagner

► **To cite this version:**

Louis-Hadrien Robert, Emmanuel Wagner. State sums for some super quantum link invariants. *Topology and Geometry*, 33 (1), EMS Press, pp.209-246, 2021, IRMA Lectures in Mathematics and Theoretical Physics, 10.4171/IRMA/33-1/12 . hal-03882732

HAL Id: hal-03882732

<https://hal.science/hal-03882732v1>

Submitted on 1 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

STATE SUMS FOR SOME SUPER QUANTUM LINK INVARIANTS

LOUIS-HADRIEN ROBERT AND EMMANUEL WAGNER

ABSTRACT. We present state sums for quantum link invariants arising from the representation theory of $U_q(\mathfrak{gl}_{N|M})$. We investigate the case of the N th exterior power of the standard representation of $U_q(\mathfrak{gl}_{N|1})$ and explicit the relation with Kashaev invariants.

CONTENTS

Introduction	1
Outline of the paper	3
1. Some q -identities	3
2. Colorings of MOY graphs	6
3. Link invariants	19
4. Non semi-simple invariants	20
Appendix A. MOY graphs, an algebraic approach	24
References	29

INTRODUCTION

In its 1990 ICM paper [Tur91], Turaev emphasized the prominent role of state sum models in low dimensional topology. Models of this kind were and remain an important part of what is called quantum topology. They are closely related to mathematical physics, quantum algebra and statistical mechanics. The most famous example of state sum model in low dimensional topology is probably the Kauffman bracket [Kau87]. Not only demonstrates it how methods of statistical mechanics and ideas of quantum field theory can be relevant in this subject as Witten explained [Wit89], but it is also the first step in a recent major development of quantum topology: categorification of quantum invariants. Indeed, Khovanov homology [Kho00] which categorifies the Jones polynomial is clearly inspired by the combinatorial definition of the Jones polynomial given by the Kauffman bracket.

In this paper, we give state sum models computing quantum invariants arising from the representation theory of the super quantum group $U_q(\mathfrak{gl}_{N|M})$. We believe these state sums will be useful in the quest of categorifying these quantum invariants. Some of the invariants described by these state sums have a non-semi-simple nature. The prototypical example of such non-semi-simple invariants is the Alexander–Conway polynomial.

Let us discuss—with a slightly biased point of view—some of the state sums and categorifications of the Alexander–Conway polynomial. The first categorification of the Alexander provided by the knot Floer homology of Ozsváth–Szabó and Rasmussen [OS04, Ras03] uses its interpretation as Reidemeister torsion. In a combinatorial version of Manolescu–Ozsváth–Szabó–Thurston using grid diagrams, they used explicitly a determinantal description [MOST07, Appendix]. They were many attempts to have a direct combinatorial approach using the state model developed by Kauffman in [Kau83], but finally Ozsváth–Szabó made the connection in [OS09] using a twisted version of the knot Floer homology. In [RW19], the authors of this paper provided a combinatorial categorification of the Alexander–Conway polynomial of knots starting from a representation theoretic interpretation (see [EPV19] for the connection between knot Floer homology and representation theory). The state sum model underlying this last categorification is one of the example of the present paper (see also [Vir06, Sar15]).

The paper [MOY98] by Murakami–Ohtsuki–Yamada is another prototypical example of a state sum model which found its full relevance and importance in the realm of the categorification process. They give an elementary description of Reshetikhin–Turaev invariant associated with exterior powers of the standard representation of $U_q(\mathfrak{gl}_N)$. The description of these quantum invariants by Murakami–Ohtsuki–Yamada proceeds in two steps. First, a link diagram is expressed as a formal linear combination of planar trivalent graphs (see also Kuperberg [Kup96]), then a positive state sum is provided for these graphs. Both of these steps were key ingredients for the categorification of $U_q(\mathfrak{gl}_N)$ -quantum invariants by Khovanov–Rozansky [KR08] (see also Mackaay–Stošić–Vaz [MSV09] and Wu [Wu14], Yonezawa [Yon11], Mazorchuk–Stroppel [MS09] and Sussan [Sus07] for colored versions).

State sum models should continue to be of importance and influential in quantum topology. For instance, they play a direct role in another version of the categorification of the previous invariants by the authors in [RW17]. A state sum model for closed 2-dimensional CW complexes called foams is given to produce a trivalent TQFT. This state sum model was reinvested by Khovanov and the first author [KR18] to investigate the four color theorem.

The present paper is firstly concerned with state sum models for exterior powers of the standard representation of $U_q(\mathfrak{gl}_{N|M})$. In particular the case $N = 0$, corresponds to invariants associated with symmetric powers of the standard representation of $U_q(\mathfrak{gl}_M)$. These invariants have been categorified in [RW18, QRS18, Cau17]. As proved in the present paper, the invariants only depend on $N - M$, providing in fact different state sum models for the same invariants. It is conjectured in [GGS18] that these various state sums could be categorified yielding potentially non-equivalent homological invariants.

Secondly, the paper investigates further the invariants associated with the N th exterior power of the standard representation of $U_q(\mathfrak{gl}_{N|1})$ which have vanishing quantum dimensions. These invariants can be renormalized, see [GPM10, QS15].

This paper provides a direct proof that they can be renormalized giving explicitly a state sum model. It also explains how these invariants are related to Kashaev invariants [Kas95] (see [GPM08] for a different proof of this latter fact). As ADO invariants [ADO92], this family of non-semisimple invariants provides interesting family of invariants generalizing the Alexander polynomial containing Kashaev invariants. The Alexander polynomial has been categorified in this framework [RW19] and the present state sum models provide a first step in the categorification of this family of invariants.

Outline of the paper.

- Section 1 is devoted to some combinatorics of q -binomials used in the rest of the paper.
- Section 2 gives the state sum models for planar graphs.
- Section 3 extend the state sum models to link diagrams.
- Section 4 investigates the non-semisimple case of the N -th exterior power of the standard representation of $U_q(\mathfrak{gl}_{N|1})$ and makes the connection with the Kashaev invariants.
- Appendix A provides the representation theoretic background; in particular the explicit maps providing the corresponding Reshetikhin–Turaev functors.

1. SOME q -IDENTITIES

In this section, we give some useful identities on quantum integers and quantum binomials. See [KC02] for a more detailed account. The first two lemmas and three next corollaries are obtained by direct computations or easy inductions and we omit their proofs. Unless otherwise specified, q is a formal parameter.

Definition 1.1. Let n be an integer, define the *quantum integer* $[n]$ by the following formula:

$$[n] := \frac{q^n - q^{-n}}{q - q^{-1}} = \begin{cases} \sum_{i=1}^n q^{-1-n+2i} & \text{if } n > 0, \\ 0 & \text{if } n = 0 \text{ and} \\ \sum_{i=-n}^{-1} -q^{1+n+2i} & \text{if } n < 0. \end{cases}$$

If k and n are two integers, define the *quantum binomial* $\begin{bmatrix} n \\ k \end{bmatrix}$ by the following formula:

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} \frac{\prod_{i=0}^{k-1} [n-i]}{\prod_{i=1}^k [i]} & \text{if } k \geq 0, \\ 0 & \text{else.} \end{cases}$$

Remark 1.2. For all integers k and n , one has:

$$[-n] = -[n] \quad \text{and} \quad \begin{bmatrix} -n \\ k \end{bmatrix} = (-1)^k \begin{bmatrix} n+k-1 \\ k \end{bmatrix}.$$

Lemma 1.3. Let n and m be two integers, the following identity holds:

$$[m+n] = q^{-n}[m] + q^m[n] = q^n[m] + q^{-m}[n].$$

Corollary 1.4. *Let k be an integer and $(a_h)_{1 \leq h \leq k}$ be a collection of k integers, then the following identity holds:*

$$\left[\sum_{h=1}^k a_h \right] = \sum_{h=1}^k q^{\sum_{i=1}^{h-1} a_i - \sum_{i=h+1}^k a_i} [a_h] = \sum_{h=1}^k q^{-\sum_{i=1}^{h-1} a_i + \sum_{i=h+1}^k a_i} [a_h].$$

Corollary 1.5. *Let k , m and n be three integers, then the following identity holds:*

$$[m+k][n] - [m][n+k] = [k][n-m].$$

Lemma 1.6. *Let n and k be two integers, then the following identity holds:*

$$\begin{bmatrix} n \\ k \end{bmatrix} = q^k \begin{bmatrix} n-1 \\ k \end{bmatrix} + q^{k-n} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}.$$

Corollary 1.7. *Let n and k be two integers with $k \geq 0$, then the following identity holds:*

$$\begin{bmatrix} n \\ k \end{bmatrix} = \sum_{i=0}^k (-1)^{k-i} q^{(i-k)(n-1)-k} \begin{bmatrix} n+1 \\ i \end{bmatrix}.$$

A version of the next proposition appears in [KC02, page 23] with a different proof only valid for $n \geq 0$ and $m \geq 0$. Hence, we provide a complete proof.

Proposition 1.8. *Let n_1 , n_2 and k be three integers. The following identity holds:*

$$\begin{bmatrix} n_1+n_2 \\ k \end{bmatrix} = \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{n_1 k_2 - n_2 k_1} \begin{bmatrix} n_1 \\ k_1 \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix}.$$

Proof. This statement is proved by induction on n_1 . If $n_1 = 0$ this is obvious. One needs to consider the case $n_1 \geq 0$ and the case $n_1 \leq 0$.

Suppose that $n_1 \geq 0$ and that the statement holds for n_1 . We use Lemma 1.6 to compute:

$$\begin{aligned} & \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{(n_1+1)k_2 - n_2 k_1} \begin{bmatrix} n_1+1 \\ k_1 \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\ &= \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{(n_1+1)k_2 - n_2 k_1} \left(q^{k_1} \begin{bmatrix} n_1 \\ k_1 \end{bmatrix} + q^{k_1 - (n_1+1)} \begin{bmatrix} n_1 \\ k_1-1 \end{bmatrix} \right) \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\ &= q^k \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{n_1 k_2 - n_2 k_1} \begin{bmatrix} n_1 \\ k_1 \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\ &\quad + q^{k - n_1 - n_2 - 1} \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{n_1 k_2 - n_2 (k_1-1)} \begin{bmatrix} n_1 \\ k_1-1 \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\ &= q^k \begin{bmatrix} n_1+n_2 \\ k \end{bmatrix} + q^{k - (n_1+n_2+1)} \begin{bmatrix} n_1+n_2 \\ k-1 \end{bmatrix} \\ &= \begin{bmatrix} n_1+n_2+1 \\ k \end{bmatrix}. \end{aligned}$$

Hence the statement holds for $n_1 + 1$, and therefore for all $n_1 \in \mathbb{Z}_{\geq 0}$.

Suppose that $n_1 < 0$ and that the statement holds for $n_1 + 1$. We use Corollary 1.7 to compute:

$$\begin{aligned}
& \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{n_1 k_2 - n_2 k_1} \begin{bmatrix} n_1 \\ k_1 \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\
&= \sum_{\substack{k_1+k_2=k \\ k_1, k_2 \geq 0}} q^{n_1 k_2 - n_2 k_1} \sum_{i=1}^{k_1} (-1)^{k_1-i} q^{(i-k_1)(n_1-1)-k_1} \begin{bmatrix} n_1+1 \\ i \end{bmatrix} \begin{bmatrix} n_2 \\ k_2 \end{bmatrix} \\
&= \sum_{j=0}^k \sum_{\substack{j_1+j_2=j \\ j_1, j_2 \geq 0}} (-1)^{k-j_1-j_2} q^{(n_1 j_2 - n_2 k + n_2 j_2) + (j-k)(n_1-1) - k + j_2} \begin{bmatrix} n_1+1 \\ j_1 \end{bmatrix} \begin{bmatrix} n_2 \\ j_2 \end{bmatrix} \\
&= \sum_{j=0}^k (-1)^{k-j} q^{(j-k)(n_1-1)-k} \sum_{\substack{j_1+j_2=j \\ j_1, j_2 \geq 0}} q^{(n_1+1)j_2 - n_2 j_1} \begin{bmatrix} n_1+1 \\ j_1 \end{bmatrix} \begin{bmatrix} n_2 \\ j_2 \end{bmatrix} \\
&= \sum_{j=0}^k (-1)^{k-j} q^{(j-k)(n_1+n_2-1)-k} \begin{bmatrix} n_1+n_2+1 \\ j \end{bmatrix} \\
&= \begin{bmatrix} n_1+n_2 \\ k \end{bmatrix}.
\end{aligned}$$

Hence, the statement holds for n_1 and therefore for all negative integers. \square

Definition 1.9. Let X be a set. A *multi-subset* of X is an application $Y : X \rightarrow \mathbb{Z}_{\geq 0}$. If $\sum_{x \in X} Y(x) < \infty$, the multi-subset Y is said to be *finite* and the sum is its *cardinal* (denoted by $\#Y$). If x is an element of X , the number $Y(x)$ is the *multiplicity* of x in Y . An element x of X is *in* Y , if its multiplicity in Y is greater than or equal to 1 (we write $x \in Y$). Let Y_1 and Y_2 be two multi-subsets of a set X .

- The *disjoint union* of Y_1 and Y_2 (denoted $Y_1 \sqcup Y_2$) is the multi-subset $Y_1 + Y_2$.
- The *union* of Y_1 and Y_2 (denoted $Y_1 \cup Y_2$) is the multi-subset $\max(Y_1, Y_2)$.
- The *intersection* of Y_1 and Y_2 (denoted $Y_1 \cap Y_2$) is the multi-subset $\min(Y_1, Y_2)$.

Remark 1.10. A subset Y of X can be thought of as a multi-subset of X by identifying it with its characteristic function.

Notation 1.11. Let X be a set. The power set of X is denoted by $\mathcal{P}(X)$. The set of finite multi-subsets of X is denoted by $\mathcal{M}(X)$. For any non-negative integer k , the set of subsets (resp. multi-subsets) of X of cardinal k is denoted by $\mathcal{P}_k(X)$ (resp. $\mathcal{M}_k(X)$).

Definition 1.12. Let X be an ordered set and Y a multi-subset of X . The *degree* of Y in X is the integer $\deg_X(Y)$ given by the following formula:

$$\deg_X(Y) := \sum_{\substack{x < y \\ x \in X \\ y \in Y \\ x \notin Y}} Y(y) - \sum_{\substack{y < x \\ x \in X \\ y \in Y \\ x \notin Y}} Y(y).$$

Notation 1.13. For N a non-negative integer, denote by $\llbracket N \rrbracket$ the ordered set of integers between 1 and N .

A version of the next proposition can be found in [KC02] (Theorem 6.1 on page 19).

Proposition 1.14. *Let N and k be two non-negative integers. The following identities hold:*

$$\sum_{Y \in \mathcal{P}_k(\llbracket N \rrbracket)} q^{\deg_{\llbracket N \rrbracket}(Y)} = \begin{bmatrix} N \\ k \end{bmatrix} \quad \text{and}$$

$$(-1)^k \sum_{Y \in \mathcal{M}_k(\llbracket N \rrbracket)} q^{\deg_{\llbracket N \rrbracket}(Y)} = \begin{bmatrix} -N \\ k \end{bmatrix}.$$

Proof. This is an easy induction. The statement is clear when $N = 0$, and one can prove that the left-hand sides of these two identities satisfy the induction formulas given in Lemma 1.6 and Corollary 1.7. \square

2. COLORINGS OF MOY GRAPHS

The aim of this section is to present a state sum à la Murakami–Ohtsuki–Yamada [MOY98] computing the Reshetikhin–Turaev invariants of links associated with the exterior powers of the standard representation of $U_q(\mathfrak{gl}_{N|M})$. These invariants are denoted $P_{N|M}$. As in [MOY98], this state sum is a combinatorial interpretation of the underlying representation theory. Details about the algebraic definition is provided in Appendix A. Note however that consistency of the MOY calculus presented here follows from its definition as a state sum and does not rely on the algebraic interpretation. The \mathfrak{sl}_N -invariants associated with exterior powers correspond to setting $M = 0$.

Symmetric powers are covered by this state sum by essentially swapping M and N . Indeed, the Hopf algebra $U_q(\mathfrak{gl}_{M|N})$ is isomorphic to $U_q(\mathfrak{gl}_{N|M})$ (see Remark A.3). The standard representation $\mathbb{C}_q^{M|N}$ of the first is isomorphic to the standard representation $\mathbb{C}_q^{N|M}$ tensored with a trivial representation of odd super degree denoted $\mathbf{1}_{\text{odd}}$. It induces isomorphisms between $\text{Sym}_q^k \mathbb{C}_q^{M|N}$ and $\Lambda^k \left(\mathbb{C}_q^{N|M} \otimes \mathbf{1}_{\text{odd}} \right)$. The latter being isomorphic to

$$\begin{cases} \Lambda_q^k \left(\mathbb{C}_q^{N|M} \right) & k \text{ even,} \\ \Lambda_q^k \left(\mathbb{C}_q^{N|M} \right) \otimes \mathbf{1}_{\text{odd}} & k \text{ odd.} \end{cases}$$

Since the $U_q(\mathfrak{gl}_{N|M})$ -invariant of a link with ℓ components associated with $\mathbf{1}_{\text{odd}}$ is equal to $(-1)^\ell$, we obtain that if L is an oriented link with ℓ components labeled by positive integers k_1, \dots, k_ℓ , interpreted as exponents of symmetric powers, then its $U_q(\mathfrak{gl}_{M|N})$ -invariant is equal to

$$(-1)^{\sum_i k_i} P_{N|M}(\bar{L}),$$

where \bar{L} denote the mirror image of L and the integers k_1, \dots, k_ℓ are now interpreted as exponents exterior powers. The mirror image appears because the isomorphism between $U_q(\mathfrak{gl}_{N|M})$ and $U_q(\mathfrak{gl}_{M|N})$ maps the universal R -matrix of $U_q(\mathfrak{gl}_{N|M})$ onto the inverse of that of $U_q(\mathfrak{gl}_{M|N})$.

The most famous invariant associated with symmetric powers is the colored Jones polynomial of links. For a link L with ℓ components, denote $J_n(L)$ the un-normalized¹

¹This means that the value of the unknot is $[n + 1]$.

colored Jones polynomial associated with the $(n + 1)$ -dimensional irreducible representation of $U_q(\mathfrak{sl}_2)$. The discussion above gives

$$P_{0|2}(L_{n,\dots,n}) = (-1)^{n\ell} J_n(\bar{L}) = J_n(L, -q^{-1}),$$

where $L_{n,\dots,n}$ means that every component of L is labeled by n . For further reference, denote by $J'_n(L) := \frac{J_n(L)}{[n+1]}$ the normalized colored Jones polynomial.

Definition 2.1 ([MOY98]). A *MOY graph* is a finite oriented planar trivalent graph $\Gamma = (V(\Gamma), E(\Gamma))$ with a labeling of its edges $\ell : E_\Gamma \rightarrow \mathbb{Z}_{\geq 0}$ such that the flow given by labels and orientations is preserved at each vertex, meaning that every trivalent vertex follows one of the two models



The first is a *merge* vertex, the second is a *split* vertex. For each vertex, there is a *thick* edge (the one labeled by $a + b$ in the depicted models) and two *thin* edges: the one labeled by a in the previous models is the *left* thin edge, the one labeled by b is the *right* thin edge.

Remark 2.2. Let Γ be a MOY graph, then the graph Γ' obtained by erasing the edges of Γ with label 0 is still a MOY graph. Moreover in all what follows the two graphs can be considered as the same. The label 0 is introduced only to make some definitions and proofs easier and more natural.

Definition 2.3. Let Γ be a MOY graphs. The *cabling* of Γ is the oriented simple multi-curve in the plane \mathbb{R}^2 obtained from Γ by replacing each e edge of Γ by $\ell(e)$ parallel oriented intervals and joining these intervals at every vertex of Γ by the only compatible oriented crossing-less matching.

Let γ be a simple multi-curve in the plane. The *rotational* of γ is the integer $\rho(\gamma)$ equal to the number of circles of γ oriented counterclockwise minus the number of circles of γ oriented clockwise.

A circle oriented counterclockwise is said *positively oriented* and a circle oriented clockwise is said *negatively oriented*.

The *rotational* of a MOY graph Γ is the integer $\rho(\Gamma)$ equal to the rotational of its cabling.

Definition 2.4. The *binomial weight* of a MOY graph Γ is the Laurent polynomial $b(\Gamma)$ defined by the following formula:

$$b(\Gamma) = \prod_{v \text{ split vertex}} \begin{bmatrix} v_l + v_r \\ v_r \end{bmatrix},$$

where for each split vertex v , v_l (resp. v_r) denotes the label of the left thin edge (resp. right thin edge) of v .

Definition 2.5. Let $\Gamma = (V(\Gamma), E(\Gamma), \ell)$ be a MOY graph. A *sub-MOY graph* of Γ is a MOY graph $\Gamma' = (V(\Gamma), E(\Gamma), \ell')$ such that for all e in $E(\Gamma)$, $\ell'(e) \leq \ell(e)$.

A sub-MOY graph Γ' is *cyclic* if for all $e \in E(\Gamma)$, $\ell'(e) \leq 1$.




FIGURE 1. A graph MOY Γ and its cabling. Its rotational $\rho(\Gamma)$ is equal to $2 - 2 = 0$.

Let I be a finite set and consider a collection $\Gamma_i = (V(\Gamma), E(\Gamma), \ell_i)_{i \in I}$ of sub-MOY graphs of Γ , the *sum* of the sub-MOY graphs $(\Gamma_i)_{i \in I}$ is the MOY graph $(V(\Gamma), E(\Gamma), \sum_{i \in I} \ell_i)$ and is denoted by $\sum_{i \in I} \Gamma_i$.

Definition 2.6. Let $\Gamma = (V(\Gamma), E(\Gamma), \ell)$ be a MOY graph and $\Gamma_1 = (V(\Gamma), E(\Gamma), \ell_1)$ and $\Gamma_2 = (V(\Gamma), E(\Gamma), \ell_2)$ be two sub-MOY graphs of Γ . For each split vertex s of Γ , define $u_s(\Gamma_1, \Gamma_2)$ to be the integer defined by the formula:

$$u_s(\Gamma_1, \Gamma_2) := a_2 b_1 - a_1 b_2,$$

where a_1 (resp. b_1) is the label of Γ_1 of the left (resp. right) small edge at the vertex s and a_2 (resp. b_2) is the label of Γ_2 of the left (resp. right) small edge at the vertex s . Define $u(\Gamma_1, \Gamma_2)$ to be the integer given by the formula:

$$u(\Gamma_1, \Gamma_2) := \sum_{s \text{ split vertex of } \Gamma} u_s(\Gamma_1, \Gamma_2).$$

Remark 2.7. Note that the sum of sub-MOY graphs of Γ is a MOY graph but not necessarily a sub-MOY graph of Γ , since it is not required that for all $e \in E(\Gamma)$, $\sum_{i \in I} \ell_i(e) \leq \ell(e)$.

For the rest of the paper, fix two non-negative integers N and M .

Definition 2.8. Let Γ be a MOY graph, an $(N|M)$ -coloring (or simply *coloring*) of Γ is a pair $(\underline{\Gamma}^E, \underline{\Gamma}^S)$ where $\underline{\Gamma}^E = (\Gamma_1^E, \dots, \Gamma_N^E)$ is an N -tuple of cyclic sub-MOY graph of Γ and $\underline{\Gamma}^S = (\Gamma_1^S, \dots, \Gamma_M^S)$ is an M -tuple of sub-MOY graph of Γ such that:

$$\sum_{j=1}^N \Gamma_j^E + \sum_{i=1}^M \Gamma_i^S = \Gamma.$$

The set of $(N|M)$ -colorings of Γ is denoted by $\text{col}_{N|M}(\Gamma)$ (or simply $\text{col}(\Gamma)$).

Remark 2.9. Let Γ be a MOY graph and $c = (\underline{\Gamma}^E, \underline{\Gamma}^S)$ a coloring of Γ . If e is an edge of Γ , consider the pair $(c^E(e), c^S(e))$: $c^E(e)$ is the subset of $\llbracket N \rrbracket$ which contains j if and only if $\ell_{\Gamma_j^E}(e) = 1$ and $c^S(e)$ is the sub-multiset of $\llbracket M \rrbracket$ where i appears with multiplicity $\ell_{\Gamma_i^S}(e)$. We have $\#c^E(e) + \#c^S(e) = \ell(e)$. A coloring can be defined by such a collection $(c^E(e), c^S(e))_{e \in E(\Gamma)}$ satisfying a flow condition at every vertex.

Definition 2.10. Let Γ be a MOY graph and $c = (\underline{\Gamma}^E, \underline{\Gamma}^S)$ an $(N|M)$ -coloring of Γ . The *s-weight* of c is the integer denoted by $w_s(c)$ and defined by the following formula:

$$w_s(c) := \sum_{1 \leq i < j \leq M} u(\Gamma_j^S, \Gamma_i^S) - \sum_{1 \leq i < j \leq N} u(\Gamma_j^E, \Gamma_i^E) + \sum_{i=1}^N \sum_{j=1}^M u(\Gamma_i^E, \Gamma_j^S).$$

The *ρ -weight* of c is the integer denoted by $w_\rho(c)$ and defined by the following formula:

$$w_\rho(c) := \sum_{i=1}^N (N + M - 2i + 1) \rho(\Gamma_i^E) + \sum_{j=1}^M (N + M - 2j + 1) \rho(\Gamma_j^S).$$

The *multiplicity* of c is the Laurent polynomial $m(c)$ with positive coefficients and symmetric in q and q^{-1} given by the following formula:

$$m(c) = \prod_{i=1}^M b(\Gamma_i^S).$$

The *parity* of c is the integer $s(c)$ given by the following formula:

$$s(c) = \sum_{i=1}^M \rho(\Gamma_i^S),$$

as the name suggests, we are only interested in the value of $s(c)$ modulo 2. Equality modulo 2 is denoted by \equiv in what follows. The *combinatorial $(N|M)$ -evaluation of the coloring c* is the Laurent polynomial $\langle \Gamma, c \rangle_{N|M}$ given by the following formula:

$$\langle \Gamma, c \rangle_{N|M} = (-1)^{s(c)} q^{w_s(c) + w_\rho(c)} m(c).$$

Finally, the *combinatorial $(N|M)$ -evaluation of Γ* is the Laurent polynomial $\langle \Gamma \rangle_{N|M}$ given by the following formula:

$$\langle \Gamma \rangle_{N|M} = \sum_{c \in \text{col}(\Gamma)} \langle \Gamma, c \rangle_{N|M}.$$

We give in Appendix A a detailed account of the Reshetikhin–Turaev invariant for MOY graphs for the quantum group $U_q(\mathfrak{gl}_{N|M})$. For any MOY graph Γ it is denoted $\langle\langle \Gamma \rangle\rangle_{N|M}$. The aim of the rest of this section is to prove the next theorem.

Theorem 2.11. *For any MOY graph Γ , the algebraic and the combinatorial evaluation of Γ agree. This means:*

$$\langle\langle \Gamma \rangle\rangle_{N|M} = \langle \Gamma \rangle_{N|M}.$$

In particular, the combinatorial evaluation depends only on $N - M$ and is symmetric in q and q^{-1} .

Following Proposition A.10, one only needs to check that $\langle \bullet \rangle_{N|M}$ satisfies the multiplicativity property and the local relations of Proposition A.9. This is the object of the rest of this section.

Lemma 2.12. *Let Γ be a MOY graph, and $\bar{\Gamma}$ be its image under the transformation $\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} -x \\ y \end{pmatrix}$. Then $\langle \Gamma \rangle_{N|M}(q) = \langle \bar{\Gamma} \rangle_{N|M}(q^{-1})$.*

Proof. There is a canonical one-to-one correspondence between colorings of Γ and colorings of $\bar{\Gamma}$. Let c be a coloring of Γ and \bar{c} the corresponding coloring of $\bar{\Gamma}$. The following identities hold:

$$m(c)(q) = m(c)(q^{-1}) = m(\bar{c})(q) = m(\bar{c})(q^{-1}),$$

$$\begin{aligned} s(c) &\equiv s(\bar{c}), \\ w_s(c) &= -w_s(\bar{c}) \quad \text{and} \\ w_\rho(c) &= -w_\rho(\bar{c}). \end{aligned}$$

From this, one immediately obtains $\langle \Gamma \rangle_{N|M}(q) = \langle \bar{\Gamma} \rangle_{N|M}(q^{-1})$. □

Lemma 2.13. *Let Γ and Υ be two MOY graphs, then*

$$\langle \Gamma \sqcup \Upsilon \rangle_{N|M} = \langle \Gamma \rangle_{N|M} \langle \Upsilon \rangle_{N|M}.$$

Proof. First of all, there is a canonical one-to-one correspondence between the set of colorings of $\Gamma \sqcup \Upsilon$ and the Cartesian product of sets of colorings of Γ and Υ . If c and d are colorings of Γ and Υ , denote by (c, d) the corresponding coloring of $\Gamma \sqcup \Upsilon$. It is enough to prove that for all colorings (c, d) ,

$$(-1)^{s(c)+s(d)} q^{w_s(c)+w_\rho(d)+w_s(c)+w_\rho(d)} m(c)m(d) = (-1)^{s((c,d))} q^{w_s((c,d))+w_\rho((c,d))} m((c,d)).$$

Let $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ (resp. $d = (\underline{\Upsilon^E}, \underline{\Upsilon^S})$) be a coloring of Γ (resp. Υ). One has $s((c, d)) = s(c) + s(d)$ and $m((c, d)) = m(c)m(d)$. Hence, it remains to prove that $w((c, d)) = w(c) + w(d)$. Let us denote by Φ the MOY graph $\Gamma \sqcup \Upsilon$. One has

$$\begin{aligned} w_s((c, d)) &= \sum_{1 \leq i < j \leq M} u(\Phi_j^S, \Phi_i^S) - \sum_{1 \leq i < j \leq N} u(\Phi_j^E, \Phi_i^E) + \sum_{i=1}^N \sum_{j=1}^M u(\Phi_i^E, \Phi_j^S) \\ &= \sum_{1 \leq i < j \leq M} u(\Gamma_j^S, \Gamma_i^S) + u(\Upsilon_j^S, \Upsilon_i^S) - \sum_{1 \leq i < j \leq N} u(\Gamma_j^E, \Gamma_i^E) + u(\Upsilon_j^E, \Upsilon_i^E) \\ &\quad + \sum_{i=1}^N \sum_{j=1}^M u(\Gamma_i^E, \Gamma_j^S) + u(\Upsilon_i^E, \Upsilon_j^S). \\ &= w_s(c) + w_s(d) \end{aligned}$$

and

$$\begin{aligned} w_\rho((c, d)) &= \sum_{i=1}^M (N + M - 2i + 1) \rho(\Phi_i^S) + \sum_{i=1}^N (N + M - 2i + 1) \rho(\Phi_i^E) \\ &= \sum_{i=1}^M (N + M - 2i + 1) (\rho(\Gamma_i^S) + \rho(\Upsilon_i^S)) \\ &\quad + \sum_{i=1}^N (N + M - 2i + 1) (\rho(\Gamma_i^E) + \rho(\Upsilon_i^E)) \\ &= w_\rho(c) + w_\rho(d). \end{aligned} \quad \square$$

Lemma 2.14. *For any non-negative integer k , the following identity holds:*

$$\left\langle \left\langle \bigcirc \right\rangle^k \right\rangle_{N|M} = \begin{bmatrix} N - M \\ k \end{bmatrix} = \left\langle \left\langle \bigcirc \right\rangle^k \right\rangle_{N|M}.$$

Proof. Let us start with the first identity. Denote by Γ the positively oriented circle of label k . A coloring $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ of Γ is completely encoded by a pair consisting of a subset X_c of $\llbracket N \rrbracket$ with k_c^E elements a multi-subset Y_c of $\llbracket M \rrbracket$ with k_c^S elements such that $k_c^E + k_c^S = k$. One has $s(c) = k_c^S$ and $m(c) = 1$. Since there is no vertex, $w_s(\Gamma) = 0$. One has

$$w_\rho(c) = \sum_{i=1}^M (N + M - 2i + 1) \rho(\Gamma_i^S) + \sum_{i=1}^N (N + M - 2i + 1) \rho(\Gamma_i^E)$$

Proof. Let Γ and Y be two MOY graphs which are identical except in a small ball where they are related by the local relation (3), Γ being on the left-hand side and Y on the right-hand side. First note that any coloring of Γ induces a coloring of Y . Let us fix a coloring $c' = (\underline{Y}^E, \underline{Y}^S)$ of Y . If a coloring c of Γ induces c' on Y , we write $c \rightarrow c'$. We will prove the following:

$$\sum_{\substack{c \rightarrow c' \\ c \in \text{col}(\Gamma)}} \langle \Gamma, c \rangle_{N|M} = [m+1] \langle Y, c' \rangle_{N|M},$$

which implies $\langle \Gamma \rangle_{N|M} = [m+1] \langle Y \rangle_{N|M}$. Denote by e the edge of Y which appears in the local relation. For i in $\llbracket M \rrbracket$ and j in $\llbracket N \rrbracket$, set

$$k_i^S := \ell_{Y^S}(e), \quad k_j^E := \ell_{Y^E}(e), \quad k^S := \sum_{i=1}^M k_i^S, \quad \text{and} \quad k^E := \sum_{j=1}^N k_j^E.$$

A coloring $c = (\underline{\Gamma}^E, \underline{\Gamma}^S)$ of Γ inducing c' is totally determined by saying which MOY graph of the collection $\underline{\Gamma}^E \sqcup \underline{\Gamma}^S$ gives label 1 to the right edge of the digon which appears in the local relation. Denote this graph by Γ_h^A . There are two possibilities: either this graph belongs to $\underline{\Gamma}^S$ (i. e. $A = S$ and $h \in \llbracket M \rrbracket$) or to $\underline{\Gamma}^E$ (i. e. $A = E$ and $h \in \llbracket N \rrbracket$). If $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$\begin{aligned} m(c) &= m(c') [k_h^S], & s(c) &= s(c'), \\ w_\rho(c) &= w_\rho(c') & \text{and} & \quad w_s(c) = w_s(c') - \sum_{i=1}^{h-1} k_i^S + \sum_{i=h+1}^M k_i^S - k^E. \end{aligned}$$

If $A = E$ and $h \in \llbracket N \rrbracket$, one has:

$$\begin{aligned} m(c) &= m(c') [k_h^E], \\ s(c) &= s(c'), \\ w_\rho(c) &= w_\rho(c') \quad \text{and} \\ w_s(c) &= w_s(c') + \sum_{i=1}^{h-1} k_i^E - \sum_{i=h+1}^N k_i^E + k^S. \end{aligned}$$

Note that for the coloring c to exist, it is necessary that $k_h^E = 1$. However, if $k_h^E = 0$, the previous formula makes the contribution of this “virtual” coloring equal to 0. This argument will be used throughout the proofs of this section. We will not repeat it. One can sum:

$$\begin{aligned} \sum_{\substack{c \rightarrow c' \\ c \in \text{col}(\Gamma)}} \langle \Gamma, c \rangle_{N|M} &= \left(\sum_{h=1}^M q^{-k^E - \sum_{i=1}^{h-1} k_i^S + \sum_{i=h+1}^M k_i^S} [k_h^S] \right. \\ &\quad \left. + \sum_{h=1}^N q^{k^S + \sum_{i=1}^{h-1} k_i^E - \sum_{i=h+1}^N k_i^E} [k_h^E] \right) \langle Y, c' \rangle_{N|M} \\ &= \left(q^{-k^E} [k^S] + q^{k^S} [k^E] \right) \langle Y, c' \rangle_{N|M} \\ &= [k^S + k^E] \langle Y, c' \rangle_{N|M} = [m+1] \langle Y, c' \rangle_{N|M}. \quad \square \end{aligned}$$

Corollary 2.17. *The combinatorial $(N|M)$ -evaluation satisfies the following local identities and their mirror images:*

$$\begin{aligned}
\left\langle \begin{array}{c} \begin{array}{c} \uparrow \\ m+1 \\ \uparrow \end{array} \\ \left(\begin{array}{c} \begin{array}{c} \uparrow \\ m+1 \\ \uparrow \end{array} \\ 1 \\ \begin{array}{c} \uparrow \\ m \\ \uparrow \end{array} \end{array} \right) \\ \begin{array}{c} \uparrow \\ m+1 \\ \uparrow \end{array} \end{array} \right\rangle_{N|M} &= [m+1] \left\langle \begin{array}{c} \uparrow \\ m+1 \\ \uparrow \end{array} \right\rangle_{N|M}, \\
\left\langle \begin{array}{c} \begin{array}{c} \uparrow \\ m+n \\ \uparrow \end{array} \\ \left(\begin{array}{c} \begin{array}{c} \uparrow \\ m+n \\ \uparrow \end{array} \\ m \\ \begin{array}{c} \uparrow \\ n \\ \uparrow \end{array} \end{array} \right) \\ \begin{array}{c} \uparrow \\ m+n \\ \uparrow \end{array} \end{array} \right\rangle_{N|M} &= \begin{bmatrix} m+n \\ m \end{bmatrix} \left\langle \begin{array}{c} \uparrow \\ m+n \\ \uparrow \end{array} \right\rangle_{N|M}, \\
\left\langle \begin{array}{c} \begin{array}{c} \uparrow \\ k-r \\ \uparrow \end{array} \\ \left(\begin{array}{c} \begin{array}{c} \uparrow \\ k \\ \uparrow \end{array} \\ \begin{array}{c} \leftarrow r \\ \rightarrow l \end{array} \\ \begin{array}{c} \uparrow \\ s \\ \uparrow \end{array} \end{array} \right) \\ \begin{array}{c} \uparrow \\ k+s \\ \uparrow \end{array} \end{array} \right\rangle_{N|M} &= \begin{bmatrix} r+s \\ r \end{bmatrix} \left\langle \begin{array}{c} \begin{array}{c} \uparrow \\ k-r \\ \uparrow \end{array} \\ \begin{array}{c} \leftarrow r+s \\ \rightarrow l+r \end{array} \\ \begin{array}{c} \uparrow \\ l-s \\ \uparrow \end{array} \end{array} \right\rangle_{N|M}.
\end{aligned}$$

Proof. The first identity is a consequence of Lemma 2.16 and Lemma 2.12. The second is an easy induction. The third is a consequence of the second and of Lemma 2.15. \square

Lemma 2.18. *The combinatorial $(N|M)$ -evaluation satisfies the following local identity:*

$$(4) \quad \left\langle \begin{array}{c} \begin{array}{c} \uparrow \\ m \\ \uparrow \end{array} \\ \left(\begin{array}{c} \begin{array}{c} \uparrow \\ m+1 \\ \uparrow \end{array} \\ 1 \\ \begin{array}{c} \uparrow \\ m \\ \uparrow \end{array} \end{array} \right) \\ \begin{array}{c} \uparrow \\ m \\ \uparrow \end{array} \end{array} \right\rangle_{N|M} = [N-M-m] \left\langle \begin{array}{c} \uparrow \\ m \\ \uparrow \end{array} \right\rangle_{N|M}.$$

Proof. Let Γ and Υ be two MOY graphs which are identical except in a small ball where they are related by the local relation (4), Γ being on the left-hand side and Υ on the right-hand side. First note that any coloring of Γ induces a coloring of Υ . Fix a coloring $c' = (\underline{\Upsilon}^E, \underline{\Upsilon}^S)$ of Υ . If a coloring c of Γ induces c' on Υ , we write $c \rightarrow c'$. We will prove the following:

$$\sum_{\substack{c \rightarrow c' \\ c \in \text{col}(\Gamma)}} \langle \Gamma, c \rangle_{N|M} = [N-M-m] \langle \Upsilon, c' \rangle_{N|M},$$

which implies $\langle \Gamma \rangle_{N|M} = [N-M-m] \langle \Upsilon \rangle_{N|M}$. Denote by e the edge of Υ which appears in the local relation. For i in $\llbracket M \rrbracket$ and j in $\llbracket N \rrbracket$, set

$$k_i^S := \ell_{\Upsilon_i^S}(e), \quad k_j^E := \ell_{\Upsilon_j^E}(e), \quad k^S := \sum_{i=1}^M k_i^S, \quad \text{and} \quad k^E := \sum_{i=1}^N k_i^E.$$

A coloring $c = (\underline{\Gamma}^S, \underline{\Gamma}^E)$ of Γ inducing c' is totally determined by saying which MOY graph of the collection $\underline{\Gamma}^E \sqcup \underline{\Gamma}^S$ gives label 1 to the right edge of the digon which appears in the local relation. Denote this graph by Γ_h^A . There are two possibilities: either this graph belongs to $\underline{\Gamma}^S$ (i. e. $A = S$ and $h \in \llbracket M \rrbracket$) or to $\underline{\Gamma}^E$ (i. e. $A = E$ and $h \in \llbracket N \rrbracket$). If $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$m(c) = m(c')[k_h^S + 1],$$

$$s(c) = s(c') + 1,$$

$$w_\rho(c) = w_\rho(c') + N + M - 2h + 1 \quad \text{and}$$

$$w_s(c) = w_s(c') - \sum_{i=1}^{h-1} k_i^S + \sum_{i=h+1}^M k_i^S - k^E.$$

Hence:

$$w_\rho(c) + w_s(c) = w_\rho(c') + w_s(c') + N - k^E - \sum_{i=1}^{h-1} (k_i^S + 1) + \sum_{i=h+1}^M (k_i^S + 1).$$

If $A = E$ and $h \in \llbracket N \rrbracket$, one has:

$$m(c) = m(c')[1 - k_h^E] \quad \text{note that one has } k_h^E = 0,$$

$$s(c) = s(c'),$$

$$w_\rho(c) = w_\rho(c') + N + M - 2h + 1 \quad \text{and}$$

$$w_s(c) = w_s(c') + \sum_{i=1}^{h-1} k_i^E - \sum_{i=h+1}^N k_i^E + k^S.$$

Hence:

$$w_\rho(c) + w_s(c) = w_\rho(c') + w_s(c') + M + k^S - \sum_{i=1}^{h-1} (1 - k_i^E) + \sum_{i=h+1}^N (1 - k_i^E).$$

Finally, one gets:

$$\begin{aligned} \sum_{\substack{c \rightarrow c' \\ c \in \text{col}(\Gamma)}} \langle \Gamma, c \rangle_{N|M} &= \left(- \sum_{h=1}^M q^{N - k^E - \sum_{i=1}^{h-1} (k_i^S + 1) + \sum_{i=h+1}^M (k_i^S + 1)} [k_h^S + 1] \right. \\ &\quad \left. + \sum_{h=1}^N q^{M + k^S - \sum_{i=1}^{h-1} (1 - k_i^E) + \sum_{i=h+1}^N (1 - k_i^E)} [1 - k_h^E] \right) \langle \Upsilon, c' \rangle \\ &= \left(-q^{N - k^E} [M + k^S] + q^{M + k^S} [N - k^E] \right) \langle \Upsilon, c' \rangle \\ &= [N - M - m] \langle \Upsilon, c' \rangle. \end{aligned}$$

□

Corollary 2.19. *The combinatorial $(N|M)$ -evaluation satisfies the following local identities:*

$$\begin{aligned} \left\langle \begin{array}{c} m \uparrow \\ 1 \circlearrowleft \\ m \downarrow \end{array} \right\rangle_{N|M} &= [N - M - m] \left\langle \begin{array}{c} m \uparrow \\ \circlearrowleft \\ m \downarrow \end{array} \right\rangle_{N|M}, \\ \left\langle \begin{array}{c} m \uparrow \\ m+n \circlearrowleft \\ n \downarrow \end{array} \right\rangle_{N|M} &= \begin{bmatrix} N - M - m \\ n \end{bmatrix} \left\langle \begin{array}{c} m \uparrow \\ \circlearrowleft \\ m \downarrow \end{array} \right\rangle_{N|M} = \left\langle \begin{array}{c} m \uparrow \\ n \circlearrowleft \\ m+n \downarrow \end{array} \right\rangle_{N|M}. \end{aligned}$$

Proof. The first identity is a consequence of Lemma 2.18 and Lemma 2.12. The second is an easy induction. □

Lemma 2.20. *The combinatorial $(N|M)$ -evaluation satisfies the following local identity:*

(5)

$$\left\langle \begin{array}{c} n \\ \uparrow \\ n+1 \\ \leftarrow 1 \\ \uparrow \\ n \\ \leftarrow 1 \\ \uparrow \\ n+k \end{array} \right\rangle_{N|M} = [k] \left\langle \begin{array}{c} n \\ \uparrow \\ n+k \end{array} \right\rangle_{N|M} + \left\langle \begin{array}{c} n \\ \uparrow \\ n-1 \\ \leftarrow 1 \\ \uparrow \\ n \\ \leftarrow 1 \\ \uparrow \\ n+k \end{array} \right\rangle_{N|M} .$$

Proof. Let Γ , Υ and Φ be three MOY graphs which are identical except in a small ball B where they are related by the local relation (5), Γ being on the left-hand side and Υ and Φ on the right-hand side (Υ the first and Φ the second). Let c be a coloring of Γ . There are two possibilities: either the two rungs receive the same color or they receive different colors. First consider a coloring $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ which gives different colors to the two rungs. There is no coloring of Υ which coincides with c outside B . On the other hand, it induces a unique coloring $c'' = (\underline{\Phi^E}, \underline{\Phi^S})$ of Φ which associates different colors to the two rungs of Φ . One has:

$$\begin{aligned} m(c) &= m(c''), & s(c) &= s(c''), \\ w_\rho(c) &= w_\rho(c'') & \text{and} & & w_s(c) &= w_s(c''). \end{aligned}$$

Hence

$$(6) \quad \langle \Gamma, c \rangle_{N|M} = \langle \Phi, c'' \rangle_{N|M}$$

Consider now the case where c gives the same color to the two rungs. It induces a coloring c' of Υ . Similarly there are colorings of Φ which induce c' on Υ . Let us write $c \rightarrow c'$ to indicate that c induces the coloring c' on Υ . We will show the following:

$$\sum_{\substack{c \in \text{col}(\Gamma) \\ c \rightarrow c'}} \langle \Gamma, c \rangle_{N|M} - \sum_{\substack{c \in \text{col}(\Phi) \\ c \rightarrow c'}} \langle \Phi, c'' \rangle_{N|M} = [k] \langle \Upsilon, c' \rangle_{N|M}$$

which together with (6) implies the lemma.

Fix $c' = (\underline{\Upsilon^E}, \underline{\Upsilon^S})$ a coloring of Υ . Let us denote by e_l (resp. e_r) the vertical edge on the left (resp. right) of the part of Υ which is in B . We need a few extra notations. For $i \in \llbracket M \rrbracket$ and $j \in \llbracket N \rrbracket$, set:

$$\begin{aligned} l_i^S &:= \ell_{\Gamma_i^S}(e_l), & r_i^S &:= \ell_{\Gamma_i^S}(e_r), & l_j^E &:= \ell_{\Gamma_j^E}(e_l), & r_j^E &:= \ell_{\Gamma_j^E}(e_r), \\ l^S &:= \sum_{i=1}^M l_i^S, & r^S &:= \sum_{i=1}^M r_i^S, & l^E &:= \sum_{j=1}^N l_j^E, & r^E &:= \sum_{j=1}^N r_j^E. \end{aligned}$$

Note that $k = r^E + r^S - l^E - l^S$. A coloring of Γ which induces c' is totally determined by the color it gives to the two rungs. Let $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ and denote by Γ_h^A the sub-MOY graph of Γ which has label 1 on the two rungs. There are two possibilities: either $A = S$ and $h \in \llbracket M \rrbracket$ or $A = E$ and $h \in \llbracket N \rrbracket$. In the case $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$\begin{aligned} m(c) &= m(c') [l_h^S + 1] [r_h^S] \\ s(c) &= s(c'), \\ w_\rho(c) &= w_\rho(c') \quad \text{and} \end{aligned}$$

$$w_s(c) = w_s(c') + r^E - l^E + \sum_{i=1}^{h-1} (l_i^S - r_i^S) - \sum_{i=h+1}^M (l_i^S - r_i^S).$$

In the case $A = E$ and $h \in \llbracket N \rrbracket$, one has:

$$m(c) = m(c')[1 - l_h^E][r_h^E]$$

$$s(c) = s(c'),$$

$$w_\rho(c) = w_\rho(c') \quad \text{and}$$

$$w_s(c) = w_s(c') + l^S - r^S + \sum_{i=1}^{h-1} (r_i^E - l_i^E) - \sum_{i=h+1}^N (r_i^E - l_i^E).$$

Note that in order c to exist it is necessary and sufficient that $r_h^E = 1$ and $l_h^E = 0$.

We perform the same analysis with a coloring $c'' = (\underline{\Phi}^E, \underline{\Phi}^S)$ inducing c' on Υ . Denote by Φ_h^A the sub-MOY graph of Φ which has label 1 on the two rungs. There are two possibilities: either $A = S$ and $h \in \llbracket M \rrbracket$ or $A = E$ and $h \in \llbracket N \rrbracket$. In the case $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$m(c'') = m(c')[l_h^S][r_h^S + 1]$$

$$s(c'') = s(c'),$$

$$w_\rho(c'') = w_\rho(c') \quad \text{and}$$

$$w_s(c'') = w_s(c') + r^E - l^E + \sum_{i=1}^{h-1} (l_i^S - r_i^S) - \sum_{i=h+1}^M (l_i^S - r_i^S).$$

In the case $A = E$ and $h \in \llbracket N \rrbracket$, one has:

$$m(c'') = m(c')[l_h^E][1 - r_h^S]$$

$$s(c'') = s(c'),$$

$$w_\rho(c'') = w_\rho(c') \quad \text{and}$$

$$w_s(c'') = w_s(c') + l^S - r^S + \sum_{i=1}^{h-1} (r_i^E - l_i^E) - \sum_{i=1}^N (r_i^E - l_i^E).$$

We compute with help of Lemma 1.5:

$$\begin{aligned} & \sum_{\substack{c \in \text{col}(\Gamma) \\ c \rightarrow c'}} \langle \Gamma, c \rangle_{N|M} - \sum_{\substack{c'' \in \text{col}(\Phi) \\ c'' \rightarrow c'}} \langle \Phi, c'' \rangle_{N|M} \\ &= \left(q^{r^E - l^E} \sum_{h=1}^M ([l_h^S + 1][r_h^S] - [l_h^S][r_h^S + 1]) q^{\sum_{i=1}^{h-1} (l_i^S - r_i^S) - \sum_{i=h+1}^M (l_i^S - r_i^S)} \right. \\ & \quad \left. + q^{l^S - r^S} \sum_{h=1}^N ([1 - l_h^E][r_h^E] - [l_h^E][1 - r_h^E]) q^{\sum_{i=1}^{h-1} (r_i^E - l_i^E) - \sum_{i=h+1}^N (r_i^E - l_i^E)} \right) \langle \Upsilon, c' \rangle_{N|M} \\ &= \left(q^{r^E - l^E} \sum_{h=1}^M [r_h^S - l_h^S] q^{\sum_{i=1}^{h-1} (l_i^S - r_i^S) - \sum_{i=h+1}^M (l_i^S - r_i^S)} \right. \\ & \quad \left. + q^{l^S - r^S} \sum_{h=1}^N [r_h^E - l_h^E] q^{\sum_{i=1}^{h-1} (r_i^E - l_i^E) - \sum_{i=h+1}^N (r_i^E - l_i^E)} \right) \langle \Upsilon, c' \rangle_{N|M} \\ &= (q^{r^E - l^E} [r^S - l^S] + q^{l^S - r^S} [r^E - l^E]) \langle \Upsilon, c' \rangle_{N|M} \\ &= [k] \langle \Upsilon, c' \rangle_{N|M}. \end{aligned}$$

□

Corollary 2.21. *The combinatorial $(N|M)$ -evaluation satisfies the following local identity and its mirror image:*

$$\left\langle \begin{array}{c} m \quad n+l \\ \leftarrow \quad \leftarrow \\ n+k \quad m+l-k \\ \rightarrow \quad \rightarrow \\ n \quad m+l \end{array} \right\rangle = \sum_{j=\max(0, m-n)}^l \begin{bmatrix} l \\ k-j \end{bmatrix} \left\langle \begin{array}{c} m \quad n+l \\ \leftarrow \quad \leftarrow \\ m-j \quad n+l+j \\ \rightarrow \quad \rightarrow \\ n \quad m+l \end{array} \right\rangle.$$

Proof. This is an easy induction using Lemma 2.20 and Corollary 2.17. \square

Lemma 2.22. *The combinatorial $(N|M)$ -evaluation satisfies the following local identity:*

(7)

$$\left\langle \begin{array}{c} m \quad m \\ \leftarrow \quad \leftarrow \\ m+1 \quad m+1 \\ \rightarrow \quad \rightarrow \\ m \quad m \end{array} \right\rangle_{N|M} = [N - M - 2m] \left\langle \begin{array}{c} \uparrow \\ m \\ \downarrow \end{array} \right\rangle_{N|M} + \left\langle \begin{array}{c} m \quad m \\ \leftarrow \quad \leftarrow \\ m-1 \quad m-1 \\ \rightarrow \quad \rightarrow \\ m \quad m \end{array} \right\rangle_{N|M}.$$

Proof. Let Γ , Υ and Φ be three MOY graphs which are identical except in a small ball B where they are related by the local relation (7), Γ being on the left-hand side and Υ and Φ on the right-hand side (Υ the first and Φ the second). Let c be a coloring of Γ . There are two possibilities: either the two rungs receive the different colors or they receive different colors.

First, consider a coloring $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ which gives different colors to the two rungs. There is no coloring of Υ which coincides with c outside B . On the other hand, it induces a unique coloring $c' = (\underline{\Phi^E}, \underline{\Phi^S})$ of Φ which associates different colors to the two rungs of Φ . One has:

$$\begin{aligned} m(c) &= m(c'), & s(c) &= s(c'), \\ w_\rho(c) &= w_\rho(c') & \text{and} & & w_s(c) &= w_s(c'). \end{aligned}$$

Hence

$$(8) \quad \langle \Gamma, c \rangle_{N|M} = (-1)^{s(c)} q^{w_\rho(c) + w_s(c)} m(c) = (-1)^{s(c')} q^{w_\rho(c') + w_s(c')} m(c') = \langle \Phi, c' \rangle_{N|M}.$$

Consider now the case where c gives the same color to the two rungs. It induces a coloring c' of Υ . Similarly there are colorings of Φ which induce c' on Υ . Let us write $c \rightarrow c'$ to indicate that c induces the coloring c' on Υ . We will show the following:

$$\sum_{\substack{c \in \text{col}(\Gamma) \\ c \rightarrow c'}} \langle \Gamma, c \rangle_{N|M} - \sum_{\substack{c'' \in \text{col}(\Phi) \\ c'' \rightarrow c'}} \langle \Phi, c'' \rangle_{N|M} = [N - M - 2m] \langle \Upsilon, c' \rangle_{N|M}$$

which together with (8) implies the lemma.

Let us fix a coloring $c' = (\underline{\Upsilon^E}, \underline{\Upsilon^S})$ of Υ . and denote by e_l (resp. e_r) the vertical edge on the left (resp. right) of the part of Υ which is in B . We need a few extra notations. For $i \in \llbracket M \rrbracket$ and $j \in \llbracket N \rrbracket$, set:

$$\begin{aligned} l_i^S &:= \ell_{\Gamma_i^S}(e_l), & r_i^S &:= \ell_{\Gamma_i^S}(e_r), & l_j^E &:= \ell_{\Gamma_j^E}(e_l), & r_j^E &:= \ell_{\Gamma_j^E}(e_r), \\ l^S &:= \sum_{i=1}^M l_i^S, & r^S &:= \sum_{i=1}^M r_i^S, & l^E &:= \sum_{j=1}^N l_j^E, & r^E &:= \sum_{j=1}^N r_j^E. \end{aligned}$$

A coloring of Γ inducing c' is totally determined by the color it gives to the two rungs. Let $c = (\underline{\Gamma^E}, \underline{\Gamma^S})$ and denote by Γ_h^A the sub-MOY graph of Γ which has label 1 on the two rungs. There are two possibilities: either $A = S$ and $h \in \llbracket M \rrbracket$ or $A = E$ and $h \in \llbracket N \rrbracket$. In the case $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$\begin{aligned} m(c) &= m(c')[l_h^S + 1][r_h^S + 1] \\ s(c) &= s(c') - 1, \\ w_\rho(c) &= w_\rho(c') + (N + M - 2h + 1) \quad \text{and} \\ w_s(c) &= w_s(c') - r^E - l^E - \sum_{i=1}^{h-1} (l_i^S + r_i^S) + \sum_{i=h+1}^M (l_i^S + r_i^S). \end{aligned}$$

In the case $A = E$ and $h \in \llbracket N \rrbracket$, one has:

$$\begin{aligned} m(c) &= m(c')[1 - l_h^E][1 - r_h^E] \\ s(c) &= s(c'), \\ w_\rho(c) &= w_\rho(c') + (N + M - 2h + 1) \quad \text{and} \\ w_s(c) &= w_s(c') + l^S + r^S + \sum_{i=1}^{h-1} (r_i^E + l_i^E) - \sum_{i=1}^N (r_i^E + l_i^E). \end{aligned}$$

Note that in order c to exist it is necessary and sufficient that $r_h^E = l_h^E = 1$.

We perform the same analysis with a coloring $c'' = (\underline{\Phi^E}, \underline{\Phi^S})$ inducing c' on Υ . Denote by Φ_h^A the sub-MOY graph of Φ which has label 1 on the two rungs. There are two possibilities: either $A = S$ and $h \in \llbracket M \rrbracket$ or $A = E$ and $h \in \llbracket N \rrbracket$. In the case $A = S$ and $h \in \llbracket M \rrbracket$, one has:

$$\begin{aligned} m(c'') &= m(c')[l_h^S][r_h^S] \\ s(c'') &\equiv s(c') + 1, \\ w_\rho(c'') &= w_\rho(c') + (N + M - 2h + 1) \quad \text{and} \\ w_s(c'') &= w_s(c') - r^E - l^E - \sum_{i=1}^{h-1} (l_i^S + r_i^S) + \sum_{i=h+1}^M (l_i^S + r_i^S). \end{aligned}$$

Note that in order c'' to exist it is necessary and sufficient for r_h^S and l_h^S to be positive.

In the case $A = E$ and $h \in \llbracket N \rrbracket$, we have:

$$\begin{aligned} m(c'') &= m(c')[l_h^E][r_h^E] \\ s(c'') &\equiv s(c'), \\ w_\rho(c'') &= w_\rho(c') + (N + M - 2h + 1) \quad \text{and} \\ w_s(c'') &= w_s(c') + l^S + r^S + \sum_{i=1}^{h-1} (r_i^E + l_i^E) - \sum_{i=1}^N (r_i^E + l_i^E). \end{aligned}$$

Hence, one has:

$$\begin{aligned} & \sum_{\substack{c \in \text{col}(\Gamma) \\ c \rightarrow c'}} \langle \Gamma, c \rangle_{N|M} - \sum_{\substack{c \in \text{col}(\Phi) \\ c'' \rightarrow c'}} \langle \Phi, c'' \rangle_{N|M} \\ &= \left(\sum_{i=1}^M \left([l_h^S][r_h^S] - [l_h^S + 1][r_h^S + 1] \right) q^{N+M-2h+1-r^E-l^E-\sum_{i=1}^{h-1}(l_i^S+r_i^S)+\sum_{i=h+1}^M(l_i^S+r_i^S)} \right. \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=1}^N \left([1 - l_h^E][1 - r_h^E] - [l_h^E][r_h^E] \right) q^{N+M-2h+1+r^S+l^S+\sum_{i=1}^{h-1}(l_i^E+r_i^E)-\sum_{i=h+1}^N(l_i^E+r_i^E)} \left\langle \Upsilon, c' \right\rangle_{N|M} \\
& = \left(\sum_{i=1}^M \left(-[l_h^S + r_h^S + 1] \right) q^{N+M-2h+1-r^E-l^E-\sum_{i=1}^{h-1}(l_i^S+r_i^S)+\sum_{i=h+1}^M(l_i^S+r_i^S)} \right. \\
& \quad \left. + \sum_{i=1}^N \left([1 - l_h^E - r_h^E] \right) q^{N+M-2h+1+r^S+l^S+\sum_{i=1}^{h-1}(l_i^E+r_i^E)-\sum_{i=h+1}^N(l_i^E+r_i^E)} \right) \left\langle \Upsilon, c' \right\rangle_{N|M} \\
& = \left(\sum_{i=1}^M \left(-[l_h^S + r_h^S + 1] \right) q^{N-r^E-l^E-\sum_{i=1}^{h-1}(l_i^S+r_i^S+1)+\sum_{i=h+1}^M(l_i^S+r_i^S+1)} \right. \\
& \quad \left. + \sum_{i=1}^N \left([1 - l_h^E - r_h^E] \right) q^{M+r^S+l^S+\sum_{i=1}^{h-1}(l_i^E+r_i^E-1)-\sum_{i=h+1}^N(l_i^E+r_i^E-1)} \right) \left\langle \Upsilon, c' \right\rangle_{N|M} \\
& = \left(-[M + l^S + r^S] q^{N-l^E-r^E} + [N - l^E - r^E] q^{M+l^S+r^S} \right) \left\langle \Upsilon, c' \right\rangle_{N|M} \\
& = [N - M - 2m] \left\langle \Upsilon, c' \right\rangle_{N|M}. \quad \square
\end{aligned}$$

3. LINK INVARIANTS

The aim of this section is to define link invariants using the combinatorics worked out in the previous section. The definition are really close from [MOY98]. The main point here is to fix normalization.

Definition 3.1. A *labeled* link diagram is an oriented link diagram whose components are labeled by non-negative integers. If D is a labeled link diagram, denote by $\times(D)$ (or simply \times) the set of crossings of D . A diagram is *unlabeled* or *trivially labeled* if all its components are labeled by 1. For each crossing x in \times , define k_x and e_x by the following formula:

$$k_x = \begin{cases} m(N - M - m + 1) & \text{if } x \text{ is positive and the two strands have label } m, \\ -m(N - M - m + 1) & \text{if } x \text{ is negative and the two strands have label } m, \\ 0 & \text{else;} \end{cases}$$

$$e_x = \begin{cases} m & \text{if the two strands of } x \text{ have label } m, \\ 0 & \text{else.} \end{cases}$$

Finally, define $k(D)$ (resp. $e(D)$) to be the sum of the k_x (resp. e_x) for all x in $\times(D)$.

Definition 3.2. Let D be a labeled link diagram. The $(N|M)$ -evaluation of D is the Laurent polynomial in q with integral coefficients denoted by $\langle D \rangle_{N|M}$ defined by the two following local relations:

$$(9) \quad \left\langle \begin{array}{c} \text{crossing} \\ \text{with strands } m, n \end{array} \right\rangle_{N|M} = \sum_{k=\max(0, m-n)}^m (-1)^{m-k} q^{k-m} \left\langle \begin{array}{c} \text{square} \\ \text{with strands } m, n, k, n+k-m \end{array} \right\rangle_{N|M},$$

$$(10) \quad \left\langle \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\rangle_{N|M} = \sum_{k=\max(0, m-n)}^m (-1)^{m-k} q^{m-k} \left\langle \begin{array}{c} m \quad n \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ n+k \quad m-k \\ k \end{array} \right\rangle_{N|M}.$$

Remark 3.3. In order to compute $\langle D \rangle_{N|M}$ one first expresses the diagram D as a linear combination of MOY graphs and then uses the definition of $\langle \bullet \rangle_{N|M}$ for graphs given in Definition 2.10.

Theorem 3.4. For any non-negative integers M and N , the Laurent polynomial $P_{N|M}(D) := (-1)^{e(D)} q^{k(D)} \langle D \rangle_{N|M}$ depends only on the oriented labeled link represented by D .

Proof. It is enough to check invariance under Reidemeister moves. This follows from the various identities satisfied by the $(N|M)$ -evaluation given in Section 2. The way to deduce invariance from these identities is given in [MOY98]. \square

Remark 3.5. From Corollary A.11, one deduces that for any link L , the polynomial $P_{N|M}(L)$ depends only on L and $N - M$.

Proposition 3.6. The polynomial $P_{N|M}$ satisfies the following skein relation:

$$q^{M-N} P_{N|M} \left(\begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right) - q^{N-M} P_{N|M} \left(\begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \right) = (q^{-1} - q) P_{N|M} \left(\begin{array}{c} \diagup \\ \diagdown \end{array} \right) \left(\begin{array}{c} \diagdown \\ \diagup \end{array} \right).$$

Proof. This follows from the local definition of $\langle \bullet \rangle_{N|M}$ on unlabeled crossings: simpler

$$\begin{aligned} \left\langle \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} \right\rangle_{N|M} &= \left\langle \begin{array}{c} 1 \quad 1 \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ 2 \\ 1 \quad 1 \end{array} \right\rangle_{N|M} - q^{-1} \left\langle \begin{array}{c} \diagup \\ \diagdown \end{array} \right\rangle_{N|M} \left(\begin{array}{c} \diagdown \\ \diagup \end{array} \right)_{N|M}, \\ \left\langle \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \right\rangle_{N|M} &= \left\langle \begin{array}{c} 1 \quad 1 \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ 2 \\ 1 \quad 1 \end{array} \right\rangle_{N|M} - q \left\langle \begin{array}{c} \diagup \\ \diagdown \end{array} \right\rangle_{N|M} \left(\begin{array}{c} \diagdown \\ \diagup \end{array} \right)_{N|M}. \end{aligned}$$

Taking in account the contribution of $k(D)$ and $(-1)^{e(D)}$ in the definition of $P_{N|M}$ gives the identity. \square

Remark 3.7. The connection between $U_q(\mathfrak{gl}_{N|M})$ -link invariants for links labeled by an arbitrary partition λ and generic versions of them known as colored HOMFLY-PT polynomials was explored by Queffelec and Sartori [QS15]. We refer the reader to this paper for a detailed account. In particular, they show that such invariants depend only on $N - M$ and the partition λ . The present paper provides an alternative proof for this well-known fact for partitions which are rows.

4. NON SEMI-SIMPLE INVARIANTS

In this part we suppose that $1 \leq M \leq N$. Let us consider a link L and suppose that one of its component has label n with $n > N - M$. One can show that in this case the $\mathfrak{gl}_{N|M}$ -invariant is equal to 0 (see for instance [QS15]. This follows from the fact that the $\mathfrak{gl}_{N|M}$ -invariant of the unknot labeled n is equal to $\begin{bmatrix} M-N \\ n \end{bmatrix}$ which happens to be 0. The aim of this part is to directly re-normalize the $\mathfrak{gl}_{N|M}$ invariant in this case in order to get a non-trivial invariant. The theory of renormalized invariants is

nowadays well developed but we refer to one the early paper by Geer and Patureau-Mirand which treats the case we are looking at [GPM10] and also to the treatment by Queffelec and Sartori [QS15]. The interested reader could also consult the paper by Geer, Patureau-Mirand and Turaev [GPMT09].

Here, we focus on the case $M = 1$ and $n = N$. Incidentally, the re-normalized $g_{1|1}$ -invariant for links uniformly colored by 1 equals the (one-variable) Alexander polynomial. Hence, this construction can be thought of a generalization of the Alexander polynomial. Let us point out that these invariants do not coincide with the colored Alexander polynomials (or ADO invariants) [ADO92].

Definition 4.1. A *marked MOY graph* Γ_\star is a MOY graph Γ with a base point \star in the interior of one of its edges. If the marked point is on an edge of label k , we say that Γ_\star has *type* k .

Definition 4.2. Let Γ_\star be a marked MOY graph and denote e the edge of Γ which contains the base point. A $(N|1)$ -coloring of Γ_\star is a $(N|1)$ -coloring $(\underline{\Gamma}^E, \underline{\Gamma}^S)$ of the underlying Γ such that for all j in $[N]$, $\ell_{\Gamma_j^E}(e) = 0$. The set of $(N|1)$ -coloring of Γ_\star is denoted by $\text{col}_{N|1}(\Gamma_\star)$ (or simply $\text{col}(\Gamma_\star)$).

If Γ_\star is a marked MOY graph and c is a $(N|1)$ -coloring of Γ_\star , define $\langle \Gamma_\star, c \rangle_{N|1} := \langle \Gamma, c \rangle_{N|1}$, and:

$$\langle \Gamma_\star \rangle_{N|1} = \sum_{c \in \text{col}_{N|1}(\Gamma_\star)} \langle \Gamma_\star, c \rangle_{N|1}.$$

Remark 4.3. (1) In the previous definition, $\underline{\Gamma}^S$ contains exactly one sub-MOY graph: Γ_1^S . One has necessarily $\ell_{\Gamma_1^S}(e) = \ell_\Gamma(e)$.

(2) Suppose that Γ_\star is a marked circle of label k with $k \geq N$, and denote by Γ the underlying not marked circle. One has:

$$\langle \Gamma_\star \rangle_{N|1} = 1 \quad \text{and} \quad \langle \Gamma \rangle_{N|1} = 0.$$

The results of the previous section extend naturally:

Proposition 4.4. *The local relations given in Proposition A.9 are still valid for the evaluation of marked MOY graphs (provided the marked point is not in the ball where the relations take place).*

Definition 4.5. Let β be a braid diagram, define $\langle \beta_\star \rangle_{N|1}$ by the following procedure:

- (1) Label all strands of β by N .
- (2) Place a marked point on the left-most strand at the bottom of β .
- (3) Close β on the right.
- (4) Use formulas (9) and (10) to get rid of crossings.
- (5) Evaluate the obtained marked MOY graphs with $\langle \bullet \rangle_{N|1}$ as given in Definition 4.2.

As in Section 3, $\langle \bullet \rangle_{N|1}$ needs to be normalized to obtain a link invariant. Define $Q_{N|1}(\beta) = (-1)^{N|\beta|} \langle \bullet \rangle_{N|1}$, where $|\beta|$ is the number of crossings of β .

Remark 4.6. Besides the appearances, the normalization used for $Q_{N|1}$ is the same as for $P_{N|1}$ (see Definition 3.1) but is especially simple for the present choice of labels.

Proposition 4.7. *The Laurent polynomial $Q_{N|1}(\beta)$ depends only on the link represented by β .*


Notation 4.8. If a link L can be obtained by closing a braid β , set $Q_{N|1}(L) := Q_{N|1}(\beta)$.

Remark 4.9. The invariant $Q_{N|1}$ vanishes on split links.


Proof of Proposition 4.7. We need to prove invariance under

- Braid relations.
- Markov moves.


Invariance under braid relations and stabilization follows from the general setting see [MOY98]. Invariance under conjugation by σ_i for $i \geq 2$ is obvious. Hence the only thing to show is invariance under conjugation by σ_1 . On the closure of β it only changes the base point in the following way:



It is enough to show that



for any MOY graph Γ with four ends. One can actually suppose (see [SW17, Section 2.3.1]) that Γ is equal to a linear combination of MOY graph of the form




with $0 \leq j \leq N$. Using relation (9), one can express $\langle \beta_1 \rangle$ and $\langle \beta_2 \rangle$ as:

$$\langle \beta_1 \rangle_{N|1} = \sum_{i=0}^n \sum_{j=0}^n \lambda_{ij} \langle \Gamma_{\star}^{ij} \rangle_{N|1} \quad \text{and}$$

$$\langle \beta_2 \rangle_{N|1} = \sum_{i=0}^n \sum_{j=0}^n \lambda_{ij} \langle \Phi_{\star}^{ij} \rangle_{N|1}$$

Where



Using the skein relations, one can compute $\langle \Gamma_\star^{ij} \rangle_{N|1}$ and $\langle \Phi_\star^{ij} \rangle_{N|1}$:

$$\begin{aligned} \langle \Gamma_\star^{ij} \rangle_{N|1} &= \begin{bmatrix} i+j \\ i \end{bmatrix} \begin{bmatrix} N-1-(i+j) \\ N-i \end{bmatrix} \begin{bmatrix} N-1-N \\ i \end{bmatrix} \begin{bmatrix} N \\ j \end{bmatrix}, \\ \langle \Phi_\star^{ij} \rangle_{N|1} &= \begin{bmatrix} i+j \\ i \end{bmatrix} \begin{bmatrix} N-1-(i+j) \\ N-j \end{bmatrix} \begin{bmatrix} N-1-N \\ j \end{bmatrix} \begin{bmatrix} N \\ i \end{bmatrix}. \end{aligned}$$

These two products of quantum binomials are equal. Hence $\langle \beta_1 \rangle_{N|1} = \langle \beta_2 \rangle_{N|1}$ and therefore $Q(\beta_1)_{N|1} = Q(\beta_2)_{N|1}$. \square

Theorem 4.10. *For any $N \in \mathbb{Z}_{\geq 0}$ and any link L , one has:*

$$Q_{N|1}(L, q = e^{\frac{i\pi}{N+1}}) = J'_N(\bar{L}, q = e^{\frac{i\pi}{N+1}}),$$

where \bar{L} denotes the mirror image of L .

Note that the sequence $(J'_n(\bar{L}, q = e^{\frac{i\pi}{n+1}}))_{n \in \mathbb{Z}_{>0}}$ appears in the formulation of the volume conjecture [Kas95, MM01]. For proving this theorem we need a technical result about quantum binomials evaluated at root of unity.

Lemma 4.11. *Let N , a and b be three integers. The following identity holds:*

$$\begin{bmatrix} N-1-a \\ b \end{bmatrix}_{q=e^{\frac{i\pi}{N+1}}} = (-1)^b \begin{bmatrix} -2-a \\ b \end{bmatrix}_{q=e^{\frac{i\pi}{N+1}}}$$

Proof. We assume here that $q = e^{\frac{i\pi}{N+1}}$.

$$\begin{aligned} \begin{bmatrix} N-1-a \\ b \end{bmatrix} &= \left(\prod_{i=1}^b \frac{q^{N-1-a+i} - q^{-N-a+1-i}}{q^i - q^{-i}} \right) \\ &= \left(\prod_{i=1}^b \frac{-q^{-2-a+i} + q^{-2-a-i}}{q^i - q^{-i}} \right) \\ &= (-1)^b \begin{bmatrix} -2-a \\ b \end{bmatrix}. \end{aligned} \quad \square$$

Proof of Theorem 4.10. Let us fix a braid diagram β representing L and denote k the braid index of β . Once every strand of β is labeled by N , the level of β is kN . We will compare $\langle \beta_\star \rangle_{N|1}$ and $\frac{\langle \beta \rangle_{0|2}}{[-N]}$ evaluated at $q = e^{\frac{i\pi}{N+1}}$. The expansion of crossings is the same in both cases. Hence it is enough to compare

$$\langle \Gamma_\star \rangle_{N|1} \Big|_{q=e^{\frac{i\pi}{N+1}}} \quad \text{and} \quad \frac{\langle \Gamma \rangle_{0|2}}{[-N]} \Big|_{q=e^{\frac{i\pi}{N+1}}}$$

for Γ_\star a marked MOY graph appearing in the expansion of β_\star . Both these quantities can be computed using the relations in Proposition A.9 for $(N|1)$ and $(0|2)$ and the fact that $\langle U_\star \rangle_{N|1} = \frac{\langle U \rangle_{0|2}}{[-N]} = 1$ for U a circle of label N . More precisely, one has:

$$\langle \Gamma_\star \rangle_{N|1} = R_\Gamma(N-1, q) \langle U_\star \rangle_{N|1} \quad \text{and} \quad \frac{\langle \Gamma \rangle_{0|2}}{[-N]} = R_\Gamma(-2, q) \frac{\langle U \rangle_{0|2}}{[-N]},$$

for a $R_\Gamma(n, q)$ a sum of product of quantum binomials of the form $\begin{bmatrix} a_i \\ b_i \end{bmatrix}$ and $\begin{bmatrix} n-c_j \\ d_j \end{bmatrix}$. The first binomials correspond to relations (13), (16) and (17) which preserve the level, the second correspond to relations (11) (14) (15) which do not preserve the

level. Note that a binomial $\begin{bmatrix} n - c_j \\ d_j \end{bmatrix}$ appears exactly when the level decreases by d_j . Since to go from Γ to U , the level is decreases by $(k - 1)N$, for each product of binomials in $R_\Gamma(n, q)$, the sum of the d_j 's equals $(k - 1)N$. This gives, in view of Lemma 4.11:

$$R_\Gamma\left(N - 1, e^{\frac{i\pi}{N+1}}\right) = (-1)^{(k-1)N} R_\Gamma\left(-2, e^{\frac{i\pi}{N+1}}\right)$$

and therefore

$$\left\langle \beta_\star, q = e^{\frac{i\pi}{N+1}} \right\rangle_{N|1} = (-1)^{(k-1)N} \frac{\left\langle \beta, q = e^{\frac{i\pi}{N+1}} \right\rangle_{0|2}}{[-N]_{q=e^{\frac{i\pi}{N+1}}}}.$$

Since all strands are labeled by N , one has:

$$\begin{aligned} P_{0|2}(L) &= (-1)^{N(c_+ + c_-)} q^{N(N+1)(c_- - c_+)} \langle \beta \rangle_{0|2} \quad \text{and} \\ Q_{N|1}(L) &= (-1)^{N(c_+ + c_-)} \langle \beta_\star \rangle_{N|1}, \end{aligned}$$

where c_+ and c_- are the number of positive and negative crossings of β respectively. Hence, for $q = \exp(\frac{i\pi}{N+1})$,

$$Q_{N|1}(L) = (-1)^{N(k-1+c_- - c_+)} \frac{P_{0|2}(L)}{[-N]} = (-1)^{N(k+c_- - c_+)} \frac{P_{0|2}(L)}{[N]}.$$

It turns out that $k + c_- - c_+$ has the same parity as the number ℓ of components of L . This gives:

$$Q_{N|1}(L, q = e^{\frac{i\pi}{N+1}}) = (-1)^{N\ell} \frac{P_{0|2}(L)}{[N]} \Big|_{q=e^{\frac{i\pi}{N+1}}} = J'_N(\bar{L}, q = e^{\frac{i\pi}{N+1}}). \quad \square$$

Remark 4.12. The previous result can also be obtained in an indirect way from the work of Geer and Patureau-Mirand [GPM08]. In this paper, they prove that the generalized Links–Gould invariants specialize to the Kashaev invariants. These invariants are constructed using typical representations $V(\alpha)$ of $U_q(\mathfrak{gl}_{N|1})$ depending on a complex parameter α . For $\alpha = -1$, these are exactly the representations considered in the present paper and this specialization $\alpha = -1$ is compatible with the ones providing the Kashaev invariants.

APPENDIX A. MOY GRAPHS, AN ALGEBRAIC APPROACH

For a general introduction to the Reshetikhin–Turaev functors, we refer to Turaev's book [Tur94]. For more details on the super setting, we refer to Geer and Patureau-Mirand [GPM10] and references therein. The later paper can also be consulted for the renormalization process (see also [QS15]). Here, we follow and expand the diagrammatic presentation of Tubbenhauer–Vaz–Wedrich [TVW17]. For the reader convenience, we made explicit the various pieces of the Reshetikhin–Turaev functor. All proofs are omitted since they are all direct verifications.

In what follows, a *super algebra* is presented. It means that it is endowed with a $\mathbb{Z}/2$ -grading and more importantly, its modules are objects of Svect the category of super vector spaces. Objects of this category are $\mathbb{Z}/2$ -graded vector spaces, morphisms are linear maps preserving the $\mathbb{Z}/2$ grading. The $\mathbb{Z}/2$ -grading, called *parity*, is denoted $|\bullet|$. Homogeneous elements with parity equal to 0 (resp. 1) are *even* (resp. *odd*). This category inherits from vect a monoidal structure and a duality. The

braiding c on Svect differs from that of vect :

$$\begin{aligned} c_{V,W}: V \otimes W &\rightarrow W \otimes V \\ v \otimes w &\mapsto (-1)^{|v||w|} w \otimes v. \end{aligned}$$

If V is an object of Svect which has finite dimension as a vector space. Its *super dimension* $\text{sdim } V$ is the integer $\dim V_0 - \dim V_1$, where $V = V_0 \oplus V_1$ is the $\mathbb{Z}/2$ -decomposition². The *classical dimension* of V is its dimension as a vector space.

Definition A.1. Let N and M be two non-negative integers. The *quantum general linear superalgebra* $U_q(\mathfrak{gl}_{N|M})$ is the associative, unital, $\mathbb{Z}/2$ -graded $\mathbb{C}(q)$ -algebra generated by L_i, L_i^{-1}, F_j and E_j , with $1 \leq i \leq N+M$ and $1 \leq j \leq N+M-1$ subject to the nonsuper relations

$$\begin{aligned} L_i L_j &= L_j L_i, & L_i L_i^{-1} &= L_i^{-1} L_i = 1, \\ L_i E_i &= q E_i L_i, & L_i E_{i-1} &= q^{-1} E_{i-1} L_i, & \text{for } i \leq N, \\ L_i F_i &= q^{-1} F_i L_i, & L_i F_{i-1} &= q F_{i-1} L_i, & \text{for } i \leq N, \\ L_i E_i &= q^{-1} E_i L_i, & L_i E_{i-1} &= q E_{i-1} L_i, & \text{for } i \geq N+1, \\ L_i F_i &= q F_i L_i, & L_i F_{i-1} &= q^{-1} F_{i-1} L_i, & \text{for } i \geq N+1, \\ L_i F_j &= F_j L_i, & L_i E_j &= E_j L_i & \text{for } j \neq i, i-1, \\ E_i F_j - F_j E_i &= \delta_{ij} \frac{L_i L_{i+1}^{-1} - L_i^{-1} L_{i+1}}{q - q^{-1}}, & \text{for } 1 \leq i \leq N-1, \\ E_i F_j - F_j E_i &= -\delta_{ij} \frac{L_i L_{i+1}^{-1} - L_i^{-1} L_{i+1}}{q - q^{-1}}, & \text{for } N+1 \leq i < N+M-1, \\ [2]_q F_i F_j F_i &= F_i^2 F_j + F_j F_i^2 & \text{if } |i-j| = 1 \text{ and } i \neq N, \\ [2]_q E_i E_j E_i &= E_i^2 E_j + E_j E_i^2 & \text{if } |i-j| = 1 \text{ and } i \neq N, \\ E_i E_j &= E_j E_i, & F_i F_j &= F_j F_i & \text{if } |i-j| > 1 \end{aligned}$$

and the super relations

$$\begin{aligned} E_N^2 = F_N^2 = 0 & \quad E_N F_N + F_N E_N = \frac{L_N L_{N+1}^{-1} - L_N^{-1} L_{N+1}}{q - q^{-1}} \\ [2] F_N F_{N-1} F_{N+1} F_N &= \\ F_N F_{N-1} F_N F_{N+1} + F_N F_{N+1} F_N F_{N-1} + F_{N-1} F_N F_{N+1} F_N + F_{N+1} F_N F_{N-1} F_N, \\ [2] E_N E_{N-1} E_{N+1} E_N &= \\ E_N E_{N-1} E_N E_{N+1} + E_N E_{N+1} E_N E_{N-1} + E_{N-1} E_N E_{N+1} E_N + E_{N+1} E_N E_{N-1} E_N. \end{aligned}$$

All these generators are $\mathbb{Z}/2$ -homogeneous and even, but E_N and F_N which are $\mathbb{Z}/2$ -homogeneous and odd.

Proposition A.2. Defining $\Delta: U_q(\mathfrak{gl}_{N|M}) \rightarrow U_q(\mathfrak{gl}_{N|M})^{\otimes 2}$, $S: U_q(\mathfrak{gl}_{N|M})^{\text{op}} \rightarrow U_q(\mathfrak{gl}_{N|M})$ and $\epsilon: U_q(\mathfrak{gl}_{N|M}) \rightarrow \mathbb{C}(q)$ to be the $\mathbb{C}(q)$ algebra maps defined by:

$$\begin{aligned} \Delta(L_i^{\pm 1}) &= L_i^{\pm 1} \otimes L_i^{\pm 1} & S(L_i^{\pm 1}) &= L_i^{\mp 1} & \epsilon(L_i^{\pm 1}) &= 1 \\ \Delta(F_i) &= F_i \otimes 1 + L_i^{-1} L_{i+1} \otimes F_i & S(F_i) &= -L_i L_{i+1}^{-1} F_i & \epsilon(F_i) &= 0 \\ \Delta(E_i) &= E_i \otimes 1 + L_i L_{i+1}^{-1} \otimes E_i & S(E_i) &= -E_i L_i^{-1} L_{i+1} & \epsilon(E_i) &= 0 \end{aligned}$$

²The super dimension is actually the categorical dimension, the sign coming from the braiding on Svect .

endows $U_q(\mathfrak{gl}_{N|M})$ with a structure of $\mathbb{Z}/2$ -graded Hopf algebra with antipode. Furthermore the category of finite-dimensional $U_q(\mathfrak{gl}_{N|M})$ -modules is braided.

Remark A.3. The Hopf algebras $U_q(\mathfrak{gl}_{N|M})$ and $U_q(\mathfrak{gl}_{M|N})$ are isomorphic. Indeed, one easily checks that the map

$$\begin{aligned} \varphi: U_q(\mathfrak{gl}_{N|M}) &\rightarrow U_q(\mathfrak{gl}_{M|N}) \\ L_i &\mapsto L_{M+N+1-i}^{-1} \\ L_i^{-1} &\mapsto L_{M+N+1-i} \\ E_i &\mapsto F_{M+N-i} \\ F_i &\mapsto e_{M+N-i} \end{aligned}$$

induces an isomorphism of Hopf algebras.

Proposition A.4. Let $\mathbb{C}_q^{N|M}$ be the $\mathbb{Z}/2$ -graded $\mathbb{C}(q)$ -vector space generated by the homogeneous basis $(b_i)_{i=1,\dots,N+M}$ (with $|b_i| = 0$ if $i \leq N$ and $|b_i| = 1$ if $i > N$). The formulas

$$\begin{aligned} L_i b_i &= q b_i, & L_i^{-1} b_i &= q^{-1} b_i, & \text{if } 1 \leq i \leq N, \\ L_j b_j &= q^{-1} b_j, & L_j^{-1} b_j &= q b_j, & \text{if } N+1 \leq j \leq N+M, \\ L_i^{\pm 1} b_j &= b_j & & & \text{if } i \neq j, \\ E_{i-1} b_i &= b_{i-1} & E_i b_j &= 0, & \text{if } i \neq j-1, \\ F_i b_i &= b_{i+1} & F_i b_j &= 0, & \text{if } i \neq j \end{aligned}$$

endow $\mathbb{C}_q^{N|M}$ with a structure of $U_q(\mathfrak{gl}_{N|M})$ -module.

Remark A.5. Since $U_q(\mathfrak{gl}_{N|M})$ is isomorphic to $U_q(\mathfrak{gl}_{M|N})$, $\mathbb{C}_q^{M|N}$ inherits a structure of $U_q(\mathfrak{gl}_{N|M})$ -module. Denote $\mathbf{1}_{\text{odd}}$ the trivial³ one-dimensional $U_q(\mathfrak{gl}_{N|M})$ -module concentrated in odd parity and $\mathbf{1}_{\text{odd}}$ a generator of this module. The $U_q(\mathfrak{gl}_{N|M})$ -modules $\mathbb{C}_q^{M|N}$ and $\mathbf{1}_{\text{odd}} \otimes \mathbb{C}_q^{N|M}$ are isomorphic. Indeed, one can check that

$$\begin{aligned} \psi: \mathbb{C}_q^{M|N} &\rightarrow \mathbb{C}_q^{N|M} \otimes \mathbf{1}_{\text{odd}} \\ b_i &\mapsto b_{N+M-i+1} \otimes \mathbf{1}_{\text{odd}} \end{aligned}$$

induces an isomorphism of $U_q(\mathfrak{gl}_{N|M})$ -modules.

Denote $T\mathbb{C}_q^{N|M}$ the $\mathbb{Z}/2 \times \mathbb{Z}$ -graded algebra generated by $\mathbb{C}_q^{N|M}$ and $\text{Sym}_q^2 \mathbb{C}_q^{N|M}$ the ideal generated by the set

$$\{b_i \otimes b_i | 1 \leq i \leq N\} \cup \{q^{-1} b_i \otimes b_j + (-1)^{|b_i||b_j|} b_j \otimes b_i | 1 \leq i < j \leq N+M\}.$$

Denote $\Lambda_q \mathbb{C}_q^{N|M}$ the space $T\mathbb{C}_q^{N|M} / \text{Sym}_q^2 \mathbb{C}_q^{N|M}$. For $k \in \mathbb{Z}_{\geq 0}$, denote $\Lambda_q^k \mathbb{C}_q^{N|M}$ the k -degree (for the \mathbb{Z} -grading) part of $\Lambda_q \mathbb{C}_q^{N|M}$.

Proposition A.6. For all k , $\Lambda_q^k \mathbb{C}_q^{N|M}$ inherits a $U_q(\mathfrak{gl}_{N|M})$ -module structure. Its super dimension over $\mathbb{C}(q)$ is $\binom{N-M}{k}$. Its classical dimension is

$$\sum_{i=0}^k \binom{N}{k-i} \binom{M+i-1}{i}.$$

³This means that the generators $L_i^{\pm 1}$ act by 1 and the other generators act by 0.

In what follows, we define some morphisms in the category of $U_q(\mathfrak{gl}_{N|M})$ -modules. For this, we need to introduce a few notations.

The image of a pure tensor $x_1 \otimes \cdots \otimes x_k$ of $T\mathbb{C}_q^{N|M}$ in $\Lambda^k \mathbb{C}_q^{N|M}$ is denoted by $x_1 \wedge \cdots \wedge x_k$. In particular one has:

$$\begin{aligned} b_i \wedge b_i &= 0 \quad \text{if } i \leq N, \\ b_i \wedge b_j &= -(-1)^{|b_i||b_j|} q b_j \wedge b_i \quad \text{for } i < j. \end{aligned}$$

The $\mathbb{C}(q)$ -vector space $\Lambda_q^k \mathbb{C}_q^{N|M}$ is spanned by the vectors

$$\left(b_{i_1} \wedge \cdots \wedge b_{i_{k^E}} \wedge b_{j_1} \wedge \cdots \wedge b_{j_{k^S}} \right)_{\substack{1 \leq i_1 < \cdots < i_{k^E} \leq N \\ N+1 \leq j_1 \leq \cdots \leq j_{k^S} \leq N+M \\ k = k^E + k^S}}.$$

If $1 \leq i_1 < \cdots < i_{k^E} \leq N$ and $N+1 \leq j_1 \leq \cdots \leq j_{k^S} \leq N+M$, denote $b_{I,J} = b_{i_1} \wedge \cdots \wedge b_{i_{k^E}} \wedge b_{j_1} \wedge \cdots \wedge b_{j_{k^S}}$, where $I = \{i_1, \dots, i_{k^E}\}$ and $J = \{j_1, \dots, j_{k^S}\}$. Note that I is a subset of $\{1, \dots, N\}$ while J is a multi-subset of $\{N+1, \dots, N+M\}$. With these notations, $(b_{I,J})_{\#I+\#J=k}$ is an homogeneous basis of $\Lambda^k \mathbb{C}_q^{N|M}$. Denote $(b^{I,J})_{\#I+\#J=k}$ its dual basis and define the following morphisms⁴:

$$\begin{aligned} \Lambda_{k,\ell}: \Lambda_q^k V_q \otimes \Lambda_q^\ell V_q &\rightarrow \Lambda_q^{k+\ell} V_q \\ b_{I_1, J_1} \otimes b_{I_2, J_2} &\rightarrow \begin{cases} q^{-|(I_2 \cup J_2) < (I_1 \cup J_1)|} b_{I_1 \sqcup I_2, J_1 \sqcup J_2} & \text{if } I_1 \cap I_2 = \emptyset, \\ 0 & \text{else.} \end{cases} \\ Y_{k,\ell}: \Lambda_q^{k+\ell} V_q &\rightarrow \Lambda_q^k V_q \otimes \Lambda_q^\ell V_q \\ b_{I,J} &\rightarrow \sum_{\substack{I_1 \sqcup I_2 = I, J_1 \sqcup J_2 = J \\ \#I_1 + \#J_1 = k, \#I_2 + \#J_2 = \ell}} [J_1, J_2] q^{|(I_2 \cup J_2) < (I_1 \cup J_1)|} b_{I_1, J_1} \otimes b_{I_2, J_2} \\ \bar{U}_k: \mathbb{C}(q) &\rightarrow \Lambda_q^k V_q \otimes (\Lambda_q^k V_q)^* \\ 1 &\rightarrow \sum_{\#I=k} b_{I,J} \otimes b^{I,J} \\ \bar{\cap}_k: (\Lambda_q^k V_q)^* \otimes \Lambda_q^k V_q &\rightarrow \mathbb{C}(q) \\ f \otimes x &\rightarrow f(x) \\ \bar{U}_k: \mathbb{C}(q) &\rightarrow (\Lambda_q^k V_q)^* \otimes \Lambda_q^k V_q \\ 1 &\rightarrow \sum_{\substack{k = k^E + k^S \\ \#I = k^E, \#J = k^S}} q^{\deg_{[M+N]}(J) - \deg_{[M+N]}(I)} (-1)^{k^S} b^{I,J} \otimes b_{I,J} \\ \bar{\cap}_k: \Lambda_q^k V_q \otimes (\Lambda_q^k V_q)^* &\rightarrow \mathbb{C}(q) \\ b_{I_1, J_1} \otimes b^{I_2, J_2} &\rightarrow q^{\deg_{[M+N]}(I_1) - \deg_{[M+N]}(J_2)} (-1)^{|J_1|} \delta_{I_1, I_2} \delta_{J_1, J_2} \end{aligned}$$


We should explain what $|I < J|$ and $[I, J]$ mean. If I and J are two multi-subsets of an ordered set X , define

$$\begin{aligned} |I < J| &= \prod_{x < y \in X} I(x)J(y) \quad \text{and} \\ [I, J] &= \prod_{x \in X} \begin{bmatrix} I(x) + J(x) \\ I(x) \quad J(x) \end{bmatrix}. \end{aligned}$$

Proposition A.7. *These maps are morphisms of $U_q(\mathfrak{gl}_{N|M})$ -modules.*

⁴See [RW18, Appendix A] for similar definitions in the symmetric case, i. e. $N = 0$.

Any MOY graph Γ is isotopic to a MOY graph Γ' *in good position*, this means that Γ' can be obtained by vertically concatenating horizontal disjoint unions of the following elementary pieces:



When a MOY graph Γ' is in good position, interpreting these elementary pieces using the morphisms

$$\text{Id}_{\Lambda_q^k \mathbb{C}^{N|M}}, \text{Id}_{(\Lambda_q^\ell \mathbb{C}^{N|M})^*}, \bar{\cap}_k, \bar{\cap}_\ell, \bar{\cup}_k, \bar{\cup}_\ell, \Lambda_{k,\ell}, Y_{k,\ell}$$

horizontal disjoint unions as tensor products and vertical concatenations as compositions, one obtains a morphism of $U_q(\mathfrak{gl}_{N|M})$ -modules $\langle\langle \Gamma' \rangle\rangle_{N|M} : \mathbb{C}(q) \rightarrow \mathbb{C}(q)$. One sees $\langle\langle \Gamma' \rangle\rangle_{N|M}$ as an element of $\mathbb{C}(q)$. Using the definition of the morphisms associated with the elementary pieces, one can show that $\langle\langle \Gamma' \rangle\rangle_{N|M}$ is a Laurent polynomial in q with integer coefficients and symmetric in q and q^{-1} . If Γ is an arbitrary MOY graph, define $\langle\langle \Gamma \rangle\rangle_{N|M}$ to be $\langle\langle \Gamma' \rangle\rangle_{N|M}$ for Γ' a MOY graph in good position and isotopic to Γ .

Proposition A.8. *The Laurent polynomial $\langle\langle \Gamma \rangle\rangle_{N|M}$ is well-defined i. e. it does not depend on the MOY graph in good position and isotopic to Γ chosen to compute it.*

Proposition A.9. *The map $\langle\langle \bullet \rangle\rangle_{N|M} : \{\text{MOY graphs}\} \rightarrow \mathbb{Z}[q, q^{-1}]$ is multiplicative under disjoint union i. e.*

$$\langle\langle \Gamma \sqcup Y \rangle\rangle_{N|M} = \langle\langle \Gamma \rangle\rangle_{N|M} \langle\langle Y \rangle\rangle_{N|M} \quad \text{for all MOY graphs } \Gamma \text{ and } Y.$$

and satisfies the following the local relations and their mirror images:

$$(11) \quad \left\langle\left\langle \begin{array}{c} \text{circle with arrow} \\ \text{dot on right} \end{array} \right\rangle\right\rangle_{N|M} = \begin{bmatrix} N-M \\ k \end{bmatrix}$$

$$(12) \quad \left\langle\left\langle \begin{array}{c} i \quad j \quad k \\ \swarrow \quad \searrow \quad \nearrow \\ \text{dot on bottom} \\ i+j+k \end{array} \right\rangle\right\rangle_{N|M} = \left\langle\left\langle \begin{array}{c} i \quad j \quad k \\ \swarrow \quad \searrow \quad \nearrow \\ \text{dot on bottom} \\ i+j+k \end{array} \right\rangle\right\rangle_{N|M}$$

$$(13) \quad \left\langle\left\langle \begin{array}{c} m+n \\ \uparrow \\ m \\ \downarrow \\ m+n \end{array} \right\rangle\right\rangle_{N|M} = \begin{bmatrix} m+n \\ m \end{bmatrix} \left\langle\left\langle \begin{array}{c} \uparrow \\ m+n \end{array} \right\rangle\right\rangle_{N|M}$$

$$(14) \quad \left\langle\left\langle \begin{array}{c} m \\ \uparrow \\ m+n \\ \downarrow \\ m \end{array} \right\rangle\right\rangle_{N|M} = \begin{bmatrix} N-M-m \\ n \end{bmatrix} \left\langle\left\langle \begin{array}{c} \uparrow \\ m \end{array} \right\rangle\right\rangle_{N|M}$$

$$(15) \quad \left\langle\left\langle \begin{array}{c} 1 \quad m \\ \uparrow \quad \downarrow \\ m+1 \\ \leftarrow \quad \rightarrow \\ m+1 \\ \downarrow \quad \uparrow \\ 1 \quad m \end{array} \right\rangle\right\rangle_{N|M} = \left\langle\left\langle \begin{array}{c} \uparrow \\ 1 \end{array} \right\rangle\right\rangle_{N|M} \left\langle\left\langle \begin{array}{c} \downarrow \\ m \end{array} \right\rangle\right\rangle_{N|M} + [N-M-m-1] \left\langle\left\langle \begin{array}{c} 1 \quad m \\ \swarrow \quad \searrow \\ m-1 \\ \swarrow \quad \searrow \\ 1 \quad m \end{array} \right\rangle\right\rangle_{N|M}$$

(16)

$$\left\langle\left\langle \begin{array}{c} l \quad m \\ \begin{array}{ccc} \xrightarrow{n} & & \xrightarrow{m-n} \\ \xleftarrow{l+n-1} & & \xleftarrow{m+l-1} \\ 1 & & m+l-1 \end{array} \end{array} \right\rangle\right\rangle_{N|M} = \begin{bmatrix} m-1 \\ n \end{bmatrix} \left\langle\left\langle \begin{array}{c} l \quad m \\ \begin{array}{ccc} \xrightarrow{l-1} & & \xrightarrow{m} \\ \xleftarrow{m+l-1} & & \xleftarrow{m+l-1} \\ 1 & & m+l-1 \end{array} \end{array} \right\rangle\right\rangle_{N|M} + \begin{bmatrix} m-1 \\ n-1 \end{bmatrix} \left\langle\left\langle \begin{array}{c} l \quad m \\ \begin{array}{ccc} \xrightarrow{l} & & \xrightarrow{m} \\ \xleftarrow{l+m} & & \xleftarrow{m+l-1} \\ 1 & & m+l-1 \end{array} \end{array} \right\rangle\right\rangle_{N|M}$$

$$(17) \quad \left\langle\left\langle \begin{array}{c} m \quad n+l \\ \begin{array}{ccc} \xrightarrow{n+k-m} & & \xrightarrow{m+l-k} \\ \xleftarrow{k} & & \xleftarrow{m+l} \\ n & & m+l \end{array} \end{array} \right\rangle\right\rangle_{N|M} = \sum_{j=\max(0, m-n)}^m \begin{bmatrix} l \\ k-j \end{bmatrix} \left\langle\left\langle \begin{array}{c} m \quad n+l \\ \begin{array}{ccc} \xrightarrow{j} & & \xrightarrow{n+l+j} \\ \xleftarrow{n+j-m} & & \xleftarrow{m+l} \\ n & & m+l \end{array} \end{array} \right\rangle\right\rangle_{N|M}$$

Proposition A.10 ([Wu14]). *The multiplicativity property and the local relations given in Proposition A.9 are enough to compute the value of $\langle\langle \Gamma \rangle\rangle_{N|M}$ for any MOY graph Γ .*

From this statement we immediately get the following corollary which should be compared with [QS15, Theorem 4.7].

Corollary A.11. *The Laurent polynomial $\langle\langle \Gamma \rangle\rangle_{N|M}$ only depends on Γ and $N - M$.*

Two special cases are especially easy to compute: $N - M = \pm 1$.

Proposition A.12. *Suppose $N - M = 1$, $\langle\langle \Gamma \rangle\rangle_{N|M} = 0$ unless Γ is a (maybe empty) collection of circles of label 1 and $\langle\langle \sqcup_i \bigcirc_1 \rangle\rangle_{N|M} = 1$.*

Proposition A.13. *Suppose $N - M = -1$, then for any MOY graph Γ , the following identity holds:*

$$\langle\langle \Gamma \rangle\rangle_{N|M} = (-1)^{\rho(\Gamma)} b(\Gamma),$$

where $b(\Gamma)$ is given in Definition 2.4.

REFERENCES

- [ADO92] Yasuhiro Akutsu, Tetsuo Deguchi, and Tomotada Ohtsuki. Invariants of colored links. *J. Knot Theory Ramifications*, 1(2):161–184, 1992. doi:10.1142/S0218216592000094.
- [Cau17] Sabin Cautis. Remarks on coloured triply graded link invariants. *Algebr. Geom. Topol.*, 17(6):3811–3836, 2017. doi:10.2140/agt.2017.17.3811.
- [EPV19] Alexander P. Ellis, Ina Petkova, and Vera Vértési. Quantum $\mathfrak{gl}_{1|1}$ and tangle Floer homology. *Adv. Math.*, 350:130–189, 2019. doi:10.1016/j.aim.2019.04.023.
- [GGS18] Eugene Gorsky, Sergei Gukov, and Marko Stošić. Quadruply-graded colored homology of knots. *Fund. Math.*, 243(3):209–299, 2018. doi:10.4064/fm30-11-2017.
- [GPM08] Nathan Geer and Bertrand Patureau-Mirand. On the colored HOMFLY-PT, multivariable and Kashaev link invariants. *Commun. Contemp. Math.*, 10(suppl. 1):993–1011, 2008. doi:10.1142/S0219199708003125.
- [GPM10] Nathan Geer and Bertrand Patureau-Mirand. Multivariable link invariants arising from Lie superalgebras of type I. *J. Knot Theory Ramifications*, 19(1):93–115, 2010. doi:10.1142/S0218216510007784.
- [GPMT09] Nathan Geer, Bertrand Patureau-Mirand, and Vladimir Turaev. Modified quantum dimensions and re-normalized link invariants. *Compos. Math.*, 145(1):196–212, 2009. doi:10.1112/S0010437X08003795.

- [Kas95] R. M. Kashaev. A link invariant from quantum dilogarithm. *Modern Phys. Lett. A*, 10(19):1409–1418, 1995. doi:10.1142/S0217732395001526.
- [Kau83] Louis H. Kauffman. *Formal knot theory*, volume 30 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1983.
- [Kau87] Louis H. Kauffman. State models and the Jones polynomial. *Topology*, 26(3):395–407, 1987. doi:10.1016/0040-9383(87)90009-7.
- [KC02] Victor Kac and Pokman Cheung. *Quantum calculus*. Universitext. Springer-Verlag, New York, 2002. doi:10.1007/978-1-4613-0071-7.
- [Kho00] Mikhail Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000. doi:10.1215/S0012-7094-00-10131-7.
- [KR08] Mikhail Khovanov and Lev Rozansky. Matrix factorizations and link homology. *Fund. Math.*, 199(1):1–91, 2008. doi:10.4064/fm199-1-1.
- [KR18] Mikhail Khovanov and Louis-Hadrien Robert. Foam evaluation and Kronheimer–Mrowka theories, 2018. arXiv:1808.09662.
- [Kup96] Greg Kuperberg. Spiders for rank 2 Lie algebras. *Comm. Math. Phys.*, 180(1):109–151, 1996. URL: <http://projecteuclid.org/getRecord?id=euclid.cmp/1104287237>.
- [MM01] Hitoshi Murakami and Jun Murakami. The colored Jones polynomials and the simplicial volume of a knot. *Acta Math.*, 186(1):85–104, 2001. doi:10.1007/BF02392716.
- [MOST07] Ciprian Manolescu, Peter Ozsváth, Zoltán Szabó, and Dylan Thurston. On combinatorial link Floer homology. *Geom. Topol.*, 11:2339–2412, 2007. doi:10.2140/gt.2007.11.2339.
- [MOY98] Hitoshi Murakami, Tomotada Ohtsuki, and Shuji Yamada. Homfly polynomial via an invariant of colored plane graphs. *Enseign. Math. (2)*, 44(3-4):325–360, 1998.
- [MS09] Volodymyr Mazorchuk and Catharina Stroppel. A combinatorial approach to functorial quantum sl_k knot invariants. *Am. J. Math.*, 131(6):1679–1713, 2009. doi:10.1353/ajm.0.0082.
- [MSV09] Marco Mackaay, Marko Stošić, and Pedro Vaz. $sl(N)$ -link homology ($N \geq 4$) using foams and the Kapustin–Li formula. *Geom. Topol.*, 13(2):1075–1128, 2009. doi:10.2140/gt.2009.13.1075.
- [OS04] Peter Ozsváth and Zoltán Szabó. Holomorphic disks and knot invariants. *Adv. Math.*, 186(1):58–116, 2004. doi:10.1016/j.aim.2003.05.001.
- [OS09] Peter Ozsváth and Zoltán Szabó. A cube of resolutions for knot Floer homology. *J. Topol.*, 2(4):865–910, 2009. doi:10.1112/jtopol/jtp032.
- [QRS18] Hoel Queffelec, David E. V. Rose, and Antonio Sartori. Annular evaluation and link homology, 2018. arXiv:1802.04131.
- [QS15] Hoel Queffelec and Antonio Sartori. A note on the $gl(m|n)$ link invariants and the HOMFLY-PT polynomial, 2015. arXiv:1506.03329.
- [Ras03] Jacob A. Rasmussen. *Floer homology and knot complements*. ProQuest LLC, Ann Arbor, MI, 2003. Thesis (Ph.D.)–Harvard University. URL: http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&
- [RW17] Louis-Hadrien Robert and Emmanuel Wagner. A closed formula for the evaluation of sl_n -foams. To appear in *Quantum Topology*, 2017. arXiv:1702.04140.
- [RW18] Louis-Hadrien Robert and Emmanuel Wagner. Symmetric Khovanov–Rozansky link homologies, 2018. arXiv:1801.02244.
- [RW19] Louis-Hadrien Robert and Emmanuel Wagner. A quantum categorification of the Alexander polynomial, 2019. arXiv:1902.05648.
- [Sar15] Antonio Sartori. The Alexander polynomial as quantum invariant of links. *Ark. Mat.*, 53(1):177–202, 2015. doi:10.1007/s11512-014-0196-5.
- [Sus07] Joshua Sussan. *Category O and $sl(k)$ link invariants*. ProQuest LLC, Ann Arbor, MI, 2007. Thesis (Ph.D.)–Yale University. URL: http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&
- [SW17] Marko Stošić and Paul Wedrich. Rational links and DT invariants of quivers, 2017. arXiv:1711.03333.
- [Tur91] Vladimir G. Turaev. State sum models in low-dimensional topology. In *Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990)*, pages 689–698. Math. Soc. Japan, Tokyo, 1991.

- [Tur94] V. G. Turaev. *Quantum invariants of knots and 3-manifolds*, volume 18 of *De Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1994.
- [TVW17] Daniel Tubbenhauer, Pedro Vaz, and Paul Wedrich. Super q -Howe duality and web categories. *Algebr. Geom. Topol.*, 17(6):3703–3749, 2017. doi:10.2140/agt.2017.17.3703.
- [Vir06] Oleg Ya. Viro. Quantum relatives of the Alexander polynomial. *Algebra i Analiz*, 18(3):63–157, 2006. doi:10.1090/S1061-0022-07-00956-9.
- [Wit89] Edward Witten. Quantum field theory and the Jones polynomial. *Comm. Math. Phys.*, 121(3):351–399, 1989.
- [Wu14] Hao Wu. A colored $\mathfrak{sl}(N)$ homology for links in S^3 . *Diss. Math.*, 499:215 p., 2014. doi:10.4064/dm499-0-1.
- [Yon11] Yasuyoshi Yonezawa. Quantum $(\mathfrak{sl}_n, \wedge v_n)$ link invariant and matrix factorizations. *Nagoya Math. J.*, 204:69–123, 12 2011. doi:10.1215/00277630-1431840.

UNIVERSITÉ DE GENÈVE, RUE DU LIÈVRE 2–4, 1227 GENÈVE, SWITZERLAND
E-mail address: louis-hadrien.robert@unige.ch

UNIV PARIS DIDEROT, IMJ-PRG, UMR 7586 CNRS, F-75013, PARIS, FRANCE
E-mail address: emmanuel.wagner@imj-prg.fr