



**HAL**  
open science

## Stereo- and Enantioselective Synthesis of Acetylenic 2-Amino-1,3-diol Stereotriads

Arnaud Voituriez, Alejandro Pérez-Luna, Franck Ferreira, Candice Botuha,  
Fabrice Chemla

► **To cite this version:**

Arnaud Voituriez, Alejandro Pérez-Luna, Franck Ferreira, Candice Botuha, Fabrice Chemla. Stereo- and Enantioselective Synthesis of Acetylenic 2-Amino-1,3-diol Stereotriads. *Organic Letters*, 2009, 11 (4), pp.931-934. 10.1021/ol802912f. hal-03883244

**HAL Id: hal-03883244**

**<https://hal.science/hal-03883244v1>**

Submitted on 3 Jan 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Stereo- and Enantioselective Synthesis of Acetylenic 2-Amino-1,3-Diol Stereotriads


Arnaud Voituriez, Alejandro Pérez-Luna, Franck Ferreira, Candice Botuha and Fabrice Chemla\*

UPMC Univ Paris 06, UMR7611, Laboratoire de Chimie Organique, Institut de Chimie Moléculaire (FR2769), case 183, 4 place Jussieu, F-75252 Paris Cedex 05, France.

fabrice.chemla@upmc.fr

Received Date (will be automatically inserted after manuscript is accepted)

## ABSTRACT



The high-yielding and highly efficient stereoselective synthesis of enantiopure *anti,anti* and *syn,anti* acetylenic 2-amino-1,3-diol stereotriads from  $\alpha$ -alkoxy-*tert*-butanesulfinylimines bearing a stereocenter  $\alpha$  to the imino group is reported. The stereoselectivity of the reaction of these *t*BS-imines with allenylzinc ( $\pm$ )-1 was found to be controlled only by the configuration of the *tert*-butanesulfinyl group. An excellent kinetic resolution of the racemic allenylzinc species was observed, allowing a high stereocontrol whatever the configuration or the protecting group of the  $\alpha$ -alkoxy group.

2-Amino-1,3-diol subunits constitute an important stereotriad pattern found in many natural molecules of biological interest, such as indolizidine and pyrrolizidine alkaloids. A straightforward synthetic approach to these alkaloids could involve the construction of the bicyclic skeleton from properly functionalized enantiopure acetylenic 2-amino-1,3-diols presenting an *anti,anti* or a *syn,anti* relationship (Figure 1). In this context we have recently shown<sup>1</sup> that allenylzinc species are highly valuable reagents to prepare acetylenic *anti* or *syn*  $\beta$ -aminoalcohols<sup>2</sup>




Figure 1. Acetylenic 2-amino-1,3-diol stereotriads

<sup>1</sup> Botuha, C.; Chemla, F.; Ferreira, F.; Pérez-Luna, A.; Roy, B. *New J. Chem.* **2007**, *31*, 1552.


<sup>2</sup> (a) Palais, L.; Chemla, F.; Ferreira, F. *Synlett* **2006**, 1039; (b) Chemla, F.; Ferreira, F.; Gaucher, X.; Palais, L. *Synthesis* **2007**, 1235.

(from the corresponding acetylenic aziridines<sup>3,4,5</sup>) and *anti* or *syn* acetylenic aminoethers.<sup>2,6</sup>

The reaction of reagent ( $\pm$ )-**1** derived from 3-methoxymethyl-1-trimethylsilyl-1-propyne with enantiopure *t*BS-imines gives the corresponding enantiopure acetylenic *anti* aminoethers **2** (Scheme 1).<sup>2b,7</sup> The key to these results is the highly efficient kinetic resolution of the racemic allenylzinc reagent brought about by the excellent stereocontrol of the *tert*-butanesulfinyl moiety for the addition onto the imine. The synthetic potential of this approach has been exploited recently for the enantiocontrolled synthesis of (-)- $\alpha$ -conhydrine<sup>8</sup> and for the formal total synthesis of the two quinolizidinyl alkaloids (-)-epiquinamide and (-)-homopumiliotoxine 223G.<sup>9</sup> In order to obtain acetylenic 2-amino-1,3-diols and thus access to the corresponding (O,N,O) stereotriads, we envisioned to apply this method to  $\alpha$ -alkoxy *t*BS-imines **3** (Scheme 1). However, this approach raises to several important difficulties which have to be examined. The presence of the oxygen functionality in the imine electrophile could modify the intrinsic stereoinduction previously observed from the *tert*butanesulfinyl group in such reactions. Another issue to be considered is the influence of the new stereocenter  $\alpha$  to the imine on the stereofacial selectivity compared to the sulfinyl chiral auxiliary. A Cram-chelate versus a Felkin-Anh model could presumably be involved. Finally the introduction of this oxygen functionality  $\alpha$  to the imine group could possibly modify the kinetic resolution efficiency of the starting allenylzinc ( $\pm$ )-**1**. Prompted by a recent report<sup>10</sup> on the addition of enolates on  $\alpha,\beta$ -dialkoxy *t*BS-imines for the assembly of the polyhydroxylated aminoacid constituents of microsclerodermins, we disclose hereafter our results concerning the reaction of allenylzinc reagent ( $\pm$ )-**1** with stereochemically pure chiral  $\alpha$ -alkoxy *t*BS-imines.


The  $\alpha$ -alkoxy *t*BS-imines necessary for our study were prepared by the reaction of the corresponding  $\alpha$ -alkoxy aldehydes with enantiopure *tert*-butanesulfinamide<sup>11</sup> under standard conditions.<sup>12,13</sup>

**Scheme 1.** Synthesis of acetylenic aminoethers and approach to acetylenic stereotriads



Following our previously reported kinetic resolution conditions (( $\pm$ )-**1**, 4 equiv., -80 °C),<sup>4</sup> the reaction of racemic allenylzinc ( $\pm$ )-**1** with enantiopure *t*BS-imines (*S,R<sub>S</sub>*)-**3a** and (*S,R<sub>S</sub>*)-**3b** proceeded in high yield and with high stereocontrol, as only one diastereomer **4a** (from **3a**) and **4b** (from **3b**) was obtained after hydrolysis. The relative and absolute configurations of the newly created stereocenters were determined unambiguously by various selective transformations into known compounds or oxazolidinones (see below). From all the correlation experiments we could determine that allenylzinc ( $\pm$ )-**1** reacts with imines (*S,R<sub>S</sub>*)-**3a,b** from their *Re* face to give with a high stereoselectivity the corresponding protected acetylenic 2-amino-1,3-diols presenting an *anti,anti* relationship.

**Scheme 2.** Reactions with imines (*S,R<sub>S</sub>*)-**3a,b**



<sup>3</sup> (a) Bernard, N.; Chemla, F.; Normant, J. *Tetrahedron Lett.* **1999**, *40*, 75; (b) Chemla, F.; Bernard, N.; Ferreira, F.; Normant, J.-F. *Eur. J. Org. Chem.* **2001**, 3295; (c) Chemla, F.; Hebbe, V.; Normant, J.-F. *Tetrahedron Lett.* **1999**, *40*, 8093; (d) Chemla, F.; Ferreira, F.; Hebbe, V.; Stercklen, E. *Eur. J. Org. Chem.* **2002**, 1385; (e) Ferreira, F.; Bejjani, J.; Denichoux, A.; Chemla, F. *Synlett* **2004**, 2051.

<sup>4</sup> (a) Chemla, F.; Ferreira, F. *Synlett* **2004**, 983; (b) Chemla, F.; Ferreira, F. *J. Org. Chem.* **2004**, *69*, 8244.

<sup>5</sup> Ferreira, F.; Audouin, M.; Chemla, F. *Chem. Eur. J.* **2005**, *11*, 5269.

<sup>6</sup> Roy, B.; Pérez-Luna, A.; Ferreira, F.; Botuha, C.; Chemla, F. *Tetrahedron Lett.* **2008**, *49*, 1534.

<sup>7</sup> Chemla, F.; Ferreira, F. *Synlett* **2006**, 2613.

<sup>8</sup> Voituriez, A.; Ferreira, F.; Chemla, F. *J. Org. Chem.* **2007**, *72*, 5358.

<sup>9</sup> Voituriez, A.; Ferreira, F.; Perez-Luna, A.; Chemla, F. *Org. Lett.* **2007**, *9*, 4705.

<sup>10</sup> Hjelmggaard, T.; Faure, S.; Lemoine, P.; Viossat, B.; Aitken, D. J. *Org. Lett.* **2008**, *10*, 841.


<sup>11</sup> (a) Weix, D. J.; Ellman, J. A. *Org. Lett.* **2003**, *5*, 1317; (b) Weix, D. J.; Ellman, J. A. *Org. Synth.* **2005**, *82*, 157.

<sup>12</sup> Evans, J. W.; Ellman, J. A. *J. Org. Chem.* **2003**, *68*, 9948.

<sup>13</sup> Liu, G.; Cogan, D. A.; Owens, T. D.; Tang, T. P.; Ellman, J. A. *J. Org. Chem.* **1999**, *64*, 1278.

The reactivity of the *t*BS-imines ( $S,S_S$ )-**3c,d** was then examined. A difference in the diastereoselectivity of the reaction could be expected, as these imines are diastereomeric to the imines **3a,b** previously used and a matched / mismatched effect might thus be expected. However, to our delight, here again, under kinetic resolution conditions, an excellent stereoselectivity in the reaction with ( $\pm$ )-**1** (4 equiv.) was observed as only one diastereomer **4c** (resp. **4d**) was obtained from imine **3c** (resp. **3d**, Scheme 3), both presenting a *syn,anti* relationship. Allenylzinc ( $\pm$ )-**1** reacts therefore with imines ( $S,S_S$ )-**3c,d** from their *Si* face to give with high stereoselectivity the corresponding protected acetylenic 2-amino-1,3-diols presenting a *syn,anti* relationship.

Scheme 3. Reactions with imines ( $S,S_S$ )-**3c, d**



The relative and absolute configurations of the stereocenters in **4a-d** were determined unambiguously after selective transformations<sup>14</sup> into the oxazolidinones **6a-c** (Figure 2) and <sup>1</sup>H NMR coupling constant measurements. The multistep transformation of compound **4a** into a known compound<sup>15</sup> was also performed.





Figure 2. Oxazolidinones **6a-c** for stereochemical correlations.

The results reported above indicate that a high stereoselection can be expected from the reaction of  $\alpha$ -alkoxy *t*BS imines with allenylzinc ( $\pm$ )-**1**. To generalize this statement, we then tried the reaction of the two

diastereomeric imines ( $R,S_S$ )-**3e** and ( $S,S_S$ )-**3f** derived from (+)- and (-)-mandelic acid respectively. The reaction occurred smoothly under the same conditions as reported above and here again we obtained only one diastereomer **4e** and **4f** respectively (scheme 4).

Scheme 4. Reaction with imines ( $R,S_S$ )-**3e** and ( $S,S_S$ )-**3f**



From structural determinations (analogous to those reported in the Figure 2)<sup>14</sup> it could be unequivocally concluded that the reaction of ( $\pm$ )-**1** with imines ( $R,S_S$ )-**3e** and ( $S,S_S$ )-**3f** occurred from the *Si* face to give stereoselectively the corresponding *anti,anti* or *syn,anti* acetylenic stereotriad.

The addition of organometallic reagents onto *t*BS-imines has been largely studied<sup>16</sup> and various models have been proposed to rationalize the stereoselectivity of such processes. More specifically, the influence of an oxygen or heteroatom functionality in the imine electrophile is a matter of interest since these groups can change the stereoselection from the *tert*butanesulfinyl group, as reported for example by Ellman who observed a reversal of the stereofacial selectivity from simple *t*BS-imines<sup>17</sup> to  $\alpha$ -alkoxy *t*BS-imines.<sup>18</sup> The origins for the inversion of the facial selectivity compared to simple alkyl *t*BS-imines remain under discussion and models involving *E/Z* isomerization of imines as well as chelate versus open transition states have been proposed to account for the atypical stereochemical behaviour. Moreover, it should be noted that this inversion in stereoselection does not seem general: it has been observed for *t*BS-imines derived from

<sup>16</sup> For a general review on chiral sulfinimines, see: (a) Morton, D.; Stockman, R. A. *Tetrahedron* **2006**, *62*, 8869; (b) Ferreira, F.; Botuha, C.; Chemla, F.; Pérez-Luna, A. *Chem. Soc. Rev.* **2009**, DOI:10.1039/B809772K

<sup>17</sup> Cogan, D. A.; Liu, G.; Ellman, J. *Tetrahedron* **1999**, *55*, 8883.

<sup>18</sup> (a) Barrow, J. C.; Ngo, P. L.; Pellicore, J. M.; Selnick, H. G.; Nantermet, P. G. *Tetrahedron Lett.* **2001**, *42*, 2051; (b) Tang, T. P.; Volkman, S. K.; Ellman, J. A. *J. Org. Chem.* **2001**, *66*, 8772.

<sup>14</sup> see Supporting Information.

<sup>15</sup> Poisson, J. F.; Normant, J.-F. *Org. Lett.* **2001**, *3*, 1889.


aldehydes bearing an  $\alpha$ -alkoxy<sup>10,18</sup> or an  $\alpha$ -chloro group,<sup>19</sup> independently of the organometallic species used (alkyl organometallics or enolates). By contrast it has not been observed in the case of *t*BS-imines derived from aldehydes bearing an  $\alpha$ -amino group,<sup>20</sup> nor in the case of *t*BS-imines derived from ketones bearing an  $\alpha$ -alkoxy group.<sup>18b</sup> The reduction of *t*BS-imines derived from ketones bearing an  $\alpha$ -chloro group proceeds with face selectivities depending on the nature of the reducing agent.<sup>21</sup> In our case no inversion of the stereochemical induction from the *t*BS group was noted, as the facial selectivity was the same in all cases we studied: imines (*S,R<sub>S</sub>*)-**3a** and (*S,R<sub>S</sub>*)-**3b** presenting a (*R*) configuration at the sulfur center reacted with ( $\pm$ )-**1** on their *Re* face, whereas products arising from the addition of imines (*S,S<sub>S</sub>*)-**3c**, (*S,S<sub>S</sub>*)-**3d**, (*R,S<sub>S</sub>*)-**3e** and (*S,S<sub>S</sub>*)-**3f** presenting a (*S*) configuration at the sulfur center were obtained through reaction on the *Si* face.

Another point of interest concerns the relative influence on the stereofacial selectivity of the stereocenter  $\alpha$  to the imine compared to the sulfinyl chiral auxiliary. A Cram-chelate versus a Felkin-Anh model could presumably be involved, and the two possibilities have also been reported by Ellman.<sup>12</sup> Former reports dealing with organometallic species addition on  $\alpha$ -alkoxy-<sup>10,12</sup> or  $\alpha$ -amino-*t*BS-imines<sup>20</sup> bearing a stereocenter  $\alpha$  to the imino group showed that generally the stereoselectivity is mainly defined by the sulfinyl moiety and not by the  $\alpha$  stereocenter, although in some cases a drop in selectivity was observed.<sup>12</sup> In our case the stereoselectivity is always excellent, and independent of the  $\alpha$  stereocenter configuration, whatever the protecting group (TBS or Bn) on the oxygen atom. Though the stereoselectivity we obtained in the case of imines (*S,R<sub>S</sub>*)-**3a**, (*S,R<sub>S</sub>*)-**3b** and (*R,S<sub>S</sub>*)-**3e** is consistent with a Felkin-type control, in the case of imines (*S,S<sub>S</sub>*)-**3c**, (*S,S<sub>S</sub>*)-**3d** and (*S,S<sub>S</sub>*)-**3f**, anti-Felkin products are observed.


Finally, it should be noted that we have always used *racemic* allenylzinc ( $\pm$ )-**1**. The excellent selectivities we obtained in all cases can only be explained by a very efficient kinetic resolution: only one enantiomer of the racemic mixture in **1** is allowed to react with the enantiopure sulfinimines. We propose the Felkin-type transition state TS1 depicted in Scheme 5 to explain the stereoselectivities observed in the reactions with (*S,R<sub>S</sub>*)-**3a** and (*S,R<sub>S</sub>*)-**3b**. In this transition state the *t*BS-imine is adopting its ground-state conformation where the S-O bond and the lone pair on the nitrogen atom are antiperiplanar.<sup>5,22</sup> The reaction of imine (*R,S<sub>S</sub>*)-**3e** to afford **4e** presumably follows a transition state enantiomeric to TS1. On the other hand, the anti-Felkin-type transition state TS2 is proposed for the reactions of imines (*S,S<sub>S</sub>*)-**3c**, (*S,S<sub>S</sub>*)-**3d** and (*S,S<sub>S</sub>*)-**3f**.

In each case only the matched reacting enantiomer of ( $\pm$ )-**1** is represented.

**Scheme 5.** Plausible transition states TS1 and TS2.



TS1 (Felkin) : R = TBS, Bn



TS2 (anti-Felkin) : R<sup>1</sup> = TBS, Bn; R<sup>2</sup> = Me, Ph

In conclusion, we have disclosed a high-yielding and highly efficient stereoselective synthesis of enantiopure *anti,anti* and *syn,anti* acetylenic 2-amino-1,3-diol stereotriads from  $\alpha$ -alkoxy-*t*BS-imines bearing a stereocenter  $\alpha$  to the imino group. The stereoselectivity was found to be controlled only by the configuration of the *tert*-butanesulfinyl group and an excellent kinetic resolution of racemic allenylzinc ( $\pm$ )-**1** was observed, whatever the configuration or the protecting group of the  $\alpha$ -alkoxy substituent. These acetylenic stereotriads should be highly versatile stereodefined building blocks for the total synthesis of indolizidine or quinolizidine alkaloids. Further studies in this area from our lab will be reported in due course.

**Acknowledgment** Thanks are due to Patrick Herson for single crystal X-ray analysis and to the CNRS for a postdoctoral fellowship grant (to AV).

**Supporting Information Available** Preparation and characterization data for imines **3e** and **3f**, acetylenic aminoethers **4a-f** and chemical correlations into known compounds **5a-c** and 2-oxazolidin-3-ones **6a-e**, including <sup>1</sup>H and <sup>13</sup>C data. This material is available free of charge via the Internet from <http://pubs.acs.org>

<sup>19</sup> (a) Denolf, B.; Mangelinckx, S.; Tornoos, K. W.; De Kimpe, N. *Org. Lett.* **2006**, *8*, 3129; (d) Hodgson, D. M.; Kloesges, J.; Evans, B. *Org. Lett.*, **2008**, *10*, 2781.

<sup>20</sup> (a) Prakash, G. K. S.; Mandal, M. *J. Am. Chem. Soc.* **2002**, *124*, 6538; (b) Liu, J.; Li, Y.; Hu, J. *J. Org. Chem.* **2007**, *72*, 3119.

<sup>21</sup> (a) Denolf, B.; Leemans, E.; De Kimpe, N. *J. Org. Chem.* **2007**, *72*, 3211; (b) Denolf, B.; Leemans, E.; De Kimpe, N. *J. Org. Chem.* **2008**, *73*, 5662.

<sup>22</sup> (a) Bharatam, P. V.; Uppal, P.; Amita; Kaur, D. *J. Chem. Soc., Perkin Trans. 2*, **2000**, 43; (b) Bharatam, P. V.; Amita; Kaur, D. *J. Phys. Org. Chem.*, **2002**, *15*, 197.