



An algorithm based on the postoperative decrease of albumin (Delta Alb) to anticipate complications after liver surgery

Ismail Labgaa, Luis Cano, Orsalia Mangana, Gaetan-Romain Joliat, Emmanuel Melloul, Nermin Halkic, Markus Schafer, Eric Vibert, Nicolas Demartines, Nicolas Golse, et al.

► To cite this version:

Ismail Labgaa, Luis Cano, Orsalia Mangana, Gaetan-Romain Joliat, Emmanuel Melloul, et al.. An algorithm based on the postoperative decrease of albumin (Delta Alb) to anticipate complications after liver surgery. *Perioperative Medicine*, 2022, 11 (1), pp.53. <10.1186/s13741-022-00285-w>. <hal-03886022>

HAL Id: hal-03886022

<https://hal.science/hal-03886022v1>

Submitted on 6 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

RESEARCH

Open Access



An algorithm based on the postoperative decrease of albumin (ΔAlb) to anticipate complications after liver surgery

Ismail Labgaa^{1*†}, Luis Cano^{2†}, Orsalia Mangana³, Gaëtan-Romain Joliat¹, Emmanuel Melloul¹, Nermin Halkic¹, Markus Schäfer¹, Eric Vibert^{3,4}, Nicolas Demartines^{1*}, Nicolas Golse^{3,4} and Martin Hübner¹

Abstract

Background: Perioperative decrease of albumin (ΔAlb) appeared as a promising predictor of complications after digestive surgery, but its role after liver surgery remains unclear. This study aimed to analyze whether and how ΔAlb can be used to predict complications after liver surgery.

Methods: A bicentric retrospective analysis of patients undergoing liver surgery (2010–2016) was performed, following TRIPOD guidelines. The preoperative and postoperative difference of albumin was calculated on POD 0 and defined as ΔAlb . Patients with any missing variable were excluded. The primary endpoint was overall complications according to the Clavien classification. A multiparametric algorithm based on ΔAlb was generated to optimize prediction performance.

Results: A total of 110 patients were analyzed. At least one complication occurred in 66 (60%) patients. Patients with and without complication showed a ΔAlb of 15.8 vs. 9.5 g/L ($p < 0.001$). Area under ROC curve (AUC) of ΔAlb was 0.75 ($p < 0.01$). The ΔAlb -based algorithm showed an AUC of 0.84 ($p < 0.01$), significantly improving performance ($p = 0.03$). Multivariable analysis identified ΔAlb as independent predictor of complications (HR, 1.12; 95% CI, 1.01–1.07; $p = 0.002$).

Conclusions: ΔAlb appeared as a promising predictor independently associated with the risk of complication after liver surgery. The study presents a novel decision-tree based on ΔAlb to anticipate complications.

Keywords: Partial hepatectomy, Liver resection, Decision-making, Morbidity, Biomarkers, Predictors

Background

Liver surgery entails major surgical procedures that trigger intensive physiological stress. Mortality and morbidity may reach 10% and 45%, respectively (Jarnagin et al. 2002; Vonlanthen et al. 2011; Breitenstein et al. 2010;

Labgaa et al. 2019; Filmann et al. 2019). Reducing the incidence of adverse outcomes is challenging, and early detection is paramount. Identification of reliable tools capable to predict and anticipate postoperative complications is the first step. Ideal candidate predictors must be performant, easy to measure and inexpensive, but also importantly, must be early indicative (Labgaa et al. 2017a, b; Longchamp et al. 2021).

The perioperative variation of albumin (ΔAlb) is a novel marker that showed promising results in digestive surgery (Labgaa et al. 2016a, 2017c, 2021; Muller et al. 2018). Unlike preoperative single values of serum albumin, it

[†]Ismail Labgaa and Luis Cano contributed equally to this work.

*Correspondence: ismail.labgaa@chuv.ch; Demartines@chuv.ch

¹ Department of Visceral Surgery, Lausanne University Hospital CHUV, University of Lausanne (UNIL), Rue du Bugnon 46, CH-1011 Lausanne, Switzerland

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

has not been extensively investigated. To date, less than 20 studies explored ΔAlb in abdominal surgery (Joliat et al. 2022). In major abdominal surgery, $\Delta\text{Alb} \geq 10$ g/L was identified as independent predictor of complications (Labgaa et al. 2017c). Müller et al. showed that ΔAlb was better prognostic marker than absolute albumin value in a cohort of patients undergoing intestinal resection for Crohn's disease (Muller et al. 2018). Nonetheless, the potential role of ΔAlb to predict complications after liver surgery remains unknown. Aside orthotopic liver transplant (OLT), data on ΔAlb remain particularly scarce for liver surgery patients. In 2016, a pilot study from our group showed that ΔAlb was an independent predictor of complication (Labgaa et al. 2016a). However, this study did not provide a specific cutoff. These results needed to be confirmed with a larger sample size and an external dataset, allowing to also establish a threshold for ΔAlb .

The present study aimed to assess the predictive value of ΔAlb for adverse outcomes and to define how it may help anticipating postoperative complications in patients undergoing liver surgery.

Methods

Design

This bicentric retrospective study was conducted in the departments of surgery at Lausanne University Hospital (CHUV) and at Paul-Brousse Hospital in Paris, between 2010 and 2016, following TRIPOD guidelines. The study was conducted in accordance with the Declaration of Helsinki, and it conformed to good clinical practice guidelines. Informed consent was obtained from all included patients. The protocol was approved by the Institutional Review Boards at University Hospital of Lausanne (CER-VD # 2017-01169) and at Paul-Brousse Hospital (CO 16-006).

Patients

Consecutive patients >18 years undergoing liver surgery during the study period were included. Very stringent selection criteria were applied: patients with any missing variable were excluded. Collected data included demographics, surgical, and perioperative variables as well as postoperative outcomes. Data were retrieved from institutional databases.

ΔAlb

Blood samples were collected by nurses preoperatively and postoperatively, allowing to obtain daily measures of serum albumin concentration [g/L]. Perioperative variation of albumin was defined as ΔAlb and calculated according to the following formula: preoperative albumin value - albumin value on POD 0 (4–8 h after the end of surgery). Timepoints were selected to tailor a predictor

being indicative as early as possible (POD 0), as well as based on data from a previous pilot study (Labgaa et al. 2016a).

Endpoints

Postoperative morbidity was graded according to the Clavien classification (Dindo et al. 2004). Minor and major complications were defined as grades I–II and III–IV, respectively. Postoperative mortality was defined as grade V. The primary endpoint was overall morbidity while secondary endpoints were Comprehensive Complication Index (CCI) (Slankamenac et al. 2013) and postoperative length of stay (LoS) (Labgaa et al. 2016b; Joliat et al. 2016).

Statistical analyses

All statistical analyses were conducted by a statistician (L.C.). Complete case study technique and bootstrap methods (1000 repetitions) were conducted to handle missing data. Single imputation method was avoided due to the risk of bias in small populations (Dziura et al. 2013; Zhang et al. 2017; Jakobsen et al. 2017).

Continuous variables were provided as median with interquartile range (IQR) or as mean with standard deviations (SD), according to their distribution pattern. They were compared with Student's *t* test or Mann-Whitney *U* test. Categorical data were provided as frequencies with percentages and compared with either χ^2 or Fisher's exact test. Receiver operating curve was generated, allowing to calculate area under the curve (AUC). Uni- and multivariable analyses were performed. Variables with $p < 0.1$ on univariable analyses were further considered in the multivariable analysis, integrating multiple potential confounding factors into the model. To increase the accuracy of variables selection, a stepwise regression in both senses was used (forward and backward). The best model was selected based on AUC values. A decision-tree was generated to optimize prediction performance. Its number of nodes were minimized on purpose, to facilitate clinical implementation. Hence, the model integrated the strongest predictors identified on univariate analyses ($p < 0.001$). A bootstrap technique with 1000 repetitions was conducted to overcome the relatively small sample size. Standard error and cross-validation error values were analyzed, in order to determine the optimal cutoff points, preserving as much accuracy as possible. DeLong test was used to compare AUC curves. The statistical significance was defined as a *p* value < 0.05 . The statistical analyses were performed with the statistical software R 4.0.1 (Core Team 2020), and the following packages were used to develop the statistical analysis: decision trees: part (4.1–15, Terry Therneau and Beth Atkinson (2019)), roc curves: ROCR (1.0–11, Sing T, Sander O,

Beerenwinkel N, Lengauer T (2005)), data manipulation: dplyr (0.8.5, Hadley Wickham, Romain François), and statistical modeling: caret (6.0.86, Max Kuhn (2020)).

Results

Patients and outcomes

After exclusion of patients with any missing variable, a total of 110 patients were included in the final analyses. Patients' characteristics and details of surgery are provided in Table 1. Regarding outcomes: minor, major, and overall complications were reported in 59 (53.6 %), 22 (20 %), and 66 (60 %) patients, respectively. Postoperative mortality was reported in 1 patient (0.9%). Median CCI was 19.14 (0–100) while median LoS was 14.3 days (3–67). Characteristics of the final cohort were compared with the excluded patients in Supplementary Table 1.

Table 1 Patients characteristics and details of surgery

	Value Mean (±SD) or n (%)
Age (years)	61.37 (10.48)
Gender (female)	39 (35.5 %)
Smoking	30 (27.7 %)
Diabetes	87 (79.1 %)
Cirrhosis	9 (8.2 %)
Cancer	85 (77.3 %)
HCC	12 (10.9 %)
CCa	14 (12.7 %)
CRLM	46 (41.8 %)
Other malignancy	13 (11.8 %)
Non-cancer	25 (22.7%)
Alveolar echinococcosis	10 (9.1%)
Cystic echinococcosis	2 (1.8%)
Adenoma	2 (1.8%)
Hemangioma	2 (1.8%)
Others	9 (8.2%)
BMI (kg/m ²)	26.51 (4.18)
Major resection (≥3 segments)	52 (47.3 %)
Laparoscopic approach	16 (14.5 %)
Surgery duration (min)	324.55 (132.55)
Blood loss (mL)	985 (956)
Complications (Clavien classification)	
Grade I	28 (25.5 %)
Grade II	48 (43.6 %)
Grade IIIa	15 (13.6 %)
Grade IIIb	10 (9.1 %)
Grade IVa	4 (3.6 %)
Grade IVb	3 (2.7 %)
Grade V	1 (0.9 %)

Abbreviations: BMI body mass index, HCC hepatocellular carcinoma, CCa cholangiocarcinoma, CRLM colorectal liver metastases

Perioperative profile of albumin and ΔAlb

Liver surgery induced a sudden decrease of albumin on POD 0, followed by a plateau phase (Fig. 1A). Although showing a similar pattern, patients who developed complication (of any grade) showed more accentuated albumin drop, compared to those without complication. Median ΔAlb was 9.5 g/L (2–30) without and 15.8 g/L (1.9–26) with complications ($p < 0.001$) (Fig. 1B). For the whole cohort, median ΔAlb was 12.5 g/L (2–30). The correlation analysis was conducted to investigate the impact of intraoperative fluid perfusion on ΔAlb, showing a low correlation coefficient (Pearson=0.341, $p = 0.001$), suggesting that the volume of fluid has a marginal effect on albumin drop (Supplementary Fig. 1).

AUC, cutoff of ΔAlb, and its ability to discriminate adverse outcomes


ROC curves of preoperative albumin and ΔAlb showed an AUC of 0.63 ($p < 0.01$.) and 0.75 ($p < 0.01$), respectively (Fig. 2A); ΔAlb curve revealed a threshold of 8.7 g/L. This yielded a sensitivity of 61.4%, specificity of 84.9%, positive predictive value of 73 %, and negative predictive value of 76.7%. It was tested whether this cutoff could discriminate patients with adverse outcomes. Minor, major, and overall complications were more prevalent in patients with ΔAlb ≥ 8.7 g/L (Fig. 2B). These patients also showed increased CCI of 20.90 compared to 7.53, $p < 0.001$ (Fig. 2C). Likewise, LoS was also prolonged in this group of patients (16.1 vs. 7 days, $p < 0.001$) (Fig. 2D).

Multivariable analysis

Univariable and multivariable analyses were performed to determine whether ΔAlb was an independent predictor of overall postoperative complications (Table 2). ΔAlb (OR, 1.12; 95% CI, 1.01–1.07; $p = 0.002$) and surgery duration (OR, 1.00; 95% CI, 1.00–1.00; $p = 0.03$) were the only factors showing a statistical significance on multivariable analysis. Preoperative value of albumin itself was not predictive in univariable analysis.

An algorithm based on ΔAlb to optimize the predictive performance

A decision-tree was generated based on the results of the univariable analyses to improve the predictive performance. The model was designed to be easy to use on purpose and to facilitate clinical implementation (Fig. 3). It integrates ΔAlb, surgery duration, and blood loss. Its performance showed an AUC of 0.84 ($p < 0.01$) (Fig. 2A), yielding a sensitivity of 70.5 %, a specificity of 86.4 %, a positive predictive value of 77.5 %, and a negative predictive value of 81.4 %. Delong test revealed a



statistically increased performance of this model compared to Δ Alb alone ($p=0.03$).


Discussion

This study adds novel data on the promising role of Δ Alb and support its use to anticipate adverse outcomes in liver surgery patients. These findings evidenced that Δ Alb was independently associated with the risk of complications and provide an easy-to-use decision-tree to predict complications after liver surgery.

As expected, liver surgery induced an abrupt fall of albumin, already on POD 0. This drop was followed by a plateau and recovery phases, which was consistent with a pilot study in liver surgery, as well as in other types of surgery (Labgaa et al. 2016a, 2017c; Ryan et al. 2007; Hubner et al. 2016). Questions are (I) what determines the amplitude of this decrease and (II) what is its clinical relevance. Regarding albumin drop, previous studies in clinical research have tried to investigate its determinants and mechanisms. Capillary leakage of albumin from plasma to tissues seemed to be a major cause (Amouzandeh et al. 2018). Dedicated investigations in basic/translational research are however needed to thoroughly decipher the precise underlying mechanisms of albumin decrease after surgery; such data as well as animal models are unfortunately not available yet. The present study was not designed to explore those elements and cannot speculate on candidate mechanisms. The amplitude of albumin decrease

seems to be clinically pertinent based on its capacity to discriminate patients who will develop adverse outcomes and independent predictive value for postoperative complications. Not only novel, the clinical impact of these results is also emphasized by the fact that Δ Alb displays all the characteristics of ideal predictor: very early indicative (POD 0), easy to measure, and inexpensive. Of note, Δ Alb was superior to preoperative value of albumin, as demonstrated by AUC and multivariable analyses; the latter showed a poor performance with an AUC of 0.63 ($p<0.01$). This is consistent with other data in colorectal surgery (Muller et al. 2018).

Similar studies have highlighted the rationale to measure perioperative albumin fluctuation in upper-GI (Ryan et al. 2007), colorectal (Muller et al. 2018; Wierdak et al. 2018), and thoracic surgery (Li et al. 2018). Beside a pilot study from our group raising the interest to monitor albumin in patients undergoing partial hepatectomy (Labgaa et al. 2016a), albumin has been more investigated in OLT. Amouzandeh et al. used OLT as a model to study albumin kinetics after major surgery, allowing to identify capillary leak as a major source of albumin decrease (Amouzandeh et al. 2018). Exploring Δ Alb impact on clinical outcomes after OLT, Hiroi et al. detected an association with Δ Alb and the need for hemodynamic support (Hiroi et al. 2019). In living donors for OLT, analyses revealed that high Δ Alb was associated with an increased risk of pleural effusion (Jeong et al. 2019). Albeit being of indisputable interest, most of these studies calculated



Δ Alb on POD2. In the present study, Δ Alb was calculated on POD 0, only 4–8 h after surgery, a striking advantage.

Limitation of this study is mainly its retrospective design. However, it is a multicenter effort, involving two tertiary centers. The relatively small sample size may also be considered as a shortcoming. However, selection was very stringent to obtain very clean data and to minimize the risk of bias. An important question is to know whether this stringent selection could have induced a bias. As showed in Supplementary Table 1, included patients only showed statistically higher values of BMI and blood loss compared to


the excluded ones. Another limitation was related to albumin perfusion. Unfortunately, these data were not available. Future studies are needed to investigate whether and how perioperative albumin perfusion may mitigate Δ Alb and improve outcomes. Dedicated studies are needed to clarify the value of albumin perfusion intra-operatively as well as postoperatively. The model displays a descent performance that might be however overfitted to this bicentric cohort. Therefore, validating this model in a future study with an independent cohort would be important. Although it would have been valuable to validate these findings in an

Table 2 Multivariable analysis for overall complications

	Univariable			Multivariable		
	HR	95% CI	p value	HR	95% CI	p value
Age (years)	1.00	0.98–1.01	0.83			
Gender (female)	0.57	0.49–1.07	0.16			
BMI (kg/m ²)	1.05	0.98–1.05	0.28			
Smoking	1.21	0.72–1.37	0.66			
Diabetes	4.04	1.12–1.59	0.01	2.63	0.88–1.57	0.14
Cirrhosis	3.28	0.6–1.48	0.98			
Cancer	0.64	0.49–1.13	0.35			
Major resection	1.39	0.80–1.41	0.39			
Surgical approach (open)	4.06	1.22–2.76	0.01	1.25	0.41–2.26	0.74
Surgery duration (min)	1.00	1.00–1.01	<0.001	1.00	1.00–1.00	0.03
Blood loss (mL)	1.00	1.00–1.00	<0.001	1.00	0.99–1.00	0.1
Preoperative albumin (g/L)	1.02	0.97–1.04	0.5			
ΔAlb (POD 0)	1.16	1.03–1.08	<0.001	1.12	1.01–1.07	0.002

Values are presented with two decimals

Abbreviations: BMI body mass index, CI confidence interval, OR odds ratio, POD postoperative days



independent cohort, one must note that the vast majority of comparable studies investigating biomarkers lacks validation cohort, upfront. Comparison with other markers can be argued but was not an objective of the present study.

ΔAlb has already been compared to other lab tests such as lactates, C-reactive protein (CRP), and procalcitonin in previous studies demonstrating a higher performance (Longchamp et al. 2021; Labgaa et al. 2016a).

In addition, the present study provided a decision-tree which is easy to use and to implement in clinical practice, offering a “precision medicine” perspective. This tool may help to tailor postoperative surveillance on an individual patient basis. As example of potential clinical application, it may help guiding decision whether to transfer patients to intensive care unit (ICU), intermediate care, or to the ward. Recent data from our group also highlighted the interest of following ERAS guidelines for liver surgery, showing a significantly lower ΔAlb in patients undergoing liver surgery with ERAS, compared to non-ERAS management (Gonvers et al. 2021). Other important potential application would be to select patients who should benefit from specific therapeutic measures which would have to be tested with randomized clinical trials.

Conclusions

These results showed an association between perioperative albumin fluctuation (ΔAlb) and the risk of postoperative complication after liver surgery, evidencing the role of ΔAlb as a predictor of adverse outcomes and the rationale to monitor it in liver surgery patients. This study also presents a novel algorithm based on ΔAlb optimizing prediction performance and helping for decision-making during the postoperative phase.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13741-022-00285-w>.

Additional file 1: Supplementary Table 1. Characteristics of the final cohort were compared with the excluded patients.

Additional file 2: Supplementary Figure 1. Intraoperative fluid [mL].

Acknowledgements

Not applicable.

Authors' contributions

Study concept and design: IL, LC, ND, EV, NG, and MH. Acquisition of data: IL, OM, GRJ, and NG. Analysis and interpretation of data: IL, LC, ND, NG, and MH. Drafting of the manuscript: IL, LC, NG, and MH. Critical revision of the manuscript for important intellectual content: IL, LC, OM, GRJ, EM, NH, MS, EV, ND, NG, and MH. The author(s) read and approved the final manuscript.

Funding

None

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The protocol was approved by the Institutional Review Boards at University Hospital of Lausanne (CER-VD # 2017-01169) and at Paul-Brousse Hospital (CO 16-006).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Visceral Surgery, Lausanne University Hospital CHUV, University of Lausanne (UNIL), Rue du Bugnon 46, CH-1011 Lausanne, Switzerland.

²Nutrition Metabolism and Cancer, INSERM, University of Rennes, INRAE, CHU Pontchaillou, UMR 1241 NUMECAN, Rennes, France. ³Department of Hepatobiliary Surgery, Paul-Brousse Hospital, Assistance Publique Hôpitaux de Paris, Centre Hépatobiliaire, 94800 Villejuif, France. ⁴Université Paris-Saclay, Inserm, Physiopathogénèse et traitement des maladies du Foie, UMR-S 1193, Paris, France.

Received: 20 May 2022 Accepted: 31 October 2022

Published online: 09 November 2022

References

- Amouzandeh M, Nowak G, Januszkiwicz A, Wernerman J, Rooyackers O, Norberg A. Albumin mass balance and kinetics in liver transplantation. *Crit Care*. 2018;22(1):152.
- Breitenstein S, DeOliveira ML, Raptis DA, Slankamenac K, Kambakamba P, Nerl J, et al. Novel and simple preoperative score predicting complications after liver resection in noncirrhotic patients. *Ann Surg*. 2010;252(5):726–34.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240(2):205–13.
- Dziura JD, Post LA, Zhao Q, Fu Z, Peduzzi P. Strategies for dealing with missing data in clinical trials: from design to analysis. *Yale J Biol Med*. 2013;86(3):343–58.
- Filmann N, Walter D, Schadde E, Bruns C, Keck T, Lang H, et al. Mortality after liver surgery in Germany. *Br J Surg*. 2019;106(11):1523–9.
- Gonvers S, Jurt J, Joliat GR, Halkic N, Melloul E, Hubner M, et al. Biological impact of an enhanced recovery after surgery programme in liver surgery. *BJS Open*. 2021;5(2):zraa015.
- Hiroi K, Matsusaki T, Kaku R, Umeda Y, Yagi T, Morimatsu H. Postoperative course of serum albumin levels and organ dysfunction after liver transplantation. *Transplant Proc*. 2019;51(8):2750–4.
- Hubner M, Mantziari S, Demartines N, Pralong F, Coti-Bertrand P, Schafer M. Postoperative albumin drop is a marker for surgical stress and a predictor for clinical outcome: a pilot study. *Gastroenterol Res Pract*. 2016;2016:8743187.
- Jakobsen JC, Gluud C, Wetterslev J, Winkel P. When and how should multiple imputation be used for handling missing data in randomised clinical trials - a practical guide with flowcharts. *BMC Med Res Methodol*. 2017;17(1):162.
- Jarnagin WR, Gonen M, Fong Y, DeMatteo RP, Ben-Porat L, Little S, et al. Improvement in perioperative outcome after hepatic resection: analysis of 1,803 consecutive cases over the past decade. *Ann Surg*. 2002;236(4):397–406 discussion -7.
- Jeong HW, Kim JW, Shin WJ, Kim SO, Moon YJ, Kwon HM, et al. Early postoperative hypoalbuminaemia is associated with pleural effusion after donor hepatectomy: a propensity score analysis of 2316 donors. *Sci Rep*. 2019;9(1):2790.
- Joliat GR, Labgaa I, Hubner M, Blanc C, Griesser AC, Schafer M, et al. Cost-benefit analysis of the implementation of an enhanced recovery program in liver surgery. *World J Surg*. 2016;40(10):2441–50.
- Joliat GR, Schoor A, Schafer M, Demartines N, Hubner M, Labgaa I. Postoperative decrease of albumin (ΔAlb) as early predictor of complications after gastrointestinal surgery: a systematic review. *Perioper Med (Lond)*. 2022;11(1):7.
- Labgaa I, Demartines N, Hubner M. Biomarkers capable to early predict postoperative complications: the grail. *Ann Surg*. 2017a;266(6):e91–e2.
- Labgaa I, Demartines N, Hubner M. Serum albumin: a promising biomarker to anticipate postoperative complications. *Dis Colon Rectum*. 2017b;60(9):e616.

- Labgaa I, Jarrar G, Joliat GR, Allemann P, Gander S, Blanc C, et al. Implementation of enhanced recovery (ERAS) in colorectal surgery has a positive impact on non-ERAS liver surgery patients. *World J Surg*. 2016b;40(5):1082–91.
- Labgaa I, Joliat GR, Demartines N, Hubner M. Serum albumin is an early predictor of complications after liver surgery. *Dig Liver Dis*. 2016a;48(5):559–61.
- Labgaa I, Joliat GR, Grass F, Jarrar G, Halkic N, Demartines N, et al. Impact of postoperative weight gain on complications after liver surgery. Oxford: HPB; 2019.
- Labgaa I, Joliat GR, Kefleyesus A, Mantziari S, Schafer M, Demartines N, et al. Is postoperative decrease of serum albumin an early predictor of complications after major abdominal surgery? A prospective cohort study in a European centre. *BMJ Open*. 2017c;7(4):e013966.
- Labgaa I, Mantziari S, Genety M, Elliott JA, Kamiya S, Kalff MC, et al. Early postoperative decrease of albumin is an independent predictor of major complications after oncological esophagectomy: a multicenter study. *J Surg Oncol*. 2021;123(2):462–9.
- Li P, Li J, Lai Y, Wang Y, Wang X, Su J, et al. Perioperative changes of serum albumin are a predictor of postoperative pulmonary complications in lung cancer patients: a retrospective cohort study. *J Thorac Dis*. 2018;10(10):5755–63.
- Longchamp G, Labgaa I, Demartines N, Joliat GR. Predictors of complications after liver surgery: a systematic review of the literature. *HPB (Oxford)*. 2021;23(5):645–55.
- Muller C, Stift A, Argeny S, Bergmann M, Gnant M, Marolt S, et al. Delta albumin is a better prognostic marker for complications following laparoscopic intestinal resection for Crohn's disease than albumin alone - a retrospective cohort study. *PLoS One*. 2018;13(11):e0206911.
- Ryan AM, Hearty A, Prichard RS, Cunningham A, Rowley SP, Reynolds JV. Association of hypoalbuminemia on the first postoperative day and complications following esophagectomy. *J Gastrointest Surg*. 2007;11(10):1355–60.
- Slankamenac K, Graf R, Barkun J, Puhon MA, Clavien PA. The comprehensive complication index: a novel continuous scale to measure surgical morbidity. *Ann Surg*. 2013;258(1):1–7.
- Vonlanthen R, Slankamenac K, Breitenstein S, Puhon MA, Muller MK, Hahnloser D, et al. The impact of complications on costs of major surgical procedures: a cost analysis of 1200 patients. *Ann Surg*. 2011;254(6):907–13.
- Wierdak M, Pisarska M, Kusnierz-Cabala B, Witowski J, Dworak J, Major P, et al. Changes in plasma albumin levels in early detection of infectious complications after laparoscopic colorectal cancer surgery with ERAS protocol. *Surg Endosc*. 2018;32(7):3225–33.
- Zhang Y, Alyass A, Vanniyasingam T, Sadeghirad B, Florez ID, Pichika SC, et al. A systematic survey of the methods literature on the reporting quality and optimal methods of handling participants with missing outcome data for continuous outcomes in randomized controlled trials. *J Clin Epidemiol*. 2017;88:67–80.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

