



**HAL**  
open science

## Temperature-mediated developmental plasticity in winter and summer larvae of *Palaemon serratus*

Jean-Baptiste Baudet, Benoît Xuereb, Aurélie Duflot, Agnès Poret, Emeline Maniez, Frank Le Foll, Romain Coulaud

► **To cite this version:**

Jean-Baptiste Baudet, Benoît Xuereb, Aurélie Duflot, Agnès Poret, Emeline Maniez, et al.. Temperature-mediated developmental plasticity in winter and summer larvae of *Palaemon serratus*. *Journal of Thermal Biology*, 2022, 110, pp.103343. 10.1016/j.jtherbio.2022.103343 . hal-03886082

**HAL Id: hal-03886082**

**<https://hal.science/hal-03886082v1>**

Submitted on 6 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Journal Pre-proof



Temperature-mediated developmental plasticity in winter and summer larvae of *Palaemon serratus*

Jean-Baptiste Baudet, Benoît Xuereb, Aurélie Duflot, Agnès Poret, Emeline Maniez, Frank Le Foll, Romain Coulaud

PII: S0306-4565(22)00157-7

DOI: <https://doi.org/10.1016/j.jtherbio.2022.103343>

Reference: TB 103343

To appear in: *Journal of Thermal Biology*

Received Date: 31 March 2022

Revised Date: 26 July 2022

Accepted Date: 20 September 2022

Please cite this article as: Baudet, J.-B., Xuereb, Benoît., Duflot, Auré., Poret, Agnès., Maniez, E., Le Foll, F., Coulaud, R., Temperature-mediated developmental plasticity in winter and summer larvae of *Palaemon serratus*, *Journal of Thermal Biology* (2022), doi: <https://doi.org/10.1016/j.jtherbio.2022.103343>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Ltd.

**Authors contributions**

**JBB** Conceptualization, Methodology, Formal analysis, Investigation, Data curation, Writing - original draft, review & editing, Visualization. **BX** Conceptualization, Methodology, Validation, Writing - review & editing, Supervision, Funding acquisition. **AD** Methodology, Investigation, Writing - review & editing. **AP** Investigation. **EM** Investigation. **FLF** Supervision. **RC** Conceptualization, Methodology, Validation, Formal analysis, Data curation, Writing - review & editing, Supervision, Project administration, Funding acquisition.

1           **Title: Temperature-mediated developmental plasticity in winter and**  
2 **summer larvae of *Palaemon serratus***

3  
4           **Authors:** Jean-Baptiste Baudet<sup>1</sup>, Benoît Xuereb<sup>1</sup>, Aurélie Duflot<sup>1</sup>, Agnès Poret<sup>1</sup>,  
5 Emeline Maniez<sup>1</sup>, Frank Le Foll<sup>1</sup>, Romain Coulaud<sup>1\*</sup>.

6 <sup>1</sup>Le Havre Normandie University (ULHN), FR CNRS 3730 SCALE, UMR-INERIS 02  
7 Environmental Stresses and Biomonitoring of Aquatic Ecosystems (SEBIO), 25 rue Philippe  
8 Le Bon - 76600 Le Havre – France.

9 \*Corresponding authors: [romain.coulaud@univ-lehavre.fr](mailto:romain.coulaud@univ-lehavre.fr)

10  
11           **Abstract**

12 In a seasonal environment, variation in larval phenotype and developmental plasticity allow  
13 crustacean larvae to maximise survival by lengthening or shortening their development. The  
14 aim of this study is to investigate the effects of temperature, laying season and their interaction  
15 on larval developmental pathways (larval instars and larval stages). We monitored the different  
16 larval stages and calculated the number of larval instars reached during the development of  
17 winter and summer larvae of *Palaemon serratus* incubated at 12, 16 or 20°C. We observed a  
18 great variability in the larval development (6 to 13 larval instars and 6 to 11 larval stages). A  
19 higher temperature decreases the development time and the number of larval instars. At a given  
20 temperature, the development time of winter and summer larvae was not different. Two larval  
21 stages were considered supernumerary (zoea 4 and 6), as they were more frequent at low  
22 temperatures. At higher temperatures, some larvae started to develop pleopods as early as the  
23 third instar, larval stage which had never been described (named here zoea 3'). This  
24 phenomenon was more common in winter larvae than in summer larvae. These results provide  
25 new insights into the expression of developmental plasticity in decapod larvae.

26           **Key words:** Larval ecology, Plasticity, Developmental pathways, Larval instars, Larval  
27 stages, *Palaemon serratus*.

Journal Pre-proof

## 28           **1. Introduction**

29   Ecological Developmental Biology (Eco-Devo) approaches focus on the interactions between  
30   organisms and their environment, and their ability to optimise their growth and development  
31   (Gilbert, 2001). For Eco-Devo studies, marine decapod crustaceans represent interesting  
32   biological models because their complex life cycle is generally biphasic (for a review see Anger  
33   2006), divided between a pelagic larval phase and a benthic adult life. Given the seasonal or  
34   spatial variability of the pelagic environment (*e.g.* food availability and quality, temperature),  
35   decapods may therefore exhibit a high degree of developmental plasticity (*i.e.* the capacity of a  
36   genotype to produce alternative developmental pathways; West-Eberhard 2003), allowing their  
37   larvae to cope with variable or even sub-optimal conditions (Anger, 2001). Several authors have  
38   studied the effects of environmental parameters on the larval development of marine decapods.  
39   Among abiotic parameters, temperature affects intermoult time, growth rate and number of  
40   larval moults achieved (Linck 1995, read in Giménez 2006; O'Connor *et al.* 2007; Oliphant *et*  
41   *al.* 2013). The amount of food ingested by the larvae determines the number of larval moult  
42   cycles and the rate of morphogenesis between them (Anger, 1987; González-Ortegón and  
43   Giménez, 2014). Thence, prolonged development has frequently been observed under poor  
44   zootechnical conditions (Críaes and Anger, 1986; Giménez and Anger, 2003; González-  
45   Ortegón *et al.*, 2013; González-Ortegón and Giménez, 2014; Knowlton, 1974; Oliphant *et al.*,  
46   2013; Ostrensky *et al.*, 1997). Conversely, abbreviated developments have been encountered  
47   under optimal conditions such as the thermal preference (Críaes and Anger, 1986; Oliphant *et*  
48   *al.*, 2013; Oliphant and Thatje, 2013). It is these different plastic responses to environmental  
49   conditions that lead to different developmental pathways (Críaes and Anger, 1986).

50   In other way, few authors have studied the interactions of environmental conditions and the  
51   offspring phenotype on the developmental plasticity of decapods (Linck 1995, read in Giménez  
52   2006; Hancock 1998; Oliphant & Thatje 2013; González-Ortegón & Giménez 2014; Oliphant

53 *et al.* 2014). The larval size is highly correlated with the nutritional reserves transferred per egg  
54 by the females during oogenesis (Giménez and Anger, 2001; Giménez and Torres, 2002;  
55 Hancock, 1998; Moland *et al.*, 2010). This increases their fitness by allowing them to better  
56 resist starvation and achieve a shorter development (Giménez *et al.*, 2004; Giménez and Anger,  
57 2003; González-Ortegón and Giménez, 2014; Hancock, 1998; Oliphant *et al.*, 2014; Oliphant  
58 and Thatje, 2013). The larval phenotype is therefore determined by maternal investment, in line  
59 with environmental variations. The largest larvae correspond to the most difficult  
60 environmental conditions, such as the beginning of the season (González-Ortegón *et al.*, 2018;  
61 Paschke, 1998; Paschke *et al.*, 2004; Urzúa and Anger, 2013). Thus, decapods living in  
62 seasonally variable environments and reproducing several times during the year show a  
63 seasonal pattern in the “fertility x offspring size trade-off” (Bas *et al.*, 2007; Boddeke, 1982;  
64 Vázquez *et al.*, 2013). The developmental pathways of the larvae of these species should be  
65 seasonally different, linked to the difference in larval size between the seasons.

66 The larval developmental pathway is therefore the result of the interaction between larval  
67 genotype, environmental conditions during development and the environment or phenotype of  
68 the mother (*i.e.* maternal effects, Marshall *et al.*, 2008). In order to study the developmental  
69 plasticity response to this interaction, it is necessary to define the developmental pathways  
70 taken. Decapod larvae develop through a succession of moults with morphological, ecological  
71 and behavioural changes, which are generally called “Larval Stages” (Anger, 2001). However,  
72 in caridean shrimps that develop anamorphically in particular (*i.e.* through a gradual pattern of  
73 development, Møller *et al.* 2020), the degree of morphological changes between moults is  
74 variable (Anger, 2001; for a review see Quinn, 2016). Additional larval stages may be  
75 “intercalated” into the developmental sequence, often called “supernumerary stages”,  
76 displaying intermediate morphology (Knowlton, 1974; Little, 1969). Each larval stage can also  
77 be “repeated”, without showing any significant change (Rothlisberg, 1980; Walsh, 1993;

78 Wehrtmann and Albornoz, 1998). We thus explain that the larvae of a species do not necessarily  
79 develop through a fixed number of moults. Therefore, the neutral term "Larval Instar" is  
80 preferred to refer to the successive moult cycles during larval development, without describing  
81 them (Martin et al., 2014; Phillips and Sastry, 1980).

82 It seems necessary to describe the number of larval instars reached and the different successive  
83 larval stages to accurately characterise the larval developmental pathway. Some authors have  
84 study the developmental plasticity of caridean shrimp larvae in response to environmental  
85 factors by focusing on larval stages, larval instars or both. Hancock (1998) described the larval  
86 stages of *Paratya australiensis* encountered during development at 2 temperatures. Oliphant *et*  
87 *al.* (2014) counted the number of larval instars achieved by *Palaemonetes varians* in relation to  
88 food availability. González-Ortegón and Giménez (2014) carried out the same type of  
89 experiment on *P. serratus* larvae adding the effects of temperature and salinity. Oliphant *et al.*  
90 (2013) monitored both the larval stages and larval instars of *P. varians* during development at  
91 3 temperatures. Críaes and Anger (1986) described the developmental pathways (larval stages  
92 and instars) of *Crangon crangon* and *C. allmanni* according to rearing temperature. Linck  
93 (1995, read in Giménez, 2006) is the only author to have studied the effects of the interaction  
94 between temperature and laying season on larval development pathways (*C. crangon*).

95 To investigate the effects of the interaction between temperature and seasonal variations in  
96 larval size on the variability of larval developmental pathways, we chose to work with the  
97 caridean shrimp, *Palaemon serratus* (Pennant, 1777). This species is abundant in European  
98 coastal environments and has a life span of 3 to 5 years (Campillo, 1979). Its reproductive cycle  
99 consists of two annual clutches (commonly called "winter laying" and "summer laying"),  
100 characterised by larger eggs, and thus larvae, in the first clutch (Campillo, 1979; González-  
101 Ortégón et al., 2018). This species is known to carry out larval development with a variable  
102 number of instars; mostly 6 to 9 but ranging from 6 to 22 in the laboratory (Campillo, 1979;



103 Fincham and Figueras, 1986; Kelly et al., 2012; Reeve, 1969a). This developmental plasticity  
104 is mainly temperature-mediated, with the larvae of this species performing fewer instars in  
105 warm water than in cold water (González-Ortegón and Giménez, 2014). However, variations  
106 in developmental pathways due to temperature-mediated plasticity have not been investigated  
107 in *P. serratus* (larval stages and instars). The effects of seasonal variation in the size of the  
108 larvae of this species on their development have also not been studied, although this should also  
109 influence their developmental pathways (González-Ortegón et al., 2018). We hypothesise the  
110 following two scenarios. Large winter larvae develop faster than small summer larvae at the  
111 same temperature, completing fewer larval instars. Furthermore, the interaction between  
112 temperature and laying season has synergistic effects on developmental pathways. To test these  
113 hypotheses, we monitor the different stages and calculate the number of instars reached during  
114 the development of larvae from winter and summer females of *P. serratus* incubated at 3  
115 temperatures, 12, 16 and 20°C.

116

## 117 **2. Material and methods**

### 118 *2.1 Collection and maintenance of females*

119 To study the larval development of the winter and summer laying, ovigerous *P. serratus*  
120 females (6 per seasons ; Fig.1) were respectively sampled between 29/04/2021 and 12/05/2021,  
121 and between 23/06/2021 and 28/06/2021 by hand-netting on Saint-Jouin-Bruneval foreshore  
122 (Seine-Maritime, France). Only the specimens carrying eggs at the end of development were  
123 selected for the study. The length of the cephalothorax was constant for both seasons, measured  
124 respectively 40.8 mm ( $\pm 1$ ) for winter females and 39.8 mm ( $\pm 1.1$ ) for summer females, in  
125 mean ( $\pm$  S.E.). The shrimps were transferred to the laboratory of the University of Le Havre  
126 (Seine-Maritime, France), and then maintained under photoperiod 12: 12 (L: D) in 100 L tank

127 temperature-controlled at 16°C, filled with filtered aerated natural seawater at 30 ppt ( $\pm 4$ ) of  
128 salinity (Fig. 1). Seawater was renewed by half 3 times per week to avoid nitrate increase. Each  
129 specimen was isolated in wire baskets covered with fine mesh and fed individually twice a week  
130 *ad libitum* with pieces of squid. Larval hatching was checked daily.

131

## 132 2.2 Larval maintenance

133 The objective was to incubate larvae from 12 clutches (6 winter and 6 summer) at three  
134 temperature treatments of 12, 16 and 20°C, as shown in Figure 1. On hatching, actively  
135 swimming larvae were separated from females using a pipette. The rostrum-telson length of 30  
136 newly hatched larvae from each clutch was measured (Campillo, 1979). For each clutch, 3  
137 groups of 104 larvae were distributed in three conical vessels, with filtered seawater at 30 ppt  
138 of salinity ( $\pm 4$ ), at a density of 100 larvae per L (Reeve, 1969b; Torres et al., 2021). The water  
139 was initially at the same temperature as for the maintenance of the females, and then the conical  
140 vessels were distributed to 3 incubators (temperature  $\pm 0.3^\circ\text{C}$ , regulated by a TK700 cooling  
141 unit by TECO), with photoperiod 12: 12 (L: D) and constant thin aeration. This resulted in a  
142 gradual change which allowed time for the larvae to acclimate to the new temperatures  
143 (Richard, 1978). Larvae were fed daily with freshly-hatched *Artemia sp.* nauplii at a density of  
144 10 nauplii per mL (Reeve, 1969b). The water was changed (100 %), and larval mortality and  
145 development, assessed by identification of stages (see section 2.4), were monitored 3 times a  
146 week. During development, the oxygen saturation was 103 % ( $\pm 4$ ), the pH was 8 ( $\pm 0.1$ ) and  
147 the ammonium concentration was 1 mg per L ( $\pm 0.3$ ), in mean ( $\pm$  S.E.). On moulting to the  
148 juvenile stage, shrimps were removed.

149

### 150 2.3 Intermoult time and number of instars

151 As several successive stages can occur without morphological change, it is necessary to  
152 determine the number of larval instars. Because the larvae are mass-reared, the larval average  
153 intermoult time cannot be individually measured and then need to be calculated. The average  
154 intermoult time of *P. serratus* larvae is known to be constant in good zootechnical conditions  
155 (González-Ortegón and Giménez, 2014; Reeve, 1969a). We compiled data on the duration of  
156 larval development of *P. serratus* under controlled temperature conditions available in the  
157 literature. We did not use data concerning larvae exposed to other stresses (salinity,  
158 contaminant, ...). This allows the construction of a function expressing the theoretical  
159 intermoult time of larval as a function of temperature. For each juvenile obtained in the  
160 experiment, its number of instars realised is estimated by dividing its developmental time by  
161 the corresponding theoretical intermoult time.

162

### 163 2.4 Identification of larval stages

164 The different larval stages are identified using morphological criteria. In the larvae of European  
165 palaemonidae, the morphological criteria corresponding to the rostrum, caudal region and  
166 pleopods are commonly used to identify the different larval stages (Fincham and Figueras,  
167 1986). Fincham and Figueras (1986) described 9 larval stages for *P. serratus*, while Campillo  
168 (1979) described 6. Description and terminology for the naming of larval stages differ between  
169 these authors, but it is possible to do correspondences and characterise 10 different larval stages  
170 (Table 1; according to Duflot et al., unpublished). According to these authors, these larval stages  
171 are potentially all observable, but are not obligatory stages through which each larvae must pass  
172 to develop. On the recent recommendations of Møller *et al.* (2020) regarding the terminology  
173 of caridean shrimp larvae, we call these stages zoea 1 to zoea 10 in chronological order of  
174 morphogenesis.

175

176 Table 1: Correspondences between the nomenclatures of the larval stages of *Palaemon serratus*  
 177 described Campillo (1979), Fincham & Figueras (1986) and us; according to Duflot et al.,  
 178 unpublished. The morphological criteria used in this study are outlined.

Authors		Zoea							Juvenile		
Campillo (1979)	1	2	3		4		5	6			
Fincham and Figueras (1986)	1	2	3	4	5	6	7	8	9		
Correspondences	1	2	3	4	5	6	7	8	9	10	
Morphological criteria	Sessile eyes	Pedunculated eyes, 1 rostral spine	2 rostral spines	3 rostral spines, telson trapezoidal shape, no pleopod buds	3 rostral spines, telson trapezoidal shape, little pleopod buds	Telson rectangular shape, little pleopod buds	Telson rectangular shape, rudimentary pleopod buds	Rudimentary biramous pleopods	Well-developed pleopods, endopod of antenna 2 not longer than exopod by more than 2 segments	3 rostral curved spines, endopod of antenna 2 longer than exopod ( $\geq 2$ segments)	Toothed rostrum ( $> 6$ dorsal teeth)

179

180

## 181 2.5 Statistical analysis

182 Statistical procedures were implemented with R software (R Core Team, 2020). The  
183 relationship between larval intermoult time and temperature (Fig. 3; Eq. 1) was obtained using  
184 the *nls* function. The difference in larval size between winter and summer laying was tested  
185 using a nonparametric Wilcoxon Rank Sum test. To test the effects of temperature and laying  
186 season on the rate of larval metamorphosis and the development time of the metamorphosed  
187 larvae, we used mixed-effects models to consider the fact that for each of the seasons, the larvae  
188 used in the different temperatures came from the same females. For the rate of larval  
189 metamorphosis, we fitted generalised linear mixed-effects model with a logit link function and  
190 a binomial error distribution (*glmer* function of the package *lme4*) and performed deviance  
191 analysis procedure including interaction terms (using the package *car*). For the development  
192 time of the metamorphosed larvae, we fitted linear mixed-effects model (using the function  
193 *lmer* of the package *lme4*) and performed deviance analysis procedure including interaction  
194 terms (using the package *car*).

195

## 196 3. Results

### 197 3.1 Zootechnical performances during larval development.

198 The average total length of hatching larvae was significantly higher in winter ( $3.68 \pm 0.07$  mm),  
199 than in summer ( $3.18 \pm 0.05$  mm) in mean  $\pm$  S.E. (Wilcoxon Rank Sum test;  $p < 0.05$ ).

200 While all females produced juveniles at 16 and 20°C, only 5 winter and 3 summer females  
201 produced juveniles at 12°C. At 12°C, only 18 of the 624 winter larvae and 5 of the 624 summer  
202 larvae metamorphosed (Fig. 2a). At 16 and 20°C respectively, 30 % ( $\pm 5$ ) and 41 % ( $\pm 7$ ) of the  
203 winter larvae metamorphosed, compared to 43 % ( $\pm 7$ ) and 55 % ( $\pm 7$ ) of the summer larvae, in  
204 mean ( $\pm$  S.E.). The rate of larval metamorphosis varied significantly with temperature and

205 laying season without interaction of these two factors (Deviance analysis, interaction term:  
 206  $Chisq = 3.18, p = 0.07$ ; temperature:  $Chisq = 16.19, p < 10^{-5}$ ; season:  $Chisq = 26.6, p < 10^{-7}$ ;  
 207  $n = 6$  per temperature and season).

208 The metamorphoses took place between days 65 and 111 at 12°C, between days 30 and 56 at  
 209 16°C and between days 19 and 42 at 20°C (Fig. 2b). The development time of the  
 210 metamorphosed larvae varied significantly between the three incubation temperatures, but  
 211 neither as a function of the laying season nor as a result of the interaction of these two factors  
 212 (Deviance analysis, interaction term:  $Chisq = 0.76, p = 0.4$ ; temperature:  $Chisq = 138.73,$   
 213  $p < 10^{-16}$ ; season:  $Chisq = 0.62, p = 0.43$ ;  $n = 6$  per temperature and season).

214

### 215 3.2 Number of instars achieved by larvae.

216 The compilation of literature data on the larval development of *P. serratus* under controlled  
 217 temperature conditions allows us to construct a function expressing the theoretical larval  
 218 intermoult time *versus* temperature (Fig. 3; Eq. 1):

$$219 \quad LIT = 147.96 e^{(-0.25 \text{ temp})} + 2.45 \quad \text{Eq. 1}$$

220 where *LIT* is the theoretical larval intermoult time and *temp* the incubation temperature.

221 This function allows us to calculate the theoretical intermoult times at 12°C (9.8 days), 16°C  
 222 (5.2 days) and 20°C (3.4 days) as well as the number of instars achieved by each juvenile  
 223 obtained in this experiment. In all experimental conditions, the larvae that metamorphosed did  
 224 so after 6 to 13 larval instars (Fig. 4). Only about 3.7 % of them developed for more than 9  
 225 instars, and about 1 % for 6 instars. While about 33 and 45 % of the larvae performed 7 instars  
 226 at 16 and 20°C, about 43 and 31 % of them performed 8 instars. At 20°C, larvae  
 227 metamorphosing after 7 instars represented about 40 % of the total in winter and 50 % in

228 summer. At 16°C, 3 larvae metamorphosed after 6 instars in summer, while none did so in  
229 winter.

230

### 231 *3.3 Succession of larval stages.*

232 Figures 5 to 7 show the number of individuals observed for each larval stage during  
233 development, separating the larvae by clutch and incubation temperature. With the exception  
234 of the first few days of development, the larvae from the same clutch incubated at the same  
235 temperature showed a diversity of larval stages, highlighting different developmental pathways  
236 within the same clutch. This diversity increased during development, so that the time of  
237 observation of a larval stage within the clutch also increased during monitoring. Regardless of  
238 the experimental conditions and the rate of metamorphosis, larval mortality is constant  
239 throughout development. Only the larvae of the females 2, 4 and 9 at 12°C experienced more  
240 than 50 % mortality from the zoea 1 stage (Fig. 5).

241 All 10 larval stages described were observed at each temperature. However, we observed  
242 differences in the proportions of the different larval stages. At 12, 16 and 20°C, respectively, at  
243 least 15.3, 5.9 and 0.4 % of the larvae passed through the zoea 4 stage, and at least 9.1, 0.6 and  
244 0.3 % of the larvae passed through the zoea 6 stage.

245 Furthermore, during the monitoring of the larval stages presented here, we frequently observed  
246 larvae with 2 rostral spines (criteria for zoea 3) and pleopod buds (criteria for zoea 5). This  
247 larval stage seems to take place instead of the zoea 3 stage, so we called it zoea 3'. This never  
248 described larval stage was not observed at 12°C. At 16 and 20°C, respectively, at least 58.1 and  
249 67.8 % of the winter larvae passed through the zoea 3' stage, compared to at least 20.8 and  
250 19.9 % of summer larvae.

251

#### 252 **4. Discussion**

253 Regardless of the incubation temperature and the laying season from which the larvae hatched,  
254 we observed a great variability in the larval development of *P. serratus* (6 to 13 larval instars  
255 and 6 to 11 larval stages). The different developmental pathways observed underline the  
256 developmental plasticity of this species.

257 In contrast to several authors studying the number of larval instars of decapods during  
258 development (Reeve, 1969a; Dalley, 1980; Linck, 1995, read in Giménez, 2006), we did not  
259 rear the larvae individually. This choice was made in view of the survival rates of *P. serratus*  
260 larvae in the laboratory available in the literature. While González-Ortegón and Giménez  
261 (2014) and Lassus and Maggi (1980) achieved respectively 40 and 45-70% metamorphosis  
262 when rearing larvae individually, Campillo (1979) and Reeve (1969b) achieved respectively 76  
263 and over 90% metamorphosis with mass-rearing. Furthermore, Reeve (1969a) observed longer  
264 larval development with lower survival when rearing *P. serratus* individually than when mass-  
265 rearing.

266 We calculated the number of instars made by the larvae from their average intermoult time  
267 according to bibliographic data. Intermoult times obtained for the same temperature may vary  
268 slightly between the 6 sources used (Fig. 3). This can be explained by the differences in  
269 experimental protocol and by maternal effects. The limited information available on this second  
270 factor complicates discussions on this subject. The females used in these 6 experiments came  
271 from populations in Ireland, Wales, Brittany and the Mediterranean (the latter being  
272 phylogenetically distant from the Atlantic populations; Weiss et al., 2018). Only González-  
273 Ortégón and Giménez (2014) specify the period of collection of females (autumn) as well as  
274 their size (between 17 and 24 mm from the postorbital margin to the end of carapace, between  
275 18 and 22 mm in our study). According to Reeve (1969a) and González-Ortegón and Giménez  
276 (2014), the intermoult time of *P. serratus* is constant during larval development under good



277 zootechnical conditions. Nevertheless, these authors reared the larvae of this species between  
278 18 and 24°C. It is possible that the intermoult time of *P. serratus* larvae increases as  
279 development proceeds at lower temperatures, as reported by Kelly et al. (2012) and Lassus and  
280 Maggi (1980) between 10 and 13°C. However, the few individuals that metamorphosed at 12°C  
281 leads us to put the interpretation of the number of instars carried out at this temperature into  
282 perspective.

283

#### 284 *4.1 Temperature-mediated developmental plasticity*

285 The larval development of *P. serratus* is temperature-driven. Here, metamorphosis rates were  
286 temperature dependent (about 2, 36 and 49 % at 12, 16 and 20°C respectively). Campillo (1979)  
287 experienced the highest rates of metamorphosis (85 %) by incubating *P. serratus* larvae from  
288 Roscoff (France) at 27°C, a temperature above the local maximum. Other authors who studied  
289 the larval development of *P. serratus* at temperatures between 10 and 13°C observed a mortality  
290 of all individuals between the third and fourth larval instar (Kelly et al., 2012; Lassus and  
291 Maggi, 1980; Yagi and Ceccaldi, 1985). We also observed very low survival rates at 12°C, but  
292 we managed to get some individuals to metamorphosis (< 2%) after 7-13 instars. We chose to  
293 incubate larvae at 12°C because this is the temperature of the sea water in the Bay of Seine  
294 (Normandy) at the beginning of May (Claquin et al., 2018), the date at which the larvae from  
295 the winter spawning are released by females of the same size as those used for this study  
296 (personal observations). Mortality at 12°C was constant during development in the laboratory.  
297 At this temperature, only the larvae from 3 of the 12 females suffered mortality rates above  
298 50 % by the end of the first instar. The mortality rate of the larvae was approximately 52 %  
299 after 30 days of incubation, when most of them were in their fourth instar. The thermal optimum  
300 of crustacean larvae experiences an ontogenic shift (Sastry and McCarthy, 1973). These  
301 variations between larval stages are due to temperature-induced changes in metabolic rates

302 (Small et al., 2015). During the month of May, the temperature of the sea water in the Bay of  
303 Seine increases by approximately 3°C (Claquin et al., 2018). A constant temperature of 12°C  
304 in the laboratory negatively affects the survival rates of *P. serratus* larvae. These effects  
305 probably increase during ontogeny.

306 In addition to decreasing survival, colder water increases the duration of development. First of  
307 all, as we know from the literature, the intermoult time increases as the temperature decreases  
308 following a negative exponential relationship (Fig. 2). Secondly, larvae tend to go through a  
309 greater number of instars in colder waters (Fig. 4). Although the duration of development is  
310 variable between individuals incubated at the same temperature, this is not so with the value of  
311 the average duration per family (Fig. S1, Supplement). The plastic response to temperature  
312 therefore appears to be the same for larvae produced by different females. However, differences  
313 in survival rates appear between larvae of different families (Fig. S1, Supplement).

314 Concerning the larval stages, zoea 4 and zoea 6 are more frequent in colder water (Fig. 5 to 7).  
315 These larval stages can be interpreted as supernumerary, or intercalated, stages occurring under  
316 unfavourable conditions such as low temperature. This would explain why Campillo (1979) did  
317 not describe them at 27°C (Table 1). Conversely, at 16 and 20°C, many larvae with zoea 3  
318 criteria (2 rostral spines) and zoea 5 criteria (pleopod buds) were observed (about 39 and 44 %  
319 at 16 and 20°C respectively). This never described stage, which we have called zoea 3', seems  
320 to replace the zoea 3 stage. This larval stage can be interpreted as a sign of abbreviated  
321 development, occurring under favourable conditions. Caridean shrimps have anamorphic  
322 development, with gradual morphogenesis. Several authors attempt to define and differentiate  
323 terminologies such as repeated instars, supernumerary and intermediate stages. These terms are  
324 the result of the conservative belief in immutable development in decapod crustaceans (Sandifer  
325 and Smith, 1979). The study of the anamorphic development of caridean shrimps allows us to  
326 consider this belief as a zombie idea. The multiple examples of developmental plasticity

327 observed in species from different groups of decapods support this view (for a review see  
328 Quinn, 2016).

329 As stated in the introduction, this temperature-mediated developmental plasticity has already  
330 been demonstrated in caridean shrimps. The larvae follow shorter developmental pathways in  
331 temperatures closer to their thermal optimum (Críaes and Anger, 1986; Linck, 1995, read in  
332 Giménez, 2006; Oliphant et al., 2013). Generally, at a given temperature, the more instars the  
333 larvae develop through, the larger the juveniles. This trend has been demonstrated in the larvae  
334 of several decapods including *Chasmagnathus granulata* (Giménez and Torres, 2002) and  
335 *P. varians* (Oliphant et al., 2013). Therefore, it would be interesting to compare the  
336 consequences of temperature-mediated developmental plasticity of *P. serratus* larvae on the  
337 phenotype of juveniles.

338 These differences in survival between thermal conditions can also be explained by differences  
339 in nutritional requirements. Crustacean larvae require long-chain polyunsaturated fatty acids  
340 (PUFAs) for development, such as eicosapentaenoic acid, or EPA, and docosahexaenoic acid,  
341 or DHA (Anger, 1998; Bergé and Barnathan, 2005). These are generally synthesised by  
342 phytoplankton and transferred into the food chain (Lubzens et al., 1995). PUFAs, DHA in  
343 particular, maintain membrane integrity and viscosity (Estévez and Kanazawa, 1996; Los and  
344 Murata, 2004; Pagliarani et al., 1986), especially in cold waters (Harwood, 1988; Munro and  
345 Thomas, 2004). Like many authors, we fed *P. serratus* larvae with artemia nauplii (Campillo,  
346 1979; Reeve, 1969b; Wickins, 1972; Yúfera and Rodríguez, 1985). Although these are  
347 traditionally used to feed decapod larvae, they are known to be deficient in PUFAs (Narciso  
348 and Morais, 2001; Navarro et al., 1999). These deficiencies may explain the very low survival  
349 of *P. serratus* larvae in cold water observed here and by the authors mentioned in the paragraph  
350 above. PUFAs also influence growth, stress and disease resistance in decapods (Anger, 1998;  
351 Kanazawa et al., 1985; Mourente and Rodríguez, 1997). It would be interesting to study the

352 effects of the interaction between temperature and diet quality on the developmental plasticity  
353 of *P. serratus*.

354

#### 355 4.2 Developmental plasticity induced by laying season and its interaction with temperature

356 The newly hatched larvae used in this study were larger in winter than in summer. These results  
357 are consistent with González-Ortegón *et al.* (2018) who observed difference in *P. serratus*  
358 larval biomass between laying seasons. Because of this difference in phenotype, we  
359 hypothesised that *P. serratus* larvae from winter females would develop faster than those from  
360 summer females. These effects were not significant in our data analyses. At a given  
361 temperature, summer larvae even appeared to perform fewer larval instars than winter larvae.  
362 These results are in contradiction with those obtained on another caridean shrimp species  
363 (Linck, 1995, read in Giménez, 2006). The author incubated larvae of *C. crangon*, from large  
364 winter eggs and small summer eggs, at two relatively cold and warm temperatures (12 and  
365 18°C). Regardless of the temperature, summer larvae metamorphosed after a greater number of  
366 larval instars than winter larvae. We need to further investigate the effects of the interaction  
367 between the laying season and environmental conditions on the developmental plasticity of  
368 *P. serratus*.

369 The winter larvae were larger than the summer ones. Few authors have studied the effects of  
370 larval size or larval biomass on developmental plasticity of decapod larvae. The number of  
371 larval instars made by larvae decreases with increasing larval size or larval biomass in  
372 *C. crangon* (Paschke *et al.*, 2004), *C. granulata* (Giménez *et al.*, 2004), *Homarus gammarus*  
373 (Moland *et al.*, 2010), *P. varians* (Oliphant *et al.*, 2014; Oliphant and Thatje, 2013) and  
374 *P. australiensis* (Hancock, 1998), a phenomenon that is accentuated under adverse conditions.  
375 Although they highlighted the decrease in the number of larval instars with increasing initial

376 larval biomass in *P. serratus* larvae under starvation conditions, González-Ortegón & Giménez  
377 (2014) did not confirm this trend under favourable feeding conditions. This result is consistent  
378 with those presented here.

379 Although the development time of the larvae in the two laying seasons did not differ, at 16 and  
380 20°C, the zoea 3' stage was observed more often in winter larvae than in summer larvae (Fig. 5  
381 to 7). The morphogenesis of the winter larvae (pleopod buds) was earlier than that of the  
382 summer larvae, at least in the early larval stages. The winter larvae may have received more  
383 available energy resources (transmitted by the mother) than summer larvae and used them  
384 during these early instars. This phenomenon may have been favoured by the choice of females  
385 used. The cephalothorax of the winter females measured on average about 40 mm. These would  
386 be more than two years old, which corresponds to less than 2 % of the female population of  
387 *P. serratus* (Campillo, 1979). The limited information available on the size of the females and  
388 the date of their sampling in previous publications does not allow comparison of our results.  
389 The lack of description of the zoea 3' stage in previous studies can be explained by the choice  
390 of females used. We need to investigate the effects of female age on maternal investment and  
391 developmental plasticity in *P. serratus*.

392 When they hatch, decapod larvae have visible vitelline reserves in the cephalothorax in the form  
393 of yolk droplets (Anger, 1996; Oliphant and Thatje, 2014). These can be used to cope with  
394 starvation conditions during the first larval instars (Anger, 1995a, 1995b; Thessalou-Legaki et  
395 al., 1999), *e.g.* when larvae are exported from a brackish environment to the marine  
396 environment (Anger and Hayd, 2009; Oliphant and Thatje, 2014). Such yolk droplets are  
397 observed in hatched larvae of *P. serratus* (personal observation). Narciso and Morais (2001)  
398 measured a decrease in fatty acid content during the first 48 hours of development of *P. serratus*  
399 larvae under starvation conditions. Their results and ours seem to confirm the use of vitelline  
400 reserves during the first larval instars in *P. serratus*. Future analyses to compare the quantity

401 and quality of resources allocated per offspring, called POI (*i.e.* per offspring investment;  
402 Oliphant, 2014), as a result of the laying season and their effects on larval development will  
403 provide a better understanding of the developmental plasticity in *P. serratus* larvae.

404 In *P. serratus*, winter larvae are larger than summer larvae. We expected that this phenotypic  
405 difference would allow winter larvae to develop faster than summer larvae at the same  
406 temperature. Although we did not observe this result, we did observe earlier morphogenesis in  
407 winter larvae at the warmest temperatures. As explained in the previous paragraphs (see section  
408 4.2), we believe that this phenomenon is related to the amount of energy allocated per offspring.  
409 This pattern of maternal investment buffers winter larvae from variable early season conditions  
410 (low temperature, Torres et al., 2020; less abundant planktonic food resources, Paschke et al.,  
411 2004). Furthermore, the negative effects of food limitation on larval survival are exacerbated  
412 by unfavourable temperatures (Torres and Giménez, 2020). The few months separating the  
413 hatching of the larvae from the two laying periods of *P. serratus* result in two cohorts of  
414 juveniles (Campillo, 1979). As these cohorts share the same habitats, they compete for  
415 resources or even cannibalise (Moksnes et al., 1998). The shift in the recruitment periods of  
416 these two cohorts results in a difference in the size of the individuals (*P. serratus* juveniles of  
417 0.5g which double or triple their weight in 2 months depending on their sex at temperatures  
418 between 18 and 29°C; Rodriguez, 1981). These traits probably favour juveniles from winter  
419 laying over those from summer laying in years when early season conditions are favourable  
420 (Temming et al., 2017). The population structure and dynamics of *P. serratus* should be studied  
421 to assess the settlement success of winter and summer cohort. This would facilitate  
422 understanding the effects of juvenile recruitment phenology on their interactions.

423

424 *Conclusion*

425 Although larval development of *P. serratus* is plastic in response to temperature, the interaction  
426 between temperature and laying season does not seem to influence its developmental pathways  
427 as expected. As the development of some groups of decapods is gradual, understanding their  
428 developmental pathways will require consideration of the number of instars and the succession  
429 of larval stages.

430

431 **Acknowledgments**

432 We would like to thank the Thalasso of Deauville for the supply of natural sea water and the  
433 University of Le Havre for the facilities. This study was supported by the Normandie Region  
434 and the European Maritime and Fisheries Fund (Action 28 “Partnerships between scientists and  
435 fishermen”; GeDuBouq project).

436

437 **Vitae**

438 Jean-Baptiste Baudet is working on crustacean ecology, developmental plasticity, phenotypic  
439 links and larval ecology during his PhD project started in December 2020. His previous  
440 experience was in penaeid shrimp research and development.

441

442 Benoît Xuereb is a researcher at the research unit for Environmental Stresses and Biomonitoring  
443 of Aquatic Ecosystems laboratory (SEBIO), Le Havre – Normandy University. His research  
444 interest focuses on the description of ecophysiology and life traits in the palaemonid shrimps,  
445 with a perspective to propose reliable effect-based methods for ecotoxicological applications.

446

447 Aurélie Duflot works in the SEBIO laboratory on various fields related to the physiology of  
448 marine organisms and is an expert on the embryo-larval development of *Palaemon serratus*.

449

450 Agnès Poret is an engineer at the SEBIO laboratory working on marine invertebrates, especially  
451 on reproduction and immunology.

452

453 Emeline Maniez has completed a master's degree in marine sciences. She specialises in marine  
454 and fisheries ecology, as well as in the functioning and management of marine ecosystems. She  
455 is interested in shrimp and plankton, for example, in relation to global changes and/or fisheries.

456

457 Frank Le Foll is a professor of organismal biology at the University of Le Havre Normandie  
458 and currently the head of the Environmental Stresses and Biomonitoring of Aquatic Ecosystems  
459 laboratory (SEBIO). SEBIO carries out basic and translational research in bioenvironmental  
460 sciences and ecotoxicology. Frank research interest focuses on invertebrate immune cell  
461 activity, host-microbe-environment interactions, multixenobiotic resistance phenotype,  
462 immunomarker research and innovative cage design for biological monitoring of polluted water  
463 bodies.

464

465 Romain Coulaud is a researcher at the research unit for Environmental Stresses and  
466 Biomonitoring of Aquatic Ecosystems laboratory (SEBIO), Le Havre – Normandy University.  
467 His research interest focuses on the ecology, ecophysiology and ecotoxicology of  
468 macroinvertebrates (particularly crustaceans), using multi-scale approaches ranging from sub-



469 individual biomarkers to community functioning and combining laboratory and field  
 470 experiments and modelling.

471

## 472 5. References

- 473 Anger, K., 2006. Contributions of larval biology to crustacean research: a review. *Invertebr.*  
 474 *Reprod. Dev.* 49, 175–205. <https://doi.org/10.1080/07924259.2006.9652207>
- 475 Anger, K., 2001. *The Biology of Decapod Crustacean Larvae*, Lisse: AA Balkema Publishers.  
 476 ed, Crustacean Issues.
- 477 Anger, K., 1998. Patterns of growth and chemical composition in decapod crustacean larvae.  
 478 *Invertebr. Reprod. Dev.* 33, 159–176. <https://doi.org/10.1080/07924259.1998.9652629>
- 479 Anger, K., 1996. Physiological and biochemical changes during lecithotrophic larval  
 480 development and early juvenile growth in the northern stone crab, *Lithodes maja*  
 481 (Decapoda: Anomura). *Mar. Biol.* 126, 283–296. <https://doi.org/10.1007/BF00347453>
- 482 Anger, K., 1995a. Starvation resistance in larvae of a semiterrestrial crab, *Sesarma curacaoense*  
 483 (Decapoda: Grapsidae). *J. Exp. Mar. Biol. Ecol.* 187, 161–174.  
 484 [https://doi.org/10.1016/0022-0981\(94\)00178-G](https://doi.org/10.1016/0022-0981(94)00178-G)
- 485 Anger, K., 1995b. Developmental biology of *Armases miersii* (Grapsidae), a crab breeding in  
 486 supratidal rock pools. I. Facultative lecithotrophy of larval stages. *Mar. Ecol. Prog. Ser.*  
 487 117, 75–81.
- 488 Anger, K., 1987. The D0 threshold: a critical point in the larval development of decapod  
 489 crustaceans. *J. Exp. Mar. Biol. Ecol.* 108, 15–30. [https://doi.org/10.1016/0022-0981\(87\)90128-6](https://doi.org/10.1016/0022-0981(87)90128-6)
- 490 Anger, K., Hayd, L., 2009. From lecithotrophy to planktotrophy: ontogeny of larval feeding in  
 491 the Amazon River prawn *Macrobrachium amazonicum*. *Aquat. Biol.* 7, 19–30.  
 492 <https://doi.org/10.3354/ab00180>
- 493 Bas, C.C., Spivak, E.D., Anger, K., 2007. Seasonal and interpopulational variability in  
 494 fecundity, egg size, and elemental composition (CHN) of eggs and larvae in a grapsoid  
 495 crab, *Chasmagnathus granulatus*. *Helgol. Mar. Res.* 61, 225–237.  
 496 <https://doi.org/10.1007/s10152-007-0070-y>
- 497 Bergé, J.-P., Barnathan, G., 2005. Fatty Acids from Lipids of Marine Organisms: Molecular  
 498 Biodiversity, Roles as Biomarkers, Biologically Active Compounds, and Economical  
 499 Aspects, in: Ulber, R., Le Gal, Y. (Eds.), *Marine Biotechnology I, Advances in*  
 500 *Biochemical Engineering/Biotechnology*. Springer, Berlin, Heidelberg, pp. 49–125.  
 501 <https://doi.org/10.1007/b135782>
- 502 Boddeke, R., 1982. The occurrence of winter and summer eggs in the brown shrimp (*Crangon*  
 503 *crangon*) and the pattern of recruitment. *Neth. J. Sea Res.* 16, 151–162.  
 504 [https://doi.org/10.1016/0077-7579\(82\)90026-6](https://doi.org/10.1016/0077-7579(82)90026-6)
- 505 Campillo, A., 1979. Contribution à l'étude de la crevette rose *Palaemon serratus* (Pennant) :  
 506 Exploitation, Biologie, Elevage, Epidémiologie (PhD Thesis). Université des sciences  
 507 et techniques du Languedoc.
- 508 Claquin, P., Jacqueline, F., Repecaud, M., Riou, P., 2018. MAREL SMILE buoy data and  
 509 metadata from coriolis Data Centre. <https://doi.org/10.17882/53689>
- 510 Criales, M.M., Anger, K., 1986. Experimental studies on the larval development of the shrimps  
 511 *Crangon crangon* and *C. allmanni*. *Helgoländer Meeresunters.* 40, 241–265.  
 512 <https://doi.org/10.1007/BF01983735>
- 513

- 514 Dalley, R., 1980. The survival and development of the shrimp *Crangon crangon* (L.), reared in  
515 the laboratory under non-circadian light-dark cycles. *J. Exp. Mar. Biol. Ecol.* 47, 101–  
516 112. [https://doi.org/10.1016/0022-0981\(80\)90105-7](https://doi.org/10.1016/0022-0981(80)90105-7)
- 517 Estévez, A., Kanazawa, A., 1996. Fatty Acid Composition of Neural Tissues of Normally  
518 Pigmented and Unpigmented Juveniles of Japanese Flounder Using Rotifer and *Artemia*  
519 Enriched in n-3 HUFA. *Fish. Sci.* 62, 88–93. <https://doi.org/10.2331/fishsci.62.88>
- 520 Fincham, A.A., Figueras, A.J., 1986. Larval keys and diagnoses for the subfamily  
521 Palaemoninae (Crustacea: Decapoda: Palaemonidae) in the north-east Atlantic and  
522 aspects of functional morphology. *J. Nat. Hist.* 20, 203–224.  
523 <https://doi.org/10.1080/00222938600770171>
- 524 Gilbert, S.F., 2001. Ecological Developmental Biology: Developmental Biology Meets the Real  
525 World. *Dev. Biol.* 233, 1–12. <https://doi.org/10.1006/dbio.2001.0210>
- 526 Giménez, L., Anger, K., 2003. Larval performance in an estuarine crab, *Chasmagnathus*  
527 *granulata*, is a consequence of both larval and embryonic experience. *Mar. Ecol. Prog.*  
528 *Ser.* 249, 251–264. <https://doi.org/10.3354/meps249251>
- 529 Giménez, L., Anger, K., 2001. Relationships among salinity, egg size, embryonic development,  
530 and larval biomass in the estuarine crab *Chasmagnathus granulata* Dana, 1851. *J. Exp.*  
531 *Mar. Biol. Ecol.* 260, 241–257. [https://doi.org/10.1016/S0022-0981\(01\)00258-1](https://doi.org/10.1016/S0022-0981(01)00258-1)
- 532 Giménez, L., Anger, K., Torres, G., 2004. Linking life history traits in successive phases of a  
533 complex life cycle: effects of larval biomass on early juvenile development in an  
534 estuarine crab, *Chasmagnathus granulata*. *Oikos* 104, 570–580.  
535 <https://doi.org/10.1111/j.0030-1299.2004.12957.x>
- 536 Giménez, L., Torres, G., 2002. Larval growth in the estuarine crab *Chasmagnathus granulata*:  
537 the importance of salinity experienced during embryonic development, and the initial  
538 larval biomass. *Mar. Biol.* 141, 877–885. <https://doi.org/10.1007/s00227-002-0887-5>
- 539 González-Ortegón, E., Blasco, J., Le Vay, L., Giménez, L., 2013. A multiple stressor approach  
540 to study the toxicity and sub-lethal effects of pharmaceutical compounds on the larval  
541 development of a marine invertebrate. *J. Hazard. Mater.* 263, 233–238.  
542 <https://doi.org/10.1016/j.jhazmat.2013.09.041>
- 543 González-Ortegón, E., Giménez, L., 2014. Environmentally mediated phenotypic links and  
544 performance in larvae of a marine invertebrate. *Mar. Ecol. Prog. Ser.* 502, 185–195.  
545 <https://doi.org/10.3354/meps10708>
- 546 González-Ortegón, E., Vay, L.L., Walton, M.E.M., Giménez, L., 2018. Maternal Trophic Status  
547 and Offspring Phenotype in a Marine Invertebrate. *Sci. Rep.* 8, 9618.  
548 <https://doi.org/10.1038/s41598-018-27709-2>
- 549 Hancock, M.A., 1998. The relationship between egg size and embryonic and larval  
550 development in the freshwater shrimp *Paratya australiensis* Kemp (Decapoda:  
551 Atyidae). *Freshw. Biol.* 39, 715–723. <https://doi.org/10.1046/j.1365-2427.1998.00323.x>
- 553 Harwood, J.L., 1988. Fatty Acid Metabolism. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 39,  
554 101–138. <https://doi.org/10.1146/annurev.pp.39.060188.000533>
- 555 Kanazawa, A., Teshima, S.-I., Sakamoto, M., 1985. Effects of dietary lipids, fatty acids, and  
556 phospholipids on growth and survival of prawn (*Penaeus japonicus*) larvae.  
557 *Aquaculture* 50, 39–49. [https://doi.org/10.1016/0044-8486\(85\)90151-6](https://doi.org/10.1016/0044-8486(85)90151-6)
- 558 Kelly, E., Tully, O., Browne, R., 2012. Effects of temperature and salinity on the survival and  
559 development of larval and juvenile *Palaemon serratus* (Decapoda: Palaemonidae) from  
560 Irish waters. *J. Mar. Biol. Assoc. U. K.* 92, 151–161.  
561 <https://doi.org/10.1017/S0025315411000415>
- 562 Knowlton, R.E., 1974. Larval developmental processes and controlling factors in decapod  
563 Crustacea, with emphasis on Caridea.

- 564 Lassus, P., Maggi, P., 1980. Etude de la sensibilité thermique des larves du crustacé décapode:  
565 *Palaemon serratus* (Pennant). Rev Trav Inst Phes Marit 44, 257–268.
- 566 Linck, B., 1995. Einfluß von Temperatur und Salzgehalt auf die Larven der Nordseegarnele  
567 *Crangon crangon*.
- 568 Little, G., 1969. The Larval Development of the Shrimp, *Palaemon macrodactylus* Rathbun,  
569 Reared in the Laboratory, and the Effect of Eystalk Extirpation on Development.  
570 Crustaceana 17, 69–87.
- 571 Los, D.A., Murata, N., 2004. Membrane fluidity and its roles in the perception of environmental  
572 signals. Biochim. Biophys. Acta BBA - Biomembr., Lipid-Protein Interactions 1666,  
573 142–157. <https://doi.org/10.1016/j.bbamem.2004.08.002>
- 574 Lubzens, E., Gibson, O., Zmora, O., Sukenik, A., 1995. Potential advantages of frozen algae  
575 (*Nannochloropsis sp.*) for rotifer (*Brachionus plicatilis*) culture. Aquaculture 133, 295–  
576 309. [https://doi.org/10.1016/0044-8486\(95\)00010-Y](https://doi.org/10.1016/0044-8486(95)00010-Y)
- 577 Marshall, D.J., Allen, R.M., Crean, A.J., 2008. The ecological and evolutionary importance of  
578 maternal effects in the sea, in: Oceanography and Marine Biology: An Annual Review.  
579 CRC Press, pp. 203–250.
- 580 Martin, J.W., Olesen, J., Høeg, J.T., Høeg, J., 2014. Atlas of Crustacean Larvae. JHU Press.
- 581 Moksnes, P.-O., Pihl, L., Montfrans, J. van, 1998. Predation on postlarvae and juveniles of the  
582 shore crab *Carcinus maenas*: importance of shelter, size and cannibalism. Mar. Ecol.  
583 Prog. Ser. 166, 211–225. <https://doi.org/10.3354/meps166211>
- 584 Moland, E., Olsen, E.M., Stenseth, N.C., 2010. Maternal influences on offspring size variation  
585 and viability in wild European lobster *Homarus gammarus*. Mar. Ecol. Prog. Ser. 400,  
586 165–173. <https://doi.org/10.3354/meps08397>
- 587 Møller, O.S., Anger, K., Guerao, G., 2020. Patterns of larval development, in: Developmental  
588 Biology and Larval Ecology : The Natural History of the Crustacea. Oxford University  
589 Press.
- 590 Mourente, G., Rodríguez, A., 1997. Effects of salinity and dietary DHA (22:6n-3) content on  
591 lipid composition and performance of *Penaeus kerathurus* postlarvae. Mar. Biol. 128,  
592 289–298. <https://doi.org/10.1007/s002270050094>
- 593 Munro, D., Thomas, D.W., 2004. The role of polyunsaturated fatty acids in the expression of  
594 torpor by mammals: a review. Zoology 107, 29–48.  
595 <https://doi.org/10.1016/j.zool.2003.12.001>
- 596 Narciso, L., Morais, S., 2001. Fatty Acid Profile of *Palaemon Serratus* (Palaemonidae) Eggs  
597 and Larvae During Embryonic and Larval Development Using Different Live Diets. J.  
598 Crustac. Biol. 21, 566–574. <https://doi.org/10.1163/20021975-99990158>
- 599 Navarro, J.C., Henderson, R.J., McEvoy, L.A., Bell, M.V., Amat, F., 1999. Lipid conversions  
600 during enrichment of *Artemia*. Aquaculture 174, 155–166.  
601 [https://doi.org/10.1016/S0044-8486\(99\)00004-6](https://doi.org/10.1016/S0044-8486(99)00004-6)
- 602 O'Connor, M.I., Bruno, J.F., Gaines, S.D., Halpern, B.S., Lester, S.E., Kinlan, B.P., Weiss,  
603 J.M., 2007. Temperature control of larval dispersal and the implications for marine  
604 ecology, evolution, and conservation. Proc. Natl. Acad. Sci. 104, 1266–1271.  
605 <https://doi.org/10.1073/pnas.0603422104>
- 606 Oliphant, A., 2014. Decapod crustacean larval developmental plasticity and the evolution of  
607 lecithotrophy and abbreviated development (PhD Thesis). University of Southampton.
- 608 Oliphant, A., Hauton, C., Thatje, S., 2013. The Implications of Temperature-Mediated  
609 Plasticity in Larval Instar Number for Development within a Marine Invertebrate, the  
610 Shrimp *Palaemonetes varians*. PLOS ONE 8, e75785.  
611 <https://doi.org/10.1371/journal.pone.0075785>
- 612 Oliphant, A., Ichino, M.C., Thatje, S., 2014. The influence of per offspring investment (POI)  
613 and starvation on larval developmental plasticity within the palaemonid shrimp,

- 614 *Palaemonetes varians*. Mar. Biol. 161, 2069–2077. <https://doi.org/10.1007/s00227->  
615 014-2486-7
- 616 Oliphant, A., Thatje, S., 2014. Energetic adaptations to larval export within the brackish-water  
617 palaemonine shrimp, *Palaemonetes varians*. Mar. Ecol. Prog. Ser. 505, 177–191.  
618 <https://doi.org/10.3354/meps10767>
- 619 Oliphant, A., Thatje, S., 2013. Per offspring investment implications for crustacean larval  
620 development: evolutionary insights into endotrophy and abbreviated development. Mar.  
621 Ecol. Prog. Ser. 493, 207–217. <https://doi.org/10.3354/meps10496>
- 622 Ostrensky, A., Pestana, D., Sternheim, U.S., 1997. Effects of different diets on the larval  
623 development and ammonia excretion rates of the crab *Chasmagnathus granulata* Dana,  
624 1851, under laboratory conditions. Ciênc. E Cult. 49, 205–210.
- 625 Pagliarani, A., Pirini, M., Trigari, G., Ventrella, V., 1986. Effect of diets containing different  
626 oils on brain fatty acid composition in sea bass (*Dicentrarchus labrax* L.). Comp.  
627 Biochem. Physiol. B 83, 277–282. [https://doi.org/10.1016/0305-0491\(86\)90366-4](https://doi.org/10.1016/0305-0491(86)90366-4)
- 628 Paschke, K., 1998. Untersuchungen zum Energiestoffwechsel der Nordsee-Garnele *Crangon*  
629 *crangon* (Linnaeus (1758) (Decapoda : Caridea). Diss. Univ. Hambg.
- 630 Paschke, K.A., Gebauer, P., Buchholz, F., Anger, K., 2004. Seasonal variation in starvation  
631 resistance of early larval North Sea shrimp *Crangon crangon* (Decapoda: Crangonidae).  
632 Mar. Ecol. Prog. Ser. 279, 183–191. <https://doi.org/10.3354/meps279183>
- 633 Phillips, B.F., Sastry, A.N., 1980. Larval ecology, in: The Biology and Management of  
634 Lobsters: Ecology and Management. Elsevier.
- 635 Quinn, B.K., 2016. Extra and “intermediate” larval stages in decapod crustacea: a review of  
636 physiological causes and ecological implications, with emphasis on lobsters, *Homarus*  
637 *sp.*, in: Crustaceans: Physiological Characteristics, Evolution and Conservation  
638 Strategies. pp. 19–80.
- 639 R Core Team, 2020. R: A language and environment for statistical computing. R Foundation  
640 for Statistical Computing, Vienna, Austria.
- 641 Reeve, M.R., 1969a. Growth, Metamorphosis and Energy Conversion in the Larvae of the  
642 Prawn, *Palaemon serratus*. J. Mar. Biol. Assoc. U. K. 49, 77–96.  
643 <https://doi.org/10.1017/S0025315400046427>
- 644 Reeve, M.R., 1969b. Laboratory culture of the prawn *Palaemon serratus*.
- 645 Richard, P., 1978. Tolérance aux températures extrêmes de *Palaemon serratus* (Pennant):  
646 Influence de la taille et de l’acclimatation. J. Exp. Mar. Biol. Ecol. 35, 137–146.  
647 [https://doi.org/10.1016/0022-0981\(78\)90072-2](https://doi.org/10.1016/0022-0981(78)90072-2)
- 648 Rodriguez, A., 1981. Growth and sexual maturation of *Penaeus kerathurus* (Forskäl, 1775) and  
649 *Palaemon serratus* (Pennant) in salt ponds. Aquaculture 24, 257–266.  
650 [https://doi.org/10.1016/0044-8486\(81\)90061-2](https://doi.org/10.1016/0044-8486(81)90061-2)
- 651 Sandifer, P.A., Smith, T.I.J., 1979. Possible significance of variation in the larval development  
652 of palaemonid shrimp. J. Exp. Mar. Biol. Ecol. 39, 55–64. [https://doi.org/10.1016/0022-](https://doi.org/10.1016/0022-0981(79)90004-2)  
653 0981(79)90004-2
- 654 Sastry, A.N., McCarthy, J.F., 1973. Diversity in metabolic adaptation of pelagic larval stages  
655 of two sympatric species of brachyuran crabs. Neth. J. Sea Res. 7, 434–446.  
656 [https://doi.org/10.1016/0077-7579\(73\)90064-1](https://doi.org/10.1016/0077-7579(73)90064-1)
- 657 Small, D.P., Calosi, P., Boothroyd, D., Widdicombe, S., Spicer, J.I., 2015. Stage-Specific  
658 Changes in Physiological and Life-History Responses to Elevated Temperature and  
659 Pco2 during the Larval Development of the European Lobster *Homarus gammarus* (L.).  
660 Physiol. Biochem. Zool. 88, 494–507. <https://doi.org/10.1086/682238>
- 661 Temming, A., Günther, C., Rückert, C., Hufnagl, M., 2017. Understanding the life cycle of  
662 North Sea brown shrimp *Crangon crangon*: a simulation model approach. Mar. Ecol.  
663 Prog. Ser. 584, 119–143. <https://doi.org/10.3354/meps12325>



- 664 Thessalou-Legaki, M., Peppas, A., Zacharaki, M., 1999. Facultative lecithotrophy during larval  
665 development of the burrowing shrimp *Callinassa tyrrensis* (Decapoda: Callinassidae).  
666 *Mar. Biol.* 133, 635–642. <https://doi.org/10.1007/s002270050503>
- 667 Torres, G., Giménez, L., 2020. Temperature modulates compensatory responses to food  
668 limitation at metamorphosis in a marine invertebrate. *Funct. Ecol.* 34, 1564–1576.  
669 <https://doi.org/10.1111/1365-2435.13607>
- 670 Torres, G., Melzer, R.R., Spitzner, F., Šargač, Z., Harzsch, S., Gimenez, L., 2021. Methods to  
671 study organogenesis in decapod crustacean larvae. I. larval rearing, preparation, and  
672 fixation. *Helgol. Mar. Res.* 75, 3. <https://doi.org/10.1186/s10152-021-00548-x>
- 673 Torres, G., Thomas, D.N., Whiteley, N.M., Wilcockson, D., Giménez, L., 2020. Maternal and  
674 cohort effects modulate offspring responses to multiple stressors. *Proc. R. Soc. B Biol.*  
675 *Sci.* 287, 20200492. <https://doi.org/10.1098/rspb.2020.0492>
- 676 Urzúa, Á., Anger, K., 2013. Seasonal variations in larval biomass and biochemical composition  
677 of brown shrimp, *Crangon crangon* (Decapoda, Caridea), at hatching. *Helgol. Mar. Res.*  
678 67, 267–277. <https://doi.org/10.1007/s10152-012-0321-4>
- 679 Vázquez, M.G., Bas, C.C., Spivak, E.D., 2013. Seasonal variation in reproductive traits of the  
680 oriental shrimp *Palaemon macrodactylus* (Crustacea: Caridea: Palaemonidae) in a non-  
681 native population. *Helgol. Mar. Res.* 67, 749–756. <https://doi.org/10.1007/s10152-013-0360-5>
- 682 Walsh, C.J., 1993. Larval Development of *Paratya australiensis* Kemp, 1917 (Decapoda:  
683 Caridea: Atyidae), Reared in the Laboratory, With Comparisons of Fecundity and Egg  
684 and Larval Size Between Estuarine and Riverine Environments. *J. Crustac. Biol.* 13,  
685 456–480. <https://doi.org/10.2307/1548788>
- 686 Wehrmann, I.S., Albornoz, L., 1998. Larval development of *Nauticaris magellanica* (A. Milne  
687 Edwards, 1891) (Decapoda: Caridea: Hippolytidae), reared under laboratory conditions.  
688 *Bull. Mar. Sci.* 62, 45–72.
- 689 Weiss, R., Torrecilla, Z., González-Ortegón, E., González-Tizón, A.M., Martínez-Lage, A.,  
690 Schubart, C.D., 2018. Genetic differentiation between Mediterranean and Atlantic  
691 populations of the common prawn *Palaemon serratus* (Crustacea: Palaemonidae) reveals  
692 uncommon phylogeographic break. *J. Mar. Biol. Assoc. U. K.* 98, 1425–1434.  
693 <https://doi.org/10.1017/S0025315417000492>
- 694 West-Eberhard, M.J., 2003. *Developmental Plasticity and Evolution*. Oxford University Press.
- 695 Wickins, J.F., 1972. The food value of Brine Shrimp, *Artemia salina* L., to larvae of the prawn,  
696 *Palaemon serratus* Pennant. *J. Exp. Mar. Biol. Ecol.* 10, 151–170.  
697 [https://doi.org/10.1016/0022-0981\(72\)90100-1](https://doi.org/10.1016/0022-0981(72)90100-1)
- 698 Yagi, H., Ceccaldi, H.J., 1985. The role of temperature and salinity on moult, metamorphosis  
699 and growth at each larval stage of *Palaemon serratus* (Pennant) (Crustacea, Decapoda,  
700 Palaemonidae). *Ann. Inst. Oceanogr. Paris Nouv. Ser. Paris* 61, 75–93.
- 701 Yúfera, M., Rodríguez, A., 1985. Effect of prey density on feeding rates during larval rearing  
702 of *Palaemon serratus* Pennant (Crustacea: Palaemonidae). *Aquaculture* 50, 31–38.  
703 [https://doi.org/10.1016/0044-8486\(85\)90150-4](https://doi.org/10.1016/0044-8486(85)90150-4)
- 704  
705

706 **Figure captions**

707 Figure 1: Experimental design to study the effects of temperature and laying season on larval  
708 developmental pathways in *Palaemon serratus*. For the 12 females (6 per season), the larvae  
709 are distributed into conical vessels in 3 water baths following the illustrated example of the  
710 larvae of females 1 and 7. The illustrations of the females are modified from the one available  
711 on the Muséum-Aquarium de Nancy website by Marion Arbona.

712

713 Figure 2: (a) Metamorphosis rate (%) and (b) development time (days) of *Palaemon serratus*  
714 larvae having reached metamorphosis. The  $\pm$  S.E. correspond to the deviation of the mean of  
715 each family from the overall mean. Larvae are grouped by laying season and incubation  
716 temperature; W12: Winter 12°C, S12: Summer 12°C, W16: Winter 16°C, S16: Summer 16°C,  
717 W20: Winter 20°C and S20: Summer 20°C (n = 104 larvae per female).

718

719 Figure 3: Theoretical larval intermoult time of *Palaemon serratus* versus rearing temperature,  
720 based on data from bibliography (Reeve, 1969a, squares; Campillo, 1979, filled circles; Lassus  
721 & Maggi, 1980, cross; Yagi & Ceccaldi, 1985, stars; Kelly *et al.*, 2012, filled squares;  
722 González-Ortegón & Giménez, 2014, filled triangle).

723

724 Figure 4: The relative frequency (%) of the total number of larval instars reached by  
725 metamorphosed *Palaemon serratus* larvae. Larvae are grouped by laying season and incubation  
726 temperature; W12: Winter 12°C (n = 18), S12: Summer 12°C (n = 5), W16: Winter 16°C (n =  
727 186), S16: Summer 16°C (n = 257), W20: Winter 20°C (n = 271) and S20: Summer 20°C (n =  
728 350).

729

730 Figure 5: Number of *Palaemon serratus* larvae observed for each larval stage during their  
731 development at 12°C. Larvae are grouped by clutch; 1 to 6: winter females, 7 to 12: summer  
732 females (n = 104 larvae per female). Juveniles are removed at metamorphosis and their  
733 cumulative numbers are shown in black on the graph. Observations stop when there are no more  
734 larvae.

735

736 Figure 6: Number of *Palaemon serratus* larvae observed for each larval stage during their  
737 development at 16°C. Larvae are grouped by clutch; 1 to 6: winter females, 7 to 12: summer  
738 females (n = 104 larvae per female). Juveniles are removed at metamorphosis and their  
739 cumulative numbers are shown in black on the graph. Observations stop when there are no more  
740 larvae.

741


742 Figure 7: Number of *Palaemon serratus* larvae observed for each larval stage during their  
743 development at 20°C. Larvae are grouped by clutch; 1 to 6: winter females, 7 to 12: summer  
744 females (n = 104 larvae per female). Juveniles are removed at metamorphosis and their  
745 cumulative numbers are shown in black on the graph. Observations stop when there are no more  
746 larvae.

747

748

**Graphical abstract**


Most frequent developmental pathways in relation to temperature with winter and summer larvae



749

750

**751 Figures**




752

753 Figure 1

754






755

756 Figure 2


757



758

759 Figure 3


760



761

762 Figure 4


763



764

765 Figure 5


766



767

768 Figure 6

769



770

771 Figure 7

### **Highlights**

- Larval development was plastic in terms of instars and stages.
- Higher temperature decreased the development time and the number of instars.
- The seasonal phenotype had no effect on the development time.
- Supernumerary stages were observed at low temperatures.
- Early morphogenesis was observed at high temperatures.

**Declaration of competing interest**

None.

Journal Pre-proof

## **Funding**

This work was supported by the EMFF Gedubouq program of the European Union.

Journal Pre-proof